Genetic diversity and evolution of rice centromeres

Lingjuan Xie[1,2,9], Yujie Huang[1,9], Wei Huang[3,9], Lianguang Shang[2,4,9], Yanqing Sun[5,6], Quanyu Chen[6], Shuangtian Bi[3], Mingyu Suo[6], Shiyu Zhang[1], Chentao Yang[7], Xiaoming Zheng[2], Weiwei Jin[3,8], Qian Qian[2], Longjiang Fan[1,2,*] & Dongya Wu[1,5,6,*]

[1] Institute of Crop Science & Institute of Bioinformatics, Zhejiang University, Hangzhou, 310058, China

[2] Yazhouwan National Laboratory, Yazhou District, Sanya, 572024, China

[3] State Key Laboratory of Maize Bio-breeding, National Maize Improvement Center, Frontiers Science Center for Molecular Design Breeding, Department of Plant Genetics and Breeding, China Agricultural University, Beijing 100193, China

[4] Shenzhen Branch, Guangdong Laboratory of Lingnan Modern Agriculture, Genome Analysis Laboratory of the Ministry of Agriculture and Rural Affairs, Agricultural Genomics Institute at Shenzhen, Chinese Academy of Agricultural Sciences, Shenzhen, 518120, China

[5] Liangzhu Laboratory, Zhejiang University Medical Center, Hangzhou, 310058, China

[6] Center for Evolutionary & Organismal Biology, Zhejiang University, Hangzhou, 310058, China

[7] State Key Laboratory of Agricultural Genomics, BGI Research, Shenzhen 518083, China

[8] Tianjin Key Laboratory of Intelligent Breeding of Major Crops, Fresh Corn Research Center of BTH, Tianjin Agricultural University, Tianjin, 300384, China

[9] These authors contributed equally: Lingjuan Xie, Yujie Huang, Wei Huang, Lianguang Shang

E-mail: fanlj@zju.edu.cn to L.F.; wudongya@zju.edu.cn to D.W.;

Abstract

Understanding the mechanisms driving centromere evolution is crucial for deciphering eukaryotic evolution and speciation processes. Despite their widely recognized characteristics of conserved function in cell division, the centromeres have showed high diversity in composition and structure between species. The mechanism underlying this paradox remain poorly understood. Here, we assembled 67 high-quality rice genomes from *Oryza* AA group, encompassing both Asian and African rice species, and conducted an extensive analysis of over 800 nearly complete centromeres. Through *de novo* annotation of satellite sequences and employing a progressive compression strategy, we quantified the local homogenization and multi-layer nested structures of rice centromeres and found that genetic innovations in rice centromeres primarily arise from internal structural variations and retrotransposon insertions, along with a certain number of non-canonical satellite repeats (*sati*). Despite these rapid structural alterations, the single-base substitution rate in rice centromeres appears relatively lower compared to the chromosome arms. Contrary to the KARMA model for *Arabidopsis* centromere evolution, our model (RICE) suggests that centrophilic LTRs contribute to the decline of progenitor centromeres composed of satellite repeats, and facilitate the formation of evolutionary neo-centromeres, which are enriched with extended CENH3 binding regions beyond the native satellite arrays in plant genomes. In summary, this study provides novel insights into genomic divergence and reproductive barriers among rice species and subspecies, and advances our understanding of plant centromere evolution.

Introduction

Centromeres are the chromosomal regions that are physically linked to the spindle microtubules during mitosis and meiosis. They are ubiquitous and essential in eukaryotic life for maintaining their genetic stability (Barra & Fachinetti, 2018; Naish & Henderson, 2024). Most centromeres are mainly composed of highly repetitive tandem satellite DNA sequences as the sites of kinetochore assembly and interspersed transposon elements (TEs), but their monomer sequences, satellite copy numbers and array organizations are diversely variable (Naish *et al*., 2021; Ahmed *et al*., 2023; Naish & Henderson, 2024). While canonical high-order repeat (HOR) structures are prevalent in human centromeres, such arrangements are less common in plant centromeres (Naish *et al*., 2021; Altemose *et al*., 2022; Wang *et al*, 2023; Logsdon *et al*., 2024). Conversely, plant centromeres frequently harbour centrophilic transposon insertions, a feature that distinguishes them from centromeres in the human and chicken (Cheng *et al*., 2002; Naish *et al*., 2021; Altemose *et al*., 2022; Chen *et al*., 2023; Huang *et al*., 2023; Lv *et al*., 2023). The functional determination of centromeres often relies on the presence of the centromere-specific histone variant CENH3 or CENP-A, which marks the sites of kinetochore assembly (Malik & Henikoff, 2009; Kursel & Malik, 2016). The genomic regions of centromeres defined by satellite DNA occupancy and CENH3 Chromatin Immunoprecipitation Sequencing (ChIP-seq) enrichment can sometimes show discrepancies, suggesting instances of centromere repositioning and the emergence of evolutionary-new centromeres (ENCs) (Liu *et al*., 2023). However, the mechanisms driving the occurrence of ENCs remain incompletely understood. Despite recent advancements in generating complete centromere assemblies using PacBio high-fidelity (HiFi) sequencing and Oxford Nanopore Technology (ONT) ultra-long sequencing, our understanding to the rapid centromere evolution is still limited, owing to the challenges in comparing highly repetitive sequences among species and the limited sampling within species (Xie *et al*., 2024). So far, comprehensive analyzes on population-scale centromeres with high accuracy and continuity have been only performed in the genomes of human and model species *Arabidopsis*, which display the undiscovered genomic diversity and provide valuable resources for centromere genomics (Wlodzimierz *et al*., 2023; Logsdon *et al*., 2024). Extending such analyses to diverse species will be crucial for capturing the trajectory of centromere evolution and understanding its implications for genome stability and speciation.

Rice is one of the major staple crops globally, serving as a primary energy source for human. It is also one of the major model plant species. Recurrent domestication, improvement and de-domestication events have been parallelly experienced for different species or subspecies in the rice AA genome group (Stein *et al*., 2018). Asian rice (*Oryza sativa*) and African rice (*O. glaberrima*) are the two main crops in *Oryza* AA group with a divergence time less than 1 million years, domesticated from their wild ancestor

Oryza rufipogon and *Oryza barthii*, respectively (**Fig. 1a**). Three telomere-to-telomere (T2T) genome

assemblies of Asian rice (varieties Zhenshan97, Minghui63 and Nipponbare) have been completely

assembled (Song *et al*., 2021; Shang *et al*., 2023). Notwithstanding the achievements in generating T2T

assemblies, the precise alignment of highly repetitive sequences within centromeres remains a

significant challenge. Consequently, systematic comparisons of centromeres across different rice

species have been limited. Here we assemble and annotate the centromeres from 70 nearly T2T rice

genomes of *Oryza* AA group. We introduce a novel framework for centromere analysis and perform

comparative investigations of rice centromeres, probing into the satellite sequence composition, satellite

array organization, the impact of transposon invasion and the dynamics of epigenetic modifications

among species, subspecies, groups, individuals and chromosomes. Through elucidating the intricate

mechanisms governing rice centromere evolution, our study aims to offer new insights into the genetic

and epigenetic factors that shape these crucial genomic regions.

92 Results

93 Assembling rice centromeres

To fully capture the genomic diversity of rice centromeres, we collected 70 nearly complete sequences

of *Oryza* AA genomes, incorporating newly assembled sequences from 67 rice genomes using PacBio

HiFi and ONT long reads and three previously released T2T assemblies of Nipponbare (NIP)(Shang *et*

al., 2023), Minghui63 (MH63) and Zhenshen97 (ZS97)(Song *et al*., 2021) (**Supplementary Table 1**).

Briefly, the 70 rice AA genomes encompassed 11 from wild *O. rufipogon* (hereafter referred to as Oruf),

18 from *O. sativa* subsp. *Geng*/*japonica* (GJ), 25 from *O. sativa* subsp. *Xian*/*indica* (XI), six from *aus*

(AUS), along with four wild from *O. barthii* (Obar) and four cultivated from *O. glaberrima* (Ogla) of

African rice, collectively covering a wide spectrum of genetic diversity in Asian and African rice (**Fig.**

1a). In detail, Asian GJ rice samples comprised 13 accessions from subgroup temperate GJ (GJ-tmp),

five from tropical GJ (GJ-trp), and XI included 12 from XI-1A, seven from XI-1B, two from XI-2 and

six from XI-3 (**Fig. 1a**). Wild *O. rufipogon* population was composed of Or-1/2 ($n = 4$, closest ancestor

of XI and AUS), Or-3 ($n = 6$, closest ancestor of GJ) and Or-4 ($n = 1$, basal group of all *O. sativa*

samples).

Genome assemblies were generated using PacBio HiFi (36.0-fold coverage in average with a N50 length

of ~19.0 Kb) and ONT data (~83.3-fold with a N50 length of ~33.3 Kb) (**Supplementary Table 1**). We

initially used Hifiasm to generate a highly accurate backbone of genome assembly, and then integrated

the Verkko assembly to fill the gaps (Cheng *et al*., 2022; Rautiainen *et al*., 2023). After polishing using

PacBio HiFi and NGS short reads, highly accurate assemblies were generated with an average

consensus quality value (QV) of 64.4 (less than one base error per Mb)(**Supplementary Fig. 1;**

Supplementary Table 1). Whole-genome synteny against the NIP reference assembly confirmed the

absence of large structural errors. Base-resolution assembly quality assessment indicators revealed the

platinum-level quality of the new assemblies. The average whole-genome GCI (Chen *et al.*, 2024), an

assembly continuity index derived from contig N50, was 71.1, surpassing previous gapless assemblies

of MH63 (49.9) and ZS97 (54.1). Assembly quality indicators (AQI) (Li *et al.*, 2023) for regional errors

(R-AQI, 99.0) and structural errors (S-AQI, 98.1) further supported the high quality of the assemblies

(**Supplementary Fig. 1; Supplementary Table 1**).

Centromere regions were delineated based on the presence of the rice centromere-specific satellite

repeat sequence *CEN155* (also known as *CentO,* ~155 bp in length) (Cheng *et al.*, 2002)

(**Supplementary Table 2**). In total of 839 centromeres were identified across the 70 genomes, with one

exception on chromosome Chr07 in wild accession CW06, where no *CEN155*-defined centromere was

found. Notably, 808 (96.2%) of these centromeres were contiguous without assembly gaps (**Fig. 1a;**

Supplementary Fig. 1). Base accuracy assessments revealed that 66.2% of assemblies had "+inf"

(infinite) QV scores for all 12 centromere sequences, indicating no erroneous bases in the centromere

assemblies (**Supplementary Fig. 2a**). At the chromosome level, 92.6% of centromeres exhibited QV

values greater than 60, corresponding to an estimated base accuracy exceeding 99.9999%

(**Supplementary Fig. 2a**). Analysis using VerityMap identified few discordant *k*-mers between the

centromere assemblies of cultivated accessions (*O. sativa* and *O. glaberrima*) and PacBio HiFi reads

(Mikheenko *et al.*, 2020), whereas a few potential erroneous *k*-mer clusters in wild accessions (*O.

rufipogon* and *O. barthii*) were observed, likely due to the higher heterozygosity in the wild populations

(**Supplementary Fig. 2b**). The average GCI score for all centromeres was 95.6 (**Supplementary Fig.**

2c), with only 39 centromeres showing potential gaps after stringent filtering of low-quality alignments

(**Supplementary Table 3**). In other words, 95.2% of assembled centromeres were confirmed to be

gapless. To avoid bias among assemblers when handling highly repetitive sequences (Logsdon *et al.*,

2024), we compared the centromere assemblies generated by Hifiasm and Verkko, and found them to be

highly consistent (**Supplementary Fig. 2d**). 414 (67.3%) out of 615 centromeres were completely

assembled by both Hifiasm and Verkko, with *CEN155* array length differences less than 0.1%, and 538

(87.5%) had differences less than 1%. In summary, we have yielded a high-quality atlas of rice

centromere assemblies with high accuracy and continuity.

Centromere length variation

The L/S ratio (long arm/short arm) and centromere region synteny against the NIP reference assembly revealed the conserved positioning of *CEN155* satellite arrays for each chromosome among individuals (**Extended Data Fig. 1a**). Most chromosomes were metacentric or subcentric, whose centromeres are located near the middle of chromosomes, except Chr04 and Chr09, generally in line with the previous cytological characterization (Cheng *et al*., 2001). Coincidentally, the rice 45S ribosomal DNA (rDNA) repeat arrays were positioned on the short arm of acrocentric Chr09, consistent with the pattern observed in the human genome, implying the potential impact of the 45S rDNA array positioning on the evolution of chromosome karyotypes (Altemose *et al*., 2022; Shang *et al*., 2023).

Centromere lengths varied significantly among rice species, subspecies, individuals and chromosomes (**Fig. 1b**). Across individual genomes, centromere sequences occupied approximately 1.11% of the total genome, ranging from 0.57% to 1.82%, considerably smaller compared to the ~3% (occupied by 171-bp alpha satellites) in the human genome (Altemose *et al*., 2022) and the ~9.6% in the *Arabidopsis thaliana* genome (occupied by 178-bp satellites) (Naish *et al*., 2021)(**Supplementary Table 2**). Among chromosomes, we observed a positive correlation between centromere length and chromosome length, indicating that longer chromosomes tend to have longer centromere sequences (**Extended Data Fig. 1b**; **Supplementary Fig. 3**). Interestingly, this correlation was not significant in wild Asian rice and wild African rice groups, in contrast to cultivated groups (**Supplementary Fig. 3**). Despite the overall positive correlation with chromosome length, centromere length contributed minimally to the variation of chromosome size within the same chromosome, and even showed a negative correlation in some cases (**Supplementary Fig. 4a**). In contrast to the significant role of centromere length in chromosome size variation in *Arabidopsis thaliana* (Lian *et al*., 2024), TEs were identified as the primary source of chromosome size variation in rice (**Supplementary Figs. 4b and 4c**).

At the whole-genome level, Asian rice exhibited longer centromeres compared to African rice, although variations were observed across chromosomes (**Fig. 1b**). Among the different groups, as the outgroup of *O. sativa* and other *O. rufipogon*, CW03 from Or-4 had the longest whole-genome centromere sequence (~6.94 Mb), followed by GJ accessions (~5.58 Mb), while XI (~3.73 Mb), AUS (~4.58 Mb), wild Oruf (~3.38 Mb) and cultivated African rice (~5.00 Mb) exhibited shorter centromeres (**Fig. 1b; Supplementary Table 2**). Within the GJ group, GJ-tmp centromeres (~5.93 Mb) were longer than those of GJ-trp (~4.73 Mb). Different chromosomes showed distinct variations in lengths (**Fig. 1b**). For instance, GJ had significantly longer centromeres than other groups on chromosomes Chr06 and Chr11, potentially due to lineage-specific expansions. The three longest centromeres were all from

chromosomes Chr11 of GJ-tmp, each exceeding 1.4 Mb in length. Interestingly, a 6-Mb inversion

spanning the centromere and one nested 1-Mb inversion near the centromere on Chr06 were observed

(**Extended Data Fig. 1c**). The smaller nested inversion likely originated in the common ancestor of

GJ-trp and GJ-tmp, while the larger inversion occurred specifically in the GJ-tmp lineage. These

structural difference in Chr06 centromeres between subspecies may contribute to the reproductive

isolation in natural hybridization and low recombination rates in artificially recombinant populations

(Wei *et al.*, 2024).

Centromere introgression and fission

Sequence similarity between intra-group centromeres was higher that that between inter-group

centromeres, highlighting the distinct evolutionary trajectories between XI and GJ (e.g. Chr02)

(**Extended Data Fig. 2a; Supplementary Fig. 5**). Notably, Chr05 centromeres showed exceptionally

high similarity between GJ and XI but lower between GJ and AUS, supporting the hypothesis that a

large Chr05 segment in the XI genomes, spanning the centromere, was introgressed from GJ during

speciation or initial domestication (Wu *et al.*, 2022) (**Extended Data Fig. 2b**). Aligning to GJ reference

assembly NIP further supported these findings, revealing reduced structural variations in the flanking

sequences of Chr05 centromeres across most XI samples, relative to other chromosomes. Additionally,

Chr10 centromeres in subgroup GJ-trp showed high similarity to a subgroup of XI, aligning with the

documented introgression events from XI to GJ-trp (Gong & Han, 2022) (**Supplementary Fig. 5**).

Considering the frequent inter-subspecies hybridization in rice breeding, centromere introgression

events affecting various chromosomes have been observed, including well-known cultivars like ZS97,

MH63 and Kasalath (**Supplementary Table 4**). For instance, the Chr04 centromere in XI cultivar

J4155S (alias Jing4155S) displayed a GJ-type centromere, in line with the historical records of the

utilization of GJ (Nongken58 from Japan and Lemont from USA) in the breeding progress of J4155S

(**Extended Data Fig. 2c**).

Although it is widely believed that the meiotic recombination in centromere regions is severely

repressed, we analyzed the centromere sequences in sliding windows to explore potential centromeric

rearrangement events. Representative centromere haplotypes were defined for each chromosome and

then we categorized all the centromere blocks into different haplotypes based on pairwise similarity

(**Supplementary Fig. 5**). Besides the whole-centromere introgression events above, rare centromere

splitting or fission events were observed. Chr04 centromeres in XI, for example, exhibited two distinct

types (X1 and X2) (**Supplementary Fig. 5**) and correspondingly the identity between X1 and X2

showed diagonal inconsistency in the heat-map plot, for which X1 was deleted from X2 with a shorter

length (**Extended Data Fig. 2c**). Similarly, the Chr12 centromere of SL044 (Nantehao, one of the most

essential backbone parents in the hybrid rice breeding), comprised a combination of two XI-type

haplotypes (X3 and X4)(**Fig. 2a**). Collinearity among SL044, X3 (e.g. CW09) and X4 (e.g. NJ11)

centromeres and internal sequence similarity within each centromere supported the structural

combination (**Fig. 2b**). An identical (99.40%) intact LTR element *RETROSAT-2C* was identified both at

the end of X3 and the beginning of X4 (**Fig. 2b; Supplementary Fig. 6**). Phylogenetic analysis using

centromere upstream and downstream SNPs (~1 Mb) indicated consistent topology among SL044 and

the two haplotypes, where the flanking sequences of SL044 centromere was both close to haplotype X4

(e.g. NJ11), suggesting that the Chr12 centromere of SL044 likely did not arise from the fusion of X3

and X4 (**Fig. 2c**). To sum up, rapid evolution and innovation in rice centromeres could be driven by

direct introgression events and large-scale rearrangements such as fissions, duplications and inversions,

although these events were comparatively rare (**Fig. 2d**).

229   Satellite repeat variation

To precisely characterize the sequence compositions of rice centromeres, we annotated the *CEN155*

satellites, TEs and potential genes. Across all centromeres, we identified a total of 1.46 million copies of

CEN155 satellites, predominantly distributed around 155 and 165 bp in length (**Supplementary Fig. 7**).

The average copy number of *CEN155* satellites varied significantly among genomes, with GJ (~28,210)

showing a significantly higher count compared to XI (~16,581), AUS (~18,011) and wild Asian rice

Oruf (~15,203), similar to African rice Ogla (~27,176) and Obar (~29,736). These variations correlated

with differences in centromere length, but markedly fewer than the abundance of *CEN180* satellites in

Arabidopsis thaliana centromeres (~66,131) (Naish *et al*., 2021). The *CEN155* satellite repeats formed

long tandem arrays, often exhibiting bias in strand orientation (**Fig. 3a**). Significant divergence in strand

preferences was observed among groups on each chromosome, with distinct patterns observed between

chromosomes in GJ and XI (**Fig. 3a**).

To comprehensively profile the structures of centromere monomers and trace the evolutionary footprints,

we clustered 6,665 representative *CEN155* sequences (140-170bp in length and >30 occurrences in the

centromere collection) into 15 families based on phylogeny (**Fig. 3b**). Low-frequency and divergent

monomers were assigned according to their least edit distances to the clustered *CEN155* sequences, and

finally 15 superfamilies (SFs) were termed as superfamily A (SF-A) to O (SF-O), with SF-N being the

most prevalent across the 70 rice genomes, comprising approximately 200 thousand occurrences, while

SF-L exhibited the least abundance with fewer than 20 thousand copies (**Fig. 3b**). Each superfamily can

be distinguished by specific motif sequences, which could serve as genetic markers for categorizing

CEN155 satellites (**Fig. 3c**). Notably, a 10-bp structural variation (CATATATTGG) was present in SF-B

to SF-E, but was mostly absent in superfamilies from SF-F to SF-O (**Supplementary Fig. 8a**). Pairwise

edit distances among monomers within the same superfamily were generally lower than those between

CEN155 satellites from different superfamilies, but the results still showed high diversity of monomers

within each superfamily (**Supplementary Fig. 8b**).

The presence of B-box-like and pJα-like motifs can be detected in rice consensus *CEN155* sequences,

analogous to motifs found in human centromeres involved in the kinetochore assembly (Rosandic *et al.*,

2006; Rice, 2020), despite human and rice belonged to different kingdoms (**Fig. 3c**). Alignment with

known B-box-like motifs from various species (marmoset, mouse, house) and rice against the canonical

B-box motif in human (5'-*TTCGTTGGAA*CGGGA-3') refined the conserved B-box motif to

5'-TT*G****A**CGGG-3' (**Fig. 3d**) (Kipling *et al.*, 1994; Kugou *et al.*, 2016; Rice, 2020; Cappelletti

et al., 2023). The presence of B-box and pJα-like elements was significantly enriched in centromere

regions (**Fig. 3e**), with divergence among superfamilies primarily attributed to variations at base

positions A and C, potentially influencing kinetochore function. Although the 10-bp PAV deleted the

TT*G of B-box in the alignment, another TTGG short motif before the 10-bp PAV was present as a

substitute for most superfamilies.

Centromeres on different chromosomes exhibited distinct compositions (**Fig. 3f**), similar as the

centromere structure of *Arabidopsis* genome (Wlodzimierz *et al.*, 2023). For instance, SF-A, D, H, J,

and O predominated on Chr11, Chr10, Chr01, Chr07 and Chr02, respectively, illustrating

chromosome-specific distribution patterns. Proportions of the 15 superfamilies also varied among

taxonomic groups. SF-I was nearly absent in Obar and GJ but comprising 17.5% in XI and AUS

genomes, primarily on Chr07 (**Fig. 3f**). Notably, size expansions of GJ centromeres on Chr06 and Chr11

were attributed to increased SF-N and SF-A monomers, respectively, compared to other groups. These

whole-genome constitutive differences in superfamilies among groups were predominately influenced

by specific chromosomes, harboring chromosome-specific *CEN155* satellites (**Fig. 3f**).

TE invasion in centromeres

Centromere satellites typically arrange head-to-tail to form satellite arrays but such homogenized

structures are occasionally disrupted by TE insertions in plant centromeres (Naish & Henderson, 2024).

We conducted an annotation of inter-*CEN155* sequences to profile the TE insertion landscape in rice

centromeres. Approximately ~85.1% of non-*CEN155* centromeric sequences were composed of TEs.

TE proportions within centromere regions (~22.2%) were significantly lower than those across the

whole genome (~51.9%), suggesting stronger purifying selection against TE insertion in centromeres

compared to chromosome arms (**Supplementary Fig. 9**; **Supplementary Table 2**). Despite shorter

centromeres, XI (~1.01 Mb) and AUS (~1.25 Mb) genomes exhibited more TEs within centromeres

compared to GJ (~0.72Mb). About 77.8% (in length) of TEs in the centromeres belonged to

long-terminal repeat (LTR) retrotransposon superfamily *Gypsy*, much more intensive than the

whole-genome *Gypsy* density (~40.7%), similar as *Arabidopsis* centromeres (Naish *et al*., 2021)

(**Supplementary Fig. 9**).

Across all the 70 genomes, we identified a total of 210,672 intact LTRs and 1,042,341 soloLTRs, with

5,128 intact LTRs and 8,366 soloLTRs located within centromeres. The ratio of intact LTRs *verse*

soloLTRs was significantly higher within centromeres than outside (0.61 *versus* 0.20), yet much lower

than in the *Arabidopsis* genomes (~2.5) (Wlodzimierz *et al*., 2023). Of all the centromeric intact LTRs,

SZ-22 ($n = 3673$, covering 8.1% of the total centromere length), *CRM* ($n = 714$, 1.8%) and *RIRE7* ($n =$

257, 0.6%) were the most abundant three families, collectively accounting for 90.6% (**Fig. 3g**;

Supplementary Table 2). Additionally, 26.0% and 22.1% of *SZ-22*, 31.9% and 27.4% of *CRM*, and

30.2% and 18.2% of *RIRE7* intact elements across the whole genome were significantly enriched within

the centromere and pericentromeric regions, respectively, suggesting evolutionary coupling and

pairwise selection between centrophilic LTRs and satellite repeats (**Fig. 3g**). Phylogeny of the two

flanking LTR sequences of intact retrotransposons revealed close relationships among the three families,

while distant when inferring from the internal sequences (**Fig. 3h**; **Supplementary Fig. 10a**). Although

SZ-22 was the most dominant LTR family in centromeres, it was non-autonomous which lacked reverse

transcriptase (*rt*), RNase H (*rh*) and integrase (*int*) genes, with a significantly shortened intact length

(~4.5 Kb) compared to *RIRE7* and *CRM* (~7.7 Kb) (**Fig. 3h**; **Supplementary Fig. 10b**).

Centromeric *SZ-22* insertions were significantly younger than those in chromosome arms (*P* value =

7.5e-4, Wilcoxon test), although a certain number of ancient *SZ-22* insertions in centromeres were

retained (**Fig. 3i**). Using the sequences flanking *SZ-22* inserted elements, we localized the integrated

sites within the *CEN155* consensus repeat. A total of 2,258 *SZ-22* insertions were found throughout the

full length of *CEN155* sequence, and the highest frequency of insertion occurred at positions 100 to 120

base, coincided with the location of B-box motif (**Fig. 3c**; **Supplementary Fig. 11**). As mentioned

above that frequent reversal events between forward and reverse strand were observed within the

CEN155 arrays (**Fig. 3a**), we noticed that the junction regions of strand reversal were enriched with TE

insertions, where TEs were observed in the 1790 out of all the 2133 (83.9%) junction regions, implying

a potential role of TE insertion in triggering strand reversal of *CEN155* satellites. Taken together, these

features reveal extensive and recurrent invasion of TEs in the *CEN155* satellite arrays, shaping the

structural and evolutionary dynamics of rice centromeres. Furthermore, we annotated a total of 535

candidate centromeric gene models within centromeres across the 70 genomes, none of which showed

evidence of gene expression.

Non-canonical satellite repeats

In addition to TE invasion within the *CEN155* arrays, we identified the presence of 6,809 short

intervening sequences dispersed between adjacent canonical *CEN155* satellites repeats (termed as *sati*

for non-canonical satellite repeat as short intervals of canonical centromeric repeats) (**Extended Data**

Fig. 3a). Some of these intervals were separated by a similar number of *CEN155* satellites, where they

formed CEN155+ arrays. The abundance and size of *sati* sequences varied among groups and

chromosomes (**Extended Data Fig. 3b**; **Supplementary Table 5**). Importantly, the high-frequency

intervals were predominantly derived from the canonical *CEN155* sequences (**Extended Data Fig. 3c**).

In three accessions of Or-3 wild Asian rice, a 207-bp interval (termed as *sati207*) between *CEN155*

satellites on Chr11 was observed 147 times, which indicated group-specific centromeric variation

(**Supplementary Table 5**). The *sati207* sequence consisted of two replicates of 104-bp satellites, but

differed from *sati104* (**Extended Data Fig. 3c**). Satellite *sati99* on Chr03, had expanded in cultivated *O.*

glaberrima compared with wild *O. barthii* (**Supplementary Table 5**).

Furthermore, in African rice, we identified 531 *sati94* sequences with an approximate length of 94 bp,

specially enriched in Chr10 centromeres, and particularly expanded in wild *O. barthii* (e.g. NH284 in

Extended Data Fig. 3d). Interestingly, structural comparison indicated that two clusters of *sati94* in

NH284 (Obar) likely emerged through duplication from a NH265-like centromere (Ogla)(**Extended**

Data Fig. 3d). CENH3 ChIP data confirmed the function of CEN155+ arrays composed of canonical

CEN155 and non-canonical *sati94* satellites(**Extended Data Fig. 3d**). Despite the similarity in the

composition and structural organization of nearby *CEN155* satellite arrays, the distance distribution

between every two adjacent *sati94* satellites within each of the three blocks revealed distinct

arrangements, suggesting differential local duplication and homogenization progresses

(**Supplementary Fig. 12a**). Phylogenetic analysis further elucidated the evolutionary trajectory of

sati94 sequences within the three CEN155+ arrays and the mixed distribution of *sati94* repeats

confirmed the sharing of common ancestor as well as individual-specific expansion (**Supplementary**

Fig. 12b).

Multi-layer nested structure of centromeres

To uncover the structural characteristics of *CEN155* arrays, we employed a progressive compression

strategy simulating the expansion progress in reverse (**Fig. 4a**). This approach revealed localized

sequence homogenization at monomer, dimer, and multimer scales, where the number of

homogenization blocks decreased exponentially with increasing repeat copy size, although the

homogenization patterns varied among groups and individuals (**Fig. 4b**; **Supplementary Fig. 13**).

Across all centromeres, 59.7% (8.77 million) of *CEN155* satellites were found within 36,661 monomer

homogenization regions (moHRs), defined as regions with >5 consecutive satellites from the same

superfamily (**Supplementary Table 6**). *S*uperfamiles A (78.2%), I (95.0%), J (80.5%), K (78.6%) and O

(83.7%) exhibited higher propensity to form moHRs, contrasting sharply with SF-C (0.5%) and G

(0.9%). Chromosomes Chr02, Chr03, Chr07 and Chr12 displayed over 75% of their *CEN155* satellites

organized in moHRs, suggesting higher homogeneity compared to other chromosomes (**Fig. 4c**).

Divergence of monomer homogenization among groups and individuals was in line with their

abundance in each family (**Fig. 3f**; **Supplementary Fig. 13a**). The longest moHR, composed of 5,288

repeats of SF-A on chromosome Chr11 in NIP, spanned approximately 865 Kb. Although *CEN155*

satellites within a moHR belonged to the same SF, finer substructures could be observed similar to the

classic HOR structure in the human genome, where an array composed of divergent satellites was

repeated for several times (Altemose *et al*., 2022). We took the annotation of two centromeres as an

example (**Fig. 4d**). In the centromere of chromosome Chr10 in GJ sample YCW03, despite SF-D being

the most abundant, interruptions by SF-C satellites within SF-D arrays hindered direct sequence

homogenization, which was not reflected from the sequence similarity directly. Satellites from SF-J

were arranged head-to-tail into the largest homogenized moHR in the centromere, composed of 271

CEN155 repeats (**Fig. 4d**). In the largest three SF-J moHRs longer than 100 satellites, the inner identity

dot plots within some moHRs exhibited a similar pattern as HOR arrays. The most typical structure (iii)

involved HORs approximately 11 satellites long, with a prominent *k*-mer distance spectrum peak at

1706 bp. In the centromere of chromosome Chr01 in CW03, SF-E satellites accounted for 65.1% of all

CEN155 repeats. Analysis of the four moHRs within this region revealed extensive local

homogenization, evident from nearly identical *CEN155* sequences observed in the identify dots and

k-mer distance spectrum among satellites (**Fig. 4d**).

Compressing each moHR into a monomer resulted in a simplified centromere string (**Fig. 4a**). We

identified 68,370 dimer arrays in total, notably 4,548 long arrays containing at least 5 consecutive dimer

repeats, totaling 135,920 dimers, termed as dimer homogenization regions (diHRs), with an average of

29.9 dimers per diHR (**Fig. 4b**; **Supplementary Fig. 13b**; **Supplementary Table 7**). The longest array

was located on chromosome Chr09 of SL187 (GJ) with 1,308 repeats of BG dimer (**Fig. 4b**). The most

frequent dimer arrays were MN arrays (40,218 dimers in 754 diHRs), followed by BG, EH and CH

dimer. Differential compositions of diHRs were evident across chromosomes (**Fig. 4c**). For instance,

MN dimers predominated on chromosomes Chr03, Chr06 and Chr08, while BG dimers were prominent

on chromosomes Chr07 and Chr09 (**Fig. 4c**). The assembly of dimers exhibited non-random patterns,

with the most abundant 15 dimers collectively accounted for 96.1% of all dimers in diHRs. A similar

pattern of dimer organization has been observed in the human alpha satellites, which was assumed to be

linked to the presence of B-box or pJα motif (Rice, 2020). We examined the divergence sites between

the two monomers for each of the top 15 dimers and identified significant divergence (pairwise allele

frequency difference greater than 0.8) at conserved B-box and pJα sites for most dimers, implying the

potential relationship between featured motif sequences and satellite dimerized organization (**Extended**

Data Fig. 4).

In human centromeres, satellites with and without a B-box motif alternate to form a B-box/no-B-box

dimer (Rice, 2020). Interestingly, we observed a significant bias in monomer copy number of dimers,

where one monomer showed a relatively constrained and stable copy number (almost always 1) while

the other exhibited variable copies (forming moHRs) within a diHR, represented as XY_m or X_mY (where

m indicates multiple copies) (**Fig. 4e**). For instance, the centromere of chromosome Chr10 in YCW03

featured four diHRs, with the largest composed of 211 CD repeats. The copy number of satellite SF-C

was fixed at one, while SF-D had variable copies ranging from one to eight (**Fig. 4f**). Similarly, another

diHR with the JK dimer displayed great variation in SF-J copy numbers but consistently one copy for

SF-K. On chromosome Chr01 in CW03, the largest dimer repeat diHR comprised 36 repeats of EH, with

monomer SF-E ranging from one to 115 copies, contrasting with a fixed one copy for SF-H (**Fig. 4f**).

However, this bias in monomer copy number variation for a dimer was not consistently observed across

the whole genome scale (**Supplementary Fig. 14**). As mentioned above, chromosomes Chr03, Chr06

and Chr08 were predominately occupied by MN dimers. However, they exhibited differential copy

number biases, with Chr03 primarily showing MN_m, while Chr06 and Chr08 displayed a mixture of

MN_m and M_mN (**Supplementary Fig. 14**). Distinct biases were also noted among groups. For instance,

the BO bias was reversed in XI compared with other groups (**Supplementary Fig. 14**). Although the

rules governing monomer homogenization within dimers were not universally unified across the

whole-genome or chromosome scale or among groups, the local bias was evident (**Supplementary Fig.**

15). Furthermore, in addition to the repetitive occurrences of the same dimer within a diHR, the copy

number bias was repeated locally and commonly, strongly suggesting localized duplication as an

approach for centromere expansion (**Fig. 4f**; **Supplementary Fig. 15**).

Compressing each diHR into a single dimer yielded a total of 1,383 multimer homogenization regions

(muHRs), comprising 418 distinct repeat structures and 8,768 multimer repeats, averaging 6.34 copies

per muHR (**Figs. 4a and 4b**; **Supplementary Fig. 13c**; **Supplementary Table 8**). Eight muHRs

contained more than 50 replicates of multimer, with the most repetitive multimer being HOJ with 81

copies (**Fig. 4b**; **Supplementary Fig. 16**). The multimers FLNL, AEAH and HOJ were the most

frequent, with 754, 590 and 373 repeats, respectively, predominantly chromosomes Chr05, Chr11 and

Chr12 (**Fig. 4c**). In summary, we propose a multi-layer nested model of the rice centromere using a

progressive compression strategy. Our findings suggest that centromere expansion occurred through

local duplication of satellites at the monomer, dimer, multimer and their combination levels,

consequently leading to the intricate complexity of the centromere.

Centromere divergence

Distinct variations in centromere composition and structure are evident among different chromosomes

and taxa, as mentioned above. However, the evolutionary dynamics and trajectories at the individual

satellite scale remained largely unexplored, either in rice or other species, due to the extremely high

similarity among satellites and the challenges in high-resolution synteny characterization among

centromeres through direct sequence alignments. Here, we developed a method, SynPan-CEN, designed

to identify syntenic satellite pairs among centromeres, by leveraging satellite clustering information and

pairwise genetic distances (**Fig. 5a; Supplementary Fig. 16**). Initially, we identified syntenic *CEN155*

pairs relative to NIP (GJ-tmp) centromeres in the other rice genomes. The synteny ratio decreased

rapidly with increasing genetic divergence, with approximately 68.0% and 55.3% of *CEN155* satellites

found to be syntenic within GJ-tmp and GJ-trp, respectively, significantly higher than between NIP and

AUS (19.0%) or XI (17.4%)(**Supplementary Fig. 17**). A substantially higher syntenic relationship was

observed between XI and GJ on chromosome Chr05 (61.7%), compared to other chromosomes (22.2%)

(**Supplementary Fig. 18**), in line with the hypothesized introgression events from GJ to XI (Wu *et al*.,

2022).

We further examined the syntenic relationships of satellites between centromeres, to investigate

genomic footprints of local expansion and contraction in GJ centromeres relative to the reference NIP

and XI centromeres compared to the reference NJ11 (**Fig. 5b**). We defined centromere structural

variations (CSVs) as instances where *CEN155* satellite copy numbers showed large expansions

(LE, >50 but <1000 copies) or contractions (LC), and small expansions (SE, <50 but >5 copies) or

contractions (SC). Across 17 and 23 samples compared to GJ and XI references, we identified 7,520 and

5,539 CSVs, encompassing approximately 140.8 thousand and 87.3 thousand copies of *CEN155*

satellites, respectively (**Fig. 5c; Supplementary Table 9**). Large LCs and LEs accounted for 5.7% of all

CSVs in count (~14.7 and ~13.7 per GJ sample, ~4.5 and ~7.2 per XI sample), yet represented 45.6% of

CSV satellite copy number variations. On average, 17.0 and 17.9 copies of *CEN155* were included per

contraction and expansion variation, respectively. The exponential distribution of CSVs in terms of size

was in line with the length distribution spectrum of SVs on the chromosome arms (Qin *et al*., 2021;

Shang *et al*., 2022)(**Fig. 5c**). Detected CSV counts ranged from 165 (SL167/Jindao 1, alias B236 in the

3000 Rice Genome Project) to 697 (SL129/Keluoduo B, alias B179 from France) for GJ, and from 158

(CX20) to 351 (Balatinaw, a glutinous and red heritage rice) for XI, respectively, depending on the

genetic distances to their respective reference genomes (**Figs. 5a & 5d**). Notably, In XI, the accession

closest to NJ11 in centromeric kinship was CX20, consistent with their closest evolutionary relationship

at the whole-genome scale (**Figs. 1a, 5a and 5d**). CX20 is a weedy rice accession, presumed to have

originated from the de-domestication of the modern cultivar NJ11, widely cultivated throughout South

China 30 years ago (Qiu *et al*., 2020).

Centromere regions are known for their accelerated evolutionary rates compared to the chromosome

arms, characterized by significant compositional differences observed among closely related species or

populations (Bensasson, 2011; Wlodzimierz *et al*., 2023; Logsdon *et al*., 2024; Minton, 2024). For

instance, human centromeres exhibit higher mutation rates (1.1~4.1 times) in the pericentromeric

regions compared with non-human primates and the active HOR arrays were presumed to have an

increased mutation rate greater than ten times (Logsdon *et al*., 2024). The high repetitiveness of

centromere regions complicates biologically meaningful sequence alignments and comparisons. Instead

of traditional sliding window alignment strategy, we utilized the satellite-resolved synteny framework

between centromeres to profile the evolutionary rates of rice centromeres. Considering the nearly

complete turnover of *CEN155* arrays between XI and GJ for certain chromosomes (**Fig. 3f**; e.g. Chr07),

comparison between GJ and XI were omitted, focusing instead on the divergence between GJ samples

and the reference assembly NIP, and between XI samples and the reference assembly NJ11, respectively.

To eliminate false positives from poor alignment, we screened 91 chromosome pairs with a syntenic

ratio exceeding 0.9 (**Supplementary Table 10**). We defined "divergence rate" to quantify the sequence

difference between alignable syntenic satellites and "mutation rate" to quantify the *de novo* changes in

absolute orthologous satellite pairs. In other words, syntenic satellites are not always orthologous. We

calculated the biological edit distances (counts of mismatches and gap openings) between DNA

sequence alignments from the gene exonic, gene intronic, intergenic regions, and the *CEN155* satellite

arrays. Due to the limited evolutionary timescale within GJ and XI groups, we focused on relative

evolution rates in centromeres compared to chromosome arms, rather than the absolute evolution rate

over generation. Among the 91 chromosome pairs, centromere divergence rate averaged 4.1 times

higher than those in chromosome arms, though varied widely (0 to 56.4), depending on chromosomes

and individuals (**Fig. 5e**). We zoomed in on the *CEN155* arrays to investigate the source of genetic

divergence. Despite similar centromere composition and structure in the compared chromosome pairs, a

few of local syntenic *CEN155* pairs were not orthologs. For instance, SL032 and NIP on chromosome

Chr02 displayed nearly identical centromeres with only two small CSVs (one 11-copy SC and one

7-copy SE) and one tiny CSV (3-copy contraction) (**Fig. 5a**). Notably, a few of syntenic *CEN155* pairs

around the CSVs exhibited higher ED values compared to other pairs. To exam whether the high

sequence divergence was contributed from high mutation rate or exaggerated genetic distance, we built

the phylogeny using all *CEN155* satellites and found that *CEN155* satellites in the high-ED syntenic

pairs were not clustered monophyletically, but instead located distantly in the phylogeny, which

suggested that such pairs were not orthologous but paralogous, and thus could not be used to measure

the mutation rate (**Fig. 5f**). Besides dispersal high-ED syntenic pairs, we also observed clustered

syntenic *CEN155* pairs with evaluated ED values among centromeres. For example, comparing the

centromeres of NJ11 and CX20 on chromosome Chr03 revealed a cluster of high-ED pairs (**Fig. 5a**).

Phylogenetic analysis confirmed these pairs were distantly related due to non-orthologous origins (**Fig.**

5g). Additionally, the *CEN155* composition differed significantly, with NJ11 containing a higher

proportion of *CEN155* satellites from super-family O, possibly due to specific gene conversion or

segmental replacement from homologous regions, as evidenced by comparison with outgroup sample

LK638S (XI-1B). Therefore, to mitigate the significant impact of overestimated genetic distances

between potential paralogous syntenic pairs in the mutation rate analyses, chromosome pairs were

excluded where syntenic *CEN155* syntenic satellite pairs were determined non-orthologous by the

CEN155 satellite phylogeny (**Fig. 5h**; **Supplementary Table 10**). Analysis of the 17 chromosome pairs

with uniformly orthologous *CEN155* satellites revealed significantly lower mutation rates (~0.11 times)

compared to chromosome arms (**Fig. 5i**). In particular, three centromeres on Chr05 and two on Chr10 in

XI samples mirrored reference centromeres in NJ11 (**Supplementary Fig. 19**). In summary, based on

the pan-centromere analysis above, rice centromeres exhibit higher sequence divergence rates than

chromosome arms, and this divergence primarily stems from non-ortholog gene conversion or

segmental substitution rather than *de novo* mutations. Conversely, *de novo* mutation rates in rice

centromeres are notably lower than those in chromosome arms, which may be explained by the

hyper-methylation nature of centromeres.

Epigenetic variation and repositioning of centromeres

We conducted ChIP-Seq experiments using specific anti-CENH3 antibody to map the functional

centromeres in rice, revealing widespread repositioning and neocentromere formation across multiple

genomes (**Supplementary Figs. 20-29**). On average, the CENH3-enriched sequences spanned

approximately 8.22 Mb across the thirteen profiled genomes, significantly exceeding the size of

centromere regions defined by *CEN155* arrays (~4.35 Mb), in contrast to the homeostasis of CENH3

loading in *Arabidopsis* (Naish *et al*., 2022; Wlodzimierz *et al*., 2023) and human centromeres (Logsdon

et al., 2024) but comparable to observations in maize centromeres (Chen *et al*., 2023) (**Supplementary

Table 11**). Detailed analysis revealed that on 73 out of 156 chromosomes, the CENH3-enriched regions

exceeded 1.5 times the size of the *CEN155*-defined centromere regions. Notably, chromosomes such as

Chr01, Chr02, Chr03 and Chr07 exhibited highly consistent positional matching between

CENH3-defined and *CEN155*-defined centromere regions, while other chromosomes showed

significantly longer CENH3-defined centromeres (**Supplementary Fig. 30**).

To investigate factors related to centromere repositioning and neocentromere formation, we examined

the pairwise correlations for each chromosome among chromosome length, *CEN155* array length,

CENH3 occupancy length, TE density within and outside the *CEN155* array, and neocentromere

formation tendency (NFT) (defined by the length ratio between CENH3 enrichment region and *CEN155*

array)(**Fig. 6a**; **Supplementary Figs. 30-36**). Generally, for a given chromosome, the diversity in

chromosome length was attributed from TE expansion (**Supplementary Fig. 4**). Longer chromosome

tended to exhibit more intense TE insertion within their centromeres, coupled with shorter *CEN155*

array (**Fig. 6a**; **Supplementary Figs. 31 and 32**), suggesting the erosion of the *CEN155* array by TE

activity. Longer chromosomes also required extended functional centromeres as defined by CENH3

occupancy, particularly noticeable in chromosomes with significant length variation (**Fig. 6a**;

Supplementary Fig. 33). We observed positive relationship between centromeric TE density and NFT

and negative correlations between the *CEN155* array length and NFT (**Fig. 6a**; **Supplementary Figs. 34

and 35**). Consequently, longer chromosomes among individuals for a given chromosome have higher

density of TE insertion, shortened *CEN155* arrays and a greater propensity for outward expansion (**Fig.

6a**; **Supplementary Fig. 36**).

On chromosome Chr04, the CENH3 binding enrichment regions of African rice Ogla (NH265) and Or-4

(CW03) aligned with the *CEN155* arrays (~400 Kb), indicative of the ancestral state of kinetochore

assembly sites (**Fig. 6b**). Conversely, other Chr04 centromeres exhibited significantly shortened

CEN155 arrays (~131 Kb) yet displayed variable and expanded CENH3 binding regions (~808 Kb) (**Fig.**

6b; **Supplementary Table 11**). Compared with CW15 (Or-3, ancestral group of GJ), GJ expanded a

subregion of CENH3 binding enrichment on the left of its *CEN155* array, accompanied with the

contraction of *CEN155* arrays from ~185 Kb to ~117Kb. Notably, despite minimal structural variations

between the centromeres of GJ-trp (SL177) and GJ-tmp (NIP), GJ-trp (SL177) had a distinct ChIP-seq

peak on the right of its *CEN155* array, suggesting either the emergence of a potential neocentromere in

GJ-trp or the inactivation of the functional centromere on that side of *CEN155* array in GJ-tmp (**Fig. 6b**).

CW11 and CW06 , from group Or-1/2 (ancestral wild group of XI and AUS), exhibited discernible

CENH3 enrichment within their *CEN155* arrays, with supplementary but robust CENH3 binding signals

observed to the right of their *CEN155* arrays in comparison to their progenitor group Or-4 (CW03).

These signals were further amplified in XI (NJ11), implying a functional transition from the *CEN155*

arrays to neocentromeres, concomitant with the shortening of *CEN155* array size from ~172 Kb to ~136

Kb. Notably, *CEN155* array of AUS (SL121) were drastically degraded (~25 Kb), with a distinct

neocentromere established and functioned on the right side of its *CEN155* array. On chromosome Chr11,

distinct neocentromeres were also observed in Or-1/2 (CW11, CW06) and Or-3 (CW15), characterized

by markedly shorter CEN155 arrays compared to other groups (**Supplementary Fig. 28**). These

findings collectively suggest a correlation between the emergence of neocentromeres and the demise of

native satellite arrays.

Chromosome Chr05 had the lowest density of centrophilic TE insertion among the 12 chromosomes

(**Supplementary Fig. 9**). In contrast to the centromeres of outgroup African rice (NH265 and NH284)

and Or-4 (CW03), where *CEN155* arrays matched the CENH3 binding regions, all other centromeres

showed additional CENH3 binding beyond the *CEN155* arrays (**Extended Data Fig. 6**). Moreover,

GJ-trp (SL177) displayed newly evolved CENH3 binding regions compared to GJ-tmp (SL170 and

NIP). Interestingly, comparative analysis of centromere structures revealed a specific *SZ-22* LTR

insertion within the *CEN155* array of GJ-trp (SL177) (**Extended Data Fig. 6**). Consequently, we

propose that TE insertion may trigger and initiate centromere repositioning events leading to

neocentromere formation.

The regions enriched with CENH3 occupancy beyond the *CEN155* arrays were predominantly situated

near the *CEN155* arrays, centrally distributed within pericentromeric regions, where centromeric LTRs

(*SZ-22*, *CRM* and *RIRE7*) were also notably enriched (**Fig. 6b; Supplementary Figs. 20-29**). These

CENH3 signals showed peaks at the centrophilic LTR loci outside the *CEN155* arrays, particularly in the

pericentromeric regions, implying these LTRs may function alternatively as *CEN155* satellites, in spite

of relatively lower enrichment intensity compared to *CEN155* satellites and corresponding LTRs within

the *CEN155* arrays (**Fig. 6c**). Above evidence implied that the function in kinetochore assembly has

driven the evolution of rice centromeric LTRs and centrophilic LTRs appear to mimic the function as

centromere satellites as an adaptive strategy to avoid eradication.

Previous studies on human and *Arabidopsis* centromeres have reported a depletion of DNA methylation

in centromeres, forming centromere dip regions (CDRs) (Naish *et al*., 2021; Wlodzimierz *et al*., 2023;

Logsdon *et al*., 2024). We profiled centromeric DNA methylation using ONT and PacBio HiFi data

(**Supplementary Fig. 39**). Across the entire chromosome, centromere and adjacent pericentromeric

regions displayed higher DNA methylation level than the chromosome arms. Within the centromere

regions, methylation levels showed no obvious difference among *CEN155* arrays and their flanking

regions, with the methylation level of CHH notably lower than CHG and CG methylation (**Fig. 6d**).

Generally, higher CHG methylation, lower CHH methylation and stable CG methylation were observed

in the centromere regions compared to the adjacent pericentromeric heterochromatin, particularly in the

homeostatic centromeres where the *CEN155* arrays aligned with CENH3 occupancy regions (**Fig. 6d**).

The methylation landscape in rice centromeres differs from that in *Arabidopsis* centromeres with a

depletion of CHG and CG methylation, possibly associated with the distinctness of satellite sequences

and the homeostasis of CENH3 loading (Wlodzimierz *et al*., 2023).

Discussion

Investigating the mechanisms driving centromere evolution and their impact on genome structure and

function is essential for understanding eukaryotic evolution and speciation processes. Based on our

comprehensive analyzes, we propose an evolution model of rice centromeres here (termed as

Retrotransposon Induced Centromere Evolution model, RICE), where genetic and epigenetic dynamics

jointly have shaped the rapid evolution of centromeres, emphasizing the importance of LTR insertions in

the centromere evolution cycle (**Fig. 7**). Genetic innovation of centromeres in RICE model primarily

arise from internal structural variations and the retrotransposon insertion from outside. Via

inter-chromosome, individual, group, subspecies and species comparisons in rice centromeres, we have

observed diverse structural variations on multiple scales from recurrently local expansions and

contractions at the monomer, dimer and multimer levels, to larger-scale centromere splitting and

duplication events. Models such as break-induced replication (BIR) model and unequal exchange model

provide plausible explanation for the formation of such complex nested structures (Rice, 2019; Talbert

& Henikoff, 2022). Despite sharing a common *CEN155* repertoire inherited from a common ancestor,

the centromeres in GJ and XI exhibit distinct compositions and structures due to lineage-specific local

homogenization progresses, which may have played an important role in the formation of reproduction

barriers between them. Contrary to the rapid structural alteration, the single-base substitution rate in

centromeres, inferred from absolutely orthologous satellite pairs, appears relatively lower compared to

the chromosome arms. Although rice centromere sizes are diverse among individuals, the variation in

chromosome length is attributed to TE or LTR insertions. Longer chromosomes typically possess

extended functional centromeres, necessary for microtubule attachment during cell division. However,

at the same time excessive LTR insertions would invade the *CEN155* arrays, disrupt CENH3 binding

continuity of kinetochore assembly and potentially trigger centromere repositioning. These centrophilic

LTRs mimic the function of centromere satellites in binding to avoid being purged. In this study, we

have observed the emergence of evolutionary neo-centromeres in rice, characterized by CENH3 binding

regions that extend significantly beyond the satellite arrays. These neo-centromeres often form in close

proximity to the progenitor centromere and are notably enriched with centrophilic LTRs. Unlike the

KARMA (kinetochore-associated recombination machine in *Arabidopsis*) mechanism, where LTR

insertions primarily contribute to satellite expansion (Wlodzimierz *et al.*, 2023), our findings propose a

different role for LTRs in plant centromere evolution. Specifically, we suggest that LTR insertions may

lead to the decline or loss of progenitor centromeres composed of canonical satellite repeats, thereby

facilitating the establishment of neo-centromeres in adjacent regions (**Fig. 7**). For instance, compared to

the centromeres of rice AA genomes investigated in this study, the centromeres of rice GG genome

(*Oryza granulata*) have no canonical tandem repeats but are primarily occupied by *gypsy*-type LTRs,

coresponding to the fact that the rice GG genome has much higher LTR content (Wu *et al.*, 2018).

Besides the rice genomes, the proposed model parallels observations in other genomes such as soybean

and maize, where high LTR retrotransposon contents have been associated with frequent centromere

repositioning (Liu *et al.*, 2023; Chen *et al.*, 2023). Particularly, wheat centromeres have no canonical

satellites but almost are filled by TEs, in line with their extremely high LTR genomic contents (Ahmed

et al., 2023). In contrast, in the human and chicken genomes, CENH3 or CENP-A binding regions are

typically embedded within arrays of higher-order repeats (HORs) at centromeres (Altemose *et al.*, 2022;

Huang *et al.*, 2023). Similarly, in *Arabidopsis*, CENH3 peaks coincide with the presence of satellite

CEN178 or within *CEN178* regions on specific chromosomes (Naish *et al.*, 2021; Wlodzimierz *et al.*,

2023). Based on these comparative insights, we hypothesize that genomes with higher LTR contents

may be more prone to evolving neo-centromeres through mechanisms involving the disruption or

replacement of existing centromeric structures. The TE expansion and polyploidization are pivotal

adaptive strategies for plants to cope with rapid environmental changes. As a by-product of TE activity

to maintain fitness, the rapid centromere alteration may have facilitated the formation of reproduction

barrier and speciation to a certain extent, which suggest a novel insight into the evolutionary trade-off

between survival and reproduction. Further investigations into the precise molecular mechanisms

underlying centromere reorganization in response to LTR insertions will be essential for validating these

hypotheses and expanding our understanding of genome evolution in diverse organisms.

Methods

Pacbio HiFi and ONT sequencing

A total of 70 rice accessions from *Oryza* AA genome group were including in this study, comprising 51 Asian rice *O. sativa*, 11 wild *O. rufipogon*, four African rice *O. glaberrima* and four wild *O. barthii* samples (**Supplementary Table 1**). The rice plants were cultivated at the China National Rice Research Institute (CNRRI), Hangzhou, China. For Pacbio HiFi sequencing, we newly generated Pacbio HiFi data for 46 accessions, and the HiFi reads for 24 accessions were adopted from previous studies (Wu *et al.*, 2023; Shang *et al.*, 2023; Zhang *et al.*, 2022; Sedeek *et al.*, 2023). High-molecular-weight DNA from a single individual was extracted from young seedlings for each accession using the CTAB method. DNA degradation and contamination was scanned on 1% agarose gels. DNA purity and concentration were quantified using a NanoDrop One UV-Vis spectrophotometer (Thermo Fisher Scientific, USA) and a Qubit 4.0 Fluorometer (Invitrogen, USA). According to the methods in the SMRTbell prep kit 3.0 kit manual, HiFi libraries for sequencing were prepared, including DNA quality control, DNA shearing and cleanup, DNA repair and A-tailing reaction, adapter ligation, nuclease treatment and library size selection. Sequencing was conducted on the PacBio Revio platform following the manufacturer's instructions..PacBio HiFi reads with quality scores over Q20 were generated from raw sequencing reads using PacBio Circular Consensus Sequencing tool ccs (min passes = 3, min RQ = 0.99). The sequencing depth ranged from 23.8× to 49.5× across genomes with an average of 30.4× (**Supplementary Table 1**). For ONT sequencing of ten samples, genomic DNA was extracted using a QIAGEN Genomic DNA extraction kit (Cat #13323, QIAGEN). DNA purity and concentration were determined using a NanoDrop One UV-Vis spectrophotometer (Thermo Fisher Scientific, USA)and a Qubit 3.0 Fluorometer (Invitrogen, USA), respectively. Long DNA fragments were selected using the PippinHT system (Sage Science, USA) via gel cutting. Subsequently, DNA repair and adapter ligation were performed using an SQK-LSK110 kit. The prepared DNA library was then loaded onto a PromethION flow cell (R10.4.1). Base modification information was called using Guppy (v6.4.6). Reads with quality scores less than seven were discarded. For 50 samples, ONT reads were from previous studies (Shang *et al.*, 2022; Zhang *et al.*, 2022) and downloaded from NCBI Sequence Read Archive (BioProjects PRJNA656318 and PRJNA692836) and National Genomics Data Center (BioProject PRJCA008812). Additionally, available whole-genome NGS data for 64 accessions were adopted from previous studies (Shang *et al.*, 2022; Shang *et al.*, 2023) (**Supplementary Table 1**).

CENH3 Chromatin immunoprecipitation and sequencing (ChIP-seq)

ChIP experiments for ten samples were performed following standard protocols (Nagaki *et al.*, 2003). Approximately 5 g of 4-week-old seedlings were well grounded in liquid nitrogen for 1 hour and then transferred into 50 ml Eppendorf tube with 20 ml nuclei isolation buffer (10 mM Tris-HCl pH 7.5, 3 mM

$CaCl_2$, 2 mM $MgCl_2$, 0.1 mM PMSF, 1 EDTA-free protease inhibitor cocktail tablets (Roche) and 0.5%

Tween 40). After incubating on ice for 15 min, the mixture was filtered through one layer of Miracloth

(Calbiochem 475855). The liquid was centrifuged at 600 rcf for 10 min at 4□. Discard the supernatant

and resuspend the nuclei pellet with 5 ml 1 × TBS buffer (10 mM Tris-HCl pH 7.5, 3 mM $CaCl_2$, 2 mM

$MgCl_2$, 0.1 mM PMSF, 1 EDTA-free protease inhibitor cocktail tablets (Roche)) with 25% sucrose, and

then added on top of 1 × TBS buffer with 50% sucrose gently and centrifuged at 1,500 rcf for 10 min at

4□. Collect the nuclei and digest the chromatin into nucleosome by MNase in MNB buffer (10%

sucrose, 50 mM Tris-HCl pH 7.5, 4 mM $MgCl_2$, 1 mM $CaCl_2$, 0.1 mM PMSF) for 6 min at 37□. The

nucleosomes were incubated with pre-immune rabbit serum or CENH3 antibody at 4□ overnight, and

was then incubated with rProtein A SepharoseTM Fast Flow (GE Healthcare) for 2 hours at 4□.

Preimmune rabbit serum was used as the negative control. The rProtein A Sepharose was collected by

centrifugation at 13,000 rpm for 1 min at 4□, and washed once using Buffer A (50 mM Tris-HCl pH 7.5,

10 mM EDTA, 0.2 mM PMSF) with 50 mM, 100 mM, and 150 mM NaCl. Immunoprecipitated

DNA-protein complexes were eluted twice from the beads with elution buffer (1% SDS, 50 mM NaCl,

20 mM Tris-HCl pH 7.5, 5 mM EDTA) at 65□ for 15 min. The DNA was extracted with equal volume of

phenol:chloroform (1:1) and the aqueous layer was transferred to new tube. The aqueous layer was then

precipitated with isopropanol, sodium acetate, and glycogen at -20□ for 2 hours, followed by washing

with 70% alcohol once and air-drying. The DNA extraction, precipitation, and washing were performed

by centrifugation at 12,000 rpm for 10 min at room temperature, then resolved with TE buffer and stored

at -20 □. The anti-CENH3 ChIP-seq library was constructed using rabbit polyclonal anti-CENH3

against the rice-specific peptide. ChIP samples were PCR amplified for 12 cycles and sequenced

(2×150bp) on the MGISEQ-200 platform.

Genome assembly

Among the 70 assemblies analyzed, 67 were newly assembled for this study. Three previously released

assemblies (NIP, MH63 and ZS97), assembled completely using extremely high sequencing depth of

long reads, were downloaded from the Rice Information GateWay (RIGW) and Rice Super Pan-genome

Information Resource Database (Shang *et al.*, 2023; Song *et al.*, 2021). Newly generated PacBio HiFi

reads with adaptor sequences were filtered using HiFiAdapterFilt (Sim *et al.*, 2022). For 50 accessions

with both PacBio HiFi and ONT reads, we integrated HiFi and Nanopore data together to generate draft

de novo assemblies using Hifiasm (0.19.5-r587)(Cheng *et al.*, 2024) and Verkko (v1.4)(Rautiainen *et al.*,

2023), respectively. For the remaining 17 accessions with only PacBio HiFi reads available, we used the

HiFi-only mode of Hifiasm and Verkko for assembly, respectively. The scaffolded assembly of Verkko

was split into contigs and used to scaffold and extend Hifiasm contigs using a in-house script. The

extended contigs (larger than 10 Kb) were then scaffolded on chromosomes using RagTag with no

further chimeric splitting (v2.1.0, scaffold -f 2000 --remove-small --aligner minimap2)(Alonge *et al.*,

2022), using T2T assembly of NIP as the reference genome (Shang *et al.*, 2023). To correct base-level

assembly errors and mitigate potential PacBio HiFi sequencing homopolymer errors, we polished the

gap-filled assemblies using both HiFi and NGS reads. High-frequency k-mers (top 0.02%) were

generated using meryl (v1.4)(Rhie *et al.*, 2020) and Winnowmap2 (v2.03)(Jain *et al.*, 2022) aligned HiFi

reads against the assemblies. NextPolish2 (Hu *et al.*, 2024) was then employed for polishing the

assembly with default parameters. Two k-mer datasets (21-mer and 31-mer) were prepared using yak

(https://github.com/lh3/yak). For five red rice accessions (Sedeek *et al.*, 2023) and one weedy accession

YCW03 (Wu *et al.* 2023), assemblies were not polished using short reads, due to their unavailability. To

prevent potential over-scaffolding based on homology, we detected and trimmed the potential mis-joins

at both ends of each chromosome. Telomere regions were identified by searching the rice telomeric

repeat motif TTTAGGG with the occurrence of at least five consecutive repeats. Non-telomeric

sequences outside the telomeric region (< 2 Mb) were then pruned from the assemblies.

Assembly quality assessment

We first manually checked and corrected the potential large structural errors in assembly by aligning to

the NIP reference. Assembly quality for each genome was evaluated using a series of indices from

completeness, correctness and continuity. For completeness, NGS short reads and long reads (PacBio

HiFi and ONT) were aligned to their corresponding chromosomes for each sample with BWA mem

(0.7.17-r1188)(Li & Durbin, 2009) and Winnowmap2 (v2.03)(Jain *et al.*, 2022) in default parameters,

respectively. The mapping statistics was generated by using samtools flagstat (v1.18-15-g9a59467) (Li

et al., 2009). Averagely 97.62% of short reads were aligned, while wild rice *O. rufipogon* had

significantly lower mapping rates (~95.23%) compared to *O. sativa* (~98.35%), due to the nature of

higher heterozygosity in wild population (**Supplementary Table 1**). ~99.92% and ~98.71 of PacBio

HiFi and ONT reads were mapped. The mapping coverage along each chromosome was calculated

using samtools depth and the coverage distribution was checked visually to detect assembly collapsed

regions in 10-Kb windows. For correctness, we used QV and discordant k-mers in VerityMap to assess

the base accuracy. NGS short reads and Pacbio HiFi reads were split into 21-mers and combined as a

k-mer library using meryl (v1.4) (Rhie *et al.*, 2020) and rare k-mers were removed (meryl gt 1).

Assembly correctness indicator QV and assembly completeness indicator k-mer completeness were

calculated using Merqury (v1.3) (Rhie *et al.*, 2020). It should be noted that NGS data used here was

sequenced after polymerase chain reaction (PCR) amplification, thus we analyzed the potential effects

of PCR on QV. We sequenced PCR-free NGS libraries for four samples (CW06, CW15, SL044 and

SL177) using Illumina Novaseq 6000. Using only NGS k-mer library to calculate QV, PCR NGS data

yielded a lower QV than PCR-free data, but varied among samples (46.4 for PCR-free QV *versus* 25.8

for PCR QV in CW06, 47.1 *versus* 23.2 in CW15, 47.4 *versus* 40.1 in SL044 and 43.5 *versus* 44.2 in

SL177). We also employed VerityMap (Bzikadze *et al.*, 2022) to detect assembly errors and

heterozygosity sites in a reference-free mode. VerityMap identifies discordant *k*-mers between the

assemblies and PacBio HiFi reads. High frequency of discordant *k*-mers indicate potential errors in local

assembly. For continuity, we detected potential assembly gaps with low-confidence read supports using

CRAQ (Li *et al.*, 2023) and GCI (Chen *et al.*, 2024)(**Supplementary Fig. 1**). We aligned the PacBio

HiFi reads and NGS reads against corresponding assemblies and calculated AQI values to identify

regional and structural assembly errors based on clipped alignment information. GCI is a continuity

inspector for complete genome assembly, by utilizing stringently filtered alignments from multiple

aligners (minimap2 and Winnowmap2 used in this study) to detect assembly gaps. We quantified the

assembly overall continuity using GCI scores at the whole-genome and chromosome levels.

Evaluation of centromere assembly completeness

To assess the assembly quality of rice centromere sequences, we first determined the centromere regions

based on the enrichment of rice *CEN155* satellites. The quality assessment metrics (e.g. QV. GCI) were

calculated for each centromere (**Supplementary Fig. 2**). We zoomed in and manually checked the raw

and curated reads mapping coverage and the issues reported by GCI and VerityMap in the centromere

regions. Additionally, given the challenges in assembling highly repetitive sequences accurately and

completely, we compared the centromere sequences solely generated by Hifiasm (Cheng *et al.*, 2024)

and Verkko (Rautiainen *et al.*, 2023) to eliminate the effects of assembly algorithms. For each accession,

two sets of contig assemblies from Hifiasm and Verkko with no gap filling and polishing were anchored

onto chromosomes using RagTag, respectively. The *CEN155* consensus sequence was then aligned to

each chromosome using BLASTN (v2.5.0+, -use_index true -task megablast -evalue 1e-6). We counted

the copy numbers of *CEN155* satellites in each centromere and compared the centromeres with no gaps

in both Hifiasm and Verkko assemblies (**Supplementary Fig. 2**).

Phylogeny and ancestry analysis

Short Illumina paired-end reads from each sample were aligned to the T2T NIP reference assembly

(Shang *et al.*, 2023) using BWA (Li & Durbin, 2009). For six samples lacking NGS short reads (YCW03,

Pulut Hitam-2, Zag, Cempo Ireng, Balatinaw, and Cempo Abang), simulated 120-bp reads were

generated from their genome assemblies and mapped against the T2T NIP assembly. Whole-genome

SNPs across all 70 samples were jointly called following the GATK best practices (McKenna *et al.*,

2010). A phylogenetic tree encompassing these 70 rice accessions was constructed using FastTreeMP

(Price *et al.*, 2009) with 1000 bootstrap replicates. This tree was utilized to elucidate the taxonomic

relationships and groupings among the rice accessions.

Gene annotation

Gene elements were annotated using a homology-based approach. We employed Gmap to align
transcripts from the MSU v7.0 annotation of Nipponbare with the parameters '-n 0
--min-trimmed-coverage 0.7 --min-identity 0.7', and kept alignments with at least 70% coverage and 90%
identity (Wu & Watanabe, 2005). Cleaned RNA-seq reads were mapped to the corresponding genome
assembly using TopHat2 (Kim *et al*., 2013), and gene expression was quantified using Cufflinks
(Trapnell *et al*., 2010).

TE annotation

To characterize repeat elements across the 70 high-quality rice genomes, a hybrid approach combining
de novo prediction and homology-based search strategies was employed. Initially, a *de novo* repeat
database was constructed using assemblies from 16 representative rice accessions (including 18XHB-57,
18XHB-83, CW03, CW06, CW11, CW15, MH63, NH271, NH285, NIP, SL028, SL039, SL044, SL121,
YZ-2 and ZS97). The *de novo* transposable element (TE) library was generated separately for each
accession using RepeatModeler with the parameter '-LTRStruct' (Tarailo-Graovac & Chen, 2009). These
individual TE libraries were integrated into a raw library consisting of 37,589 records, and subsequently
clustered and merged into a non-redundant library. To remove redundant records, pairwise alignment of
TEs from each accession and TE library of rice in repbase (Repbase 28.09) using BLAST were
performed. Records showing alignment lengths over 60% of the element's length and identities over 60%
were considered as redundant records. A total of 23,541 redundant records were eliminated from the
final library. For further filtering, specific accession de novo TEs library and that from Repbase was
used to mask 16 representative accessions by RepeatMasker separately (Tarailo-Graovac *et al*., 2009).
TEs, that did not overlap with records in other libraries, had an accumulated length over 5 Kb and more
than 10 instances in presence, were selected as candidate records. Subsequently, 341 non-redundant TEs
identified in 2 to 15 accessions were clustered based on similarity using CD-HIT (Li & Godzik, 2006).
The final non-redundant library comprised 3,847 sequences, including 3,321 records from the rice TE
library (Repbase 28.09), 341 homologous TEs and 185 accession-specific TEs. All genome assemblies
were masked against the newly-built non-redundant TE library using RepeatMasker. For elements
classified as 'UNKNOWN' by RepeatModeler in more than 50% of cases, a convolutional neural
network-based approach (deepTE) was applied for reclassification (Yan *et al*., 2020). Full-length
elements were defined based on the annotation results, initially merged based on identity and alignment
length to obtain non-redundant or overlapping entries. Specifically, full-length LTR retrotransposons
were characterized by the canonical LTR-Internal-LTR structure, with the intervals between LTR and
internal units being less than 10% of the consensus length, and the overall length deviating by no more

than 10% from the consensus. To ensure accuracy, candidate full-length LTRs meeting these criteria

were extracted and compared against consensus sequences, applying a minimum coverage threshold of

90%. Solo-LTRs were defined as units lacking TE annotations of the same family within a distance of 10%

of the consensus length.

Centromere annotation

The Tandem Repeat Annotation and Structural Hierarchy (TRASH) pipeline (Wlodzimierz *et al*., 2023),

available at https://github.com/vlothec/TRASH, was utilized for the prediction of centromeric satellite

repeats across genomes. The pipeline was configured with parameters "--simpleplot --horclass *CEN155*

--frep 2 --par 5 --k 1000". Genomic sequences were segmented into 1-Kb windows to assess local *k*-mer

counts and identify repetitive regions. Score of each window was determined based on the proportion of

repeated *k*-mers, with regions scoring above a defined threshold considered to contain repeats. Tandem

repeats within these windows were characterized by the distances between identical *k*-mers, aiding in

the identification of consensus sequences such as *CEN155*. To pinpoint centromeric regions within each

rice chromosome, all satellite repeats identified as *CEN155* from the TRASH results were extracted and

mapped according to their chromosomal positions. *CEN155* arrays, indicative of centromeric blocks,

were defined where the spacing between repeat units did not exceed 150 Kb. The largest contiguous

block on each chromosome was considered as the candidate centromeric region. Subsequently, the

CEN155 consensus sequence for each rice accession was aligned back to its respective genome

assembly using BLASTN (v2.5.0+) under default settings. These alignments were integrated into the

centromeric blocks based on their proximity to adjacent *CEN155* arrays. Final validation of centromeric

regions involved manual inspection, confirming regions supported by both TRASH predictions and

BLASTN alignments as definitive centromere locations. Synteny analysis was performed to validate the

predicted centromere boundaries across 69 rice accessions by extracting and examining the 800-kb

flanking regions surrounding the centromere in the NIP reference genome for collinearity with other rice

centromere assemblies. Sequence identity heat maps were plotted for each chromosome using

StainedGlass (Vollger *et al*., 2022), providing a visualization strategy to confirm centromere boundaries.

Centromere similarity and haplotyping

Centromere classification was initiated based on pairwise similarity. Firstly, pairwise alignments were

performed between all centromeres divided into 2-Kb windows using minimap2 (v.2.24) with

parameters (-f 5000 -s 400 -ax ava-ont --dual=yes --eqx) (Li, 2018). Segment windows with identities

over 95% were merged, and the cumulative length was designated as the identical length between

centromeres. The cumulative length between different centromeres was normalized by the length of the

query centromere, highlighting structural variations such as deletions causing asymmetry between the

upper and lower triangles in the heat map (**Fig. 2**). Next, specific haplotypes were assigned to each

centromere. Haplotypes were manually defined based on identity heat maps generated from 2-Kb

windows of different centromeres. Candidate reference haplotypes were established, comprising

sequences from XI and GJ that shared at least 95% of full length with two or more accessions. The

longest regions among candidate centromeres for each haplotype were selected as reference haplotype

sequences. Subsequently, windows split into 2-Kb segments from each centromere were classified into

corresponding haplotypes based on the identity among all haplotypes, when the alignment covered at

least 85% of the window size. In cases where multiple haplotypes exhibit the same identity, the window

was assigned to the haplotype with the fewest mismatches against reference haplotype sequences.

Structural variations and SNPs around centromeres

HiFi reads from each accession were aligned to the reference assembly of NIP using minimap2 (Li, 2018)

with parameters (-ax map-hifi -R "@RG\tID:${pre}\tLB\tSM:${pre}") and SVision (v.1.4)

(--min_mapq 10 --min_sv_size 50) (Lin *et al*., 2022) was utilized to generate structural variation (SV)

calls. SVs within a 500-Kb upstream and downstream region of each centromere were extracted for

statistical analysis. Phylogenetic trees of all accessions were constructed based on SNPs located in the

500-Kb upstream and downstream regions of each centromere with FasttreeMP (Price *et al*., 2009).

Satellite clustering

The *CEN155* repeats identified by TRASH were classified as canonical satellites if their lengths ranged

from 140 bp to 170 bp. Across the 70 rice genomes, a total of 249,952 unique *CEN155* sequences were

detected. Initially, satellites occurring frequently (>30 times) across all accessions were selected.

Additionally, satellites predominantly (>90%) found in specific populations (GJ, XI, AUS, Oruf, Obar,

and Ogla) were included, despite lower frequencies ranged from 5 to 30. A collection of 6,665

representative *CEN155* repeats were finally obtained. These selected satellites were aligned using

MUSCLE (Edgar, 2022) with default parameters, followed by phylogenetic categorization into 15

subfamilies using FastTreeMP (Price *et al*., 2009). To further classify the remaining 243,287 unique

satellites, Levenshtein edit distances were calculated in an all-vs-all comparison against the 6,665

categorized satellites. Each unique satellite was assigned to the superfamily of the assigned *CEN155*

with the closest edit distance.

CENP-B box and pJα motif analysis

To investigate the DNA binding sites of rice centromeres, we introduced a scoring matrix based on the

CENP-B box sequence (5'-PyTTCGTTGGAAPuCGGGA-3') and the pJα motif

(5'-TTCCTTTTPyCACCPuTAG-3'), originally characterized in the human centromeres. Each k-mer (k

= 17) from the query sequences underwent global alignment scoring using a dynamic programming

algorithm against the two binding motif sequences. Additional penalties and bonuses were applied at

conserved loci. All match scores of all monomer in NIP were used to explore the criteria for defining

B-box-like elements. The screening score was set to be top 1% of all scores with a threshold set at 24,

denoting the match of all conserved sites. According to the score distribution, the k-mers with scores

above 25/27 were considered as candidate B-box/pJα-like elements. To enable rapid scanning of

genome-wide elements across multiple genomes, the k-mers harboring B-box-like elements from NIP

genome were integrated using MEME v5.4.1 with parameters '-w 17 -dna -revcomp' to construct a

B-box-like motif library (Bailey *et al*., 2015). Subsequently, B-box-like elements were identified across

the whole genome with e-value < 1e-05 against the motif library for B-box and pJα motif separately,

utilizing FIMO in MEME with the parameters '--max-strand'.

Structural organization analysis of *CEN155* arrays

We resolved the structure of satellite arrays in a progressive compression strategy, merging identical

units iteratively (**Fig. 4a**). Homogenization regions of monomer (moHRs) were identified where more

than five consecutive satellites from the same family were present. TEs embedded in the satellite arrays

were masked. Fine analysis of satellite organization within each moHR was performed using identity

dot plots and NTRprism. Pairs of identical satellites within moHRs were selected to investigate

internally local collinearity using DAGchainer (Haas *et al*., 2004). NTRprism (Altemose *et al*., 2022)

was employed to infer the most abundant tandem repeat periodicities by counting the interval lengths

between adjacent instances of each 6-mer. For a canonical HOR array, a distinct peak in the interval

spectra was expected at the HOR unit length (Altemose *et al*., 2022). Using the moHR-compressed

satellite array, we defined homogenization regions of dimer (diHRs) containing at least five replicates of

dimers. For determining homogenization regions of multimer (muHRs), a de Bruijn graph was initially

constructed based on the dimer-compressed satellite string of each centromere, where the vertex set

comprises all monomer superfamilies and the edge set consists of all pairs of its consecutive monomers

superfamily. For instance, for a dimer-compressed centromere sequence "ABCDABCD", the edge set

would include the pairs A-B{2}, B-C{2}, C-D{2}, D-A{1}. Candidate multimers were identified by

exhaustively enumerating all cycles in the graph using a depth-first search (DFS) strategy. We observed

that ancient multimers might not exhibit a unified head-to-tail pattern like in the human centromeres,

due to local expansion, random insertion or deletion. Candidate multimer calls were filtered out based

on criteria of multiplicity less than 2, traversal less than 3 and adjacent multimer spacing greater than

three times the multimer length. The multiplicity of an edge (A, B) in the graph was defined as the

number of times the monomers family A followed the monomer family B in each dimer-compressed

centromere. Traversal represented the occurrence frequency of a cycle (multimer) in each

dimer-compressed centromere string. To identify multimer hierarchically, candidate nested-multimers

were mapped onto the discrete dimer-compressed centromere and detect overlapping regions with

continuity and occurrence. If the merged multimer presented better continuity and higher occurrence, it

would replace the nested one.

Centrophilic LTR analysis

To identify centrophilic LTRs, which preferentially localize within centromeres, we conducted a

statistical analysis based on the annotation of full-length LTRs. LTRs that constituted more than 90% of

all full-length LTRs in centromeres and exhibited a significantly higher preference for the centromere

region compared to the chromosome arms were designated as centrophilic LTRs. To characterize the

core domains of these LTRs, ORFs within the internal domain of consensus elements from the pan-TE

database were extracted using getorf in EMBOSS (Rice *et al*., 2000). Subsequently, domain predictions

were performed using HMMER (v3.4) (-E 1e-5) against Pfam hidden Markov models, including

PF00026, PF13650, PF08284, PF13975, PF00077, PF09668 for protease; PF03732 for GAG; PF00665,

PF13683, PF17921, PF02022, PF09337, PF00552 for integrase; PF17917, PF17919, PF13456 for

RNase H; and PF00078 for reverse transcriptase (Mistry *et al*., 2021). Given the presence of

non-autonomous LTRs in the pan-TE library, the phylogenetic relationships of LTRs were resolved

separately based on the 5' LTR, 3' LTR, and internal sequences. Multiple sequence alignments were

conducted using MAFFT (--anysymbol −maxiterate 1000), and conserved sites were extracted using

trimal (Katoh *et al*., 2013; Capella-Gutiérrez *et al*., 2009). Phylogenetic trees were constructed using

FastTreeMP (-nt -gtr) and visualized with iTOL (Price *et al*., 2009; Letunic & Bork, 2021). Pairwise

alignments among full-length centrophilic LTRs were performed and visualized using VISTA (Mayor *et*

al., 2000). Identity between 5' LTR and 3' LTR of centrophilic LTRs was calculated based on full-length

LTRs existing within centromeres. Alignments for each LTR pair were conducted using MAFFT, and

identities were computed using ape (Paradis & Schliep, 2019). To map the insertion sites of SZ-22

elements, 150-bp regions flanking each element were extracted and queried against the *CEN155*

database using BLASTn. Centromeric elements were filtered based on close alignments to the same

monomer (2,258 out of 3,673 sequences). The midpoint between aligned flanking sequences determined

the insertion positions, and their distribution along the *CEN155* consensus sequence was visualized.

Pairwise comparison of rice centromeres

We introduced a new algorithm termed SynPan-CEN, to compared satellite arrays in single monomer

scale. This approach leverages satellite clustering information and pairwise genetic distances to identify

syntenic pairs of *CEN155* satellites between rice centromeres. Initially, we conducted an exhaustive

all-to-all comparison of monomer sequences from two satellite arrays using pairwise edit distance (ED)

to quantify similarity. We refined synteny construction by conservatively selecting monomer pairs in a

correct order, incrementally increasing the ED threshold until the framework encompassed more than 20%

of total monomers or 100 monomers. Using DAGchainer software (Haas *et al*., 2004), which computes

chains of syntenic genes based on sequence homology and genomic coordinates, we identified syntenic

satellite regions and syntelogs. DAGchainer employs a scoring function considering the distance

between neighboring elements on each DNA molecule and BLAST E-value scores between homologs,

facilitating the identification of maximally scoring chains of ordered element pairs. For the analysis here,

DAGchainer parameters (-g 1 -D 20 -A 6 -Z 50 -e -3f) were set to retain blocks with the highest scores,

with non-overlapping monomer pairs added to complement the chain backbone. Using a window of ten

syntenic pairs, we applied sliding windows along the fixed chain backbone to extract each region. Local

syntenic pairs were identified using DAGchainer, maintaining consistency with the parameters

mentioned above. Blocks with the highest scores were preserved, and additional non-overlapping

syntenic monomer pairs were incorporated. To refine our results, we incrementally tested various ED

thresholds and observed a significant impact on identifying optimal syntenic pairs of *CEN155* satellites.

Syntenic satellite pairs increased as ED values were below 15, but decreased beyond this threshold. This

phenomenon is likely due to the rapid accumulation of potential syntenic paths after exceeding the ED

threshold, which complicates the determination of the optimal paths by DAGchainer (Haas *et al*., 2004).

In cases where sliding windows presented excessive alignment possibilities, we subdivided windows

into two sub-regions, each containing five syntenic pairs, mitigating potential mismatches.

1000

Sequence divergence and mutation rate in centromeres

Edit distances between syntenic *CEN155* satellite pairs inferred by SynPan-CEN were calculated.

Continuous gaps at the beginning or end of a monomer alignment were systematically removed to

minimize potential mismatches. Within alignments, consecutive gaps were treated as a single transition,

and insertion counts were used exclusively to quantify edit distances. To optimize the alignment process,

we compared three widely-used programs—MUSCLE (Edgar, 2022), MAFFT (Rozewicki *et al*., 2019),

and ClustalW (Thompson *et al*., 2002)—using default parameters for pairwise monomer sequence

alignments. Based on alignment performance and computational efficiency, MAFFT and ClustalW

consistently demonstrated superior reliability compared to MUSCLE. Consequently, MAFFT was

selected for following analyses. A phylogenetic tree was constructed using all syntenic satellites

between chromosomes, where the synteny information was visualized with links in the circle tree plot.

Orthologous pairs were identified based on their proximity to each other in the tree structure. The

satellite pair, where the two *CEN155* were clustered as a monoclade, were confirmed as orthologs, while

pairs lacking such proximity were more likely to be paralogs.

1015

DNA methylation analysis using PacBio HiFi data

For the 46 samples whose HiFi reads were newly generated in this study, 5mC methylation probability

was determined by combining kinetics from multiple passes in HiFi sequencing. HiFi reads with

methylation MM and ML tags were aligned against the corresponding assembly using pbmm2 (v1.2.0)

(https://github.com/PacificBiosciences/pbmm2). We employed pb-CpG-tools

(https://github.com/PacificBiosciences/pb-CpG-tools) to call base-level methylation status in the model

'pileup_calling_model.v1'. Given that HiFi-based 5mC signals were predominantly enriched in the CG

islands, we focused exclusively on the methylation signals at CG sites along chromosomes, therefore

excluding methylation information for CHH and CHG contexts.

1025

DNA methylation analysis using ONT data

ONT sequencing data was utilized for methylation analysis. Methylation probabilities in CG, CHH, and

CHG contexts were called using the ONT reads basecaller Dorado (v.0.5.3) from the raw pod5 files,

with the basecalling model options: '--modified-bases-models dna_r10.4.1_e8.2_400bps_hac

@v4.3.0_5mC_5hmC@v1 --emit-moves'. The base-called read sequences were aligned against their

corresponding genome assemblies. Modkit (v.1.0, https://github.com/nanoporetech/modkit) was

utilized to generate summary counts of modified and unmodified bases in bedMethyl files. DNA

methylation levels were quantified using 'modkit pileup' based on only primary alignments. The

consistency of methylation quantification between PacBio HiFi reads and ONT reads was assessed by

randomly selecting 10,000 CG sites, using Pearson correlation as the evaluation metric.

1036

ChIP-seq data alignment and processing

CENH3 ChIP-seq reads from 13 accessions (ten newly generated in this study and three previously

released) were used to determine the functional centromere regions. Paired-end raw reads were

subjected to adapter trimming and filtering using fastp (v.0.23.4) (Chen *et al*., 2023), with the following

options: '--cut_front_window_size 4 --cut_front_mean_quality 20 -3 --cut_tail_window_size 4

--cut_tail_mean_quality 20 --cut_right --cut_right_window_size 4 --cut_right_mean_quality 20'.

Filtered read sequences were aligned to their corresponding genome assembly using Bowtie2 (v.2.3.5.1)

(Langmead *et al*., 2019) with parameters '--very-sensitive --no-mixed --no-discordant --maxins 800'.

Reads that aligned to multiple locations were filtered using samtools (v.1.7) (Li *et al*., 2009). The

enrichment level of CENH3 for each 1-Kb window was calculated using bamCompare from the

Deeptools package (v.3.5.4.post1) (Ramírez *et al*., 2016) with the parameters '--binSize 1000

--outFileFormat bedgraph --operation log2 -p 5 --extendReads'. Base-level CENH3 enrichment was

calculated using bamCompare with the parameters '--binSize 1 --numberOfProcessors 40 --operation

ratio --outFileFormat bedgraph'. Custom R scripts were used to visualize CENH3 occupancy profiles

across the 12 chromosomes and *CEN155* satellite repeat arrays.

1052

1053

Data availability

Raw PacBio HiFi reads for 46 rice accessions, raw ONT sequencing reads for ten accessions, and

CENH3 ChIP-seq NGS reads for ten accessions generated in this study have been deposited in NGDC

(https://ngdc.cncb.ac.cn/) under the BioProject accession PRJCA025388, with the GSA numbers of

CRA016014, CRA016014 and CRA017653, respectively. The newly-generated genome assemblies in

this study are available at Zenodo (https://doi.org/10.5281/zenodo.12770803). The genome assemblies

of NIP, MH63 and ZS97 are available at the RiceSuperPIRdb (http://www.ricesuperpir.com/) and RIGW

(http://rice.hzau.edu.cn/rice_rs3/). The TE and gene annotation files of 70 rice genomes are deposited at

Zenodo (https://doi.org/10.5281/zenodo.12698984). Rice centromere annotation and comparison plots

for all accessions and chromosomes are supplemented at Zenodo

(https://doi.org/10.5281/zenodo.12702715), including similarity heat-map plots generated by

StainedGlass for each centromere, whole-genome synteny to the NIP reference assembly, centromere

synteny against NIP and NJ11 assemblies and centromere composition for all chromosomes.

1067

1068

Code availability

The SynPan-CEN code is available at https://github.com/Darlene1997/SynPan-CEN and the scripts for

the progressive compression strategy in deciphering the satellite organization and additional in-house

codes associated with this study (including assembly, annotation and visualization) are available at

https://github.com/dongyawu/CenTools. The similarity heat-map within each centromere was drawn by

StainedGlass (https://github.com/mrvollger/StainedGlass). The visualization of centromere annotation

and synteny tracks was performed using ggplot2 (https://github.com/tidyverse/ggplot2) in R (v4.3.1,

https://www.r-project.org/).

1077

1078

Reference

Ahmed, H.I., Heuberger, M., Schoen, A., Koo, D.H., Quiroz-Chavez, J., Adhikari, L., Raupp, J.,
Cauet, S., Rodde, N., and Cravero, C., et al. (2023). Einkorn genomics sheds light on history of the
oldest domesticated wheat. Nature **620**:830-838. https://doi.org/10.1038/s41586-023-06389-7.
Alonge, M., Lebeigle, L., Kirsche, M., Jenike, K., Ou, S., Aganezov, S., Wang, X., Lippman, Z.B.,
Schatz, M.C., and Soyk, S. (2022). Automated assembly scaffolding using ragtag elevates a new
tomato system for high-throughput genome editing. Genome Biol **23**:258.
https://doi.org/10.1186/s13059-022-02823-7.
Altemose, N., Logsdon, G.A., Bzikadze, A.V., Sidhwani, P., Langley, S.A., Caldas, G.V., Hoyt, S.J.,
Uralsky, L., Ryabov, F.D., and Shew, C.J., et al. (2022). Complete genomic and epigenetic maps
of human centromeres. Science **376**:eabl4178. https://doi.org/10.1126/science.abl4178.
Bailey, T.L., Johnson, J., Grant, C.E., and Noble, W.S. (2015). The MEME Suite. Nucleic Acids Res
43:W39-W49. https://doi.org/10.1093/nar/gkv416.
Barra, V., and Fachinetti, D. (2018). The dark side of centromeres: types, causes and consequences of
structural abnormalities implicating centromeric DNA. Nat Commun **9**:4340.
https://doi.org/10.1038/s41467-018-06545-y.
Bensasson, D. (2011). Evidence for a high mutation rate at rapidly evolving yeast centromeres. BMC
Evol Biol **11**:211. https://doi.org/10.1186/1471-2148-11-211.
Bzikadze, A.V., Mikheenko, A., and Pevzner, P.A. (2022). Fast and accurate mapping of long reads to
complete genome assemblies with VerityMap. Genome Res **32**:2107-2118.
https://doi.org/10.1101/gr.276871.122.
Cappelletti, E., Piras, F.M., Sola, L., Santagostino, M., Petersen, J.L., Bellone, R.R., Finno, C.J.,
Peng, S., Kalbfleisch, T.S., and Bailey, E., et al. (2023). The localization of centromere protein a is
conserved among tissues. Commun Biol **6**:963. https://doi.org/10.1038/s42003-023-05335-7.
Chen, J., Wang, Z., Tan, K., Huang, W., Shi, J., Li, T., Hu, J., Wang, K., Wang, C., and Xin, B., et
al. (2023). A complete telomere-to-telomere assembly of the maize genome. Nat Genet
55:1221-1231. https://doi.org/10.1038/s41588-023-01419-6.
Chen, Q., Yang, C., Zhang, G., and Wu, D. (2024). GCI: a continuity inspector for complete genome
assembly. bioRxiv. https://doi.org/10.1101/2024.04.06.588431.

Chen, S. (2023). Ultrafast one-pass fastq data preprocessing, quality control, and deduplication using
fastp. iMeta **2**:e107. https://doi.org/10.1002/imt2.107.
Cheng, H., Asri, M., Lucas, J., Koren, S., and Li, H. (2024). Scalable telomere-to-telomere assembly
for diploid and polyploid genomes with double graph. Nat Methods **21**:967-970.
https://doi.org/10.1038/s41592-024-02269-8.
Cheng, H., Jarvis, E.D., Fedrigo, O., Koepfli, K.P., Urban, L., Gemmell, N.J., and Li, H. (2022).
Haplotype-resolved assembly of diploid genomes without parental data. Nat Biotechnol
40:1332-1335. https://doi.org/10.1038/s41587-022-01261-x.
Cheng, Z., Buell, C.R., Wing, R.A., Gu, M., and Jiang, J.M. Toward a cytological characterization of
the rice genome. Genome Res **11**: 2133-2141. https://doi.org/10.1101/gr.194601.
Cheng, Z., Dong, F., Langdon, T., Ouyang, S., Buell, C.R., Gu, M., Blattber F.R., and Jiang, J.M.
(2002). Functional rice centromeres are marked by a satellite repeat and a centromere-specific
retrotransposon. Plant Cell **14**: 1691-1704. https://doi.org/10.1105/tpc.003079.
Edgar, R.C. (2022). Muscle5: high-accuracy alignment ensembles enable unbiased assessments of
sequence homology and phylogeny. Nat Commun **13**:6968.
https://doi.org/10.1038/s41467-022-34630-w.
Gong, H., and Han, B. (2022). Genetic introgression between different groups reveals the differential
process of Asian cultivated rice. Sci Rep **12**:17662. https://doi.org/10.1038/s41598-022-22674-3.
Haas, B.J., Delcher, A.L., Wortman, J.R., and Salzberg, S.L. (2004). DAGchainer: a tool for mining
segmental genome duplications and synteny. Bioinformatics **20**:3643-3646.
https://doi.org/10.1093/bioinformatics/bth397.
Hu, J., Wang, Z., Liang, F., Liu, S.L., Ye, K., and Wang, D.P. (2024). NextPolish2: a repeat-aware
polishing tool for genomes assembled using HiFi long reads. Genom Proteom Bioinf **22**: qzad009.
https://doi.org/10.1093/gpbjnl/qzad009.
Huang, Z., Xu, Z., Bai, H., Huang, Y., Kang, N., Ding, X., Liu, J., Luo, H., Yang, C., and Chen, W.,

et al. (2023). Evolutionary analysis of a complete chicken genome. Proc Natl Acad Sci U S A
120:e2078326176. https://doi.org/10.1073/pnas.2216641120.

Jain, C., Rhie, A., Hansen, N.F., Koren, S., and Phillippy, A.M. (2022). Long-read mapping to
repetitive reference sequences using Winnowmap2. Nat Methods **19**:705-710.
https://doi.org/10.1038/s41592-022-01457-8.

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019). MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief Bioinform **20**:1160-1166.
https://doi.org/10.1093/bib/bbx108.

Katoh, K., and Standley, D.M. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol **30**:772-780.
https://doi.org/10.1093/molbev/mst010.

Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S.L. (2013). TopHat2:
accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions.
Genome Biol **14**:R36. https://doi.org/10.1186/gb-2013-14-4-r36.

Kipling, D., Wilson, H.E., Mitchell, A.R., Taylor, B.A., and Cooke, H.J. (1994). Mouse centromere
mapping using oligonucleotide probes that detect variants of the minor satellite. Chromosoma
103:46-55. https://doi.org/10.1007/BF00364725.

Kugou, K., Hirai, H., Masumoto, H., Koga, A., Gong, J., Liu, J., and Yong, K. (2016). Formation of
functional CENP-B boxes at diverse locations in repeat units of centromeric DNA in new world
monkeys. Sci Rep **6**:27833. https://doi.org/10.1038/srep27833.

Kursel, L.E., and Malik, H.S. (2016). Centromeres. Curr Biol **26**:R481-R492.
https://doi.org/10.1016/j.cub.2016.05.031.

Langmead, B., Wilks, C., Antonescu, V., and Charles, R. (2019). Scaling read aligners to hundreds
of threads on general-purpose processors. Bioinformatics **35**:421-432.
https://doi.org/10.1093/bioinformatics/bty648.

Letunic, I., and Bork, P. (2021). Interactive tree of life (iTol) v5: an online tool for phylogenetic tree
display and annotation. Nucleic Acids Res **49**:W293-W296. https://doi.org/10.1093/nar/gkab301.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics **34**:3094-3100.
https://doi.org/10.1093/bioinformatics/bty191.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and
Durbin, R. (2009). The sequence alignment/map format and SAMtools. Bioinformatics
25:2078-2079. https://doi.org/10.1093/bioinformatics/btp352.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics **25**:1754-1760. https://doi.org/10.1093/bioinformatics/btp324.

Li, K., Xu, P., Wang, J., Yi, X., and Jiao, Y. (2023). Identification of errors in draft genome
assemblies at single-nucleotide resolution for quality assessment and improvement. Nat Commun
14:6556. https://doi.org/10.1038/s41467-023-42336-w.

Li, W., and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of protein
or nucleotide sequences. Bioinformatics. https://doi.org/10.1093/bioinformatics/btl158.

Lian, Q., Huettel, B., Walkemeier, B., Mayjonade, B., Lopez-Roques, C., Gil, L., Roux, F.,
Schneeberger, K., and Mercier, R. (2024). A pan-genome of 69 *Arabidopsis thaliana* accessions
reveals a conserved genome structure throughout the global species range. Nat Genet **56**:982-991.
https://doi.org/10.1038/s41588-024-01715-9.

Lin, J., Wang, S., Audano, P.A., Meng, D., Flores, J.I., Kosters, W., Yang, X., Jia, P., Marschall,
T., and Beck, C.R., et al. (2022). SVision: a deep learning approach to resolve complex structural
variants. Nat Methods **19**:1230-1233. https://doi.org/10.1038/s41592-022-01609-w.

Liu, Y., Yi, C., Fan, C., Liu, Q., Liu, S., Shen, L., Zhang, K., Huang, Y., Liu, C., and Wang, Y., et
al. (2023). Pan-centromere reveals widespread centromere repositioning of soybean genomes. Proc
Natl Acad Sci U S A **120**:e1984790176. https://doi.org/10.1073/pnas.2310177120.

Logsdon, G.A., Rozanski, A.N., Ryabov, F., Potapova, T., Shepelev, V.A., Catacchio, C.R.,
Porubsky, D., Mao, Y., Yoo, D., and Rautiainen, M., et al. (2024). The variation and evolution of
complete human centromeres. Nature **629**:136-145. https://doi.org/10.1038/s41586-024-07278-3.

Lv, Y., Liu, C., Li, X., Wang, Y., He, H., He, W., Chen, W., Yang, L., Dai, X., and Cao, X., et al.
(2024). A centromere map based on super pan genome highlights the structure and function of rice
centromeres. J Integr Plant Biol **66**:196-207. https://doi.org/10.1111/jipb.13607.

Malik, H.S., and Henikoff, S. (2009). Major evolutionary transitions in centromere complexity. Cell
138:1067-1082. https://doi.org/10.1016/j.cell.2009.08.036.
Mayor, C., Brudno, M., Schwartz, J.R., Poliakov, A., Rubin, E.M., Frazer, K.A., Pachter, L.S.,
and Dubchak, I. (2000). VISTA: visualizing global DNA sequence alignments of arbitrary length.
Bioinformatics **16**:1046-1047. https://doi.org/10.1093/bioinformatics/16.11.1046.
Mckenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,
Altshuler, D., Gabriel, S., and Daly, M., et al. (2010). The genome analysis toolkit: a mapreduce
framework for analyzing next-generation DNA sequencing data. Genome Res **20**:1297-1303.
https://doi.org/10.1101/gr.107524.110.
Mikheenko, A., Bzikadze, A.V., Gurevich, A., Miga, K.H., and Pevzner, P.A. (2020). TandemTools:
mapping long reads and assessing/improving assembly quality in extra-long tandem repeats.
Bioinformatics **36**:i75-i83. https://doi.org/10.1093/bioinformatics/btaa440.
Minton, K. (2024). Tandem repeat variation of human centromeres. Nat Rev Genet **25**:455.
https://doi.org/10.1038/s41576-024-00741-x.
Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G.A., Sonnhammer, E., Tosatto,
S., Paladin, L., Raj, S., and Richardson, L.J., et al. (2021). Pfam: the protein families database in
2021. Nucleic Acids Res **49**:D412-D419. https://doi.org/10.1093/nar/gkaa913.
Nagaki, K., Talbert, P.B., Zhong, G.X., Dawe, R.K., Henikoff, S., and Jiang, J. (2003). Chromatin
immunoprecipitation reveals that the 180-bp satellite repeat is the key functional DNA element of
Arabidopsis thaliana centromeres. Genetics **163**:1221-1225.
Naish, M., Alonge, M., Wlodzimierz, P., Tock, A.J., Abramson, B.W., Schmücker, A., Mandáková,
T., Jamge, B., Lambing, C., and Kuo, P., et al. (2021). The genetic and epigenetic landscape of the
Arabidopsis centromeres. Science **374**. https://doi.org/10.1126/science.abi7489.
Naish, M., and Henderson, I.R. (2024). The structure, function, and evolution of plant centromeres.
Genome Res **34**:161-178. https://doi.org/10.1101/gr.278409.123.
Paradis, E., and Schliep, K. (2019). Ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics **35**:526-528.
https://doi.org/10.1093/bioinformatics/bty633.
Price, M.N., Dehal, P.S., and Arkin, A.P. (2009). FastTree: computing large minimum evolution trees
with profiles instead of a distance matrix. Mol Biol Evol **26**:1641-1650.
https://doi.org/10.1093/molbev/msp077.
Qin, P., Lu, H., Du H, Wang, H., Chen, W., Chen, Z., He, Q., Ou, S., Zhang, H., and Li, X., et al.
(2021). Pan-genome analysis of 33 genetically diverse rice accessions reveals hidden genomic
variations. Cell **184**:3542-3558. https://doi.org/10.1016/j.cell.2021.04.046.
Qiu, J., Jia, L., Wu, D., Weng, X., Chen, L., Sun, J., Chen, M., Mao, L., Jiang, B., and Ye, C., et al.
(2020). Diverse genetic mechanisms underlie worldwide convergent rice feralization. Genome Biol
21:70. https://doi.org/10.1186/s13059-020-01980-x.
Ramirez, F., Ryan, D.P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dundar,
F., and Manke, T. (2016). DeepTools2: a next generation web server for deep-sequencing data
analysis. Nucleic Acids Res **44**:W160-W165. https://doi.org/10.1093/nar/gkw257.
Rautiainen, M., Nurk, S., Walenz, B.P., Logsdon, G.A., Porubsky, D., Rhie, A., Eichler, E.E.,
Phillippy, A.M., and Koren, S. (2023). Telomere-to-telomere assembly of diploid chromosomes
with Verkko. Nat Biotechnol **41**:1474-1482. https://doi.org/10.1038/s41587-023-01662-6.
Rhie, A., Walenz, B.P., Koren, S., and Phillippy, A.M. (2020). Merqury: reference-free quality,
completeness, and phasing assessment for genome assemblies. Genome Biol **21**:245.
https://doi.org/10.1186/s13059-020-02134-9.
Rice, P., Longden, L., and Bleasby, A. (2000). EMBOSS: the European molecular biology open
software suite. The European Molecular Biology Open Software Suite.
Rice, W.R. (2019). A game of thrones at human centromeres ii. A new molecular/evolutionary model.
bioRvix. https://doi.org/10.1101/731471.
Rice, W.R. (2020). A game of thrones at human centromeres i. Multifarious structure necessitates a new
molecular/evolutionary model. bioRxiv. https://doi.org/10.1101/731430.
Sedeek, K., Zuccolo, A., Fornasiero, A., Weber, A.M., Sanikommu, K., Sampathkumar, S.,
Rivera, L.F., Butt, H., Mussurova, S., and Alhabsi, A., et al. (2023). Multi-omics resources for
targeted agronomic improvement of pigmented rice. Nat Food **4**:366-371.

https://doi.org/10.1038/s43016-023-00742-9.

Shang, L., He, W., Wang, T., Yang, Y., Xu, Q., Zhao, X., Yang, L., Zhang, H., Li, X., and Lv, Y., et
al. (2023). A complete assembly of the rice Nipponbare reference genome. Mol Plant **16**:1232-1236.
https://doi.org/10.1016/j.molp.2023.08.003.

Shang, L., Li, X., He, H., Yuan, Q., Song, Y., Wei, Z., Lin, H., Hu, M., Zhao, F., and Zhang, C., et
al. (2022). A super pan-genomic landscape of rice. Cell Res **32**:878-896.
https://doi.org/10.1038/s41422-022-00685-z.

Sim, S.B., Corpuz, R.L., Simmonds, T.J., and Geib, S.M. (2022). HiFiAdapterFilt, a memory
efficient read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and
their negative impacts on genome assembly. BMC Genom **23**:157.
https://doi.org/10.1186/s12864-022-08375-1.

Song, J., Xie, W., Wang, S., Guo, Y., Koo, D., Kudrna, D., Gong, C., Huang, Y., Feng, J., and
Zhang, W., et al. (2021). Two gap-free reference genomes and a global view of the centromere
architecture in rice. Mol Plant **14**:1757-1767. https://doi.org/10.1016/j.molp.2021.06.018.

Stein, J.C., Yu, Y., Copetti, D., Zwickl, D.J., Zhang, L., Zhang, C., Chougule, K., Gao, D., Iwata,
A., and Goicoechea, J.L., et al. (2018). Genomes of 13 domesticated and wild rice relatives
highlight genetic conservation, turnover and innovation across the genus *Oryza*. Nat Genet **50**:1618.
https://doi.org/10.1038/s41588-018-0261-2.

Talbert, P., and Henikoff, S. (2022). Centromeres organize (epi)genome architecture. Cell
185:3083-3085. https://doi.org/10.1126/science.1069609.

Tarailo Graovac, M., and Chen, N. (2009). Using RepeatMasker to identify repetitive elements in
genomic sequences. Curr Protoc Bioinformatics **25**. https://doi.org/10.1002/0471250953.bi0410s25.

Thompson, J.D., Gibson, T.J., and Higgins, D.G. (2002). Multiple sequence alignment using
ClustalW and ClustalX. Curr Protoc Bioinformatics **Chapter 2**:2-3.
https://doi.org/10.1002/0471250953.bi0203s00.

Trapnell, C., Williams, B.A., Pertea, G., Mortazavi, A., Kwan, G., van Baren, M.J., Salzberg, S.L.,
Wold, B.J., and Pachter, L. (2010). Transcript assembly and quantification by RNA-seq reveals
unannotated transcripts and isoform switching during cell differentiation. Nat Biotechnol
28:511-515. https://doi.org/10.1038/nbt.1621.

Vollger, M.R., Kerpedjiev, P., Phillippy, A.M., and Eichler, E.E. (2022). StainedGlass: interactive
visualization of massive tandem repeat structures with identity heatmaps. Bioinformatics
38:2049-2051. https://doi.org/10.1093/bioinformatics/btac018.

Wang, T., Wang, B., Hua, X., Tang, H., Zhang, Z., Gao, R., Qi, Y., Zhang, Q., Wang, G., and Yu,
Z., et al. (2023). A complete gap-free diploid genome in *Saccharum* complex and the genomic
footprints of evolution in the highly polyploid *Saccharum* genus. Nat Plants **9**:554-571.
https://doi.org/10.1038/s41477-023-01378-0.

Wei, X., Chen, M., Zhang, Q., Gong, J., Liu, J., Yong, K., Wang, Q., Fan, J., Chen, S., and Hua, H.,
et al. (2024). Genomic investigation of 18,421 lines reveals the genetic architecture of rice. Science
385:eadm8762. https://doi.org/10.1126/science.adm8762.

Wlodzimierz, P., Rabanal, F.A., Burns, R., Naish, M., Primetis, E., Scott, A., Mandáková, T.,
Gorringe, N., Tock, A.J., and Holland, D., et al. (2023). Cycles of satellite and transposon
evolution in *Arabidopsis* centromeres. Nature **618**:557-565.
https://doi.org/10.1038/s41586-023-06062-z.

Wu, D., Xie, L., Sun, Y., Huang, Y., Jia, L., Dong, C., Shen, E., Ye, C.Y., Qian, Q., and Fan, L.
(2023). A syntelog-based pan-genome provides insights into rice domestication and
de-domestication. Genome Biol **24**:179. https://doi.org/10.1186/s13059-023-03017-5.

Wu, T.D., and Watanabe, C.K. (2005). GMAP: a genomic mapping and alignment program for
mRNA and EST sequences. Bioinformatics **21**:1859-1875.
https://doi.org/10.1093/bioinformatics/bti310.

Wu, Z., Fang, D., Yang, R., Gao, F., An, X., Zhou, X., Li, Y., Yi, C., Zhang, T., and Liang, C., et al.
(2018) De novo genome assembly of *Oryza granulata* reveals rapid genome expansion and adaptive
evolution. Commun Biol **1**:84. https://doi.org/10.1038/s42003-018-0089-4.

Xie, L., Gong, X., Yang, K., Huang, Y., Zhang, S., Shen, L., Sun, Y., Wu, D., Ye, C., and Zhu, Q.H.,
et al. (2024). Technology-enabled great leap in deciphering plant genomes. Nat Plants **10**:551-566.
https://doi.org/10.1038/s41477-024-01655-6.

Yan, H., Bombarely, A., and Li, S. (2020). DeepTE: a computational method for *de novo* classification
of transposons with convolutional neural network. Bioinformatics **36**:4269-4275.
https://doi.org/10.1093/bioinformatics/btaa519.
Zhang, Y., Fu, J., Wang, K., Han, X., Yan, T., Su, Y., Li, Y., Lin, Z., Qin, P., and Fu, C., et al.
(2022). The telomere-to-telomere gap-free genome of four rice parents reveals SV and PAV patterns in
hybrid rice breeding. Plant Biotechnol J **20**:1642-1644. https://doi.org/10.1111/pbi.13880.

1304

1305

Acknowledgements

This work was supported by China National Postdoctoral Program for Innovative Talents

(BX20220269), China Postdoctoral Science Foundation (No. 2023M743045), Young Scientists Fund of

the National Natural Science Foundation of China (No. 32300490) to D.W.; National Key Research and

Development Program of China (No. 2019YFA0903904) to W.H.; Biological Breeding-Major Projects

(No. 2023ZD04076), Hainan Province Science and Technology Special Fund (No.

ZDYF2022XDNY271) and CIC-MIC to L.F. We thanks to Guojie Zhang (Zhejiang University), Yafei

Mao (Shanghai Jiao Tong University) and Kun Wu (Zhejiang University) for constructive suggestions.

1314

1315

Author contribution

D.W. conceived and initiated this study. L.S. collected the samples. D.W., Q.C. and M.S. performed the

sequencing data quality control, centromere assembly and quality evaluation. L.X., D.W. and Y.S.

performed the analysis of satellite sequence identification and clustering and satellite array organization.

Y.H. and D.W. performed the annotation and centromeric insertion analysis of TEs. W.H., L.X. and S.B.

conducted the CENH3 chromatin immunoprecipitation (ChIP) experiments. L.X., D.W. and S.Z.

processed the ChIP-seq data and analyzed the epigenetic profiling of rice centromeres. L.F. and D.W.

supervised all analyses. Q.Q., W.J., C.Y., L.S. and X.Z. provided suggestions on analysis, organization

and writing. D.W., L.X. and Y.H. wrote the manuscripts with the inputs from all co-authors. All authors

discussed the results and commented on the manuscript.

1326

1327

Competing interests

1329 All authors declare no competing interests.

1330

Figure legends

Figure 1 | Centromere diversity of *Oryza* AA genomes.
a, Geographic origin of the 70 rice accessions in the *Oryza* AA group, coloured by taxonomic group membership. A phylogenetic tree constructed using whole-genome SNPs is embedded in the map, with the branches and labels colored according to taxonomic information. Samples selected for performing CENH3 ChIP-seq are highlighted with bold labels on a gray background. Divergence times were adapted from Stein *et al*., 2018. **b**, Variation in centromere length across the 70 accessions per chromosome.

Figure 2 | Centromere introgression and splitting.
a,Centromere sequence similarity, structural variations in centromere flanking regions (compared to NIP), and siding-window haplotyping of Chr12 centromeres. Haplotypes are shown in different colors. **b**, StainedGlass sequence similarity heat maps of Chr12 centromeres from SL044, CW09 and NJ11, and their synteny. TEs (blue) and gene-like elements (red) are shown. Their commonly shared retrotransposon *RETROSAT-2C* is highlighted. **c**, Phylogeny of Asian rice accessions based on the upstream and downstream SNPs of the Chr12 centromere. Taxonomic information is represented by colored circles, with the position of SL044 marked by a dashed line. **d**, Schematic of centromere rapid alterations in structure, including introgression, duplication and fission or splitting.

Figure 3 | Variation of *CEN155* sequences and centrophilic TEs in centromere regions.
a, Pie charts illustrating the *CEN155* proportion on the forward (red, +) or reverse (blue, -) strand across chromosomes (rows) and accessions (columns). **b**, A maximum-likelihood phylogenetic tree of representative *CEN155* sequences ($n = 6,665$). Fifteen superfamilies are clustered and colored. **c**, Consensus sequences of the fifteen *CEN155* superfamilies. B-box and pJα-like motifs are highlighted. **d**. Alignment of B-box motifs across human, marmoset, mouse, horse and rice. **e**, Enrichment analysis of B-box in different genomic regions of the rice genome. Significance values were calculated using Student's *t* test. **f**, Centromere compositions across the 70 rice genomes. Branch colors in the phylogeny tree (left) indicate taxonomic groups. **g**, Abundance enrichment of centrophilic retrotransposons (*SZ-22*, *CRM*, *RIRE7*) in centromere, peri-centromere and chromosome arm regions. Each chromosome is shown in 100 bins, where the window sizes are adjusted depending on the length of each region. **h**, Phylogenetic relationship and sequence similarity among centrophilic LTRs. Bootstrap values of 1000 replicates are shown near the branches. **i**, Insertion ages, measured by sequence identity between the two flanking LTR sequences within each intact *SZ-22*, *CRM* and *RIRE7*, comparing inside and outside the *CEN155* arrays.

Figure 4 | Multi-layer nested structure of rice centromeres
a, Schematic illustrating a progressive compression strategy to decompose the structure of *CEN155* arrays. Identical adjacent monomers, dimers and multimers are compressed stepwise, defining local homogenization blocks as moHRs, diHRs and muHRs, respectively. **b**, Distribution of replicate times for monomers, dimers and multimers. **c**, Structural composition of moHR (left), diHR (middle) and muHR (right) in centromeres across chromosomes. The asterisks denote the superfamily composition of all *CEN155* and the other stacked columns represent *CEN155* composition within moHRs. **d**, Centromere structures of two exemplar chromosomes. Annotation tracks display information including superfamily, LTR, strand, moHR, diHR and muHR. Zoomed-in views of seven moHRs illustrate fine HOR-like structures. **e**. Biased copy number distribution for the two monomers within a dimer. **f**, Regularity in the copy number for the two monomers from a dimer along the diHR.

Figure 5 | Sequence divergence and mutation rate of rice centromeres
a, Centromere synteny inferred using SynPan-CEN. **b**, Schematic of centromere structural variations (CSVs), including contraction and expansion. **c**, CSVs in centromeres of GJ and XI compared to the corresponding reference assemblies NIP and NJ11, respectively. **d**. CSVs in individual genomes

associated with the phylogenetic kinship. **e**, Relative sequence divergence in centromeres compared to
intergenic regions, estimated from syntenic *CEN155* pairs. **f**, Phylogenetic analysis of *CEN155*
sequences from Chr02 centromeres of SL032 and NIP, indicating non-ortholog syntenic pairs (red, blue
and green links). **g**, Phylogenetic analysis of *CEN155* sequences from Chr03 centromeres of CX20 and
NJ11, indicating non-ortholog syntenic pairs (red and blue links). **h**, Phylogenetic analysis of *CEN155*
sequences from Chr04 centromeres of CX20 and NJ11, indicating absolutely orthologous pairs of
syntenic *CEN155* satellites. **i**, Relative mutation rates for 17 centromere pairs with absolutely
orthologous *CEN155* satellites, compared to non-centromere regions.

1390

Figure 6 | Epigenetic profiling and centromere repositioning
a, Correlation analysis showing relationships among variations in chromosome length, *CEN155* array
length, CENH3 occupancy signal, TE density and neocentromere formation tendency (NFT). **b**, CENH3
ChIP enrichment and element annotation across Chr04 centromeres. **c**, CENH3 enrichment levels in the
CEN155 satellites, centromeric LTRs, pericentromeric LTRs and LTRs in chromosome arms. **d**, DNA
methylation levels in the CG, CHG and CHH sequence contexts within centromeric and flanking
regions of chromosomes Chr10 and Chr11 in a GJ accession SL170.

1398

1399 **Figure 7 | The RICE (Retrotransposon Induced Centromere Evolution) model**

1400

Extended Data Figure 1 | Variation in rice centromere positioning. **a**, Chromosome length and the ratio
of long arm length to short arm length across each chromosome. **b**, Relationships between chromosome
length and centromere length (defined by the presence of *CEN155* satellites) across different taxonomic
groups. **c**, Two megabase-scale inversions observed in the centromere region of chromosome Chr06.

Extended Data Figure 2 | Centromere divergence and introgression in rice evolution. **a-c**, Centromere
similarity and structural variations compared to NIP on chromosomes Chr02 (**a**), Chr05 (**b**), and Chr04
(**c**).

Extended Data Figure 3 | Non-canonical satellites in *CEN155* arrays. a, Schematic representation of
a centromeric region with non-canonical satellites (*sati*). **b**, Length distribution of *sati* between
canonical *CEN155* satellites showing the high-frequency non-canonical satellites. **c**, Origin of *sati*
sequences derived from *CEN155* repeats. **d**, Comparison of the two Chr10 centromeres indicating the
functional and evolutionary dynamics of CEN155+ arrays incorporating *sati94*. StainedGlass similarity
heat maps suggest centromere duplication in NH284 (*O. barthii*), compared to NH265 (*O. glaberrima*),
where the CEN155+ arrays composed of *CEN155* and *sati94* are marked from #1 to #3. CENH3 ChIP
occupancy plots for the two centromeres indicate the functional consistency of *sati94* as *CEN155*.

Extended Data Figure 4 | Divergence sites between *CEN155* superfamilies. The CENP-B box-like and
pJα-like motif regions are shown.

Extended Data Figure 5 | Inference of multimers and muHRs in rice centromeres. The de Bruijn graphs
are constructed based on the dimer-compressed satellite string of each centromere.

Extended Data Figure 6 | CENH3 ChIP enrichment and element annotation across Chr05 centromeres.

1421

Supplementary Figure Legends

Supplementary Fig. 1 | Genomic features and quality assessments of rice genomes in this study. QV, consensus quality value; AQI, assembly quality indicator value; GCI, assembly continuity index, derived from contig N50.

Supplementary Fig. 2 | Accuracy and continuity assessments of rice centromere assemblies. **a**, Base-level accuracy of centromere sequences represented by QV, where "+inf" indicates no erroneous bases in the centromere assemblies. **b**, Potential *k*-mer discordance in centromere assemblies of CW03 (Or-4), NH265 (Ogla), and NJ11 (XI1A), detected by VerityMap. **c**, CGI scores for rice centromere assemblies. **d**, Comparison of centromere assemblies generated by Hifiasm and Verkko, showing high consistency.

Supplementary Fig. 3 | Relationship between centromere length and chromosome length across 12 chromosomes among different taxa.

Supplementary Fig. 4 | Impacts of TEs on chromosome size variation. **a**, Relationship between centromere length and chromosome size within each chromosome. **b**, Relationship between the length of TE sequences outside of centromeres and chromosome size. **c**, Relationship between centromere length and chromosome size in *Arabidopsis thaliana* genomes.

Supplementary Fig. 5 | Sequence similarity and haplotypes of rice centromeres revealing intra-species introgression and rearrangement events across chromosomes. Left, phylogenetic tree based on chromosome arm SNPs. Middle, heat maps showing the percentage of sequence similarity shared by pairwise centromeres. Right, haplotype assignments of rice centromeres.

Supplementary Fig. 6 | StainedGlass sequence similarity heat maps comparing within- and between the Chr12 centromeres of CW09, SL044 and NJ11.

Supplementary Fig. 7 | Distribution of satellite repeat length across the 70 rice genomes in each taxonomic group (Obar, Ogla, Oruf, AUS, GJ, and XI).

Supplementary Fig. 8 | Characteristics of the fifteen *CEN155* superfamilies. **a**, Distribution of *CEN155* length within each superfamily. **b**, Edit distances among superfamilies.

Supplementary Fig. 9 | TE invasion landscape in rice centromeres. **a**, TE proportions and accumulated lengths in centromere regions across chromosomes, colored by different TE super-families. **b**, TE proportions among rice genomes, shown according to the phylogenetic tree. From left to right, the circled diagrams represent repeat proportions in centromere, composition in centromeric TEs, and LTR composition in centromeric TEs, respectively. The size of the circle represents the sum of cumulative length. **c**, TE proportions across chromosomes with orange point representing centromere region and blue points stand for genome-wide region.

Supplementary Fig. 10 | Features of intact centrophilic LTR families (*SZ-22*, *RIRE7* and *CRM*) in rice

centromeres. **a**, Phylogeny of all LTRs in the pan-TE library based on interval sequences, showing

distant relationship between *SZ-22* and other two centrophilic LTRs. Triangles in different colors

represent the domains of gag, RVT, RNaseH, Integrase_H2C2 and rve, respectively. **b**, Length

distributions of *SZ-22*, *RIRE7* and *CRM* identified within the centromere, pericentromere, and

chromosome arm regions.

Supplementary Fig. 11 | Integration frequency of centrophilic LTR *SZ-22* along the *CEN155* consensus

sequence.

Supplementary Fig. 12 | Organization features of non-canonical satellites *sati94*. **a**, The genomic

distance (measured by physical distance and *CEN155* copy number) between adjacent non-canonical

satellite *sati94* in the three CEN155+ arrays indicating local homogenization. **b**, The

maximum-likelihood phylogeny of *sati94* sequences from the three CEN155+ arrays in Chr10

centromeres of NH284 and NH265. Bootstrap values are shown at the branches.

Supplementary Fig. 13 | Local homogenization at the monomer, dimer, and multimer scales. **a**,

Distribution of *CEN155* satellites exhibiting monomer homogenization across the fifteen superfamilies

in 70 rice accessions from various taxa. Numbers of dimers and multimers for the top fifteen groups are

shown in **b** and **c**. Each rice accession is colored by taxonomic groups at the top, and summed up at the

bottom. The total numbers of monomers, dimers, and multimers are showed on the right.

Supplementary Fig. 14 | Biased monomer copy numbers across dimers and taxonomic groups. One

monomer has a relatively constrained and stabilized copy number (nearly 1), while the other exhibits

variable copies (forming moHRs) within a diHR, represented by XY_m or X_mY.

Supplementary Fig. 15 | Local homogenization of copy number bias within a dimer through tandem

duplication. **a**-**d**, Repeat of bias pattern in monomer copy number of dimers in NIP(GJ), NJ11(XI),

SL022(AUS) and NH265(Ogla).

Supplementary Fig. 16 | Workflow of SynPan-CEN. (1) Stepwise backbone building. All vs all

pairwise satellite edit distance (ED) values were used to construct the Directed Acyclic Graph (DAG).

To enhance accuracy in constructing synteny, the chain-backbone was built with the most conservative

and highly-ordered monomer pairs by gradually increasing the ED value thresholds until the framework

contained more than 20% of the total monomers or 100 monomers. This framework was then split by 10

monomer pairs within one window for downstream analysis. (2) Local satellite rescuing. DAGs were

used again to rescue chains of syntenic satellite pairs within each window, employing relaxed EDs of 15.

The sub-blocks with the highest scores were retained, and any non-overlapping monomer pairs were

added to complete the matches. (3) Blocks from the chain-backbone were iteratively merged.

Supplementary Fig. 17 | Synteny ratio of *CEN155* satellites in rice centromeres against NIP (GJ-tmp)

centromeres.

Supplementary Fig. 18 | Heat map showing the synteny ratio between NIP (GJ-tmp) and other rice

centromeres across 12 chromosomes, with Chr05 centromeres showing a significantly higher syntenic

relationship between XI and GJ.

Supplementary Fig. 19 | Satellite syntenic relationships between centromeres of J4155S, SL148, and

XL628S with NJ11 on Chr05, and CX20, LK638S on Chr10. Tracks show centromere annotations,

including TE families, *CEN155* strands, *CEN155* superfamilies, and edit distances, respectively.

Supplementary Fig. 20 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr01. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 21 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr02. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 22 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr03. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 23 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr06. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 24 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr07. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 25 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr08. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 26 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr09. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 27 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr10. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 28 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr11. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 29 | Genomic features of the functional centromere and its flanking regions on

chromosome Chr12. Top, CENH3 ChIP-seq enrichment ($\log_2$(ChIP/input), two replicates) in 10-Kb

windows. Track 1, satellite superfamilies; Track 2, LTRs *SZ-22*; Track 3, LTRs *CRM*; Track 4, LTRs

RIRE7; Track 5, *sati* (non-canonical satellites).

Supplementary Fig. 30 | Correlation between the length of CENH3 enriched regions and

CEN155-defined centromere size for each chromosome.

Supplementary Fig. 31 | Correlation between chromosome length and TE density across the 12

chromosomes.

Supplementary Fig. 32 | Correlation between the chromosome length and *CEN155* array length across

the 12 chromosomes.

Supplementary Fig. 33 | Correlation between the chromosome length and CENH3 enrichment length

across the 12 chromosomes.

Supplementary Fig. 34 | Correlation between the *CEN155* array length and neocentromere formation

tendency (NFT) across 12 chromosomes.

Supplementary Fig. 35 | Correlation between the TE density in the *CEN155* array and neocentromere

formation tendency (NFT) across 12 chromosomes.

Supplementary Fig. 36 | Correlation between the chromosome length and neocentromere formation

tendency (NFT) across 12 chromosomes.

Supplementary Fig. 37 | Comparison of profiled DNA methylation probability from ONT sequencing

data and HiFi data. Each point represents a randomly selected 5mC site.

1552

Supplementary Table legends

Supplementary Table 1 | The source, sequencing, assembly and QC information of 70 rice genomes in

this study.

Supplementary Table 2 | Detailed annotations of centromeres and chromosome arms in the 70 rice

genomes.

Supplementary Table 3 | Gaps in the rice centromere assemblies.

Supplementary Table 4 | Putative centromere introgression events in rice evolution.

Supplementary Table 5 | Frequency of non-canonical satellites (*sati*) in rice centromeres.

Supplementary Table 6 | Satellite annotation in moHRs.

Supplementary Table 7 | Satellite annotation in diHRs.

Supplementary Table 8 | Satellite annotation in muHRs.

Supplementary Table 9 | Centromere structural variations (CSVs) in rice genomes.

Supplementary Table 10 | Relative divergence and mutation rate estimation for rice centromeres.

Supplementary Table 11 | Functional centromeres in rice genomes defined by CENH3 ChIP data.

1567

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

CEN155 satellites ▮ Centrophilic LTR ⬠⬠⬠ CENH3 ChIP signal (high to low)
— Centromere — CEN155 (+) — CEN155 (−) ⋯ Peri-centromere

Extended Data Figure 1 | Variation in rice centromere positioning. **a**, Chromosome length and the ratio of long arm length to short arm length across each chromosome. **b**, Relationships between chromosome length and centromere length (defined by the presence of *CEN155* satellites) across different taxonomic groups. **c**, Two megabase-scale inversions observed in the centromere region of chromosome Chr06.

Extended Data Figure 2 | Centromere divergence and introgression in rice evolution. **a-c**, Centromere similarity and structural variations compared to NIP on chromosomes Chr02 (**a**), Chr05 (**b**), and Chr04 (**c**).

Extended Data Figure 3 | Non-canonical satellites in *CEN155* arrays. **a**, Schematic representation of a centromeric region with non-canonical satellites (*sati*). **b**, Length distribution of *sati* between canonical *CEN155* satellites showing the high-frequency non-canonical satellites. **c**, Origin of *sati* sequences derived from *CEN155* repeats. **d**, Comparison of the two Chr10 centromeres indicating the functional and evolutionary dynamics of CEN155+ arrays incorporating *sati94*. StainedGlass similarity heat maps suggest centromere duplication in NH284 (*O. barthii*), compared to NH265 (*O. glaberrima*), where the CEN155+ arrays composed of *CEN155* and *sati94* are marked from #1 to #3. CENH3 ChIP occupancy plots for the two centromeres indicate the functional consistency of *sati94* as *CEN155*.

Extended Data Figure 4 | Divergence sites between *CEN155* superfamilies. The CENP-B box-like and Pjα-like motif regions are shown.

Extended Data Figure 5 | Inference of multimers and muHRs in rice centromeres. The de Bruijn graphs are constructed based on the dimer-compressed satellite string of each centromere.

Extended Data Figure 6 | CENH3 ChIP enrichment and element annotation across the Chr05 centromeres.