Ecology and abundant occurrence of *Genea* (Pyronemataceae) in western Oregon, U.S.A.

Authors: Hilary Rose Dawson[1,2,3†]* and Heather A. Dawson[2,3†]

Affiliations: 1. Australian National University, Acton, ACT 2601, Australia; 2. Institute of

Ecology and Evolution, University of Oregon, Eugene, OR 97402, USA; 3. Department of

Biology, University of Oregon, Eugene, OR 97402, USA; † indicates authors contributed equally

Corresponding author: *hilary.dawson@anu.edu.au

ORCIDs: H.R.D 0000-0003-4613-762X; H.A.D. 0009-0002-4110-1257

Keywords: hypogeous fungi, truffles, Pezizales, cup fungi, U.S. Pacific Northwest, conservation

detection dogs

Abstract

Genea (Pyronemataceae, Ascomycota) develops hypogeous ascomata (truffles) that are

often small and soil-colored. This group is considered common but challenging to find. We used

a truffle diversity dog to locate *Genea* truffles in the southern Willamette Valley and associated

mountains in western Oregon, U.S.A, recording when the sporocarps were present at a wide

range of elevations (113 to 1879 m). We found *Genea* was present in half of our surveys, and

noted that it rarely fruited in areas that had experienced wildfire. This study demonstrates the

value of truffle diversity dog surveys for documenting fungi that are difficult to locate visually,

and provides further evidence for the abundance of *Genea*.

Introduction

The truffle genus *Genea* (Pyronemataceae, Ascomycota) is characterized by lobed and

often inconspicuous hypogeous ascomata and was originally circumscribed in Italy in the 1800s

(Vittadini 1831). It is considered common in both Europe and North America (Alvarado et al.

2016), but because it fruits underground and is often difficult to differentiate from soil

aggregates, we have limited knowledge of its ecology and distribution. One of the most

comprehensive studies in the U.S. Pacific Northwest focused on oak grasslands (Smith et al.

2006) and described two new species. However, *Genea* forms ectomycorrhizal associations with

a wide variety of plants (Smith et al. 2006), suggesting that it may be widely distributed

throughout the Douglas-fir (*Pseudotsuga menziesii*) forests of the Cascades and Coast Range.

We tested this possibility using a novel tool: a truffle dog with an uncanny ability to find *Genea*

truffles.

Truffle dogs are most commonly trained to find mature culinary truffles that command a

high commercial price. However, some dogs have a natural inclination towards generalizing on

any hypogeous odorous sporocarps. If the trainer rewards all finds (rather than just culinarily

valuable species), they can train "diversity dogs" who can locate a broad range of truffles that

may be of interest to science. Most truffle surveys are currently performed by raking, where a

researcher uses a small garden hand rake to peel back the upper layer of the soil. This method

has a visual bias towards what the researcher spots under the duff, and often misses rare and

inconspicuous taxa. In contrast, truffle dogs have a scent bias driven by mature sporocarps of any

size or color, can find truffles deeper than most people rake, and do not have preconceived ideas

of where truffles 'should' be. By documenting which truffle species are found throughout a

range of sites and seasons, we can use truffle diversity dogs to broaden our understanding of

hypogeous fungal ecology.

Methods

All truffles were located by trained truffle diversity dog Rye, who had found 23 genera of

truffles in two years of truffling as of February 2023. Rye demonstrated particular persistence in

locating *Genea* truffles as of the time of this study. He received a reward for each *Genea* truffle

located, reinforcing his pursuit of truffles with this type of aroma.

We performed 87 general truffle surveys and forays in all seasons between October 2021

and November 2022. We visited 66 sites, 12 of which we visited more than once throughout the

year (Figure 1; Table S1). Most surveys were conducted along existing public trails that allowed

us easy access to varied landscapes. We performed these surveys at a range of elevations from

113 m to 1879 m (elevation data extracted from WorldClim (Fick and Hijmans 2017)). On each

survey, we primed Rye to find truffles of any species and rewarded his finds with praise and a

brief game of fetch with a tennis ball (his favorite reward). He was given freedom to roam off-

leash with voice recall during these surveys so that he could cover more ground and follow his

nose.

Figure 1: Sites and methods used to document *Genea* ecology. A) Map of all sites surveyed as
part of this study. Blue points are sites where we found at least one *Genea* truffle. Symbols
indicate elevation category (circle = high elevation, triangle = mid elevation, square = low
elevation). B) Example of truffle dog Rye finding a truffle. C-E) Photographs of *Genea* truffles
demonstrating lobing, dark peridium and light gleba, and sometimes very small size and
resemblance to soil clusters.

We conducted our surveys primarily in conifer-dominated forests of varying age classes

and densities, with the exception of the site at Mt. Pisgah which was oak savanna (Table S1).

Our sites had a variety of fire histories, from no recent fire to recently burned (1-11 years). Most

low-mid elevation conifer forest sites were primarily composed of Douglas-fir (*Pseudotsuga*

menziesii) and western hemlock (*Tsuga heterophylla*), and high elevation sites often included

mountain hemlock (*Tsuga mertensiana*), true fir (*Abies* spp.), and pine (*Pinus* spp.). All sites

were located in Benton, Deschutes, Lane, and Linn Counties in southern Willamette Valley,

Oregon, USA. These counties encompass the western Cascades, southern Coast Range, and

Willamette Valley. We excluded observations from coastal Lane County because we performed

only spatiotemporally limited surveys on the coast.

We only recorded truffle presence/absence at each site. Where multiple truffles were

found, this counted as a single 'presence' for a site. If we surveyed multiple habitat types at a

single location (for example, inside and outside a burn area), this was counted as multiple sites.

Elevation was categorized as 'low' (0-750 m), 'mid' (750-1350 m), and 'high' (>1350 m) based

on observed changes in plant communities and general ecology for the area.

Results

We found *Genea* fruiting bodies in slightly over half (51.7%) of the site surveys. We

reported occurrences of *Genea* nearly year-round, except the months of August and September

(Figure 2) which are the driest months of the year due to western Oregon's Mediterranean

climate. Low and mid elevation sites had the most consistent *Genea* presence. The highest site

we found *Genea* at was at 1716 m elevation. *Genea* truffles were not present in any burned

conifer forests, but were found during two oak savanna surveys that had burned three years prior.

Figure 2: *Genea* **presence in the southern Willamette Valley and associated mountains throughout a single year.** Numbers within the bars indicate the number of surveys where *Genea* was found (blue) or not found (gray). Parenthetical numbers indicate how many of those sites were burned.

Odor and morphology of the truffles suggested that there are likely multiple species present. We noted that some visually similar truffles collected from shared locations had a mellow cheese odor, while others had a sharp garlic-truffle odor. Microscopic examination of these particular variants showed ellipsoid spores in the former and subglobose spores in the latter. Subsequent examinations of additional collections suggest extensive hidden diversity in the dark-colored *Genea* found in this region.

Discussion

Using a dog trained to find truffle diversity, we found *Genea* grew abundantly nearly
year-round. In the literature, *Genea* is known to be common (Alvarado et al. 2016); however,
there can be an impression of local scarcity due to the difficulty of finding small and soil-like
sporocarps. The commonness of *Genea* suggests that this genus may be a key ectomycorrhizal
partner for trees in the area, especially the dominant Douglas-fir timber trees. Although the
decrease in abundance with elevation may be partially due to a sampling bias, our findings
suggest that *Genea* is most prevalent at lower elevations. Douglas-fir trees have limited
distribution at higher elevations (Case and Peterson 2005), where the landscape is more likely to
be dominated by mountain hemlock (*Tsuga mertensiana*) or true firs (*Abies*), enforcing the
likelihood of a strong mycorrhizal association between Douglas-fir and *Genea*.

Our surveys demonstrate that human truffle surveys have a visual bias that can be
overcome by complementing with scent surveys. The dog strongly indicates, requiring the
accompanying human to look more closely and find the "hidden" *Genea*. However, our surveys
were limited to the abilities of the truffle dog. He cannot find truffles in hot weather, so our
summer surveys were focused at higher elevations where the temperatures were cooler.
Similarly, we were unable to access higher elevation sites until after snowmelt and the roads had
opened, but that also would have been true of the more traditional truffle-finding method of
raking.

Genea truffles are difficult to identify from morphological features alone, and even
spores can be indistinct. Most *Genea* collections from conifer forests in the Pacific Northwest
have typically been identified as *G. harknessii* (species complex) or *G. gardneri*, as the only two
species of dark *Genea* currently known from this habitat (Smith et al. 2006; Trappe et al. 2007).

We have observed that there are likely several additional species that are morphologically similar

and difficult to distinguish without microscopic examination and sequence data. In some

instances we noted the change in odor and texture of the *Genea* we found, but we did not

consistently track these traits given that these characteristics do not reliably separate different

species in many hypogeous fungi. This group would benefit from further taxonomic study using

modern molecular methods to determine which species are present in the area in the different

habitats and elevations.

Eleven of our sites were burned, including eight conifer sites that had experienced

wildfire in the past 1 to 11 years. In light of the increasing frequency and severity of wildfire in

western Oregon (Halofsky et al. 2020; Dye et al. 2024), the lack of *Genea* at these sites is

notable and suggests that some species of *Genea* may be sensitive to wildfire, at least in the

earlier years of recovery. On the other hand, we did find *Genea* during both surveys at the

burned oak savanna site, suggesting that not all *Genea* species may be equally affected by fire.

This genus is relatively abundant in the Willamette Valley; however, it may decline in

abundance as Douglas-fir forests are affected by climate change and high severity fire events.

Our observations of *Genea* as ubiquitous in modified and disturbed habitat suggests it is resilient

to anthropogenic effects, but extreme fire may be an exception for some species. We must

prioritize document fungal species distributions, particularly ones that are difficult to locate like

Genea, to understand what can be lost through climate change.

Acknowledgements

Bitty Roy provided valuable feedback on this manuscript. This research would not have

been possible without Heather's truffle dog–good boy, Rye.

Disclosure Statement

The authors report there are no competing interests to declare.

References

Alvarado P, Cabero J, Moreno G, Bratek Z, Vooren N van, Kaounas V, Konstantinidis G, Agnello C, Merényi Z, Smith ME, Vizzini A, Trappe JM. 2016. Phylogenetic overview of the genus *Genea* (Pezizales, Ascomycota) with an emphasis on European taxa. Mycologia 108:441–456.

Case MJ, Peterson DL. 2005. Fine-scale variability in growth climate relationships of Douglas-fir, North Cascade Range, Washington. Canadian Journal of Forest Research 35:2743–2755.

Dye AW, Reilly MJ, McEvoy A, Lemons R, Riley KL, Kim JB, Kerns BK. 2024. Simulated future shifts in wildfire regimes in moist forests of Pacific Northwest, USA. Journal of Geophysical Research: Biogeosciences 129:e2023JG007722.

Fick SE, Hijmans RJ. 2017. WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. International Journal of Climatology 37:4302–4315.

Halofsky JE, Peterson DL, Harvey BJ. 2020. Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA. Fire Ecology 16:4.

Smith ME, Trappe JM, Rizzo DM. 2006. *Genea, Genabea* and *Gilkeya* gen. nov.: ascomata and ectomycorrhiza formation in a *Quercus* woodland. Mycologia 98:699–716.

Vittadini C. 1831. Monographia Tuberacearum. Mediolani.

174