Direct quantitative assessment using digital droplet PCR and field-scale spatial

distribution of *Plasmopara viticola* oospores in vineyard soil

Charlotte Poeydebat[1]#, Eva Courchinoux[2], Isabelle D. Mazet[2], Marie Rodriguez[2],

Alexandre Chataigner[2], Mélanie Lelièvre[3], Jean-Pascal Goutouly[4], Jean-Pierre

Rossi[5], Marc Raynal[6], Laurent Delière[2], François Delmotte[2]

[1] Bordeaux Sciences Agro, INRAE, SAVE, ISVV, Gradignan, France

[2] INRAE, Bordeaux Sciences Agro, SAVE, ISVV, Villenave d'Ornon, France

[3] Université Bourgogne Franche-Comté, INRAE, AgroSup Dijon, Agroécologie, Dijon,

France

[4] INRAE, Université de Bordeaux, Bordeaux Sciences Agro, EGFV, ISVV, Villenave

d'Ornon, France

[5] INRAE, CIRAD, IRD, Institut Agro, CBGP, Montferrier-sur-Lez, France

[6] IFV, Bordeaux Nouvelle Aquitaine, UMT SEVEN, Villenave d'Ornon, France

Running title: Quantitative assessment of *P. viticola* oospores in soil

#Address correspondence to Charlotte Poeydebat, charlotte.poeydebat@agro-

bordeaux.fr

Abstract

Grapevine downy mildew, caused by the oomycete *Plasmopara viticola*, is one of the most devastating diseases of grapevine worldwide. Primary inoculum is increasingly thought to play a decisive role in downy mildew epidemics, but we still know very little about its abundance in vineyard soil. We present a direct and sensitive method for the quantification of *P. viticola* oospores (i.e. primary inoculum) in vineyard soil, based on digital droplet PCR and an estimation of the number of copies of the ITS sequence used for amplification in the genome of *P. viticola*. We used this method to describe the spatial distribution of *P. viticola* oospores in a vineyard. We found oospores in all samples except one, and concentration ranged from 0 to 1858 oospores per gram of soil (303 ± 308 on average). The distribution of oospores at field scale was not random but characterized by 25 m-diameter patches of concentrically increasing oospore concentration. Oospores accumulated 5 times more below the vine stocks than in the inter-row. We conducted a leaf disc bioassay to assess soil infectious potential on a subset of our samples. Soil infectious potential significantly increased with oospore concentration, suggesting that DNA-based methods can have an epidemiological significance. Quantitative data on *P. viticola* oospores could be useful to design sampling strategy for future research or to improve epidemic risk models.

Importance

Grapevine downy mildew is an airborne disease caused by the oomycete Plasmopara viticola affecting leaves and bunches and leading to important economic losses for viticulturists. Its control mainly relies on the use of fungicides during the growing season. Due to their harmful effects on human health and biodiversity,

European and national policies encourage pesticide use reduction either by

developing alternative control strategies or by optimizing use efficiency. Recently,

evidences have accumulated that primary inoculum (i.e. oospores in the soil)

importantly contributes to disease progress. The significance of our work is in

presenting a direct and sensitive method for assessing soil oospore concentration,

as well as quantitative and spatially-explicit data on downy mildew primary inoculum,

opening the way to new research, the evaluation of new disease control strategies

based on primary inoculum management and the improvement of epidemic risk

models, that will in fine contribute to lower fungicide use in viticulture.

Keywords: oomycete, downy mildew, primary inoculum, geostatistics, ddPCR,

bioassay, soil infectious potential

Introduction

While crop protection from pest and diseases is largely ensured by the use of

pesticides worldwide (Sharma et al., 2019), there is accumulating evidence that they

are responsible for environmental pollution threatening both biodiversity (Wan et al.,

2025; Beaumelle *et al.*, 2023; Beketov *et al.*, 2013) and human health (*e.g.*

Dall'Agnol *et al.*, 2021). In Europe and France, policies are clearly encouraging the

reduction of the use of chemical plant protection products (European Green Deal

and National Action Plan Ecophyto; Lamichhane *et al.*, 2019), either by improving

use efficiency or by developing alternative pest and disease management strategies.

Grapevine downy mildew, caused by the obligate biotrophic oomycete *Plasmopara*

viticola, is one of the most devastating diseases of grapevine worldwide. Its

management still relies mainly on intensive use of fungicides throughout the growing

season. In 2006, the French wine-growing area covered 3.3% of utilized agricultural

land but contributed up to 14.4% of national fungicide use (Butault et al., 2011). In

2019, more than 80% of the treatment frequency index was made up of fungicides

used against downy mildew and powdery mildew (Agreste, 2023).

In temperate regions, the life cycle of *P. viticola* comprises a phase of asexual

multiplication on its host tissues during the growing season and a phase of sexual

reproduction, which occurs in infected grapevine leaves between thalli of sexually

compatible strains (as *P. viticola* is heterothallic, Wong *et al*., 2001) leading to the

formation of oospores. Leaves then fall on the ground and oospores overwinter in

leaf debris or as free propagules in the soil. The stock of oospores in the soil

constitutes the primary inoculum for downy mildew epidemics in the next growing

season. In contrast to the widely-held assumption that primary contaminations rarely

occur after flowering, primary inoculum is increasingly considered to play a major

role in grapevine downy mildew epidemics (Gessler *et al*., 2011) for several reasons.

First, population genetic studies evidenced that grapevine downy mildew populations

are panmictic across European vineyards (Fontaine *et al*., 2021, 2013) showing that

genetic mixing resulting from sexual reproduction largely structures pathogen

populations. Second, it has been shown that oospores are able to germinate and

cause infections throughout the growing season (until late September in the northern

hemisphere; Kennelly *et al*., 2007; Maddalena *et al*., 2021) and that a substantial

amount of infections, including in the middle of the growing season, can arise from

oospores (up to 50% of contaminations; Gobbin *et al*., 2003; Hong *et al*., 2020).

Third, while the epidemic continues to progress after harvest (from what we observe

in surveyed vineyards), the incidence of typical mosaic-like downy mildew symptoms

on grapevine leaves was shown to be proportional to the amount of oospores formed

in the leaves (Si Ammour *et al.*, 2020). Finally, downy mildew incidence at the end of

the season was found to explain the precocity and intensity of the disease the

following season in Canadian vineyards (Carisse, 2016).

While actual control of grapevine downy mildew primarily targets the asexual

propagation of the disease, the notable contribution of *P. viticola* primary inoculum to

epidemic dynamics suggests that measures based on primary inoculum

management or the limitation of primary contaminations could make a significant

contribution. At least, decisions on the application of fungicides should better

account for the abundance of primary inoculum to better assess the real epidemic

risk. Although the stages of the sexual reproduction cycle of *P. viticola* are well

known (see for example Gessler *et al.*, 2011), their quantitative description remains

incomplete. The study of disease inoculum abundance and spatial distribution in

agricultural soils are generally restricted to soilborne pathogens, including

oomycetes (see for example Botero-Ramirez *et al.*, 2021; Moussart *et al.*, 2009). To

our knowledge, these questions have rarely been addressed regarding airborne

pathogens with a telluric conservation life stage. Molecular approaches have been

developed and used to quantify airborne asexual inoculum of grapevine downy

mildew, as well as oospores in grapevine leaves (Carisse *et al.*, 2021; Douillet *et al.*,

2022; Si Ammour *et al.*, 2020; Valsesia *et al.*, 2005). Direct indicators of the

pathogen abundance in the vineyards, such as spore concentration in the air, are

progressively included in decision tools or regional monitoring programs (Brischetto

et al., 2020; Douillet, 2023). Yet, the prediction of downy mildew epidemic risk in

decision support systems is mainly based on meteorological variables that control

the development of *P. viticola* (Dubuis *et al.*, 2012; Park *et al.*, 1997; Rossi *et al.*,

2008a; Strizyk *et al.*, 1983). Indicators of primary inoculum reservoir in the soil

compartment have never been accounted for until now, partly because quantitative

data is scarce. Instead, predictive models assume a fictitious quantity of primary

inoculum and calculates rates of change (associated with maturation, dispersal or

germination; *e.g.* Rossi *et al.*, 2008a). Recently, a molecular method based on real-

time qPCR was used to quantify oospores in soil of vineyards in China (Yang *et al.*,

2023). Quantitative data on oospores that could be gained from such methods could

contribute to the transition of wine-growing towards more sustainable downy mildew

management strategies by improving epidemiological models for epidemic risk

prediction, and allowing evaluation of disease control measures based on primary

inoculum management.

Here, we present a direct and sensitive method for the quantification of *P.viticola*

oospores in vineyard soil based on the quantification of *P. viticola* DNA by digital

droplet polymerase chain reaction (ddPCR). We provide an estimation of the number

of copy of the ITS sequence used for amplification (ITS1/5.8S) in the genome of

Plasmopara viticola that we used to convert the ddPCR output into a number of

oospores per gram of soil. By means of geostatistics, we described the spatial

distribution of downy mildew primary inoculum in a vineyard at the onset of the

growing season. If *P. viticola* oospores can persist in the soil and remain infectious

for several years (Caffi *et al.*, 2011), the determinants of oospore germination rate

are poorly understood in natural conditions and the effect of aging of oospores on

germination probability remains unstudied to date. However, we hypothesized that

the DNA-based quantification of primary inoculum in the soil is positively related to

grapevine primary contaminations. Building on previous works, we finally conducted

a bioassay to assess the infectious potential of soil samples and discuss the

epidemiological significance of our DNA-based assessments of downy mildew

primary inoculum.

Material & Methods

<u>Soil sampling and preparation</u>

We conducted our study in an experimental vineyard with a sandy-gravel soil located

at the INRAE Bordeaux Nouvelle-Aquitaine research center (Villenave d'Ornon,

France; 44.79194, -0.57702) within the Pessac-Léognan appellation area. The

surveyed vineyard was planted in 2011 with 18 rows of 68 Merlot grapevine stocks

over a 0.22 ha surface (70 x 32 m), corresponding to a density of 6600 vines per ha

(*i.e.*, 1.6 m between rows and 0.95m between vines along the row). It was managed

according to French organic farming specifications, meaning that Bordeaux mixture

(copper) was the only chemical fungicide allowed to control downy mildew. Inter-

rows were either sown with a mix of plant species selected for green manuring or

covered by spontaneous vegetation in an alternating arrangement rotating every

year. All inter-rows were tilled before grapevine budburst. For winter protection and

weed management purposes, soil from both sides of the row was turned over on the

lower part of the vines in the fall. The ridge formed under the row was then reopened

in the spring. The surveyed vineyard was bordered on each of its long sides by a

diversified shrubby hedgerow (*ca*. 1.5-2 m high) that separated it from other

experimental vineyards.

Soil sampling was done in February 2022. We collected 198 soil samples at the

nodes of a 2.85 x 3.2 m regular grid covering the entire surface of the vineyard; *i.e.*

one sample every three vine stocks in the row, in one row out of two (Figure 1). To

explore the spatial structure of primary inoculum of downy mildew in the soil at a finer scale, we delineated five plots in the experimental vineyard within which we collected soil samples following a nested sampling design with three distance levels. Each plot consisted of a 2.85 x 3.2 m rectangle containing 28 sampling points (Figure 1). The first distance level corresponded to the four corners of the plot (n_3=4), hence sampling points were separated by about 3 m (samples were the same as for the regular grid sampling). The second distance level was set at 0.8 m (half the inter-row width), and sampling points were located in the corners of four 0.8 m squares located in each corner of the plot ($n_{0.8}$=16, including 4 points in common with the first distance level; see Figure 1). The third distance level was set at 0.2 m with sampling points corresponding to the corners of four 0.2 m squares located in the corners of one of the four 0.8 m squares ($n_{0.2}$=16, including 1 point in common with the first distance level and 4 points in common with the second distance level; see Figure 1). We also considered that the points from the regular grid were located in the row, while the points from the 2nd and 3rd levels of the nested sampling were located in the inter-row space. At each sampling point of both the regular grid and the nested sampling plots, we collected 500 g of soil from the top 15 cm using an auger. We ended with 318 soil samples, including 198 samples from the regular grid and 120 additional samples from the five nested sampling plots.

To investigate the vertical distribution of *P. viticola* primary inoculum in the soil, we dug a hole 40 cm deep and 30 cm in diameter at six points located in the inter-row space in the experimental vineyard (Figure 1). We divided the soil profile into four strata of 10 cm (0-10 excluding leaf debris on soil surface, 11-20, 21-30 and 31-40 cm) and sampled 500 g of soil from each stratum, starting at the bottom of the profile to avoid contaminating deeper layers with upper ones. This corresponded to 24

194   samples in total (4 samples per hole or sampling location). All soil samples were

195   stored at 5°C before processing.

196
197   **Figure 1.** Map of the soil sampling points within the experimental vineyard.
198

<u>Quantification of *Plasmopara viticola* primary inoculum in vineyard soil samples</u>

DNA extraction

Each sample was passed through a 4 mm mesh sieve and thoroughly homogenized

for 30 s using an electric mixer set at medium speed. Directly after homogenization,

ca. 6 g of the sample was collected and placed in a 5 ml tube to be freeze-dried.

Then, a 2 g subsample of dry soil was placed in a 15 ml Falcon tube containing 4 g

of 0.1 mm diameter silica beads, 5 g of 1.4 mm diameter ceramic beads and eight 4

mm diameter glass beads (as recommended by MP-Biomedicals for TeenPrep™

Lysing Matrix E). For the grinding and lysis extraction step, a volume of 8mL of lysis

buffer composed of 100 mM Tris-HCl (pH 8), 100 mM EDTA (pH 8), 100 mM NaCl,

20% SDS and ultra-pure water was added in each tube. Mechanical cell lysis was

achieved placing the tubes in a FastPrep-24 classic instrument (MP-Biomedicals) set

for three runs of 30 s at 4 m.sec^{-1}. Tubes were then incubated for 30 min at 70°C to

activate chemical lysis and centrifuged at 5000 g for 7 min at 20°C to eliminate

particles in suspension. DNA extraction was carried out using 1 ml of supernatant

obtained from the centrifugation, as follows: 1) *Deproteinization:* addition of 100 µl

potassium acetate 3M pH5.5 to the sample, homogenization and incubation on ice

for 15 min to precipitate proteins, centrifugation at 14000 g for 5 min at 4°C to

eliminate precipitated proteins (supernatant recovery and pellet disposal); 2) *DNA*

precipitation: addition of 900µl cold isopropanol (v/v) to the supernatant from the

deproteinization, gentle agitation of the tubes that were then placed at -20°C for 30

min, centrifugation at 13000 rpm for 30 min at 4°C, elimination of the supernatant; 3)

DNA Washing: rinsing of the pellet with 400µl 70° cold ethanol, centrifugation at

13000 rpm for 5 min at 4°C, elimination of the ethanol supernatant, drying of the

pellet by placing the tubes in a drying oven at 60°C for 20 min; 4) *DNA Purification:*

resuspension of the pellet in 500 µl of ultra-pure water. Raw DNA extracts were

further purified using a NucleoSpin Soil kit (Macherey-Nagel) following the

manufacturer's protocol from step 4 (addition of SL3 buffer). Finally, purified DNA

extracts were eluted in 60 µL of buffer (TrisHCL, 5mM, pH 8.5) and stored at –20°C.

Digital droplet polymerase chain reaction (ddPCR)

To avoid common PCR inhibition problems with the soil matrix and to obtain direct

absolute DNA quantification (no standard curve needed), samples were analyzed by

droplet digital PCR (ddPCR) using the QX200 ddPCR system (Bio-Rad) composed

of a droplet generator, a thermal cycler and a droplet reader. We used the specific

PCR primers and TaqMan probe (Giop set) designed by Valsesia *et al.* (2005) that

target the ITS1/5.8S to amplify *P. viticola* DNA in our samples.

For each sample, we added 4 µl of DNA extract diluted to the tenth to 18 µl of a PCR

mix composed of 11 µL of 2x ddPCR Supermix for Probes (Bio-Rad), 2.2 µL of 7.5

µM Giop F, 2.2 µL of 7.5 µM Giop R, 2.2 µL of 5 µM Giop P-VIC and 0.4 µL of ultra-

pure water. The samples (20 µl) were first placed in the droplet generator to proceed

to the breakdown of the samples into 15000 to 20000 micro-droplets using oil.

Samples were then carefully transferred to 96-well plates with a pipette, with one

sample per well (one single analysis per sample). The sample plates were placed in

the thermal cycler for DNA to be amplified according to the following program: 1) an

initial phase at 95°C for 5 min; 2) 40 cycles of 30 s at 95°C then 1 min at 60°C then

30 s at 72°C; and 3) a final phase at 98°C for 10 min. Samples were cooled at 12°C

before being transferred to the droplet reader for DNA quantification (ratio of positive

vs negative droplets). Finally, we obtained one value per sample that corresponded

to the number of ITS1 copies per µl of reaction mix.

Estimation of ITS1/5.8S copy number in Plasmopara viticola genome

In order to estimate the ITS1/5.8S copy number in the genome of *P. viticola*, we

compared the ddPCR signals obtained with the ITS primer pairs (Valsesia *et al.*,

2005) *vs* primers targeting the single copy β-tubulin gene (Weerakoon *et al.*, 1998).

First, sporangia of seven *P. viticola* strains frequently used in laboratory for genetic

applications were collected with a paintbrush from grapevine (*Vitis vinifera* cv.

Cabernet Sauvignon) leaves infected beforehand on purpose and placed into

separate 2 mL microcentrifuge tubes with 1 mL Tris-EDTA buffer. Second, freeze-

dried samples were extracted following a SDS based method (adapted from Möller

et al., 1992) and DNA quantity and quality were checked by dosage with either Qubit

4 or Denovix-11. About 35 sequences from NCBI (Fontaine *et al.*, 2021; Rouxel *et*

al., 2013) were aligned with *P. viticola* genome using Geneious Prime. Primers

(Forward: ACAGCCAATTTTCGTAAGTCC,   Reverse:

GCCTTGTACGACATTTGCTTT)   and   a   TaqMan   probe   (5'   FAM-

TGTCCGGGAAATCGAAGGCA-MGB 3') were designed using the consensus

sequence with 100% similarity. Specificity was tested using NCBI Blast

(blast.ncbi.nlm.nih.gov) and by running quantitative PCRs on either pure *P. viticola*

strain, *P. viticola* with grapevine leaf or grapevine leaf only (see Figure S1 in

Supplementary Materials) using a QuantStudio 5 (Life Technologies). Reaction mix

was composed of 10 µL of 2X PCR Buffer (Takyon Low Rox Probe MasterMix -

Eurogentec), 1,6 µL of each primer at 10 µM, 0,4 µL of probe at 10 µM, 3,9 µL of

molecular biology water (Sigma) and 2,5 µL of DNA template for a 20 µL final

volume. Due to a large copy number difference between both genes and ddPCR

sensitivity, we run the reactions for each marker in separate reaction wells. In

consequence, each sample were normalized with a final quantity of 5 pg for ITS

samples and 5 ng for β-tubulin samples. All reactions were run with a cycling

program starting with 3 min at 95°C followed by 40 cycles of 10 s at 94°C alternating

with 60 s at 60°C. For each strain, the number of ITS1/5.8S copies was estimated as

the ratio of β-tubulin to ITS copy number per ng.

<u>Spatial structure of *Plasmopara viticola* primary inoculum in a vineyard</u>

The spatial structure of *P. viticola* primary inoculum in the experimental vineyard was

investigated at both field-scale and plot-scale by analyzing the regular grid data and

the nested sampling plots data, respectively, using a geostatistical approach.

Hereafter, we describe the field-scale analysis only, but the same methodology

(except model fitting and kriging) was applied for the plot-scale analysis (see Table

S1 in Supplementary Materials for details).

Plasmopara viticola DNA in soil samples, expressed as the number of ITS copies per

µl of reaction mix, was log-transformed before analysis to reduce the skewness of

the distribution and limit variance inflation and variogram distortion (Oliver and

Webster, 2014). First, all pairs of data (N=19503) were segregated into distance

classes. The maximal distance between two points in our dataset was 65.09 m. In

our analysis, however, we considered a maximal separation distance of 60 m to

exclude highest distance classes with a number of point pairs below the empirical

threshold for semivariance estimates accuracy (at least 30 to 50 pairs required;

Chiles and Delfiner, 1999; Journel and Huijbregts, 1976). We considered a spatial

lag of 2 m, which led to a total of 29 distance classes, with 79 to 1498 pairs each.

The sampling points in the first distance class were separated by a median distance

of 3.02 m. We then computed the semivariance for each distance class using the

Matheron estimator (Matheron, 1963), defined as follows:

$$\gamma(h) = \frac{1}{2n(h)}\sum_{i=1}^{n(h)}[z(x_i) - z(x_{i+h})]^2,$$

where $z(xi)$ is the primary inoculum at the first point of the ith pair, $z(xi + h)$ is the

primary inoculum at the second point of the ith pair, h is the median of the class of

distance considered and n is the number of point pairs in the distance class

considered. We represented the omnidirectional experimental semivariogram of the

primary inoculum of *P. viticola* in the vineyard by plotting the semivariance against

the sampling point separation distance h. A semivariogram shows how the similarity

between samples changes with increasing separating distance *i.e.* spatial

dependence. It can take various forms, and semi-variance often increases with

distance until it reaches a plateau at which point it remains stable. The variogram is

flat in the absence of spatial structure.

A visual inspection of the experimental semivariogram allows one to determine its

general shape. We fitted a spherical model (linear-sill shape) to the experimental

semivariogram using ordinary least square residuals method (Cressie, 1985). The

spherical model is as follows:

$$\gamma(h) = C_0 + C\left(\frac{3h}{2a} - \frac{1}{2}\left(\frac{h}{a}\right)^3\right) \quad 0 < h \leq a$$

$$\gamma(h) = C_0 + C \quad h > a$$

The model parameter estimates were regarded as quantitative descriptors of the

spatial pattern of the data. They included i) the spatial range a, or the distance

beyond which variance is constant; ii) the nugget variance C_0, an estimate of error

measurement plus error due to uncontrolled variability at distance lags shorter than

the shortest lag considered; iii) the structural variance C; and iv) the sill $C_0 + C$, or

sample variance.

We used the shape and parameter estimates of our semivariogram model to

interpolate *P. viticola* primary inoculum values at unsampled locations over a 0.05 m

× 0.05 m grid by ordinary kriging, which is an optimal interpolation procedure

described in detail in various sources (*e.g.*, Goovaerts, 1998; Oliver and Webster,

2015). After back-transforming the resultant values, we produced a contour map of

the spatial distribution of the primary inoculum in the field, as interpolated from our

data (*i.e.*, given the local conditions and our sampling plan).

All geostatistical analyses were performed using the package gstat (Pebesma, 2004)

in the R statistical software (v4.3.1; R Core Team, 2023). The contour map was

designed using the ggplot2 package (Wickham, 2016).

Plasmopara viticola primary inoculum along depth and distance-to-row

gradients

In the nested sampling plots, the sampling points were located either below the vine

stocks in the row (10 samples per plot) or in the inter-row space at 20, 60 or 80 cm

from the nearest row (respectively 4, 4 and 10 samples per plot). We used this data

to test whether the amount of primary inoculum in the soil varied along a gradient of

distance to the row. We fitted a linear mixed model with the log-transformed number

of ITS copies per µl as a response variable, the distance to the row as a fixed effect

and the plot as a random intercept effect (to account for the dependence of the data

within one plot). Model parameters were estimated by restricted likelihood estimation

and the significance ($\alpha = 0.05$) of the regression coefficients was tested with Student

t tests and Satterthwaite's approximation for degrees of freedom, using the lmer

function of the lmerTest package (Kuznetsova *et al.*, 2017). We evaluated model fit

by calculating the percentage of variance explained by fixed (R^2m) and by fixed plus

random effects (R^2c) (Nakagawa and Schielzeth, 2013). Finally, we tested whether

the amount of primary inoculum varied with the depth of sampling using a Kruskal-

wallis test and a Dunn test of multiple comparison. Statistical analyses were

conducted using the R statistical software (v4.3.1; R Core Team, 2023).

A bioassay to assess soil infectious potential

The principle of the bioassay is to place soil samples in optimal temperature and humidity conditions for oospore germination, to trap the released zoospores using susceptible grapevine leaf discs, and finally to approximate the infectious potential by evaluating the extent to which leaf discs are infected. The method described below was adapted from previous works (Hill, 1998; Pertot and Zulini, 2003; Rossi *et al.*, 2008b).

We performed the bioassay on one eighth of our 318 soil samples (n=40) using a 1 g portion per sample. Each sample was placed in the bottom of a 10 cm diameter circular plastic box half-filled with osmosed water and incubated at 20°C in the dark to stimulate oospore germination (day 0). The sample was held at the bottom of the box using a piece of blotting cloth (80µm mesh). We collected leaves from untreated *Vitis vinifera* cv. Cabernet Sauvignon grafted plants (SO4 rootstock) grown in greenhouses. We specifically picked the third or fourth leaves from the apex, as leaves at this stage of development are particularly susceptible to downy mildew infection (Isabelle Demeaux, personal communication). We cut 10 mm diameter discs in the leaf blade with a manual round cutter. On the first day of the incubation (day 0), we placed 10 discs from 10 different leaves (to avoid biases due to differences in downy mildew susceptibility between leaves) on the surface of the water (abaxial face down) in each box. The leaf discs were used to trap the zoospores released into the water following germination of oospores present in the soil sample, *i.e.* the zoospores swam to the leaf discs and infected them via the stomata. The blotting cloth allowed the zoospores to pass through but prevented the discs coming into direct contact with the soil and limited contamination of the discs

by saprophytic microorganisms. A separation structure was used to prevent the discs

in a box from overlapping and maintain the same trapping surface in each box (see

Figure S2 in Supplementary Materials). After 48 h, we placed the 10 discs abaxial

face up on a moist Whatman® filter paper sheet in a transparent box sealed with

plastic film (to maintain high humidity) and incubated at 20°C with a 12h-12h day-

night light cycle for 6 days. After 6 days of incubation, we visually inspected the discs

and evaluated the percentage of the area covered with *P. viticola* sporangiophores

for each disc. The germination of a cohort of oospores under optimal conditions

(immersed in water, in the dark and at 20°C) extend over 6 to 8 days, with a peak

around the second and third days of incubation (Isabelle Demeaux, personal

communication). To account for germination dynamics, we repeated zoospore

trapping using leaf discs on day 2, 4 and 6 from incubation start (day 0). We used

the cumulative number of infected discs and the mean percentage of leaf area

infected as indicators of the infectious potential of the soil.

We assessed the relationship between the abundance of primary inoculum in the soil

(as estimated with the ddPCR approach) and soil infectious potential by fitting one

generalized linear model per indicator (response variable following Poisson

distribution in both cases) with the number of *P. viticola* ITS copies per µl as a

predictor. In both cases, model parameters were estimated by maximum likelihood

and the significance ($\alpha = 0.05$) of the regression coefficients was tested with z tests.

We computed the McFadden's pseudo R^2 of the two models to determine the

indicator best fitting the primary inoculum data. Statistical analyses were conducted

using the R statistical software (v4.3.1; R Core Team, 2023).

Results

Quantification of *Plasmopara viticola* primary inoculum in vineyard soil samples

Considering the regular grid and nested sampling plot data, we found *P. viticola* DNA

in all samples except one, with 6.62 ± 6.68 ITS copies per µl on average. The value

for positive samples varied from 0.05 to 40.4 ITS copies per µl. A map of the raw

data is available in Supplementary materials (Figure S3).

By comparing the ITS1/5.8S to the β-tubulin signal obtained by ddPCR for seven *P.*

viticola strains, we found that the genome of *P. viticola* contains on average 215 ± 37

copies of the ITS sequence, but varied from 175 to 281 copies depending on the

strain (see Table S2 in Supplementary Materials for details). Using the average

estimate for the number of ITS1 copies, we estimated that the number of oospores

per gram of soil ranged from 0 to 1858 and that there was on average 303 ± 308

oospores per gram of soil in the experimental vineyard (see Figure S4 in

Supplementary materials for computation details). Using the minimal and maximal

estimates for the number of ITS1 copy, the average number of oospores per gram of

soil was 372 ± 378 or 232 ± 235, respectively.

Spatial structure of *Plasmopara viticola* primary inoculum in a vineyard

After log-transformation, the skewness of the distribution of *P. viticola* DNA data was

reduced from 1.69 to -0.35 for the regular grid, and from 2.40 to 0.42 for the nested

sampling plots. The shape of the fitted spherical model (regular grid data) revealed

that the downy mildew primary inoculum was spatially structured at the field scale

(Figure 2). The range was 25.8m. The nugget variance, the structural variance and

the sill were 0.36, 0.27 and 0.61 respectively. The high nugget value indicated that

very close points could be quite dissimilar in terms of primary inoculum, suggesting

the existence of a source of variation operating at a scale finer than that of the

regular grid. The map of the values interpolated by kriging using the fitted model is

shown in Figure 2.

The pattern of the plot-scale experimental semivariogram was not typical, with

semivariance alternatively increasing and decreasing along the gradient of

separation distance (Figure S5 in Supplementary Materials). Semivariance peaks

corresponded to distance classes with a higher proportion of heterogeneous pairs of

points, *i.e.* pairs consisting of one sampling point in a row and one sampling point in

an inter-row. On the other hand, lowest semivariance values corresponded to

distance classes with a higher proportion of homogeneous pairs of points, *i.e.* pairs

with two sampling points in a row or pairs with two sampling points in an inter-row

(Figure S5 in Supplementary Materials). We consistently found that semivariance

was higher for heterogeneous pairs of points than for homogeneous ones, on

average (anova P <0.0001, Figure S5 and Table S3 in Supplementary Materials).

Figure 2. Empirical semivariogram of downy mildew primary inoculum in the soil of the experimental vineyard (bottom left panel; open circles size is proportional to the number of data pairs; solid line corresponds to the fitted spherical model). Map of the experimental vineyard with interpolated values of downy mildew primary inoculum (right panel), expressed as the number of ITS copies per μl.

<u>*Plasmopara viticola* primary inoculum along depth and distance-to-row</u>

<u>gradients</u>

Our results clearly indicated that there was a decreasing gradient of primary

inoculum from the row to the middle of the inter-row (Figure 3; β = -0.0263 ± 0.0023;

df = 133; t value = -11.54; P < 0.001; R²m = 0.48; R²c = 0.50). Downy mildew

primary inoculum was on average five to six times more concentrated in the ridge of

soil below the vine stocks (9.32 ± 6.32 ITS copies per μl) than in the inter-row space

(1.72 ± 1.66 ITS copies per μl). These results were in line with the results from the

geostatistical analysis of nested sampling plot data (see above). We also found that

the primary inoculum concentration decreased with soil depth (Figure 3; Kruskal-

Wallis test's statistic = 12.1; df = 3; P < 0.01). In particular, it was significantly lower

between 21 and 30 cm and between 31 and 40 cm than in the first 10 cm from the

soil surface (see results of Dunn's post-hoc test in Table S4 in Supplementary

Materials).

Although we were not seeking to address this point, we also found that primary

inoculum concentration in the soil varied between rows, with significantly less *P.*

viticola DNA on average in the westernmost row compared to others in the vineyard

(Figure S6 Supplementary Materials; $F = 6.522$; $P < 0.001$; see Table S5 in

Supplementary Materials for results of Tukey's multiple comparison test). This

unexpected result was consistent with the distribution of primary inoculum as

predicted by kriging of observed data (Figure 2).

454
455   **Figure 3.** Abundance of downy mildew primary
456   inoculum (as number of ITS copies per µl estimated by
457   ddPCR) along gradients of distance to the row (upper

458   panel) and soil depth (lower panel). Solid line in the
459   upper panel corresponds to model predictions. Different
460   letters between depths in the lower panel indicate a
461   significant difference.

<u>A bioassay to assess soil infectious potential</u>

The cumulated number of leaf discs infected by *Plasmopara viticola* from 1g of soil

sample over the duration of the assay varied from 0 to 12 discs (out of 40). The

average proportion of disc area covered with *P. viticola* sporulation varied from 0 to

10.3%. The cumulated number of infected discs significantly increased with the

number of ITS copies per µl (Figure 4; $\square$ = 0.029 ± 0.008; P < 0.0001; McFadden's

pseudo R² = 0.05). In addition, the average infected area significantly increased with

the number of ITS copies per µl (Figure 4; $\square$ = 0.06 ± 0.01; P < 0.0001; McFadden's

pseudo R² = 0.16).

Figure 4. Comparison of bioassay and digital droplet PCR. Relationships between the abundance of DM primary inoculum in the soil (measured as number of ITS copies per µl by ddPCR) and the infectious potential of the soil measured by a bioassay, as either the cumulated number of infected discs (upper panel) or the average trap disc infected area (lower panel). Solid lines represent fitted model predictions.

Discussion

We present a direct, sensitive and specific method for the quantification of *P. viticola*

oospores in vineyard soil samples based on a ddPCR approach. We used this

method to describe the spatial distribution of downy mildew primary inoculum

(oospores) at different spatial scales in one vineyard. We also evaluated whether the

ddPCR-based assessment of inoculum concentration was related to an indicator with

an epidemiological significance, *i.e.* the soil infectious potential as assessed using a

bioassay. The method presented here could be used to survey primary inoculum in

vineyards and obtain quantitative data on soil oospore content, which would be

useful to improve models for epidemic risk prediction.

The method used for inoculum assessment in soil is based on the ddPCR technique,

which provides an absolute quantification of DNA and is particularly well suited for

rare targets in complex matrices. We combined these advantages to the use of an

ITS sequence occurring in numerous copies which increases the probability of

detection of *P. viticola* DNA in small quantities in the sample. The partition of the

samples into thousands of droplets improves the yield of amplification and the

sensitivity of the analysis as compared to qPCR, including in the case of oomycete

quantification in soil samples (Gossen *et al.*, 2019; Rački *et al.*, 2014; Wen *et al.*,

2020). Using ddPCR and the average estimate for the number of ITS1 copy in the

genome of *P. viticola* (215 copies), we found that the number of oospores per gram

of soil ranged from 0 to 1858 in the experimental vineyard and averaged 303

oospores per gram of soil. This result falls within the range of values (*i.e.* from a few

dozen to 4000 oospores per gram of soil approximately) previously estimated by

Yang *et al.* (2023) in Chinese vineyards using qPCR.

A potential drawback of the oospore estimate from a multicopy DNA sequence lies in

the accuracy of the assessment of the number of copy of the sequence in the

genome of the target species. Variability in ITS copy numbers among isolates from

the same oomycete species has been poorly studied but can potentially vary among

strains (McLeod *et al.* 2024). In our case, we found that the number of ITS1/5.8S

copies in *P. viticola* genome varied from 175 to 281 copies depending on the strain

(n=7), which caused the primary inoculum concentration estimates to vary from 232

to 372 oospores per gram of soil on average. However, the number of ITS copies

found in our study is consistent with what was found for other oomycetes (95 copies

per haploid genome on average; Gangneux *et al.*, 2014). Further analyses should be

undertaken to better assess the occurrence of the ITS1/5.8S on a population level.

Another possible drawback for the quantification of oospores may arise from

suboptimal DNA extraction and amplification efficiency, which would lead to

underestimation of oospore concentration. This bias need to be further addressed for

a more accurate estimate of oospores concentration in future studies. However, DNA

was extracted from 2g of soil in our protocol, which is eight times greater than with

extraction kits (*e.g.* 250 mg for the DNeasy PowerSoil Pro kit from QIAGEN).

Together with the sample homogenization step, we believe that this may have two

important advantages: i) limit estimation bias due to small-scale spatial variability in

oospore abundance and improve the repeatability of the method (which we

assessed; see Supplementary Materials) and ii) allows increasing the sensitivity of

the method because higher amount of oospores is present initially in the sample.

In our study, *P. viticola* was encountered at all sampled locations in the field contrary

to soilborne pathogens whose presence is generally restricted to some areas

(Botero-Ramirez *et al.*, 2021; Moussart *et al.*, 2009). This difference may be

explained by the greater dispersal ability of the airborne form of *P. viticola* (as

compared to strictly soilborne pathogens) resulting in a more even distribution of the

disease in the field. Still, we showed that the distribution of downy mildew oospores

was not random or regular on a field scale, but that it was characterized by 25 m-

diameter patches of concentrically increasing oospore concentration. This pattern

could be related to the distribution of the disease in the vineyard at the end of the

growing season, which we unfortunately did not assess the year before our soil

survey (but see Supplementary materials for comparisons of the spatial distribution

of primary inoculum against soil electrical resistivity and grapevine NDVI

distributions, as indirect proxies for disease incidence). Alternatively, environmental

factors could affect grapevine leaf litter (containing the oospores) distribution or the

survival of the inoculum during autumn and winter. Future studies should investigate

the relationship between the spatial distribution of the symptoms at the end of the

growing season and the distribution of the inoculum in the soil the following year.

We found that, on average, oospores accumulates preferentially in the ridge of soil

below the vine stocks (or "row location") at a rate of 4 to 5 times the inter-row space,

and in the first 10 cm of soil. Although oospore concentration varied on a wider

spatial scale within the field (25m patches), this row *vs* inter-row contrast was

observed at all locations in the field. Moreover, oospore concentration was more

similar at two row locations, whatever the distance between the two points, than

when comparing row *vs* inter-row locations. This suggests that the distribution of

oospores is primarily determined by the distribution of grapevine leaf litter that carry

them, which first falls and accumulates at the foot of the grapevines. However, the

distribution of grapevine leaf litter may change during autumn and winter under the

influence of the wind and of cultural practices. For instance, tillage was found to

modify the spatio-temporal distribution of crop residues in annual cropping systems,

and in particular the distribution between ridges and furrows (Grigera *et al.*, 2007;

Guérif *et al.*, 2001; Schneider *et al.*, 2006). Here, we hypothesize that soil ridging at

the foot of the vines contributes to "trap" the fallen leaves and could prevent their

dispersion by the wind (see Figure S7 in Supplementary Materials for an illustration),

resulting in a higher concentration of inoculum locally. Future research should study

the dispersion of grapevine leaf litter (or crop residues) during winter and how it is

affected by cultural practices to better understand oospores distribution in vineyards.

This could help identify cultural practices that prevent inoculum from accumulating in

the vineyard below the grapevines, where it is more likely to reach the host tissue in

the event of rain splashes. Finally, data on oospore distribution within a vineyard

could be useful to determine an optimal sampling strategy (number and distribution

of soil sampling points within a field) for the design of operational, cost-effective field-

scale indicators of primary inoculum reservoir. This could be done using methods

such as hierarchical variance analysis based on mixed models with nested random

effects (Webster *et al.*, 2006) for example. Nevertheless, data from various vineyards

should be gathered for such an indicator to be as generic as possible.

Our study also revealed that soil oospore concentration can vary between rows

within a field and be subjected to border effects. Indeed, we found that the primary

inoculum was 7 to 8 times less concentrated in the westernmost row compared to

the rest. This difference might arise from various mechanisms mediated by the

presence of a tree hedgerow alongside the experimental field, possibly limiting

disease incidence of leaf litter abundance locally. These mechanisms include i) wind

erosion and microclimate modification in the vicinity of the hedgerow (Campi *et al.*,

2009; Jacobs *et al.*, 2022), ii) modulation of grapevine vigor and susceptibility to

downy mildew potentially due to competition for resources (Favor and Udawatta,

2021; but see Bourgade *et al.*, 2020), and iii) interception of airborne spores

(Delatouche *et al.*, 2024). In our case, the absence of a border effect on the east

side of the field (also bordered by a hedgerow) suggests that mechanisms at play

operate asymmetrically, which restricts the choice to mechanisms associated to wind

erosion and microclimate modification. In any case, these results raise questions

regarding the effects of agroecological infrastructures on downy mildew

epidemiology, as well as their possible contribution to preventive disease

management strategies.

We further demonstrated that the concentration of *P. viticola* oospores in the soil is

related to soil infectious potential as estimated in controlled conditions using a leaf

disc bioassay. In particular, we found that primary inoculum abundance is the soil

was positively related to the proportion of leaf discs infected during the bioassay.

Positive relationships between inoculum abundance and disease severity were found

for soilborne root pathogens (Chatterton *et al.*, 2023; Hwang *et al.*, 2011; Moussart

et al., 2009). However, we found no study that addressed the inoculum density-

disease severity relationship for soil-to-aerial plant parts infection pathway, such as

in the case of downy mildew primary contaminations. Contrary to DNA-based

methods, the bioassay discriminate between viable and unviable cells, as only viable

oospores will be able to cause leaf disc infections. However, considering that the

oospore germination period in a population extends over several months (Maddalena

et al., 2021), our bioassay (carried out on a single date) "captured" only part of the

oospore population present in the samples, which may explain that the relationship

was not strong (McFadden's R^2 = 0.16). Future works should establish whether this

method for soil infectious potential assessment is a good proxy for the "true"

infectious potential, *i.e.* accounting for the entire oospore population. Nevertheless,

our result suggests that DNA-based quantification of primary inoculum have an

epidemiological significance and could be used in models to improve epidemic risk

predictions.

As agriculture moves away from chemical biocides, the design of sustainable,

adaptative and cost-effective crop disease management strategies will rely more and

more on the prevention of epidemic risks and a predictive evaluation of pathogen

population dynamics. In this context, the quantitative and spatially-explicit survey of

inoculum reservoirs is crucial. The insights gained into downy mildew primary

inoculum from our methods could contribute to the design of monitoring and

decision-making tools to accompany the implementation of sanitation measures

preventing primary inoculum accumulation in the vineyard from one season to the

next.

Acknowledgements

The authors thank the Experimental Viticultural Unit of Bordeaux 1442, INRAE, F-

33883 Villenave d'Ornon, for the maintenance of the experimental vineyard that was

used in this study, and the GenoSol platform (DOI 10.15454/L7QN45) for sharing its

expertise and helping with methodological development for DNA extraction from soil

samples. Part of the experiments (ddPCR analysis) were performed at the Genome

Transcriptome Platform of Bordeaux (PGTB;

doi:10.15454/1.5572396583599417E12) with the help of Adline Delcamp and Emilie

Chancerel. Finally, the authors are grateful to the sponsors of the Chaire Alexis

Millardet for their support to research activities, namely Château Ausone, Château

Cheval Blanc, Petrus and Château Montrose.

Fundings

This work was initiated thanks to the Ministries for an Ecological Transition, for

Agriculture and Food, for Solidarity and Health and of Higher Education, Research

and Innovation, with the financial support of the French Office for Biodiversity, as

part of the call for research projects "Overall approaches to limit the use of

phytosanitary products : coupling preventive and curative solutions within agricultural

sectors from farmers to consumers", with the fees for diffuse pollution coming from

the Ecophyto II+ plan (PROFIL project 2020-2024). This work was also financially

supported by the ISVV Fund of the Fondation Bordeaux Université, through the

Chaire Alexis Millardet program. The authors acknowledge the financial support of

the French National Research Agency (ANR) under the grant 20‑PCPA‑0010 (Vitae

project from the PPR Cultiver et Protéger Autrement). The GenoSol platform (DOI

10.15454/L7QN45) of the infrastructure ANAEE-Services received a grant from the

French state through the National Agency for Research under the program

"Investments for the Future" (reference ANR-11-INBS-0001), as well as a grant from

the Regional Council of Bourgogne Franche-Comté. The BRC GenoSol is a part of

BRC4Env (10.15454/TRBJTB), the pillar « Environmental Resources » of the

Research Infrastructure AgroBRC-RARe 10.15454/b4ec-tf49.

Literature Cited

Agreste (2023). Enquête pratiques culturales en viticulture en 2019, IFT et nombre

de traitements, données révisées. Chiffres et Données. 4:1-12.

Beaumelle, L., Tison, L., Eisenhauer, N., Hines, J., Malladi, S., Pelosi, C., et al.

(2023). Pesticide effects on soil fauna communities—A meta‑analysis. Journal of

Applied Ecology. 60:1239–1253.

Beketov, M. A., Kefford, B. J., Schäfer, R. B., & Liess, M. (2013). Pesticides reduce

regional biodiversity of stream invertebrates. Proceedings of the National Academy

of Science U.S.A. 110:11039–11043.

Botero-Ramirez, A., Hwang, S.-F., & Strelkov, S. E. (2021). *Plasmodiophora

brassicae* inoculum density and spatial patterns at the field level and relation to soil

characteristics. Pathogens. 10:499.

Bourgade, E., Alonso Ugaglia, A., Bustillo, V., Dufourcq, T., Grimaldi, J., Guenser, J.,

et al. (2020). VITIFOREST☐: Evaluation de l'impact de l'arbre agroforestier en

contexte viticole. Innovations Agronomiques. 79:471–497.

Brischetto, C., Bove, F., Languasco, L., & Rossi, V. (2020). Can spore sampler data

be used to predict *Plasmopara viticola* infection in vineyards? Frontiers in Plant

Science. 11:1187.

Butault, J.-P., Delame, N., Jacquet, F., & Zardet, G. (2011). L'utilisation des

pesticides en France : état des lieux et perspectives de réduction. Notes et Études

Socio-économiques. 35:7-26.

Caffi, T., Rossi, V., & Lusitani, M. (2011). Long-term survival of *Plasmopara viticola*

oospores. Integrated protection and production in viticulture. IOBC WPRS Bulletin.

67:111–114.

Campi, P., Palumbo, A. D., & Mastrorilli, M. (2009). Effects of tree windbreak on

microclimate and wheat productivity in a Mediterranean environment. European

Journal of Agronomy. 30:220–227.

Carisse, O. (2016). Development of grape downy mildew (*Plasmopara viticola*) under

northern viticulture conditions: influence of fall disease incidence. European Journal

of Plant Pathology. 144:773–783.

Carisse, O., Van der Heyden, H., Tremblay, D. M., Hébert, P. O. & Delmotte F.

(2021). Evidence for differences in the temporal progress of *Plasmopara viticola*

clades *riparia* and *aestivalis* airborne inoculum monitored in vineyards in eastern

Canada using a specific multiplex quantitative PCR assay. Plant Disease 105:1666-

1676.

Chatterton, S., Schwinghamer, T. D., Pagé, A., Davidson, R. B., Harding, M. W., &

Banniza, S. (2023). Inoculum dose–disease response relationships for the pea root

rot pathogen, *Aphanomyces euteiches*, are dependent on soil type and other

pathogens. Frontiers in Plant Science. 14:1115420.

Chiles, J.-P., & Delfiner, P. (1999). Geostatistics: modeling spatial uncertainty. New

York: John Wiley & Sons.

Dall'Agnol, J. C., Pezzini, M. F., Uribe, N. S., & Joveleviths, D. (2021). Systemic

effects of the pesticide mancozeb – A literature review. European Review for Medical

and Pharmacological Sciences. 25:4113–4120.

Delatouche, L., Tixier, P., Sainte-Rose, J., Daribo, M.-O., & de Lapeyre de Bellaire,

L. 2024. How do hedgerow characteristics alter the dispersal of Pseudocercospora

fijiensis propagules? Pest Management Science, 80:1454-1464.

Douillet, A., Laurent, B., Beslay, J., Massot, M., Raynal, M., & Delmotte, F. (2022).

LAMP for in-field quantitative assessments of airborne grapevine downy mildew

inoculum. Journal of Applied Microbiology. 133:3404–3412.

Douillet, A. (2023). Étude pluriannuelle de la sporée aérienne de *Plasmopara viticola*

par quantification moléculaire : un nouvel indicateur de risque des épidémies du

mildiou de la vigne. PhD thesis.

Dubuis, P. H., Viret, O., Bloesch, B., Fabre, A. L., Naef, A., Bleyer, G., et al. (2012).

Lutte contre le mildiou de la vigne avec le modèle VitiMeteo-Plasmopara. Revue

Suisse de Viticulture Arboriculture Horticulture. 44 (3), 192-198.

Favor, K., & Udawatta, R. P. (2021). Belowground services in vineyard agroforestry

systems. In *Agroforestry and ecosystem services*, eds. Ranjith P. Udawatta and

Jose Shibu. Springer International Publishing, p. 65–94.

Fontaine, M. C., Austerlitz, F., Giraud, T., Labbé, F., Papura, D., Richart-Cervera, S.,

et al. (2013). Genetic signature of a range expansion and leap-frog event after the

recent invasion of Europe by the grapevine downy mildew pathogen *Plasmopara*

viticola. Molecular Ecology. 22:2771–2786.

Fontaine, M. C., Labbé, F., Dussert, Y., Delière, L., Richart-Cervera, S., Giraud, T.,

et al. (2021). Europe as a bridgehead in the worldwide invasion history of grapevine

downy mildew, *Plasmopara viticola*. Current Biology. 31:2155-2166.e4.

Gangneux, C., Cannesan, M.-A., Bressan, M., Castel, L., Moussart, A., Vicré-

Gibouin, M., et al. (2014). A Sensitive Assay for Rapid Detection and Quantification

of *Aphanomyces euteiches* in Soil. Phytopathology. 104:1138–1147.

Gessler, C., Pertot, I., & Perazzolli, M. (2011). *Plasmopara viticola*: a review of

knowledge on downy mildew of grapevine and effective disease management.

Phytopathologia Mediterranea. 50:3–44.

Gobbin, D., Pertot, I., & Gessler, C. (2003). Identification of microsatellite markers for

Plasmopara viticola and establishment of high throughput method for SSR analysis.

European Journal of Plant Pathology. 109:153–164.

Goovaerts, P. (1998). Geostatistical tools for characterizing the spatial variability of

microbiological and physico-chemical soil properties. Biology and Fertility of Soils.

27:315–334.

Gossen, B. D., Al-Daoud, F., Dumonceaux, T., Dalton, J. A., Peng, G., Pageau, D.,

et al. (2019). Comparison of techniques for estimation of resting spores of

Plasmodiophora brassicae in soil. Plant Pathology. 68:954–961.

Grigera, M., Drijber, R., & Wienhold, B. (2007). Redistribution of crop residues during

row cultivation creates a biologically enhanced environment for soil microorganisms.

Soil and Tillage Research. 94:550–554.

Guérif, J., Richard, G., Dürr, C., Machet, J. M., Recous, S., & Roger-Estrade, J.

(2001). A review of tillage effects on crop residue management, seedbed conditions

and seedling establishment. Soil and Tillage Research. 61:13–32.

Hill, G. K. (1998). Studies on the germination of *Plasmopara viticola* oospores with a

floating disc test. IOBC WPRS Bulletin, 21.

Hong, C., Brewer, M. T., Brannen, P. M., & Scherm, H. (2020). Temporal disease

dynamics and relative importance of sexual and asexual reproduction of grape

downy mildew (*Plasmopara viticola*) in an isolated vineyard in the North Georgia

Mountains, USA. Plant Pathology. 69:1721–1730.

Hwang, S. F., Ahmed, H. U., Strelkov, S. E., Gossen, B. D., Turnbull, G. D., Peng,

G., et al. (2011). Seedling age and inoculum density affect clubroot severity and

seed yield in canola. Canadian Journal of Plant Science. 91:183–190.

Jacobs, S. R., Webber, H., Niether, W., Grahmann, K., Lüttschwager, D., Schwartz,

C., et al. (2022). Modification of the microclimate and water balance through the

integration of trees into temperate cropping systems. Agricultural and Forest

Meteorology. 323:109065.

Journel, A. G., and Huijbregts, C. J. (1976). Mining geostatistics. London: Academic

Press.

Kennelly, M. M., Gadoury, D. M., Wilcox, W. F., Magarey, P. A., & Seem, R. C.

(2007). Primary infection, lesion productivity, and survival of sporangia in the

grapevine downy mildew pathogen *Plasmopara viticola*. Phytopathology. 97:512–

522.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package:

Tests in Linear Mixed Effects Models. Journal of Statistical Software, 82(13):1–26.

Lamichhane, J. R., Messéan, A., & Ricci, P. (2019). Chapter Two - Research and

innovation priorities as defined by the Ecophyto plan to address current crop

protection transformation challenges in France. In Advances in Agronomy, ed.

Donald L. Sparks. Academic Press, p. 81–152.

Maddalena, G., Russo, G., & Toffolatti, S. L. (2021). The study of the germination

dynamics of *Plasmopara viticola* oospores highlights the presence of phenotypic

synchrony with the host. Frontiers in Microbiology. 12:698586.

Matheron, G. (1963). Principles of geostatistics. Economic geology. 58:1246–1266.

McLeod, A., Hamza, Z., Schoeman, M. H., Moyo, L., & Akinsanmi, O. A. (2024).

Temporal colonization of macadamia tree roots by *Phytophthora cinnamomi* based

on a newly developed quantitative real-time PCR assay. Plant Pathology. 73(2), 366-

377.

Möller, E. M., Bahnweg, G., Sandermann, H., & Geiger, H. H. (1992). A simple and

efficient protocol for isolation of high molecular weight DNA from filamentous fungi,

fruit bodies, and infected plant tissues. Nucleic acids research. 20(22), 6115.

Moussart, A., Wicker, E., Le Delliou, B., Abelard, J.-M., Esnault, R., Lemarchand, E.,

et al. (2009). Spatial distribution of *Aphanomyces euteiches* inoculum in a naturally

infested pea field. European Journal of Plant Pathology. 123:153–158.

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining

R2 from generalized linear mixed-effects models. Methods in Ecology and Evolution.

4(2), 133–142.

Oliver, M. A., & Webster, R. (2014). A tutorial guide to geostatistics: Computing and

modelling variograms and kriging. CATENA. 113:56–69.

Oliver, M. A., & Webster, R. (2015). Basic steps in geostatistics: the variogram and

kriging. Cham: Springer International Publishing.

Park, E. W., Seem, R. C., Gadoury, D. M., & Pearson, R. C. (1997). DMCAST: a

prediction model for grape downy mildew development. Wein-Wissenschaft. 52(3-4),

182-189.

Pebesma, E. J. (2004). Multivariable geostatistics in S: the gstat package.

Computers & Geosciences. 30:683–691.

Pertot, I., & Zulini, L. (2003). Studies on *Plasmopara viticola* oospore germination in

Trentino, Italy. IOBC WPRS Bulletin. 26:43–46.

R Core Team (2023). R: A language and environment for statistical computing.

Available at: https://www.R-project.org/.

Rački, N., Dreo, T., Gutierrez-Aguirre, I., Blejec, A., & Ravnikar, M. (2014). Reverse

transcriptase droplet digital PCR shows high resilience to PCR inhibitors from plant,

soil and water samples. Plant Methods. 10:42.

Rossi, V., Caffi, T., Giosuè, S., & Bugiani, R. (2008a). A mechanistic model

simulating primary infections of downy mildew in grapevine. Ecological modelling.

212(3-4), 480-491.

Rossi, V., Caffi, T., Bugiani, R., Spanna, F., & Della Valle, D. (2008b). Estimating the

germination dynamics of *Plasmopara viticola* oospores using hydro-thermal time.

Plant Pathology. 57, 216-226.

Rouxel, M., Mestre, P., Comont, G., Lehman, B. L., Schilder, A., & Delmotte, F.

2013. Phylogenetic and experimental evidence for host‐specialized cryptic species in

a biotrophic oomycete. New Phytologist. 197:251–263.

Schneider, O., Roger-Estrade, J., Aubertot, J.-N., & Doré, T. (2006). Effect of

seeders and tillage equipment on vertical distribution of oilseed rape stubble. Soil

and Tillage Research. 85:115–122.

Sharma, A., Kumar, V., Shahzad, B., Tanveer, M., Sidhu, G. P. S., Handa, N., et al.

(2019). Worldwide pesticide usage and its impacts on ecosystem. SN Applied Sciences.

1:1-16.

Si Ammour, M., Bove, F., Toffolatti, S. L., & Rossi, V. (2020). A Real-Time PCR

assay for the quantification of *Plasmopara viticola* oospores in grapevine leaves.

Frontiers in Plant Science. 11:1202.

Strizyk, S. (1983). Modèle d'état potentiel d'infection. Application à *Plasmopara*
viticola. ACTA.

Valsesia, G., Gobbin, D., Patocchi, A., Vecchione, A., Pertot, I., & Gessler, C.
(2005). Development of a high-throughput method for quantification of *Plasmopara*
viticola DNA in grapevine leaves by means of quantitative real-time polymerase
chain reaction. Phytopathology. 95:672–678.

Wan, N.-F., Fu, L., Dainese, M., Kiaer, L. P., Hu, Y.-Q., Xin, F., et al. (2025).
Pesticides have negative effects on non-target organisms. Nature Communications.
16:1360.

Webster, R., Welham, S. J., Potts, J. M., & Oliver, M. A. (2006). Estimating the
spatial scales of regionalized variables by nested sampling, hierarchical analysis of
variance and residual maximum likelihood. Computers & Geosciences. 32(9), 1320-
1333.

Weerakoon, N. D., Roberts, J. K., Lehnen, L. P., Wilkinson, J. M., Marshall, J. S., &
Hardham, A. R. (1998). Isolation and characterization of the single β-tubulin gene in
Phytophthora cinnamomi. Mycologia. 90:85–95.

Wen, R., Lee, J., Chu, M., Tonu, N., Dumonceaux, T., Gossen, B. D., et al. (2020).
Quantification of *Plasmodiophora brassicae* resting spores in soils using droplet
digital PCR (ddPCR). Plant Disease. 104:1188–1194.

Wickham, H. (2016). ggplot2: Elegant graphics for data analysis. New York:
Springer-Verlag.

Wong, F. P., Burr, H. N., & Wilcox, W. F. (2001). Heterothallism in *Plasmopara*

viticola. Plant Pathology. 50:427–432.

Yang, L., Chu, B., Deng, J., Yuan, K., Sun, Q., Jiang, C., & Ma, Z. (2023). Use of a

real-time PCR method to quantify the primary infection of *Plasmopara viticola* in

commercial vineyards. Phytopathology Research. 5(1), 19.