# Effect of curcumin and 5-bromoindole-3-carboxaldehyde on the inhibition of Quorum sensing-dependent virulence factors in *Pseudomonas aeruginosa*

Gabriel Martínez González[1], Rodolfo García Contreras[2], Juan Andres Alvarado-Salazar[3*], Israel García Valdés[4], Nilda García Valdés[4], Denisse Alejandra Lugo Gutiérrez[4], Ingrid Lizeth Bustamante Martínez[4], Alonso Ruben Tescucano Alonso[1], Alejandra Ramirez-Villalva[1,5], Jorge Angel Almeida Villegas[4,6*].

[1]Instituto de Investigación y Estudios en Salud (INIES), Universidad de Ixtlahuaca CUI, Ixtlahuaca, Mexico.
[2]Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad Nacional Autónoma de Mexico, Ciudad de Mexico, Mexico.
[3]Facultad de Estudios Superiores "Zaragoza", Universidad Nacional Autónoma de México, Ciudad de Mexico, Mexico.
[4]Licenciatura en Química Farmacéutica Biológica, Universidad de Ixtlahuaca CUI, Ixtlahuaca, Mexico.
[5]Facultad de Química, Universidad Autónoma del Estado de México, Toluca, Mexico.
[6]Escuela Profesional de Medicina, Universidad de Ixtlahuaca CUI, Ixtlahuaca, Mexico.

Corresponding author
angel.almeida@uicui.edu.mx (JAAV), andres.alvarado.salazar@comunidad.unam.mx (JAAS)

## Abstract

*Pseudomonas aerugoinosa* is a bacterium that causes important nosocomial diseases with high mortality and is classified within the bacteria of the acronym ESKAPE. The virulence of *Psuedomonas aeruginosa* is mediated by Quorum sensing, a cellular communication system that allows it to modulate the expression of virulence factors such as elastase, protease, siderophores, biofilm, alginate among others, through the LasI / LasR, RhlI / RhlR and PQS systems. Quorum inhibition allows an antivirulence strategy, where it has been shown that molecules such as curmunin and carboxaldehydes have considerable activity for this purpose. In the present work, the ability to inhibit Quorum Sensing-dependent virulence factors in *Pseudomonas aeruginosa* was evaluated using curcumin and 5-bromoindole-3-carboxaldehyde. The strains evaluated are Pa01, Pa14 and ATCC 27853 at concentrations (1 and 5 µg/ml) for curcumin and (30, 50 and 100 µg/ml) for 5-bromoindole-3-carboxaldehyde. 5-Bromoindole-3-carboxaldehyde significantly inhibited the production of pyocyanin in Pa01 (92.5% at 100 µg/ml), elastase in Pa01 (79.85% at 100 µg/ml), protease in Pa01 (92.8% at 50 µg/ml) and alginate in ATCC 27853 (76.16% at 100 µg/ml). Curcumin showed substantial inhibition of pyocyanin in ATCC 27853 (100% at 5 µg/ml), elastase in Pa14 (56.19% at 5 µg/ml), protease in Pa01 (87.04% at 5 µg/ml) and alginate in Pa01 (48.26% at 5 µg/ml). These findings suggest the potential of these compounds as antivirulence agents, to combat *P. aeruginosa* infections.

## Keywords

Quorum sensing, curcumin, antivirulence, bromoindole, *Pseudomonas aeruginosa*

**Introduction**

*Pseudomonas aeruginosa* is a Gram-negative aerobic rod-shaped bacterium that can be isolated from most environments, including soil, plants, and mammal tissue [1]. Has become an important cause of infection in humans and can be associated with significant morbidity and mortality [2]. Is a common cause of nosocomial infections, particularly pneumonia, infection in immunocompromised hosts, and in those with structural lung disease such as cystic fibrosis [3,4].

The World Health Organization (WHO) listed *P. aeruginosa* as one of three bacterial species in which there is a critical need for the development of new antibiotics to treat infections [5]. Importantly, *P. aeruginosa* is one of the MDR ESKAPE pathogens [6].

*P. aeruginosa* virulence regulation uses both chemical factors such as quorum sensing (QS) and mechanical cues such as surface association to modulate a large suite of virulence factors [7]. Essential function for QS in *P. aeruginosa* is to regulate the production of multiple virulence factors, such as extracellular proteases, iron chelators, efflux pump expression, biofilm development, swarming motility, phospholipases C, alginate, DNase and the response to host immunity [8,9].

Resistance of *Pseudomonas aeruginosa* to antibiotics is a major problem. Targeting virulence factors is an alternative option to avoid the emergence of resistance to antibiotics and treat infections that this pathogen causes [10]. QS system an attractive target for antibacterial therapy [11]. Inhibition of this system may cause the attenuation of virulence and protect against infection. In fact, an anti-QS approach has already shown promise in the battle against *P. aeruginosa* infections [12].

The use of QS inhibitors would be of particular interest in inhibiting bacterial pathogenicity and infections [13]. Turmeric (*Curcuma longa*) has traditionally been used as anti-inflammatory, antimicrobial and antifungal, antioxidant, anticancer, anti-inflammatory, antiobesity and antidepressant [14], but reports on its anti-QS properties are scarce [15]. Several studies reported that curcumin inhibits bacterial QS systems/biofilm formation and prevents bacterial adhesion to host receptors in various species, including *S. aureus, E. faecalis, E. coli, S. mutans, L. monocytogenes, H. pylori, P. aeruginosa, S. marcescens, A. hydrophila* and *A. baumannii* [16].

Another molecule whose reports indicate antibacterial activity are indoles, which, although they use bacteria as signalling, have shown important antibiofilm activity [17]. Functional alteration of indole carboxaldehydes may increase their effectiveness as quorum sensing inhibitors (QSIs). The bromination of carboxaldehydes represents an increase in their activity [18].

Given the background of the anti-quorum sensing activity of molecules such as curcumin and carboxaldehydes, the objective of this work is to evaluate in three strains of *Pseudomonas aeruginosa* (Pa01, Pa14 and ATCC 27853) the activity of curcumin and 5-bromoindole-3-carboxyaldehyde on quorum-sensing-dependent virulence factors, such as alginate, pyocyanin, elastase and protease.

The proposal of new antibacterial treatments often requires the use of computer software that allows the calculation of molecular properties and the prediction of bioactivity, with which the cheminformatics properties, previously obtained, can be correlated with the viability of the new treatment proposal. Chemoinformatic properties have advantages such as predicting the site of action, probable route of administration, physicochemical, pharmacokinetic, pharmacodynamic behaviour and possible toxic effects. The use of simulation techniques and online servers allow us to know the behaviour of a compound to propose an effective and safe therapy.

## Methods and materials

### Preparation of the strains and addition of evaluated molecules

To carry out the tests, 3 different strains of *P. aeruginosa* were used; Pa01, Pa14 and ATCC 27853, all belonging to the microbiology laboratory of the University of Ixtlahuaca CUI. To activate these strains, they were reseeded in boxes of LB medium with 1.5% agar for at least 20 hours at 37 °. Subsequently, ON (overnight) cultures of each strain were inoculated in 50 ml flasks with 25 ml of LB medium, cultivated for 18 hours at 37 °. After the elapsed time, the growth of the cultures was measured at OD of 600 nm in a 1:10 ratio (dilution in LB).

Subsequently, new cultures of the strains were inoculated from the ON cultures with an OD of 600 nm. The inoculums of each strain of interest were added to 3 series of flasks (each with 8 flasks with 25 ml of LB medium) since three different strains were evaluated, different concentrations of each molecule were added to each series: For curcumin (Sigma-Aldrich), they evaluated concentrations of 1 µg/ml and 5 µg/ml, 5-bromoindole-3-carboxaldehyde (Sigma-Aldrich) 30, 50 and 100 µg/ml, both molecules dissolved in DMSO previously filtered with acrodiscs 0.22 µm . They were incubated at 37 ° with constant shaking for 24 hours. To maintain control, a culture that did not have any of the added molecules was used. After 24 hours, growth was measured at 600 nm of both the controls and the flasks with the evaluated molecules. As the molecules were dissolved in DMSO, negative controls were done with this solvent.

### Determination of Pyocyanin

From the mother flasks with ON, 1000 µl of each culture were taken. They were subsequently centrifuged at 13,000 rpm for 1.5 minutes. 800 µl of the supernatant were taken. Adding 420 µl of chloroform, it was vortexed for 1 minute at speed 10 and centrifuged at 13,000 rpm for 5 minutes. We transferred 300 µl of the bottom phase together with 800 µl of 0.1 N HCl. It was shaken for 1 minute at vortex speed 10, and centrifuged at 13,000 rpm for 5 minutes. It was determined quantitatively by absorbance, measured in a quartz cell, at 520 nm [19].

### Protease Determination

1000 µl were taken from the stock flasks with ON. The samples were centrifuged at 13000 rpm for 2 minutes. 875 µl of buffer with protease reagent and 125 µl of culture supernatant were used. It was vortexed for 1 minute. Samples were incubated at 37 ° at 200 rpm for 1 h. At the end of the incubation time, aliquots were taken and placed in 1% $HNO_3$. Shaking

with the help of vortex for 1 minute at speed 10, we then centrifuged at 13,000 rpm for 2 minutes. The supernatant was placed in the presence of 0.5% NaOH. We stir lightly with the help of the vortex. Transferring into quartz cells, the samples were read at 595 nm [20,21].

## Determination of alginate

1000 µl were taken from the stock flasks. 1 ml of the cell culture from the cultures was centrifuged and subsequently heated at 80° for 30 min. The resulting supernatant was centrifuged for 30 min at 13,000 rpm, the pellet was removed, and the alginate was precipitated with ice-cold 99% ethanol. The precipitated alginate was collected and dissolved in 1 ml of 0.9% sterile saline. A mixture of 118 µl of sample plus 1 ml of $H_3BO_3/H_2SO_4$ and 34 µl of carbazole is made. This mixture was heated for 30 min at 55°C and the absorbance (OD 530) was measured [22].

## Determination of Elastase

1000 µl were taken from the mother flasks, then centrifuged for 1.5 minutes at 13000 rpm and the supernatant was separated. Subsequently, 100 µl of the supernatant from each of the cultures was placed and 5 mg of elastin-Congo red from Sigma- Aldrich and 900 µl of pH 7.5 buffer were added. They were incubated for 2 h at 37 °, centrifuged at 13,500 rpm for 5 minutes, and absorbance was measured at 495 nm [23].

## Molecular docking

The 2D structure of curcumin and 5-bromoindole-3-carboxaldehyde were drawn with ChemSketch ACD/Labs and protonation states were calculated at pH 7.4 with MarvinSketch Chemaxon. The 3D structure was obtained and optimized to a semi-empirical PM3 level with Spartan 08 V1.2.0. The crystals of the LasR, PQS and RhlR proteins were obtained from the Protein Data Bank repository with the codes 3IX3, 4JVI and 8DQ0, respectively. The preparation of the proteins was carried out in Molegro Virtual Docker, the water molecules were removed, the druggable cavities were detected and the ligand-binding site was selected for molecular docking studies with a search sphere of 10 Å radius, grid of 0.30 Å, with the coordinates X:  10.54, Y: 4.32, Z: 20.98 for LasR, X: -34.71, Y: 56.02, Z: 9.17 for PQS and X: 116.01, Y: 96.69, Z: 137.48 for RhlR [24-26].

The validation of the molecular docking method was carried out using 12 methods based on the combinations of 4 scoring functions (MolDock Score, MolDock Score GRID, Plant Score and Plant Score GRID) and 3 search algorithms (MolDock Optimizer, MolDock SE and Iterated Simplex). In each method, 10 runs were performed, a maximum of 1500 interactions, and an initial population of 50 poses [24, 25]. The method for performing molecular docking, which consists of the scoring function and search algorithm, was chosen considering an RMSD value ≤ 2.0 as inclusion criteria [27], obtained for coupled ligand with respect to the position of the co-crystallized ligand. The binding energy was obtained in the online server PROtein binDIng enerGy prediction (PRODIGY) [28] and the interactions were visualized with the BIOVIA Discovery Studio program [29].

**Statistical analysis.**

Two-way analysis of variance (ANOVA) with replication was performed to evaluate the effect of different treatments on bacterial strains Pa01, Pa14, and ATCC27853. The treatments included 5-bromoindole-3-carboxaldehyde at three different concentrations (30 µg/ml, 50 µg/ml, and 100 µg/ml), and curcumin at two different concentrations (1 µg/ml and 5 µg/ml). Both the ANOVA analysis and the calculation of standard deviation (SD) were performed using IBM SPSS Statistics for Windows, version 21.0 (IBM Corp., Armonk, NY).

**Results**

**Experimental evaluation**

In the present study were the effectiveness of two compounds, 5-Bromoindole-3-carboxaldehyde and curcumin at different concentrations, in the inhibition of pyocyanin, alginate, elastase and protease quorum-dependent virulence factors in three *Pseudomonas aeruginosa* strains was evaluated: Pa01, Pa14 and ATCC 27853.

**Pyocyanin:** Bromoindole: Strain Pa01 showed significant inhibition at concentrations of 30 µg/ml (85.42%), 50 µg/ml (87.77%) and 100 µg/ml (92.5%). The Pa14 strain also presented a reduction, although smaller, with values of 60.3%, 61.67% and 67.65% at the same concentrations, respectively. Strain ATCC 27853 showed lower inhibition at 30 µg/ml (8.7%) and 50 µg/ml (13.05%), but higher inhibition at 100 µg/ml (72.47%). Regarding the Pa14 strain at the same concentration. Although the Pa01 strain was the one that produced the highest concentration of pyocyanin in its control, it was the strain in which the molecule had the highest activity at all the concentrations studied. On the other hand, the control production of the Pa14 strains and the ATCC 27853 strain varies by 0.001, showing greater activity in the Pa14 strain at concentrations of 30 and 50 µg/ml, but not for the concentration of 100 µg/ml where it is seen. greater inhibition in strain ATCC 27853.

When curcumin was added, strain Pa01 showed an inhibition of 60.68% at 1 µg/ml and 90.63% at 5 µg/ml. The Pa14 strain presented inhibitions of 19.12% at 1 µg/ml and 55.89% at 5 µg/ml. The ATCC 27853 strain showed inhibitions of 81.16% at 1 µg/ml and 100% at 5 µg/ml. Although total inhibition is achieved in strain ATCC 27853, it is the one that showed the lowest pyocyanin production, however, it is the strain with the highest activity, followed by Pa01 and Pa14 respectively. Graph 1.

The two-factor analysis of variance found no significant differences in the inhibition of pyocyanin production between the different treatments with Bromoindole and Curcumin ($F_{(5, 5)} = 1.448$, $p = 0.347$) and between the different bacterial strains evaluated ($F_{(1, 5)} = 0.003$, $p = 0.961$).

**Graph 1.** Pyocyanin inhibition in the presence of different concentrations of curcumin and 5-bromoindole-3-carboxaldehyde

**Elastase:** Bromoindole: For strain Pa01, a significant inhibition was observed with values of 69.77% at 30 μg/ml, 70.64% at 50 μg/ml and 79.85% at 100 μg/ml. The Pa14 strain showed a variable behaviour since when adding 30 μg/ml no inhibition was observed, on the contrary it seems that it favoured the expression of the virulence factor with 21.11% more compared to the control, an inhibition of 30.42% was shown at 50 μg/ml and 32.99% at 100 μg/ml. The ATCC 27853 strain showed an initial inhibition at 100 μg/ml (31.35%) since, as happened with the Pa14 strain, at concentrations of 30 and 50, greater production of the virulence factor was shown, which suggests that at low concentrations it does not It inhibits but it does stimulate production. Therefore, the molecule showed a high inhibitory effect at all concentrations studied in the Pa01 strain, for the Pa14 strain it is the second in terms of activity and in which it showed the least activity was in the ATCC 27853 strain.

The Pa01 strain presented inhibitions when adding curcumin of 3.6% at 1 μg/ml and 32.08% at 5 μg/ml. Strain Pa14 showed minor inhibitions of 5.93% at 1 μg/ml and 56.19% at 5 μg/ml. The ATCC 27853 strain presented inhibitions of 5.65% at 1 μg/ml and 18.5% at 5 μg/ml. The molecule at 1 μg/ml presented the least inhibition in the Pa01 strain, and greater activity at both concentrations in the Pa14 strain. Graph 2.

The two-factor analysis of variance did not find significant differences in the inhibition of elastase production between the different treatments with Bromoindole and Curcumin ($F_{(5, 5)} = 3.844$, $p = 0.083$) and between the different strains evaluated ($F_{(1, 5)} = 1.076$, $p = 0.347$). The variability due to error was relatively low (0.0092), indicating that most of the total variability observed in the data can be explained by the molecules, although not significantly in terms of the strains.

**Graph 2.** Elastase inhibition in the presence of different concentrations of curcumin and 5-bromoindole-3-carboxaldehyde

**Protease:** Bromoindole: Inhibition in strain Pa01 was 92.51% at 30 µg/ml, 92.8% at 50 µg/ml and 89.92% at 100 µg/ml. Being the strain where at low concentrations of 30 µg/ml and 50 µg/ml a considerable inhibition is seen, however at 100 µg/m no greater inhibition is seen, on the contrary it seems that it decreases by about 3% compared to concentration No. 2. Strain Pa14 showed variable inhibition, with an initial increase at 30 µg/ml (66.6%), a reduction at 50 µg/ml (24.25%), and almost complete inhibition at 100 µg/ml (96.22%). The ATCC 27853 strain presented an inhibition of 85.56% at 50 µg/ml and 97.48% at 100 µg/ml, although at 30 µg/ml it also showed an increase in the expression of the virulence factor. However, for this virulence factor at concentrations of 30 and 50 µg/ml it is more effective in the Pa01 strain, while at the concentration of 100 µg/ml in the strain that showed the greatest activity in ATCC 27853 and in the one that showed the least activity at the same concentration was Pa01.

For curcumin, the inhibition in strain Pa01 was 79.83% at 1 µg/ml and 87.04% at 5 µg/ml. The Pa14 strain presented inhibitions of 1.52% at 1 µg/ml and 38.64% at 5 µg/ml. The ATCC 27853 strain showed inhibitions of 41.51% at 1 µg/ml and 71.12% at 5 µg/ml. The strain that showed the greatest inhibition of this virulence factor is Pa01, followed by ATCC 27853 and the one that showed the least activity against the molecule was Pa14. Graph 3.

The analysis of variance showed significant differences between the applied treatments in terms of protease activity ($F(5, 5) = 6.786$, $p = 0.028$). This suggests that at least one of the treatments (at different concentrations of Indole and Curcumin) has a significant effect in reducing protease activity compared to the control. The analysis did not indicate significant differences in the response to treatment between the bacterial strains Pa01, Pa14 and ATCC 27853 ($F(1, 5) = 1.408$, $p = 0.289$). This implies that the strains do not respond differentially to the molecules tested. The variability due to error was relatively low (0.0027), indicating that most of the total variability observed in the data can be explained by the molecules.

**Graph 3.** Protease inhibition in the presence of different concentrations of curcumin and 5-bromoindole-3-carboxaldehyde

**Alginate:** Bromoindole: Strain Pa01 showed an inhibition of 35.87% at 30 μg/ml, 48.4% at 50 μg/ml and 61.4% at 100 μg/ml. The Pa14 strain presented inhibitions of 16.27% at 30 μg/ml, 18.17% at 50 μg/ml and 40.67% at 100 μg/ml. Strain ATCC 27853 showed an initial increase in alginate production at 30 μg/ml (12.22%), but a significant inhibition at 50 μg/ml (52.78%) and 100 μg/ml (76.16%). The molecule was more active in strain Pa01 and ATCC 27853 at concentrations of 50 and 100 μg/ml, where the greatest activity is seen in the latter strain. For Pa14 it showed the least activity, but with significant inhibition.

With curcumin, strain Pa01 showed inhibitions of 37.91% at 1 μg/mL and 48.26% at 5 μg/ml. The Pa14 strain presented inhibitions of 32.3% at 1 μg/ml and 41.63% at 5 μg/ml. The ATCC 27853 strain showed inhibitions of 2.78% at 1 μg/ml and 15% at 5 μg/ml. Graph 4.

The analysis of variance did not find significant differences in alginate activity between the different treatments with Bromoindole and Curcumin ($F_{(5, 5)}$ = 1.637, p = 0.301). Likewise, no significant differences were observed in the response to treatment between the strains evaluated ($F_{(1, 5)}$ = 0.442, p = 0.536).

**Graph 4.** Alginate inhibition in the presence of different concentrations of curcumin and 5-bromoindole-3-carboxaldehyde

Table 1 shows the results in percentage and inhibition concentration of the virulence factors; pyocyanin, elastase, protease and alginate studied with bromoindole. And table 2 shows the results in percentage and inhibition concentration of the virulence factors with curcumin.

**Table 1.** Inhibition percentages of 5-bromoindole-3-carboxaldehyde on Quorum-dependent virulence factors

| Strain / [] | Virulence factors | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | Pyocyanin | | | | Elastase | | | | Protease | | | | Alginate | | | |
| | Control | 30 µg/ml | 50 µg/ml | 100 µg/ml | Control | 30 µg/ml | 50 µg/ml | 100 µg/ml | Control | 30 µg/ml | 50 µg/ml | 100 µg/ml | Control | 30 µg/ml | 50 µg/ml | 100 µg/ml |
| **Pa01** | 0 | 85.42 | 87.77 | 92.5 | 0 | 69.77 | 70.64 | 79.85 | 0 | 92.51 | 92.8 | 89.92 | 0 | 35.87 | 48.4 | 61.4 |
| | (0.384) | (0.56) | (0.047) | (0.029) | (2.282) | (0.69) | (0.67) | (0.46) | (0.694) | (0.052) | (0.05) | (0.07) | (0.686) | (0.44) | (0.354) | (0.265) |
| **Pa14** | 0 | 60.3 | 61.67 | 67.65 | 0 | +21.11 | 30.42 | 32.99 | 0 | +66.6 | 24.25 | 96.22 | 0 | 16.27 | 18.17 | 40.67 |
| | (0.068) | (0.027) | (0.026) | (0.022) | (0.388) | (0.47) | (0.27) | (0.26) | (0.132) | (0.22) | (0.1) | (0.005) | (0.418) | (0.35) | (0.34) | (0.284) |
| **ATCC 27853** | 0 | 8.7 | 13.05 | 72.47 | 0 | +121.1 | +44.22 | 31.35 | 0 | +14.44 | 85.56 | 97.48 | 0 | +12.22 | 52.78 | 76.16 |
| | (0.069) | (0.063) | (0.06) | (0.019) | (0.319) | (0.708) | (0.46) | (0.219) | (0.277) | (0.317) | (0.04) | (0.007) | (0.36) | (0.44) | (0.17) | (0.086) |

**Table 2**. Inhibition percentages of curcumin on Quorum-dependent virulence factors

| | Virulence factors | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| **Strain/ []** | Pyocianin | | | Elastase | | | Protease | | | Alginate | | |
| | Control | 1 µg/ml | 5 µg/ml | Control | 1 µg/ml | 5 µg/ml | Control | 1 µg/ml | 5 µg/ml | Control | 1 µg/ml | 5 µg/ml |
| **Pa01** | 0 | 60.68 | 90.63 | 0 | 3.6 | 32.08 | 0 | 79.83 | 87.04 | 0 | 37.91 | 48.26 |
| | (0.384) | (0.151) | (0.036) | (2.282) | (2.2) | (1.55) | (0.695) | (0.14) | (0.09) | (0.686) | (0.426) | (0.355) |
| **Pa14** | 0 | 19.12 | 55.89 | 0 | 5.93 | 56.19 | 0 | 1.52 | 38.64 | 0 | 32.3 | 41.63 |
| | (0.068) | (0.055) | (0.03) | (0.388) | (0.365) | (0.17) | (0.132) | (0.13) | (0.081) | (0.418) | (0.283) | (0.244) |
| **ATCC 27853** | 0 | 81.16 | 100 | 0 | 5.65 | 18.5 | 0 | 41.51 | 71.12 | 0 | 2.78 | 15 |
| | (0.069) | (0.013) | (0) | (0.319) | (0.301) | (0.26) | (0.277) | (0.162) | (0.08) | (0.36) | (0.35) | (0.306) |

## Computational Studies

Molecular and cheminformatics properties were obtained using the Molinspiration, Molsoft, ChemSketch, swissADME and Osiris Property Explorer simulators. The property values of curcumin and 5-bromoindole-3-carboxaldehyde (Figure 1) are summarized in Table 3. Lipinski's rules indicate that a compound must meet the following criteria to have good bioavailability and to be preferably administered orally: a) LogP value <5; b) TPSA <140 Å; c) molecular weight <500 Da; d) Hydrogen bond donors (-NH, -OH) <5; e) Hydrogen bond acceptors (N, O) <10; When these conditions are met, the molecule under study tends to present a good therapeutic potential. As can be seen in the table, curcumin and 5-bromoindole-3-carboxaldehyde do not present violations of Lipinsky's rules, this would indicate that both compounds are candidates for oral administration, which is an advantage in antibacterial therapy.

**Figure 1.** Structure of curcumin and carboxaldehyde

**Table 3.** Molecular and chemoinformatic properties of curcumin and 5-bromoindole-3-carboxaldehyde

| Properties | Curcumin | 5-bromoindole-3-carboxaldehyde |
|---|---|---|
| MF | $C_{21}H_{20}O_6$ | $C_9H_6BrNO$ |
| MW | 368.3799 | 224.05404 |
| No. HBA | 6 | 2 |
| No. HBD | 2 | 1 |
| PSA | 93.07 | 32.86 |
| LogP | 2.30 | 2.66 |
| δ | $1.279 \pm 0.06$ g/cm$^3$ | $1.727 \pm 0.06$ g/cm$^3$ |
| ST | $54.3 \pm 3.0$ dyne/cm | $63.5 \pm 3.0$ dyne/cm |
| Polarizability | $41.24 \pm 0.5 \times 10^{-24}$ cm$^3$ | $20.99 \pm 0.5 \times 10^{-24}$ cm$^3$ |
| MR | $104.04 \pm 0.3$ cm$^3$ | $52.97 \pm 0.3$ cm$^3$ |
| MV | $287.8 \pm 3.0$ cm$^3$ | $129.7 \pm 3.0$ cm$^3$ |
| Nrotb | 8 | 1 |
| Drug likeness | -3.95 | -6.29 |
| Lipinski's rules Violations | 0 | 0 |
| SMILES | COc2cc(C=CC(=O)CC(=O)C=Cc1ccc(O)c(OC)c1)ccc2O (E/Z)-Curcumin | [H]C(=O)c1c[nH]c2ccc(Br)cc12 |

MF= Molecular formula, MW= Molecular weight (g/mol), HBA= Hydrogen bond acceptors, HBD: Hydrogen bond donors, PSA= Polar Surface Area (Å$^2$), LogP= Partition coefficient, δ= density (g/cm$^3$), ST= Surface tension, MR= Molar refractivity (cm$^3$), nrotb= No. of rotatable bonds, MV= Molar volume (cm$^3$), SMILES= Simplified Molecular Input Line Entry System.

The characteristics to highlight include solubility, which can be determined by the logP of the compounds, as seen in Table 3, curcumin has a LogP of 2.30 and 5-BIC of 2.66, therefore, both compounds are water-soluble, the PSA depends on the MW and indicates the polar area surface of a molecule and the ease with which they can penetrate the cell, for which it must be less than 140 Å, curcumin has a PSA of 93.07 and 5-BIC of 32.86, this is indicative that the molecules could easily penetrate the cell membrane to carry out an effect on the cellular organelles. The HBD and HBA atoms are useful to know if the molecule is specific, and thus, reduce the risks of low bioavailability of the compounds. The topological descriptors of hydrophobicity (Log P), molar refractivity (MR), polarizability (P) and polar surface area (PSA) play an important role when estimating their activity and are generally related to the drug-active site interaction. Compounds with a water-soluble LogP usually present greater biological activity. Likewise, molar refractivity values close to 70 coincide with an increase in activity. Low polarizability (less than 50) indicates that the compound is specific, that is, it does not have the capacity to bind to several ligands, but rather its specificity for a receptor or target site increases. The data in Table 3 indicate that the presence of polar atoms in the molecule (but not polarizable) is necessary for these to be active. These results strengthen the proposal to be administered orally [30].

Molinspiration and Osiris Property Explorer simulators were used to predict the bioactivity of the compounds against important drug target sites and to predict their potential tumorigenic, irritant, reproductive and mutagenic risks, these results are found in Table 4. The cheminformatics and bioactivity properties together indicate that curcumin is the

compound with the greatest pharmacological interest, mainly at the target sites of nuclear receptor ligands and as an enzyme inhibitor, it is important that the values have a positive value to consider them as bioactive at a target site.

**Table 4.** Prediction of bioactivity and toxicity of compounds **1** and **2**

| Compound | NCL | IE | GPCR | KI | PI | RE | T | M | I |
|---|---|---|---|---|---|---|---|---|---|
| Curcumin | 0.12 | 0.08 | -0.06 | -0.26 | -0.14 | No | No | No | No |
| 5-bromoindole-3-carboxaldehyde | -1.22 | -0.53 | -0.77 | -0.57 | -1.53 | No | No | No | No |

* NCL= Nuclear receptor ligand, IE= Enzyme inhibitor, GPCR= GPCR ligand, KI= Kinase inhibitor, RE= Reproductive effective, T= Tumorigenic, M= Mutagenic, I= Irritant.

The greatest failure in the drug discovery process is toxicity and according to the prediction's curcumin and 5-bromoindole-3-carboxaldehyde should not present tumorigenic, irritant, reproductive or mutagenic effects. Based on the combined results of the cheminformatics properties, curcumin is the most promising compound to have the desired biological activity, which would be reflected in a possible better availability and distribution of the compound in the body, this is corroborated by the absorption and metabolism data in Table 5.

**Table 5.** Data of chemoinformatic properties of ADME processes.

| ADMET | | Curcumin | 5-bromoindole-3-carboxaldehyde |
|---|---|---|---|
| **Absorption** | WS | -4.048 | -2.891 |
| | IA (%) | 87.577 | 92.361 |
| | SP | -2.755 | -2.425 |
| | SGP | Yes | Yes |
| | P-GP-1-In | Yes | No |
| **Distribution** | BBBP | -0.511 | Yes |
| | CNSP | -3.07 | -1.758 |
| **Metabolism** | CYP3A4 inhibitor | Yes | No |
| | CYP1A2 inhibitior | Yes | Yes |
| **Excretion** | TC | 0.06 | 0.204 |

*WS: water solubility (log mol/L), IA: intestinal absorption, SP: skin permeability (logKp), SGP: P-glycoprotein substrate, P-GP-1-In: P-glycoprotein I inhibitor, BBBP: blood brain barrier permeability (logBB), CNSP: CNS permeability (logPS), TC: total clearance (log mL/min/kg).

## Molecular docking studies

We performed molecular docking studies to understand the binding interactions of curcumin and 5-bromoindole-3-carboxaldehyde with the LasR, PQS, and RhlR proteins of the Quorum sensing of *Pseudomona aeruginosa*. The molecular docking protocols were validated with re-docking, coupling the co-crystallized ligands in the proteins obtained in the Protein Data Bank using 12 methods from the combination of search algorithms and scoring functions, which are described in the materials and methods section. Figure 2 shows the results of the re-docking of the LasR, PQS and RhlR proteins with the co-crystallized ligands, as well as the method that was selected to perform the molecular docking of the two compounds under study from an RMSD less than 2 Å, which indicates that these methods can predict the experimental binding mode in the Quorum sensing proteins.

**Figure 2.** Re-docking analysis, the selected methods reproduce the pose of co-crystallized ligands (yellow colour) with RMSD less than 2 Å in the superposition of the re-coupled ligands (red colour). Docking protocols, each method is implemented by a scoring function and a search algorithm: LasR (MolDock Score/MolDock Optimizer), PQS (PLANTS Score/Iterated Simplex) and RhlR MolDock Score GRID/MolDock Optimizer).

Table 6 shows the affinity energies of curcumin and 5-bromoindole-3-carboxyaldehyde on QS proteins, as well as the binding affinity of the endogenous inducers of the proteins LasR (3O-C12-HSL) and RhlR (C4-HSL) and the PQS inhibitor (3NH2-7Cl-C9-QZN) [31]. The amino acids of the ligand binding domain (LBD) with which the endogenous ligands interact are also shown, and the two compounds studied; as well as the types of ligand-receptor interaction. It should be noted that curcumin is the compound that has the highest affinity energy on the LBD in the 3 proteins, while 5-Bromoindole-3-carboxaldehyde is the molecule

with the lowest affinity energy; however, it is inferred that it could be a Quorum sensing inhibitor since it binds within the LBDs of the LasR, PQS and RhlR proteins, as can be seen in Figures 3, 4 and 5.

**Table 6.** Affinity energy and binding mode of curcumin and 5-bromoindole-3-carboxaldehyde with *P. aeruginosa* Quorum sensing proteins

| Target | Compounds | $\Delta G$ Kcal/mol | Interaction |
|---|---|---|---|
| LasR | $3O\text{-}C_{12}\text{-}HSL$ | -9.29 | Hydrogen bond (Tyr56, Asp73, Thr75, Ser129), Carbon hydrogen bond (Ser29, Tyr93), alkyl (Leu36, Ala50, Val76, Cys79, Ala105, Leu110, Leu125, Ala127), pi-alkyl (Tyr64, Trp88, Phe101). |
| | Curcumin | -11.07 | Hydrogen bond (Arg61), pi-anion (Asp73), pi-donor hydrogen bond (Tyr56), pi-pi stacked (Tyr64), pi-pi T-shaped (Tyr56), alkyl (Ala105, Leu110), pi-alkyl (Leu36, Trp60, Phe101, Leu110). |
| | 5-Bromoindole-3-carboxaldehyde | -7.66 | Hydrogen bond (Arg61, Tyr93, Leu125), pi-anion (Asp73), pi-donor hydrogen bond (Tyr56), alkyl (Val76, Cys79, Leu125), pi-alkyl (Leu40, Ala50, Trp60, Val76, Phe101, Leu110). |
| PQS | $3NH_2\text{-}7Cl\text{-}C9\text{-}QZN$ | -8.23 | Hydrogen bond (Leu207), alkyl (Ala102, Ile149, Ile186, Ile189, Leu207, Ile236, Pro238, Ile263), pi-alkyl (Ile149, Ala168, Leu207, Leu208, Ile236, Tyr258). |
| | Curcumin | -9.23 | Hydrogen bond (Gln194, Leu207), carbon hydrogen bond (Ser196, Arg209, Pro210), pi-sigma (Leu208), alkyl (Ile149, Ile236, Pro238), pi-alkyl (Leu208, Phe221, Ile236). |
| | 5-Bromoindole-3-carboxaldehyde | -7.14 | Hydrogen bond (Ile236, Tyr258), carbon hydrogen bond (Ile186, Leu207), pi-pi T-shaped (Tyr258), alkyl (Ile186, |

| | | | |
|---|---|---|---|
| | | | Leu189, Leu208, Ile236), pi-alkyl (Ala168, Val170, Leu208, Ile236). |
| RhlR | C4-HSL | -7.12 | Hydrogen bond (Try64, Trp68, Asp81, Ser135), alkyl (Ile84, Leu107, Ala111, Val133), pi-alkyl (Trp96, Phe101). |
| | Curcumin | -10.27 | Hydrogen bond (Trp68), pi-anion (Asp81), pi-pi stacked (Trp96), alkyl (Leu107, Ala111), pi-alkyl (Ala44, Ala83, Phe101, Trp108). |
| | 5-Bromoindole-3-carboxaldehyde | -7.68 | Hydrogen bond (Trp68), carbon hydrogen bond (Pro65, Leu69), pi-donor hydrogen bond (Ser135), pi-pi T-shaped (Tyr72), alkyl (Val60, Leu69, Ile84), pi-alkyl (Ala44, Tyr64, Val60, Ala83). |

The ligand binding domain of LasR is housed in a large hydrophobic pocket formed by the amino acids Leu36, Gly38, Leu39, Leu40, Tyr47, Glu48, Ala50, Ile52, Tyr56, Trp60, Arg61, Tyr64, Asp65, Gly68, Tyr69, Ala70, Asp73, Pro74, Thr75, Val76, Cys79, Thr80, Trp88, Tyr93, Phe101, Phe102, Ala105, Leu110, Thr115, Leu125, Gly126, Ala127, and Ser129 [32]. Figure 3 shows the interactions between curcumin and 5-Bromoindole-3-carboxaldehyde with LBD amino acids and that they share with the endogenous ligand 3O-C12-HSL, such as Tyr56, Asp73, Leu110 and Phe101. In addition, curcumin also interacts with Leu36 and Tyr64 as the endogenous ligand. Meanwhile, 5-Bromoindole-3-carboxaldehyde as well as 3O-C12-HSL interact with the residues Ala50, Val76, Cys79, Tyr93 and Leu125.

**Figure 3.** Results of molecular docking on LasR. The image shows the position of the ligand binding domain (red) on the LasR protein and the 2D interactions of curcumin (A) and 5-Bromoindole-3-carboxaldehyde (B).

Figure 4 shows the structure of the LysR-like transcription factor that responds to the quinolone signal in Pseudomonas (PQS) present in *Pseudomona aeruginosa*, as well as the ligand binding domain in yellow that covers from residue 91 to 319 [33]. Ilangovan et al., in 2013 evaluated quinazolinone analogues and reported the compound 3NH2-7Cl-C9-QZN (QZN 23) as a novel PQS inhibitor with an IC50 of 5.0 mM [31], which binds to the LBD with the amino acids reported in Table 6. The interactions between curcumin and 5-Bromoindole-3-carboxaldehyde in the LBD are shown in Figure 4. QZN-23 inhibitor, curcumin, and 5-Bromoindole-3-carboxaldehyde interact with Leu207, Leu208, and Ile 236 residues. Curcumin also interacts with the amino acids Ile149 and Pro238 as well as the inhibitor QZN 23. While 5-Bromoindole-3-carboxaldehyde like QZN 23 binds with Ala168, Ile186, Ile189 and Tyr258.

**Figure 4.** Results of molecular docking on PQS. The image shows the position of the ligand binding domain (yellow) on the PQS protein and the 2D interactions of curcumin (A) and 5-Bromoindole-3-carboxaldehyde (B).

It has been reported that for the multimerization and activity of RhlR, its autoinducer C4-HSL is necessary, which binds in the LBD efficiently and that, in its absence, homodimerizes decreasing its capacity as a transcriptional activator [34]. The residues of RhlR LBD with which C4-HSL interacts are reported in Table 6. Both C4-HSL, curcumin and 5-Bromoindole-3-carboxaldehyde bind in LBD and interact commonly with Trp68. Curcumin also binds with other amino acids with which it interacts with C4-HSL, Asp81, Phe101, Leu107 and Ala111. Meanwhile, 5-Bromoindole-3-carboxaldehyde binds with Tyr64, Ile84 and Ser135 residues as well as C4-HSL, as shown in Figure 5.

**Figure 5.** Results of molecular docking on RhlR. The image shows the position of the LBD (grey) on the RhlR protein and the 2D interactions of curcumin (A) and 5-Bromoindole-3-carboxaldehyde (B).

## Discussion

It has been shown that *P. aeruginosa* employs quorum sensing in the regulation of genes encoding extracellular virulence factors. Quorum sensing is an intercellular communication system that allows bacteria to control gene expression in a cell population density-dependent manner [35].

In *P. aeruginosa*, four fundamental QS pathways have been identified. The first is known as the LasI/LasR cascade. The second includes the RhlI/RhlR pathway. The third is the *Pseudomonas* quinolone signal (PQS). The fourth is the AI-based integrated quorum sensing system [36].

In particular, QS blockade abrogates the virulence without affecting bacterial growth. Therefore, QS represents an attractive target for anti-virulence therapy [37].

In a study conducted by Rudrappa in Pa01 where they evaluated different concentrations of curcumin, at concentrations of 1.5 to 3 µg/ml they achieved an inhibition of pyocyanin of between 60 and 80% of the production with respect to the control, and at 5 µg/ml they achieved an inhibition of 3 times compared to the control. Not only did they achieve an inhibition of pyocyanin, other factors regulated by the QS systems lasI-lasR and rhlI-rhlR were also studied, where they observed a double decrease in the production of elastase and protease, where they also observed that curcumin directly affected the production of the lactones 3-oxoC12-HSL and C4-HSL. In our study we observed that at 1 µg/ml of curcumin we obtained an inhibition for Pa01 of 60% which is similar to that observed for Rudrappa, a value similar to that observed in the ATCC 27853 strain [38].

Sethupathy and collaborators demonstrated the activity of curcumin at 5 µg/ml on the inhibition of pyocyanin, elastase, protease, alginate and biofilm in 32 clinical isolates and using Pa01 as a control, finding inhibitions that they considered as effective inhibition (50-90%), minimal inhibition (20-49%), no inhibition (0-19%) and induced. Where most of the strains had an effectiveness of between 50 and 90% at said concentration, the factors that showed said inhibition are mainly pyocyanin and elastase. The rest of the strains were maintained with minimal inhibition [39]. In our study at a concentration of 1 µg/ml for Pa01, an inhibition of 60.60% was obtained and for 5 mg of 90.63%, for Pa14 an inhibition of 55.89% was achieved at that concentration, where the greatest activity is seen is in the ATCC 27853 strain where it is achieves 100% pyocyanin inhibition. In addition to similar results in the rest of the virulence factors.

A published study reported for Fernandes, that curcumin has inhibitory effects on the LuxS/AI-2 and LasI/LasR QS systems, resulting in a reduction of 33%–77% and 21%, respectively. Significantly, curcumin at high concentration of 200 µg/ml resulted in a 21% decrease in the generation of 3-oxo-C12-HSL by *P. aeruginosa* Pa14 strain [40].

Liu and collaborators, they monitored pyocyanin production and biofilm formation in *Pseudomonas aeruginosa* when adding curcumin, and they showed a significant decrease in both virulence factors with the aforementioned molecule [41].

Gholami and collaborators studied the activity of curcumin and curcumin with metal complexes to inhibit quorum and found important decreases in pyocyanin and considerable alginate. Their study shows that curcumin alone reduces pyocyanin production by more than half with a 67.5 % compared to the control in Pa01, and to a lesser extent with metals. On the other hand, the activity in alginate with curcumin alone is lower than that shown in combination with metals with only 21.3% [42].

Jaduan tested the effect of curcumin mixed with honey to inhibit quorum in *Pseudomonas aeruginosa* in strain Pa01 with the following results: Treatment with curcumin (50 µg/ml) and honey (1%) only showed a weak inhibitory effect on siderophore production, a result similar to that observed in treatment with curcumin alone. Curcumin (50 µg/ml) and honey (1%) alone were not found to significantly reduce the activity of Pa01 metalloproteases elastase y protease. Interestingly, curcumin (50 µg/ml) and honey (1%) alone were ineffective in decreasing alginate production [43]. This synergy shows lower activity than what we obtained in our study for the virulence factors evaluated.

It is known that curcumin acts on the las system. This signalling system is regulated by virulence factors such as the expression of alkaline protease and elastase, which represents the decrease in the production of the autoinducer lactone and therefore the inhibition of production of said factors. As shown in the results [44]. If lasR/lasI system transcriptional regulators expression is affected, other systems such as the PQS and all virulence factors in *P. aeruginosa* will be affected too [45].

Several studies have described the antibiofilm and antivirulence activities of indole derivatives against clinically important pathogens such as *Pseudomonas aeruginosa* [46]. Halogenation of indoles demonstrates a decrease in IC50 and also greater potency as inhibitors of quorum sensing, as described by Odularu et. al., [17]. The antimicrobial activity of some indolecarboxaldehydes has been demonstrated in strains of *Pseudomonas sp*, but there is no information on quorum inhibition in *Pseudomonas aeruginosa* [47].

Among the most studied bacterial species, *C. violaceum* has been shown to affect the production of AHL with good results at concentrations similar to those studied by us [18].

Indolecarboxaldehydes such as 5-bromoindole-3-carboxaldehyde and derivatives in *Vibrio fischeri* have also been shown to inhibit the LuxI/LuxR system, thereby quenching its bioluminescence or biofilm inhibition in *Vibrio parahaemolyticus* [18,48].

In *Pseudomonas aeruginosa*, the following indole derivatives have been found Indole, 3-indolylacetonitrile, 7-fluoroindole, indole-based N-acylated L-homoserine lactones and 2-Aminobenzimidazoles, which can inhibit the production of virulence factors and biofilm formation, however, carboxaldehyde derivatives with this activity are not reported [49].

The solvent did not affect the production of virulence factors at concentrations of 0.01%, 0.5% and 1% (v/v), according to the prepared stocks of the molecules, since equal results were obtained in control and negative control, however some authors report that DMSO can act on quorum sensing at higher concentrations such as 2% (v/v).

The analyses of variance performed for elastase, protease and alginate activities in *Pseudomonas aeruginosa* with Curcumin and 5-bromoindole-3-carboxaldehyde revealed significant differences only in protease activity between treatments. These results suggest that while some compounds may influence the activity of certain enzymes, not all treatments have a uniform impact on all enzyme activities evaluated.

Some limitations of our work are molecular studies to know exactly which gene is repressed and to what extent it is done, however, this represents an opportunity for the future, especially in the case of bromoindole studied, since so far, we are the first to report this activity in three different strains of *Pseudomonas aeruginosa*.

Chemoinformatic properties related to ADME processes were predicted using the online predictors pkCSM and SwissADME (Table 5). Absorption properties indicate that the compounds can be administered orally according to Lipinsky rules. Other predicted properties, such as LogP data, absorption, highlighting intestinal absorption which is 87.5% for curcumin and 92.3% for 5-bromoindole-3-carboxaldehyde, the skin permeability of both compounds is relatively similar (-2.755 and -2.425 respectively), in addition to the fact that curcumin could inhibit P-GP-1-In, which would cause the compound not to be expelled from

the cell and could promote biological activity. On the other hand, 5-bromoindole-3-carboxaldehyde appears to permeate the blood-brain barrier while curcumin does not, this is to be expected due to the difference in size, polar atoms and solubility, but opens the possibility that 5-bromoindole-3-carboxaldehyde could be used in the treatment of infections affecting the Central Nervous System.

*In silico* analysis was used to support the hypothesis that curcumin and 5-bromoindole-3-carboxaldehyde interact with the three major Quorum sensing proteins of *P. aeruginosa* (Table 6 and Figures 3-5). Molecular docking studies were performed to obtain the affinity energy and binding mode of the molecules under study in the LBDs of the pharmacological targets.

Molecular docking with the LasR protein showed that curcumin has a stronger binding (-11.07 Kcal/mol) compared to its endogenous ligand (-9.29 Kcal/mol), while 5-bromoindole-3-carboxaldehyde was weaker in affinity (-7.66 Kcal/mol). Similarly, curcumin has a higher energy affinity for the PQS protein (-9.23 Kcal/mol) compared to the QZN 23 ligand that has an $IC_{50}$ of 5.0 μM (-8.23 Kcal/mol), which we rely on to infer that curcumin could be a PQS inhibitor, while 5-bromoindole-3-carboxaldehyde has a weaker binding (-7.14 Kcal/mol). Finally, with the RhlR protein, curcumin also presented a higher affinity energy (-10.27 Kcal/mol) compared to the affinity presented by the endogenous ligand C4-HSL (-7.12 Kcal/mol), while 5-bromoindole-3-carboxaldehyde presented an affinity like the endogenous ligand (-7.68 Kcal/mol). It is important to mention that the result of the affinity energy is only indicative of a higher probability of a more stable bond. Therefore, the binding mode of the ligands under study should be compared with endogenous ligands or other inhibitors or antagonists of the Quorum sensing proteins.

In this sense, Figures 3, 4 and 5 show the binding modes of curcumin and 5-bromoindole-3-carboxaldehyde in the ligand binding domains of the LasR, PQS and RhlR proteins. It is shown that both compounds under study can accommodate at the site of activation of LasR and RhlR, suggesting a possible competition or antagonism with their autoinducers in the LBD and by interacting with essential residues they promote a disruption of the QS signalling cascade. Likewise, curcumin and 5-bromoindole-3-carboxaldehyde are accommodated in the same LBD cavity as the QZN 23 inhibitor and bind with amino acids relevant to PQS activity, preventing their interaction with DNA and impeding their transcriptional activity. Taken together, the *in silico* analysis supports experimental results showing that both curcumin and 5-bromoindole-3-carboxdehyde are potential inhibitors of QS.

### Conclusion

This study provides strong evidence that both 5-bromoindole-3-carboxaldehyde and curcumin are effective in inhibiting quorum-sensing-dependent virulence factors in *Pseudomonas aeruginosa* in reference strains such as Pa01, Pa14 and ATCC 27853.

5-bromoindole-3-carboxaldehyde showed significant inhibition of pyocyanin, elastase, protease and alginate production, particularly in strain Pa01, which showed the greatest reductions in the levels of these virulence factors. The inhibition observed in strains Pa14

and ATCC 27853, although notable, was lower, suggesting further studies to elucidate the mechanism.

On the other hand, curcumin is a potent inhibitor of the same virulence factors in all strains studied, with a complete inhibition of pyocyanin production in strain ATCC 27853 at a concentration of 5 μg/ml. However, the efficacy of curcumin is variable between strains, with strain Pa14 showing the lowest overall sensitivity to this compound.

These findings support the hypothesis that inhibition of the quorum sensing system may be an effective strategy to reduce the virulence of *P. aeruginosa* without affecting its growth. The variability observed in the response of the different strains underlines the importance of considering bacterial heterogeneity when developing therapies based on quorum sensing inhibition.

**Conflicts of interests**

The authors declare that they have no conflict of interest.

**Sources of funding**

This research work did not receive funding.

**Code of ethics**

In the present study, reference strains of *Pseudomonas aeruginosa* (Pa01, Pa14 and ATCC 27853) were used to investigate the quorum sensing inhibitory activity of curcumin and 5-bromoindole-3-carboxyaldehyde. All experimental activities were carried out in compliance with BSL-1 level biosafety regulations, strictly following institutional protocols for the safe handling of microorganisms. The use of reference strains in this study does not involve humans or animals, and the samples used are of non-clinical origin and have been widely characterized and documented in the scientific literature. Therefore, ethics committee approval is not required for the performance of this work.

**Acknowledgment**

The researchers thank the academic direction of the degree in Biological Pharmaceutical Chemistry and laboratory technicians for their support in carrying out this research work and Cristian Sadalis Santos López for their technical support.

**References**

[1]   Tuon, F. F.; Dantas, L. R.; Suss, P. H.; Tasca Ribeiro, V. S. Pathogenesis of the *Pseudomonas aeruginosa* Biofilm: A Review. *Pathogens* **2022**, *11* (3), 300. https://doi.org/10.3390/pathogens11030300.

[2]   Spagnolo, A. M.; Sartini, M.; Cristina, M. L. *Pseudomonas aeruginosa* in the Healthcare Facility Setting. *Reviews in Medical Microbiology* **2021**, *32* (3), 169–175. https://doi.org/10.1097/MRM.0000000000000271.

[3]    Moradali, M. F.; Ghods, S.; Rehm, B. H. A. *Pseudomonas aeruginosa* Lifestyle: A Paradigm for Adaptation, Survival, and Persistence. *Front. Cell. Infect. Microbiol.* **2017**, *7*. https://doi.org/10.3389/fcimb.2017.00039.

[4]    Reynolds, D.; Kollef, M. The Epidemiology and Pathogenesis and Treatment of *Pseudomonas aeruginosa* Infections: An Update. *Drugs* **2021**, *81* (18), 2117–2131. https://doi.org/10.1007/s40265-021-01635-6.

[5]    Pang, Z.; Raudonis, R.; Glick, B. R.; Lin, T.-J.; Cheng, Z. Antibiotic Resistance in *Pseudomonas aeruginosa*: Mechanisms and Alternative Therapeutic Strategies. *Biotechnology Advances* **2019**, *37* (1), 177–192. https://doi.org/10.1016/j.biotechadv.2018.11.013.

[6]    Qin, S.; Xiao, W.; Zhou, C.; Pu, Q.; Deng, X.; Lan, L.; Liang, H.; Song, X.; Wu, M. *Pseudomonas aeruginosa*: Pathogenesis, Virulence Factors, Antibiotic Resistance, Interaction with Host, Technology Advances and Emerging Therapeutics. *Sig Transduct Target Ther* **2022**, *7* (1), 199. https://doi.org/10.1038/s41392-022-01056-1.

[7]    Scheffler, R. J.; Bratton, B. P.; Gitai, Z. *Pseudomonas aeruginosa* Clinical Blood Isolates Display Significant Phenotypic Variability. *PLoS ONE* **2022**, *17* (7), e0270576. https://doi.org/10.1371/journal.pone.0270576.

[8]    Edward, E. A.; El Shehawy, M. R.; Abouelfetouh, A.; Aboulmagd, E. Prevalence of Different Virulence Factors and Their Association with Antimicrobial Resistance among *Pseudomonas aeruginosa* Clinical Isolates from Egypt. *BMC Microbiol* **2023**, *23* (1), 161. https://doi.org/10.1186/s12866-023-02897-8.

[9]    Jurado-Martín, I.; Sainz-Mejías, M.; McClean, S. *Pseudomonas aeruginosa*: An Audacious Pathogen with an Adaptable Arsenal of Virulence Factors. *IJMS* **2021**, *22* (6), 3128. https://doi.org/10.3390/ijms22063128.

[10]   Abbas, H. A. Inhibition of Virulence Factors of *Pseudomonas aeruginosa* by Diclofenac Sodium. *Roum Arch Microbiol Immunol* **2015**, *74* (3–4), 79–85.

[11]   Tajani, A. S.; Jangi, E.; Davodi, M.; Golmakaniyoon, S.; Ghodsi, R.; Soheili, V.; Fazly Bazzaz, B. S. Anti-Quorum Sensing Potential of Ketoprofen and Its Derivatives against *Pseudomonas aeruginosa*: Insights to in Silico and in Vitro Studies. *Arch Microbiol* **2021**, *203* (8), 5123–5132. https://doi.org/10.1007/s00203-021-02499-w.

[12]   Adonizio, A.; Kong, K.-F.; Mathee, K. Inhibition of Quorum Sensing-Controlled Virulence Factor Production in *Pseudomonas aeruginosa* by South Florida Plant Extracts. *Antimicrob Agents Chemother* **2008**, *52* (1), 198–203. https://doi.org/10.1128/AAC.00612-07.

[13]   Tanhay Mangoudehi, H.; Zamani, H.; Shahangian, S. S.; Mirzanejad, L. Effect of Curcumin on the Expression of ahyI/R Quorum Sensing Genes and Some Associated Phenotypes in Pathogenic *Aeromonas hydrophila* Fish Isolates. *World J Microbiol Biotechnol* **2020**, *36* (5), 70. https://doi.org/10.1007/s11274-020-02846-x.

[14] Luiza Koop, B.; Nascimento da Silva, M.; Diniz da Silva, F.; Thayres dos Santos Lima, K.; Santos Soares, L.; José de Andrade, C.; Ayala Valencia, G.; Rodrigues Monteiro, A. Flavonoids, Anthocyanins, Betalains, Curcumin, and Carotenoids: Sources, Classification and Enhanced Stabilization by Encapsulation and Adsorption. *Food Research International* **2022**, *153*, 110929. https://doi.org/10.1016/j.foodres.2021.110929.

[15] Packiavathy, I. A. S. V.; Priya, S.; Pandian, S. K.; Ravi, A. V. Inhibition of Biofilm Development of Uropathogens by Curcumin – An Anti-Quorum Sensing Agent from Curcuma Longa. *Food Chemistry* **2014**, *148*, 453–460. https://doi.org/10.1016/j.foodchem.2012.08.002.

[16] Dai, C.; Lin, J.; Li, H.; Shen, Z.; Wang, Y.; Velkov, T.; Shen, J. The Natural Product Curcumin as an Antibacterial Agent: Current Achievements and Problems. *Antioxidants* **2022**, *11* (3), 459. https://doi.org/10.3390/antiox11030459.

[17] Odularu, A. T.; Afolayan, A. J.; Sadimenko, A. P.; Ajibade, P. A.; Mbese, J. Z. Multidrug-Resistant Biofilm, Quorum Sensing, Quorum Quenching, and Antibacterial Activities of Indole Derivatives as Potential Eradication Approaches. *BioMed Research International* **2022**, *2022*, 1–9. https://doi.org/10.1155/2022/9048245.

[18] Kemp, C. A.; McCullough, D. K.; Bialonska, D.; Johnson, P. J. T. Effect of Bromination on the Quorum Sensing-Inhibiting Properties of Indole-3-Carboxaldehydes in *Chromobacterium violaceum* AHL System. *Microbiology Research* **2021**, *12* (2), 376–382. https://doi.org/10.3390/microbiolres12020025.

[19] Essar, D. W.; Eberly, L.; Hadero, A.; Crawford, I. P. Identification and Characterization of Genes for a Second Anthranilate Synthase in *Pseudomonas aeruginosa*: Interchangeability of the Two Anthranilate Synthases and Evolutionary Implications. *J Bacteriol* **1990**, *172* (2), 884–900. https://doi.org/10.1128/jb.172.2.884-900.1990.

[20] Huelgas, D.; Cazares, D.; Alcaraz, L. D.; Ceapã, C. D.; Cocotl-Yañez, M.; Shotaro, T.; Maeda, T.; Fernández-Presas, A. M.; Tostado-Islas, O.; González-Vadillo, A. L.; Limones-Martínez, A.; Hernandez-Cuevas, C. E.; González-García, K.; Jiménez-García, L. F.; Martínez, R.-L.; Santos-López, C. S.; Husain, F. M.; Khan, A.; Arshad, M.; Kokila, K.; Wood, T. K.; García-Contreras, R. Exoprotease Exploitation and Social Cheating in a *Pseudomonas aeruginosa* Environmental Lysogenic Strain with a Noncanonical Quorum Sensing System. *FEMS Microbiology Ecology* **2023**, *99* (9), fiad086. https://doi.org/10.1093/femsec/fiad086.

[21] Loarca, D.; Díaz, D.; Quezada, H.; Guzmán-Ortiz, A. L.; Rebollar-Ruiz, A.; Presas, A. M. F.; Ramírez-Peris, J.; Franco-Cendejas, R.; Maeda, T.; Wood, T. K.; García-Contreras, R. Seeding Public Goods Is Essential for Maintaining Cooperation in *Pseudomonas aeruginosa*. *Front. Microbiol.* **2019**, *10*, 2322. https://doi.org/10.3389/fmicb.2019.02322.

[22] Knutson, C. A.; Jeanes, A. A New Modification of the Carbazole Analysis: Application to Heteropolysaccharides. *Analytical Biochemistry* **1968**, *24* (3), 470–481. https://doi.org/10.1016/0003-2697(68)90154-1.

[23] Ohman, D. E.; Cryz, S. J.; Iglewski, B. H. Isolation and Characterization *of Pseudomonas aeruginosa* PaO1 Mutant That Produces Altered Elastase. *J Bacteriol* **1980**, *142* (3), 836–842. https://doi.org/10.1128/jb.142.3.836-842.1980.

[24] Córdova-Bahena, L.; Sánchez-Álvarez, A.A.; Ruiz-Moreno, A.J.; Velasco-Velázquez, M.A. Repositioning of Etravirine as a Potential CK1e Inhibitor by Virtual Screening. Pharmaceuticals **2022**, 15(1): 8. DOI: 10.3390/ph15010008

[25] Alvarado Salazar, J. A.; Valdes, M.; Cruz, A.; Moreno De Jesús, B.; Patiño González, D.; Olivares-Corichi, I. M.; Tamay Cach, F.; Mendieta Wejebe, J. E. In Silico and In Vivo Evaluation of Novel 2-Aminobenzothiazole Derivative Compounds as Antidiabetic Agents. Preprints **2024**, 2024082205. DOI: 10.20944/preprints202408.2205.v1

[26] Bhagwat, S.K.; Janardan Pawar, T.; Kulkarni, S.A.; Patil, A.A.; Ashokrao More, R.; Jimenez-Halla, J.O.C.; Alvardo-Salazar, J.A.; Olivares-Romero, J.L.; Muteeb, G.; Delgado-Alvarado, E.; Patil, S.V. Synthesis, characterization, biological activities, and computational studies of pyrazolyl-thiazole derivatives of thiophene. RSC Adv. **2024**, 14: 39004-39016. DOI: 10.1039/D4RA06228K

[27] Houston, DR & Walkinshaw, MD. Consensus docking: improving the reliability of docking in a virtual screening context. J. Chem. Inf. Model. **2013**, 53(2): 384-390. DOI: 10.1021/ci300399w

[28] Xue, L.C.; Rodrigues, J.P.; Kastritis, P.L.; Bonvin, A.M.; Vangone, A. PRODIGY: a web server for predicting the binding affinity of protein-protein complexes. Bioinformatics. 2016, 32(23): 3676-3678. DOI: 10.1093/bioinformatics/btw514

[29] Pawar, S.S. & Rohane, S.H. Review on Discovery Studio: An important Tool for Molecular Docking. Asian J. Research Chem. **2021**, 14(1): 86-88. DOI: 10.5958/0974-4150.2021.00014.6

[30] Malavé, J. A.; Márquez Brazón, E.; Lezama, J. A. Cálculo de descriptores moleculares topológicos en flavonoides con actividad anti-VIH-1. *Saber* **2015**, *27* (1), 102–109.

[31] Ilangovan, A.; Fletcher, M.; Rampioni, G.; Pustelny, C.; Rumbaugh, K.; Heeb, S.; Cámara, M.; Truman, A.; Chhabra, S.R.; Emsley, J.; Williams, P. Structural basis for native agonist and synthetic inhibitor recognition by the *Pseudomonas aeruginosa* Quorum Sensing regulator PqsR (MvfR). PLOS Pathog. **2013**, 9(7): e1003508. DOI: 10.1371/journal.ppat.1003508

[32] Bottomley, M.J.; Muraglia, E.; Bazzo, R.; Carfi, A. Molecular insights into Quorum sensing in the human pathogen *Pseudomonas aeruginosa* from the structure of the virulence regulator LasR bound to Its autoinducer. *J Biol. Chem*. **2007**, 282(18): 13592-13600. DOI: 10.1074/jbc.M700556200

[33] Xu, N.; Yu, S.; Moniot, S.; Weyand, M.; Blankenfeld, W. Crystallization and preliminary crystal structure analysis of the ligand-binding domain of PqsR (MvfR), the Pseudomonas quinolone signal (PQS) responsive quorum-sensing transcription factor of *Pseudomonas aeruginosa*. *Acta Cryst.* **2012**, 68(9): 1034-1039. DOI: 10.1107/S1744309112032538

[34] Lamb, J.R.; Patel, H.; Montminy, T.; Wagner, V.E.; Iglewski, B.H. Functional domains of the RhlR transcriptional regulator of *Pseudomonas aeruginosa*. *J. Bacteriol*. **2003**, 185(24): 7129-7139. DOI: 10.1128/jb.185.24.7129-7139.2003

[35] Ishida, T.; Ikeda, T.; Takiguchi, N.; Kuroda, A.; Ohtake, H.; Kato, J. Inhibition of Quorum Sensing in *Pseudomonas aeruginosa* by *N* -Acyl Cyclopentylamides. *Appl Environ Microbiol* **2007**, *73* (10), 3183–3188. https://doi.org/10.1128/AEM.02233-06.

[36] Naga, N. G.; Zaki, A. A.; El-Badan, D. E.; Rateb, H. S.; Ghanem, K. M.; Shaaban, M. I. Inhibition of *Pseudomonas aeruginosa* Quorum Sensing by Methyl Gallate from Mangifera Indica. *Sci Rep* **2023**, *13* (1), 17942. https://doi.org/10.1038/s41598-023-44063-0.

[37] Xie, Y.; Chen, J.; Wang, B.; Peng, A.-Y.; Mao, Z.-W.; Xia, W. Inhibition of Quorum-Sensing Regulator from *Pseudomonas aeruginosa* Using a Flavone Derivative. *Molecules* **2022**, *27* (8), 2439. https://doi.org/10.3390/molecules27082439.

[38] Rudrappa, T.; Bais, H. P. Curcumin, a Known Phenolic from Curcuma Longa, Attenuates the Virulence of *Pseudomonas aeruginosa* PaO1 in Whole Plant and Animal Pathogenicity Models. *J. Agric. Food Chem.* **2008**, *56* (6), 1955–1962. https://doi.org/10.1021/jf072591j.

[39] Sethupathy, S.; Prasath, K. G.; Ananthi, S.; Mahalingam, S.; Balan, S. Y.; Pandian, S. K. Proteomic Analysis Reveals Modulation of Iron Homeostasis and Oxidative Stress Response in *Pseudomonas aeruginosa* PaO1 by Curcumin Inhibiting Quorum Sensing Regulated Virulence Factors and Biofilm Production. *J Proteomics* **2016**, *145*, 112–126. https://doi.org/10.1016/j.jprot.2016.04.019.

[40] Fernandes, S.; Borges, A.; Gomes, I. B.; Sousa, S. F.; Simões, M. Curcumin and 10-Undecenoic Acid as Natural Quorum Sensing Inhibitors of LuxS/AI-2 of *Bacillus Subtilis* and LasI/LasR of *Pseudomonas aeruginosa*. *Food Research International* **2023**, *165*, 112519. https://doi.org/10.1016/j.foodres.2023.112519.

[41] Liu, L.; Cao, X.; Ma, W.; Chen, L.; Li, S.; Hu, B.; Xu, Y. In-Situ and Continuous Monitoring of Pyocyanin in the Formation Process of *Pseudomonas aeruginosa* Biofilms by an Electrochemical Biosensor Chip. *Sensors and Actuators B: Chemical* **2021**, *327*, 128945. https://doi.org/10.1016/j.snb.2020.128945.

[42] Gholami, M.; Zeighami, H.; Bikas, R.; Heidari, A.; Rafiee, F.; Haghi, F. Inhibitory Activity of Metal-Curcumin Complexes on Quorum Sensing Related Virulence Factors of *Pseudomonas aeruginosa* PaO1. *AMB Expr* **2020**, *10* (1), 111. https://doi.org/10.1186/s13568-020-01045-z.

[43] Jadaun, V.; Prateeksha, P.; Singh, B. R.; Paliya, B. S.; Upreti, D. K.; Rao, Ch. V.; Rawat, A. K. S.; Singh, B. N. Honey Enhances the Anti-Quorum Sensing Activity and Anti-Biofilm Potential of Curcumin. *RSC Adv.* **2015**, *5* (87), 71060–71070. https://doi.org/10.1039/C5RA14427B.

[44] Li, Q.; Mao, S.; Wang, H.; Ye, X. The Molecular Architecture of *Pseudomonas aeruginosa* Quorum-Sensing Inhibitors. *Marine Drugs* **2022**, *20* (8), 488. https://doi.org/10.3390/md20080488.

[45] Rúgeles, D.; Gámez-Castillo, B.; Gómez, V.; Rodríguez, P. H. Effects of Natural Compounds and Commercial Antibiotics on *Pseudomonas aeruginosa* Quorum Sensing: Ingles. *Universitas Scientiarum* **2022**, *27* (3), 357–372. https://doi.org/10.11144/Javeriana.SC273.eonc.

[46] Sethupathy, S.; Sathiyamoorthi, E.; Kim, Y.-G.; Lee, J.-H.; Lee, J. Antibiofilm and Antivirulence Properties of Indoles Against *Serratia marcescens*. *Front. Microbiol.* **2020**, *11*, 584812. https://doi.org/10.3389/fmicb.2020.584812.

[47] Sinha, D.; Tiwari, A. K.; Singh, S.; Shukla, G.; Mishra, P.; Chandra, H.; Mishra, A. K. Synthesis, Characterization and Biological Activity of Schiff Base Analogues of Indole-3-Carboxaldehyde. *European Journal of Medicinal Chemistry* **2008**, *43* (1), 160–165. https://doi.org/10.1016/j.ejmech.2007.03.022.

[48] Rajalaxmi, M.; Amsa Devi, V.; Pandian, S. K. In Vitro Evaluation of Indole-3-Carboxaldehyde on *Vibrio parahaemolyticus* Biofilms. *Biologia* **2016**, *71* (3), 247–255. https://doi.org/10.1515/biolog-2016-0043.

[49] Lee, J.-H.; Wood, T. K.; Lee, J. Roles of Indole as an Interspecies and Interkingdom Signaling Molecule. *Trends in Microbiology* **2015**, *23* (11), 707–718. https://doi.org/10.1016/j.tim.2015.08.001.