Title

'Tuning' of ribosome levels mediated by RNase I and hibernating ribosomes

Authors

Atsushi Minami[1,†], Takehito Tanzawa[2,†], Zhuohao Yang[3], Takashi Funatsu[3], Tomohisa

Kuzuyama[1,5], Hideji Yoshida[4], Takayuki Kato[2], Tetsuhiro Ogawa[1,5]

Affiliations

1 Graduate School of Agricultural and Life Sciences, The University of Tokyo, Yayoi,

Bunkyo-ku, Tokyo 113-8657, Japan

2 Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

3 Graduate School of Pharmaceutical Sciences, The University of Tokyo, Bunkyo-ku,

Tokyo 113-0033, Japan

4 Faculty of Medicine, Osaka Medical and Pharmaceutical University, Takatsuki, Osaka

569-0801, Japan

5 Collaborative Research Institute for Innovative Microbiology, The University of Tokyo,

Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

† These authors contributed equally: Atsushi Minami, Takehito Tanzawa

Abstract

Ribosomes consume vast energy to synthesize proteins, so controlling the ribosome abundance is a significant concern for cells. Ribonucleases mediate ribosome degradation in response to stresses, while some ribosomes deactivate translational activity and protect themselves from degradation, called ribosome hibernation. RNase T2 is an endoribonuclease found in almost all organisms, and they are thought to be involved in the degradation of ribosomal RNA. Although it was recently reported that the activity of *Escherichia coli* RNase T2, called RNase I, depends on the environmental conditions, the regulation mechanism remains elusive. Here, we report how rRNA degradation by RNase I is regulated by hibernating ribosomes. Combining the biochemical, cryo-electron microscopy, and single-molecule analyses, we found that hibernating ribosome is an inhibitor by forming a complex with RNase I. Moreover, RNase I does not bind to the translating ribosome, so rRNA is protected. On the other hand, RNase I degrades the rRNA of each subunit dissociated from stalled ribosomes on aberrant mRNA by *trans*-translation. Under stress conditions, and even in the actively growing phase, some ribosomes are stalling or pausing. Although such ribosomes were thought to be recycled after being rescued, our results add a new insight that they are not recycled but degraded. These findings have broad implications for understanding the regulation of ribosome levels, which is critical for cellular homeostasis and response to environmental stresses.

Main

Ribosomes, a platform of translation, are sophisticated molecular machinery. In the living cell, protein synthesis on ribosomes consumes much energy, and translational activity generally depends on the number of ribosomes. Hence, the cell monitors their environmental conditions and synchronizes the intracellular ribosome level. For example, under stress conditions such as nutrient starvation, heat shock, or oxidative stress, ribosome biogenesis and translation initiation are promptly inhibited[1]. In addition, excess ribosomes are degraded[2–5].

On the contrary, the remaining ribosomes are protected from degradation. This protection is essential for resuming protein synthesis upon recovery from stress conditions. One of the protection mechanisms is 'ribosome hibernation'. This mechanism also represses the translational activity of residual ribosomes to avoid energy waste[6–10]. In bacteria, ribosomes typically dimerize and form 100S ribosomes with the involvement of hibernation-advancing factors. Several hibernation factors have been found, such as RMF (Ribosome Modulation Factor), HPF (Hibernation Promoting Factor), and a newly identified factor, Balon, in bacteria[6,11]; ribosome hibernation is also observed in eukaryotes such as yeast and human[12,13].

Ribonucleases play a vital role in halting the activity of ribosomes since one of the essential functions of ribosomes, peptide bond formation, is catalyzed by rRNA. Among many ribonucleases, RNase T2 has been known as one of the most crucial endoribonucleases degrading rRNA in various organisms[14–16]. RNase T2 was first discovered in *Aspergillus oryzae*[17], and later, it was found to exist in almost all organisms. It cleaves single-stranded RNA non-specifically[18], using two histidine residues as catalytic residues. for example, yeast RNase T2, Rny1, degrades rRNA in the process of

autophagy[19,20]. The reaction mechanism consists of two steps: Firstly, the phosphodiester

bond is cleaved by phosphate transfer, which forms an intermediate 2′,3′-cyclic nucleotide

monophosphate (2′,3′-cNMPs). Subsequently, the 3′-phosphate group is formed by

hydrolysis.

While functional analysis of RNase T2 is ongoing in eukaryotes[14–16,21–23], little

is known about it in bacteria. Several reports have been made about *Escherichia coli*

RNase T2, called RNase I, which is a primary ribonuclease that degrades rRNA[24,25].

Moreover, it has been known to bind to the ribosomal 30S subunit[26,27], but the biological

significance of this interaction remains elusive. Recently, Weinert's group showed that

2′,3′-cNMPs, generated by RNase I, accumulate in the cytoplasm and inhibit biofilm

formation. Biofilms are aggregate structures of various microorganisms formed in poor

nutritional conditions and exposure to stress[28]. Under such conditions, the activity of

RNase I must be inhibited for biofilm formation. However, the regulatory mechanisms of

RNase I in response to its environment were utterly unknown. As described above, RNase

I interacts with ribosomes, which are converted into hibernating ribosomes under stress

conditions. Based on these facts, we hypothesized that the hibernating ribosome binds to

RNase I and inhibits its activity to form biofilm.

In this study, we demonstrated our hypothesis by investigating the inhibition

activity of hibernating ribosomes to RNase I *in vivo* through rRNA degradation and

biofilm formation *in vivo* and *in vitro*. Next, to understand the mechanism by which the

hibernating ribosome interacts with RNase I activity, we determined the cryo-EM

structure of hibernating ribosome complexed with RNase I mutant (catalytically inactive,

H55F/H133F) at a nominal resolution of 2.8 Å. Moreover, using single-molecule imaging

we revealed how RNase I discriminates between ribosomes that can bind and those that

cannot. Finally, we verified the state of ribosomes RNase I can degrade by biochemical

analyses. Taken together, the molecular mechanism and physiological significance of

RNase I binding to hibernating ribosomes are discussed, including molecular

evolutionary considerations of RNase I.

93   RNase I interacts with the hibernating ribosome

We first investigated whether the action of the hibernating ribosome on RNase I

inhibits this activity, resulting in biofilm formation in amino acid starvation condition.

Since RaiA, another hibernating factor in *E. coli*, is homologous with HPF, the binding

to ribosomes is competitive. Therefore, the RaiA-deficient strain (Δ*raiA*) highly

accumulates 100S ribosomes in the cytoplasm[29,30] (Extended Data Fig. 1a). These wild-

type, RNase I-deficient (Δ*rna*), and Δ*raiA* strains were grown under an amino acid

starvation condition for 4 h. The total RNA was then prepared and electrophoresed to

assess the extent of RNA degradation. Following the amino acid starvation, the rRNA of

the wild-type strain was the most degraded, approximately half that of the Δ*raiA* strain,

suggesting that the RNase I activity is down-regulated in the Δ*raiA* strain. In the wild-

type and Δ*raiA* strains, 23S rRNA was more susceptible to RNase I than 16S rRNA. On

the other hand, rRNA was not degraded at all; instead, it increased in the Δ*rna* strain,

which is consistent with previous reports[24,31–33] (Fig. 1a and Extended Data Fig. 1b). Next,

the biofilm-forming ability of these strains was quantified. We found that the Δ*raiA* strain,

like the Δ*rna* strain, showed a higher biofilm-forming ability than the wild-type strain

(Fig. 1b). Moreover, the high biofilm-forming ability observed in the Δ*raiA* strain was

reduced when RNase I was overexpressed in Δ*raiA* strain, showing that the high biofilm

formation observed in Δ*raiA* strain is due to the inhibition of the RNase I activity by

hibernating ribosomes.

We analyzed the interaction between RNase I and ribosomes to verify the result

in Fig. 1a and b. The cytoplasmic lysate was obtained from the spheroplast of *E. coli* to

exclude RNase I localized in the periplasm. After collecting the cytoplasmic lysate,

ribosomes were fractionated by ultrafiltration, followed by western blotting. As a result,

crude ribosome fraction extracted from the Δ*raiA* strain contained about twice as much

RNase I than the wild-type strain (Extended Data Fig. 2), showing that hibernating

ribosomes interact more readily with RNase I than 70S ribosomes. Furthermore, we

investigated the ability of hibernating ribosomes to inhibit RNase I activity. Δ*raiA*Δ*rna*

and Δ*raiA*Δ*hpf*Δ*rmf*Δ*rna* strains were constructed from the Δ*rna* strain to eliminate

RNase I in the cells. The Δ*raiA*Δ*rna* strain highly accumulates hibernating ribosomes in

the cells, whereas the Δ*raiA*Δ*hpf*Δ*rmf*Δ*rna* strain lacks all hibernating factors and,

therefore, does not make hibernating ribosomes. Crude ribosomes were extracted and

added to RNase I, and the RNase I activity was measured as the previous report[33]. We

found that crude ribosomes from the Δ*raiA*Δ*rna* strain inhibited RNase I activity more

strongly than those from the Δ*raiA*Δ*hpf*Δ*rmf*Δ*rna* strain (Fig. 1c). Together, we

confirmed that hibernating ribosomes interact with RNase I to inhibit its activity.

Fig. 1

Fig. 1 | *E. coli* strain accumulating hibernating ribosomes exhibits suppression of rRNA degradation by RNase I

a, Total RNAs were extracted from indicated strains cultured before and after 4 hours of amino acid starvation, and 16S and 23S rRNA were analyzed using a Bioanalyzer. **b,** (left) The biofilm formation of *E. coli* wild-type, Δ*rna,* and Δ*raiA* strains was quantified by the crystal violet staining. (right) RNase I was overexpressed in the Δ*raiA* strain, and the biofilm formation was quantified. **c,** RNase I was added to ribosomes extracted from each strain, and the enzymatic activity of free RNase I was quantified. The experiments in **b** and **c** were five and three independent experiments, respectively, and the data are shown as the means ± standard error. Asterisks indicate significant differences (*:$P < 0.05$; **:$P < 0.01$, by Welch's t-test).

Structure of hibernating ribosome and RNase I complex

To unveil the mechanism of how RNase I recognizes hibernating ribosome, we reconstituted the complex of RNase I H55F/H133F double mutant, which lacks the catalytic activity almost completely, bound to the *E. coli* dimerized 100S ribosome. The purified RNase I mutant was added to the 100S hibernating ribosomes isolated from *E. coli* W3110 cells in the stationary phase, and the mixture was applied to the EM-grids for cryo-EM observation. We collected the cryo-EM movies and operated the single particle analysis of 100S•RNase complex with two or three-dimensional classification (Extended Data Fig. 3). Because it has been reported that the 100S ribosome is easily split into two 70S ribosomes at the low concentration and that it also tends to exhibit the preferred orientation on the EM-grid[30,34], we picked and extracted the RNase I-bound 70S ribosome particles from micrographs, and the density map of 70S•RNase I, in which the maps of both hibernation factors HPF and RMF were visible, were selected and refined. The first complex chosen provided 351,633 particles, yielding a density map at a nominal resolution of 2.4 Å. However, as the density of RMF was not fully occupied by this complex, the 3D classification was repeated to enrich the density, finally yielding a refined map to a nominal resolution of 2.8 Å consisting of 50,955 particles (Fig. 2a, Extended Data Table 1, Extended Data Figs. 3, 4a and 4b).

The RNase I-bound hibernating complex structure revealed that the RNase I binds to the surface of the head domain of the 30S subunit (Fig. 2a and Extended Data Fig. 4c). As well as previous structural studies of hibernating ribosome complex, the HPF interacts with the channel between the 30S head and body domains. At the same time, the RMF does the anti-Shine Dalgarno (anti-SD) region between the 30S head and platform domains, respectively[30,34]. The binding site of RNase I is located on the opposite side of

the binding regions of HPF and RMF, hence RNase I and the hibernation factors do not

interfere with each other (Fig. 2a). Overall, the shape of this RNase I is almost the same

as previous crystal structure[35]. The *E. coli* RNase I has a characteristic arch-like

architecture, with the active residues, H55 and H133, are located at the "keystone" of the

arch[35], and both ends of the arch interact with the ribosome (Fig. 2b). The active center

of RNase I is not in physical contact with the ribosome and is sufficiently far away from

the 16S rRNA, indicating that rRNA escapes degradation by RNase I (Fig. 2b). In addition

to helix 41 (h41) of the 16S rRNA, suggested in previous report[33], the ribosomal protein

uS14 interacts with the bottoms of the arch in our structure. Estimation of the binding

interfaces and areas obtained by PDBe-PISA[36] and APBS[37] suggest that the RNase I and

its binding partners are contacted mainly by electrostatic and hydrogen bonds and partly

by hydrophobic interactions, and that the RNase I binds primarily to the h41 and

secondarily to the uS14, through shape complementarity (Fig. 2c, Extended Data Fig. 5

and Extended Data Table 2).

The stem-loop A1257–G1278 of h41 binds by bridging both arch bottoms of the

RNase I (Fig. 2b and Extended Data Fig. 5). The polar amino acid residues R31, R33,

Q198, and N233 of the RNase I form hydrogen bonds with the nucleotides G1272–A1274

in the h41 stem region. The binding residues of RNase I do not recognize the groove of

the mini-helix formed by G1272–A1274 and their base-pair partners, but take advantage

of the arch-shaped features of RNase I to bind only one bank of the groove (Fig. 2c and

2d [h-(i)]). In the other binding region, the RNase I recognizes A1257 and G1278 at the

kink of the h41 stem. This kink does not give both bases any base-pairing partners and

leaves them free, allowing the RNase I to stabilize on the stem-loop. The R83 mainly

forms hydrogen bonds with both A1257 and G1278 and exhibits stacking interactions

with the base of A1257 (Fig. 2c, 2d [h-(ii)]). Interestingly, compared to a typical *E. coli*

70S ribosome[38], RNase I binding causes the A1257 to rotate approximately 100 degrees

to the h41 stem-loop side to avoid collision with the R74 (Fig. 2e). This conformation

forms a pseudo-major groove consisting of A1257–G1260 and A1274–G1278, where

helix α4 of the RNase I is placed in a shape complementary manner (Fig. 2d [h-(ii)] and

Extended Data Fig. 6). In detail, the F78 has a π-stacking interaction with the base of

A1257 at the other side of R83, and the R73 and R74 make hydrogen bonds with its main

chain (Fig. 2d [h-(ii)]). The conformational change of A1257 by the recruiting to RNase

I promotes the loss of interaction between A1257 and F21 of uS14 (Fig. 2e), and the cleft

created by this local dissociation between h41 and uS14 is filled by the R74 of RNase I

(Fig. 2d [s-(i)]). Thereby, the R74 recognizes the F21 instead of the A1257, also the loop

containing G69–D71 of RNase I contacts the residues D18–A22, located in the bent of

the N-terminal helix of uS14 (Fig. 2b, 2d [s-(i)]). As well as h41, the N-terminal helix

also binds to bridge both arches of the RNase I, with another binding site on the opposite

side of the above (Fig. 2b and Extended Data Fig. 5). The residues P229–P231 in the

Pro/Gly-rich long loop between β5 and β6 of RNase I exhibit electrostatic interactions

with the residues I30–N35 of uS14, with hydrogen bond between the H230 and D33 (Fig.

2c, 2d [s-(ii)]). Collectively, our structure of the RNase I•hibernating ribosome complex

provides an understanding of the unique interaction mechanism, allowing hibernating

ribosomes to protect themselves from degradation by RNase I.

Fig. 2 | Structure of the RNase I-bound hibernating ribosome complex.
a, Overall view of the RNase I-bound hibernating ribosome showing the structure
of the *E. coli* 70S ribosome in complex with RNase I (dark blue), HPF (green),
and RMF (orange). The 50S subunit is shown in light blue and the 30S subunit in
light yellow (left panel). Overview from a different angle with the 50S subunit
omitted for clarity (right panel). **b,** Close-up view of the interaction between the
RNase I and the h41 (light green) of the 16S rRNA or the ribosomal protein uS14
(light pink), viewed from two different angles. The black dashed rectangles
indicate the estimated binding interfaces between RNase I and h41 or uS14
calculated using PDBe-PISA. The h41 and uS14 interact at both ends of the arch
architecture of the RNase I, respectively, and these are shown by h-(i) and h-(ii)
for h41 and s-(i) and s-(ii) for uS14. **c,** The electrostatic potential surface of the
RNase I, h41, and uS14 as a single mass is shown from the same direction as **b**.
The mass is split into the RNase I and h41•uS14, each viewed from two different
perspectives. Both positive, negative, and neutral electrostatic surface potentials
are shown in blue, red, and white. The yellow dashed rectangle illustrates each
binding region. **d,** Detailed views of the major binding regions of RNase I to h41
(h-(i) and h-(ii)) or uS14 (s-(i) and s-(ii)). Chains of RNase I, h41, and uS14 are
colored the same as **b**. Stick models represent amino acid residues and
nucleotides involved in these bindings, and the N, O, and P-atom are colored

blue, red, and orange, respectively. Dashed lines indicate hydrogen bonds. **e,**
Structural comparison of the A1257 of h41 and the F21 of uS14 by superposing
the stem-loop of h41 between RNase I bound hibernating complex and a typical
70S ribosome (PDB 8SYL). The 8SYL ribosome is colored by whitish blue and
the ternary complex is shown in the same as **b**. The base of A1257 (PDB 8SYL)
and the side chains of R74 (RNase I) and F21 (our model and PDB 8SYL) are
represented by stick and sphere models. The curving arrow illustrates the rotation
of the A1257 base.
240
241

RNase I does not interact with the translating ribosome

Our *in vivo* binding assay showed that RNase I interacts with hibernating ribosomes more tightly than 70S ribosomes, and the cryo-EM structure revealed the interaction mechanism between RNase I and ribosomes. However, the *in vitro* binding assay of hibernating ribosomes, reconstituted by adding RMF and HPF (hereafter referred to as "HFs") and 70S ribosomes almost comparably bound to RNase I mutant (Fig. 3a, b). The binding affinity of RNase I to ribosomes decreased in the exponential phase (Extended Data Fig. 2), where most ribosomes are engaged in translation. Given these results, we assumed that RNase I cannot interact with the translating ribosomes. We then performed the binding assay between RNase I and translating ribosomes *in vitro* using the PURE system[39], the reconstituted translation system derived from *E. coli*. Translating ribosomes had a reduced ability to interact with RNase I compared to the non-translating 70S ribosomes or reconstituted hibernating ribosomes (Fig. 3c).

It is known that not all ribosomes are engaged in translation; instead, limited ones synthesize proteins in an *in vitro* translation such as the PURE system. We considered that this fact might influence our experimental results. Therefore, we performed a single-molecule imaging method to solve this issue. In recent years, several groups have developed real-time imaging techniques of translation at the single-molecule level[40–43]. In the experimental system, multi-epitope protein tags are translated on the ribosome, and fluorescently labeled antibodies recognize and bind to the tags, enabling the translational visualization of ribosomes. However, almost all single-molecule imaging methods are performed in mammalian cells. To our knowledge, only one *in vitro* experiment has been reported using yeast lysate.

Here, we attempted to perform antibody-based single-molecule translation

imaging *in vitro* using the PURE system. This allowed real-time translation and binding

of RNase I to the ribosome at the single-molecule level. The experimental design is

summarized in Fig. 3d. A single ribosome particle was defined as a spot where mRNA-

derived fluorescence was observed. Here, a fluorescently labeled nanobody, a small

single peptide recognizing multi-epitope protein tags, is used instead of whole antibodies

or Fab fragments. Nanobody-derived fluorescence appeared in some ribosome particles,

indicating that translation elongation was in progress ("translating region"). In other

regions, ribosomes remain arrested as nanobody-derived fluorescence was not observed

("arrested region"). Instead, RNase I mutant (H55F/H133F)-derived fluorescence was

observed, and fluorescence of nanobody and RNase I were never detected in the same

spot (Fig. 3e). Together, we could observe that RNase I does not interact with translating

ribosomes as a single-molecule level. Notably, spots of RNase I-derived fluorescence

observed in the arrested region were not retained long; most of RNase I was released

within 1 or 2 min (Fig. 3f), indicating that RNase I may be unable to maintain a stable

interaction with ribosome.

Fig. 3

Fig. 3 | RNase I binds to 70S and 100S ribosomes but does not bind to translating 70S ribosomes.

a, HFs were added to 70S ribosomes, and 70S and 100S ribosomes were detected using sucrose density gradient centrifugation. **b,** RNase I (H55F/H133F) was added to the ribosome with or without hibernation factors. Mixtures were fractionated by ultrafiltration with a 300 kDa cut-off. RNase I remained in the large fraction and was detected by western blotting. "T" indicates the total mixtures, and "L" indicates the large fraction, while "S" indicates the small fraction. **c,** RNase I was added during translation using an *in vitro* reconstituted translation system. Mixtures were fractionated as described in **(b)**, and the enzymatic activity

of RNase I that did not bind to ribosomes was detected in Fig. **1(c)**. The enzymatic
activities of RNase I were compared between when RNase I was added to
translating ribosomes (black, solid line) and to ribosomes with added HFs (blue,
solid line) (left) or to not translating ribosomes (black, dashed line) (right). Data
are presented as the means ± standard error of 3 independent experiments. **d,**
(upper) Schematic depicting the experiments performed in **(e)** and **(f)**. The
template mRNA is marked by the oligo DNA probe that hybridizes to repeated
sequences located in 3′ end (green). At the start of the observation, the mRNA
and the ribosome complex is fixed on the glass base via HaloTag. The ribosome
is arrested on the mRNA due to the SecM arrest sequence (AS), so labeled
RNase I mutant can bind to the ribosome ("arrested"). After the arrested ribosome
is released, the nascent polypeptide chain is labeled by nanobodies (blue) that
bind to peptide epitopes, MoonTag ("translating"). (lower) Example co-
localization of nanobodies-bound nascent chain and mRNA. **e,** The RNase I-
derived fluorescence intensity distribution at the coordinates where translation
spots were observed and the overall fluorescence intensity of the RNase I-
derived spots. The vertical axis is the relative value with the mean of the overall
fluorescence intensity of the entire RNase I-derived fluorescence spot (right) as
1. **f,** RNase I-derived fluorescence spots localized in the "arrested" regions were
detected, and the number of consecutive frames detected was counted. The
frame interval is approximately 1 minute.

RNase I degrades rRNA within dissociated ribosomal subunits derived from *trans*-translation

Finally, we considered how RNase I cleaves rRNA to produce 2′,3′-cNMPs in cells. The previous report showed that 2′,3′-cNMPs were detected during the actively growing phase[44]. However, it is known that the 70S ribosome is highly resistant to RNases[45]. Indeed, RNase I hardly degraded the rRNA in equimolar amounts with the ribosomes *in vitro* (Fig. 4b). This low susceptibility of rRNA to RNases is due to ribosomal proteins wrapping around the rRNA, which results in protecting rRNA from degradation[46]. As shown above, our results indicate that the target of RNase I is non-translating ribosomes; however, RNase I cannot attack rRNAs within 70S ribosomes. Then, how does RNase I degrade the rRNA of non-translating ribosomes? Compared to our complex structure, we found that the RNase I bound to the ribosome spatially shares a region with a large pseudoknot ring (PK ring) of transfer-messenger RNA (tmRNA) (Fig. 4a). In bacteria, *trans*-translation is one of the primary 'rescue system' for ribosomes stalled on mRNA, mediated by tmRNA and small protein B (SmpB). The tmRNA is composed of a PK ring, a tRNA-like domain (TLD), and a mRNA-like domain (MLD), and it enters the vacant A-site of ribosomes stalled at the 3′ ends of mRNA, which lacks a stop codon. Then, a degradation tag, encoded by the MLD, was added to the C-terminus of a nascent polypeptide chain, thereby degrading the synthesized polypeptide. In addition, the stalled ribosomes are split into the 30S and 50S subunits, resulting in release from the aberrant mRNA[47]. Taken together, we hypothesized that RNase I could cooperate with tmRNA to degrade rRNA within the split ribosomal subunits.

To validate the hypothesis that rRNAs within split ribosomal subunits become susceptible to RNase I, we separated the ribosome into the 30S and 50S subunits[48] and

then incubated with RNase I. As expected, rRNAs of subunits were more degraded by

RNase I than that of 70S ribosomes (Fig. 4b). Even in the actively growing phase, a

certain number of ribosomes are stalled on mRNA, e.g., due to aberrant mRNA or

ribosomes[49], resulting in the generation of 2′,3′-cNMPs. Notably, the rRNAs were

fragmented by the degradation, suggesting that the first cleavage by RNase I may have

occurred in the helix 69 (h69) of 23S rRNA[50]. If this assumption is correct, a fragment of

approximately 1900 nt should accumulate, roughly consistent with the fragment length

visualized by electrophoresis shown in Fig. 4b.

Amino acid starvation also induces intracellular accumulation of 2′,3′-cNMPs[44];

that is, RNase I degrades rRNA. However, ribosomes are engaged in translation, as cells

proliferate even under such condition, albeit at a reduced growth rate. During amino acid

starvation, ribosomes are generally prone to stall and resume translation repeatedly[51,52].

Swivels in the head domain, where RNase I has been found to bind, are important in

translation. Therefore, we surmised that the binding of RNase I to the head domain

interferes with translation. Thus, we measured the translational activity of RNase I-bound

ribosomes. Ribosomes and RNase I mutant (H55F/H133F) were mixed *in vitro*, and

translation was initiated using the PURE system (Fig. 4c). As a result, the translational

activity was critically reduced when RNase I was added before translation initiation.

Although inhibition of translation initiation by RNase I seems to be beneficial for *trans*-

translation as tmRNA acts on ribosomes stalled on mRNA, our complex structure

suggests that RNase I and tmRNA spatially share a binding region as described above

(Fig. 4a). We measured *in vivo trans*-translation activities in the presence or absence of

RNase I to verify the result obtained from the cryo-EM analysis. Here, *trans*-translation

was induced by expressing a gene encoding mRNA lacking the stop codon of GFP and

transcribing a tmRNA mutant that adds a non-degradation tag to the nascent polypeptides.

Then, *trans*-translation activity can be quantified by measuring the GFP fluorescence.

Consequently, *trans*-translation activity increased in the RNase I-deficient strain,

indicating that binding sites of RNase I and tmRNA are overlapped (Fig. 4d). As

mentioned, subunit dissociation by *trans*-translation is required for the rRNA degradation

by RNase I. However, if RNase I continues to interact with the head domain stably, the

efficiency of tmRNA entering the A-site will severely decrease, and RNase I will never

produce 2′,3′-cNMPs. Considering that RNase I efficiently degrades rRNAs under amino

acid starvation condition (Fig. 1a, Extended data Fig. 1b), the binding of RNase I to the

head domain is not so stable, which is consistent with the results arose from the single-

molecule imaging that RNase I binding with the ribosome is transient (Fig. 3f). Taken

together, we concluded that this transient binding allows tmRNA to enter the ribosomal

A-site, leading to RNase I degrading rRNA in the subunit after the dissociation by *trans*-

translation.

Fig. 4

a, The hibernating ribosome•RNase I complex and the trans-translating 70S•tmRNA complex (PDB 6Q98) were superposed by 70S, based on the interaction sites shown in Fig. 2. The PK ring of tmRNA extends outside the head domain of the 30S subunit, which spatially overlaps with RNase I. **b,** The 30S/50S subunits dissociated by adding a dissociation buffer to 70S ribosomes, and 70S ribosomes were prepared. Total RNAs were extracted from each of them after incubation with or without RNase I, and electrophoresed using a formalin-denaturing gel. "70" indicates rRNAs from intact 70S ribosomes, while "50+30" indicates rRNAs from dissociated subunits. **c,** RNase I mutant was added at the beginning of *in vitro* translation, and the translational activity was measured. **d,** The *trans*-translation activities in the Δ*ssrA* strain and the Δ*ssrA*Δ*rna* strain carrying the two plasmids (see Methods) were compared. Data are presented as the means ± standard error of 3 independent experiments. Asterisks indicate significant differences (*:P < 0.05, by Welch's t-test).

Fig. 4 | RNase I degrades rRNA from subunits dissociated by *trans*-translation.
a, The hibernating ribosome•RNase I complex and the trans-translating 70S•tmRNA complex (PDB 6Q98) were superposed by 70S, based on the interaction sites shown in Fig. 2. The PK ring of tmRNA extends outside the head domain of the 30S subunit, which spatially overlaps with RNase I. **b,** The 30S/50S subunits dissociated by adding a dissociation buffer to 70S ribosomes, and 70S ribosomes were prepared. Total RNAs were extracted from each of them after incubation with or without RNase I, and electrophoresed using a formalin-denaturing gel. "70" indicates rRNAs from intact 70S ribosomes, while "50+30" indicates rRNAs from dissociated subunits. **c,** RNase I mutant was added at the beginning of *in vitro* translation, and the translational activity was measured. **d,** The *trans*-translation activities in the Δ*ssrA* strain and the Δ*ssrA*Δ*rna* strain carrying the two plasmids (see Methods) were compared. Data are presented as the means ± standard error of 3 independent experiments. Asterisks indicate significant differences (*:P < 0.05, by Welch's t-test).

Discussion

Ribosome degradation and protection are central concerns for all living things, especially under stress conditions. In *E. coli*, RNase I is the most critical ribonuclease in the mechanism for the control of intracellular ribosome levels. Although RNase I was discovered over 60 years ago, its regulatory mechanism has been poorly understood. This study shows the specific interaction between RNase I and hibernating ribosomes to inhibit RNase I activity under stress conditions. These results explain the biological significance of the past findings on why RNase I binds to the 30S subunit[26,27]. As mentioned, hibernating ribosomes protect themselves from the attack of RNase I. However, hibernating ribosomes are gradually degraded after prolonged exposure to stresses[29]. Therefore, the biological significance of interactions between RNase I and hibernating ribosomes may be not only in the inhibition of RNase I activity but also in rRNA degradation under prolonged stresses. Fessler *et al*. demonstrated that rRNA is degraded by several starvation conditions and suggested that phosphate and nucleotides are recycled[53]; this is consistent with our results that RNase I degrades hibernating ribosomes when cells are exposed to long-term stresses (Extended Data Fig. 1b). In addition, our result indicated that RNase I degrades rRNA to several fragments (Fig. 4b). It has been suggested that these rRNA fragments are ultimately degraded by the exoribonucleases RNase R or PNPase[45], whose product may be utilized to survive prolonged nutrient starvation.

Our study indicated that rRNA degradation by RNase I, responsible for the first step for degradation (Fig. 1a, Extended Data Fig. 1b), targets the stalled ribosome on the aberrant mRNA. Such stalled ribosomes are possible to induce ribosome collision[54]. SmrB in *E. coli* was recently found to cleave mRNA upstream of collided ribosomes,

enabling *trans*-translation[55]. Furthermore, ribosome dissociation into ribosomal subunits,

commonly referred to as ribosome rescue, is required for the rRNA degradation by RNase

I. This study provides an opportunity to reconsider whether ribosomes are 'recycled' after

trans-translation. To our knowledge, no reports show that ribosomes rescued from mRNA

by *trans*-translation are recycled. On the other hand, a mechanism called nonfunctional

rRNA decay (NRD)[56,57], which is present not in bacteria but in eukaryotes, recognizes

and degrades stalled ribosomes during the translation initiation or elongation. Our

research proposes a model in which the rescued ribosomes are degraded rather than

recycled, similar to NRD. Moreover, taking into account that RNase I can only bind to

stalled ribosomes and that RNase I-bound ribosomes cannot initiate translation, we

further extend the existing molecular mechanism of ribosome rescue (Fig. 5b). Even

during the exponential phase, some ribosomes are stalled on mRNA for various reasons,

such as damage to rRNA or the absence of stop codons on mRNA. Our results indicate

that RNase I needs to leave the ribosome for tmRNA to enter the ribosomal A-site. One

possible reason why RNase I binds to the 30S ribosomal subunit is that it gathers near

arrested ribosomes, which are dissociated to each subunit by *trans*-translation, allowing

RNase I to degrade rRNA in dissociated subunits efficiently. This assumption seems

reasonable given the intracellular ratio of ribosomes to RNase I. The location where

RNase I binds may be coincidentally suitable for destabilizing RNase I binding by turning

the head domain, although this could not be proven experimentally. Rare codons on the

mRNA reduce the translation elongation rate while it does not stall the ribosome

completely. It may be possible that RNase I binds to these ribosomes, which results in the

inhibition of translation initiation as the presence of a rare codon induces *trans*-

translation[58].

Fig. 5

Fig. 5 | Model of the interaction between RNase I and ribosomes.
a, When the ribosome is translating, the 30S head domain rotates relatively to
the body, called "swivel," so the interacting site of RNase I is unstable (left). On
the other hand, the domain in the stalled ribosome (center) or hibernating
ribosome (right) is stable, so RNase I easily binds to the ribosome. **b,** Extended
Model of ribosome recycling and RNase I-mediated rRNA degradation. RNase I
binds to abnormally arrested ribosomes that occur at a specific rate during

translation **(i)**. Binding at this point is also thought to play a role in completely
stopping translation by binding to ribosomes that are prone to stall. However, the
binding affinity of RNase I to the arrested ribosome is not so high, and the binding
and dissociation are thought to be repeated **(ii)**. The stably stopped ribosome
may return to the normal cycle of translation **(iii)**, but with the help of some RNase,
tmRNA can act on the arrested ribosome (**iv**), and the ribosome is rescued and
split **(v)**. Those split ribosomal subunits were thought to be reused for translation,
but RNase I can degrade the dissociated ribosome-derived rRNA after *trans*-
translation **(vi)**.
460

RNase I localizes mainly in the periplasm, but the biological role of the

localization is still unknown. One possible physiological significance of RNase I

accumulating in the periplasm lies in oxidative folding, as RNase I has four disulfide

bonds[59,60]. As described, RNase I degrades rRNA in the cytoplasm (Fig. 1a and Extended

Data Fig. 1b). However, the mechanism which RNase I, generally localized in the

periplasm, enters the cytoplasm is unclear. Previous reports discussed the entrance of

RNase I in the cytoplasm by such as changes in cell membrane permeability and

involvement of F factor-derived toxin, SrnB[61,62]. Further verification is needed to

determine the cause of RNase I influx into the cytoplasm.

One of the most remarkable findings in the present study is the elucidation of

where and how RNase I interacts with the hibernating ribosome. Our cryo-EM structure

shows that the RNase I binding region is located on the head domain of the 30S subunit,

which constantly fluctuates during translation due to swivel and rotation. In contrast, the

head domain becomes stable when the ribosome stops translating or transits the

hibernation state (Fig. 5a). In particular, the early cryo-EM study showed that h41, to

which RNase I binds, rotates approximately 10 degrees during EF-G•GTP-mediated

ratchet rotation[63]. As a result, translation may reduce the structural and thermodynamic

stabilities of the head domain, destabilizing the interaction with RNase I. Moreover, the

molecular dynamic simulation of the translation initiation complex predicted that the

region of C1254-C1284 of h41 would undergo conformational changes[64]. Together with

our results (Fig. 4c), the fluctuation of the 30S head may be suppressed in RNase I-bound

ribosomes, resulting in the inhibition of translation. Moreover, the RNase I interacts with

the uS14, which may be vital to unlocking the diverse functionality and evolutionary

significance of the uS14 protein. Despite the gene for uS14 being essential, the function

of uS14 in translation remains elusive. The uS14 protein is classified into three groups

according to the presence or absence of a zinc-binding motif, and *E. coli* uS14 belongs to

the C- long type[65]. Compared to the crystal structure of uS14 in *Thermus thermophilus*

70S ribosome (PDB 4V8H[34]) , which belongs to the C+ type, *E. coli* uS14 has a distinctive

domain with an internal insertion containing the contact site to RNase I (Extended Data

Fig. 7a). No homologous protein of RNase I has been found in *T. thermophilus*,

suggesting that *E. coli* uS14 may have evolved independently, along with its interaction

with RNase I. In addition, C+/- paralogue pairs for uS14 are conserved in some bacteria,

and their functionality differences have been reported[65–67]. These results support that

uS14 is a susceptible region in the evolutionary process for gaining diverse functions.

RNase I shows the diversity of their sequences and structures among species.

The RNase T2 family exists in a wide range of species, as mentioned above. We focused

on *E. coli* RNase I in this study, and previously, we reported on *Saccharomyces cerevisiae*

RNase T2, Rny1[20]. In that study, molecular phylogenetic analysis of RNase T2 shows

that fungal RNase T2 evolved independently from other species in the early stage of

evolution. Moreover, it was also suggested that *E. coli* RNase I also evolved apart from

mammals and plants. We thus compared the structure of *E. coli* RNase I with that of

human RNase T2 (PDB 3T0O[68]). Then, we found a region-specific to *E. coli* RNase I,

which is one of the binding sites to the ribosome (Extended Data Fig. 7b). This may mean

that during the evolution of bacteria, RNase I acquired its binding function to the

ribosome. Together with the diversity of uS14, RNase I and ribosomes have evolved to

'checkmate' each other by the interaction. In particular, the structure of ribosomes is often

highlighted for its commonality between species, but many areas differ between species;

ribosomes have likely optimized their structures in response to their relationship with

trans-acting diverse protein factors, such as RNase I and hibernation factors.

Recently, it was reported that the RNase R binds to the 30S platform to proceed

with degrading the anti-SD and the helix 44 at the decoding center using cryo-EM single

particle analysis[69]. Compared with our structure of the RNase I•hibernating ribosome

complex, the binding region of RNase R is entirely different from that of RNase I. Rather,

it overlaps in that of the hibernation factor RMF (Extended Data Fig. 7c), indicating that

hibernating ribosomes also inhibit the activity of RNase R in *E. coli*[9]. Our research

revealed that the inhibitory mechanism of RNase I by hibernating ribosomes is also quite

different from that of RNase R, in which hibernating ribosomes bind directly to RNase I,

thereby keeping it away from its active site. This suggests that hibernating ribosomes

have acquired multiple protection mechanisms to resist the attacks of diverse intracellular

ribonucleases.

Methods

Construction of strains and plasmids

The strains and plasmids used in this study are listed in Supplementary Table S1. The RNase I, RaiA, RMF, or HPF-deficient strains (Δ*rna*, Δ*raiA*, Δ*rmf*, or Δ*hpf*, respectively) were obtained from the Keio Collection, which was provided by the National BioResource Project (NBRP) at the National Institute of Genetics, Japan[70]. The kanamycin-resistance gene of strains from the Keio Collection was eliminated by homologous recombination using plasmid 707-FLPe (Gene Bridges). The double mutant strain (Δ*raiA*Δ*rna*) or quadruple mutant strain (Δ*raiA*Δ*hpf*Δ*rmf*Δ*rna*) were produced by homologous recombination using the Red/ET system.

The open reading frame (ORF) and the promotor region of the *rna* gene were PCR-amplified from *E. coli* genomic DNA extracted from BW25113 wild-type strain and inserted into the pUC18 vector with the seamless cloning method (In-Fusion HD Cloning Kit, TaKaRa Bio). For the expression of recombinant RNase I wild type and mutants, RMF, HPF, and the nanobody of MoonTag, PCR-amplified ORFs were inserted into the pET24d(+) vector with the same method. The plasmid coding the nanobody of MoonTag was kindly gifted by Dr. Wataru Fujii (The University of Tokyo, Japan). The plasmid pET-*gfp* was created in this study to measure the *in vitro* translation activity. The plasmid pMW-*ssrA*^{DD} expressing mutant tmRNA was kindly gifted by Dr. Hyota Himeno (Hirosaki University, Japan). The plasmid harboring the *trans*-translation inducing sequence was created as follows: First, the ampicillin resistance cassette of pAcGFP1 plasmid purchased from TaKaRa Bio was replaced with the kanamycin resistance cassette by the seamless cloning method. Next, the stop codon of the ORF was replaced with the *trpA* terminator sequence using the same method. The resulting plasmid was designated

pAcGFP-woSTOP-*trpAt*. The template sequence for the single-molecule imaging was

created as follows: First, the template 12xMoonTag-12xSunTag-kif18b-24xPP7 was

purchased from Addgene (#128606)[71]. Second, the HaloTag-SecM sequence was inserted

into the spacer region between 12xMoonTag and 12xSunTag by the seamless cloning

method (NEBuilder HiFi DNA Assembly, NEB). Third, the plasmid was digested with

NcoI and ApeI, and inserted into pET24d(+). Subsequently, PCR-amplified HaloTag-

L26-SecM[133-170] sequence[72] was inserted between the start codon and 12xMoonTag of

the plasmid.

Preparation of RNase I, hibernation factors

E. coli BLR(DE3) strain transformed with the expression plasmids were grown

in the lysogeny broth (L-broth) supplemented with 30 μg/mL of kanamycin. When an

OD_{660} reached 0.5, isopropyl β-D-1-thiogalactopyranoside (IPTG) was added to the

culture at a final condition of 1 mM to induce the expression of RNase I wild type and

mutants, RMF, and HPF, and the culture was further incubated for 3 h at 37 °C. The

purification of RNase I wild type and mutants was performed as described[20] with slight

modifications. The cells were harvested by centrifugation, and the cell pellet was

resuspended in 20 mL of lysis/wash buffer (20 mM Tris-HCl (pH 7.5), 500 mM NaCl,

and 20 mM imidazole (pH 8.0)), followed by sonication on ice to lyse the cells. Cell

debris was removed by centrifugation at 17,000 rpm for 20 min. The supernatant was

applied to the Ni-NTA column, and the resin was washed with 20 mL of lysis/wash buffer.

Then, RNase I was eluted in 5 mL of elution buffer (20 mM Tris-HCl (pH 7.5), 500 mM

NaCl, and 250 mM imidazole (pH 8.0)). The purified RNase I was concentrated using a

Vivaspin 20 device (10,000 MWCO; Sartorius) and stored in the storage buffer (20 mM

Tris-HCl (pH 7.5), 50 mM NaCl)[60]. For the purification of RMF and HPF protein, lysis

buffer (20 mM Tris-HCl (pH 7.6), 100 mM NH$_4$Cl, 10 mM MgCl$_2$, 1 mM β-

mercaptoethanol), wash buffer (20 mM Tris-HCl (pH 7.6), 1 M NH$_4$Cl, 10 mM MgCl$_2$, 1

mM β-mercapthoethanol), elution buffer (20 mM Tris-HCl (pH 7.6), 100 mM NH$_4$Cl, 10

mM MgCl$_2$, 2 M imidazole (pH 8.0), 1 mM β-mercapthoethanol), and storage buffer (20

mM Tris-HCl (pH 7.6), 100 mM NH$_4$Cl, 10 mM MgCl$_2$)[34] was used instead of each buffer

described above. For the purification of RMF protein, an equal volume of 2 M LiCl was

added to the supernatant after sonication and stirred overnight at 4 °C, followed by

elution[73].

Isolation of ribosomes and sucrose density gradient centrifugation analyses

E. coli 70S ribosomes were prepared from the *ΔraiAΔhpfΔrmfΔrna* strain as

described[74]. *E. coli* 100S ribosomes for the sucrose density gradient centrifugation

(SDGC) analysis, the cell pellets were resuspended in 1 mL of the ribosome gradient

buffer (20 mM Tris-HCl (pH 7.5), 60 mM NH$_4$Cl, 7.5 mM Mg(OAc)$_2$, 0.5 mM EDTA, 6

mM β-mercaptoethanol)[75] and disrupted by shaking for 5 min with 300 mg of zirconium

silicate beads in a homogenizer (ShakeMaster Neo, Bio Medical Science Inc.). The

sample was then centrifuged for 10 min at 18,500 g and 4 °C, and the resultant supernatant

was collected. The absorbance at 260 nm (A$_{260}$) for lysates was determined by Nanodrop,

and 300 μL of normalized lysate was loaded on a 10% to 40 % sucrose gradient in the

ribosome gradient buffer. After 3 h of centrifugation at 35,000 rpm and 4 °C in a Beckman

SW41 rotor, the sample was collected with a BioComp gradient station monitoring

continuous absorbance at 254 nm. The *in vitro* reconstituted hibernating ribosomes were

prepared by incubating 23 pmol of 70S ribosomes with 240 pmol of RMF and 830 pmol

of HPF in the reaction buffer (20 mM Tris-HCl (pH 8.0), 100 mM NH$_4$Cl, 10 mM MgCl$_2$)

for 30 min and 37 °C. The SDGC analysis profiled the resulting mixture as above.

Extraction of total RNA from E. coli cells

E. coli each strain was cultivated in L-broth until an OD_{660} reached 0.5. The

medium was changed to M9 minimum medium (supplemented with 0.4% glucose and

0.2% casamino acids) and further cultivated for 2 or 4 h. The cells were subsequently

collected, and total RNA was extracted using an RNeasy Mini Kit (QIAGEN). The

extracted total RNA was analyzed with electrophoresis using MOPS-formaldehyde

denaturing gel or a Bioanalyzer (Agilent).

Quantification of biofilm formation activity

Biofilm formation was quantified by crystal violet staining following a published

procedure[76].

Preparation of 100S ribosomes for cryo-electron microscopy

E. coli W3110, a wild-type K-12 strain, was used to prepare 100S ribosomes.

Overnight 2-mL precultures were obtained in medium E containing $MgSO_4$, citric acid,

K_2HPO_4, and $NaNH_4HPO_4$, and were supplemented with 2% polypeptone and 0.5%

glucose at 37 °C with shaking at 180 cycles per minute[77]. Mass cultures were performed

in 100 mL of the same medium at 37 °C, shaking at 130 cycles per minute. The cells were

harvested 24 h after the start of mass culture. The harvested cell pellets were suspended

in an association buffer [100 mM CH_3COONH_4, 15 mM $(CH_3COO)_2Mg:4H_2O$, 20 mM

Tris–HCl at pH 7.6, and 6 mM 2-mercaptoethanol] and then vortexed with an

approximately equal volume of glass beads (212–300 mm) for 3 minutes (Merck KGaA,

Germany). The homogenate was centrifuged at 12,000 g for 8 min at 4 °C. The

supernatant was layered on top of a 5–20% linear sucrose density gradient in the

association buffer. It was centrifuged in an SW 41 Ti rotor (Beckman Coulter, Inc., United

States) at 285,000 g for 1.5 h at 4 °C. After centrifugation, the absorbance of the sucrose

gradient was measured at 260 nm using a UV-1800 spectrophotometer (Shimadzu Co.,

Japan) to obtain ribosome profiles. The 100S ribosomal fractions were identified from

the measured ribosomal profiles, and these fractions were stored. The solution containing

100S ribosomes was concentrated (final concentration: about 895 pmol), and sucrose was

removed using Amicon-100K (Merck KGaA, Germany).

Sample preparation and cryo-electron microscopy acquisition

The ribosome complex was program-reconstructed using 1.2 μM 100S

hibernating ribosomes complexed with HPF and RMF from *E. coli* W3110 strain and 7.2

μM *E. coli* RNase I, incubated in the reaction buffer (20 mM Hepes-KOH pH 7.6, 10 mM

$MgOAc_2$, 30 mM NH_4Cl, 6 mM β-mercaptoethanol) at 37°C for 30 min. After 1.5-fold

diluting the complex, it was applied onto an EM-grid and plunge-frozen in liquid

nitrogen-cooled ethane. Quantifoil R2/2 gold 200 mesh grid (Quantifoil Micro Tools

GmbH, Germany) glow-discharged for 30 sec in a PIB-10 Ion Bombarder (Vacuum

Device Inc, Japan) was used to apply 2.7 μL sample, pre-incubated for 10 sec, blotted for

15 sec, and post-incubated for 2 sec at 4°C and 100% humidity before freezing in a

Vitrobot Mark IV (ThemoScientific, United States). The grids were subsequently clipped

into an autogrid cartridge and transferred into a CRYO ARM 300 II field emission cryo-

electron microscope (JEOL, Japan) operating at 300 keV, and data acquired with a K3

camera (Gatan, United States) using SerialEM software[78]. The cryo-EM movies were

acquired at a calibrated magnification of 80,460 and a pixel size of 0.87 Å/pixel to record

movies with 70 frames with a total accumulated dose of 50 $e^-/Å^2$/movie.

Cryo-electron microscopy image processing

Image processing steps are summarized in Extended Data Fig. 3. All image

processing steps were performed by cryoSPARC 3.3.1 and 4.0.0 software[79]. 2,000 of

33,302 movies were initially motion-corrected by the patch motion correction job,

followed by the patch CTF estimation. 258,744 particles were picked by the circular blob,

which is 260 – 300 Å, using the blob picker job and extracted at a 4-fold decimated pixel

size (3.48 Å per pixel). The extracted particles were two-dimensional (2D) classified to

remove the classes of 30S subunit and debris. The 70S particles on the micrographs were

then re-picked at a size of 360 Å using the classes selected as templates, followed by the

extract from the micrographs job and 2D classification jobs. Five ab-initio volumes were

reconstructed using selected 146,282 particles to obtain the main class average of the 70S

ribosome and discard the classes of 50S subunit and debris by the heterogenous

refinement job. After the 70S-like particles were combined using the homogeneous

refinement, followed by the local refinement with signal subtraction on the 30S head, the

three-dimensional (3D) classification focused on the head region was performed,

obtaining RNase I binding particles. Selected 33,823 particles were extracted at 1.25-fold

decimated pixel size (2.175 Å per pixel). The complex could not find the RMF in the

volume at a nominal resolution of 2.8 Å. Next, the remaining movies were processed in

the same way as described above. The heterogeneous refinement was performed to extract

the 70S-like classes composed of 1,501,256 particles from the 2D classes of all 33,294

micrographs. The soft mask focused on the 30S head with the hibernation factors (HFs),

and RNase I was generated using UCSF ChimeraX 1.5[80], before 351,633 particles

containing hibernating 70S with RNase I were obtained by the 3D classification, followed

by the local refinement with signal subtraction to improve the local resolution of the

focused region. The local 3D volume was refined to a nominal resolution of 2.4 Å, yet

the density of RMF remained poor. Thereupon, two rounds of the 3D classification

discarded particles containing the weak density map of RMF. 3D classification using the

mask on the 30S platform, including S1 further sorted out 50,955 particles, including

RMF and part of S1. The final reconstruction was refined to a nominal resolution of 2.8

Å in a homogeneous map and to 2.8 Å in a local map of the mask on the 30S head, taking

into account the global/local CTF refinements after the particles were re-extracted at 1.4-

fold decimated pixel size (1.21 Å per pixel). Since the orientation of both homogeneous

and local volumes were slightly different, the local volume was docked into the

homogeneous volume using the 'Fit in Map' function and resampled to the grid of

homogeneous refinement using trilinear interpolation by the 'volume resample' operation

in ChimeraX 1.6.1[80].

Model building and refinement

The *E. coli* 70S ribosome structure in this study was built in the 2.8 Å density

map obtained by homogeneous refinement in cryoSPARC, using a model of the *E. coli*

70S ribosome structure from PDB 8SYL as follows. After removing all molecules and

ions other than ribosome components of the 8SYL[5] model, the ribosomal subunits, split

into 50S and 30S using PyMOL, were docked as rigid bodies into the density map using

the 'Fit in Map' function in ChimeraX 1.6.1[80]. The model of each subunit was refined

with jelly-body restraints by Servalcat 0.4.74[81]. The models were manually corrected for

their ribosomal proteins according to secondary structure and Ramachandran restraints

for better fitting into the density map using the 'Real Space Refine Zone' function in Coot

0.9.8.95[82,83]. Also, nucleotides of the rRNA chains in the models were manually corrected

for their syn or anti-conformations with reference to the differential Fourier map produced

by Servalcat. The models of the HPF derived from the crystal structure of *T. thermophilus*

70S•*E. coli* HPF complex (PDB 4V8H[34]) and the RMF derived from the cryo-EM

structure of *E. coli* hibernating ribosome complex (PDB 6H4N[30]) were rigid-body fitted

into the homogenous density map, respectively. The crystal structure of *E. coli* RNase I

(PDB 2Z70[35]) was rigid-body fitted into the EM density map of the local refinement

focused on the 30S head region in cryoSPARC. Two mutated residues H55F and H133F

at the active site of the RNase I model were manually modified and corrected in Coot.

These models were merged into the 30S model before it was jelly-body refined by

Servalcat, then the 30S binding partners, especially, were manually corrected, residue by

residue, in Coot, together with Ramachandran and secondary structure restraints. After

merging the models of the 30S complex and 50S into hibernating 70S•RNase I complex,

a series of operations in jelly-body refinement and manual correction were repeated

several rounds until the model satisfied its geometry. Magnesium ions and water

molecules from 8SYL were docked into the model of the 70S•RNase I complex in Coot.

After refining the small molecules merged with the complex by Servalcat, their atomic

coordinates were manually corrected or removed to better fit the density map and the

differential Fourier map. The resulting model was validated using a comprehensive cryo-

EM validation tool in PHENIX 1.21.1[84,85] (Extended Data Table 1).

APBS and PISA calculations

The surface potential values were produced by APBS[37]. The interfacing residues

and the buried surface areas were calculated using the protein interface, surfaces, and

assemblies   (PISA)   provided   by   Protein   Data   Bank   Europe   (PDBe)[36]

(https://www.ebi.ac.uk/pdbe/pisa/). Color coding based on BSA values was done on

PyMOL   using   the   original   Python   program   colorByBSA.py

(https://github.com/minami1009/colorByBSA).

In vitro RNA degradation and binding assays

The *in vitro* assays were performed based on the previous study[33]. For the

inhibition assay, 0.075 pmol of the recombinant RNase I was mixed with 0.030 A_{260} of

crude, purified, or reconstituted ribosomes. The mixtures were incubated for 30 min and

30 °C, and the RNase activity of each mixture was quantified using the RNase Activity

Assay Kit (Abcam).

For the binding assay, 20 µM each of ribosomes (70S ribosome or reconstituted

hibernating ribosome) and RNase I mutant (H55F/H133F) was mixed and incubated for

30 min and 37 °C. The mixture was then loaded onto the ultrafiltration device (300,000

MWCO; PALL) and centrifuged at 14,000 g. Following centrifugation, the retentate and

the filtrate were collected separately. RNase I and the ribosomal protein in each sample

were detected by western blotting.

The binding assay following *in vitro* translation was performed as follows: *In*

vitro translation using PURE*frex* 1.0 (GeneFrontier) with or without 1 µg of pET-*gfp* as

a template was performed for 30 min and 37 °C. When hibernating ribosomes were used,

RMF and HPF were added to ribosomes before *in vitro* translation. RNase I mutant

(H55F/H133F) was added to the translation mixture at a final concentration of 4.8 µM

and incubated for a further 30 min. After ultrafiltration, the filtrate was collected, and the

RNase activity of the filtrate was detected as described above.

For the *in vitro* translation inhibition assays, *in vitro* translation was performed

as described above with modifications: 6 µM of RNase I mutant (H55F/H133F) was

added to the translation mixture with 1 µM of ribosomes at the beginning of incubation.

Single-molecule imaging analyses

The translation mRNA template was transcribed from the template vector using

an *in vitro* transcription kit (Promega) and purified using NucleoSpin RNA

(MACHEREY-NAGEL). The mRNA template was hybridized with the 5′ Cy3-labeled

oligo DNA as follows: 1 µM of oligo DNA was denatured at 1 min and 95 °C in the

phosphate buffer (25 mM Na_2HPO_4, 75 mM NaH_2PO_4, 100 mM NaCl, pH 6.4), 300 nM

of the mRNA template was added to the mixture, and hybridization was performed by

lowering the temperature to 25 °C at a rate of 0.1 °C/s. The mixture was added to the

PURE*frex* 1.0 solution I, II, 200 nM of ribosomes, and 50 nM of HaloTag-biotin ligand.

The resulting solution is described as translation reaction solution A. The translation

reaction solution A was pre-incubated at 25 min and 30 °C.

Purified RNase I mutant (H55F/H133F) and nanobody of MoonTag were

buffered to 100 mM $NaHCO_3$, and fluorescence-labeled NHS esters (Alexa Fluor 488 for

nanobody of MoonTag and Alexa Fluor 660 for RNase I mutant, respectively) were added

and incubated at room temperature for 1 h. To remove the unreacted fluorescent dye, the

mixtures were ultrafiltrated using Amicon Ultra 0.5 (3,000 MWCO; Merck Millipore)

and purified using a gel filtration column NAP-5 (Cytiva). These fluorescence-labeled

proteins were added to the PURE*frex* 1.0 solution I and II at a final concentration of 250

nM (nanobody of MoonTag) and 20 nM (RNase I mutant), respectively. The resulting

solution is described as translation reaction solution B.

Single-molecule imaging analyses for the detection of translating ribosomes

were carried out as described previously[86] with the following modifications. 3 mg/mL

biotinylated BSA was poured into a flow cell and washed with Polymix buffer (50 mM

Tris-OAc (pH 7.5), 5 mM $Mg(OAc)_2$, 1 mM EDTA, 100 mM KCl, 5 mM NH_4OAc, 0.5

mM $Ca(OAc)_2$, 5 mM putrescine, 1 mM spermidine, 5 mM β-mercaptoethanol)

containing 0.2% Tween 20. After blocking, 0.33 mg/mL streptavidin was loaded,

followed by washing with Polymix buffer. After streptavidin treatment, 10 µL of

translation reaction solution A, which was pre-incubated at 30°C for 25 min, was loaded,

followed by washing with Polymix buffer twice. At last, translation reaction solution B

containing fluorescence-labeled RNase I mutant (H55F/H133F) and nanobody was added

into the flow cell, and the flow cell was sealed. The flow cell was placed on inverted

microscopy (Nikon), and Alexa Fluor 488, Cy3, and Alexa Fluor 660 fluorescences were

detected upon 488, 532, and 647 nm excitation, respectively. The images were analyzed

with TrackMate plugin of Fiji, after background subtraction. Analysis of the tracking data

and creation of figures were performed using an original Python program.

In vivo trans-translation activity assays

The *trans*-translation activity in *E. coli* cells was quantified based on the

previous research[87]. In brief, *E. coli* Δ*ssrA* or Δ*ssrA*Δ*rna* cells harboring pMW-*ssrA*[DD]

and pAcGFP-woSTOP-*trpAt* were cultivated in L-broth at 37 °C until an OD660 reached

0.3, IPTG was added to the culture at a final condition of 1 mM and further cultivated for

1.5 h. The culture was collected and resuspended in PBS, and GFP intensity was

quantified.

In vitro rRNA degradation from split ribosome subunits

The susceptibility of rRNA to degradation by RNase I was compared for rRNA

derived from 70S ribosomes and 30S/50S subunits. The dissociation of 70S ribosomes

into 30S/50S subunits was performed as follows[48]: 20 pmol of ribosomes was

resuspended to the prokaryotic ribosome dissociation buffer (10 mM Tris-HCl (pH 7.5),

1 mM Mg(OAc)$_2$, 100 mM NH$_4$Cl, 1 mM DTT, 0.5 mM EDTA) and allowed to stand

overnight at 4 °C. 124 pmol of RNase I was added to the solution and incubated for 30

min and 37 °C, and rRNA was extracted using RNeasy Mini Kit as described above. The

rRNAs were detected with electrophoresis using MOPS-formaldehyde denaturing gel.

Preparation of figures

All figures were generated by PyMOL (The PyMOL Molecular Graphics System,

Version 3.0.0 Schrödinger, LLC), Chimera X 1.7.1, Matplotlib (Version 3.7.1), and

Seaborn (Version 0.13.1) libraries of Python (Version 3.10.12), OmniGraffle, and

Affinity Designer (Version 1.10.5).

Data availability

The atomic coordinate of hibernating 70S ribosome in complex with RNase I

mutant was deposited in the Protein Data Bank Japan (PDBj) under accession code 9IOT.

The cryo-EM map was deposited in the Electron Microscopy Data Bank (EMDB) under

accession code EMD-60747, and the local map focused on the 30S head was deposited

as an additional map.

Acknowledgments

We thank K. Namba, T. Miyata, and F. Makino for their work supporting the

cryo-EM facility of Osaka University; the members of the Laboratory for Cryo-EM

structural biology (the T. Kato group) for supporting and advising the cryo-EM single

particle analysis; K. Yamashita for advising model refinement using Servalcat; T. Suzuki,

A. Nagao and N. Akiyama for performing the SDGC and the isolation of ribosomes; W.

Fujii for providing the plasmid encoding the antibody of MoonTag; and H. Himeno for

providing the plasmid pMW-ssrA[DD]. This work was financed by the JSPS KAKENHI

(Grant numbers 20K21266 and 23H02122) to T.O. and by JST SPRING (Grant number

JPMJSP2108) to A.M. This research was [partially] supported by Research Support

Project for Life Science and Drug Discovery (Basis for Supporting Innovative Drug

Discovery and Life Science Research (BINDS)) from AMED under Grant Number

JP23ama121001.

Author contributions

A.M. performed biochemical assays. A.M., T.T. and H.Y. prepared cryo-EM

samples. T.T. prepared cryo-EM grids, collected cryo-EM data, and generated and refined

the molecular models. Z.Y. and T.F. prepared the microscopy. A.M. and Z.Y. performed

the single-molecule imaging assays and analyses. A.M., T.T., Ta.K., To.K. and T.O.

analysed and interpreted data. A.M. and T.T. prepared the figures and tables. A.M., T.T.

and T.O. wrote the manuscript. T.O. conceived and supervised the project. All authors

discussed and commented on the final manuscript.

Corresponding authors

Correspondence to Tetsuhiro Ogawa (atetsu@g.ecc.u-tokyo.ac.jp) or Takehito

Tanzawa (takehito.tanzawa@protein.osaka-u.ac.jp).

Competing interest declaration

The authors declare no competing interests.

References

1.   Irving, S. E., Choudhury, N. R. & Corrigan, R. M. The stringent response and

physiological roles of (pp)pGpp in bacteria. *Nat. Rev. Microbiol.* **19**, 256–271 (2021).

2.   Bechhofer, D. H. & Deutscher, M. P. Bacterial ribonucleases and their roles in RNA

metabolism. *Crit. Rev. Biochem. Mol. Biol.* **54**, 242–300 (2019).

3.   Sulthana, S., Basturea, G. N. & Deutscher, M. P. Elucidation of pathways of ribosomal

RNA degradation: an essential role for RNase E. *RNA* **22**, 1163–1171 (2016).

4.   Jacob, A. I., Köhrer, C., Davies, B. W., RajBhandary, U. L. & Walker, G. C. Conserved

bacterial RNase YbeY plays key roles in 70S ribosome quality control and 16S rRNA

maturation. *Mol. Cell* **49**, 427–438 (2013).

5.   Cheng, Z.-F. & Deutscher, M. P. Quality control of ribosomal RNA mediated by

polynucleotide phosphorylase and RNase R. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 6388–

6393 (2003).

6.   Prossliner, T., Skovbo Winther, K., Sørensen, M. A. & Gerdes, K. Ribosome Hibernation.

Annu. Rev. Genet. **52**, 321–348 (2018).

7.   Wada, A., Yamazaki, Y., Fujita, N. & Ishihama, A. Structure and probable genetic

location of a "ribosome modulation factor" associated with 100S ribosomes in stationary-

phase Escherichia coli cells. *Proc. Natl. Acad. Sci. U. S. A.* **87**, 2657–2661 (1990).

8.   Wada, A., Igarashi, K., Yoshimura, S., Aimoto, S. & Ishihama, A. Ribosome modulation

factor: stationary growth phase-specific inhibitor of ribosome functions from Escherichia

coli. *Biochem. Biophys. Res. Commun.* **214**, 410–417 (1995).

9.   Prossliner, T., Gerdes, K., Sørensen, M. A. & Winther, K. S. Hibernation factors directly

block ribonucleases from entering the ribosome in response to starvation. *Nucleic Acids*

Res. **49**, 2226–2239 (2021).

10. Lipońska, A. & Yap, M.-N. F. Hibernation-Promoting Factor Sequesters Staphylococcus

aureus Ribosomes to Antagonize RNase R-Mediated Nucleolytic Degradation. *MBio* **12**,

e0033421 (2021).

11. Helena-Bueno, K. *et al.* A new family of bacterial ribosome hibernation factors. *Nature*

626, 1125–1132 (2024).

12. Van Dyke, N., Baby, J. & Van Dyke, M. W. Stm1p, a ribosome-associated protein, is

important for protein synthesis in Saccharomyces cerevisiae under nutritional stress

conditions. *J. Mol. Biol.* **358**, 1023–1031 (2006).

13. Wang, Y. J. *et al.* Lso2 is a conserved ribosome-bound protein required for translational

recovery in yeast. *PLoS Biol.* **16**, e2005903 (2018).

14. Andersen, K. L. & Collins, K. Several RNase T2 enzymes function in induced tRNA and

rRNA turnover in the ciliate Tetrahymena. *Mol. Biol. Cell* **23**, 36–44 (2012).

15. Hillwig, M. S. *et al.* RNS2, a conserved member of the RNase T2 family, is necessary for

ribosomal RNA decay in plants. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 1093–1098 (2011).

16. Haud, N. *et al. rnaset2* mutant zebrafish model familial cystic leukoencephalopathy and

reveal a role for RNase T2 in degrading ribosomal RNA. *Proceedings of the National*

Academy of Sciences **108**, 1099–1103 (2011).

17. Sato, K. & Egami, F. STUDIES ON RIBONUCLEASES IN TAKADIASTASE. I. *J.*

Biochem. **44**, 753–767 (1957).

18. Luhtala, N. & Parker, R. T2 Family ribonucleases: ancient enzymes with diverse roles.

Trends Biochem. Sci. **35**, 253–259 (2010).

19. Huang, H. *et al.* Bulk RNA degradation by nitrogen starvation-induced autophagy in yeast.

EMBO J. **34**, 154–168 (2015).

20. Minami, A. *et al.* RNase T2-involved selective autophagy of ribosomes induced by

starvation in yeast. *bioRxiv* 2023.07.14.548783 (2023) doi:10.1101/2023.07.14.548783.

21. Henneke, M. *et al.* RNASET2-deficient cystic leukoencephalopathy resembles congenital

cytomegalovirus brain infection. *Nat. Genet.* **41**, 773–775 (2009).

22. Greulich, W. *et al.* TLR8 Is a Sensor of RNase T2 Degradation Products. *Cell* **179**, 1264-

1275.e13 (2019).

23. Takayama, S. & Isogai, A. Self-incompatibility in plants. *Annu. Rev. Plant Biol.* **56**, 467–

489 (2005).

24. Beppu, T. & Arima, K. Induction by mercuric ion of extensive degradation of cellular

ribonucleic acid in Escherichia coli. *J. Bacteriol.* **98**, 888–897 (1969).

25. Meador, J., 3rd & Kennell, D. Cloning and sequencing the gene encoding Escherichia coli

ribonuclease I: exact physical mapping using the genome library. *Gene* **95**, 1–7 (1990).

26.   Neu, H. C. & Heppel, L. A. THE RELEASE OF RIBONUCLEASE INTO THE

MEDIUM WHEN ESCHERICHIA COLI CELLS ARE CONVERTED TO

SPEROPLASTS. *J. Biol. Chem.* **239**, 3893–3900 (1964).

27.   Datta, A. K. & Burma, D. P. Association of ribonuclease I with ribosomes and their

subunits. *J. Biol. Chem.* **247**, 6795–6801 (1972).

28.   Flemming, H.-C. & Wuertz, S. Bacteria and archaea on Earth and their abundance in

biofilms. *Nat. Rev. Microbiol.* **17**, 247–260 (2019).

29.   Ueta, M. *et al.* Ribosome binding proteins YhbH and YfiA have opposite functions during

100S formation in the stationary phase of Escherichia coli. *Genes Cells* **10**, 1103–1112

(2005).

30.   Beckert, B. *et al.* Structure of a hibernating 100S ribosome reveals an inactive

conformation of the ribosomal protein S1. *Nature Microbiology* **3**, 1115–1121 (2018).

31.   Maruyama, H. & Mizuno, D. The participation of ribonuclease in the degradation of

Escherichia coli ribosomal ribonucleic acid as revealed by oligonucleotides accumulation

in the phorphorus-deficient stage. *Biochim. Biophys. Acta* **108**, 593–604 (1965).

32.   Wade, H. E., Lovett, S. & Robinson, H. K. The autodegradation of 32-P-labelled

ribosomes from Escherichia coli. *Biochem. J* **93**, 121–128 (1964).

33.   Kitahara, K. & Miyazaki, K. Specific inhibition of bacterial RNase T2 by helix 41 of 16S

ribosomal RNA. *Nat. Commun.* **2**, 1–7 (2011).

34.   Polikanov, Y. S., Blaha, G. M. & Steitz, T. A. How Hibernation Factors RMF, HPF, and

YfiA Turn Off Protein Synthesis. *Science* **336**, 915–918 (2012).

35.   Rodriguez, S. M. *et al.* Nonspecific base recognition mediated by water bridges and

hydrophobic stacking in ribonuclease I from Escherichia coli. *Protein Sci.* **17**, 681–690

(2008).

36.   Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state.

J. Mol. Biol. **372**, 774–797 (2007).

37.   Jurrus, E. *et al.* Improvements to the APBS biomolecular solvation software suite. *Protein*

Sci. **27**, 112–128 (2018).

38.   Seely, S. M. *et al.* Molecular basis of the pleiotropic effects by the antibiotic amikacin on

the ribosome. *Nat. Commun.* **14**, 4666 (2023).

39.   Shimizu, Y. *et al.* Cell-free translation reconstituted with purified components. *Nat.*

Biotechnol. **19**, 751–755 (2001).

40.   Wu, B., Eliscovich, C., Yoon, Y. J. & Singer, R. H. Translation dynamics of single

mRNAs in live cells and neurons. *Science* **352**, 1430–1435 (2016).

41.   Morisaki, T. *et al.* Real-time quantification of single RNA translation dynamics in living

cells. *Science* **352**, 1425–1429 (2016).

42.   Yan, X., Hoek, T. A., Vale, R. D. & Tanenbaum, M. E. Dynamics of Translation of Single

mRNA Molecules In Vivo. *Cell* **165**, 976–989 (2016).

43.   Wang, C., Han, B., Zhou, R. & Zhuang, X. Real-Time Imaging of Translation on Single

mRNA Transcripts in Live Cells. *Cell* **165**, 990–1001 (2016).

44.   Fontaine, B. M. *et al.* RNase I regulates Escherichia coli 2',3'-cyclic nucleotide

monophosphate levels and biofilm formation. *Biochem. J* **475**, 1491–1506 (2018).

45.   Deutscher, Murray P. Chapter 9 Maturation and Degradation of Ribosomal RNA in

Bacteria. in *Progress in Molecular Biology and Translational Science* vol. 85 369–391

(Academic Press, 2009).

46.   Zundel, M. A., Basturea, G. N. & Deutscher, M. P. Initiation of ribosome degradation

during starvation in Escherichia coli. *RNA* **15**, 977–983 (2009).

47.   Himeno, H., Nameki, N., Kurita, D., Muto, A. & Abo, T. Ribosome rescue systems in

bacteria. *Biochimie* **114**, 102–112 (2015).

48.   Rivera, M. C., Maguire, B. & Lake, J. A. Dissociation of ribosomes into large and small

subunits. *Cold Spring Harb. Protoc.* **2015**, 363–367 (2015).

49. Keiler, K. C. Mechanisms of ribosome rescue in bacteria. *Nat. Rev. Microbiol.* **13**, 285–

297 (2015).

50. Paier, A., Leppik, M., Soosaar, A., Tenson, T. & Maiväli, Ü. The effects of disruptions in

ribosomal active sites and in intersubunit contacts on ribosomal degradation in Escherichia

coli. *Sci. Rep.* **5**, 7712 (2015).

51. Ron, E. Z. Polysome turnover during amino acid starvation in Escherichia coli. *J.*

Bacteriol. **108**, 263–268 (1971).

52. Cozzone, A. J. How do bacteria synthesize proteins during amino acid starvation? *Trends*

Biochem. Sci. **6**, 108–110 (1981).

53. Fessler, M., Gummesson, B., Charbon, G., Svenningsen, S. L. & Sørensen, M. A. Short-

term kinetics of rRNA degradation in Escherichia coli upon starvation for carbon, amino

acid or phosphate. *Mol. Microbiol.* **113**, 951–963 (2020).

54. Filbeck, S., Cerullo, F., Pfeffer, S. & Joazeiro, C. A. P. Ribosome-associated quality-

control mechanisms from bacteria to humans. *Mol. Cell* **82**, 1451–1466 (2022).

55. Saito, K. *et al.* Ribosome collisions induce mRNA cleavage and ribosome rescue in

bacteria. *Nature* **603**, 503–508 (2022).

56. LaRiviere, F. J., Cole, S. E., Ferullo, D. J. & Moore, M. J. A late-acting quality control

process for mature eukaryotic rRNAs. *Mol. Cell* **24**, 619–626 (2006).

57. Cole, S. E., LaRiviere, F. J., Merrikh, C. N. & Moore, M. J. A convergence of rRNA and

mRNA quality control pathways revealed by mechanistic analysis of nonfunctional rRNA

decay. *Mol. Cell* **34**, 440–450 (2009).

58. Garza-Sánchez, F., Gin, J. G. & Hayes, C. S. Amino acid starvation and colicin D

treatment induce A-site mRNA cleavage in Escherichia coli. *J. Mol. Biol.* **378**, 505–519

(2008).

59. Messens, J. *et al.* The Oxidase DsbA Folds a Protein with a Nonconsecutive Disulfide*. *J.*

Biol. Chem. **282**, 31302-31307 (2007).

60.   Padmanabhan, S., Zhou, K., Chu, C. Y., Lim, R. W. & Lim, L. W. Overexpression,

biophysical characterization, and crystallization of ribonuclease I from Escherichia coli, a

broad-specificity enzyme in the RNase T2 family. *Arch. Biochem. Biophys.* **390**, 42–50

(2001).

61.   Ohnishi, Y., Iguma, H., Ono, T., Nagaishi, H. & Clark, A. J. Genetic mapping of the F

plasmid gene that promotes degradation of stable ribonucleic acid in Escherichia coli. *J.*

Bacteriol. **132**, 784–789 (1977).

62.   Onishi, Y. F factor promotes turnover of stable RNA in escherichia coli. *Science* **187**,

257–258 (1975).

63.   Gao, H. *et al.* Study of the structural dynamics of the E coli 70S ribosome using real-space

refinement. *Cell* **113**, 789–801 (2003).

64.   Yapici, I. *et al.* 4D Crystallography Captures Transient IF1-Ribosome Dynamics in

Translation Initiation. *bioRxiv* 2023.10.27.564398 (2023) doi:10.1101/2023.10.27.564398.

65.   Akanuma, G. *et al.* Evolution of Ribosomal Protein S14 Demonstrated by the

Reconstruction of Chimeric Ribosomes in Bacillus subtilis. *J. Bacteriol.* **203**, (2021).

66.   Xaplanteri, M. A., Papadopoulos, G., Leontiadou, F., Choli-Papadopoulou, T. & Kalpaxis,

D. L. The contribution of the zinc-finger motif to the function of Thermus thermophilus

ribosomal protein S14. *J. Mol. Biol.* **369**, 489–497 (2007).

67.   Li, Y. *et al.* Zinc depletion induces ribosome hibernation in mycobacteria. *Proc. Natl.*

Acad. Sci. U. S. A. **115**, 8191–8196 (2018).

68.   Thorn, A. *et al.* Structure and activity of the only human RNase T2. *Nucleic Acids Res.* **40**,

8733–8742 (2012).

69.   Dimitrova-Paternoga, L. *et al.* Structural basis of ribosomal 30S subunit degradation by

RNase R. *Nature* **626**, 1133–1140 (2024).

70.   Baba, T. *et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout

mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

71. Boersma, S. *et al.* Multi-Color Single-Molecule Imaging Uncovers Extensive

Heterogeneity in mRNA Decoding. *Cell* **178**, 458-472.e19 (2019).

72. Yang, Z., Iizuka, R. & Funatsu, T. Nascent SecM chain outside the ribosome reinforces

translation arrest. *PLoS One* **10**, e0122017 (2015).

73. Dijk, J. & Littlechild, J. Purification of ribosomal proteins from Escherichia coli under

nondenaturing conditions. *Methods Enzymol.* **59**, 481–502 (1979).

74. Akiyama, N. *et al.* Structural insights into the decoding capability of isoleucine tRNAs

with lysidine and agmatidine. *Nat. Struct. Mol. Biol.* **31**, 817–825 (2024).

75. Feaga, H. A., Kopylov, M., Kim, J. K., Jovanovic, M. & Dworkin, J. Ribosome

Dimerization Protects the Small Subunit. *J. Bacteriol.* **202**, (2020).

76. Minami, A. *et al.* Establishment of the improved colonization of Escherichia coli

laboratory strain in the intestine mediated by single gene deletion. *Biochem. Biophys. Res.*

Commun. 150448 (2024).

77. Vogel, H. J. & Bonner, D. M. ACETYLORNITHINASE OF ESCHERICHIA COLI:

PARTIAL PURIFICATION AND SOME PROPERTIES. *J. Biol. Chem.* **218**, 97–106

(1956).

78. Mastronarde, D. N. Automated electron microscope tomography using robust prediction of

specimen movements. *J. Struct. Biol.* **152**, 36–51 (2005).

79. Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for

rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

80. Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,

and developers. *Protein Sci.* **30**, 70–82 (2021).

81. Yamashita, K., Palmer, C. M., Burnley, T. & Murshudov, G. N. Cryo-EM single-particle

structure refinement and map calculation using Servalcat. *Acta Crystallogr D Struct Biol*

77, 1282–1291 (2021).

82. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.

Acta Crystallogr. D Biol. Crystallogr. **66**, 486–501 (2010).

83. Casañal, A., Lohkamp, B. & Emsley, P. Current developments in Coot for macromolecular

model building of Electron Cryo-microscopy and Crystallographic Data. *Protein Sci.* **29**,

1069–1078 (2020).

84. Liebschner, D. *et al.* Macromolecular structure determination using X-rays, neutrons and

electrons: recent developments in Phenix. *Acta Crystallogr D Struct Biol* **75**, 861–877

(2019).

85. Chen, V. B. *et al.* MolProbity: all-atom structure validation for macromolecular

crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12–21 (2010).

86. Masuda, T. *et al.* Initiation factor 2, tRNA, and 50S subunits cooperatively stabilize

mRNAs on the ribosome during initiation. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 4881–4885

(2012).

87. Ramadoss, N. S. *et al.* Small molecule inhibitors of *trans*-translation have broad-spectrum

antibiotic activity. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 10282–10287 (2013).

1030