EFFECTS OF TWO DIFFERENT COMPOUNDS ON SEIZURE SUPPRESSION USING THE ZEBRAFISH PTZ-SEIZURE MODEL

Jhonathan Angel Araujo Fernández[1,2], Thatiane Cristina de Moura[1,2], Sabela Fernández Vila[3],

Juan Andrés Rubiolo Gaytán[3], Iñaki López-Díaz[5], Soraya Learte-Aymamí[4], M. Eugenio

Vázquez [4], Maria D. Mayán[5], Laura Sánchez[3], Claudia Vianna Maurer-Morelli[1,2*]

[1]Departamento de Medicina Translacional, Faculdade de Ciências Médicas, Universidade Estadual de
Campinas, UNICAMP, Campinas, Brazil

[2]Brazilian Institute of Neuroscience and Neurotechnology, Campinas, Brazil

[3]Departamento de Zoología Genética y Antropología Física, Facultad de Veterinaria, Universidad de
Santiago de Compostela, Campus de Lugo, 27002 Lugo, Spain

[4]Centro Singular de Investigación en Química Biolóxica e Materiais Moleculares (CiQUS),
Departamento de Química Orgánica, Universidade de Santiago de Compostela, Santiago de
Compostela, Spain

[5]CELLCOM Research Group. CINBIO and Galicia Sur Health Research Institute (IIS Galicia Sur).
SERGAS. University of Vigo. Edificio Olimpia Valencia, Campus Universitario Lagoas Marcosende,
36310, Pontevedra, Spain

*Corresponding author

Cláudia Vianna Maurer-Morelli

E-mail: cmorelli@unicamp.br

Abstract

Epilepsies are a common and severe neurological condition characterized by spontaneous and recurrent seizures. Although anti-seizure medications are effective for most patients, about 30% remain pharmacoresistant. Moreover, uncontrolled seizures are associated with risk factors and shortened life expectancy for individuals with refractory epilepsy. Preclinical studies are an essential step for drug discovery and the zebrafish (*Danio rerio*) has been successfully employed for this purpose. In this study, we applied the zebrafish PTZ-seizure model to investigate the effect of two compounds on seizure suppression, Tripeptide (p-BTX-I) and the Cx43 peptide CX2. Zebrafish larvae at 6 days post-fertilization (dpf) were exposed to both compounds, according to their group, 24h prior to PTZ-seizure induction. We quantified the compounds' effect on seizure latency, number of seizures and transcript levels of genes related to inflammation, oxidative stress, and apoptosis (*il1b, tnfa, cox1, cox2a, il6, casp3a, casp9, baxa, bcl2a, nox1, sod1* and *cat*).

Our results showed that CX2 at a concentration of 0.1 µM/mL yielded the best outcome for seizure suppression as it reduced the number of seizures and increased the seizure latency. Additionally, CX2 treatment before PTZ-induced seizures decreased the transcript of *il1b, il6, tnfa* and *cox1* genes, all related to inflammation. A bio-distribution study showed that the CX2 reached the zebrafish brain at both times investigated, 1h and 6h. Similarly, the tripeptide exhibited anti-inflammatory and anti-apoptotic action, reducing mRNA expression of the *il1b* and *casp9* genes. Our findings suggest that both Tripeptide and CX2 hold translational potential for seizure suppression.

Keywords: zebrafish[1], drug screening[2], seizure[3], compounds[4], pentylenetetrazole[5].

Introduction

Epilepsies are a common neurological disease characterized by recurrent unprovoked seizures, affecting approximately 50 million people worldwide [1,2]. Currently, anti-seizure medications are effective for 66% of people with epilepsy in developed countries [3,4]. However, more than 30% of people with epilepsy do not respond well to conventional therapies, making them pharmacoresistant [5]. Uncontrolled seizures can lead to various risks for people with epilepsy, such as risk of injury, neuropsychological impairment, and shortened lifespan [6]. Therefore, the search for new drugs or substances that could improve the treatment of people with epilepsy is urgent.

Pre-clinical trials are crucial in testing the therapeutic or toxicological effects of new substances, using *in vivo*, *in vitro*, or *ex-vivo* strategies. The scalability and rapid access to the results significantly contribute to the value of such studies. In this context, the zebrafish emerges as a pivotal model, offering distinct advantages over other animal models for pre-clinical trials [7]. These include its small size, low maintenance cost, high fecundity, and optical transparency during embryogenesis [8]. Furthermore, the zebrafish genome exhibits approximately 70% homology with the human genome, with 84% of known genes associated with human diseases, including epilepsy [9,10]. In recent years, several platforms for automatic data acquisition and analysis in the zebrafish model have been developed, enabling a variety of multiplexed phenotypic assays with minimal human intervention [11,12]. Moreover, the zebrafish aligns with the 3Rs (reduction, refinement, and replacement) philosophy [13]. Beyond its investigation of molecular pathways and behaviors relevant to various diseases, the zebrafish offers a time and cost-effective model due to its small size, among other attributes, which is advantageous for scalable studies involving simultaneous analysis of animals and drugs [14].

In the context of epilepsy, the zebrafish is a well-characterized seizure model in adults and larvae, as it is used for modeling epilepsy disorders by genetic manipulation [15–17]. In 2005, Baraban and

colleagues used the pro-convulsant agent pentylenetetrazol (PTZ) to induce seizures in zebrafish larvae [16]. They observed a specific seizure behavior, up-regulation of the *c-fos* gene in the brain, and electrographic discharges in the optic tectum after exposing the larvae to 15 mM PTZ at seven days post-fertilization. Moreover, these responses were attenuated by common anti-seizure medications. Overall, these findings suggest that zebrafish exhibit many similarities to traditional models such as rodents.

Historically, natural substances from plants and animals are a source of numerous medicinal preparations, widely employed for preventive and therapeutic care [18,19]. This fact is corroborated by the Food and Drug Administration (FDA), which reports that between 1981 and 2019, 34% of drugs were based on substances from natural products [20].

Considering the advantages of zebrafish for drug screening based on phenotype and the need to find new treatments for controlling seizures, we aimed to evaluate the effect of different natural compounds on seizure suppression using the PTZ-zebrafish model.

Materials and Methods

Zebrafish maintenance and embryo acquisition

Wild-type adult zebrafish were obtained from the Laboratory of Zebrafish and Husbandry at the School of Medical Sciences, Unicamp. The animals were housed in 30 – 50 liter tanks, accommodating two animals per liter of water. The tanks were maintained under controlled physicochemical conditions of temperature (26±2°C), pH (7-7.5), levels of ammonia (< 0.1 ppm), nitrite (< 0.2 ppm) and dissolved oxygen (4-11 ppm). A photoperiod cycle of 14 h of light and 10 h of darkness was maintained. Adult fish were fed three times a day with commercial flake food (Tetramin, Tetra, Blacksburg, VA, USA) and once a day with brine shrimp and paramecium. Embryos were collected following natural

93 spawning and nurtured in Petri dishes containing water from the aquariums maintaining consistent

94 temperature and photoperiod with the adults. From the 5th day onwards, the larvae were fed with

95 paramecium. Ethical approval for all experimental protocols was obtained from the Ethics Committee

96 for Animal Research of the State University of Campinas (CEUA 4895-1/2019 and CEUA 5757-

97 1/2021).

98 Compounds

99 Tripeptide (p-BTX-I)

The compound sequence (Glu-Val-Trp) was obtained from AminoTech – Research and Development

(São Paulo, Brazil). Subsequently, it was diluted in Milli-Q water (Merck KGaA, Darmstadt, Germany)

and then aliquoted. These aliquots were stored at -20°C. The employed concentrations (as listed in

Table 1) were determined based on research conducted in cell models [21,22].

Table 1. Compounds and their concentrations were analyzed to determine their effect on the

suppression of seizures induced by the proconvulsant agent pentylenetetrazole.

Compound	Concentration
Tripeptide (p-BTX-I)	25, 10 and 5 µg/mL
CX2	0.5, 0.1 and 0.05 µM

110 CX2

The CX2 (ARG-Cx43p) sequence corresponds to the C-terminal domain of the zebrafish Cx43 gene

(PCT/EP2020/071242). CX2 was assembled following standard Fmoc/tBu solid-phase MW-assisted

peptide synthesis protocols [23,24], purified by reverse-phase HPLC, and their identity confirmed by

HPLC-MS(ESI). The concentrations used (as listed in Table 1) were determined through studies

involving zebrafish models for regeneration and senescence. Like Tripeptide, CX2 was diluted in Milli-

Q water, aliquoted, and stored at -20°C.

Compounds pretreatment

For all compounds, zebrafish larvae at 6 days post-fertilization (dpf) were randomly placed into Petri

dishes, each one containing 25 larvae. The larvae were incubated in the respective solutions and

concentrations for 24h (as specified in Table 1) prior to seizure induction. The temperature and

photoperiod conditions were consistent with those described for the adult zebrafish. The study groups

consisted of the Control Group (CG), PTZ Group (PTZ), Treatment 1, Treatment 2, and Treatment 3.

Treatments 1 to 3 employed the previously determined concentrations mentioned above.

Seizure induction by pentylenetetrazol

Seizure induction followed the pretreatment of the compounds. Larvae with 7 dpf were carefully

transferred to a 96-well plate filled with 100 µL of aquarium water, with one larva per well.

Subsequently, 100 µL of 30 mM of the proconvulsant agent pentylenetetrazol (PTZ) (Sigma-Aldrich,

St. Louis, MO, USA) was added to each well, resulting in a final concentration of 15 mM of PTZ. The

larvae were exposed to PTZ for 20 minutes. The control group underwent the same procedure but in

PTZ-free water.

Latency and number of seizures

132 The latency and number of seizures were monitored through visual observation, following the method

133 described by Barbalho et al., 2016a [25]. Specifically, latency was defined as the period between the

134 initiation of PTZ exposure and the larva reaching stage 3 of seizure-like activity, according to the

135 criteria established by Baraban et al., 2005 [16]. Larvae were evaluated for 10 minutes of PTZ

136 exposure, and a complete seizure was considered when they reached stage 3.

RNA extraction

Immediately after PTZ exposure, fish were cryo-anesthetized and their heads were cut, collected, and

then incubated in TRIzol® (Invitrogen, Carlsbad, CA, USA). The biological material was lysed using

the TissueLyzer equipment (QIAGEN, GmbH, Germany) at 25 beats per second (BPS) for 2 minutes.

Subsequently, the heads were stored at -80 °C until further processing. Each group consisted of five

samples (n = 5); however, each sample was composed by pooling five larval heads, to obtain sufficient

biological material for RNA extraction. Total RNA extraction was performed using TRIzol® according

to the manufacturer's protocol. The concentration and quality were determined using the Epoch™

spectrophotometer (BioTek, Winooski, VT, USA) and gel electrophoresis.

RT-qPCR

The synthesis of cDNA was carried out using the High-Capacity cDNA Reverse Transcription kit

(Applied Biosystems™, Califórnia, USA) following the manufacturer's instructions. Quantitative PCR

(qPCR) was performed using the SYBR ® Green Master Mix reagent (Bio-Rad) using the ABI 7500

system (Applied Biosystems, Foster City, CA, USA). Target genes (as listed in Table 2) were designed

using the Primer-BLAST online tool (https://www.ncbi.nlm.nih.gov/tools /primer-blast/) from NCBI

(National Center for Biotechnology Information) zebrafish database. Runs were carried out in triplicate

using *eef1a1l1* as a housekeeping gene [26] to normalize the genes from the inflammation pathway

(*il1b, tnfa, cox1, cox2a* and *il6*), the apoptosis pathway (*casp3a, casp9, baxa* and *bcl2a*), and the

oxidative stress pathway (*nox1, sod* and *cat*) were analyzed. Data were evaluated using 7500 software

v2.03 (Applied Biosystems). Relative gene expression analysis was calculated using the Livak and

Schmittgen equation RQ = 2-$\Delta\Delta$CT [27].

The three pathways (inflammation, apoptosis, and oxidative stress) are all closely linked to the

underlying mechanisms of seizure generation and progression, which is why we selected representative

genes from these pathways for analysis.

Table 2: Primer sequences designed using the Primer-Blast online tool.

Gene	Seq.Ref.NCBI	Forward	Reverse	Size (pb)
eef1a1l1	NM_131263.1	AGCAGCAGCTGAGGAGTGAT	CCGCATTTGTAGATCAGATGG	140
il1b	NM_212844.2	GCTGGAGATCCAAACGGATA	ATTTGACGGACTCGAAGGTG	85
tnfa	NM_212859.2	TCGGGTGTATGGAGGGTGTT	TTGATTGCCCTGGGTCTTATGG	96
cox1	NM_153656.2	CTGGGAGGCTTATTCCAACA	CCAGAAGTTTAGGGTCTGGAAG	119
cox2a	NM_153657.1	ACCAGGGCGTGTGTTTATCC	GTGAGAAGCTCAGGGGTAGTG	100
il6	NM_001261449.1	GGCATTTGAAGGGGTCAGGA	GCGTTAGACATCTTTCCGTGC	92
casp3a	NM_131877.3	CAGCTTGAACTACCCCAACA	AAGCTTTCGAAACACGTTCA	133
casp9	NM_001007404.2	GCGACAAGCTGGAGAAAAGA	GATGACCACACAGCAGTCGTA	140
baxa	NM_131562.2	GAGCTGCACTTCTCAACAACTTT	GAAGATCTCACGGGCCACTC	245
bcl2a	NM_001030253.2	ACTACCTGAACGGGCCACT	AAAACGGGTGGAACACAGAG	105

nox1	XM_009291119.3	TCGTCAAGAAAACCCATCGTCT	TGTTGATCTTGCAGCACAGTCT	213
sod1	NM_131294.1	GGCCTTACTCCAGGAAAACA	TCTCCGACGTGTCTCACACT	140
cat	NM_130912.2	CAACGCTGAAGGGAAAAAGAACA	GCATTTGCTAAGATCACTTGTGTTG	250

Statistical analysis

164 Statistical analysis was performed using GraphPad Prism version 5.0 (GraphPad Software, San Diego,

165 CA, USA). One-way ANOVA was performed, followed by the Bonferroni method for multiple

166 comparisons. Differences were considered significant if $p<0.05$. Results are presented as mean ± SEM.

Bio-distribution

Bio-distribution analysis was conducted on Tripeptide and CX2, labeled with tetramethylrhodamine-

TMR and carboxytetramethylrhodamine – TAMRA, respectively.

Larvae at 6 dpf were transferred to a 96-well plate, with one larva per well, containing 100 µL of

aquarium water. Subsequently, 100 µL of the Tripeptide compound at 20 µg/mL or CX2 at 1 µM was

added according to the respective group, totalizing 200 ml of solution at final concentration of 10 µg/ml

for Tripeptide and 0.5 µM for CX2. Exposure times for both compound groups were 1, 6, 18, and 24

hours (n=3 each time). The control group underwent similar manipulation but in water only. The

incubation temperature during the experiments was maintained at 26±2°C. upon completion of

exposure, larvae were rapidly and carefully transferred to a Becker containing 0.02% tricaine (Tricaine,

Sigma) for approximately 2 minutes. Subsequently, the larvae were transferred to a Petri dish

containing 1% agarose as a base, along with a drop of water. Larvae were examined and photographed

using the Multizoon AZ100 microscope (Nikon, Tokyo, Japan) at 2X magnification. Two photographs

of each larva were taken, one in black and white, and the other capturing only the fluorescence

emission. Imaging merging was performed using the NIS-Elements software (Nikon, Tokyo, Japan).

The experiments and protocols were approved by the animal care and use committee of the Universidad

de Santiago de Compostela and the standard protocols of Spain (CEEA-LU-003 and Directive 2012-

63-EU).

Results

Behavioral assay

We analyzed the effects of tripeptide and CX2 on seizure behavior. We found an increase in latency

only for animals pretreated with the CX2 at 0.1 µM, ($p \leq 0.05$) as well as a significant decrease in the

number of seizures ($p \leq 0.001$) when compared to the PTZ group (Fig 1B). No statistical significance

was found for the tripeptide (Fig 1A).

Fig 1: The effect of compound treatment prior to pentylenetetrazol (PTZ)-induced seizures on

the number of seizure-like behaviors and latency was investigated. The number of seizures and

latency were determined through visual inspection during a 10-minute exposure to PTZ (15 mM). (A)

Animals exposed to the tripeptide for 24 hours: Pentylenetetrazol (PTZ) group; treatment group 25

µg/mL (T25); 10 µg/ml treatment group (T10); treatment group 5 µg/ml (T5). (B) Animals exposed to

CX2 for 24 hours: Pentylenetetrazol (PTZ) group; treatment group 0.5 µM (T0.5); treatment group 0.1

µM (T0.1); treatment group 0.05 µM (T0.01). Data are presented as mean ± SEM. Statistical analyses

were performed using one-way ANOVA, followed by the Bonferroni method for multiple

comparisons. Differences were considered significant if $p < 0.05$. An asterisk (*) indicates that $p \leq 0.05$;

three asterisks (***), $p \leq 0.001$.

Molecular assay

Regarding the tripeptide, our results indicated downregulation of the *il1b* and *casp9* genes (Fig 2A and

2B), with significant effects at low concentrations (10 and 5 µg/mL). Additionally, we observed that

the expression of the *casp3a*, *baxa,* and *bcl2a* genes was most significantly reduced at a concentration

of 25 µg/mL ($p<0.05$), compared to the other concentrations. Similarly, at a concentration of 10 µg/mL,

the *casp3a*, *baxa,* and *bcl2a* genes exhibited notably high expression levels, roughly equivalent to or

slightly higher than those of the PTZ group. However, no statistical difference was detected.

Fig 2: Expression of *il1b*, *casp3a*, *casp9*, *baxa* and *bcl2a* genes in zebrafish brain after

pentylenetetrazol (PTZ)-induced seizures. Each treatment group was initially exposed to the

tripeptide for 24 h and subsequently to 15 mM PTZ for 20 min, and the control and PTZ groups were

handled identically, but with exposure to water (n = 5 per group). Data are presented as mean ± SEM.

Statistical analyses were performed with the one-way ANOVA, followed by the Bonferroni method

for multiple comparisons. Differences were considered significant if p<0.05. An asterisk (*) indicates

that p≤0.05; two asterisks (**), p≤0.01. Control group (Control); pentylenetetrazol (PTZ) group;

treatment group 25 µg/mL (T25); 10 µg/ml treatment group (T10); treatment group 5 µg/mL (T5).

In the case of the CX2 compounds, we found down-regulating of the *il1b, cox1, il16, and tnfa* genes

for most of the analyzed concentrations (Fig 3) compared to the PTZ group. The most significant down-

regulation was observed at a concentration of 0.1 µM. Conversely, the *cox2a* gene displayed increased

expression across all the three concentrations tested, with the most significant increase occurring at a

concentration of 0.05 µM (p≤0.001).

Fig 3: Expression of *il1b*, *il6*, *tnfa*, *cox1* and *cox2a* genes in zebrafish brain after pentylenetetrazol

(PTZ)-induced seizures. Each treatment group was initially exposed to the CX2 for 24 h and

subsequently to 15 mM PTZ for 20 min, and the control and PTZ groups were treated identically but

with water exposure (n = 5 per group). Data are presented as mean ± SEM. Statistical analyses were

performed with the one-way ANOVA, followed by the Bonferroni method for multiple comparisons.

Differences were considered significant if p<0.05. An asterisk (*) indicates that p≤0.05; two asterisks

(**), p≤0.01; three asterisks (***), p≤0.001; four asterisks (****), p≤0.0001. Control group (Control);

pentylenetetrazol (PTZ) group; treatment group 0.5 µM (T0.5); treatment group 0.1 µM (T0.1);

treatment group 0.05 µM (T0.01).

Bio-distribution

By tagging both the Tripeptide and CX2 with fluorescence, we tracked their biodistribution at four

time points, 1h, 6h, 18h and 24h. The CX2 compound exhibited robust detection in the brain after 1h

and persisted at 6 hours (Fig 4). Additionally, the compound was detected throughout the body,

indicating successful absorption by the zebrafish. In contrast, the tripeptide compound did not reach

the brain, and it was predominantly retained in the zebrafish intestine (S1 Fig).

Figure 4: Time course of CX2 (8ARG-Cx43p, corresponding to the C-terminal domain sequence

of Cx43) efficiently crossing the blood-brain barrier (BBB) in a 7-day-old zebrafish larva. The

carboxytetramethylrhodamine-TAMRA labeled peptide exhibited fluorescence microscopy detection

in the brain and throughout the body (indicated by red color) at 1 hour and 6 hours post-treatment.

Discussion

Despite the availability of pharmacological treatments, uncontrolled seizures remain a concern,

requiring further investigations for new seizure-suppression approaches [28]. Seizures occur due to a

complex process involving multiple factors and affecting various cellular pathways [29]. This study

focused on targeting inflammation, oxidative stress, and cell death pathways for seizure modulation

[30–33]. To achieve our goal, we used tripeptide (p-BTX-I) to address inflammation and cell death,

and the CX2 for its anti-inflammatory action[21,22].

Zebrafish offer many advantages in drug screening and phenotype assessment. The zebrafish PTZ-

seizure model mimics human seizure behavior and electrographic patterns and is responsive to anti-

seizure drugs. The scalability and rapid results of the zebrafish make them an advantageous model for

discovering new anti-seizure compounds [8,9,16,34,35].

The process of inflammation involves the recruitment of many inflammatory mediators such as

interleukins (ILs), interferons (IFNs), tumor necrosis factors (TNFs), and growth factors. *IL1B*, *TNF*-

α, and *IL6* genes are among the most studied inflammatory cytokines in the CNS [32]. After a seizure,

genes such as *IL1B*, *IL2*, *IL6*, *TNF*-α, and *VEGF,* which are normally present in low concentrations,

increase rapidly, leading to damaging changes in synapses and heightened neuronal excitability

[33,36].

Prostaglandins/*COX-2* are also produced, potentially disrupting the blood-brain barrier and

significantly contributing to seizure onset and recurrence [37].

Among the compounds analyzed, CX2 at a concentration of 0.1 μM showed the most significant results

in seizure suppression, reducing seizure frequency and increasing latency (Fig 1B). Treatment with

CX2 prior to PTZ-induced seizures down-regulated the *il1b, cox1, tnfa, and il16* genes, indicating its

role in regulating inflammation during seizures [33]. In contrast, the *cox2a* gene showed up-regulation

compared to the PTZ group (Fig 3E). *COX2,* which is responsible for prostaglandin production [38],

increases during seizures [39]. While *COX2* inhibitors have been explored as therapeutic agents;

controversies exist due to types of inhibitors and timing of administration [39,40]

Studies in rats and zebrafish suggest that *cox1* may play a more crucial role in inhibiting seizure [41–

43]. Barbalho et al. found that inhibiting *cox-1* had a positive impact on seizure in zebrafish larvae,

while *cox-2* inhibition did not affect seizures [25].

Our results align with these findings. CX2 decreased the levels of *cox1* transcript (Fig 3D) and reduced

the occurrence of seizures by increasing latency and decreasing their frequency (Fig 1B). The

compound was quickly absorbed and distributed, as it was detected in the brain at the earliest time

point examined (Fig 4), and also in the optic tectum, a region associated with significant neuronal

activity during seizures [44].

As for the Tripeptide compound (Fig 2), we observed various gene responses, particularly the down-

regulation of genes related to inflammation. Neuroinflammation, triggered by factors such as tissue

damage, infection, stress, and seizures, has been associated with epilepsy [28,45]. Pathways

connecting epilepsy to neuroinflammation have been identified, with animal models of seizure

induction showing increased mRNA levels of genes involved in inflammatory cascades [46–48]. Thus,

efforts have focused on low molecular weight compounds acting on these pathways. Tripeptide (10

µg/mL) downregulated *il1b* gene and *casp9* gene indicating its positive effect on these pathways.

Regarding the bio-distribution of the Tripeptide compound, it was primarily detected in the zebrafish

larvae intestine (S1 Fig). This could be attributed to the fluorophore, tetramethylrhodamine-TMR,

which along with the compound, may interfere with absorption or crossing the blood-brain barrier for

CNS delivery.

Conclusion

Our study provides evidence that Tripeptide and CX2 – Cx43 peptide have the potential to modulate

gene expression and reduce epileptic seizures in zebrafish. Tripeptide increased the expression of genes

associated with antioxidant functions while decreasing the expression of inflammatory genes, with no

observed behavioral changes. The compound CX2 peptide demonstrated high effectiveness in reducing

the expression of inflammatory genes, lowering the number of seizures, and increasing latency time.

CX2 was widely distributed in the zebrafish brain and the body. These compounds could represent a

promising avenue for further research in the development of novel anti-seizure medications.

Nevertheless, further studies are needed to assess their effects in other animal models and determine

their efficacy in humans.

Reference

1.   Fisher RS, Acevedo C, Arzimanoglou A, Bogacz A, Cross JH, Elger CE, et al. ILAE official
report: a practical clinical definition of epilepsy. Epilepsia. 2014;55: 475–482.
doi:10.1111/EPI.12550

2.   Fisher RS, Cross JH, French JA, Higurashi N, Hirsch E, Jansen FE, et al. Operational
classification of seizure types by the International League Against Epilepsy: Position Paper of
the ILAE Commission for Classification and Terminology. Epilepsia. 2017;58: 522–530.
doi:10.1111/EPI.13670

3.   Duncan JS, Sander JW, Sisodiya SM, Walker MC. Adult epilepsy. Lancet. 2006;367: 1087–
1100. doi:10.1016/S0140-6736(06)68477-8

4.   Thijs RD, Surges R, O'Brien TJ, Sander JW. Epilepsy in adults. Lancet. 2019;393: 689–701.
doi:10.1016/S0140-6736(18)32596-0

5.   Laxer KD, Trinka E, Hirsch LJ, Cendes F, Langfitt J, Delanty N, et al. The consequences of
refractory epilepsy and its treatment. Epilepsy Behav. 2014;37: 59–70.
doi:10.1016/J.YEBEH.2014.05.031

6.   Sirven JI. Epilepsy: A Spectrum Disorder. Cold Spring Harb Perspect Med. 2015;5: a022848.
doi:10.1101/CSHPERSPECT.A022848

7.   Patton EE, Zon LI, Langenau DM. Zebrafish disease models in drug discovery: from
preclinical modelling to clinical trials. Nature Reviews Drug Discovery 2021 20:8. 2021;20:
611–628. doi:10.1038/s41573-021-00210-8

8.   Olivares CI, Field JA, Simonich M, Tanguay RL, Sierra-Alvarez R. Arsenic (III, V), indium
(III), and gallium (III) toxicity to zebrafish embryos using a high-throughput multi-endpoint in
vivo developmental and behavioral assay. Chemosphere. 2016;148: 361–368.
doi:10.1016/J.CHEMOSPHERE.2016.01.050

9.   Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al. The zebrafish
325   reference genome sequence and its relationship to the human genome. Nature. 2013;496: 498–
326   503. doi:10.1038/NATURE12111

327   10.   Hortopan GA, Dinday MT, Baraban SC. Zebrafish as a model for studying genetic aspects of
328   epilepsy. Dis Model Mech. 2010;3: 144–148. doi:10.1242/DMM.002139

329   11.   Lubin A, Otterstrom J, Hoade Y, Bjedov I, Stead E, Whelan M, et al. A versatile, automated
330   and high-throughput drug screening platform for zebrafish embryos. Biol Open. 2021;10.
331   doi:10.1242/BIO.058513

332   12.   Patton EE, Zon LI, Langenau DM. Zebrafish disease models in drug discovery: from
333   preclinical modelling to clinical trials. Nat Rev Drug Discov. 2021;20: 611.
334   doi:10.1038/S41573-021-00210-8

335   13.   Vincent F, Loria P, Pregel M, Stanton R, Kitching L, Nocka K, et al. Developing predictive
336   assays: the phenotypic screening "rule of 3." Sci Transl Med. 2015;7.
337   doi:10.1126/SCITRANSLMED.AAB1201

338   14.   MacRae CA, Peterson RT. Zebrafish as tools for drug discovery. Nat Rev Drug Discov.
339   2015;14: 721–731. doi:10.1038/NRD4627

340   15.   Baraban SC. A zebrafish-centric approach to antiepileptic drug development. Dis Model
341   Mech. 2021;14. doi:10.1242/DMM.049080

342   16.   Baraban SC, Taylor MR, Castro PA, Baier H. Pentylenetetrazole induced changes in zebrafish
343   behavior, neural activity and c-fos expression. Neuroscience. 2005;131: 759–768.
344   doi:10.1016/J.NEUROSCIENCE.2004.11.031

345   17.   Mussulini BHM, Leite CE, Zenki KC, Moro L, Baggio S, Rico EP, et al. Seizures induced by
346   pentylenetetrazole in the adult zebrafish: a detailed behavioral characterization. PLoS One.
347   2013;8. doi:10.1371/JOURNAL.PONE.0054515

348   18.   De Figueiredo SG, De Lima ME, Cordeiro MN, Diniz CR, Patten D, Halliwell RF, et al.
349   Purification and amino acid sequence of a highly insecticidal toxin from the venom of the
350   brazilian spider Phoneutria nigriventer which inhibits NMDA-evoked currents in rat
351   hippocampal neurones. Toxicon. 2001;39: 309–317. doi:10.1016/S0041-0101(00)00129-X

352   19.   Harvey AL, Edrada-Ebel R, Quinn RJ. The re-emergence of natural products for drug
353   discovery in the genomics era. Nat Rev Drug Discov. 2015;14: 111–129.
354   doi:10.1038/NRD4510

355   20.   Newman DJ, Cragg GM. Natural Products as Sources of New Drugs over the Nearly Four
356   Decades from 01/1981 to 09/2019. J Nat Prod. 2020;83: 770–803.
357   doi:10.1021/ACS.JNATPROD.9B01285

358   21.   Bernardes CP, Santos NAG, Sisti FM, Ferreira RS, Santos-Filho NA, Cintra ACO, et al. A
359   synthetic snake-venom-based tripeptide (Glu-Val-Trp) protects PC12 cells from MPP +
360   toxicity by activating the NGF-signaling pathway. Peptides (NY). 2018;104: 24–34.
361   doi:10.1016/J.PEPTIDES.2018.04.012

22.   Martins NM, Santos NAG, Sartim MA, Cintra ACO, Sampaio S V., Santos AC. A tripeptide
isolated from Bothrops atrox venom has neuroprotective and neurotrophic effects on a cellular
model of Parkinson's disease. Chem Biol Interact. 2015;235: 10–16.
doi:10.1016/J.CBI.2015.04.004

23.   Coin I, Beyermann M, Bienert M. Solid-phase peptide synthesis: from standard procedures to
the synthesis of difficult sequences. Nat Protoc. 2007;2: 3247–3256.
doi:10.1038/NPROT.2007.454

24.   Vanier GS. Microwave-assisted solid-phase peptide synthesis based on the Fmoc protecting
group strategy (CEM). Methods Mol Biol. 2013;1047: 235–249. doi:10.1007/978-1-62703-
544-6_17

25.   Barbalho PG, Lopes-Cendes I, Maurer-Morelli CV. Indomethacin treatment prior to
pentylenetetrazole-induced seizures downregulates the expression of il1b and cox2 and
decreases seizure-like behavior in zebrafish larvae. BMC Neurosci. 2016;17.
doi:10.1186/S12868-016-0246-Y

26.   Jarque S, Ibarra J, Rubio-Brotons M, García-Fernández J, Terriente J. Multiplex Analysis
Platform for Endocrine Disruption Prediction Using Zebrafish. Int J Mol Sci. 2019;20.
doi:10.3390/IJMS20071739

27.   Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods. 2001;25: 402–408.
doi:10.1006/METH.2001.1262

28.   Vezzani A, Balosso S, Ravizza T. Neuroinflammatory pathways as treatment targets and
biomarkers in epilepsy. Nat Rev Neurol. 2019;15: 459–472. doi:10.1038/S41582-019-0217-X

29.   Becker AJ. Review: Animal models of acquired epilepsy: insights into mechanisms of human
epileptogenesis. Neuropathol Appl Neurobiol. 2018;44: 112–129. doi:10.1111/NAN.12451

30.   Henshall DC, Simon RP. Epilepsy and apoptosis pathways. J Cereb Blood Flow Metab.
2005;25: 1557–1572. doi:10.1038/SJ.JCBFM.9600149

31.   Aguiar CCT, Almeida AB, Arajo PVP, Abreu RNDC De, Chaves EMC, Vale OC Do, et al.
Oxidative Stress and Epilepsy: Literature Review. Oxid Med Cell Longev. 2012;2012.
doi:10.1155/2012/795259

32.   Vezzani A, French J, Bartfai T, Baram TZ. The role of inflammation in epilepsy. Nat Rev
Neurol. 2011;7: 31–40. doi:10.1038/nrneurol.2010.178

33.   Rana A, Musto AE. The role of inflammation in the development of epilepsy. J
Neuroinflammation. 2018;15. doi:10.1186/S12974-018-1192-7

34.   Jirsa VK, Stacey WC, Quilichini PP, Ivanov AI, Bernard C. On the nature of seizure
dynamics. Brain. 2014;137: 2210–2230. doi:10.1093/BRAIN/AWU133

35. Dinday MT, Baraban SC. Disorders of the Nervous System Large-Scale Phenotype-Based
Antiepileptic Drug Screening in a Zebrafish Model of Dravet Syndrome 1,2,3. 2015 [cited 29
Jul 2022]. doi:10.1523/ENEURO.0068-15.2015

36. Scorza CA, Marques MJG, Gomes da Silva S, Naffah-Mazzacoratti M da G, Scorza FA,
Cavalheiro EA. Status epilepticus does not induce acute brain inflammatory response in the
Amazon rodent Proechimys, an animal model resistant to epileptogenesis. Neurosci Lett.
2018;668: 169–173. doi:10.1016/J.NEULET.2017.02.049

37. Vezzani A, Dingledine R, Rossetti AO. Immunity and inflammation in status epilepticus and
its sequelae: possibilities for therapeutic application.
http://dx.doi.org/101586/1473717520151079130. 2015;15: 1081–1092.
doi:10.1586/14737175.2015.1079130

38. Yagami T, Koma H, Yamamoto Y. Pathophysiological Roles of Cyclooxygenases and
Prostaglandins in the Central Nervous System. Mol Neurobiol. 2016;53: 4754–4771.
doi:10.1007/S12035-015-9355-3

39. Rojas A, Jiang J, Ganesh T, Yang MS, Lelutiu N, Gueorguieva P, et al. Cyclooxygenase-2 in
epilepsy. Epilepsia. 2014;55: 17–25. doi:10.1111/EPI.12461

40. Dey A, Kang X, Qiu J, Du Y, Jiang J. Anti-Inflammatory Small Molecules To Treat Seizures
and Epilepsy: From Bench to Bedside. Trends Pharmacol Sci. 2016;37: 463–484.
doi:10.1016/J.TIPS.2016.03.001

41. Ma L, Cui XL, Wang Y, Li XW, Yang F, Wei D, et al. Aspirin attenuates spontaneous
recurrent seizures and inhibits hippocampal neuronal loss, mossy fiber sprouting and aberrant
neurogenesis following pilocarpine-induced status epilepticus in rats. Brain Res. 2012;1469:
103–113. doi:10.1016/J.BRAINRES.2012.05.058

42. Tanaka S, Nakamura T, Sumitani K, Takahashi F, Konishi R, Itano T, et al. Stage- and region-
specific cyclooxygenase expression and effects of a selective COX-1 inhibitor in the mouse
amygdala kindling model. Neurosci Res. 2009;65: 79–87.
doi:10.1016/J.NEURES.2009.05.013

43. Barbalho PG, Lopes-Cendes I, Maurer-Morelli CV. Indomethacin treatment prior to
pentylenetetrazole-induced seizures downregulates the expression of il1b and cox2 and
decreases seizure-like behavior in zebrafish larvae. BMC Neurosci. 2016;17.
doi:10.1186/S12868-016-0246-Y

44. Turrini L, Fornetto C, Marchetto G, Müllenbroich MC, Tiso N, Vettori A, et al. Optical
mapping of neuronal activity during seizures in zebrafish. Sci Rep. 2017;7.
doi:10.1038/S41598-017-03087-Z

45. Pracucci E, Pillai V, Lamers D, Parra R, Landi S. Neuroinflammation: A Signature or a Cause
of Epilepsy? Int J Mol Sci. 2021;22. doi:10.3390/IJMS22136981

46. Löscher W, Potschka H, Sisodiya SM, Vezzani A. Drug Resistance in Epilepsy: Clinical
Impact, Potential Mechanisms, and New Innovative Treatment Options. Pharmacol Rev.
2020;72: 606–638. doi:10.1124/PR.120.019539

47. Ravizza T, Gagliardi B, Noé F, Boer K, Aronica E, Vezzani A. Innate and adaptive immunity
during epileptogenesis and spontaneous seizures: evidence from experimental models and
human temporal lobe epilepsy. Neurobiol Dis. 2008;29: 142–160.
doi:10.1016/J.NBD.2007.08.012

48. Kim I, Mlsna LM, Yoon S, Le B, Yu S, Xu D, et al. A postnatal peak in microglial
development in the mouse hippocampus is correlated with heightened sensitivity to seizure
triggers. Brain Behav. 2015;5. doi:10.1002/BRB3.403

49. Mukhtar I. Inflammatory and immune mechanisms underlying epileptogenesis and epilepsy:
From pathogenesis to treatment target. Seizure. 2020;82: 65–79.
doi:10.1016/J.SEIZURE.2020.09.015

Supporting information

S1 Fig. Time course of the tripeptide in a 7-days post fertilization zebrafish larva. The tetramethylrhodamine-TMR labeled peptide was not detected in the brain through fluorescence microscopy.

Financial Support

C.V.M.M. received research financial support from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) 2014/15640-8 and the Brazilian Institute of Neuroscience and Neurotechnology (BRAINN) CEPID-FAPESP 2013-07559-3.

L.S. thanks the GI-1251-ACUIGEN research group of the Santiago de Compostela University (Spain) and the ED431C 2018/28 project of the Xunta de Galicia for funding this project.

M.D.M - This work was supported in part through funding from a grant from the Ministry of Science, Innovation and Universities (MICIU/AEI/10.13039/501100011033): PID2022-137027OB-I00 ERDF/EU to MDM. M.D.M also was granted from HORIZON-MSCA-2023-SE-01 (101183034) and EU HORIZON-CSA 101079489.

M.E.V. thanks PID2021-127702NB-I00 funded by MCIN/AEI/10.13039/501100011033 and by ERDF A way of making Europe, the Xunta de Galicia (Centro de investigación do Sistema Universitario de Galicia accreditation 2023-2027, ED431G 2023/03) and the European Union (European Regional Development Fund - ERDF), and the GRC grant ED431C 2021/29 funded by the Conselleria de Cultura, Educación e Universidades da Xunta de Galicia.

467 J.A.A.F. and T.C.M. received a fellowship from Coordenação de Aperfeiçoamento de Pessoal de Nível
468 Superior (CAPES), Brazil, grant number 001. J.A.A.F. also received CAPES/PRINT Proc.
469 88887.470075/2019-00. No sponsors or funders were involved in the study design, data collection
470 and analysis, decision to publish, or preparation of the manuscript.

Competing interests

473 The authors declare that the research was conducted in the absence of any commercial or financial
474 relationships that could be construed as a potential conflict of interest.