White matter connections of human ventral temporal cortex are organized by cytoarchitecture, eccentricity, and category-selectivity from birth

Emily Kubota[1*], Xiaoqian Yan[2], Sarah Tung[1], Bella Fascendini[3], Christina Tyagi[1], Sophie Duhameau[1], Danya Ortiz[1], Mareike Grotheer[4,5], Vaidehi S. Natu[1], Boris Keil[5,6,7,8], & Kalanit Grill-Spector[1,9]

1. Department of Psychology, Stanford University, 450 Jane Stanford Way, Stanford, CA 94305, USA
2. Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai 200433, China
3. Department of Psychology, Princeton University, 40 Woodlands Way, Princeton, NJ 08544, USA
4. Department of Psychology, Philipps-Universität Marburg, Frankfurter Str. 35, Marburg 35037, Germany
5. Center for Mind, Brain and Behavior – CMBB, Universities of Marburg, Giessen, and Darmstadt, Marburg 35039, Germany
6. Institute of Medical Physics and Radiation Protection, TH Mittelhessen University of Applied Sciences, Giessen 35390, Germany
7. Department of Diagnostic and Interventional Radiology, University Hospital Marburg, Philipps-Universität Marburg, Baldinger Str., Marburg 35043, Germany.
8. LOEWE Research Cluster for Advanced Medical Physics in Imaging and Therapy (ADMIT), TH Mittelhessen University of Applied Sciences, Giessen 35390, Germany.
9. Wu Tsai Neurosciences Institute, 288 Campus Drive, Stanford, CA 94305 USA

Corresponding author*: emily.kubota@gmail.com

Abstract

Category-selective regions in ventral temporal cortex (VTC) have a consistent anatomical organization, which is hypothesized to be scaffolded by white matter connections. However, it is unknown how white matter connections are organized from birth. Here, we scanned newborn to 6-month-old infants and adults to determine the organization of the white matter connections of VTC. We find that white matter connections are organized by cytoarchitecture, eccentricity, and category from birth. Connectivity profiles of functional regions in the same cytoarchitectonic area are similar from birth and develop in parallel, with decreases in endpoint connectivity to lateral occipital, and parietal, and somatosensory cortex, and increases to lateral prefrontal cortex. Additionally, connections between VTC and early visual cortex are organized topographically by eccentricity bands and predict eccentricity biases in VTC. These data show that there are both innate organizing principles of white matter connections of VTC, and the capacity for white matter connections to change over development.

Introduction

Human ventral temporal cortex (VTC) contains regions that are selective for categories that are
important for our everyday lives such as faces[1], bodies[2], words[3], and places[4]. A key debate in
cognitive neuroscience is what structural and functional factors contribute to the consistent
functional organization of VTC[5–10]. One central theory is that innate white matter connections
between VTC and other parts of the brain may lead to the emergence of category-selective regions
in consistent anatomical locations relative to cortical folds[11–16]. Indeed, recent research in adults
has revealed that the white matter connections of VTC are highly regular with respect to cortical
folds[17–19], cytoarchitecture[20], eccentricity[21], and category-selectivity[11,12,22]. However, it is unknown
which of these organizing principles of white matter connections of VTC are already present in
infancy and if white matter connections remain stable from infancy to adulthood. Here, we address
these gaps in knowledge by using anatomical and diffusion MRI in infants and adults to elucidate
the organization principles of white matter connections of VTC from birth to adulthood.

We first consider multiple hypotheses of how white matter connections at birth may constrain the
development of VTC function. The *category hypothesis* suggests that category-selective regions in
VTC have category-specific connections forming specialized networks that enable processing of
category-specific information[15,23,24]. While some studies have found that face-selective regions in
VTC only emerge with visual experience[25,26], and develop during the first year of life[27], the existence
of category-selective regions in the congenitally blind[28,29], has led to the hypothesis that innate
white matter connections might scaffold the development of category-selective regions in VTC.
This theory is supported by studies showing that, in adults, white matter connectivity profiles
predict the location of face, place, and word-selective regions in VTC[11,22,30] and in children, white
matter connectivity profiles at age five predict where word-selective regions will emerge at age
eight[13]. Further, functional connectivity of resting state data in infants show category-specific
patterns[31,32], suggesting that category-specific networks may be present from birth. The category
hypothesis thus predicts that white matter connections of VTC will be organized by category from
birth.

The *cytoarchitecture hypothesis* suggests that white matter connections are linked to
cytoarchitectonic areas – areas defined by their distribution of cell density across cortical layers[33–
35]. VTC contains four cytoarchitectonic areas: FG1-FG4[36,37], which have different neural hardware
that is thought to support different functional computations. Notably, in both children and adults,
regions with different category-selectivity within the same cytoarchitectonic area, e.g., face-
selective and word-selective regions located in FG4, have similar white matter connectivity
profiles[20]. But regions with the same category-selectivity located in different cytoarchitectonic
regions, e.g., face-selective regions located in FG2 and FG4, respectively, have different white
matter connectivity profiles[20]. The cytoarchitectonic hypothesis therefore predicts that white
matter connections of VTC will be organized by cytoarchitecture from birth.

The *eccentricity-bias hypothesis* suggests that eccentricity biases in VTC are due to innate white
matter connections with early visual cortex (EVC, union of V1-V2-V3), where faces and word-
selective regions have more connections with foveal EVC and place-selective regions have more
connections with peripheral EVC. This theory is supported by findings that in both children and
adults, face and word-selective regions in VTC have a foveal bias[5,7,8,38], whereas place-selective
regions in VTC have a peripheral bias[5,7,8,38]. Additionally, patterns of functional connectivity between
face- and place-selective regions in VTC and EVC in infant humans, neonate monkeys, and
congenitally blind adults follow eccentricity bands[39-43]. That is, face-selective regions have higher
functional connectivity to central eccentricity bands in EVC and place-selective regions have higher
functional connectivity to peripheral eccentricity bands in EVC. Finally, in adults, white matter
connections between VTC and EVC also correspond to eccentricity bands, with face-selective
regions having more connections to central EVC eccentricities and place-selective regions having
more connections to peripheral ones[21]. The eccentricity-bias hypothesis thus predicts that the
subset of white matter connections between VTC to EVC will have a non-uniform distribution,
where regions in VTC that have foveal bias in adults (faces and words) will have more connections
with central bands in EVC from birth, and regions that have a peripheral bias (places) will have more
connections with peripheral bands in EVC from birth. Crucially, when experiments only include the
categories of faces and places, it is not possible to distinguish between the category and
eccentricity hypotheses because there is one-one-mapping between category and eccentricity
bias. In the present study, we include multiple categories that have a foveal bias (faces and words),
which will allow us to distinguish the category from the eccentricity hypotheses.

In addition to the unknown organizational principles of white matter connections in infancy, it is
also unknown if white matter connectivity profiles are stable or change over development.
Accumulating evidence reveals that the large fascicles (axonal bundles that travel in parallel
connecting distant parts of the brain) are present at birth[44-48]. Nonetheless, some fascicles
continue to develop after birth. For example, the arcuate fasciculus (AF), which connects the VTC
with lateral prefrontal cortex is not fully developed in infancy: it has a smaller cross section in
children compared to adults[49] and, and in infants, it reaches premotor cortex but not lateral
prefrontal cortex as it does in adults[50]. Additionally, mature white matter connections in adulthood
may depend on visual experience as visual deprivation during infancy and childhood leads to
degradation of white matter tracts of the visual system[51-55]. However, it is unknown which aspects
of VTC white matter connectivity profiles are innate and which aspects may develop from infancy
to adulthood.

To address these gaps in knowledge, we obtained diffusion magnetic resonance imaging (dMRI)
and anatomical MRI in infants and adults and evaluated the organization and development of white
matter connections of category-selective regions of VTC.

Results

We collected MRI data from 43 newborn to 6-month-old infants (26 longitudinally) during natural sleep using 3T MRI over 75 sessions, as well as from 21 adults (28.21 ± 5.51 years), obtaining whole brain anatomical MRI and multishell dMRI data in each participant. We implemented several quality assurance measures to ensure high quality data (Methods: Quality Assurance). Two infant sessions were excluded because of missing diffusion data, and six infant sessions were excluded after quality assurance. We report data from 67 infant sessions (42 infants (16 female, 26 male); 21 longitudinal infants; timepoints in Supplementary Figure 1) and 21 adult sessions (17 female, 4 male) with no significant differences in data quality (see Methods). In total, we report data from 88 sessions: 23 sessions from newborns (mean age$\pm$standard deviation: 28.6 ± 10.2 days), 23 sessions from 3-month-olds (106.9 ± 19.3 days), 21 sessions from 6-month-olds: (189.0 ± 15.8 days), and 21 sessions from adults (28.21 ± 5.51 years). Anatomical MRIs were used to segment the brain to gray and white matter, define the gray-white matter boundary to seed the tractography, and create cortical surface reconstructions. dMRI data was used to derive the whole brain white matter connectome of each individual and session.

To determine how white matter connections are organized in infants and adults and how they may constrain the development of regions in VTC, we project adult functional regions of interest (fROIs) to the native space of each individual and determine the white matter connections of each fROI. To do so, we use cortex based alignment[56] to project maximal probability maps (MPMs) of six adult category-selective regions from independent data[20] into each individual brain and session. The fROIs are: mFus-faces (face-selective region located near anterior end of the mid fusiform sulcus), pFus-faces (posterior fusiform face-selective region), OTS-bodies (body-selective region located on the occipital temporal sulcus), CoS-places (place-selective region located on the collateral sulcus), mOTS-words (word-selective region located on the occipital temporal sulcus), and pOTS-words (posterior OTS word-selective region). MPMs of fROIs selective for faces, bodies, and places (mFus-faces, pFus-faces, OTS-bodies and CoS-places) are bilateral, and MPM fROIs selective for words (mOTS-words and pOTS-words) are in the left hemisphere only (Fig 1A,B).

To verify that fROIs are not proportionally different in size across age groups, which could affect the resulting white matter connectivity profiles, we measured the surface area of fROIs relative to the surface area of each hemisphere in all infant and adult sessions. We tested if the relative fROI size changes over development (as development plateaus, we use a logarithmic model: fROI size/surface area ~ log10(age in days)). We find no relationship between the surface area of fROIs relative to hemisphere surface area and age ($t(878) = 8.77\times10^{-6}$, $p = 0.88$, adjusted $R^2 = -0.001$, 95% CI = [-0.0001,0.0001]).

Figure 1. Functional atlas and analysis pipeline. A) mFus-faces aligns to the mid-fusiform sulcus in the fsaverage brain space (left) and B) the newborn brain space (right). C) *Left:* Functional region of interest (fROI) atlas from 28 adults in[20] and Atlas of cytoarchitectonic areas from[59] projected to the fsaverage brain; *Right:* the same atlases projected into an example newborn's brain (17 days old). Solid colors represent fROIs. Salmon *pink:* mFus-faces (mid fusiform face-selective region); *Light blue:* mOTS-words (mid occipital temporal sulcus word-selective region); *Yellow:* OTS-bodies (occipital temporal sulcus body and limb-selective region); *Red:* pFus-faces (posterior fusiform face-selective region); *Blue: p*OTS-words (posterior occipital temporal sulcus word-selective region). Green: CoS-places (collateral sulcus place-selective region). Outlines represent cytoarchitectonic areas[59]. *Cyan:* FG2; *Green:* FG3; *Magenta:* FG4. Both fROIs and cytoarchitectonic areas align to the expected anatomical landmarks in the newborn. All individuals in Supplementary Figs 2-9. D) Pipeline to define white matter connectivity profiles for each fROI. *Left:* example connections of mFus-faces in an individual newborn; *Middle*: endpoint map on the cortical surface; *Right:* quantification of endpoint density within each Glasser ROI. E) Principal Components Analysis plot across the first two principal components: explaining 38% and 20% of the variance, respectively. Each dot represents a single connectivity profile in a single subject; colored dot represents the connectivity profile shown in (D). (F-H) Schematics illustrating what the data may look like in PC space according to the predictions of three hypotheses. Each dot represents a connectivity profile for a given fROI and session. (F) Category hypothesis. *Green:* place-selective, *blue:* word-selective, *red:* face-selective, *yellow:* body-selective. G) Cytoarchitectonic

To ensure that fROIs and cytoarchitectonic areas align to the expected anatomical landmarks in
each infant and adult session, we visually inspected each session to ensure that there is a
consistent relationship between projected ROIs and anatomical landmarks. First, we checked
whether the collateral sulcus (CoS) and mid fusiform sulcus (MFS) are present in both
hemispheres. We found that the anatomical landmarks were present in nearly each infant (CoS:
67/67, MFS: 66/67) and adult (CoS: 21/21, MFS: 19/21) session. Next, we tested whether CoS-
places aligns to the intersection of the anterior lingual sulcus and CoS[6,9], mFus-faces aligns to the
anterior tip of the MFS[57], and whether the boundaries between FG1 and FG2 and between FG3 and
FG4 are aligned to the MFS, as in prior studies[20,58]. We find that both fROIs and cytoarchitectonic
areas are aligned to the expected anatomical landmarks in nearly each infant (fROIs; LH: 66/67,
RH: 65/67, cytoarchitectonic areas; LH: 67/67, RH: 62/67) and adult (fROIs: LH: 21/21, RH: 21/21,
cytoarchitectonic areas; LH: 20/21, RH: 21/21) (Supplementary Figs 2-9). Therefore, in both infants
and adults there is a consistent relationship between MPMs of category-selective regions and
cytoarchitectonic areas[20,58], where pOTS-words and pFus-faces are within FG2, CoS-places is
within FG3, and mFus-faces, OTS-bodies, and mOTS-words are within FG4 (Fig 1C).

We use a data-driven approach to determine the organizing principles of white matter connections
of VTC. As the category and cytoarchitecture hypotheses make predictions about connectivity
across the whole brain, whereas the eccentricity-bias hypothesis makes predictions only about
connectivity with EVC, we first test if whole brain connectivity profiles are organized by category-
selectivity or cytoarchitecture. In addition, we test if whole brain connectivity profiles are organized
by age to determine if connectivity profiles are stable or changing over development. We separately
examine the eccentricity-bias hypothesis in the next section.

We derived each fROI's white matter connections by intersecting each fROI with the whole brain
white matter connectome derived from dMRI in each session (Fig 1D). We quantified the white
matter connectivity profile by measuring the endpoint density of each fROI's connections on the
whole brain as the proportion of connections ending in each of 169 Glasser Atlas ROIs[60] (180
Glasser ROIs excluding 11 VTC ROIs, Fig 1D). This resulted in 880 unique connectivity profiles (10
fROIs x 88 sessions across infants and adults). Then we used principal component analysis (PCA)
to reduce the dimensionality of the connectivity profiles. The first 10 principal components (PCs)
explain 98% of the variance in the data, with the first and second components explaining 38% and
20%, respectively.

To visualize how connectivity profiles cluster, we plot the connectivity profiles of all fROIs and
sessions in PC space (Fig 1E). The plot consists of 880 dots including data from all infants and
adults, where each dot represents a single connectivity profile of an fROI from one session. Note
that each white matter connectivity profile has several non-mutually exclusive features: the
category-selectivity of the fROI, the cytoarchitectonic area in which the fROI is located, and the age

of the participant. Therefore, we can label each connectivity profile by each of these features
(category, cytoarchitecture, age group) and test if connectivity profiles cluster by one or more of
these features. For example, if connectivity profiles are organized by category, then connectivity
profiles of fROIs with the same category-selectivity would be nearby in PC space and separate from
connectivity profiles of fROIs with a different category-selectivity (Fig 1F). However, if connectivity
profiles are organized by cytoarchitecture, then connectivity profiles of fROIs in the same
cytoarchitectonic area would be nearby in PC space (Fig 1G). Further, if connectivity profiles change
across development, then connectivity profiles of newborns may be different than other age groups
and connectivity profiles of adults may be most distinct (Fig 1H).

Figure 2. White matter connections are organized by cytoarchitecture and category from infancy. (A-C) White matter
connectivity profiles of all participants (n = 88) and fROIs (n=10) projected to the first two principal components (PC1
and PC2); each dot is a connectivity profile of a single fROI and session. The same data is shown in all three panels,
except that they are colored by different features. (A) Category: *Red*: faces, *green:* places, *blue*: words, *yellow:* bodies, (B)
Cytoarchitectonic area: *Cyan:* FG2, *green:* FG3, *magenta:* FG4. *FG:* fusiform gyrus. (C) Age group: *Salmon:* newborn, *green:*
3 months, *teal:* 6 months, *purple*: adults. (D-F) Classification accuracy of a feature from white matter connectivity profiles,
using multinomial logistic regression with leave-one-out-participant cross validation. In D-F, Rows sum to 100. Color
depicts the percentage of samples classified within each bin; brighter colors indicate higher percentage of samples (see
color bar). On diagonal values are percentage correct classification and off diagonal values are percentage incorrect
classification (D) Confusion matrix for classification of category. (E) Confusion matrix for classification of
cytoarchitectonic area. (F) Confusion matrix for classification of age group. Overall classification accuracy is found in
Supplementary Figure 11.

Labeling connectivity profiles by fROIs' preferred category reveals some clustering by category (Fig
2A). For example, connectivity profiles of face- and place-selective ROIs tend to be clustered and

separate from others. However, clustering by category is imperfect, as connectivity profiles of
word-selective regions split into two clusters, and there is overlap between the connectivity profiles
of word- and body-selective fROIs. In contrast, labeling connectivity profiles by fROIs'
cytoarchitectonic area reveals a strikingly clear organization whereby connectivity profiles of each
of the fusiform gyrus (FG) cytoarchitectonic areas FG2, FG3, FG4 form a different cluster with a
distinct connectivity profile (Fig 2B). Examination of the PC loadings (Supplementary Fig 10) and
the coefficients across the first 2 PCs suggests that connectivity profiles of fROIs in different
cytoarchitectonic areas largely vary in their connections to early visual, lateral occipito-temporal,
and anterior temporal cortex. Finally, labeling connectivity profiles by participant's age group
reveals that connectivity profiles are intermixed across age groups with no clear organization by
age across the first 2 PCs (Fig 2C).

To utilize the information across all 10 PCs, we use a leave-one-out classification approach to test
if a classifier trained on white matter connectivity profiles of n-1 participants can predict the
category preference, cytoarchitectonic area, or the participant's age group from the left-out
participant's connectivity profiles. We find that category (mean accuracy±SD = 83%±10%, 95% CI =
[81%, 85%], t(87) = 54.68, p < 0.001, Cohen's d = 5.83), cytoarchitecture (mean accuracy±SD=
97%±5%, 95% CI = [95%, 98%], t(87) = 109.6, p < 0.001, Cohen's d = 11.68), and age (mean
accuracy±SD = 41%± 27%, 95% CI = [35%, 46%], $t(87)$ = 5.48, p < 0.001, Cohen's d = 0.58) are all
classified significantly above chance (Supplementary Fig 11). Classification of cytoarchitecture is
significantly higher than classification of category preference (odds ratio= 5.96, 95% CI [3.98 8.93],
binomial logistic regression) and age (odds ratio = 41.51, 95% CI [28.16, 61.19], binary logistic
regression). Confusion matrices reveal almost no error for cytoarchitecture classification (Fig 2E),
but some category errors that arise from confusion of word-and body-selectivity as well as
confusion of body and face-selectivity (Fig 2D). The confusion matrix for age classification (Fig 2F)
reveals that newborns and adults are classified the best, that 3-month-olds are often confused for
newborns, and that 6-month-olds are equally confused across all age groups. This suggests that
there is heterogeneity in the connectivity profiles of 6-months-olds, and that around 6 months of
age, the white matter connections of some infants may start to become more adult-like, while
connections in others resemble 3-month-olds and even newborns.

As there are unbalanced training sets (e.g., two face-selective vs. one body-selective fROI), we
performed an additional analysis where we created a training set that had an equal number of
examples for each label by sampling the training set with replacement. We find similar
classification results with balanced examples in the training data (Supplementary Fig 12).

To assess if the high classification performance of cytoarchitecture is due to higher anatomical
proximity of fROIs within the same cytoarchitectonic area compared to those in distinct
cytoarchitectonic areas, we repeated the classification analysis on connectivity profiles of
equidistant disk ROIs that are either in the same (FG4) or different (FG2/FG4) cytoarchitectonic
areas (Supplementary Fig13A,B). We reasoned that if anatomical proximity explains classification
results, then classification accuracy would be reduced once distance is controlled. However, even

when distance was held constant, cytoarchitecture is classified from the connectivity profiles with
high accuracy (Supplementary Fig13C, mean accuracy±SD=95%±22%, t(87) = 31.62, p < 0.001,
Cohen's d = 3.37, 95% CI = [0.92,0.98]), suggesting that organization of connectivity profiles by
cytoarchitecture is not just due to anatomical distance between fROIs.

Interestingly, both category and cytoarchitecture classification are not significantly different across
age groups (comparison between newborns, 3-month-olds, and 6-month-olds compared to adults
(newborns: odds ratio =1.20, 95% CI [0.77, 1.89], 3-months: odds ratio =1.01, 95% CI [0.65, 1.56], 6-
months: odds ratio =1.05, 95% CI [0.67, 1.65], binomial logistic regression). Indeed, examination of
the organization structure across the first 2 PCs separately for each age group reveals similar
organization across age groups (Supplementary Fig 14). These analyses suggest that
organizational features of white matter connections by cytoarchitecture and category persist
across development.

Are Connections Between VTC and EVC Organized Retinotopically?

To test the eccentricity-bias hypothesis we next examined the subset of connections between each
fROI and early visual cortex (EVC). We reasoned that if connections between VTC and EVC are
organized by eccentricity from birth, then fROIs that overlap with foveal representations in adults
(pFus-faces, pOTS-faces, mFus-faces, OTS-bodies, mOTS-words) would have more white matter
connections to central than peripheral eccentricities in EVC in infancy, and fROIs that overlap with
peripheral representations in adults (CoS-places) would have more connections to peripheral than
central eccentricities in VTC in infancy. To test these predictions, we measured the distribution of
white matter endpoints between each of the VTC category-selective fROIs and three eccentricity
bands in EVC: 0-5°, 5-10°, and 10-20° and compared across age groups.

We first visualized the white matter connections between each fROI and EVC, coloring the white
matter connections according to the endpoint eccentricity in EVC. This visualization reveals a
striking orderly arrangement of the connections from each eccentricity band to each of the fROIs
in both infants and adults (Fig 3A and 3B; 3D visualization: Supplementary Videos 1-4). Specifically,
in all age groups, newborns to adults, there is a lateral to medial arrangement of connections such
that connections to EVC central eccentricities (red, 0-5°) are more lateral, and connections to more
peripheral EVC eccentricities are more medial (green, 5-10° to blue, 10-20°; Fig 3A,B).

Figure 3. White matter connections of VTC are organized by eccentricity from birth. Connections between each fROI and early visual cortex (EVC, union of V1, V2, V3) in an example newborn (A) and an example adult (B). Here we show data for the left hemisphere as it includes all 6 fROIs; data for the right hemisphere is in Supplementary Fig 56. Connections are colored by the endpoint eccentricity band in EVC: *Red:* 0-5°, *green*: 5-10°, *blue:* 10-20°. (C) Quantification of the percentage of endpoints in each eccentricity band by age group for each fROI. Violin plots indicate distributions of the percentage of connections for each age group and eccentricity band. *Black dot and error bars*: mean± standard error of the mean. *Red:* 0-5°, *green*: 5-10°, blue: 10-20°, see inset. *0M:* newborns (n = 23), *3M:* 3 month-olds (n = 23), *6M:* 6 month-olds (n = 21), *A*: adults (n = 21). *Bottom horizontal lines:* cytoarchitectonic area.

We then quantified the distributions of endpoint connections to EVC for each fROI and age group. Consistent with the predictions of the eccentricity-bias hypothesis, in all age groups, pFus-faces, pOTS-words, mFus-faces, OTS-bodies, and mOTS-words have more connections to the EVC 0-5° eccentricity band than the more peripheral eccentricities (Fig 3C) and CoS-places has more connections to the EVC 10-20° eccentricity band than to the more central eccentricities, which may be supported by the medial occipital longitudinal tract[61] (Fig 3C). Additionally, the connections from VTC to EVC by eccentricity band also mirror the cytoarchitectonic organization of the VTC connectivity profiles. Specifically, across all age groups: (i) pFus-faces and pOTS-words, which are located in cytoarchitectonic area FG2, have ~60% of their connections to the EVC 0-5° eccentricity band, whereas (iii) CoS-places, which is located in FG3, has ~40% of its connections to the EVC 10-20° eccentricity band, and (iii) mFus-faces, OTS-bodies, and mOTS-words in FG4 have 40-50% of their EVC connections in the central 5° (Fig 4C).

We further tested how eccentricity is related to the features of cytoarchitecture, category, and age that we examined in the whole brain analysis. To do so, for each fROI we fit a linear model predicting the percentage of connections to the central 5° by cytoarchitecture, category-selectivity, and age group. As the percentage of connections to each eccentricity band is proportional, we use the percentage of connections to the central 5° as the independent variable, where a greater

percentage connections to the central 5° indicate more connections to central eccentricity bands,
and fewer connections to the central 5° indicate more connections to peripheral eccentricity bands.

We find a main effect of cytoarchitecture ($F(2,504) = 390.02$, $p < 0.001$, $\eta^2 = 0.61$), suggesting that
regions that are in distinct cytoarchitectonic areas have different percentages of connections with
the central 5° of EVC. In particular, the majority of connections between FG2 and EVC are in the
central 5°, but FG3 has a minority of connections in the central 5° (mean±sd: FG2: 63.57%±14.41%,
FG3: 26.58%±8.33%, 4: 43.97%±9.34%). Additionally, there is a main effect of category-selectivity
($F(2,504) = 8.32$, $p < 0.001$, $\eta^2 = 0.03$), reflecting that regions with different category-selectivity also
have different percentages of connections with the central 5° of EVC. Face- and word-selective
regions have a majority of connections with the central 5°, but place-selective regions have only a
minority (mean±sd: faces: 51.70%±15.02%, words: 56.21%±15.73%, places: 26.58%±8.33%,
bodies: 43.23%±9.05%). In contrast, there is no main effect of age group ($F(3,504) = 1.14$, $p = 0.33$,
$\eta^2 = 6.77 \times 10^{-3}$), and no significant interaction between category and age ($F(6,504) = 0.93$, $p = 0.47$,
$\eta^2 = 0.01$). There are, however, significant interactions between cytoarchitecture and category
($F(1,504) = 24.55$, $p < 0.001$, $\eta^2 = 0.05$), cytoarchitecture and age ($F(6,504) = 3.59$, $p = 0.002$, $\eta^2 =$
0.04), and between cytoarchitecture, category, and age ($F(3,504) = 2.76$, $p = 0.04$, $\eta^2 = 0.02$). Post-
hoc tests reveal that the cytoarchitecture by category interaction is driven by differences in
percentage of connections with the central 5° between pFus-faces and pOTS-words in FG2 ($t(87)$
$= 5.68$, $p < 0.001$, Bonferroni corrected, Cohen's d = 0.75, mean difference = -10.09, 95% CI = [-13.63,
-6.56]), as well as differences in percentage of connections to the central 5° between mFus-faces
and OTS-bodies in FG4 ($t(87) = 3.68$, $p = .0016$, Bonferroni corrected, Cohen's d = 0.17, mean
difference = 1.08, 95% CI = [0.76, 2.55]). Further post-hoc tests reveal that differential development
is due to a decrease in the percentage of connections from FG4 with the EVC central 5° over
development ($\beta = -2.75$, 95% CI = [-4.34, -1.16], $t(163.09) = -3.39$, $p = 0.0027$, Bonferroni corrected),
and a differential development within FG2 where there is a significant increase in the connections
of pOTS-words with central 5° of EVC over development ($\beta = 3.95$, 95% CI = [1.75, 6.14], $t(81.26) =$
3.52, $p = 0.0042$, Bonferroni corrected), but no significant development in the percentage of pFus-
faces connections with the central 5° ($\beta = -0.78$, 95% CI = [-4.05, 2.49], $t(85.90) = -0.47$, $p = 1$,
Bonferroni corrected). These results suggest that in addition to cytoarchitecture, and category,
white matter connections of VTC are organized topographically by eccentricity from birth and that
the differential proportion of connections between VTC and different eccentricity bands from birth
mirrors the eccentricity bias of both cytoarchitectonic areas and category-selective regions in VTC.

Do Connections of VTC Develop from Infancy to Adulthood?

In addition to revealing the organizing principles of white matter connections by cytoarchitecture,
category, and eccentricity, we also found evidence for development: we were able to classify the
age group of the participant above chance from the whole connectivity profile, and we found some
developmental changes in connectivity with EVC. To qualitatively assess how white matter
connections of VTC change over development, we visualized the white matter connections of all
category-selective fROIs in all participants. Fig 4 shows an example infant at newborn, 3 months,

Figure 4. Connectivity of VTC develops from infancy to adulthood. (A) Connections of each fROI in an individual infant across three time points (newborn, 3 months, 6 months) next to an example adult shown in a sagittal cross section. Brains are shown to scale, scale bar, bottom left. Each row is a different fROI ordered by cytoarchitectonic area; from top to bottom: FG2 to FG4. (B) Change in endpoint density over development for each fROI's connectivity profile. The color of the Glasser ROI indicates the development in endpoint density over time, namely the slope of the regression: endpoint density ~ log10(age in days). *Green:* decreasing endpoint density over development, *magenta:* increasing endpoint density over development. *Black arrow:* primary visual area (V1); *Green arrow:* primary somatosensory area (S1). Each row shows the development of white matter connections of an fROI; rows ordered identical to (A). Here we show data for the left hemisphere; data for the right hemisphere is in Supplementary Fig 55.

Consistent with the prior analyses, within an age group, white matter connections of fROIs in the same cytoarchitectonic area were similar. For example, in the example newborn, pFus-faces and

pOTS-words, both in FG2, are characterized by vertical connections to dorsal occipital cortex and
longitudinal connections to both the anterior temporal lobe and early visual cortex (Fig 3A, top 2
rows). However, we also observe developmental changes. For example, pFus-faces and pOTS-
words appear to have more abundant vertical connections in infants than adults, but more
connections to the frontal cortex in adults than infants (Fig 3A, Supplementary Figs 23-30, 51-54
all participants). We see a similar pattern for the white matter connections of fROIs in FG4: mFus-
faces, OTS-bodies, and OTS-words show similar connectivity profiles at each timepoint (e.g.,
newborn), yet we also observe developmental differences: vertical connections appear more
abundant in newborns than adults, but connections to the frontal cortex appear to be more
abundant in adults than infants (Fig 3A, bottom row, Supplementary Figs 15-22, 31-38,47-50). For
CoS-places, located medially in FG3, there is a different connectivity profile and a different
developmental pattern: in newborns, CoS-places has abundant frontal, vertical, and longitudinal
connections to both occipital and anterior temporal lobes, but in adults there appear to be fewer
frontal and vertical connections (Fig 3A-third row, Supplementary Figs 39-46).

To quantify developmental changes, we tested whether each fROI's endpoint density, quantified as
the proportion of connections in each Glasser ROI, varies as a function of age (fROI endpoint
density ~ log10(age in days) x Glasser ROI, linear model). As expected, fROIs' endpoint density vary
across the brain (significant main effect of Glasser ROI; Fs > 74.94, ps < 0.001; statistics in
Supplementary Table 1) reflecting that fROIs have non-uniform connectivity across the brain.
Additionally, fROIs' endpoint density differentially vary with age across the brain (significant age by
Glasser ROI interaction; Fs > 3.54, ps < 0.001; statistics in Supplementary Table 1; repeated with
correction for gestational age in Supplementary Table 2). As we find a differential development
across the brain, for each fROI, we calculate the change in endpoint density across age per Glasser
ROI (endpoint density ~ log10(age in days), linear model). Then, we visualize the slope of this model
for each Glasser ROI (Fig 4D). A positive slope (magenta) indicates that the endpoint density within
the Glasser ROI increases from infancy to adulthood, whereas a negative slope (green) indicates
that endpoint density decreases with age.

This analysis reveals both developmental increases and decreases in endpoint density with distinct
spatial characteristics. First, there are common developmental patterns for face, word, and body
fROIs (Fig 4-right): endpoint densities to lateral occipital and dorsal occipito-parietal visual Glasser
ROIs as well as somatosensory areas decrease (Fig 4-green arrows), but endpoint densities to
lateral prefrontal Glasser ROIs increase (Fig 4). There is also differential development of white
matter connections of fROIs in FG2 (pOTS-words, pFus-faces) compared to FG4 (mFus-faces
mOTS-words, and OTS-bodies): the former show developmental decreases in endpoint densities in
V1 and early visual cortex more broadly (Fig 4-medial view) but the latter show developmental
increases. Finally, for CoS-places, which is in FG3, we find mostly developmental decreases in
endpoint densities in both visual and orbitofrontal cortex with increases in endpoint densities in
ventral temporal lateral regions.

Finally, we tested whether development was more similar for fROIs with the same cytoarchitectonic
area or same category-selectivity. We used a bootstrapping procedure to estimate the
developmental slopes (Methods) and evaluated the similarity between white matter development
by calculating the pairwise correlation between the development slopes across Glasser ROI for
each pair of fROIs. We then fit a linear model to test whether the correlation was higher for fROIs
with the same cytoarchitecture or category-selectivity (linear model: Fisher transformed correlation
~ cytoarchitecture + category, adjusted R^2 = 0.81). We find a significant main effect of
cytoarchitecture (t(14997) = 248.0, p < 0.001, 95% CI = [0.88, 0.89]), reflecting that regions within
the same cytoarchitectonic area had more similar developmental slopes (correlation±sd: .78±11)
than regions in different cytoarchitectonic areas (correlation±sd: .25±13). There was also a
significant main effect of category (t(14997) = 32.7, p < 0.001, 95% CI = [0.14, 0.16]). However, here
we find that regions with the same category-selectivity had less similar developmental slopes
(correlation±sd: .36±16) than regions with different category-selectivity (correlation±sd: .40±.28).
These analyses suggest that although the relationship between white matter and cytoarchitecture
is consistent over development, connectivity profiles develop in parallel within each
cytoarchitectonic area, with both developmental increases and decreases in endpoint densities.

Discussion

Here, we use anatomical and diffusion MRI in infants within the first six months of life and in adults
to determine the organizing principles and developmental trajectories of the white matter
connections of regions in VTC. We find that white matter connections are organized by
cytoarchitecture, eccentricity, and category from birth. We also find evidence for development of
white matter connections of VTC, with increasing endpoint density to lateral frontal cortex, and
decreasing endpoint density to lateral occipital, parietal, somatosensory, and orbitofrontal cortex.
These findings have important implications for understanding the interplay between nature and
nurture on both white matter connections and functional brain development.

Cytoarchitecture and eccentricity are organizing principles

Here, we tested theories that suggest that innate white matter connections support the
organization of VTC. When we use the term innate, we refer to the organization of white matter
connections at birth, prior to the experience of patterned vision outside the womb. A predominant
theory suggests that the consistent organization of category-selective regions in VTC is due to
innate white matter connections that support category-specific processing[11,12,15,31]. While we find
that in infancy white matter connections are organized to a certain extent by category, especially
for faces and places, several of our empirical findings suggest that cytoarchitecture and
eccentricity are more parsimonious organizing principles of white matter connectivity from infancy
to adulthood. Even as both category and cytoarchitecture can be classified from white matter
connectivity profiles irrespective of age, classification of cytoarchitecture is higher than category-
selectivity. In addition, the subset of connections between VTC and early visual cortex are
organized topographically by eccentricity bands and show similar eccentricity biases for functional

regions that are selective for different categories but located in the same cytoarchitectonic area.
Finally, connectivity profiles of functional regions in the same cytoarchitectonic area largely
develop in parallel from infancy to adulthood.

This organization by cytoarchitecture and eccentricity suggests that these are general principles
underlying the white matter organization of the visual system, which may predate and constrain
the coupling of white matter connections and brain function. That is, our data suggest a theoretical
shift: rather than humans being born with specialized connections to support category-specific
processing, cytoarchitectonic boundaries and eccentricity gradients are the innate organizing
principles of white matter connections of the visual system. Indeed, in children and adults
cytoarchitecture and eccentricity are linked as the boundary between lateral and medial
cytoarchitectonic areas in VTC align to the transition between foveal and peripheral
representations in VTC[20,57,58,62]. As we find that white matter connections are organized by
eccentricity from birth, one functional prediction is that eccentricity biases in VTC[7,8] will also be
present from birth. Nonetheless, we underscore that white matter connections start developing
before birth during the second and third trimesters of gestation[63]. Thus, an important direction for
future research is understanding how the prenatal environment in the womb[63] and retinal waves[64]
prior to birth contribute to the organization of white matter connections of the visual system at
birth.

As cytoarchitectonic[35,59,65–68] and retinotopic organization[7,8,21,69–71] are prevalent across the entire
visual system, we hypothesize that these two organizing principles may underlie the white matter
organization of the visual system more broadly. This hypothesis can be tested in future research,
for example, in the parietal cortex that contains a series of retinotopic[72,73] and cytoarchitectonic[66]
areas. Critically, as cytoarchitecture[35,60,7435,75–78], and topography[79,80] are also key features
distinguishing brain areas beyond the visual system, e.g., language[81], our findings raise the
possibility that white matter connections are innately organized by cytoarchitecture and
topographic gradients even beyond the visual system and throughout the entire brain. This
hypothesis can be tested in future research leveraging large anatomical and diffusion MRI datasets
that are being collected in infants[82–86].

Development of white matter connections of VTC

While we find innate organization of white matter connections, we also find that the connectivity
profiles of fROIs within the same cytoarchitectonic area develop in parallel from infancy to
adulthood. Endpoint density from VTC to the frontal lobe showed both developmental increases
and decreases. Endpoint density from FG2 and FG4 in lateral VTC to prefrontal cortex increased
over development but from FG3 in medial VTC to orbitofrontal cortex decreased. The former finding
is consistent with observations that the arcuate fasciculus[87] is underdeveloped in infants, only
reaching the precentral gyrus and premotor cortex in newborns, and not reaching lateral prefrontal
cortex as it does in adults[50,88].

Additionally, we find developmental decreases in endpoint density between VTC and lateral
occipital, parietal, and somatosensory cortex. Our observations in the human visual system mirror
results from a large body of tracer studies in cats and macaques that documented exuberant
connections between V1, V2, and V3[89−91] as well as between early visual areas and somatosensory
areas[92]. These exuberant connections are present early in development and then eliminated by
adulthood[90,92−94]. Our data not only suggest that exuberant connections may exist in the human
visual system, but also suggest the possibility that connections between VTC and both the lateral
and dorsal visual streams[17,95−98] may decrease over development.

We acknowledge that our *in-vivo* dMRI measurements do not enable us to make inferences about
the cellular and molecular underpinnings of the development of white matter connectivity profiles,
and there are multiple possible underlying causes of the developmental effects that we observed.
For example, increases in connectivity may reflect a pathway that exists at birth and becomes
myelinated over development[45], or a pathway that exists at birth and becomes expanded over
development[49]. Likewise, the observation of decreasing connections may reflect decreases in
connectivity or increases in crossing fibers. Future studies can examine if developmental increases
in endpoint density may be related to increased myelination[99−102], axonal sprouting[103−105], and glial
proliferation[100,102] that are associated with activity-dependent white matter plasticity[106,107].
Additionally, future studies can test whether developmental decreases in endpoint density may be
related to elimination of exuberant long range connections[90,92−94].

Implications for theories of functional development of VTC

Our findings that the white matter connections of VTC have both innate organization and the
capacity to change over development raise questions about the relationship between white matter
and function in atypical development. One question is whether lack of visual experience may
reshape white matter connectivity profiles of VTC and consequently function[14]. For example, it is
unknown if lack of visual experience may lead to the preservation of connections between VTC and
somatosensory cortex, which in turn, may enable VTC to respond to haptic inputs in individuals
who are congenitally blind[108]. Critically, our framework enables measuring the fine-grained white
matter connections of VTC in individual infants longitudinally[109], increasing accuracy and precision
in measuring the interplay between white matter connections, functional regions, and anatomy
across development. Our framework thus opens opportunities not only for evaluating development
of white matter associated with functional regions in large infant datasets[82−86] but also for early
identification and assessment of developmental disorders associated with VTC such as autism[110−
113], Williams syndrome[114], congenital prosopagnosia[115,116], and dyslexia[117,**110−113**].

Together, the present study advances our understanding of white matter organization in the visual
system suggesting that cytoarchitecture, eccentricity, and category are organizing principles from
birth, even as aspects of the white matter develop from infancy to adulthood. These data have
implications not only for theories of cortical functional development, but also have ramifications
for early identification of atypical white matter development.

Methods

Participants

The study was approved by the Institutional Review Board of Stanford University and complies with all ethical regulations. Adult participants and parents of infant participants provided written informed consent prior to their scan session. Both infant and adult participants were paid $25/hour for participation.

Expectant parent and infant screening procedure

Expectant parents and their infants in our study were recruited from the San Francisco Bay Area using social media platforms. We performed a two-step screening process. First, parents were screened over the phone for eligibility based on exclusionary criteria designed to recruit a sample of typically developing infants. Second, eligible expectant mothers were screened once again after giving birth. Exclusionary criteria were as follows: recreational drug use during pregnancy, significant alcohol use during pregnancy (> 3 instances of alcohol consumption per trimester; more than 1 drink/instance), taking prescription medications for a disorder involving psychosis or mania during pregnancy, insufficient written and spoken English ability to understand study instructions, or learning disabilities. Exclusionary criteria for infants were preterm birth (<37 gestational weeks), low birthweight (<5 lbs 8 oz), congenital, genetic, and neurological disorders, visual problems, complications during birth that involved the infant (e.g., NICU stay), history of head trauma, and contraindications for MRI (e.g., metal implants).

43 full term infants following a typical pregnancy (gestational weeks: 39.31±1.32, mean±sd) participated in a total of 75 MRI sessions. Two sessions were excluded for missing diffusion data, and six sessions were excluded for too much motion (see Quality Assurance). We report data from 42 infants over 67 sessions (21 longitudinal): sex: 16 female, 26 male; race and ethnicity: 4 Asian, 5 Hispanic, 11 Multiracial, and 22 White participants; age: n=23 newborns (28.56 ± 10.21days, mean±sd), n=23 3-month-olds (106.91 ± 19.33 days), and n=21 6-month-olds (189.05 ±15.77 days).

We also collected data from 21 adults: sex: 17 female, 4 male; race and ethnicity: 12 Asian, 1 Hispanic, 2 Multiracial, and 6 White participants; age: M=28.21 years, SD=5.51 years. All adult sessions met inclusion criteria. No statistical methods were used to pre-determine sample sizes, but our sample sizes are similar to previous publications[45].

MRI acquisitions

Scanning sessions were scheduled in the evenings around infants' bedtime and were done during
natural sleep. Infant data were acquired on a 3T GE Ultra High Performance (UHP) scanner (GE
Healthcare, Waukesha, WI) equipped with a customized 32-channel infant head-coil[118].

Hearing protection included soft wax earplugs, and MRI compatible neonatal noise attenuators
(https://newborncare.natus.com/products-services/newborn-care-products/nursery-
essentials/minimuffs-neonatal-noise-attenuators), and headphones
(https://www.alpinehearingprotection.com/products/muffy-baby) that covered the infant's ears.
During sessions with newborns, an MR-safe plastic immobilizer (MedVac, www.supertechx-
ray.com) was used to stabilize the infant and their head position. When the infant was asleep, the
caregiver placed the infant on the scanner bed. Weighted bags were placed at the edges of the bed
to prevent side-to-side movements. Pads were also placed around the infant's head and body to
stabilize head position. An experimenter stayed inside the MR suite with the infant during the entire
scan.
562
563   During scan, experimenters monitored the infant using an infrared camera that was affixed to the
564   head coil and positioned for viewing the infant's face. Experimenters stopped the scan if the infant
565   showed signs of waking or distress or excessively moved; scans were repeated if there was
566   excessive head motion.
567
568
569   *Anatomical MRI acquisition*
570
We obtained T1-weighted MRI data (GE's BRAVO sequence) for each infant and adult session with
the following parameters: TE = 2.9ms; TR = 6.9ms; voxel size = 0.8x 0.8 x0.8 [mm^3]; FOV = 20.5 cm;
Scan time: 3:05 min. We obtained T2-weighted MRI data for each infant session as the T2-weighted
contrast between different tissue types is better than T1-weighted images for young infants. T2-
weighted (GE's CUBE sequence) parameters: TE = 124ms; TR = 3650 ms; voxel size = 0.8x 0.8x 0.8
[mm^3]; FOV = 20.5 cm; Scan time: 4:05 min.

Anatomical MRI processing for infant sessions

The T1-weighted and T2-weighted images from each individual were aligned to a plane running
through the commissures (AC-PC transformed)using rigid body alignment (FSL FLIRT[119];
https://web.mit.edu/fsl_v5.0.10/fsl/doc/wiki/FLIRT(2f)UserGuide.html). Then, we used iBEAT
V2.0[120] to segment the white and gray matter. White matter segmentation was then manually fixed
for errors using ITKgray[121]. We used the manually edited segmentation file to reconstruct the
cortical surface with Infant FreeSurfer[122].

Anatomical MRI processing for adult sessions

The T1-weighted images were aligned to a plane running through the commissures (AC-PC
transformed). Tissue segmentation was done with Freesurfer v7.2[123] .

Diffusion MRI (dMRI) acquisition

dMRI used the following parameters: multi-shell, #diffusion directions/b-value = 9/0, 30/700,
64/2000; TE = 75.7 ms; TR = 2800 ms; voxel size = 2x2x2 [mm^3]; number of slices = 60;
FOV = 20 cm; in-plane/through-plane acceleration = 1/3; scan time: 5:08 min. We also acquired a
short dMRI scan with reverse phase encoding direction and only 6 b = 0 images.

Diffusion MRI processing

dMRI data was preprocessed using MRtrix3[124] (https://github.com/MRtrix3/mrtrix3) in accordance
with prior work from the human connectome project and our lab [45,125,126]. Data were denoised using
principal component analysis[127]. We used FSL's top-up tool (https://fsl.fmrib.ox.ac.uk/) and one
image with reverse phase-encoding to correct for susceptibility-induced distortions. We used FSL's
eddy tool to perform eddy current and motion correction, where outlier slices were detected and
replaced[128].   Finally   we   performed   bias   correction   using   ANTS[129]
(https://picsl.upenn.edu/software/ants/). The preprocessed dMRI data were aligned to the T2-
weighted anatomy for infants, and T1-weighted anatomy for adults using whole-brain rigid body
registration. Alignment was checked manually.

Generating White Matter Connectomes

We generated a whole brain white matter connectome in each session using MRTrix3[124]. Voxel-
wise fiber orientation distributions (FODs) were calculated using constrained spherical
deconvolution (CSD). We used the Dhollander algorithm[130] to estimate the three-tissue response
function. We computed FODs separately for the white matter and CSF. As in past work[45,125], the
gray matter was not modeled separately, as white and gray matter do not have sufficiently distinct
b-value dependencies to allow for a clean separation of the signals. Finally, we generated a whole
brain white matter connectome for each session using MRTrix3 tckgen. We used a FOD amplitude
threshold of 0.05. Tractography was optimized using the gray/white matter segmentation from
anatomical MRI data (Anatomically Constrained Tractography; ACT[131]). For each connectome, we
used probabilistic fiber tracking with the following parameters: iFOD2 algorithm, step size of 0.2
mm, minimum length of 4 mm, maximum length of 200 mm, and maximum angle of 15°.
Streamlines were randomly seeded on this gray-matter/white-matter interface, and each
connectome consisted of 5 million streamlines. This procedure is critical for accurately identifying
the connections that reach fROIs, which are in the gray matter[132].

Quality Assurance

To evaluate the quality of dMRI data we implemented three quality assurance measures: (i) We
measured the number of outliers (dMRI volumes with signal dropout measured by FSL's eddy tool).
The exclusion criterion was >5% outlier volumes as in past work[20]. (ii) We visualized in mrView the
fractional anisotropy (FA) colored by direction of the diffusion tensors to validate the expected
maps. That is, the existence of between hemispheric connections through the corpus callosum,
and the inferior-superior directionality of the cerebral spinal tract. (iii) We used babyAFQ[22] to identify
the bundles of the brain and validate that the major bundles known to be present at birth[45,46,133] can
be found in each session.

These quality assurance measures lead to the following data exclusions:

(i) No adult sessions were excluded for outliers. Six infant sessions were excluded because they
had >5% outliers. After exclusion, there were no significant differences in outlier volumes between
infants and adults (infants: mean percent outliers±SD = 0.45% ± 0.43%; adults: mean percent
outliers±SD = 0.29% ± 0.10%; $t(86) = 1.70$, $p = .092$).

(ii) We identified the corpus callosum with left-right directionality and corticospinal tract with
inferior-superior directionality in all infant and adult sessions; no data was excluded by this
measure.

(iii) We identified the major bundles in the infant brain using babyAFQ, and in adults using AFQ. In
each infant and adult session, we identified the arcuate (AF), posterior arcuate (pAF), vertical
occipital (VOF), inferior longitudinal (ILF), superior longitudinal (SLF), corticospinal (CST), cingulate
(CC), forceps major (FcMa), and forceps minor (FcMi), as such, no data were excluded. This
confirms the quality of dMRI data and our tracking procedure as we could identify the major white
matter bundles that are known to exist from birth in all sessions.

Cytoarchitectonic Areas of VTC

We used the Rosenke Atlas[59] that contains maximum probability maps (MPMs) of 8
cytoarchitectonic regions of the human ventral visual stream, created from 10 post mortem adult
samples published in[37,134]. We used cortex based alignment in FreeSurfer[56] (https://freesurfer.net/,
mri_label2label) to map the MPM of four cytoarchitectonic areas of the ventral stream FG1, FG2,
FG3, FG4 to each cortical surface for each of the 88 sessions in the study. Two independent raters
(SD, DO) manually checked each brain to examine if the mid fusiform sulcus (MFS) was the
boundary between FG3/FG4 and FG1/FG2 as reported in prior studies[5,58]. In places where there
were disagreements between the raters, data was checked by KGS. The analysis confirmed that
the boundaries between the cytoarchitectonic areas aligned with the MFS (see examples in Fig 1c)
except for five infants in the right hemisphere, and one adult in the left hemisphere.

Category-Selective Regions of VTC

A functional atlas of category-selective regions of VTC was created from independent fMRI data in
28 adults (11 females and 17 males, ages 22.1–28.6 years, mean±SD = 24.1±1.6 years)from a
previous study[20]. Category-selective regions were defined in each of the individuals using the fLoc
experiment containing low-level and familiarity-controlled gray level images of items from 10 visual
categories[135]. Category-selective regions were defined using a voxel-level t-statistic contrasting
each category of interest with all other categories (t > 3, no spatial smoothing), and anatomical
criteria[5,6,135]. We defined two face-selective regions on the fusiform gyrus of each hemisphere,
mFus-faces and pFus-faces (contrast: faces > bodies, limbs, characters, objects, places). mFus-
faces is located near the anterior tip of the mid fusiform sulcus (MFS), pFus-faces is in the posterior
fusiform. We defined OTS-bodies in both hemispheres as the region on the occipital temporal
sulcus selective for bodies (bodies and limbs > characters, faces, objects, places). We defined CoS-
places in both hemispheres as the region in the intersection between the collateral sulcus (CoS)
and anterior lingual sulcus (ALS) that was selective for places (places > faces, characters, objects,
bodies, limbs). Finally, we identified mOTS-words and pOTS-words as the regions in the occipital
temporal sulcus (OTS) that were selective for words (psuedowords > faces, scenes, objects, bodies,
limbs), where pOTS-words in on the posterior end of the OTS lateral to pFus-faces, and mOTS-
words is more anterior and lateral to mFus-faces. After identifying the functional regions within
each participant, we mapped the regions to the fsaverage template brain space using cortex based
alignment[56]. We then created probabilistic maps for each fROI where each vertex was the
probability of a participant having the fROI at that location. We thresholded the probabilistic maps
at .2 and then created MPMs, where in the case that two fROIs had probabilistic values > .2 at the
same vertex, the vertex was assigned to the fROI with the highest probability (as in[136]). MPM-fROIs
of word-selective regions were only found in the left hemisphere[137,138].

To map fROIs to individual participants' cortical space for the 88 sessions of the main experiment,
we used cortex based alignment in Freesurfer[56] (https://freesurfer.net/, mri_label2label). Two
independent raters (DO and SD) then visually checked each surface to ensure that the fROIs were
aligned to the expected anatomical landmarks in each individual participant. The raters checked
whether mFus-faces aligned to the mid fusiform sulcus (MFS) and whether CoS-places aligned to
the junction of the ALS and CoS. When there was disagreement, KGS then checked whether the
fROIs aligned to anatomical landmarks. Category-selective fROIs aligned to these anatomical
landmarks in all hemispheres except for one infant in which left hemisphere CoS-places did not
align to the junction or the ALS and CoS, and two infants in which right hemisphere mFus-faces did
not align to the MFS. Supplementary Figs. 2-9 show the mapping of each fROI in each participant.

We tested whether fROI surface area relative to brain surface varied with age as follows:

(1) (fROI surface area)/(brain surface area) ~ log10(age in days).

We found no significant differences between infants and adults on fROI surface area relative to
brain surface area (effect of age: t(878) = 8.77x10^{-6}, p = 0.88)), reflecting that fROIs were the same
size relative to the size of the brain regardless of age.

Identifying functionally defined white matter connections

To identify the white matter connections of each fROI, we intersected it with each individual's whole
brain connectome using an open source software package, FSuB-Extractor[109]
(https://github.com/smeisler/fsub_extractor). The software takes in fROIs in the native space of
each participant and projects them along the surface normal into the gray-matter-white-matter
interface. It then restricts the fROI to the gray-matter-white-matter interface and then selects all
streamlines that intersect with the fROI using a radial search with a search distance of 3mm.

Defining connectivity profiles

For each fROI, we define its connectivity profile. To do so, we took the white matter connections of
each fROI, projected their endpoints to the cortical surface using tract density imaging (TDI) with
MRTrix3[124], and calculated the distribution of these white matter endpoints across the cortical
surface. We transformed the TDI output into a distribution of endpoints by dividing the endpoint
map by the total number of endpoints. This results in an endpoint density map that sums to 1 for
each fROI and participant. We then used the Glasser Atlas[60] to quantify how the endpoints were
distributed across the brain. We chose to use the Glasser Atlas because it covers the whole cortical
surface of each hemisphere, and because it divides cortex into meaningful parcellations according
to functional and anatomical criteria. For each fROI, we define the white matter connectivity profile
or the endpoint density in each region in the Glasser Atlas. The Glasser Atlas consists of 180
regions per hemisphere. As in prior work[20], we excluded 11 VTC regions of the Glasser Atlas to
avoid quantifying looping fibers from the seed fROI. Therefore, each connectivity profile consisted
of the endpoint density across 169 regions in the Glasser Atlas within the same hemisphere of the
fROI.

Principal Components Analysis

Because our connectivity profiles are high-dimensional (169), we used principal component
analysis (PCA function in MATLAB) to reduce the dimensionality of the connectivity profiles. We
conducted PCA on the 880 (10 fROIs x 88 sessions connectivity profiles) x 169 (Glasser ROIs)
endpoint connectivity matrix. We used find_curve_elbow in R
(https://rdrr.io/cran/pathviewr/man/find_curve_elbow.html) to find the elbow of the curve of
principal components vs. variance explained. We found that the elbow of the curve was at 10
principal components, and that the first 10 principal components explained 98% of the variance in
the data. In Figure 2, we plot the first principal component (explains 38% of the variance) vs the
second principal component (explains 20% of the variance). Each dot represents a single

connectivity profile in a single subject and session; The connectivity profiles (dots) can be coded
by different features (cytoarchitectonic area, category-preference, or participant's age).

Classification Analysis

We used a n-way leave-one-out classifier to test if we could predict different features
(cytoarchitecture, category, or age) of a held-out connectivity profile. For each feature
(cytoarchitecture, category, age), we used multinomial logistic regression (multinom function from
the nnet package in R[139]) fit on all data, excluding all sessions from the held-out subject to account
for the longitudinal nature of the data to predict the feature of the held out connectivity profile (e.g.,
probability of belonging to FG2, FG3, or FG4 for the cytoarchitecture classification). We used a
winner-take-all approach and assigned the connectivity profile to the label with the highest
probability. We repeated the process for each connectivity profile (leave-one-out cross-validation).
We then calculated classification accuracy by comparing the predicted classification to the ground
truth. We performed three separate classifications: one predicting cytoarchitecture
(FG2/FG3/FG4), one predicting category (face/word/body/place), and one predicting age group:
(newborn/3 months/6 months/adult).

We compared classification accuracy across classification tasks by using a binomial logistic
regression (glm in R
(https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/glm)) with the following
model:

(1) classification accuracy(1/0) ~ classification task (cytoarchitecture/category/age)

As our results could be affected by training sets with different numbers of examples for each label,
we used a bootstrapping procedure to create a balanced training set. To do so, we sampled the
training set (all data except for the held-out subject) with replacement 200 times for each label (e.g.,
FG2/FG3/FG4) to create a balanced training set. We then fit a multinomial logistic regression on
the balanced training set to predict the feature of the held-out connectivity profile. We then
calculated classification accuracy by comparing the predicted classification to the ground truth.
We repeated this procedure for the cytoarchitecture, category, and age classification.

Connections to Eccentricity Bands in Early Visual Cortex

To test how each fROI connects to different eccentricity bands in early visual cortex (EVC), we
created a region of interest corresponding to EVC from the union of V1, V2, and V3 ROIs from the
Benson Atlas[140] (https://github.com/noahbenson/neuropythy). We then used the average
retinotopic map in 21 adults from a previously published paper[141] to divide EVC into three
eccentricity bands, 0-5°, 5-10°, and 10-20°. Using cortex-based alignment in FreeSurfer

(mri_label2label) we aligned each of these EVC eccentricity band ROIs to each individual
participant's native space. We then used FSuB-Extractor [109]
(https://github.com/smeisler/fsub_extractor), to identify the white matter connections between
each fROI and each eccentricity band in VTC. We divided the number of streamlines connecting to
each eccentricity band by the total number of streamlines between the fROI and EVC to estimate
the percentage of streamlines connected to each eccentricity band. To test if the percentage of
streamlines to each eccentricity band differs by cytoarchitecture, category, and age, we fit the
following model using the lm function in R
(https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/lm):

(2) percentage of streamlines in central 5° ~ cytoarchitecture x category x age group

After finding a significant cytoarchitecture by category interaction, we performed post-hoc tests to
determine which fROIs were driving the interaction. To do so, we used paired t-tests to test whether
the percentage of streamlines in the central 5° differed for fROIs within the same cytoarchitectonic
area for the following pairs of fROIs: pFus-faces and pOTS-words in FG2, mFus-faces and OTS-
bodies in FG4, mFus-faces and mOTS-words in FG4, and OTS-bodies and mOTS-words in FG4.

After finding a significant cytoarchitecture by age interaction, we performed post-hoc tests to test
whether the percentage of streamlines in the central 5° differed as a function of age for fROIs within
each cytoarchitectonic area using the following linear mixed model (using lmer function from the
lmerTest package[142] in R) separately for each cytoarchitectonic area (FG2, FG3, FG4), using a
random intercept for each subject to account for the longitudinal nature of the data:

(3) percentage of streamlines in central 5° ~ log10(age) + (1|subject)

After finding a significant interaction between cytoarchitecture, category, and age we performed
post-hoc tests to test whether the percentage of streamlines to the central 5° changed over
development using the following linear mixed model (using lmer function from the lmerTest
package[142] in R) separately for each fROI (pFus-faces, pOTS-words, CoS-places, mFus-faces, OTS-
bodies, mOTS-words), using a random intercept for each subject to account for the longitudinal
nature of the data:

(4) percentage of streamlines in central 5° ~ log10(age) + (1|subject)

Quantifying development

As development is expected to asymptote across the lifespan and typically follows a logarithmic
function of age, to quantify how white matter connectivity profiles change with age, we fit a linear
model (lm in R (https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/lm))

predicting endpoint density of each fROI as a function of Glasser ROI and log10(age in days) across
the brain. Age is a continuous variable and Glasser ROI is a categorical variable. As endpoint density
was normalized to sum to one across the brain, it is not appropriate to model a random intercept
for each subject.

(5) fROI endpoint density ~ log10(age in days) x Glasser ROI

After finding a significant Glasser ROI by age interaction (Supplementary Table 1), for each fROI,
we fit linear models (lm in R
https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/lm) relating endpoint
density vs log10(age) separately for each Glasser ROI to quantify the development of endpoint
density within each Glasser ROI. The regression slope quantifies the rate of the development:

(6) fROI endpoint density ~ log10(age in days).

We visualize the regression results in Fig 4B, where each Glasser ROI is colored by the slope of the
regression. Data is reported for the left hemisphere in the main text and Fig 3 because all 6 fROIs
are found consistently in the left hemisphere. Data from 4 fROIs in the right hemisphere is reported
in Supplementary Figure 55. Supplementary Tables 3-8 provide all the developmental slopes per
Glasser ROI.

Calculating the similarity between developmental slopes

To test whether development was more similar for regions within the same cytoarchitectonic area
or category-selectivity we estimated the similarity between developmental slopes of white matter
connectivity of fROI pairs using a bootstrapping procedure.

First, we calculated the developmental slopes of endpoint connectivity across the brain using a
bootstrapping procedure. For each fROI, and for each of 1000 iterations we used 75% of the data
(66 randomly selected connectivity profiles), and calculated the developmental slope within each
of the 169 Glasser ROIs using the following model (lm in R
https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/lm):

(7) fROI endpoint density ~ log10(age in days)

Then for each pair of fROIs we calculated the pairwise correlation between the vector of 169 slopes
across the bootstrap iterations. Finally, we fit a linear model (lm in R
https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/lm)) to test whether the
Fisher transformed correlation between development slopes was higher for fROIs with the same
cytoarchitecture and category-selectivity than for fROIs with different cytoarchitecture and

category-selectivity. We calculated the Fisher transform to ensure normality (as correlations are
bounded between -1 and 1); category and cytoarchitecture are categorical variables (same: 1;
different: 0).

(8) FisherZ(correlation) ~ category + cytoarchitecture

Data availability

823 The data to make the figures, tables, and statistics associated with this manuscript is available
824 here: https://github.com/VPNL/bbVTCwm/tree/main/data

Code availability

The code to analyze the data, compute statistics, and make the individual figure elements is
available here: https://github.com/VPNL/bbVTCwm/. The code folder contains the R code used to
generate all other figures and statistics in the figures/ and statistics/ subdirectories. The code used
to preprocess the data and perform the analyses are included in the analyses/ subdirectory. The
label files for the fROIs and the EVC ROIs are provided in the labels folder. The supplement folder
contains code to generate Supplementary Figures.

Acknowledgements

This work was funded by Stanford Wu Tsai Neurodevelopment big idea and accelerator grants, as
well as NIH grants R01EY033835 and R01EY022318 to KGS; the National Science Foundation
Graduate Research Fellowship (grant number DGE-1656518) to EK; the Deutsche
Forschungsgemeinschaft (DFG, German Research Foundation – project number 222641018 –
SFB/TRR 135 TP C10), as well as "The Adaptive Mind", funded by the Excellence Program of the
Hessian Ministry of Higher Education, Science, Research and Art to MG; the Deutsche
Forschungsgemeinschaft (DFG, German Research Foundation – grant INST 169/22–1), the
Excellence Program of the Hessian Ministry of Higher Education, Science, Research and Art
(grants: 2/16/519/03/09.001(0001)/101 and LOEWE/4TP//519/05/02.002(0004)/107) to BK. **The
funders had no role in study design, data collection and analysis, decision to publish or
preparation of the manuscript.**

Author Contributions

EK: designed the analyses, wrote the code and data analysis pipelines, analyzed the data, and wrote
the manuscript. XY: participated in the design and data analysis and collected the data. ST, BF, CT

collected the data, segmented each brain anatomy image into gray and white matter, and created
cortical surface reconstructions. SF and DO: validated the alignment between fROIs,
cytoarchitectonic areas and anatomical landmarks. MG: participated in the data analyses. VSN:
participated in the design and data analysis and collected the data. BK: designed the infant coil
used for data collection. KGS: oversaw all parts of the research: design, data analysis, and wrote
the manuscript. All authors read and gave feedback on the manuscript.

Competing Interests

The authors declare no competing interests.

References

1. Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human

extrastriate cortex specialized for face perception. *J. Neurosci.* **17**, 4302–4311 (1997).

2. Peelen, M. V. & Downing, P. E. Selectivity for the human body in the fusiform gyrus. *J.*

Neurophysiol. **93**, 603–608 (2005).

3. Cohen, L. *et al.* Language-specific tuning of visual cortex? Functional properties of the Visual

Word Form Area. *Brain* **125**, 1054–1069 (2002).

4. Epstein, R., Harris, A., Stanley, D. & Kanwisher, N. The parahippocampal place area:

recognition, navigation, or encoding? *Neuron* **23**, 115–125 (1999).

5. Weiner, K. S. *et al.* The mid-fusiform sulcus: a landmark identifying both cytoarchitectonic

and functional divisions of human ventral temporal cortex. *Neuroimage* **84**, 453–465 (2014).

6. Weiner, K. S. *et al.* Defining the most probable location of the parahippocampal place area

using cortex-based alignment and cross-validation. *Neuroimage* **170**, 373–384 (2018).

7. Levy, I., Hasson, U., Avidan, G., Hendler, T. & Malach, R. Center–periphery organization of

human object areas. *Nat. Neurosci.* **4**, 533–539 (2001).

8. Hasson, U., Levy, I., Behrmann, M., Hendler, T. & Malach, R. Eccentricity Bias as an Organizing

Principle for Human High-Order Object Areas. *Neuron* **34**, 479–490 (2002).

9. Natu, V. S. *et al.* Infants' cortex undergoes microstructural growth coupled with myelination

during development. *Commun Biol* **4**, 1191 (2021).

10. Cachia, A. *et al.* How interindividual differences in brain anatomy shape reading accuracy.

Brain Struct. Funct. **223**, 701–712 (2018).

11. Saygin, Z. M. *et al.* Anatomical connectivity patterns predict face selectivity in the fusiform

gyrus. *Nat. Neurosci.* **15**, 321–327 (2011).

12. Osher, D. E. *et al.* Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple

Visual Categories across Cortex. *Cereb. Cortex* **26 4**, 1668–1683 (2016).

13. Saygin, Z. M. *et al.* Connectivity precedes function in the development of the visual word form

area. *Nat. Neurosci.* **19**, 1250–1255 (2016).

14. Bi, Y., Wang, X. & Caramazza, A. Object Domain and Modality in the Ventral Visual Pathway.

Trends Cogn. Sci. **20**, 282–290 (2016).

15. Mahon, B. Z. & Caramazza, A. What drives the organization of object knowledge in the brain?

Trends Cogn. Sci. **15**, 97–103 (2011).

16. Kanwisher, N. Functional specificity in the human brain: a window into the functional

architecture of the mind. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 11163–11170 (2010).

17. Weiner, K. S., Yeatman, J. D. & Wandell, B. A. The posterior arcuate fasciculus and the vertical

occipital fasciculus. *Cortex* **97**, 274–276 (2017).

18. Caffarra, S., Karipidis, I. I., Yablonski, M. & Yeatman, J. D. Anatomy and physiology of word-

selective visual cortex: from visual features to lexical processing. *Brain Struct. Funct.* **226**,

3051–3065 (2021).

19. Lerma-Usabiaga, G., Carreiras, M. & Paz-Alonso, P. M. Converging evidence for functional and

structural segregation within the left ventral occipitotemporal cortex in reading. *Proceedings*

of the National Academy of Sciences **115**, E9981–E9990 (2018).

20. Kubota, E. *et al.* White matter connections of high-level visual areas predict cytoarchitecture

better than category-selectivity in childhood, but not adulthood. *Cereb. Cortex* **33**, 2485–2506

(2023).

21. Finzi, D. *et al.* Differential spatial computations in ventral and lateral face-selective regions

are scaffolded by structural connections. *Nat. Commun.* **12**, (2021).

22. Grotheer, M., Yeatman, J. & Grill-Spector, K. White matter fascicles and cortical

microstructure predict reading-related responses in human ventral temporal cortex.

Neuroimage **227**, 117669 (2021).

23. Hannagan, T., Amedi, A., Cohen, L., Dehaene-Lambertz, G. & Dehaene, S. Origins of the

specialization for letters and numbers in ventral occipitotemporal cortex. *Trends Cogn. Sci.*

19, 374–382 (2015).

24. Op de Beeck, H. P., Pillet, I. & Ritchie, J. B. Factors Determining Where Category-Selective

Areas Emerge in Visual Cortex. *Trends Cogn. Sci.* **23**, 784–797 (2019).

25. Arcaro, M. J., Schade, P. F., Vincent, J. L., Ponce, C. R. & Livingstone, M. S. Seeing faces is

necessary for face-domain formation. *Nat. Neurosci.* **20**, 1404–1412 (2017).

26. Sugita, Y. Face perception in monkeys reared with no exposure to faces. *Proc. Natl. Acad. Sci.*

U. S. A. **105**, 394–398 (2008).

27. Yan, X. *et al.* When do visual category representations emerge in infants' brains? (2024)

doi:10.7554/elife.100260.

28. van den Hurk, J., Van Baelen, M. & Op de Beeck, H. P. Development of visual category

selectivity in ventral visual cortex does not require visual experience. *Proc. Natl. Acad. Sci. U.*

S. A. **114**, E4501–E4510 (2017).

29. Ratan Murty, N. A. *et al.* Visual experience is not necessary for the development of face-

selectivity in the lateral fusiform gyrus. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 23011–23020

(2020).

30. Osher, D. E. *et al.* Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple

Visual Categories across Cortex. *Cereb. Cortex* **26**, 1668–1683 (2016).

31.   Li, J., Osher, D. E., Hansen, H. A. & Saygin, Z. M. Innate connectivity patterns drive the

development of the visual word form area. *Sci. Rep.* **10**, 18039 (2020).

32.   Kamps, F. S., Hendrix, C. L., Brennan, P. A. & Dilks, D. D. Connectivity at the origins of domain

specificity in the cortical face and place networks. *Proceedings of the National Academy of*

Sciences **117**, 6163–6169 (2020).

33.   Zeki, S. & Shipp, S. The functional logic of cortical connections. *Nature* **335**, 311–317 (1988).

34.   Van Essen, D. C., Anderson, C. H. & Felleman, D. J. Information processing in the primate

visual system: an integrated systems perspective. *Science* **255**, 419–423 (1992).

35.   Amunts, K. & Zilles, K. Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron*

88, 1086–1107 (2015).

36.   Caspers, J. *et al.* Cytoarchitectonical analysis and probabilistic mapping of two extrastriate

areas of the human posterior fusiform gyrus. *Brain Struct. Funct.* **218**, 511–526 (2012).

37.   Lorenz, S. *et al.* Two New Cytoarchitectonic Areas on the Human Mid-Fusiform Gyrus. *Cereb.*

Cortex **27**, 373–385 (2017).

38.   Gomez, J., Natu, V. S., Jeska, B., Barnett, M. & Grill-Spector, K. Development differentially

sculpts receptive fields across early and high-level human visual cortex. *Nat. Commun.* **9**,

(2018).

39.   Ellis, C. T. *et al.* Retinotopic organization of visual cortex in human infants. *Neuron* **109**,

2616–2626.e6 (2021).

40.   Butt, O. H., Benson, N. C., Datta, R. & Aguirre, G. K. The fine-scale functional correlation of

striate cortex in sighted and blind people. *J. Neurosci.* **33**, 16209–16219 (2013).

41.   Butt, O. H., Benson, N. C., Datta, R. & Aguirre, G. K. Hierarchical and homotopic correlations of

spontaneous neural activity within the visual cortex of the sighted and blind. *Front. Hum.*

Neurosci. **9**, 25 (2015).

42. Arcaro, M. J. & Livingstone, M. S. A hierarchical, retinotopic proto-organization of the primate

visual system at birth. *Elife* **6**, (2017).

43. Arcaro, M. J. & Livingstone, M. S. On the relationship between maps and domains in

inferotemporal cortex. *Nat. Rev. Neurosci.* **22**, 573–583 (2021).

44. Dudink, J., Kerr, J. L., Paterson, K. & Counsell, S. J. Connecting the developing preterm brain.

Early Hum. Dev. **84**, 777–782 (2008).

45. Grotheer, M. *et al.* White matter myelination during early infancy is linked to spatial gradients

and myelin content at birth. *Nat. Commun.* **13**, 997 (2022).

46. Grotheer, M. *et al.* Human white matter myelinates faster in utero than ex utero. *Proc. Natl.*

Acad. Sci. U. S. A. **120**, e2303491120 (2023).

47. Dubois, J., Hertz-Pannier, L., Dehaene-Lambertz, G., Cointepas, Y. & Le Bihan, D. Assessment

of the early organization and maturation of infants' cerebral white matter fiber bundles: A

feasibility study using quantitative diffusion tensor imaging and tractography. *Neuroimage*

30, 1121–1132 (2006).

48. Zöllei, L., Jaimes, C., Saliba, E., Grant, P. E. & Yendiki, A. TRActs constrained by UnderLying

INfant anatomy (TRACULInA): An automated probabilistic tractography tool with anatomical

priors for use in the newborn brain. *Neuroimage* **199**, 1–17 (2019).

49. Dimond, D. *et al.* Early childhood development of white matter fiber density and morphology.

Neuroimage **210**, 116552 (2020).

50. Perani, D. *et al.* Neural language networks at birth. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 16056–

16061 (2011).

51. Levin, N., Dumoulin, S. O., Winawer, J., Dougherty, R. F. & Wandell, B. A. Cortical maps and

white matter tracts following long period of visual deprivation and retinal image restoration.

Neuron **65**, 21–31 (2010).

52. Bauer, C. M. *et al.* Abnormal white matter tractography of visual pathways detected by high-

angular-resolution diffusion imaging (HARDI) corresponds to visual dysfunction in

cortical/cerebral visual impairment. *J. AAPOS* **18**, 398–401 (2014).

53.   Ortibus, E. *et al.* Integrity of the inferior longitudinal fasciculus and impaired object

recognition in children: a diffusion tensor imaging study. *Dev. Med. Child Neurol.* **54**, 38–43

(2012).

54.   Lennartsson, F., Nilsson, M., Flodmark, O. & Jacobson, L. Damage to the immature optic

radiation causes severe reduction of the retinal nerve fiber layer, resulting in predictable

visual field defects. *Invest. Ophthalmol. Vis. Sci.* **55**, 8278–8288 (2014).

55.   Zufferey, P. D., Jin, F., Nakamura, H., Tettoni, L. & Innocenti, G. M. The role of pattern vision in

the development of cortico-cortical connections. *Eur. J. Neurosci.* **11**, 2669–2688 (1999).

56.   Fischl, B., Sereno, M. I. & Dale, A. M. Cortical surface-based analysis. II: Inflation, flattening,

and a surface-based coordinate system. *Neuroimage* **9**, 195–207 (1999).

57.   Weiner, K. S. *et al.* The mid-fusiform sulcus: A landmark identifying both cytoarchitectonic

and functional divisions of human ventral temporal cortex. *Neuroimage* **84**, 453–465 (2014).

58.   Weiner, K. S. *et al.* The Cytoarchitecture of Domain-specific Regions in Human High-level

Visual Cortex. *Cereb. Cortex* **27**, 146–161 (2016).

59.   Rosenke, M. *et al.* A cross-validated cytoarchitectonic atlas of the human ventral visual

stream. *Neuroimage* **170**, 257–270 (2018).

60.   Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171–

178 (2016).

61.   Beyh, A. *et al.* The medial occipital longitudinal tract supports early stage encoding of

visuospatial information. *Commun. Biol.* **5**, 318 (2022).

62.   Grill-Spector, K. & Weiner, K. S. The functional architecture of the ventral temporal cortex and

its role in categorization. *Nat. Rev. Neurosci.* **15**, 536–548 (2014).

63.   Dubois, J. *et al.* MRI of the Neonatal Brain: A Review of Methodological Challenges and

Neuroscientific Advances. *J. Magn. Reson. Imaging* **53**, 1318–1343 (2021).

64. Katz, L. C. & Shatz, C. J. Synaptic activity and the construction of cortical circuits. *Science*

274, 1133–1138 (1996).

65. Caspers, S. *et al.* The human inferior parietal cortex: cytoarchitectonic parcellation and

interindividual variability. *Neuroimage* **33**, 430–448 (2006).

66. Richter, M. *et al.* Cytoarchitectonic segregation of human posterior intraparietal and adjacent

parieto-occipital sulcus and its relation to visuomotor and cognitive functions. *Cereb. Cortex*

29, 1305–1327 (2019).

67. Malikovic, A. *et al.* Cytoarchitecture of the human lateral occipital cortex: mapping of two

extrastriate areas hOc4la and hOc4lp. *Brain Struct. Funct.* **221**, 1877–1897 (2016).

68. Rottschy, C. *et al.* Ventral visual cortex in humans: Cytoarchitectonic mapping of two

extrastriate areas. *Hum. Brain Mapp.* **28**, (2007).

69. Groen, I. I. A., Dekker, T. M., Knapen, T. & Silson, E. H. Visuospatial coding as ubiquitous

scaffolding for human cognition. *Trends Cogn. Sci.* **26**, 81–96 (2022).

70. Gomez, J., Barnett, M. & Grill-Spector, K. Extensive childhood experience with Pokémon

suggests eccentricity drives organization of visual cortex. *Nat Hum Behav* **3**, 611–624 (2019).

71. Silson, E. H., Chan, A. W.-Y., Reynolds, R. C., Kravitz, D. J. & Baker, C. I. A Retinotopic Basis for

the Division of High-Level Scene Processing between Lateral and Ventral Human

Occipitotemporal Cortex. *J. Neurosci.* **35**, 11921–11935 (2015).

72. Somers, D. C. & Sheremata, S. L. Attention maps in the brain. *Wiley Interdiscip. Rev. Cogn. Sci.*

4, 327–340 (2013).

73. Silver, M. A. & Kastner, S. Topographic maps in human frontal and parietal cortex. *Trends*

Cogn. Sci. **13**, 488–495 (2009).

74. Amunts, K., Mohlberg, H., Bludau, S. & Zilles, K. Julich-Brain: A 3D probabilistic atlas of the

human brain's cytoarchitecture. *Science* **369**, 988–992 (2020).

75. Sylvester, C. M. *et al.* Network-specific selectivity of functional connections in the neonatal

brain. *Cereb. Cortex* **33**, 2200–2214 (2023).

76. Huang, H. *et al.* White and gray matter development in human fetal, newborn and pediatric

brains. *Neuroimage* **33**, 27–38 (2006).

77. Thiebaut de Schotten, M. & Forkel, S. J. The emergent properties of the connected brain.

Science **378**, 505–510 (2022).

78. Dubois, J. *et al.* The early development of brain white matter: a review of imaging studies in

fetuses, newborns and infants. *Neuroscience* **276**, 48–71 (2014).

79. Harding-Forrester, S. & Feldman, D. E. Somatosensory maps. *Handb. Clin. Neurol.* **151**, 73–

102 (2018).

80. Humphries, C., Liebenthal, E. & Binder, J. R. Tonotopic organization of human auditory cortex.

Neuroimage **50**, 1202–1211 (2010).

81. Amunts, K. *et al.* Broca's region revisited: cytoarchitecture and intersubject variability. *J.*

Comp. Neurol. **412**, 319–341 (1999).

82. Lebel, C., Treit, S. & Beaulieu, C. A review of diffusion MRI of typical white matter

development from early childhood to young adulthood. *NMR Biomed.* **32**, e3778 (2019).

83. Edwards, A. D. *et al.* The Developing Human Connectome Project Neonatal Data Release.

Front. Neurosci. **16**, 886772 (2022).

84. Howell, B. R. *et al.* The UNC/UMN Baby Connectome Project (BCP): An overview of the study

design and protocol development. *Neuroimage* **185**, 891–905 (2019).

85. Walker, L. *et al.* The diffusion tensor imaging (DTI) component of the NIH MRI study of

normal brain development (PedsDTI). *Neuroimage* **124**, 1125–1130 (2016).

86. Volkow, N. D., Gordon, J. A. & Freund, M. P. The Healthy Brain and Child Development Study—

Shedding Light on Opioid Exposure, COVID-19, and Health Disparities. *JAMA Psychiatry* **78**,

471–472 (2021).

87. Eichert, N. *et al.* What is special about the human arcuate fasciculus? Lateralization,

projections, and expansion. *Cortex* **118**, 107–115 (2019).

88. Brauer, J., Anwander, A., Perani, D. & Friederici, A. D. Dorsal and ventral pathways in language

development. *Brain Lang.* **127**, 289–295 (2013).

89. Innocenti, G. M. & Frost, D. O. Effects of visual experience on the maturation of the efferent

system to the corpus callosum. *Nature* **280**, 231–234 (1979).

90. LaMantia, A. S. & Rakic, P. Axon overproduction and elimination in the corpus callosum of the

developing rhesus monkey. *J. Neurosci.* **10**, 2156–2175 (1990).

91. Price, D. J., Ferrer, J. M., Blakemore, C. & Kato, N. Postnatal development and plasticity of

corticocortical projections from area 17 to area 18 in the cat's visual cortex. *J. Neurosci.* **14**,

2747–2762 (1994).

92. Dehay, C., Kennedy, H. & Bullier, J. Characterization of transient cortical projections from

auditory, somatosensory, and motor cortices to visual areas 17, 18, and 19 in the kitten. *J.*

Comp. Neurol. **272**, 68–89 (1988).

93. Innocenti, G. M. & Price, D. J. Exuberance in the development of cortical networks. *Nat. Rev.*

Neurosci. **6**, 955–965 (2005).

94. Webster, M. J., Bachevalier, J. & Ungerleider, L. G. Transient subcortical connections of

inferior temporal areas TE and TEO in infant macaque monkeys. *J. Comp. Neurol.* **352**, 213–

226 (1995).

95. Kaneko, T. *et al.* Spatial organization of occipital white matter tracts in the common

marmoset. *Brain Struct. Funct.* **225**, 1313–1326 (2020).

96. Takemura, H., Pestilli, F. & Weiner, K. S. Comparative neuroanatomy: Integrating classic and

modern methods to understand association fibers connecting dorsal and ventral visual

cortex. *Neurosci. Res.* **146**, 1–12 (2019).

97. Takemura, H. *et al.* Occipital White Matter Tracts in Human and Macaque. *Cereb. Cortex* **27**,

98. Takemura, H. *et al.* A prominent vertical occipital white matter fasciculus unique to primate brains. *Curr. Biol.* (2024) doi:10.1016/j.cub.2024.06.034.

99. Zatorre, R. J., Fields, R. D. & Johansen-Berg, H. Plasticity in gray and white: neuroimaging changes in brain structure during learning. *Nat. Neurosci.* **15**, 528–536 (2012).

100. Blumenfeld-Katzir, T., Pasternak, O., Dagan, M. & Assaf, Y. Diffusion MRI of structural brain plasticity induced by a learning and memory task. *PLoS One* **6**, e20678 (2011).

101. Fields, R. D. A new mechanism of nervous system plasticity: activity-dependent myelination. *Nat. Rev. Neurosci.* **16**, 756–767 (2015).

102. Fields, R. D. Neuroscience. Change in the brain's white matter. *Science* **330**, 768–769 (2010).

103. Florence, S. L., Taub, H. B. & Kaas, J. H. Large-scale sprouting of cortical connections after peripheral injury in adult macaque monkeys. *Science* **282**, 1117–1121 (1998).

104. Jain, N., Florence, S. L., Qi, H. X. & Kaas, J. H. Growth of new brainstem connections in adult monkeys with massive sensory loss. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 5546–5550 (2000).

105. Dancause, N. *et al.* Extensive cortical rewiring after brain injury. *J. Neurosci.* **25**, 10167–10179 (2005).

106. Monje, M. Myelin Plasticity and Nervous System Function. *Annu. Rev. Neurosci.* **41**, 61–76 (2018).

107. Bacmeister, C. M. *et al.* Motor learning drives dynamic patterns of intermittent myelination on learning-activated axons. *Nat. Neurosci.* **25**, 1300–1313 (2022).

108. Murty, N. A. R. *et al.* Visual experience is not necessary for the development of face-selectivity in the lateral fusiform gyrus. *Proceedings of the National Academy of Sciences* **117**, 23011–23020 (2020).

109. Meisler, S. L., Kubota, E., Grotheer, M. & Gabrieli, J. A practical guide for combining functional regions of interest and white matter bundles. (2024).

110. Ewbank, M. P. *et al.* Repetition Suppression in Ventral Visual Cortex Is Diminished as a

Function of Increasing Autistic Traits. *Cereb. Cortex* **25**, 3381–3393 (2015).

111. Dakin, S. & Frith, U. Vagaries of visual perception in autism. *Neuron* **48**, 497–507 (2005).

112. Conti, E. *et al.* Network over-connectivity differentiates autism spectrum disorder from other

developmental disorders in toddlers: A diffusion MRI study. *Hum. Brain Mapp.* **38**, 2333–2344

(2017).

113. Picci, G., Gotts, S. J. & Scherf, K. S. A theoretical rut: revisiting and critically evaluating the

generalized under/over-connectivity hypothesis of autism. *Dev. Sci.* **19**, 524–549 (2016).

114. Golarai, G. *et al.* The fusiform face area is enlarged in Williams syndrome. *J. Neurosci.* **30**,

6700–6712 (2010).

115. Behrmann, M. & Avidan, G. Congenital prosopagnosia: face-blind from birth. *Trends Cogn. Sci.*

9, 180–187 (2005).

116. Duchaine, B. C. & Nakayama, K. Developmental prosopagnosia: a window to content-specific

face processing. *Curr. Opin. Neurobiol.* **16**, 166–173 (2006).

117. Yeatman, J. D. & White, A. L. Reading: The Confluence of Vision and Language. *Annu Rev Vis*

Sci **7**, 487–517 (2021).

118. Ghotra, A. *et al.* A size-adaptive 32-channel array coil for awake infant neuroimaging at 3

Tesla MRI. *Magn. Reson. Med.* **86**, 1773–1785 (2021).

119. Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W. & Smith, S. M. FSL.

Neuroimage **62**, 782–790 (2012).

120. Wang, L. *et al.* iBEAT V2.0: a multisite-applicable, deep learning-based pipeline for infant

cerebral cortical surface reconstruction. *Nat. Protoc.* **18**, 1488–1509 (2023).

121. Yushkevich, P. A. *et al.* User-guided 3D active contour segmentation of anatomical

structures: significantly improved efficiency and reliability. *Neuroimage* **31**, 1116–1128

(2006).

122. Zöllei, L., Iglesias, J. E., Ou, Y., Grant, P. E. & Fischl, B. Infant FreeSurfer: An automated

segmentation and surface extraction pipeline for T1-weighted neuroimaging data of infants

0–2 years. *Neuroimage* **218**, 116946 (2020).

123. Fischl, B. FreeSurfer. *Neuroimage* **62**, 774–781 (2012).

124. Tournier, J.-D. *et al.* MRtrix3: A fast, flexible and open software framework for medical image

processing and visualisation. *Neuroimage* **202**, 116137 (2019).

125. Pietsch, M. *et al.* A framework for multi-component analysis of diffusion MRI data over the

neonatal period. *Neuroimage* **186**, 321–337 (2019).

126. Bastiani, M. *et al.* Automated processing pipeline for neonatal diffusion MRI in the developing

Human Connectome Project. *Neuroimage* **185**, 750–763 (2019).

127. Veraart, J. *et al.* Denoising of diffusion MRI using random matrix theory. *Neuroimage* **142**,

394–406 (2016).

128. Andersson, J. L. R., Graham, M. S., Zsoldos, E. & Sotiropoulos, S. N. Incorporating outlier

detection and replacement into a non-parametric framework for movement and distortion

correction of diffusion MR images. *Neuroimage* **141**, 556–572 (2016).

129. Tustison, N. J. *et al.* N4ITK: improved N3 bias correction. *IEEE Trans. Med. Imaging* **29**, 1310–

1320 (2010).

130. Dhollander, T., Raffelt, D. & Connelly, A. Unsupervised 3-tissue response function estimation

from single-shell or multi-shell diffusion MR data without a co-registered T1 image. in (09

2016).

131. Smith, R. E., Tournier, J.-D., Calamante, F. & Connelly, A. Anatomically-constrained

tractography: Improved diffusion MRI streamlines tractography through effective use of

anatomical information. *Neuroimage* **62**, 1924–1938 (2012).

132. Grotheer, M., Kubota, E. & Grill-Spector, K. Establishing the functional relevancy of white

matter connections in the visual system and beyond. *Brain Struct. Funct.* **227**, 1347–1356

(2022).

133. Warrington, S. *et al.* Concurrent mapping of brain ontogeny and phylogeny within a common

space: Standardized tractography and applications. *Science Advances* **8**, eabq2022 (2022).

134. Caspers, J. *et al.* Cytoarchitectonical analysis and probabilistic mapping of two extrastriate

areas of the human posterior fusiform gyrus. *Brain Struct. Funct.* **218**, 511–526 (2013).

135. Stigliani, A., Weiner, K. S. & Grill-Spector, K. Temporal Processing Capacity in High-Level

Visual Cortex Is Domain Specific. *J. Neurosci.* **35**, 12412–12424 (2015).

136. Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K. & Goebel, R. A Probabilistic

Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cereb. Cortex* **31**, 603–619

(2020).

137. Cai, Q., Paulignan, Y., Brysbaert, M., Ibarrola, D. & Nazir, T. A. The left ventral occipito-

temporal response to words depends on language lateralization but not on visual familiarity.

Cereb. Cortex **20**, 1153–1163 (2010).

138. Petersen, S. E., Fox, P. T., Posner, M. I., Mintun, M. A. & Raichle, M. E. Positron emission

tomographic studies of the cortical anatomy of single-word processing. *Nature* **331**, 585–

589 (1988).

139. Venables, W. N. & Ripley, B. D. Modern Applied Statistics with S. Preprint at

https://www.stats.ox.ac.uk/pub/MASS4/ (2002).

140. Benson, N. C., Butt, O. H., Brainard, D. H. & Aguirre, G. K. Correction of distortion in flattened

representations of the cortical surface allows prediction of V1-V3 functional organization

from anatomy. *PLoS Comput. Biol.* **10**, e1003538 (2014).

141. Finzi, D. *et al.* Differential spatial computations in ventral and lateral face-selective regions

are scaffolded by structural connections. *Nat. Commun.* **12**, 2278 (2021).

142. Kuznetsova, A., Brockhoff, P. B. & Christensen, R. H. B. lmerTest Package: Tests in Linear

Mixed Effects Models. *J. Stat. Softw.* **82**, 1–26 (2017).

1174
1175

1176