White matter connections of human ventral temporal cortex are organized by
cytoarchitecture, eccentricity, and category-selectivity from birth

Emily Kubota[1]*, Xiaoqian Yan[2], Sarah Tung[1], Bella Fascendini[3], Christina Tyagi[1], Sophie Duhameau[1], Danya
Ortiz[1], Mareike Grotheer[4,5], Vaidehi S. Natu[1], Boris Keil[5,6,7,8], & Kalanit Grill-Spector[1,9]

1. Department of Psychology, Stanford University, 450 Jane Stanford Way, Stanford, CA 94305, USA
2. Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai, China
3. Department of Psychology, Princeton University, Peretsmfan Scully Hall, Princeton, NJ 08540, USA
4. Department of Psychology, Philipps-Universität Marburg, Frankfurter Str. 35, Marburg 35037, Germany
5. Center for Mind, Brain and Behavior – CMBB, Universities of Marburg, Giessen, and Darmstadt, Marburg 35039, Germany
6. Institute of Medical Physics and Radiation Protection, TH Mittelhessen University of Applied Sciences, Giessen 35390, Germany
7. Department of Diagnostic and Interventional Radiology, University Hospital Marburg, Philipps-Universität Marburg, Baldinger Str.,
Marburg 35043, Germany.
8. LOEWE Research Cluster for Advanced Medical Physics in Imaging and Therapy (ADMIT), TH Mittelhessen University of Applied
Sciences, Giessen 35390, Germany.
9. Wu Tsai Neurosciences Institute, 288 Campus Drive, Stanford, CA 94305 USA

Corresponding author*: emily.kubota@gmail.com

18
Abstract

Category-selective regions in ventral temporal cortex (VTC) have a consistent anatomical
organization, which is hypothesized to be scaffolded by white matter connections. However, it is
unknown how white matter connections are organized from birth. Here, we scanned newborn to 6-
month-old infants and adults and used a data-driven approach to determine the organization of the
white matter connections of VTC. We find that white matter connections are organized by
cytoarchitecture, eccentricity, and category from birth. Connectivity profiles of functional regions
in the same cytoarchitectonic area are similar from birth and develop in parallel, with decreases in
endpoint connectivity to lateral occipital, and parietal, and somatosensory cortex, and increases to
lateral prefrontal cortex. Additionally, connections between VTC and early visual cortex are
organized topographically by eccentricity bands and predict eccentricity biases in VTC. These data
have important implications for theories of cortical functional development and open new
possibilities for understanding typical and atypical white matter development.

Introduction

Human ventral temporal cortex (VTC) contains regions that are selective for categories that are important for our everyday lives such as faces[1], bodies[2], words[3], and places[4]. These regions are located in consistent anatomical locations across individuals[5–10]. A key debate in cognitive neuroscience is what structural and functional factors contribute to the consistent functional organization of VTC. One central theory is that innate anatomical constraints and, particularly, white matter connections between VTC and other parts of the brain may lead to the emergence of category-selective regions in consistent anatomical locations relative to cortical folding[11–16]. Indeed, recent research in adults has revealed that the white matter connections of VTC are highly regular with respect to cortical folds[17–19], cytoarchitecture[20], eccentricity[21], and category-selectivity[11,12,22]. However, it is unknown which of these organizing principles of white matter connections of VTC are already present in infancy and if white matter connections remain stable from infancy to adulthood. Here, we address this gap in knowledge by using anatomical and diffusion MRI in infants and adults to elucidate the organization principles of white matter connections of VTC from birth to adulthood.

We first consider multiple hypotheses of how white matter connections at birth may constrain the development of VTC function. The *category hypothesis* suggests that category-selective regions in VTC have category-specific connections forming specialized networks that enable processing of category-specific information[15,23,24].This theory is supported by studies showing that, in adults, white matter connectivity profiles predict the location of face, place, and word-selective regions in VTC[11,22,25] and in children, white matter connectivity profiles at age five predict where word-selective regions will emerge at age eight[13]. Further, functional connectivity of resting state data in infants show category-specific patterns[26,27], suggesting that category-specific networks may be present from birth. The category hypothesis thus predicts that white matter connections of VTC will be organized by category from birth.

The *cytoarchitecture hypothesis* suggests that white matter connections are linked to cytoarchitectonic areas – areas defined by their distribution of cell density across cortical layers[28–30]. VTC contains four cytoarchitectonic areas: FG1-FG4[31,32], which have different neural hardware that is thought to support functional computations. Notably, in both children and adults, regions with different category-selectivity within the same cytoarchitectonic area, e.g., face-selective and word-selective regions located in FG4, have similar white matter connectivity profiles[20]. But regions with the same category-selectivity located in different cytoarchitectonic regions, e.g., face-selective regions located in FG2 and FG4, respectively, have different white matter connectivity profiles[20]. The cytoarchitectonic hypothesis therefore predicts that white matter connections of VTC will be organized by cytoarchitecture from birth.

The *eccentricity-bias hypothesis* suggests that eccentricity biases in VTC are due to innate white matter connections with early visual cortex (EVC, union of V1-V2-V3), where faces and word-selective regions have more connections with foveal EVC and place-selective regions have innate

white matter connections with peripheral EVC. This theory is supported by findings that in both
children and adults, face and word-selective regions in VTC have a foveal bias[5,7,8,33], whereas place-
selective regions in VTC have a peripheral bias[5,7,8,33]. Additionally, patterns of functional connectivity
between face- and place-selective regions in VTC and EVC in infant humans, neonate monkeys,
and congenitally blind adults follow eccentricity bands[34−38]. That is, face-selective regions have
higher functional connectivity to central eccentricity bands in EVC and place-selective regions have
higher functional connectivity to peripheral eccentricity bands in EVC. Finally, in adults, white matter
connections between VTC and EVC also correspond to eccentricity bands, with face-selective
regions having more connections to central EVC eccentricities and place-selective regions having
more connections to peripheral ones[21]. Therefore, the eccentricity-bias hypothesis predicts that the
subset of white matter connections between VTC to EVC will have a non-uniform distribution,
where regions in VTC that have foveal bias in adults (faces and words) will have more connections
with central bands in EVC from birth, and regions that have a peripheral bias (places) will have more
connections with peripheral bands in EVC from birth.

In addition to the unknown organizational principles of white matter connections in infancy, it is
also unknown if white matter connectivity profiles are stable or change over development.
Accumulating evidence reveals that the large fascicles (axonal bundles that travel in parallel
connecting distant parts of the brain) are present at birth[39−43]. Nonetheless, some fascicles
continue to develop after birth. For example, the arcuate fasciculus (AF), which connects the VTC
with lateral prefrontal cortex is not fully developed in infancy: it has a smaller cross section in
children compared to adults[44] and, and in infants, it reaches premotor cortex but not lateral
prefrontal cortex as it does in adults[45]. Additionally, mature white matter connections in adulthood
may depend on visual experience as visual deprivation during infancy and childhood leads to
degradation of white matter tracts of the visual system[46−50]. However, it is unknown which aspects
of VTC white matter connectivity profiles are innate and which aspects may develop from infancy
to adulthood.

To address these key gaps in knowledge, we obtained diffusion magnetic resonance imaging
(dMRI) and anatomical MRI in infants and adults and evaluated the organization and development
of white matter connections of category-selective regions of VTC.

Results

We collected MRI data from 44 newborn to 6-month-old infants (26 longitudinally) during natural
sleep using 3T MRI over 77 sessions, as well as from 21 adults (28.21±5.51 years), obtaining whole
brain anatomical MRI (T1-weighted, T2-weighted) and multishell dMRI data in each participant. We
implemented several quality assurance measures to ensure high quality data (see Methods: Quality
Assurance). Four infant sessions were excluded because of missing diffusion data, and six infant

sessions were excluded after quality assurance. We report data from 67 infant sessions (21
longitudinal infants) and 21 adult sessions with no significant differences in data quality (see
Methods). In total, we report data from 88 sessions: 23 sessions from newborns (mean
age±standard deviation: 28.6±10.2 days), 23 sessions from 3-month-olds (106.9±19.3 days), 21
sessions from 6-month-olds: (189.0±15.8 days), and 21 sessions from adults (28.21±5.51 years).
Anatomical MRIs were used to segment the brain to gray and white matter, define the gray-white
matter boundary to seed the tractography, and create cortical surface reconstructions. dMRI data
was used to derive the whole brain white matter connectome of each individual and session.

To determine how white matter connections are organized in infants and adults and how they may
constrain the development of regions in VTC, we project adult functional regions of interest (fROIs)
to the native space of each individual and determine the white matter connections of each fROI.
We reason that this procedure enables determining the relationship between white matter
connectivity in infancy and the eventual location of each fROI in adulthood. To do so, we use cortex
based alignment[51] to project maximal probability maps (MPMs) of adult category-selective regions
from independent data[20] into each individual brain and session. MPMs of fROIs selective for faces,
bodies, and places (mFus-faces, pFus-faces, OTS-bodies and CoS-places) are bilateral, and MPM
fROIs selective for words (mOTS-words and pOTS-words) are in the left hemisphere only (Fig 1A,B).
We measure the surface area of fROIs relative to the surface area of each hemisphere in all infant
and adult sessions and test if the relative fROI size changes over development (as developmental
effects plateau, we use a logarithmic model: fROI size/surface area ~ log10(age in days)). We find
no development as the relationship between the surface area of fROIs relative to hemisphere
surface area does not change with age (t(878) = 8.77x10^{-6}, p = 0.88). We also visually inspected in
each infant and adult session the location of fROIs and cytoarchitectonic areas relative to the
cortical folding to test if there is a consistent relationship across age. We find that both fROIs and
cytoarchitectonic areas are aligned to the expected anatomical landmarks in nearly each infant
(fROIs; LH: 66/67, RH: 65/67, cytoarchitectonic areas; LH: 67/67, RH: 62/67) and adult (fROIs: LH:
21/21, RH: 21/21, cytoarchitectonic areas; LH: 20/21, RH: 21/21) (Supplementary Figs 1-8). For
example, as in prior studies[20,52] CoS-places aligns to the intersection of the anterior lingual sulcus
and collateral sulcus[6,9], and the boundaries between F1 and FG2 and between FG3 and FG4 are
aligned to the MFS. Therefore, in both infants and adults there is a consistent relationship between
category-selective regions and cytoarchitectonic areas[20,52], where MPMs for pOTS-words and
pFus-faces are within FG2, CoS-places is within FG3, and mFus-faces, OTS-bodies, and mOTS-
words are within FG4 (Fig 1C).

Figure 1. Functional atlas and analysis pipeline. A) mFus-faces aligns to the mid-fusiform sulcus in the fsaverage brain space (left) and B) the newborn brain space (right). C) *Left:* Functional region of interest (fROI) atlas from 28 adults in[20] and Atlas of cytoarchitectonic areas from[53] projected to the fsaverage brain; *Right:* the same atlases projected into an example newborn's brain (17 days old). Solid colors represent fROIs. Salmon *pink:* mFus-faces (mid fusiform face-selective region); *Light blue:* mOTS-words (mid occipital temporal sulcus word-selective region); *Yellow:* OTS-bodies (occipital temporal sulcus body and limb-selective region); *Red:* pFus-faces (posterior fusiform face-selective region); *Blue: p*OTS-words (posterior occipital temporal sulcus word-selective region). Green: CoS-places (collateral sulcus place-selective region). Outlines represent cytoarchitectonic areas[53]. *Cyan:* FG2; *Green:* FG3; *Magenta:* FG4. Both fROIs and cytoarchitectonic areas align to the expected anatomical landmarks in the newborn. All individuals data in Supplementary Figs 1-8. D) Pipeline to define white matter connectivity profiles for each fROI. *Left:* example connections of mFus-faces in an individual newborn participant; *Middle*: endpoint map on the cortical surface; *Right:* quantification of endpoint density within each Glasser ROI. E) Principal Components Analysis (PCA) plot across the first two PCs: PC1 explains 38% percent of the variance, and PC2 explains 20% percent of the variance. Each dot represents a single connectivity profile in a single subject, the salmon pink dot represents the connectivity profile shown in (D). (F-H) Schematic illustrating predictions of three hypotheses in PC space. Each dot represents a connectivity profile for a given fROI and session. (F) Category hypothesis: schematic demonstrating what the data might look like if the connectivity profiles were organized by category. *Green:* place-selective, *blue:* word-selective, *red:* face-selective, *yellow:* body-selective. G) Cytoarchitectonic

hypothesis: schematic demonstrating what the data might look like if the connectivity profiles were organized by
cytoarchitecture. *Cyan:* FG2, *green:* FG3, *magenta:* FG4. H) Age hypothesis: schematic demonstrating what the data might
look like if the connectivity profiles were organized by age. *Salmon pink:* 0-months, *green:* 3 months, *blue:* 6 months, *purple:*
adults.

We use a data-driven approach to determine the organizing principles of white matter connections
of VTC. As the category and cytoarchitecture hypotheses make predictions about connectivity
across the whole brain, whereas the eccentricity-bias hypothesis makes predictions only about
connectivity with EVC, we first test if white matter connections of whole brain connectivity profiles
are organized by category-selectivity or cytoarchitecture. In addition, we test if whole brain
connectivity profiles are organized by age to determine if connectivity profiles are stable or
changing over development. We separately examine the eccentricity-bias hypothesis in the next
section.

We derive each fROI's white matter connections by intersecting each fROI with the whole brain
white matter connectome derived from dMRI in each session (Fig 1D). We quantify the white matter
connectivity profile by measuring the endpoint density of each fROI's connections on the whole
brain as the proportion of connections ending in each of 169 Glasser Atlas ROIs[54] (180 Glasser
ROIs excluding 11 VTC ROIs, Fig 1D). This results in 880 unique connectivity profiles (10 fROIs x 88
sessions across infants and adults). Then we use principal component analysis (PCA) to reduce
the dimensionality of the connectivity profiles. The first 10 principal components (PCs,) explain 98%
of the variance in the data, with the first and second components explaining 38% and 20%,
respectively.

To visualize how connectivity profiles cluster, we plot the connectivity profiles of all fROIs and
sessions in PC space (Fig 1E). The plot consists of 880 dots including data from all infant and adult
participants, where each dot represents a single connectivity profile of an fROI from one session.
Note that each white matter connectivity profile has several non-mutually exclusive features: the
category-selectivity of the fROI, the cytoarchitectonic area in which the fROI is located, and the age
of the participant. Therefore, we can label each connectivity profile by each of these features
(category, cytoarchitecture, age group) and test if connectivity profiles cluster by one or more of
these features. For example, if connectivity profiles are organized by category, then connectivity
profiles of fROIs with the same category-selectivity would be nearby in PC space, and separate
from connectivity profiles of fROIs with a different category-selectivity (Fig 1F). However, if
connectivity profiles are organized by cytoarchitecture, then connectivity profiles of fROIs in the
same cytoarchitectonic area would be nearby in PC space (Fig 1G). Further, if connectivity profiles
change across development, then connectivity profiles of newborns may be different than other
age groups and connectivity profiles of adults may be most distinct (Fig 1H).

Labeling connectivity profiles by fROIs' preferred category reveals some clustering by category (Fig
2A). For example, connectivity profiles of face- and place-selective ROIs tend to be clustered and
separate from others. However, clustering by category is imperfect, as connectivity profiles of

word-selective regions split into two clusters, and there is overlap between the connectivity profiles
of word- and body-selective fROIs. In contrast, labeling connectivity profiles by fROIs'
cytoarchitectonic area reveals a strikingly clear organization whereby connectivity profiles of each
of the cytoarchitectonic areas FG2, FG3, FG4 form a different cluster with a distinct connectivity
profile (Fig 2B). Examination of the PC loadings (Supplementary Fig 9) and the coefficients across
the first 2 PCs suggests that connectivity profiles of fROIs in different cytoarchitectonic areas
largely vary in their connections to early visual, lateral occipito-temporal, and anterior temporal
cortex. Finally, labeling connectivity profiles by participant's age group reveals that connectivity
profiles are intermixed across age groups with no clear organization by age across the first 2 PCs
(Fig 2C).

206
Figure 2. White matter connections are organized by cytoarchitecture and category from infancy. (A-C) White matter
connectivity profiles of all participants and fROIs projected to the first two principal components (PC1 and PC2); each
dot is a connectivity profile of a single fROI and session. The same data is shown in all three panels, except that they are
labeled in color by different features. (A) Category: *red*: faces, *green:* places, *blue*: words, *yellow:* bodies, (B)
Cytoarchitectonic area: *cyan:* FG2, *green:* FG3, *magenta:* FG4. (C) Age group: *salmon:* newborn, *green:* 3 months, *teal:* 6
months, *purple*: adults. (D-E) Classification accuracy of a feature from white matter connectivity profiles, using
multinomial logistic regression with leave-one-out-participant cross validation. (D) Classification accuracy of category.
(E) Classification accuracy of cytoarchitectonic area. (F) Classification accuracy of age group. In D-F, *bar:* mean
performance across participants; *Error bars*: standard error of the mean across participants; *Dashed line*: chance level.

To utilize the information across all 10 PCs, we use a leave-one-out classification approach to test
if a classifier trained on white matter connectivity profiles of n-1 participants can predict the
category preference, cytoarchitectonic area, or the participant's age group from the left out
participant's connectivity profiles. We find that category (Fig 2D, mean accuracy±SD = 83%±10%,
t(87) = 47.09, p < 2.2x10^{-16}), cytoarchitecture (Fig 2E, mean accuracy±SD= 97%±5%, t(87) = 109.6,
p < 2.2x10^{-16}), and age (Fig 2F, mean accuracy±SD = 41%± 27%, t(87) = 5.48, p < 3.99x10^{-7}) are all
classified significantly above chance. Classification of cytoarchitecture is significantly higher than
classification of category preference (odds ratio= 5.96, 95% CI [3.98 8.93], binomial logistic
regression) and age (odds ratio = 41.51, 95% CI [28.16, 61.19], binary logistic regression). Confusion
matrices reveal almost no error for cytoarchitecture classification (Supplementary Fig 10A), but
some category errors that arise from confusion of word-and body-selectivity as well as confusion
of body and face-selectivity (Supplementary Fig 10B). The confusion matrix for age classification
(Supplementary Fig 10C) reveals that newborns and adults are classified the best, that 3 month-
olds are often confused for newborns, and that 6 month-olds are equally confused across all age
groups. This suggests that there is heterogeneity in the connectivity profiles of 6-months-olds, and
that around 6 months of age, the white matter connections of some infants may start to become
more adult-like.

To assess if the high classification performance of cytoarchitecture is due to higher anatomical
proximity of fROIs within the same cytoarchitectonic area compared to those in distinct
cytoarchitectonic areas, we repeat the classification analysis on connectivity profiles of equidistant
disk ROIs that are either in the same (FG4) or different (FG2/FG4) cytoarchitectonic areas
(Supplementary Fig11A,B). We reason that if anatomical proximity drives classification results,
then classification accuracy will be reduced once distance is controlled. However, even when
distance was held constant, cytoarchitecture is classified from the connectivity profiles with high
accuracy (Supplementary Fig11C, mean accuracy±SD=95%±22%, t(87) = 31.62, p < 2.2x10^{-16}),
suggesting that organization of connectivity profiles by cytoarchitecture is not just due to
anatomical distance between fROIs.

Interestingly, both category and cytoarchitecture classification are not significantly different across
age groups (comparison between newborns, 3-month-olds, and 6-month-olds compared to adults
(newborns: odds ratio =1.20, 95% CI [0.77, 1.89], 3-months: odds ratio =1.01, 95% CI [0.65, 1.56], 6-
months: odds ratio =1.05, 95% CI [0.67, 1.65], binomial logistic regression). Indeed, examination of
the organization structure across the first 2 PCs separately for each age group reveals similar
organization across age groups (Supplementary Fig 12). These analyses suggest that
organizational features of white matter connections by cytoarchitecture and category persist
across development.

To test the eccentricity-bias hypothesis we next examined the subset of connections between each
fROI and early visual cortex (EVC). We reasoned that if connections between VTC and EVC are
organized by eccentricity from birth, then fROIs that overlap with foveal representations in adults
(pFus-faces, pOTS-faces, mFus-faces, OTS-bodies, mOTS-words) would have more white matter
connections to central than peripheral eccentricities in EVC in infancy, and fROIs that overlap with
peripheral representations in adults (CoS-places) would have more connections to peripheral than
central eccentricities in VTC in infancy. To test these predictions, we measured the distribution of
white matter endpoints between each of the VTC category-selective fROIs and three eccentricity
bands in EVC: 0-5°, 5-10°, and 10-20° and compared across age groups.

We first visualized the white matter connections between each fROI and EVC, coloring the white
matter connections according to the endpoint eccentricity in EVC. This visualization reveals a
striking orderly arrangement of the connections from each eccentricity band to each of the fROIs
in both infants and adults (Fig 3A and 3B; see Supplementary Videos 1-4 for 3D visualization).
Specifically, in all age groups, newborns to adults, there is a lateral to medial arrangement of
connections such that connections to EVC central eccentricities (red, 0-5°) are more lateral, and
connections to more peripheral EVC eccentricities are more medial (green, 5-10° to blue, 10-20°;
Fig 3A,B).

We then quantified the distributions of endpoint connections to EVC for each fROI and age group.
Consistent with the predictions of the eccentricity-bias hypothesis, in all age groups, pFus-faces,
pOTS-words, mFus-faces, OTS-bodies, and mOTS-words had more connections to the EVC 0-5°
eccentricity band than the more peripheral eccentricities (Fig 3C) and CoS-places had more
connections to the EVC 10-20° eccentricity band than to the more central eccentricities (Fig 3C).
Additionally, the connections from VTC to EVC by eccentricity band also mirror the
cytoarchitectonic organization of the VTC connectivity profile. Specifically, across all age groups:
(i) pFus-faces and pOTS-words, which are located in cytoarchitectonic area FG2, have ~60% of
their connections to the EVC 0-5° eccentricity band, whereas (iii) CoS-places, which is located in
FG3, has ~40% of its connections to the EVC 10-20° eccentricity band, and (iii) mFus-faces, OTS-
bodies, and mOTS-words in FG4 have 40-50% of their EVC connections in the central 5° (Fig 4C).

We further tested how eccentricity is related to the features of cytoarchitecture, category, and age
that we examined in the whole brain analysis. To do so, for each fROI we fit a linear model predicting
the percentage of connections to the central 5° by cytoarchitecture, category-selectivity, and age
group. As the percentage of connections to each eccentricity band is proportional, we use the
percentage of connections to the central 5° as the independent variable, where a greater
percentage connections to the central 5° indicate more connections to central eccentricity bands,
and fewer connections to the central 5° indicate more connections to peripheral eccentricity bands.

289
Figure 3. White matter connections of VTC are organized by eccentricity from birth. Connections between each fROI and
early visual cortex (EVC, union of V1, V2, V3) in an example newborn (A) and an example adult (B). Here we report data
for the left hemisphere as it includes all 6 fROIs; data for the right hemisphere is reported in Supplementary Fig 53.
Connections are colored by the endpoint eccentricity band in EVC: *red:* 0-5°, *green*: 5-10°, *blue*: 10-20°. (C) Quantification
of the percentage of endpoints in each eccentricity band by age group for each fROI: Violin plots indicate distributions of
the percentage of connections for each age group and eccentricity band. Black dot and error bar indicates mean±
standard error of the mean. *Red:* 0-5°, *green*: 5-10°, blue: 10-20°, see inset. *0M:* newborns, *3M:* 3 month-olds, *6M:* 6 month-
olds, A: adults. Horizontal lines in the bottom indicate cytoarchitectonic area..

We find a main effect of cytoarchitecture (F(2,504) = 390.02, *p* < 2.2x10⁻¹⁶), suggesting that regions
that are in distinct cytoarchitectonic areas have different percentages of connections going to the
EVC central 5°. In particular, FG2 has a majority of its connections going to the central 5° in EVC,
but FG3 has a minority (mean±sd: FG2: 63.57%±14.41%, FG3: 26.58%±8.33%, FG4:
43.97%±9.34%). Additionally, there is a main effect of category-selectivity (F(2,504) = 8.32, *p*=
0.0003), reflecting that regions with different category-selectivity also have different percentages
of connections going to the EVC central 5°. Face- and word-selective regions have a majority of
connections going to the central 5°, but place-selective regions have only a minority (mean±sd:
faces: 51.70%±15.02%, words: 56.21%±15.73%, places: 26.58%±8.33%, bodies: 43.23%±9.05%).
In contrast, there is no main effect of age group (F(3,504) = 1.14, p = 0.33), and no significant
interaction between category and age (F(6,504) = 0.93, p = 0.47). There are, however, significant
interactions between cytoarchitecture and category (F(1,504) = 24.55, p = 9.91x10⁻⁷),
cytoarchitecture and age (F(6,504) = 3.59, p = 0.002), and between cytoarchitecture, category, and
age (F(3,504) = 2.76, p = 0.04). Post-hoc tests reveal that the cytoarchitecture by category
interaction is driven by differences in percentage of connections to the central 5° between pFus-
faces and pOTS-words in FG2 (t(87) = 5.68, p = 1.77 x 10⁻⁷), as well as differences in percentage of
connections to the central 5° between mFus-faces and OTS-bodies in FG4 (t(87) = 3.68, p = .0004).
Further post-hoc tests reveal that differential development is due to a decrease in the percentage
of connections from FG4 to the EVC central 5° over development (β = -2.75, *t*(163.09) = -3.39, *p* =
0.0009), and a differential development within FG2 where there is a significant increase in the

connections of pOTS-words to the central 5° over development (β = 3.95, t(81.26) = 3.52, p =
0.0007), but no significant development in the percentage of pFus-faces connections to the central
5° (β = -0.78, *t*(85.90) = -0.47, *p* = 0.64). These results suggest that in addition to cytoarchitecture,
and category, white matter connections of VTC are organized topographically by eccentricity from
birth and that the differential proportion of connections between VTC to different eccentricity bands
from birth mirrors the eccentricity bias of both cytoarchitectonic areas and category-selective
regions in VTC.

Do White Matter Connectivity Profiles of VTC Develop from Infancy to Adulthood?

In addition to revealing the organizing principles of white matter connections by cytoarchitecture,
category, and eccentricity, we also found evidence for development as we were able to classify the
age group of the participant above chance from the whole connectivity profile, and we find some
developmental changes in connectivity with EVC. To qualitatively assess how the whole brain white
matter connections of VTC change over development, we visualized the white matter connections
of all category-selective fROIs in all participants. Fig 4 shows an example infant at newborn, 3
months, and 6 months as well as an example adult (connections of all participants in
Supplementary Figs 12-50).

Consistent with the prior analyses, within an age group, white matter connections of fROIs in the
same cytoarchitectonic area were similar. For example, in the newborn, pFus-faces and pOTS-
words, both in FG2, are characterized by vertical connections to dorsal occipital cortex and
longitudinal connections to both the anterior temporal lobe and early visual cortex (Fig 3A, top 2
rows). However, we also observe developmental changes. For example, pFus-faces and pOTS-
words appear to have more abundant vertical connections in infants than adults, but more
connections to the frontal cortex in adults than infants (Fig 3A, Supplementary Figs 20-27, 47-50
all participants). We see a similar pattern for the white matter connections of fROIs in FG4: mFus-
faces, OTS-bodies, and OTS-words show similar connectivity profiles at each timepoint (e.g.,
newborn), yet we also observe developmental differences: vertical connections appear more
abundant in newborns than adults, but connections to the frontal cortex appear to be more
abundant in adults than infants (Fig 3A, bottom row, Supplementary Figs 12-19, 28-35,44-46). For
CoS-places, located medially in FG3, there is a different connectivity profile and a different
developmental pattern: in newborns, CoS-places has abundant frontal, vertical, and longitudinal
connections to both occipital and anterior temporal lobes, but in adults there appear to be fewer
frontal and vertical connections (Fig 3A-third row, Supplementary Figs 36-43).

Figure 4. Connectivity of VTC develops from infancy to adulthood . (A) Connections of each category-selective ROI in an individual infant (bb05) across three time points (newborn, 3 months, 6 months) next to an adult control (A11) shown in a sagittal cross section. Brains are shown to scale, see scale bar bottom left. Each row is a different fROI (indicated by color) and ordered by cytoarchitectonic area, from top to bottom: FG2 to FG4. (B) Change in endpoint density over development for each fROI's connectivity profile. The color of the Glasser ROI indicates the development in endpoint density over time, namely the slope of the regression: endpoint density ~ log10(age in days) for 88 sessions. Here we report data for the left hemisphere because it includes all 6 fROIs. Data for the right hemisphere is in Supplementary Fig 51. *Green*: decreasing endpoint density over development, *magenta:* increasing endpoint density over development. *Black arrow: primary visual area* (V1), *Green arrow:* primary somatosensory area(S1). Each row shows the development of white matter connections of an fROI; rows are ordered identically to (A).

To quantify developmental changes, we test whether each fROI's endpoint density, quantified as the proportion of connections in each Glasser ROI, varies as a function of age (fROI endpoint density ~ log10(age in days) x Glasser ROI, linear model). As expected, fROIs' endpoint density vary across the brain (significant main effect of Glasser ROI; $Fs > 3.54$, $ps < 2.2 \times 10^{-16}$; statistics in Supplementary Table 1) reflecting that fROIs have non-uniform connectivity across the brain.

Additionally, fROIs' endpoint density differentially vary with age across the brain (significant age by
Glasser ROI interaction; $Fs > 74.94$, $ps < 2.2 \times 10^{-16}$; statistics in Supplementary Table 1). As we find
a differential development across the brain, for each fROI, we calculate the change in endpoint
density across age per Glasser ROI (endpoint density ~ log10(age in days), linear model). Then, we
visualize the slope of this model for each Glasser ROI (Fig 4D). A positive slope (magenta) indicates
that the endpoint density increases from infancy to adulthood within the Glasser ROI, whereas a
negative slope (green) indicates that endpoint density decreases with age. This analysis reveals
both developmental increases and decreases in endpoint density with distinct spatial
characteristics. First, there are common developmental patterns for face, word, and body fROIs
(Fig 4-right): endpoint densities to lateral occipital and dorsal occipito-parietal visual Glasser ROIs
as well as somatosensory areas decrease (Fig 4-green arrows), but endpoint densities to lateral
prefrontal Glasser ROIs increase (Fig 4). There are also differential developmental patterns of
connectivities of fROIs in FG2 (pOTS-words, pFus-faces) compared to FG4 (mFus-faces mOTS-
words, and OTS-bodies): the former show developmental decreases in endpoint densities in V1 and
early visual cortex more broadly (Fig 4-medial view) but the latter show developmental increases.
Finally, for CoS-places, which is in FG3, we find mostly developmental decreases in endpoint
densities in both visual and orbitofrontal cortex with increases in endpoint densities in ventral
temporal lateral regions.

Finally, we test whether development was more similar for fROIs with the same cytoarchitectonic
area or same category-selectivity. We use a bootstrapping procedure to estimate the
developmental slopes (see Methods), and evaluate the similarity between white matter
development by calculating the pairwise correlation between the development slopes across
Glasser ROI for each pair of fROIs. We then fit a linear model to test whether the correlation was
higher for fROIs with the same cytoarchitecture or category-selectivity (linear model: Fisher
Transformed correlation ~ cytoarchitecture + category). We find a significant main effect of
cytoarchitecture ($F(1,14997) = 61,161.5$, $p < 2.2 \times 10^{-16}$), reflecting that regions within the same
cytoarchitectonic area had more similar developmental slopes (correlation$\pm$sd: .78$\pm$11) than
regions in different cytoarchitectonic areas (correlation$\pm$sd: .25$\pm$13). There was also a significant
main effect of category ($F(1,14997) = 1,069.1$ $p < 2.2 \times 10^{-16}$). However, here we find that regions
with the same category-selectivity had less similar developmental slopes (correlation$\pm$sd: .36$\pm$16)
than regions with different category-selectivity (correlation$\pm$sd: .40$\pm$.28). These analyses suggest
that although the relationship between white matter and cytoarchitecture is consistent over
development, connectivity profiles develop in parallel within each cytoarchitectonic area, with both
developmental increases and decreases in endpoint densities.

Discussion

Here, we use anatomical and diffusion MRI in infants within the first six months of life and in adults to determine the organizing principles and developmental trajectories of the white matter connections of regions in VTC. We find that white matter connections are organized by cytoarchitecture, eccentricity, and category from birth. We also find evidence for development of white matter connections of VTC, with increasing endpoint density to lateral frontal cortex, and decreasing endpoint density to lateral occipital, parietal, somatosensory, and orbitofrontal cortex. These findings have important implications for understanding the interplay between nature and nurture on both white matter connections and functional brain development.

Innate aspects of white matter connections of VTC: cytoarchitecture and eccentricity are parsimonious principles

A predominant theory suggests that the consistent organization of category-selective regions in VTC is due to innate white matter connections that support category-specific processing[11,12,15,26]. While we find that in infancy white matter connections are organized to a certain extent by category, especially for faces and places, several of our empirical findings suggest that cytoarchitecture and eccentricity are more parsimonious organizing principles of white matter connectivity from infancy to adulthood. Even as both category and cytoarchitecture can be classified from whole brain white matter connectivity profiles irrespective of age, classification of cytoarchitecture is higher than category-selectivity. In addition, the subset of connections betweenVTC and early visual cortex are organized topographically by eccentricity bands and show similar eccentricity biases for functional regions that are selective for different categories but located in the same cytoarchitectonic area. Finally, connectivity profiles of functional regions in the same cytoarchitectonic area largely develop in parallel from infancy to adulthood. This organization by cytoarchitecture and eccentricity suggests that these are general principles underlying the white matter organization of the visual system, which may predate and constrain the coupling of white matter connections and brain function. That is, our data suggest a theoretical shift: rather than humans being born with specialized connections to support category-specific processing, there may be more general principles underlying the white matter organization of the visual system that correspond to cytoarchitectonic boundaries and eccentricity gradients. Indeed, in children and adults cytoarchitecture and eccentricity are linked as the boundary between lateral and medial cytoarchitectonic areas in VTC align to the transition between foveal and peripheral representations in VTC[20,52,55,56].

As cytoarchitectonic[30,53,57−60] and retinotopic organization[7,8,21,61−63] are prevalent across the entire visual system, we hypothesize that these two organizing principles may underlie the white matter organization of the visual system more broadly. This hypothesis can be tested in future research, for example, in the parietal cortex that contains a series of retinotopic[64,65] and cytoarchitectonic[58] areas. Critically, as cytoarchitecture[30,54,6630,67−70], and topography[71,72] are also key features distinguishing brain areas beyond the visual system, e.g., language[73], our findings raise the

442 possibility that white matter connections are innately organized by cytoarchitecture and
443 topographic gradients even beyond the visual system and throughout the entire brain. This
444 hypothesis can be tested in future research leveraging large anatomical and diffusion MRI datasets
445 that are being collected in infants[74–78].

Development of white matter connections of VTC

While we find innate organization of white matter connections, we also find that the connectivity
profiles of fROIs within the same cytoarchitectonic area develop in parallel from infancy to
adulthood. Endpoint density from VTC to the frontal lobe showed both developmental increases
and decreases. Endpoint density from FG2 and FG4 in lateral VTC to prefrontal cortex increased
over development but from FG3 in medial VTC to orbitofrontal cortex decreased. The former finding
is consistent with observations that the arcuate fasciculus[79] is underdeveloped in infants, only
reaching the precentral gyrus and premotor cortex in newborns, and not reaching lateral prefrontal
cortex as it does in adults[45,80].

Additionally, we find developmental decreases in endpoint density between VTC and lateral
occipital, parietal, and somatosensory cortex. Our observations in the human visual system mirror
results from a large body of tracer studies in cats and macaques that documented exuberant
connections between V1, V2, and V3[81–83] as well as between early visual areas and somatosensory
areas[84]. These exuberant connections are present early in development and then eliminated by
adulthood[82,84–86]. Our data not only suggest that exuberant connections may exist in the human
visual system, but also suggest the possibility that connections between VTC and both the lateral
and dorsal visual streams[17,87–90] may decrease over development.

We acknowledge that our *in-vivo* dMRI measurements do not enable us to make inferences about
the cellular and molecular underpinnings of the development of white matter connectivity profiles.
However, our findings provide new ideas that can be investigated in future histological and animal
research. Future studies can examine if developmental increases in endpoint density may be
related to increased myelination[91–94], axonal sprouting[95–97], and glial proliferation[92,94] that are
associated with activity-dependent white matter plasticity[98,99]. In addition, it can be tested whether
developmental decreases in endpoint density may be related to elimination of exuberant long range
projections[82,84–86].

Implications for theories of functional development of VTC

Several theories have posited that innate white matter connections of VTC constrain its function[13–
15]. Here, we provide support for these theories by showing that the main principles of white matter
organization persist from infancy to adulthood. At the same time, new questions regarding the
functional development of VTC emerge from our findings.

First, our data demonstrate that anatomical locations corresponding to adult face and place-
selective regions already have distinct whole brain connections at birth. This finding suggests that
white matter connections are already organized by category at birth. Questions for future studies
will be at what age face and place-selective regions first appear and whether they emerge in a
stable anatomical location from infancy to adulthood, as predicted by the white matter.

Second, we find that white matter connections between EVC and VTC are organized by eccentricity
from birth. This discovery is consistent with studies in infant human and nonhuman primates[26,33]
finding stronger functional connectivity between face-selective regions and foveal V1, and place-
selective regions and peripheral V1. Together, these observations suggest that white matter
connectivity has functional ramifications, and lead to the prediction that eccentricity biases in
VTC[7,8] will be present from birth.

Third, our findings that the white matter connections of VTC have both innate organization and the
capacity to change over development raise new questions about the relationship between white
matter and function in atypical development. One question is whether lack of visual experience may
reshape white matter connectivity profiles of VTC and consequently function[14]. For example, it is
unknown if lack of visual experience may lead to the preservation of connections between VTC and
somatosensory cortex, which in turn, may enable VTC to respond to haptic inputs in individuals
who are congenitally blind[100]. Critically, our framework enables measuring the fine-grained white
matter connections of VTC in individual infants longitudinally[101], increasing accuracy and precision
in measuring the interplay between white matter connections, functional regions, and anatomy
across development. Our framework opens new opportunities not only for evaluating development
of white matter associated with functional regions in large infant datasets[74–78] but also for early
identification and assessment of developmental disorders associated with VTC such as autism[102–
105], Williams syndrome[106], congenital prosopagnosia[107,108], and dyslexia[109,**102–105**].

Together, the present study advances our understanding of white matter organization in the visual
system suggesting that cytoarchitecture, eccentricity, and category are organizing principles from
birth, even as aspects of the white matter develop from infancy to adulthood. These data have
implications not only for theories of cortical functional development, but also have ramifications
for early identification of atypical white matter development.

Methods

Participants

The study was approved by the Institutional Review Board of Stanford University. Parents of the
infant participants provided written informed consent before the first scan session, and also
provided written informed consent prior to each session if they came for multiple sessions.
Participants were paid $25/hour for participation.

Expectant parent and infant screening procedure

Expectant parents and their infants in our study were recruited from the San Francisco Bay Area
using social media platforms. We performed a two-step screening process. First, parents were
screened over the phone for eligibility based on exclusionary criteria designed to recruit a sample
of typically developing infants. Second, eligible expectant mothers were screened once again after
giving birth. Exclusionary criteria were as follows: recreational drug use during pregnancy,
significant alcohol use during pregnancy (more than three instances of alcohol consumption per
trimester; more than 1 drink per occasion), taking prescription medications for a disorder involving
psychosis or mania during pregnancy, insufficient written and spoken English ability to understand
the instructions of the study, or learning disabilities. Exclusionary criteria for infants were preterm
birth (<37 gestational weeks), low birthweight (<5 lbs 8 oz), any congenital, genetic, and
neurological disorders, visual problems, complications during birth that involved the infant (e.g.,
NICU stay), history of head trauma, and contraindications for MRI (e.g., metal implants).

44 full term infants following a typical pregnancy participated in a total of 77 MRI sessions. Four
sessions were excluded for missing diffusion data, and six sessions were excluded for too much
motion (see Quality Assurance section). We report data from 42 infants over 67 sessions (21
longitudinal): sex: 16 female, 26 male; race and ethnicity: 4 Asian, 5 Hispanic, 11 Multiracial, and 22
White participants; age: n=23 newborns (M=28.56 days, SD= 10.21days), n=23 3-month-olds
(M=106.91 days, SD=19.33 days), and n=21 6-month-olds (M=189.05 days, SD=15.77 days).

We also collected data from 21 adults: sex: 17 female, 4 male; race and ethnicity: 12 Asian, 1
Hispanic, 2 Multiracial, and 6 White participants; age: M=28.21 years, SD=5.51 years. All adult
sessions met inclusion criteria.

MRI acquisitions

Scanning sessions were scheduled in the evenings around the infants' typical bedtime and were
done during natural sleep. Experimenters waited till the infant fell asleep to place the infant in the
MRI scanner and start the session. Infant data were acquired on a 3T GE Ultra High Performance

(UHP) scanner (GE Healthcare, Waukesha, WI) equipped with a customized 32-channel infant head-
coil [110].

Hearing protection included soft wax earplugs inserted to the infant's ears and then, MRI
compatible neonatal noise attenuators (https://newborncare.natus.com/products-
services/newborn-care-products/nursery-essentials/minimuffs-neonatal-noise-attenuators), and
headphones. (https://www.alpinehearingprotection.com/products/muffy-baby) that covered the
infant's ears. During sessions with newborns, an MR-safe plastic immobilizer (MedVac,
www.supertechx-ray.com) was used to stabilize the infant and their head position. When the infant
was asleep, the caregiver placed the infant on the scanner bed. Weighted bags were placed at the
edges of the bed to prevent any side-to-side movement. Additional pads were also placed around
the infant's head and body to stabilize head position. An experimenter stayed inside the MR suite
with the infant during the entire scan.
553
For additional monitoring of the infant's safety and tracking of the infant's head motion, an infrared
camera was affixed to the head coil and positioned for viewing the infant's face in the scanner. The
researcher operating the scanner monitored the infant via the camera feed, which allowed for the
scan to be stopped immediately if the infant showed signs of waking or distress. This setup also
allowed tracking the infant's motion; scans were stopped and repeated if there was excessive head
motion.
560
561 *Anatomical MRI acquisition*
562
We obtained both T1-weighted and T2-weighted MRI data for each infant and session with the
following parameters: T1-weighted image: TE = 2.9ms; TR = 6.9ms; voxel size = 0.8x 0.8 x0.8 [mm^3];
FOV = 20.5 cm; Scan time: 3:05 min; T2-weighted image: TE = 124ms; TR = 3650 ms; voxel size =
0.8x 0.8x 0.8 [mm^3]; FOV = 20.5 cm; Scan time: 4:05 min.

Anatomical MRI processing

The T1-weighted and T2-weighted images from each individual were first aligned with each other,
then aligned to a plane running through the commissures (AC-PC transformed), and then
processed with iBEAT V2.0[111] tissue segmentation. White matter segmentation was, and then
manually fixed for errors using ITKgray[112]. We used the manually edited segmentation file to
reconstruct the cortical surface with Infant FreeSurfer[113].

Diffusion MRI (dMRI) acquisition

We obtained dMRI data with the following parameters: multi-shell, #diffusion directions/b-
value = 9/0, 30/700, 64/2000; TE = 75.7 ms; TR = 2800 ms; voxel size = 2x2x2 mm^3; number of
slices = 60; FOV = 20 cm; in-plane/through-plane acceleration = 1/3; scan time: 5:08 min. We also

acquired a short dMRI scan with reverse phase encoding direction and only 6 b = 0 images (scan
time 0:20 min).

Diffusion MRI processing

The dMRI data were preprocessed using MRtrix3[114] (https://github.com/MRtrix3/mrtrix3) and in
accordance with prior work from the human connectome project and our lab [40,115,116]. Data were
denoised using principal component analysis[117]. We used FSL's top-up tool
(https://fsl.fmrib.ox.ac.uk/) and one image with reverse phase-encoding to correct for
susceptibility-induced distortions. We used FSL's eddy tool to perform eddy current and motion
correction, where outlier slices were detected and replaced[118]. Finally we performed bias correction
using ANTs[119] (https://picsl.upenn.edu/software/ants/). The preprocessed dMRI data were aligned
to the T2 weighted anatomy using whole-brain rigid body registration. Alignment was checked
manually for all images.

Generating White Matter Connectomes

We generated a whole brain white matter connectome in each session using MRTrix3[114]. Voxel-
wise fiber orientation distributions (FODs) were calculated using constrained spherical
deconvolution (CSD). We used the Dhollander algorithm[120] to estimate the three-tissue response
function. We lowered the FA threshold to 0.1 to account for the generally lower FA in infant brains.
We computed FODs separately for the white matter and CSF. As in past work[40,115], the gray matter
was not modeled separately, as white and gray matter do not have sufficiently distinct b-value
dependencies to allow for a clean separation of the signals. Finally, we generated a whole brain
white matter connectome for each session. Tractography was optimized using the gray/white
matter segmentation from anatomical MRI data (Anatomically Constrained Tractography; ACT[121]).
For each connectome, we used probabilistic fiber tracking with the following parameters: IFOD2
algorithm, step size of 0.2 mm, minimum length of 4 mm, maximum length of 200 mm, and
maximum angle of 15°. Streamlines were randomly seeded on this gray-matter/white-matter
interface, and each connectome consisted of 5 million streamlines. This procedure is critical for
accurately identifying the connections that reach fROIs, which are located in the gray matter[122].

Quality Assurance

To evaluate the quality of the diffusion data we implemented three quality assurance measures: (i)
We measured the number of outliers (dMRI volumes with signal dropout measured by FSL's eddy
tool). The exclusion criterion was >5% outlier volumes as in past work[20]. (ii) We visualized in mrView
the fractional anisotropy (FA) colored by direction of the diffusion tensors to validate the expected
maps. That is, the existence of between hemispheric connections through the corpus callosum,
and the inferior-superior directionality of the cerebral spinal tract.(iii) We used babyAFQ[22] to identify

the bundles of the brain and validate that the major bundles known to be present at birth[40,41,123] can
be found in each individual and session.

These quality assurance measures lead to the following data exclusions:

(i) No adult sessions were excluded for outliers. Six infant sessions were excluded because they
had >5% outliers. After exclusion, there were no significant differences in outlier volumes between
the infants and the adults (infants: mean percent outliers±SD = 0.45% ± 0.43%; adults: mean percent
outliers±SD = 0.29% ± 0.10%; $t(86) = 1.70$, $p = .092$).

(ii) We identified the corpus callosum with left-right directionality and corticospinal tract with
inferior-superior directionality in all infant and adult sessions, so no data was excluded due to this
measure.

(iii) Using babyAFQ, we identified the major bundles in the infant brain and using AFQ, we identified
these bundles in adults. In each infant and adult session we identified the arcuate (AF), posterior
arcuate (pAF), vertical occipital (VOF), inferior longitudinal (ILF), superior longitudinal (SLF),
corticospinal (CST), cingulate (CC), forceps major (FcMa), and forceps minor (FcMi), as such no
data were excluded. This confirms the quality of dMRI data and our tracking procedure as we could
identify the major white matter bundles that are known to exist from birth in all sessions.

Cytoarchitectonic Areas of VTC

We used the Rosenke Atlas[53] that contains maximum probability maps (MPMs) of 8
cytoarchitectonic regions of the human ventral visual stream, created from 10 post mortem adult
samples published in[32,124]. We used cortex based alignment in FreeSurfer[51] (https://freesurfer.net/,
mri_label2label) to map the MPM of four cytoarchitectonic areas of the ventral stream FG1, FG2,
FG3, FG4 to each cortical surface for each of the 88 sessions in the study. Two independent raters
(SD, DO) manually checked each brain to examine if the mid fusiform sulcus (MFS) was the
boundary between FG3/FG4 and FG1/FG2 as reported in prior studies[5,52]. In places where there
were disagreements between the raters, alignment was checked by KGS. The analysis confirmed
that the boundaries between the cytoarchitectonic areas aligned with the MFS (see examples in
Fig 1c) except for five infants in the right hemisphere, and one adult in the left hemisphere.

Category-Selective Regions of VTC

A functional atlas of category-selective regions of VTC was created from independent fMRI data in
28 adults from a previously published study[20]. Participants included 11 females and 17 males ages
22.1–28.6 years, (mean±SD = 24.1±1.6 years. Category-selective regions were defined in each of
the 28 individuals using the fLoc experiment containing low-level and familiarity-controlled gray

level images of items from 10 visual categories[125]. Category-selective regions were defined using
a voxel-level t-statistic contrasting each category of interest with all other categories (t > 3, no
spatial smoothing), and anatomical criteria[5,6,125]. We defined two face-selective regions on the
fusiform gyrus of each hemisphere, mFus-faces and pFus-faces using the contrast of (faces >
bodies, limbs, characters, objects, places). We defined OTS-bodies in both hemispheres as the
region on the occipital temporal sulcus selective for bodies (bodies and limbs > characters, faces,
objects, places). We defined CoS-places in both hemispheres as the region in the intersection
between the collateral sulcus (CoS) and anterior lingual sulcus (ALS) that was selective for places
(places > faces, characters, objects, bodies, limbs). Finally, we identified mOTS-words and pOTS-
words as the regions in the occipital temporal sulcus (OTS) that were selective for words
(psuedowords > faces, scenes, objects, bodies, limbs). After identifying the functional regions
within each participant we mapped the regions to the fsaverage template brain space using cortex
based alignment[51]. We then created probabilistic maps for each fROI where each vertex was the
probability of a participant having the fROI at that location (a value of 1 meant that all participants
had the region at that location and a value of .5 meant that half of participants had a value at that
location). We thresholded the probabilistic maps at .2 and then created MPMs, where in the case
that two fROIs had probabilistic values > .2 at the same vertex, the vertex was assigned to the fROI
with the highest probability (as in[126]). MPM-fROIs of word-selective regions were only found in the
left hemisphere due to their left lateralization[127,128].

To map fROIs to individual participants' cortical space for the 88 sessions of the main experiment,
we used cortex based alignment in Freesurfer[51] (https://freesurfer.net/, mri_label2label). Two
independent raters (DO and SD) then visually checked each surface to ensure that the fROIs were
aligned to the expected anatomical landmarks in each individual participant. Specifically, the raters
checked whether mFus-faces aligned to the mid fusiform sulcus (MFS) and whether CoS-places
aligned to the junction of the ALS and CoS. When there was disagreement, KGS then checked
whether the fROIs aligned to anatomical landmarks. Category-selective fROIs aligned to these
anatomical landmarks in all hemisphere except for one infant in which left hemisphere CoS-places
did not align to the junction or the ALS and CoS, and two infants in which right hemisphere mFus-
faces did not align to the MFS. We show the mapping of each fROI in each participant in
Supplementary Figs. 1-8.

We tested whether there were size differences in how the fROIs mapped to each individual brain.
To do so, tested if fROI surface area relative to brain surface varied with age:

(1) (fROI surface area)/(brain surface area) $\sim$ log10(age in days).

We found no significant differences between infants and adults on fROI surface area relative to
brain surface area (effect of age: $t(878) = 8.77 \times 10^{-6}$, p = 0.88)), reflecting that fROIs were the same
size relative to the size of the brain regardless of age.

Identifying functionally defined white matter connections

To identify the white matter connections of each fROI, we intersected the whole brain connectome
generated from dMRI and tractography in each participant. We used an open source software
package, FSuB-Extractor[101] (https://github.com/smeisler/fsub_extractor) to intersect each fROI
with the whole brain connectome of each participant and identify the white matter connections of
the fROI. In brief, the software takes in fROIs in the native space of each participant and projects
them along the surface normal into the gray-matter-white-matter interface. It then restricts the fROI
to the gray-matter-white-matter interface and then selects all streamlines that intersect with the
fROI.

Defining connectivity profiles

For each fROI, we define its connectivity profile or how it connects to the rest of the brain. To do
so, we took the white matter connections of each fROI (see above), projected their endpoints to the
cortical surface using tract density imaging (TDI) with mrTrix3[114], and calculated the distribution of
these white matter endpoints across all cortical surface vertices. We transformed the TDI output
into a distribution of endpoints by dividing the endpoint map by the total number of endpoints. This
results in an endpoint density map that sums to 1 for each fROI and participant. We then used the
Glasser Atlas[54] to quantify how the endpoints were distributed across the brain. We chose to use
the Glasser Atlas because it covers the whole cortical surface of each hemisphere, and because it
divides cortex into meaningful parcellations according to functional and anatomical criteria. For
each fROI, we define the white matter connectivity profile or the endpoint density in each region in
the Glasser Atlas. The Glasser Atlas consists of 180 regions per hemisphere. As in prior work[20], we
excluded 11 VTC regions of the Glasser Atlas to avoid quantifying looping fibers from the seed
fROI. Therefore, each connectivity profile consisted of the endpoint density across 169 regions in
the Glasser Atlas within the same hemisphere of the fROI.

Principal Components Analysis

Because our connectivity profiles are high-dimensional (169), we used principal component
analysis (PCA) to reduce the dimensionality of the connectivity profiles. We conducted PCA on the
880 (10 fROIs x 88 sessions connectivity profiles) x 169 (Glasser ROIs) endpoint connectivity
matrix.   We   used   find_curve_elbow   in   R
(https://rdrr.io/cran/pathviewr/man/find_curve_elbow.html) to find the elbow of the curve of
principal components vs variance explained and found that the first 10 principal components
explained 98% of the variance in the data. In Figure 2, we plot the first principal component (explains
38% of the variance) vs the second principal component (explains 20% of the variance). Each dot
represents a single connectivity profile in a single subject and session; The connectivity profiles
(dots) can be coded by different features (the cytoarchitectonic area, category-preference, or
participant's age).

Classification of Cytoarchitecture, Category and Age from Connectivity Profile

We used a n-way leave-one-out classifier to test if we could predict different features
(cytoarchitecture, category, or age) of a held-out connectivity profile. For each feature
(cytoarchitecture, category, age), we used multinomial logistic regression (fit on all data except for
the held out subject) to predict the feature of the held out connectivity profile (e.g., probability of
belonging to FG2, FG3, or FG4 for the cytoarchitecture classification). We used a winner-take-all
approach and assigned the connectivity profile to the label with the highest probability. We
repeated the process for each connectivity profile (leave-one-out cross-validation). We then
calculated classification accuracy by comparing the predicted classification to the ground truth.
We performed three separate classifications: one predicting cytoarchitecture (FG2/FG3/FG4), one
predicting category (face/word/body/place), and one predicting age group: (newborn/3 months/6
months/adult). Additionally, we performed separate classifications predicting category
(face/word/body/place) and age group (newborn/3 months/6 months/adults) within each
cytoarchitectonic area.

Connections to Eccentricity Bands in Early Visual Cortex

To test how each fROI connects to different eccentricity bands in early visual cortex (EVC), we
created a region of interest corresponding to EVC from the union of V1, V2, and V3 ROIs from the
Benson Atlas[129]. We then used the average retinotopic map in 21 adults from a previously published
paper[130] to divide the EVC fROI into three eccentricity bands, 0-5°, 5-10°, and 10-20°. Using cortex
based alignment in FreeSurfer (mri_label2label) we aligned each of these EVC eccentricity band
ROIs to each individual participant's native space. We then used FSuB-Extractor [101]
(https://github.com/smeisler/fsub_extractor), to identify the white matter connections between
each fROI and each eccentricity band in VTC. We divided the number of streamlines connecting to
each eccentricity band by the total number of streamlines between the fROI and EVC to estimate
the percentage of streamlines connected to each eccentricity band. To test if the percentage of
streamlines to each eccentricity band differs by cytoarchitecture, category, and age, we fit the
following model:

(2) percentage of streamlines in central 5° ~ cytoarchitecture x category x age group

After finding a significant cytoarchitecture by category interaction, we performed post-hoc tests to
determine which fROIs were driving the interaction. To do so, we used paired t-tests to test whether
the percentage of streamlines in the central 5° differed for fROIs within the same cytoarchitectonic
area for the following pairs of fROIs: pFus-faces and pOTS-words in FG2, mFus-faces and OTS-
bodies in FG4, mFus-faces and mOTS-words in FG4, and OTS-bodies and mOTS-words in FG4.

After finding a significant cytoarchitecture by age interaction, we performed post-hoc tests to test
whether the percentage of streamlines in the central 5° differed as a function of age for fROIs within

each cytoarchitectonic area using the following model separately for each cytoarchitectonic area
(FG2, FG3, FG4):

(3) percentage of streamlines in central 5° ~ log10(age) + (1|subject)

After finding a significant interaction between cytoarchitecture, category, and age we performed
post-hoc tests to test whether the percentage of streamlines to the central 5° changed over
development using the following model separately for each fROI (pFus-faces, pOTS-words, CoS-
places, mFus-faces, OTS-bodies, mOTS-words):

(4) percentage of streamlines in central 5° ~ log10(age) + (1|subject)

Quantifying development

As development is expected to asymptote across the lifespan and typically follows a logarithmic
function of age, to quantify how white matter connectivity profiles change with age, we fit a linear
model predicting endpoint density of each fROI as a function of Glasser ROI and log10(age in days)
across the brain. Age is a continuous variable and Glasser ROI is a categorical variable. As endpoint
density was normalized to sum to one across the brain, it is not appropriate to model a random
intercept for each subject.

(5) fROI endpoint density ~ log10(age in days)xGlasser ROI

After finding a significant Glasser ROI by age interaction (Supplementary Table 4), for each fROI,
we fit linear models relating endpoint density vs log10(age) separately for each Glasser ROI to
quantify the development of endpoint density within each Glasser ROI. The regression slope
quantifies the rate of the development:

(6) fROI endpoint density ~ log10(age in days).

We visualize the regression results in Fig 4B, where each Glasser ROI is colored by the slope of the
regression. Data is reported for the left hemisphere in the main text and Fig 3 because all 6 fROIs
are found consistently in the left hemisphere. Data from 4 fROIs in the right hemisphere is reported
in Supplementary Figure 53. Supplementary Tables 2-7 provide all the developmental slopes per
Glasser ROI.

Calculating the similarity between developmental slopes

To test whether development was more similar for regions within the same cytoarchitectonic area
or category-selectivity we estimated the similarity between developmental slopes of white matter
connectivity of fROI pairs using a bootstrapping procedure.

First, we calculated the developmental slopes of endpoint connectivity across the brain using a
bootstrapping procedure. For each fROI, and for each of 1000 iterations we used 75% of the data
(66 randomly selected connectivity profiles), and calculated the developmental slope within each
of the 169 Glasser ROIs using the following model:

(7) fROI endpoint density ~ log10(age in days)

Then for each pair of fROIs we calculated the pairwise correlation between the vector of 169 slopes
across the bootstrap iterations. Finally, we fit a linear model to test whether the Fisher transformed
correlation between development slopes was higher for fROIs with the same cytoarchitecture and
category-selectivity than for fROIs with different cytoarchitecture and category-selectivity. We
calculated the Fisher transform to ensure normality (as correlations are bounded between -1 and
1); category and cytoarchitecture are categorical variables (same: 1; different: 0).

(8) FisherZ(correlation) ~ category + cytoarchitecture

Data availability

The data to make the figures, tables, and statistics associated with this manuscript is available
here: https://github.com/VPNL/bbVTCwm/tree/main/data

Code availability

The code to analyze the data, compute statistics, and make the individual figure elements is
available here: https://github.com/VPNL/bbVTCwm/. The code folder contains the R code used to
generate all other figures and statistics in the figures/ and statistics/ subdirectories. The code used
to preprocess the data and perform the analyses are included in the analyses/ subdirectory. The
label files for the fROIs and the EVC ROIs are provided in the labels folder. The supplement folder
contains code to generate Supplementary Figures.
804

Author Contributions

EK: designed the analyses, wrote the code and data analysis pipelines, analyzed the data, and wrote
the manuscript. XY: participated in the design and data analysis, and collected the data. ST, BF, CT
collected the data, segmented each brain anatomy image into gray and white matter and created
cortical surface reconstructions. SF and DO: validated the alignment between fROIs,
cytoarchitectonic areas and anatomical landmarks. MG: participated in the data analyses. VSN:
participated in the design and data analysis, and collected the data. BK: designed the infant coil
used for data collection. KGS: oversaw all parts of the research: design, data analysis, and wrote
the manuscript. All authors read and gave feedback on the manuscript.

Acknowledgements

This work was funded by Stanford Wu Tsai Neurodevelopment big idea and accelerator grants, as
well as NIH grants R01EY033835 and R01EY022318 to KGS. EK was supported by the National
Science Foundation Graduate Research Fellowship (grant number DGE-1656518); MG was funded
by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation – project number
222641018 – SFB/TRR 135 TP C10), as well as by "The Adaptive Mind", funded by the Excellence
Program of the Hessian Ministry of Higher Education, Science, Research and Art.

References

1. Kanwisher, N., McDermott, J. & Chun, M. M. The fusiform face area: a module in human
extrastriate cortex specialized for face perception. *J. Neurosci.* **17**, 4302–4311 (1997).

2. Peelen, M. V. & Downing, P. E. Selectivity for the human body in the fusiform gyrus. *J.*
Neurophysiol. **93**, 603–608 (2005).

3. Cohen, L. *et al.* Language-specific tuning of visual cortex? Functional properties of the Visual
Word Form Area. *Brain* **125**, 1054–1069 (2002).

4. Epstein, R., Harris, A., Stanley, D. & Kanwisher, N. The parahippocampal place area:
recognition, navigation, or encoding? *Neuron* **23**, 115–125 (1999).

5. Weiner, K. S. *et al.* The mid-fusiform sulcus: a landmark identifying both cytoarchitectonic
and functional divisions of human ventral temporal cortex. *Neuroimage* **84**, 453–465 (2014).

6. Weiner, K. S. *et al.* Defining the most probable location of the parahippocampal place area
using cortex-based alignment and cross-validation. *Neuroimage* **170**, 373–384 (2018).

7. Levy, I., Hasson, U., Avidan, G., Hendler, T. & Malach, R. Center–periphery organization of
human object areas. *Nat. Neurosci.* **4**, 533–539 (2001).

8. Hasson, U., Levy, I., Behrmann, M., Hendler, T. & Malach, R. Eccentricity Bias as an Organizing
Principle for Human High-Order Object Areas. *Neuron* **34**, 479–490 (2002).

9. Natu, V. S. *et al.* Infants' cortex undergoes microstructural growth coupled with myelination
during development. *Commun Biol* **4**, 1191 (2021).

843 10. Cachia, A. *et al.* How interindividual differences in brain anatomy shape reading accuracy.
844   *Brain Struct. Funct.* **223**, 701–712 (2018).

845 11. Saygin, Z. M. *et al.* Anatomical connectivity patterns predict face selectivity in the fusiform
846   gyrus. *Nat. Neurosci.* **15**, 321–327 (2011).

847 12. Osher, D. E. *et al.* Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple
848   Visual Categories across Cortex. *Cereb. Cortex* **26 4**, 1668–1683 (2016).

849 13. Saygin, Z. M. *et al.* Connectivity precedes function in the development of the visual word form
850   area. *Nat. Neurosci.* **19**, 1250–1255 (2016).

851 14. Bi, Y., Wang, X. & Caramazza, A. Object Domain and Modality in the Ventral Visual Pathway.
852   *Trends Cogn. Sci.* **20**, 282–290 (2016).

853 15. Mahon, B. Z. & Caramazza, A. What drives the organization of object knowledge in the brain?
854   *Trends Cogn. Sci.* **15**, 97–103 (2011).

855 16. Kanwisher, N. Functional specificity in the human brain: a window into the functional
856   architecture of the mind. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 11163–11170 (2010).

857 17. Weiner, K. S., Yeatman, J. D. & Wandell, B. A. The posterior arcuate fasciculus and the vertical
858   occipital fasciculus. *Cortex* **97**, 274–276 (2017).

859 18. Caffarra, S., Karipidis, I. I., Yablonski, M. & Yeatman, J. D. Anatomy and physiology of word-
860   selective visual cortex: from visual features to lexical processing. *Brain Struct. Funct.* **226**,
861   3051–3065 (2021).

862 19. Lerma-Usabiaga, G., Carreiras, M. & Paz-Alonso, P. M. Converging evidence for functional and
863   structural segregation within the left ventral occipitotemporal cortex in reading. *Proceedings*
864   *of the National Academy of Sciences* **115**, E9981–E9990 (2018).

865 20. Kubota, E. *et al.* White matter connections of high-level visual areas predict cytoarchitecture
866   better than category-selectivity in childhood, but not adulthood. *Cereb. Cortex* **33**, 2485–2506
867   (2023).

868 21. Finzi, D. *et al.* Differential spatial computations in ventral and lateral face-selective regions
869   are scaffolded by structural connections. *Nat. Commun.* **12**, (2021).

870 22. Grotheer, M., Yeatman, J. & Grill-Spector, K. White matter fascicles and cortical
871   microstructure predict reading-related responses in human ventral temporal cortex.
872   *Neuroimage* **227**, 117669 (2021).

873 23. Hannagan, T., Amedi, A., Cohen, L., Dehaene-Lambertz, G. & Dehaene, S. Origins of the
874   specialization for letters and numbers in ventral occipitotemporal cortex. *Trends Cogn. Sci.*
875   **19**, 374–382 (2015).

876 24. Op de Beeck, H. P., Pillet, I. & Ritchie, J. B. Factors Determining Where Category-Selective
877   Areas Emerge in Visual Cortex. *Trends Cogn. Sci.* **23**, 784–797 (2019).

878 25. Osher, D. E. *et al.* Structural Connectivity Fingerprints Predict Cortical Selectivity for Multiple
879   Visual Categories across Cortex. *Cereb. Cortex* **26**, 1668–1683 (2016).

880 26. Li, J., Osher, D. E., Hansen, H. A. & Saygin, Z. M. Innate connectivity patterns drive the
881   development of the visual word form area. *Sci. Rep.* **10**, 18039 (2020).

882 27. Kamps, F. S., Hendrix, C. L., Brennan, P. A. & Dilks, D. D. Connectivity at the origins of domain
883   specificity in the cortical face and place networks. *Proceedings of the National Academy of*
884   *Sciences* **117**, 6163–6169 (2020).

885 28. Zeki, S. & Shipp, S. The functional logic of cortical connections. *Nature* **335**, 311–317 (1988).

29. Van Essen, D. C., Anderson, C. H. & Felleman, D. J. Information processing in the primate
visual system: an integrated systems perspective. *Science* **255**, 419–423 (1992).

30. Amunts, K. & Zilles, K. Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron*
88, 1086–1107 (2015).

31. Caspers, J. *et al.* Cytoarchitectonical analysis and probabilistic mapping of two extrastriate
areas of the human posterior fusiform gyrus. *Brain Struct. Funct.* **218**, 511–526 (2012).

32. Lorenz, S. *et al.* Two New Cytoarchitectonic Areas on the Human Mid-Fusiform Gyrus. *Cereb.*
Cortex **27**, 373–385 (2017).

33. Gomez, J., Natu, V. S., Jeska, B., Barnett, M. & Grill-Spector, K. Development differentially
sculpts receptive fields across early and high-level human visual cortex. *Nat. Commun.* **9**,
(2018).

34. Ellis, C. T. *et al.* Retinotopic organization of visual cortex in human infants. *Neuron* **109**,
2616–2626.e6 (2021).

35. Butt, O. H., Benson, N. C., Datta, R. & Aguirre, G. K. The fine-scale functional correlation of
striate cortex in sighted and blind people. *J. Neurosci.* **33**, 16209–16219 (2013).

36. Butt, O. H., Benson, N. C., Datta, R. & Aguirre, G. K. Hierarchical and homotopic correlations of
spontaneous neural activity within the visual cortex of the sighted and blind. *Front. Hum.*
Neurosci. **9**, 25 (2015).

37. Arcaro, M. J. & Livingstone, M. S. A hierarchical, retinotopic proto-organization of the primate
visual system at birth. *Elife* **6**, (2017).

38. Arcaro, M. J. & Livingstone, M. S. On the relationship between maps and domains in
inferotemporal cortex. *Nat. Rev. Neurosci.* **22**, 573–583 (2021).

39. Dudink, J., Kerr, J. L., Paterson, K. & Counsell, S. J. Connecting the developing preterm brain.
Early Hum. Dev. **84**, 777–782 (2008).

40. Grotheer, M. *et al.* White matter myelination during early infancy is linked to spatial gradients
and myelin content at birth. *Nat. Commun.* **13**, 997 (2022).

41. Grotheer, M. *et al.* Human white matter myelinates faster in utero than ex utero. *Proc. Natl.*
Acad. Sci. U. S. A. **120**, e2303491120 (2023).

42. Dubois, J., Hertz-Pannier, L., Dehaene-Lambertz, G., Cointepas, Y. & Le Bihan, D. Assessment
of the early organization and maturation of infants' cerebral white matter fiber bundles: A
feasibility study using quantitative diffusion tensor imaging and tractography. *Neuroimage*
30, 1121–1132 (2006).

43. Zöllei, L., Jaimes, C., Saliba, E., Grant, P. E. & Yendiki, A. TRActs constrained by UnderLying
INfant anatomy (TRACULInA): An automated probabilistic tractography tool with anatomical
priors for use in the newborn brain. *Neuroimage* **199**, 1–17 (2019).

44. Dimond, D. *et al.* Early childhood development of white matter fiber density and morphology.
Neuroimage **210**, 116552 (2020).

45. Perani, D. *et al.* Neural language networks at birth. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 16056–
16061 (2011).

46. Levin, N., Dumoulin, S. O., Winawer, J., Dougherty, R. F. & Wandell, B. A. Cortical maps and
white matter tracts following long period of visual deprivation and retinal image restoration.
Neuron **65**, 21–31 (2010).

47. Bauer, C. M. *et al.* Abnormal white matter tractography of visual pathways detected by high-

angular-resolution diffusion imaging (HARDI) corresponds to visual dysfunction in
cortical/cerebral visual impairment. *J. AAPOS* **18**, 398–401 (2014).

48. Ortibus, E. *et al.* Integrity of the inferior longitudinal fasciculus and impaired object
recognition in children: a diffusion tensor imaging study. *Dev. Med. Child Neurol.* **54**, 38–43
(2012).

49. Lennartsson, F., Nilsson, M., Flodmark, O. & Jacobson, L. Damage to the immature optic
radiation causes severe reduction of the retinal nerve fiber layer, resulting in predictable
visual field defects. *Invest. Ophthalmol. Vis. Sci.* **55**, 8278–8288 (2014).

50. Zufferey, P. D., Jin, F., Nakamura, H., Tettoni, L. & Innocenti, G. M. The role of pattern vision in
the development of cortico-cortical connections. *Eur. J. Neurosci.* **11**, 2669–2688 (1999).

51. Fischl, B., Sereno, M. I. & Dale, A. M. Cortical surface-based analysis. II: Inflation, flattening,
and a surface-based coordinate system. *Neuroimage* **9**, 195–207 (1999).

52. Weiner, K. S. *et al.* The Cytoarchitecture of Domain-specific Regions in Human High-level
Visual Cortex. *Cereb. Cortex* **27**, 146–161 (2016).

53. Rosenke, M. *et al.* A cross-validated cytoarchitectonic atlas of the human ventral visual
stream. *Neuroimage* **170**, 257–270 (2018).

54. Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**, 171–
178 (2016).

55. Grill-Spector, K. & Weiner, K. S. The functional architecture of the ventral temporal cortex and
its role in categorization. *Nat. Rev. Neurosci.* **15**, 536–548 (2014).

56. Weiner, K. S. *et al.* The mid-fusiform sulcus: A landmark identifying both cytoarchitectonic
and functional divisions of human ventral temporal cortex. *Neuroimage* **84**, 453–465 (2014).

57. Caspers, S. *et al.* The human inferior parietal cortex: cytoarchitectonic parcellation and
interindividual variability. *Neuroimage* **33**, 430–448 (2006).

58. Richter, M. *et al.* Cytoarchitectonic segregation of human posterior intraparietal and adjacent
parieto-occipital sulcus and its relation to visuomotor and cognitive functions. *Cereb. Cortex*
29, 1305–1327 (2019).

59. Malikovic, A. *et al.* Cytoarchitecture of the human lateral occipital cortex: mapping of two
extrastriate areas hOc4la and hOc4lp. *Brain Struct. Funct.* **221**, 1877–1897 (2016).

60. Rottschy, C. *et al.* Ventral visual cortex in humans: Cytoarchitectonic mapping of two
extrastriate areas. *Hum. Brain Mapp.* **28**, (2007).

61. Groen, I. I. A., Dekker, T. M., Knapen, T. & Silson, E. H. Visuospatial coding as ubiquitous
scaffolding for human cognition. *Trends Cogn. Sci.* **26**, 81–96 (2022).

62. Gomez, J., Barnett, M. & Grill-Spector, K. Extensive childhood experience with Pokémon
suggests eccentricity drives organization of visual cortex. *Nat Hum Behav* **3**, 611–624 (2019).

63. Silson, E. H., Chan, A. W.-Y., Reynolds, R. C., Kravitz, D. J. & Baker, C. I. A Retinotopic Basis for
the Division of High-Level Scene Processing between Lateral and Ventral Human
Occipitotemporal Cortex. *J. Neurosci.* **35**, 11921–11935 (2015).

64. Somers, D. C. & Sheremata, S. L. Attention maps in the brain. *Wiley Interdiscip. Rev. Cogn. Sci.*
4, 327–340 (2013).

65. Silver, M. A. & Kastner, S. Topographic maps in human frontal and parietal cortex. *Trends*
Cogn. Sci. **13**, 488–495 (2009).

66. Amunts, K., Mohlberg, H., Bludau, S. & Zilles, K. Julich-Brain: A 3D probabilistic atlas of the

human brain's cytoarchitecture. *Science* **369**, 988–992 (2020).

67. Sylvester, C. M. *et al.* Network-specific selectivity of functional connections in the neonatal
brain. *Cereb. Cortex* **33**, 2200–2214 (2023).

68. Huang, H. *et al.* White and gray matter development in human fetal, newborn and pediatric
brains. *Neuroimage* **33**, 27–38 (2006).

69. Thiebaut de Schotten, M. & Forkel, S. J. The emergent properties of the connected brain.
Science **378**, 505–510 (2022).

70. Dubois, J. *et al.* The early development of brain white matter: a review of imaging studies in
fetuses, newborns and infants. *Neuroscience* **276**, 48–71 (2014).

71. Harding-Forrester, S. & Feldman, D. E. Somatosensory maps. *Handb. Clin. Neurol.* **151**, 73–
102 (2018).

72. Humphries, C., Liebenthal, E. & Binder, J. R. Tonotopic organization of human auditory cortex.
Neuroimage **50**, 1202–1211 (2010).

73. Amunts, K. *et al.* Broca's region revisited: cytoarchitecture and intersubject variability. *J.*
Comp. Neurol. **412**, 319–341 (1999).

74. Lebel, C., Treit, S. & Beaulieu, C. A review of diffusion MRI of typical white matter
development from early childhood to young adulthood. *NMR Biomed.* **32**, e3778 (2019).

75. Edwards, A. D. *et al.* The Developing Human Connectome Project Neonatal Data Release.
Front. Neurosci. **16**, 886772 (2022).

76. Howell, B. R. *et al.* The UNC/UMN Baby Connectome Project (BCP): An overview of the study
design and protocol development. *Neuroimage* **185**, 891–905 (2019).

77. Walker, L. *et al.* The diffusion tensor imaging (DTI) component of the NIH MRI study of
normal brain development (PedsDTI). *Neuroimage* **124**, 1125–1130 (2016).

78. Volkow, N. D., Gordon, J. A. & Freund, M. P. The Healthy Brain and Child Development Study—
Shedding Light on Opioid Exposure, COVID-19, and Health Disparities. *JAMA Psychiatry* **78**,
471–472 (2021).

79. Eichert, N. *et al.* What is special about the human arcuate fasciculus? Lateralization,
projections, and expansion. *Cortex* **118**, 107–115 (2019).

80. Brauer, J., Anwander, A., Perani, D. & Friederici, A. D. Dorsal and ventral pathways in language
development. *Brain Lang.* **127**, 289–295 (2013).

81. Innocenti, G. M. & Frost, D. O. Effects of visual experience on the maturation of the efferent
system to the corpus callosum. *Nature* **280**, 231–234 (1979).

82. LaMantia, A. S. & Rakic, P. Axon overproduction and elimination in the corpus callosum of the
developing rhesus monkey. *J. Neurosci.* **10**, 2156–2175 (1990).

83. Price, D. J., Ferrer, J. M., Blakemore, C. & Kato, N. Postnatal development and plasticity of
corticocortical projections from area 17 to area 18 in the cat's visual cortex. *J. Neurosci.* **14**,
2747–2762 (1994).

84. Dehay, C., Kennedy, H. & Bullier, J. Characterization of transient cortical projections from
auditory, somatosensory, and motor cortices to visual areas 17, 18, and 19 in the kitten. *J.*
Comp. Neurol. **272**, 68–89 (1988).

85. Innocenti, G. M. & Price, D. J. Exuberance in the development of cortical networks. *Nat. Rev.*
Neurosci. **6**, 955–965 (2005).

86.   Webster, M. J., Bachevalier, J. & Ungerleider, L. G. Transient subcortical connections of
1015   inferior temporal areas TE and TEO in infant macaque monkeys. *J. Comp. Neurol.* **352**, 213–
1016   226 (1995).

1017   87.   Kaneko, T. *et al.* Spatial organization of occipital white matter tracts in the common
1018   marmoset. *Brain Struct. Funct.* **225**, 1313–1326 (2020).

1019   88.   Takemura, H., Pestilli, F. & Weiner, K. S. Comparative neuroanatomy: Integrating classic and
1020   modern methods to understand association fibers connecting dorsal and ventral visual
1021   cortex. *Neurosci. Res.* **146**, 1–12 (2019).

1022   89.   Takemura, H. *et al.* Occipital White Matter Tracts in Human and Macaque. *Cereb. Cortex* **27**,
1023   3346–3359 (2017).

1024   90.   Takemura, H. *et al.* A prominent vertical occipital white matter fasciculus unique to primate
1025   brains. *Curr. Biol.* (2024) doi:10.1016/j.cub.2024.06.034.

1026   91.   Zatorre, R. J., Fields, R. D. & Johansen-Berg, H. Plasticity in gray and white: neuroimaging
1027   changes in brain structure during learning. *Nat. Neurosci.* **15**, 528–536 (2012).

1028   92.   Blumenfeld-Katzir, T., Pasternak, O., Dagan, M. & Assaf, Y. Diffusion MRI of structural brain
1029   plasticity induced by a learning and memory task. *PLoS One* **6**, e20678 (2011).

1030   93.   Fields, R. D. A new mechanism of nervous system plasticity: activity-dependent myelination.
1031   *Nat. Rev. Neurosci.* **16**, 756–767 (2015).

1032   94.   Fields, R. D. Neuroscience. Change in the brain's white matter. *Science* **330**, 768–769 (2010).

1033   95.   Florence, S. L., Taub, H. B. & Kaas, J. H. Large-scale sprouting of cortical connections after
1034   peripheral injury in adult macaque monkeys. *Science* **282**, 1117–1121 (1998).

1035   96.   Jain, N., Florence, S. L., Qi, H. X. & Kaas, J. H. Growth of new brainstem connections in adult
1036   monkeys with massive sensory loss. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 5546–5550 (2000).

1037   97.   Dancause, N. *et al.* Extensive cortical rewiring after brain injury. *J. Neurosci.* **25**, 10167–
1038   10179 (2005).

1039   98.   Monje, M. Myelin Plasticity and Nervous System Function. *Annu. Rev. Neurosci.* **41**, 61–76
1040   (2018).

1041   99.   Bacmeister, C. M. *et al.* Motor learning drives dynamic patterns of intermittent myelination on
1042   learning-activated axons. *Nat. Neurosci.* **25**, 1300–1313 (2022).

1043   100.   Murty, N. A. R. *et al.* Visual experience is not necessary for the development of face-
1044   selectivity in the lateral fusiform gyrus. *Proceedings of the National Academy of Sciences* **117**,
1045   23011–23020 (2020).

1046   101.   Meisler, S. L., Kubota, E., Grotheer, M. & Gabrieli, J. A practical guide for combining functional
1047   regions of interest and white matter bundles. (2024).

1048   102.   Ewbank, M. P. *et al.* Repetition Suppression in Ventral Visual Cortex Is Diminished as a
1049   Function of Increasing Autistic Traits. *Cereb. Cortex* **25**, 3381–3393 (2015).

1050   103.   Dakin, S. & Frith, U. Vagaries of visual perception in autism. *Neuron* **48**, 497–507 (2005).

1051   104.   Conti, E. *et al.* Network over-connectivity differentiates autism spectrum disorder from other
1052   developmental disorders in toddlers: A diffusion MRI study. *Hum. Brain Mapp.* **38**, 2333–2344
1053   (2017).

1054   105.   Picci, G., Gotts, S. J. & Scherf, K. S. A theoretical rut: revisiting and critically evaluating the
1055   generalized under/over-connectivity hypothesis of autism. *Dev. Sci.* **19**, 524–549 (2016).

106. Golarai, G. *et al.* The fusiform face area is enlarged in Williams syndrome. *J. Neurosci.* **30**,
6700–6712 (2010).

107. Behrmann, M. & Avidan, G. Congenital prosopagnosia: face-blind from birth. *Trends Cogn. Sci.*
9, 180–187 (2005).

108. Duchaine, B. C. & Nakayama, K. Developmental prosopagnosia: a window to content-specific
face processing. *Curr. Opin. Neurobiol.* **16**, 166–173 (2006).

109. Yeatman, J. D. & White, A. L. Reading: The Confluence of Vision and Language. *Annu Rev Vis*
Sci **7**, 487–517 (2021).

110. Ghotra, A. *et al.* A size-adaptive 32-channel array coil for awake infant neuroimaging at 3
Tesla MRI. *Magn. Reson. Med.* **86**, 1773–1785 (2021).

111. Wang, L. *et al.* iBEAT V2.0: a multisite-applicable, deep learning-based pipeline for infant
cerebral cortical surface reconstruction. *Nat. Protoc.* **18**, 1488–1509 (2023).

112. Yushkevich, P. A. *et al.* User-guided 3D active contour segmentation of anatomical
structures: significantly improved efficiency and reliability. *Neuroimage* **31**, 1116–1128
(2006).

113. Zöllei, L., Iglesias, J. E., Ou, Y., Grant, P. E. & Fischl, B. Infant FreeSurfer: An automated
segmentation and surface extraction pipeline for T1-weighted neuroimaging data of infants
0–2 years. *Neuroimage* **218**, 116946 (2020).

114. Tournier, J.-D. *et al.* MRtrix3: A fast, flexible and open software framework for medical image
processing and visualisation. *Neuroimage* **202**, 116137 (2019).

115. Pietsch, M. *et al.* A framework for multi-component analysis of diffusion MRI data over the
neonatal period. *Neuroimage* **186**, 321–337 (2019).

116. Bastiani, M. *et al.* Automated processing pipeline for neonatal diffusion MRI in the developing
Human Connectome Project. *Neuroimage* **185**, 750–763 (2019).

117. Veraart, J. *et al.* Denoising of diffusion MRI using random matrix theory. *Neuroimage* **142**,
394–406 (2016).

118. Andersson, J. L. R., Graham, M. S., Zsoldos, E. & Sotiropoulos, S. N. Incorporating outlier
detection and replacement into a non-parametric framework for movement and distortion
correction of diffusion MR images. *Neuroimage* **141**, 556–572 (2016).

119. Tustison, N. J. *et al.* N4ITK: improved N3 bias correction. *IEEE Trans. Med. Imaging* **29**, 1310–
1320 (2010).

120. Dhollander, T., Raffelt, D. & Connelly, A. Unsupervised 3-tissue response function estimation
from single-shell or multi-shell diffusion MR data without a co-registered T1 image. in (09
2016).

121. Smith, R. E., Tournier, J.-D., Calamante, F. & Connelly, A. Anatomically-constrained
tractography: Improved diffusion MRI streamlines tractography through effective use of
anatomical information. *Neuroimage* **62**, 1924–1938 (2012).

122. Grotheer, M., Kubota, E. & Grill-Spector, K. Establishing the functional relevancy of white
matter connections in the visual system and beyond. *Brain Struct. Funct.* **227**, 1347–1356
(2022).

123. Warrington, S. *et al.* Concurrent mapping of brain ontogeny and phylogeny within a common
space: Standardized tractography and applications. *Science Advances* **8**, eabq2022 (2022).

124. Caspers, J. *et al.* Cytoarchitectonical analysis and probabilistic mapping of two extrastriate

areas of the human posterior fusiform gyrus. *Brain Struct. Funct.* **218**, 511–526 (2013).

125. Stigliani, A., Weiner, K. S. & Grill-Spector, K. Temporal Processing Capacity in High-Level
Visual Cortex Is Domain Specific. *J. Neurosci.* **35**, 12412–12424 (2015).

126. Rosenke, M., van Hoof, R., van den Hurk, J., Grill-Spector, K. & Goebel, R. A Probabilistic
Functional Atlas of Human Occipito-Temporal Visual Cortex. *Cereb. Cortex* **31**, 603–619
(2020).

127. Cai, Q., Paulignan, Y., Brysbaert, M., Ibarrola, D. & Nazir, T. A. The left ventral occipito-
temporal response to words depends on language lateralization but not on visual familiarity.
Cereb. Cortex **20**, 1153–1163 (2010).

128. Petersen, S. E., Fox, P. T., Posner, M. I., Mintun, M. A. & Raichle, M. E. Positron emission
tomographic studies of the cortical anatomy of single-word processing. *Nature* **331**, 585–
589 (1988).

129. Benson, N. C. & Winawer, J. Bayesian analysis of retinotopic maps. *Elife* **7**, (2018).

130. Finzi, D. *et al.* Differential spatial computations in ventral and lateral face-selective regions
are scaffolded by structural connections. *Nat. Commun.* **12**, 2278 (2021).

1114
1115

1116