A new tuna specimen (Genus *Auxis*) from the Duho Formation (middle Miocene) of South Korea

Dayun Suh[1], Su-Hwan Kim[*2], Gi-Soo Nam[3]

[1]University of California, Los Angeles, Los Angeles, California, 90095, USA <msdianesuh@gmail.com>

[2]School of Earth and Environmental Sciences, Seoul National University, Seoul, 08826, South Korea <shwan0@snu.ac.kr>

[3]Gongju National University of Education, Gongju, South Chungcheong Province, 32553, South Korea <nks33@hanmail.net>

[*]Corresponding author

ABSTRACT

The partially articulated caudal vertebrae of a tuna were discovered from the Duho Formation (middle Miocene) of South Korea. This specimen was assigned to the genus *Auxis* and represents the second record of fossil *Auxis* found in South Korea. We compared the vertebral morphology of the studied specimen to that of currently known species of *Auxis,* including extinct taxa, but the specimen is not assigned to a new or existing species of *Auxis* due to poor preservation and a lack of diagnostic features. The discovery of a new specimen of *Auxis* supports theories of high marine biodiversity in the East Sea (Sea of Japan) and the opening of the East Sea in the early to middle Miocene. A widely opened East Sea might have increased the abundance and diversity of large oceanic fishes such as tunas during the deposition of the Duho Formation. A taphonomic scenario of the specimen was inferred based on the lack of anal pterygiophores and the leaf imprint on the matrix. The specimen would have been exposed for

27 at least a month in a low-energy sedimentary environment at the deep-sea bottom and

28 undergone disintegration before being buried.

Introduction

The family Scombridae includes mostly epipelagic marine fishes, such as tunas, a large,

epipelagic predator (Yemmen and Gargouri, 2022). All five genera of tuna form the tribe

Thunnini. Among Thunnini, the genus *Auxis* is an epipelagic, neritic, and oceanic genus found

worldwide in tropical and subtropical oceans (Collette and Nauen, 1983). *Auxis* consumes

various fishes, crustaceans, cephalopods, and other prey and is preyed upon by large tunas,

billfishes, barracudas, sharks, and more (Collette and Nauen, 1983). *Auxis* comprises two extant

species: the frigate and bullet tunas (*Auxis thazard* and *Auxis rochei*). They exhibit significant

morphological similarities (Vieira et al., 2022) and little osteological differences. Few fossil

specimens of *Auxis* are reported, making for an unreliable fossil history of the genus.

Additionally, many fossil *Auxis* identifications have been revised throughout the decades as

some previously identified as *Auxis* underwent multiple modifications within the Scombridae

(Nam et al., 2021). With such lack of reports and many invalidations of fossil *Auxis* specimens,

currently, the only valid fossil record of *Auxis* dates back to the Miocene, reported from the

same formation as the specimen described in this paper (†*Auxis koreanus*, Nam et al., 2021).

Moreover, the detailed study of the vertebral anatomy of *Auxis* has been hindered by the

paucity of recovered specimens including both skulls and vertebrae.

An imprint of tuna vertebrae was collected from the Duho Formation, Pohang City,

South Korea, in 2020 (Fig. 1). The new specimen (GNUE322001, Gongju National University of

Education) represents the second discovery of *Auxis* from the Duho Formation of the Korean

Peninsula. Although the specimen is preserved poorly and lacks cranial elements, it possesses

diagnostic characters of the vertebrae of the genus *Auxis*. This paper describes the new

specimen and discusses the paleogeographic and palaeoecological implications of tunas in the

middle Miocene of South Korea.

Figure 1. Geologic map of the northern part of the Pohang area with Tertiary basins in South

Korea (inset), depicting the fossil locality where GNUE322001 was collected.

Geological setting

The Pohang Basin is the largest Tertiary basin in South Korea (Yoon, 1975; Fig. 1) and is a

pull-apart basin that started to form by post-volcanism subsidence at ~17 Ma (Sohn et al., 2001).

The Yeonil Group, in the Pohang Basin, is a more than 1 km thick non-marine to deep-marine

strata that consists predominantly of clastic sediments of marine origin (Sohn et al., 2001; Kim,

2008). This group comprises conglomerates and sandstones along the basin margin and

hemipelagic mudstones and sandstones towards the basin center (Sohn et al., 2001; Woo and

Khim, 2006). The Duho Formation, where the studied specimen was collected, occurs in the

uppermost part of the Yeonil Group and is about 250 m thick (Yun, 1986). A pale grey to light

brown homogeneous mudstone with intercalated sandstone is the main deposit of the Duho

Formation (Hwang et al., 1995; Kim and Paik, 2013). The Duho Formation produces a variety of

marine invertebrate and vertebrate fossils, including mollusks (Kim and Lee, 2011; Kong and

Lee, 2012), fishes (Ko, 2016; Ko and Nam, 2016; Kim et al., 2018; Nam et al., 2021; Malyshkina

et al., 2022), and whales (Lim, 2005; Lee et al., 2012). Such a diverse fossil record has produced

equally diverse paleoenvironmental interpretations during the deposition of the Duho

Formation. The paleoenvironmental interpretation of the Duho Formation ranges between

shallow marine (Kim, 1965; Yun, 1985), offshore (Lee, 1992; Yoon, 1975; Yoon, 1976), low

energy (Seong et al., 2009; Kim and Lee, 2011), hemipelagic (Chough et al., 1990; Kim and Paik,

2013), and deep-sea environments (Chough et al., 1990; Kim and Paik, 2013). Various studies

on the age of the Duho Formation additionally resulted in diverse interpretations (Kim et al.,

2018), ranging from the early Miocene based on Zircon dating (Lee et al., 2014), middle

Miocene based on paleomagnetic dating and volcanic rocks (Kim et al., 1993; Chung and Koh,

2005), and late Miocene based on dinoflagellate and radiolarian fossils (Byun and Yun, 1992;

Bak et al., 1996).

86   **Materials and methods**

The specimen was photographed using a digital camera (Sony A7R4A). Image processing

and line drawings of the specimen were done using Adobe Photoshop v 23.4.2. and Adobe

Illustrator v 26.4.1. All measurements were taken using a digital caliper.

Anatomical nomenclature

We follow the terminology of Starks (1910), which was applied to *Auxis*, to describe peculiar

vertebral structures of the studied specimen and occasionally refer to the terminology of

Romeo and Mansueti (1962) for efficient comparison between *Auxis*, *Euthynnus*, and

Katsuwonus.

Repositories and institutional abbreviation

The specimen is deposited in the Gongju National University of Education (GNUE), Gongju City,

South Korea.

Results

Systematic Paleontology

105   Order Perciformes Nelson, 2006

106   Suborder Scombroidei Nelson, 2006

107   Family Scombridae Rafinesque, 1815

108   Tribe Thunnini Starks, 1910

109   Genus *Auxis* Cuvier, 1829

Type species

Scomber rochei Risso, 1810

Figure 2. Photographs and drawings of GNUE322001. (A) Photographs of GNUE322001. Each
counterpart mold shows a lateral side of the vertebrae without the original bones. (B) Drawings
of GNUE322001. Black arrows point towards the anterior direction of the vertebrae. Dashed
lines indicate a broken edge. Grey areas indicate an unidentified leaf imprint. Vertebral
terminology follows Starks (1910). Abbreviations: ds, dorsal spine; iaz, inferior antero-
zygapophysis; ipz, inferior postero-zygapophysis; saz, superior antero-zygapophysis; spz,
superior postero-zygapophysis.

Occurrence
Duho Formation, Hwanho-dong, Buk-gu, Pohang City, North Gyeongsang Province, South Korea
(N36°3'49.10", E129°23'47.07") (Fig. 1), preserved in a massive grey mudstone in the Duho
Formation (Fig. 2).

Description
Due to the dissolution of the original bones, only the molds of the eight articulated vertebrae
are partially preserved (Fig. 2). In particular, due to the breakage of the matrix, only small
fragments of the first and last vertebrae are preserved. The centra have an amphicoelous shape,
consisting of two robust cones. Each counterpart was split along a parasagittal plane, making
both cones appear strongly connected by a wide notochordal foramen. However, the centra of
Thunnini generally are not pierced through by a notochordal foramen, and the notochord is
segmented (Starks, 1910; Graham and Dickson, 2000). The anteroposterior length and
dorsoventral height of the centrum are subequal, and the dorsal and ventral margins of the
centrum are slightly concave in lateral view.
The superior antero-zygapophysis is quite large and dorsoventrally deep, covering most
of the posterodorsal margin of the preceding centrum from the posterior margin of the
centrum to the posterior edge of its neural spine (Fig. 2). In contrast, the superior postero-
zygapophysis is weakly developed and is barely discerned in lateral view due to the overlapping
superior antero-zygapophysis of the following vertebra.

The dorsal spine originates from the centrum at mid-length, and is slightly angled

posteriorly, forming an angle of ~80-85° with the posterodorsal margin of the centrum (Fig. 2).

It slightly curves posteriorly at a third of the total length of the preserved spine from its base.

On the fourth to seventh vertebrae, the preserved inferior antero- and postero-

zygapophyses project from the centrum ventroposteriorly at an angle of ~70-80° (Fig. 2). The

length of these ventral processes of the vertebrae progressively decreases in more posterior

vertebral positions. The length of these processes in the first to third vertebrae cannot be

assessed due to incomplete preservation.

All preserved inferior antero-zygapophyses are bifurcated into an anterior and posterior

branch, and the latter tends to be longer (Fig. 2). The inferior antero-zygapophysis of the fourth

vertebra is much longer than that of the other vertebrae. It extends nearly to the level of the

posterior tip of that of the following vertebra. The inferior postero-zygapophysis almost

extends to the ventral tip of the anterior branch of the inferior antero-zygapophysis of the

following vertebra. They firmly attach to each other along the entire posterior margin of the

inferior postero-zygapophysis.

Remarks

The classification of extant _Auxis_ is based primarily on the relative body depth, corset width, the

number of gill rankers and color pattern (Collette and Aadland, 1996). The extinct _Auxis_, †_A._

koreanus, is distinguished from extant _Auxis_ by the osteological differences in skull elements

(Nam et al., 2021). Because only the caudal vertebrae are preserved in GNUE322001, the skull

is not a character available for comparison between GNUE322001 and other species of _Auxis_.

However, GNUE322001 exhibits several morphological differences in the caudal vertebrae, so

we compared its caudal vertebral morphology with that of other extant _Auxis_ species in the

following discussion.

Figure 3. Comparative diagram of the middle vertebrae of *Auxis*, *Euthynnus*, *Katsuwonus*, and

GNUE322001. (A) *Euthynnus.* (B) *Katsuwonus.* (C) *A. rochei.* (D) *A. thazard.* (E) GNUE322001

(Godsil and Byers, 1944; Yoshida, 1979; Uchida, 1981).

<div align="center">**Discussion**</div>

Anatomical comparisons

Among the Thunnini, the genera *Auxis*, *Euthynnus*, and *Katsuwonus* share a morphological

similarity in the inferior antero-zygapophysis in that it is bifurcated into anterior and posterior

branches, a unique characteristic only observed in these three genera. However, *Auxis* exhibits

ventral bifurcation only in the caudal vertebrae, whereas this character begins from the

posterior abdominal vertebrae in *Euthynnus* and *Katsuwonus* (see Godsil and Byers 1944: fig. 19;

Godsil, 1954: fig. 83; Yoshida and Nakamura, 1965: fig. 3). Furthermore, the pedicle of *Auxis*, a

median rod formed by the fusion of both sides of the inferior antero-zygapophyses below the

centrum and above the haemal canal (Kishinouye, 1923), is far longer than in *Euthynnus* and

Katsuwonus (Godsil, 1954; Fig. 3). Most significantly, *Euthynnus* and *Katsuwonus* are

characterized by the trellis pattern and inferior foramen, formed by ventral processes of the

vertebrae. In these two taxa, the posterior branch of the inferior antero-zygapophysis

(prehaemapophysis of Romeo and Mansueti, 1962) fuses with the anterior branch of the

inferior postero-zygapophysis (posthaemapophysis of Romeo and Mansueti, 1962), forming a

completely enclosed inferior foramen under the centrum (see Romeo and Mansueti, 1962: fig. 2D; Fig. 3A, B). In *Auxis*, the trellis pattern and inferior foramen are scarcely developed and only observed in the posterior most vertebrae (when present) (Kishinouye, 1923; Godsill, 1954).

Figure 4. Reconstruction of GNUE322001. The red-lined vertebrae indicate the possible position of the vertebrae of GNUE322001 in the vertebral column.

Although the cranial elements are not preserved in GNUE322001, this specimen was identified as *Auxis* primarily based on having the bifurcated inferior antero-zygapophysis with a long pedicle and no trellis pattern. Based on the vertebral column of extant *Auxis* (see Uchida, 1981: fig. 24; Jawad et al., 2013: fig. 1), it is suggested that GNUE322001 represents the anterior to the middle caudal vertebral series (Fig. 4) as indicated by the length of ventral processes, which progressively decreases throughout the vertebral series of this taxon.

There are three valid taxa within *Auxis,* including an extinct species (*A. thazard*, *A. rochei*, and †*A. koreanus*) (Collette and Aadland, 1996; Nam et al., 2021). GNUE322001 is morphologically similar to the vertebrae of *A. rochei* in that the anterior branch of the inferior antero-zygapophysis is short and does not reach the preceding inferior antero-zygapophysis (Yoshida and Nakamura, 1965; Uchida, 1981; Fig. 3C, E). In *A. thazard*, the anterior branches of the inferior antero-zygapophyses are long enough to contact the preceding inferior antero-zygapophyses (Fig. 3D). Meanwhile, the extinct taxon †*A. koreanus* is limited to comparison with GNUE322001 because only the abdominal vertebrae are preserved in the specimens of †*A.*

koreanus (Nam et al., 2021). Although †*A. koreanus* is also discovered from the Duho Formation

like GNUE322001, it is challenging to assign GNUE322001 to †*A. koreanus* based solely on their

shared occurrence within the same formation. Furthermore, the vertebrae of †*A. koreanus* and

GNUE322001 exhibit a significant size difference with lengths of approximately 0.5 and 1.5 cm

respectively (Nam et al., 2021; Fig. 2). However, there are no suitable diagnostic features to

assign GNUE322001 to a new species. Therefore, additional study and discovery of *Auxis*

specimens from the Duho Formation are necessary to determine the relationship between

GNUE322001 and †*A. koreanus,* as well as the other extant species of *Auxis.*

Paleoenvironmental perspectives

The major opening of the East Sea between 23 and 18 Ma widened the gap between the

Japanese Arc and the Korean Peninsula by 200-250 km (Sohn et al., 2001). A diverse fossil

record of large oceanic animals such as the tunas (Nam et al., 2021; GNUE322001 in this paper),

sharks (Kim et al., 2018), and whales (Lim, 2005; Lee et al., 2012) indicates high marine

biodiversity in the East Sea during this period.

Upwelling regions, although only constituting 0.1 % of the total ocean areas (Wang and

Lee, 2019), are where fishes are most abundant due to high production rates (Lalli and Parsons,

1997). One of such fishes is the tuna, which are attracted by the zones of foraging availability

created by upwelling zones (Grandperrin, 1978; Nicol et al., 2014). Additionally, based on the

record of the fossilized diatom resting spores, which indicate an upwelling activity from the

Duho Formation (Hargraves, 1979; Lee, 1993), Kim et al. (2018) hypothesized that biodiversity

of the East Sea increased due to the influence of upwelling during the deposition of the Duho

Formation. Thus, it can be concluded that upwelling activity during the middle Miocene

increased pelagic fishes' and their preys' biodiversity in the East Sea.

The absence of anal pterygiophores in GNUE322001, which in tunas are located directly

under the prehaemapophyses (Fig. 2), suggests that the specimen underwent significant

decomposition underwater. The first steps of decomposition of a fish involve the disarticulation

of the jaw and external scales as soft tissues (muscles, skins) decompose (Burrow and Turner,

2012). However, body parts are often disarticulated but still loosely connected (Burrow and

Turner, 2012). At this stage, invertebrate and vertebrate scavengers completely disconnect the

bones by feeding on the soft tissue or the bones themselves (Burrow and Turner, 2012). In

GNUE322001, the absent anal pterygiophores would have been disconnected and/or consumed

by marine scavengers, indicating that the vertebrae have been underwater for a long time.

However, the exact taphonomic time frame cannot be determined with the partially preserved

vertebrae.

246   An unidentified leaf imprint is preserved on the anterior portion of the vertebrae of

GNUE322001 (Fig. 2). Since the fine-grained matrix indicates that the specimen was buried in a

low-energy sedimentary environment at the deep-sea bottom, the leaf associated with

GNUE322001 would have traveled from shore to the depths of the sea. The leaf exhibits tears

on its edges, a characteristic of the fragmentation stage of decomposition where marine

detritivorous invertebrates feed on deposited leaves (Bridgham and Lamberti, 2009). The

decomposition rate during fragmentation varies depending on salinity; aquatic ecosystems with

lower salinity are correlated with faster decomposition (Quintino et al., 2009). Thus, decay rates

are highest in freshwater ecosystems, followed by transitional communities, and slowest in

marine ecosystems (Quintino et al., 2009). While the torn edges of the leaf imprint associated

with GNUE322001 resemble those resulting from a two-week decomposition in transitional

communities (Bridgham and Lamberti, 2009: fig. 15.2), leaves deposited in marine ecosystems

take more than twice the time to exhibit a similar amount of biomass remain (Quintino et al.,

2009: fig. 4). Thus the leaf associated with GNUE322001 would have decomposed after a month

of being exposed to water. Although the vertebrae and leaf have experienced different

decompositions in isolated conditions, based on the taphonomic time frame inferred from the

preservation of the leaf imprint, it can be estimated that the decomposition of GNUE322001

took at least a month. However, perfectly preserved leaves were also reported from the Duho

Formation (Jung and Lee, 2009); therefore, the taphonomic scenario inferred from

GNUE322001 does not represent a general depositional condition of the Duho Formation.

267   **Acknowledgments**

269 We thank Professor Yuong-Nam Lee for helping with the identification, discussion, and writing
270 of the paper.
271
Data, scripts, code, and supplementary information availability
273
274 High resolution version of Figure 2: https://doi.org/10.6084/m9.figshare.26355148
275

Funding
277

278 This research was supported by Basic Science Research Program through the National Research
279 Foundation of Korea (NRF), funded by the Ministry of Education (grant number
280 2022R1I1A2060919).

Conflict of interest disclosure
283
284 The authors declare that they comply with the PCI rule of having no financial conflicts of
285 interest in relation to the content of the article.
286

Author contributions
288
289 Dayun Suh contributed to conceptualization, formal analysis, investigation, visualization, writing
290 of the original draft, and writing of review and editing. Su-Hwan Kim contributed to
291 conceptualization, formal analysis, investigation, methodology, supervision, validation,
292 visualization, and writing of review and editing. Gi-Soo Nam contributed to resources, and
293 validation, and writing of review and editing.
294

References
296

Bak, Y., Lee, J.D., and Yun, H., 1996, Middle Miocene radiolarians from the Duho Formation in

the Pohang Basin, Korea: Journal of the Paleontological Society of Korea, v. 12, p. 225–

261.

Bridgham, S.D., and Lamberti, G.A., 2009, Ecological dynamics III: Decomposition in wetlands, *in*

Maltby, E., and Barker, T., eds., The Wetlands Handbook: Blackwell Publishing, Oxford, p.

326–345. https://doi.org/10.1002/9781444315813.ch15

Burrow, C.J., and Turner, S., 2012, Fossil fish taphonomy and the contribution of microfossils in

documenting Devonian vertebrate history, *in* Talent, J.A., ed., Earth and Life: Springer,

New York, p. 184–223. https://doi.org/10.1007/978-90-481-3428-1_8

Byun, H., and Yun, H., 1992, Miocene dinoflagellate cysts from the central part of the Pohang

Basin, Korea: Journal of the Paleontological Society of Korea, v. 8, p. 164–235.

Chough, S.K., Hwang, I.G., and Choe, M.Y., 1990, The Miocene Doumsan fan-delta, southeast

Korea: a composite fan-delta system in back-arc margin: Journal of Sedimentary

Research, v. 60, p. 445–455. https://doi.org/10.1306/212f91ba-2b24-11d7-

8648000102c1865d

Chung, C.-H., and Koh, Y.-K., 2005, Palynostratigraphic and palaeoclimatic investigations on the

Miocene deposits in the Pohang area, South Korea: Review of Palaeobotany and

Palynology, v. 135, p. 1–11. https://doi.org/10.1016/j.revpalbo.2005.02.002

Collette, B.B., and Nauen, C.E., 1983, Scombrids of the world, an annotated and illustrated

catalogue of tunas, mackerels, bonitos, and related species known to date: Food and

Agriculture Organization of the United Nations, Rome, v. 2, 137 p.

Collette, B.B., and Aadland, C.R., 1996, Revision of the frigate tunas (Scombridae, *Auxis*), with

descriptions of two new subspecies from the eastern Pacific. Fishery Bulletin: v. 94, p.

423–441.

Cuvier, G., 1829, Le Règne Animal, distribué d'après son organisation, pour servie de base à

l'histoire naturelle des animaux et d'introduction à l'anatomie comparé: Paris, 532 p.

Godsil, H.C., and Byers, R.D., 1944, A systematic study of the Pacific tunas: California

Department of Fish and Game, Fish Bulletin, v. 60, p. 1–131.

Godsil, H.C., 1954, A descriptive study of certain tuna-like fishes: California Department of Fish
and Game, Fish Bulletin, v. 97, p. 1–185.

Graham, J. B., and Dickson, K. A., 2000, The evolution of thunniform locomotion and heat
conservation in scombrid fishes: new insights based on the morphology of *Allothunnus*
fallai: Zoological Journal of the Linnean Society, v. 129, p. 419–466.
https://doi.org/10.1111/j.1096-3642.2000.tb00612.x

Grandperrin R., 1978, Influence of currents on the production of tropical seas: consequences
for fisheries: SPC Fisheries Newsletter, v. 17, p. 14–20.

Hargraves, P.E., 1979, Studies on marine plankton diatoms, IV. Morphology of *Chatoceros*
resting spores: Beihefte zur Nova Hedwigia, v. 64, p. 99–120.

Hwang, I.G., Chough, S.K., Hong, S.W., and Choe, M.Y., 1995, Controls and evolution of fan delta
systems in the Miocene Pohang Basin, SE Korea: Sedimentary Geology, v. 98, p. 147–179.
https://doi.org/10.1016/0037-0738(95)00031-3

Jawad, L.A., Al-Hassani, L., and Al-Kharusi, L.H., 2013, On the morphology of the vertebral
column of the frigate tuna, *Auxis thazard* (Lacepedea, 1800)(Family: Scombridae)
collected from the sea of Oman: Acta Musei Nationalis Pragae, v. 69, p. 101–105.
https://doi.org/10.14446/amnp.2013.101

Kim, B.K., 1965, The stratigraphic and paleontologic studies on the Tertiary (Miocene) of the
Pohang area, Korea: Seoul University Journal Science and Technology Series, v. 15, p.
32–121.

Kim, D.H., and Lee, S.-J., 2011, Fossil scallops from the Hagjeon Formation and the Duho
Formation, Pohang Basin: Journal of the Geological Society of Korea, v. 47, p. 235–244
[in Korean with English abstract].

Kim, J., and Paik, I.S., 2013, Chondrites from the Duho Formation (Miocene) in the Yeonil Group,
Pohang Basin, Korea: Occurrences and paleoenvironmental implications: Journal of the
Geological Society of Korea, v. 49, p. 407–416 [in Korean with English abstract].
https://doi.org/10.14770/jgsk.2013.49.3.407

Kim, J.H., 2008, A new species of *Acer* samaras from the Miocene Yeonil Group in the Pohang
Basin, Korea: Geosciences Journal, v. 12, p. 331–336. https://doi.org/10.1007/s12303-
008-0033-6

Kim, J.-H., Nam, K.-S., and Jeon, Y.-S., 2017, Diversity of Miocene fossil *Acer* from the Pohang
Basin, Korea: Journal of the Geological Society of Korea, v. 53, p. 387–405 [in Korean
with English abstract].

Kim, K.H., Doh, S.-J., Hwang, C.-S., and Lim, D.S., 1993, Paleomagnetic study of the Yeonil Group
in Pohang basin: Journal of Korean Society of Economic and Environmental Geology, v.
26, p. 507–518 [in Korean with English abstract].

Kim, S.-H., Park, J.-Y., and Lee, Y.-N., 2018, A tooth of *Cosmopolitodus hastalis* (Elasmobranchii:
Lamnidae) from the Duho Formation (middle Miocene) of Pohang-si, Gyeongsangbuk-do,
South Korea. Journal of the Geological Society of Korea: v. 54, p. 121–131 [in Korean
with English abstract]. https://doi.org/10.14770/jgsk.2018.54.2.121

Kishinouye, K., 1923, Contributions to the comparative study of the so-called scombrid fishes:
Journal of the College of Agriculture, Imperial University of Tokyo, v. 8, p. 293–475.

Ko, J.Y., 2016, The description of the flat fish (Pleuronectiformes) fossils from the Miocene
Duho Formation, Pohang Yeonam-dong in Korea and its implication: Journal of the
Korean Earth Science Society, v. 37, p. 1–10 [in Korean with English abstract].
https://doi.org/10.5467/jkess.2016.37.1.1

Ko, J.Y., and Nam, K.-S., 2016, *Pleuronichthys* sp. fossils (Pleuronectidae) from the Duho
Formation, Pohang Uhyeon-dong in Korea: Journal of the Korean Earth Science Society,
v. 37, p. 133–142 [in Korean with English abstract].
https://doi.org/10.5467/jkess.2016.37.3.133

Kong, D.Y., and Lee, S.J., 2012, Fossil scaphopods from the Hagjeon Formation and the Duho
Formation, the Cenozoic Pohang Basin, Korea: Korean Journal of Cultural Heritage
Studies, v. 45, p. 218–231 [in Korean].

Lalli, C. and Parsons, T., 1997, Biological oceanography: an introduction: Oxford, Elsevier
Butterworth-Heinemann, 314 p.

Lee, T.-H., Yi, K., Cheong, C.-S., Jeong, Y.-J., Kim, N., and Kim, M.-J., 2014, SHRIMP U-Pb zircon
geochronology and geochemistry of drill cores from the Pohang Basin: Journal of the
Korean Earth Science Society, v. 23, p. 167–185 [in Korean with English abstract].
https://doi.org/10.7854/jpsk.2014.23.3.167

Lee, Y.G., 1992, Paleontological study of the Tertiary molluscan fauna in Korea: Science Reports
of the Institution of Geosciences University of Tsukuba, Section B (geological science), v.
13, p. 15–125.

Lee, Y.G., 1993, The marine diatom genus *Chaetoceros* Ehrenberg flora and some resting spores
of the Neogene Yeonil Group in the Pohang Basin, Korea: Journal of the Paleontological
Society of Korea, v. 9, p. 24–52.

Lee, Y.-N., Ichishima, H., and Choi, D.K., 2012, First record of a platanistoid cetacean from the
middle Miocene of South Korea: Journal of Vertebrate Paleontology, v. 32, p. 231–234.
https://doi.org/10.1080/02724634.2012.626005

Lim, J.-D., 2005, The first dolphin fossil from the Miocene of Korea: Current Science, v. 89, p.
939–940.

Malyshkina, T.P., Ward, D.J., Nazarkin, M.V., Nam, G.-S., Kwon, S.-H., Lee, J.-H., Kim, T.-W., Kim,
D.-K., and Baek, D.-S., 2022, Miocene Elasmobranchii from the Duho Formation, South
Korea: Historical Biology. https://doi.org/10.1080/08912963.2022.21 10870

Nam, G.-S., Nazarkin, M.V., and Bannikov, A.F., 2021, First discovery of the genus *Auxis*
(Actinopterygii: Scombridae) in the Neogene of South Korea: Bollettino della Societa
Paleontologica Italiana, v. 60, p. 61–67. https://doi.org/10.4435/BSPI.2021.05

Nelson, J.S., 2006, Fishes of the World (fourth edition): Hoboken, John Wiley and Sons, 601 p.

Nicol, S., Menkes, C., Jurado-Molina, J., Lehodey, P., Usu, T., Kumasi, B., Muller, B., Bell, J.,
Tremblay-Boyer, L., and Briand, K., 2014, Oceanographic characterisation of the Pacific
Ocean and the potential impact of climate variability on tuna stocks and tuna fisheries:
SPC Fisheries Newsletter, v. 145, p. 37–48.

Quintino, V., Sangiorgio, F., Ricardo, F., Mamede, R., Pires, A., Freitas, R., Rodrigues, A.M., and
Basset, A., 2009, In situ experimental study of reed leaf decomposition along a full

408   salinity gradient: Estuarine, Coastal and Shelf Science, v. 85, p. 497–506.

409   https://doi.org/10.1016/j.ecss.2009.09.016

410   Rafinesque, C.S., 1815, Analyse de la nature, ou tableau de l'univers et des corps organisés:

411   Palermo, 224 p. https://doi.org/10.5962/bhl.title.106607

412   Risso, A., 1810, Ichthyologie de Nice ou histoire naturelle des poissons du département des

413   Alpes Maritimes: Paris, F. Schoell, 388 p. https://doi.org/10.5962/bhl.title.7052

414   Romeo, J., and Mansueti, A.J., 1962, Little tuna, *Euthynnus alletteratus*, in northern Chesapeake

415   Bay, Maryland, with an illustration of its skeleton: Chesapeake Science, v. 3, p. 257–263.

416   https://doi.org/10.2307/1350633

417   Seong, M.N., Kong, D.-Y., Lee, B.J., and Lee, S.-J., 2009, Cenozoic brittle stars (Ophiuroidea) from

418   the Hagjeon Formation and the Duho Formation, Pohang Basin, Korea: Journal of

419   Korean Society of Economic and Environmental Geology, v. 42, p. 367–376 [in Korean

420   with English abstract].

421   Sohn, Y.K., Rhee, C.W., and Shon, H., 2001, Revised stratigraphy and reinterpretation of the

422   Miocene Pohang basinfill, SE Korea: sequence development in response to tectonism

423   and eustasy in a back-arc basin margin: Sedimentary Geology, v. 143, p. 265–285.

424   https://doi.org/10.1016/s0037-0738(01)00100-2

425   Starks, E.C., 1910, The osteology and mutual relationships of the fishes belonging to the family

426   Scombridae: Journal of Morphology, v. 21, p. 77–99.

427   https://doi.org/10.1002/jmor.1050210103

428   Uchida, R.N., 1981, Synopsis of biological data on frigate tuna, *Auxis thazard*, and bullet tuna, *A.*

429   *rochei*: Silver Spring, US Department of Commerce, National Oceanic and Atmospheric

430   Administration, National Marine Fisheries Service, 63 p.

431   https://doi.org/10.5962/bhl.title.63109

432   Vieira, J.M., Costa, P.A., Braga, A.C., São-Clemente, R.R., Ferreira, C.E., and Silva, J.P., 2022, Age,

433   growth and maturity of frigate tuna (*Auxis thazard* Lacepède, 1800) in the Southeastern

434   Brazilian coast: Aquatic Living Resources, v. 35, p. 11.

435   https://doi.org/10.1051/alr/2022010

Wang, Y.-C., and Lee, M.-A., 2019, Composition and distribution of fish larvae surrounding the
upwelling zone in the waters of northeastern Taiwan in summer: Journal of Marine
Science and Technology, v. 27, p. 8.

Woo, K.S., and Khim, B.-K., 2006, Stable oxygen and carbon isotopes of carbonate concretions
of the Miocene Yeonil Group in the Pohang Basin, Korea: Types of concretions and
formation condition: Sedimentary Geology, v. 183, p. 15–30.
https://doi.org/10.1016/j.sedgeo.2005.09.005

Yemmen, C., and Gargouri, M., 2022, Potential hazards associated with the consumption of
Scombridae fish: Infection and toxicity from raw material and processing: Journal of
Applied Microbiology, v. 132, p. 4077–4096. https://doi.org/10.1111/jam.15499

Yoon, S., 1975, Geology and paleontology of the Tertiary Pohang Basin, Pohang District, Korea
Part 1, Geology: Journal of the Geological Society of Korea, v. 11, p. 187–214.

Yoon, S., 1976, Geology and paleontology of the Tertiary Pohang Basin, Pohang District, Korea
Part 2, Paleontology (Mollusca): Journal of the Geological Society of Korea, v. 12, p. 1–22.

Yoshida, H.O., and Nakamura, E.L., 1965, Notes on schooling behavior, spawning, and
morphology of Hawaiian frigate mackerels, *Auxis thazard* and *Auxis rochei*: Copeia, v. 1,
p. 111–114. https://doi.org/10.2307/1441254

Yoshida, H.O., 1979, Synopsis of biological data on tunas of the genus *Euthynnus*: Silver Spring,
US Department of Commerce, National Oceanic and Atmospheric Administration,
National Marine Fisheries Service, 9 p.

Yun, H., 1985, Some fossil Squillidae (Stomatopoda) from the Pohang Tetiary Basin, Korea:
Journal of the Paleontological Society of Korea, v. 1, p. 19–31.

Yun, H., 1986, Emended stratigraphy of the Miocene formations in the Pohang Basin, Part I:
Journal of the Paleontological Society of Korea, v. 2, p. 54–69.