A new tuna specimen (Genus *Auxis*) from the Duho Formation (Miocene) of South Korea

Dayun Suh[1], Su-Hwan Kim[*2], Gi-Soo Nam[3]

[1]University of California, Los Angeles, Los Angeles, California, 90095, USA <msdianesuh@gmail.com>

[2]School of Earth and Environmental Sciences, Seoul National University, Seoul, 08826, South Korea <shwan0@snu.ac.kr>

[3]Gongju National University of Education, Gongju, South Chungcheong Province, 32553, South Korea <nks33@hanmail.net>

[*]Corresponding author

ABSTRACT

A partially preserved caudal vertebrae imprint of a tuna was discovered from the Duho Formation (Miocene) of South Korea. This specimen was assigned to the genus *Auxis* and represents the second record of fossil *Auxis* found in South Korea and in the world. We compared the vertebral morphology of the studied specimen to that of currently known species of *Auxis*, including extinct taxa. However, the specimen could not be assigned to any extant or new species of *Auxis* due to anatomical differences and a lack of comparability. The discovery of a new specimen of *Auxis* aligns with theories of high marine biodiversity in the East Sea (Sea of Japan) and its opening in the Early to Middle Miocene. A widely opened East Sea and upwelling activities might have increased the abundance and diversity of large oceanic fishes such as tunas during the deposition of the Duho Formation. The specimen supports paleoenvironmental interpretations of the Duho Formation as pelagic and subtropical. A taphonomic scenario of the specimen was inferred based on the lack of anal pterygiophores and the leaf imprint on the matrix. The specimen would have been exposed for at least a month

28 in a low-energy sedimentary environment at the deep-sea bottom and would have undergone

29 disintegration before being buried.

Introduction

The family Scombridae includes mostly epipelagic marine fishes, such as tunas, a large,

epipelagic predator (Collette and Nauen, 1983). The four genera of tuna, *Auxis*, *Euthynnus*,

Katsuwonus, and *Thunnus*, form the tribe Thunnini. Among the Thunnini, the genus *Auxis* is an

epipelagic, neritic, and oceanic genus found worldwide in tropical and subtropical oceans

(Collette and Nauen, 1983). *Auxis* consumes various fishes, crustaceans, cephalopods, and

other prey and is preyed upon by large tunas, billfishes, barracudas, sharks, and more (Collette

and Nauen, 1983). *Auxis* comprises two extant species: the frigate and bullet tunas (*Auxis*

thazard and *Auxis rochei*). They exhibit significant morphological similarities (Vieira et al., 2022)

and little osteological differences. The fossil history of *Auxis* is incomplete due to a lack of fossil

records and many invalidations of fossil specimens. The fossilization potential is extremely low

for taxa in deep-sea and pelagic environments with a 15% and 3% possibility of fossilization

respectively due to factors like the fragile skeleton and high environmental stress (Shaw et al.,

2020). As a result, *Auxis* fossils are extremely rare, with very few reported specimens.

Meanwhile, several fossil specimens previously identified as *Auxis* have undergone multiple

taxonomic changes within the Scombridae (Nam et al., 2021). Several cases of *Auxis*

misidentification have been noted in the literature. Five scombrid species from the Miocene in

Europe were initially identified as *Auxis* by Kramberger-Gorjanović (1882) and Gorjanović-

Kramberger (1895); however, Nam et al. (2021) later disputed these identifications based on

differences in traits such as vertebral count, lack of the haemal arch, and gill cover and

dentition size. Additionally, Woodward (1901) reclassified a scombrid fossil originally described

by Agassiz (1833–1844) as *Auxis*, representing what was considered the earliest record of *Auxis*

from the Eocene. Bannikov and Sorbini (1984) later corrected this identification, noting

discrepancies in vertebral number and shape of haemal arches. Another specimen, a Middle to

Upper Miocene scombrid from Russia identified as *Auxis* by Bogatshov (1933), was reclassified

by Bannikov (1985) due to an inconsistent vertebral count. With these invalidations, the only

currently accepted fossil record of *Auxis* dates back to the Miocene and is reported from the

same formation as the specimen described in this paper (†*Auxis koreanus*, Nam et al., 2021).

Moreover, the detailed study of the vertebral anatomy of *Auxis* has been hindered by the

paucity of recovered specimens including both skulls and vertebrae.

An imprint of tuna vertebrae was collected from the Duho Formation, Pohang City,

South Korea, in 2020 (Fig. 1). The new specimen (GNUE322001, Gongju National University of

Education) represents the second discovery of *Auxis* from the Duho Formation of the Korean

Peninsula and the second valid *Auxis* specimen in the world. Although the specimen is

preserved poorly and lacks cranial elements, it possesses diagnostic characters of the vertebrae

of the genus *Auxis*: the bifurcated inferior antero-zygapophysis with a long pedicle and no trellis.

This paper describes this new tuna specimen and discusses the palaeoecological implications of

the presence of tunas in the Miocene of South Korea.

Figure 1. Geologic map of the northern part of the Pohang area with Cenozoic basins in South

Korea (inset), depicting the fossil locality where GNUE322001 was collected.

Geological setting

The Pohang Basin is the largest Cenozoic basin in South Korea (Yoon, 1975; Fig. 1) and is

a pull-apart basin that started to form by post-volcanism subsidence at ~17 Ma (Sohn et al.,

2001). The Yeonil Group, in the Pohang Basin, includes more than 1 km thick non-marine to

deep-marine strata that are characterized predominantly of clastic sediments of marine origin

(Sohn et al., 2001; Kim, 2008). This group comprises conglomerates and sandstones along the

basin margin and hemipelagic mudstones and sandstones towards the basin center (Sohn et al.,

2001; Woo and Khim, 2006). The Yeonil Group consists of the Duho, Idong, Heunghae, Hagjeon,

and Chunbuk Formations (Yoon, 1975; Yun, 1986; Fig. 1). The Duho Formation, where the

studied specimen was collected, occurs in the uppermost part of the Yeonil Group and is about

250 m thick (Yun, 1986). A pale grey to light brown homogeneous mudstone with intercalated

sandstone is the main deposit of the Duho Formation (Hwang et al., 1995; Kim and Paik, 2013).

The Duho Formation produces a variety of marine invertebrate and vertebrate fossils, including

mollusks (Kim and Lee, 2011; Kong and Lee, 2012), fishes (Ko, 2016; Ko and Nam, 2016; Kim et

al., 2018; Nam et al., 2019; Nam et al., 2021; Nam and Nazarkin, 2022; Nazarkin and Nam, 2022;

Malyshkina et al., 2023; tab. 1), and whales (Lim, 2005; Lee et al., 2012). Such a diverse fossil

record has produced equally diverse paleoenvironmental interpretations of the depositional

environment of the Duho Formation. The paleoenvironmental interpretation of the Duho

Formation ranges between shallow marine (Kim, 1965; Yun, 1985), offshore (Lee, 1992; Yoon,

1975; Yoon, 1976), low energy (Seong et al., 2009; Kim and Lee, 2011), hemipelagic (Chough et

al., 1990; Kim and Paik, 2013), and deep-sea environments (Chough et al., 1990; Kim and Paik,

2013). Various studies on the age of the Duho Formation additionally resulted in diverse

interpretations (Kim et al., 2018), ranging from the Early Miocene based on Zircon dating (Lee

et al., 2014), Middle Miocene based on paleomagnetic dating and volcanic rocks (Kim et al.,

1993; Chung and Koh, 2005), and Late Miocene based on dinoflagellate and radiolarian fossils

(Byun and Yun, 1992; Bak et al., 1996).

Table 1. List of fish taxa from the Duho Formation.

Taxa		References
Actinopterygii	Pleuronectiformes	Ko (2016)
	Pleuronichthys sp.	Ko and Nam (2016)
	†*Vinciguerria orientalis*	Nam et al. (2019)
	†*Stenobrachius sangsunii*	Nam and Nazarkin (2022)
	Vinciguerria sp.	Nazarkin and Nam (2022)
	†*Auxis koreanus*	Nam *et al.* (2021)
Elasmobranchii	†*Carcharodon hastalis*	Kim *et al.* (2018) Malyshkina *et al.* (2023)
	Hexanchus griseus	Malyshkina *et al.* (2023)
	†*Dalatias orientalis*	Malyshkina *et al.* (2023)
	Mitsukurina owstoni	Malyshkina *et al.* (2023)
	†*Otodus megalodon*	Malyshkina *et al.* (2023)
	†*Parotodus benedenii*	Malyshkina *et al.* (2023)
	†*'Isurus' planus*	Malyshkina *et al.* (2023)
	Isurus sp.	Malyshkina *et al.* (2023)
	†*Cetorhinus huddlestoni,*	Malyshkina *et al.* (2023)
	Carcharhinus aff. *C. plumbeus*	Malyshkina *et al.* (2023)
	Carcharhinus aff. *C. amblyrhynchos*	Malyshkina *et al.* (2023)
	Carcharhinus aff. *C. altimus*	Malyshkina *et al.* (2023)
	†*Galeocerdo aduncus*	Malyshkina *et al.* (2023)

105   **Materials and methods**

The specimen GNUE322001, a partially preserved caudal tuna vertebrae imprint, is

housed in the Gongju National University of Education (GNUE), Gongju City, South Korea. The

specimen was photographed using a digital camera (Sony A7R4A). Image processing and line

drawings of the specimen were done using Adobe Photoshop v 23.4.2. and Adobe Illustrator v

26.4.1. All measurements were taken using a digital caliper.

Anatomical nomenclature

We follow the terminology of Starks (1910), which was applied to *Auxis*, to describe

peculiar vertebral structures of the studied specimen and occasionally refer to the terminology

of Romeo and Mansueti (1962) for efficient comparison between *Auxis*, *Euthynnus*, and

Katsuwonus of the tribe Thunnini.

Repositories and institutional abbreviation

The specimen is deposited in the Gongju National University of Education (GNUE),

Gongju City, South Korea.

123   **Results**

125 ***Systematic Paleontology***

126   Order Perciformes Nelson, 2006

127   Suborder Scombroidei Nelson, 2006

128   Family Scombridae Rafinesque, 1815

129   Tribe Thunnini Starks, 1910

130   Genus *Auxis* Cuvier, 1829

131 ***Type species***

132   *Scomber rochei* Risso, 1810

Figure 2. Photographs and drawings of GNUE322001. (A) Photographs of GNUE322001. Each counterpart mold shows a lateral side of the vertebrae without the original bones. (B) Drawings of GNUE322001. Black arrows point towards the anterior direction of the vertebrae. Dashed lines indicate a broken edge. Grey areas indicate an unidentified leaf imprint. Vertebral terminology follows Starks (1910). Abbreviations: ds, dorsal spine; iaz, inferior antero-zygapophysis; ipz, inferior postero-zygapophysis; saz, superior antero-zygapophysis; spz, superior postero-zygapophysis.

Occurrence

Duho Formation, Hwanho-dong, Buk-gu, Pohang City, North Gyeongsang Province, South Korea (N36°3'49.10", E129°23'47.07") (Fig. 1), preserved in a massive grey mudstone in the Duho Formation (Fig. 2).

Description

Due to the dissolution of the original bones, only the molds of the eight articulated

caudal vertebrae are partially preserved (Fig. 2). In particular, due to the breakage of the matrix,

only small fragments of the first and last vertebrae are preserved. The centra have an

amphicoelous shape, consisting of two robust cones, and both cones are connected by what

appears as a wide foramen. In all Thunnini and certain fish vertebrae, the centrum is not

pierced thus lacks a notochordal foramen (Starks, 1910; Graham and Dickson, 2000). Therefore,

the apparent wide foramen in GNUE322001 is interpreted as the result of the split of the

specimen along a parasagittal plane, and does not represent a real anatomical trait. The

anteroposterior length and dorsoventral height of the centrum are subequal, and the dorsal

and ventral margins of the centrum are slightly concave in lateral view.

The superior antero-zygapophysis is quite large and dorsoventrally deep, covering most

of the posterodorsal margin of the preceding centrum from the posterior margin of the

centrum to the posterior edge of the base of its dorsal spine (Fig. 2). In contrast, the superior

postero-zygapophysis is weakly developed and is barely discernable in lateral view due to the

overlapping superior antero-zygapophysis of the following vertebra.

The dorsal spine originates from the centrum at mid-length, and is slightly angled

posteriorly, forming an angle of ~80-85° with the posterodorsal margin of the centrum (Fig. 2).

It slightly curves posteriorly at a third of the total length of the preserved spine from its base.

On the fourth to seventh vertebrae, the preserved inferior antero- and postero-

zygapophyses project from the centrum ventroposteriorly at an angle of ~70-80° (Fig. 2). The

length of these ventral processes of the vertebrae progressively decreases in more posterior

vertebral positions. The length of these processes in the first to third vertebrae cannot be

assessed due to incomplete preservation.

All preserved inferior antero-zygapophyses are bifurcated into anterior and posterior

branches, and the latter tends to be longer (Fig. 2). The inferior antero-zygapophysis of the

fourth vertebra is much longer than that of the other vertebrae. It extends nearly to the level of

the posterior tip of that of the following vertebra. The inferior postero-zygapophysis extends

almost to the ventral tip of the anterior branch of the inferior antero-zygapophysis of the

following vertebra. They firmly attach to each other along the entire posterior margin of the

inferior postero-zygapophysis.

Remarks

We classify GNUE322001 as anterior to the middle caudal vertebrae based on the

progressively decreasing lengths of the ventral processes, a pattern also observed in the caudal

vertebrae of extant *Auxis* (see Uchida, 1981: fig. 24; Jawad et al., 2013: fig. 1; Fig. 4).

The classification of extant *Auxis* is based primarily on the relative body depth, corset

width, the number of gill rankers and color pattern (Collette and Aadland, 1996). The extinct

Auxis, †*A. koreanus*, is distinguished from extant *Auxis* by the osteological differences in skull

elements (Nam et al., 2021). Because only the caudal vertebrae are preserved in GNUE322001,

the skull is not available for comparison with other species of *Auxis*. However, GNUE322001

exhibits several distinct morphological features in the caudal vertebrae, which we compare

with that of other extant *Auxis* species in the following discussion.

Figure 3. Comparative diagram of the anterior caudal vertebrae of *Auxis*, *Euthynnus*,

Katsuwonus, and GNUE322001. (A) *Euthynnus*. (B) *Katsuwonus*. (C) *A. rochei*. (D) *A. thazard*. (E)

GNUE322001 (Godsil and Byers, 1944; Yoshida, 1979; Uchida, 1981). Vertebral terminology

follows Starks (1910). Abbreviations: iaz/a, anterior branch of inferior antero-zygapophysis;

iaz/p, posterior branch of inferior antero-zygapophysis; ipz/a, anterior branch of inferior

postero-zygapophysis; ipz/p, posterior branch of inferior postero-zygapophysis; if, inferior

foramen. Note that in *Euthynnus* (A) and *Katsuwonus* (B), the posterior branch of inferior

antero-zygapophysis and the anterior branch of inferior postero-zygapophysis are fused,

forming the inferior foramen and trellis. Size not to scale.

203   **Discussion**

Anatomical comparisons

Among the Thunnini, the genera *Auxis*, *Euthynnus*, and *Katsuwonus* share a

morphological similarity in the inferior antero-zygapophysis in that it is bifurcated into anterior

and posterior branches, a unique characteristic only observed in these three genera. However,

Auxis exhibits ventral bifurcation only in the caudal vertebrae, whereas this trait is also found in

the posterior abdominal vertebrae in *Euthynnus* and *Katsuwonus* (see Godsil and Byers 1944:

fig. 19; Godsil, 1954: fig. 83; Yoshida and Nakamura, 1965: fig. 3). Furthermore, the pedicle of

Auxis, a median rod formed by the fusion of both sides of the inferior antero-zygapophyses

below the centrum and above the haemal canal (Kishinouye, 1923), is far longer than that of

Euthynnus and *Katsuwonus* (Godsil, 1954; Fig. 3).

Most significantly, the anterior caudal vertebrae in *Euthynnus* and *Katsuwonus* are

characterized by the presence of an inferior foramen and trellis pattern. The inferior foramen is

a hole-like structure fenestra created by the complete fusion of the posterior branch of the

inferior antero-zygapophysis (prehaemapophysis of Romeo and Mansueti, 1962) and the

anterior branch of the inferior postero-zygapophysis (posthaemapophysis of Romeo and

Mansueti, 1962) under the centrum (see Romeo and Mansueti, 1962: fig. 2D; Fig. 3A, B). The

trellis pattern is formed by the continuous repetition of the inferior foramen across the

vertebrae (Fig. 3A, B). In *Auxis*, the inferior foramen and trellis pattern are scarcely developed,

and even when present, only occur in the posterior caudal vertebrae, unlike in *Euthynnus* and

Katsuwonus (Kishinouye, 1923; Godsill, 1954; Fig. 3).

Figure 4. Reconstruction of GNUE322001. The red-lined vertebrae indicate the possible position

of the vertebrae of GNUE322001 in the vertebral column.

Although the cranial elements are not preserved in GNUE322001, this specimen was

identified as *Auxis* primarily based on having a bifurcated inferior antero-zygapophysis with a

long pedicle and no trellis. Based on the vertebral column of extant *Auxis* (see Uchida, 1981: fig.

24; Jawad et al., 2013: fig. 1), it is suggested that GNUE322001 represents the anterior to the

middle caudal vertebral series (Fig. 4) as indicated by the lengths of ventral processes that

progressively decrease throughout the vertebral series in this taxon.

There are three valid taxa within *Auxis,* including an extinct species (*A. thazard*, *A. rochei*,

and †*A. koreanus*) (Collette and Aadland, 1996; Nam et al., 2021). GNUE322001 is

morphologically similar to the vertebrae of *A. rochei* in that the anterior branch of the inferior

antero-zygapophysis is short and does not reach the preceding inferior antero-zygapophysis

(Yoshida and Nakamura, 1965; Uchida, 1981; Fig. 3C, E). In *A. thazard*, the anterior branches of

the inferior antero-zygapophyses are long enough to contact the preceding inferior antero-

zygapophyses (Fig. 3D). Comparisons with the extinct taxon †*A. koreanus* are not possible

because the caudal vertebrae are unknown in this species, as they are not preserved in its only

known specimens (Nam et al., 2021). Although †*A. koreanus* was also discovered in the Duho

Formation, as was GNUE322001, it is challenging to assign GNUE322001 to †*A. koreanus* based

solely on their shared occurrence within the same formation. Furthermore, the centra of †*A.*

koreanus and GNUE322001 differ significantly in size, with lengths of approximately 0.5 and 1.5

cm respectively (Nam et al., 2021; Fig. 2). Therefore, additional study and discovery of *Auxis*

specimens from the Duho Formation are necessary to determine the relationship between

GNUE322001 and †*A. koreanus,* as well as other extant species of *Auxis.*

Paleoenvironmental perspectives

The major opening of the East Sea that occurred between 23 and 18 Ma widened the

gap between the Japanese Arc and the Korean Peninsula by 200-250 km (Sohn et al., 2001). The

opening of the East Sea would have facilitated the creation of a variety of marine environments.

This increased environmental complexity likely provided habitats that were able to support a

wider range of marine species. A diverse fossil record of large oceanic animals in the East Sea

during this period such as the tunas (Nam et al., 2021; GNUE322001 in this paper), sharks (Kim

et al., 2018), and whales (Lim, 2005; Lee et al., 2012) aligns with this theory.

Upwelling regions, although only constituting 0.1% of the total ocean areas (Wang and

Lee, 2019), are where fishes are most abundant due to high production rates (Lalli and Parsons,

1997). One of such fishes is the tuna, which are attracted by the zones of foraging availability

created by upwelling zones (Grandperrin, 1978; Nicol et al., 2014). Additionally, based on the

record of the fossilized diatom resting spores, which indicate an upwelling activity in the Duho

Formation (Hargraves, 1979; Lee, 1993), Kim et al. (2018) hypothesized that the biodiversity of

the East Sea increased due to the influence of upwelling during the deposition of the Duho

Formation. Thus, it can be concluded that upwelling activity during the Miocene increased

pelagic fishes' and their preys' biodiversity in the East Sea.

Today, tunas inhabit tropical and subtropical epipelagic ocean ecosystems (Collette and

Nauen, 1983). During warm seasons, tunas move into coastal areas, whereas in colder seasons,

they exclusively occupy deeper offshore waters (Kishinouye, 1923). In deep epipelagic

environments, tunas specifically inhabit deep rocky banks and forage in the deep scattering

layer (Kishinouye, 1923; Graham and Dickson, 2004). Graham and Dickson (2004) argue that

most oceanic physical and biological features observed in ocean ecosystems today have been in

place since the Miocene. This suggests that the East Sea during the Miocene likely supported a

pelagic and subtropical environment, as tuna, which inhabit such ecosystems, were present.

This interpretation is supported by researchers who consider the Duho Formation to represent

a deep-sea and subtropical environment, among the various interpretations of its

environmental context (see the Geological Setting section). Recent discoveries of pelagic sharks

(Malyshkina et al., 2023), lightfishes (Nam et al., 2019; Nazarkin and Nam, 2022), and

lanternfish (Nam and Nazarkin, 2022) provide evidence that the Duho Formation represents a

deep-sea environment. Moreover, based on discoveries of certain plant fossils, some

researchers interpret the Duho Formation to have been a subtropical environment (Kim, 2008,

2010; Kim et al., 2009, 2017; Jung and Lee, 2009). The discovery of GNUE322001 supports both

of these interpretations.

Taphonomic interpretations

The absence of anal pterygiophores in GNUE322001, which in tunas are located directly

under the prehaemapophyses (Fig. 2), suggests that the specimen underwent significant

decomposition underwater. The first steps of decomposition of a fish involve the disarticulation

of the jaw and external scales as soft tissues (muscles, skins) decompose (Burrow and Turner,

2012). However, body parts are often disarticulated but still loosely connected (Burrow and

Turner, 2012). At this stage, invertebrate and vertebrate scavengers completely disconnect the

bones by feeding on the soft tissue or the bones themselves (Burrow and Turner, 2012). In

GNUE322001, the absent anal pterygiophores would have been disconnected and/or consumed

by marine scavengers, indicating that the vertebrae would have been underwater for a long

time. However, the exact taphonomic time frame cannot be determined with the partially

preserved vertebrae.

An unidentified leaf imprint is preserved on the anterior portion of the vertebrae of

GNUE322001 (Fig. 2). Since the fine-grained matrix indicates that the specimen was buried in a

low-energy sedimentary environment at the deep-sea bottom, the leaf associated with

GNUE322001 would have traveled from shore to the depths of the sea. The leaf exhibits tears

on its edges, a characteristic of the fragmentation stage of decomposition where marine

detritivorous invertebrates feed on deposited leaves (Bridgham and Lamberti, 2009). The

decomposition rate during fragmentation varies depending on salinity; aquatic ecosystems with

lower salinity are correlated with faster decomposition (Quintino et al., 2009). Thus, decay rates

are greatest in freshwater ecosystems, followed by transitional communities, and slowest in

marine ecosystems (Quintino et al., 2009). While the torn edges of the leaf imprint associated

with GNUE322001 resemble those of leaves that underwent a two-week decomposition in

transitional communities (Bridgham and Lamberti, 2009: fig. 15.2), leaves deposited in marine

ecosystems take more than twice the time to exhibit a similar amount of biomass remain

(Quintino et al., 2009: fig. 4). Thus the leaf associated with GNUE322001 would have

decomposed after a month of being exposed to water. Although the vertebrae and leaf have

experienced different decompositions in isolated conditions, based on the taphonomic time

frame inferred from the preservation of the leaf imprint, it can be estimated that the

decomposition of GNUE322001 took at least a month. However, perfectly preserved leaves

were also reported from the Duho Formation (Jung and Lee, 2009); therefore, the taphonomic

scenario inferred from GNUE322001 does not represent a general depositional condition of the

Duho Formation.

Acknowledgments

We thank Professor Yuong-Nam Lee for improving the quality of this manuscript. We

also thank two anonymous reviewers for their valuable feedback and Dr. Adriana López-

Arbarello for overseeing the review process and providing valuable suggestions.

Funding

This work was not supported by any funding source.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of

interest in relation to the content of the article.

336   **Author contributions**

338   Dayun Suh contributed to conceptualization, formal analysis, investigation, visualization,

339   writing of the original draft, and writing of review and editing. Su-Hwan Kim contributed to

340   conceptualization, formal analysis, investigation, methodology, supervision, validation,

341   visualization, and writing of review and editing. Gi-Soo Nam contributed to resources, and

342   validation, and writing of review and editing.

344   **References**

Agassiz, L., 1833-1844, Recherches sur les poissons fossiles, 5 Volumes: Neuchâtel, Imprimerie

de Petitpierre, 1420 p.

Bak, Y., Lee, J.D., and Yun, H., 1996, Middle Miocene radiolarians from the Duho Formation in

the Pohang Basin, Korea: Journal of the Paleontological Society of Korea, v. 12, p. 225–

261.

Bannikov, A.F., 1985, Iskopayemye skumbriyevye SSSR. Trudy Paleontologicheskogo instituta

AN SSSR, v. 210, p, 1–111 [In Russian].

Bannikov, A.F. and Sorbini, L., 1984, On the occurrence of genus Scombrosarda in the Eocene of

Bolca, v. 4, p. 307–317.

Bogatshov, V.V., 1933, Materialy po izucheniyu tretichnoi ikhtiofauny Kavkaza. Trudy

Azerbaidzhanskogo Neftyanogo Issledovatelskogo Instituta, Geologicheskii Otdel, v. 15, p.

1–62 [In Russian].

Bridgham, S.D., and Lamberti, G.A., 2009, Ecological dynamics III: Decomposition in wetlands, *in*

Maltby, E., and Barker, T., eds., The Wetlands Handbook: Blackwell Publishing, Oxford, p.

326–345. https://doi.org/10.1002/9781444315813.ch15

Burrow, C.J., and Turner, S., 2012, Fossil fish taphonomy and the contribution of microfossils in

documenting Devonian vertebrate history, *in* Talent, J.A., ed., Earth and Life: Springer,

New York, p. 184–223. https://doi.org/10.1007/978-90-481-3428-1_8

Byun, H., and Yun, H., 1992, Miocene dinoflagellate cysts from the central part of the Pohang

Basin, Korea: Journal of the Paleontological Society of Korea, v. 8, p. 164–235.

Chough, S.K., Hwang, I.G., and Choe, M.Y., 1990, The Miocene Doumsan fan-delta, southeast

Korea: a composite fan-delta system in back-arc margin: Journal of Sedimentary

Research, v. 60, p. 445–455. https://doi.org/10.1306/212f91ba-2b24-11d7-

8648000102c1865d

Chung, C.-H., and Koh, Y.-K., 2005, Palynostratigraphic and palaeoclimatic investigations on the

Miocene deposits in the Pohang area, South Korea: Review of Palaeobotany and

Palynology, v. 135, p. 1–11. https://doi.org/10.1016/j.revpalbo.2005.02.002

Collette, B.B., and Nauen, C.E., 1983, Scombrids of the world, an annotated and illustrated

catalogue of tunas, mackerels, bonitos, and related species known to date: Food and

Agriculture Organization of the United Nations, Rome, v. 2, 137 p.

Collette, B.B., and Aadland, C.R., 1996, Revision of the frigate tunas (Scombridae, *Auxis*), with

descriptions of two new subspecies from the eastern Pacific. Fishery Bulletin: v. 94, p.

423–441.

Cuvier, G., 1829, Le Règne Animal, distribué d'après son organisation, pour servie de base à

l'histoire naturelle des animaux et d'introduction à l'anatomie comparé: Paris, 532 p.

Godsil, H.C., and Byers, R.D., 1944, A systematic study of the Pacific tunas: California

Department of Fish and Game, Fish Bulletin, v. 60, p. 1–131.

Godsil, H.C., 1954, A descriptive study of certain tuna-like fishes: California Department of Fish

and Game, Fish Bulletin, v. 97, p. 1–185.

Gorjanović-Kramberger, D., 1895, Fosilne ribe Komena, Mrzleka, Hvara i M. Libanona i dodatak

o oligocenskim ribama Tüffera, Zagora i Trifalja. Djela Jugoslavenske Akademije Znanosti

i Umjetnosti, v. 16, p. 1–67.

Graham, J. B., and Dickson, K. A., 2000, The evolution of thunniform locomotion and heat

conservation in scombrid fishes: new insights based on the morphology of *Allothunnus*

390   *fallai*: Zoological Journal of the Linnean Society, v. 129, p. 419–466.

391   https://doi.org/10.1111/j.1096-3642.2000.tb00612.x

392 Grandperrin R., 1978, Influence of currents on the production of tropical seas: consequences

393   for fisheries: SPC Fisheries Newsletter, v. 17, p. 14–20.

394 Hargraves, P.E., 1979, Studies on marine plankton diatoms, IV. Morphology of *Chatoceros*

395   resting spores: Beihefte zur Nova Hedwigia, v. 64, p. 99–120.

396 Hwang, I.G., Chough, S.K., Hong, S.W., and Choe, M.Y., 1995, Controls and evolution of fan delta

397   systems in the Miocene Pohang Basin, SE Korea: Sedimentary Geology, v. 98, p. 147–179.

398 Jawad, L.A., Al-Hassani, L., and Al-Kharusi, L.H., 2013, On the morphology of the vertebral

399   column of the frigate tuna, *Auxis thazard* (Lacepedea, 1800)(Family: Scombridae)

400   collected from the sea of Oman: Acta Musei Nationalis Pragae, v. 69, p. 101–105.

401   https://doi.org/10.14446/amnp.2013.101

402 Jung, S.H., and Lee, S.J., 2009, Fossil-Winged Fruits of *Fraxinus* (Oleaceae) and *Liriodendron*

403   (Magnoliaceae) from the Duho Formation, Pohang Basin, Korea. Acta Geologica Sinica, v.

404   83, p. 845–852. https://doi.org/10.1111/j.1755-6724.2009.00113.x

405 Kim, B.K., 1965, The stratigraphic and paleontologic studies on the Tertiary (Miocene) of the

406   Pohang area, Korea: Seoul University Journal Science and Technology Series, v. 15, p.

407   32–121.

408 Kim, D.H., and Lee, S.-J., 2011, Fossil scallops from the Hagjeon Formation and the Duho

409   Formation, Pohang Basin: Journal of the Geological Society of Korea, v. 47, p. 235–244

410   [in Korean with English abstract].

411 Kim, J., and Paik, I.S., 2013, Chondrites from the Duho Formation (Miocene) in the Yeonil Group,

412   Pohang Basin, Korea: Occurrences and paleoenvironmental implications: Journal of the

413   Geological Society of Korea, v. 49, p. 407–416 [in Korean with English abstract].

414   https://doi.org/10.14770/jgsk.2013.49.3.407

415 Kim, J.H., 2008, A new species of *Acer* samaras from the Miocene Yeonil Group in the Pohang

416   Basin, Korea: Geosciences Journal, v. 12, p. 331–336. https://doi.org/10.1007/s12303-

417   008-0033-6

Kim, J.H., Lee, S., An, J., Lee, H., and Hong, H., 2009, Albizia Fruit Fossils from the Miocene Duho
Formation of Yeonil Group in the Pohang Basin, Korea. Journal of the Korean Earth
Science Society, v. 30, p. 10–18 [in Korean with English abstract].
https://doi.org/10.5467/jkess.2009.30.1.010
Kim, J.-H., Nam, K.-S., and Jeon, Y.-S., 2017, Diversity of Miocene fossil *Acer* from the Pohang
Basin, Korea: Journal of the Geological Society of Korea, v. 53, p. 387–405 [in Korean
with English abstract].
Kim, K.H., Doh, S.-J., Hwang, C.-S., and Lim, D.S., 1993, Paleomagnetic study of the Yeonil Group
in Pohang basin: Journal of Korean Society of Economic and Environmental Geology, v.
26, p. 507–518 [in Korean with English abstract].
Kim, S.-H., Park, J.-Y., and Lee, Y.-N., 2018, A tooth of *Cosmopolitodus hastalis* (Elasmobranchii:
Lamnidae) from the Duho Formation (middle Miocene) of Pohang-si, Gyeongsangbuk-do,
South Korea. Journal of the Geological Society of Korea: v. 54, p. 121–131 [in Korean
with English abstract]. https://doi.org/10.14770/jgsk.2018.54.2.121
Kishinouye, K., 1923, Contributions to the comparative study of the so-called scombrid fishes:
Journal of the College of Agriculture, Imperial University of Tokyo, v. 8, p. 293–475.
Ko, J.Y., 2016, The description of the flat fish (Pleuronectiformes) fossils from the Miocene
Duho Formation, Pohang Yeonam-dong in Korea and its implication: Journal of the
Korean Earth Science Society, v. 37, p. 1–10 [in Korean with English abstract].
https://doi.org/10.5467/jkess.2016.37.1.1
Ko, J.Y., and Nam, K.-S., 2016, *Pleuronichthys* sp. fossils (Pleuronectidae) from the Duho
Formation, Pohang Uhyeon-dong in Korea: Journal of the Korean Earth Science Society,
v. 37, p. 133–142 [in Korean with English abstract].
https://doi.org/10.5467/jkess.2016.37.3.133
Kong, D.Y., and Lee, S.J., 2012, Fossil scaphopods from the Hagjeon Formation and the Duho
Formation, the Cenozoic Pohang Basin, Korea: Korean Journal of Cultural Heritage
Studies, v. 45, p. 218–231 [in Korean].
Kramberger-Gorjanović, D., 1882, Die jungtertiäre Fischfauna Croatiens. Beitrag zur
Paläontologie und Geologie Österreich-Ungarns und des Orients, v. 2, p. 86–136.

Lalli, C. and Parsons, T., 1997, Biological oceanography: an introduction: Oxford, Elsevier

Butterworth-Heinemann, 314 p.

Lee, T.-H., Yi, K., Cheong, C.-S., Jeong, Y.-J., Kim, N., and Kim, M.-J., 2014, SHRIMP U-Pb zircon

geochronology and geochemistry of drill cores from the Pohang Basin: Journal of the

Korean Earth Science Society, v. 23, p. 167–185 [in Korean with English abstract].

https://doi.org/10.7854/jpsk.2014.23.3.167

Lee, Y.G., 1992, Paleontological study of the Tertiary molluscan fauna in Korea: Science Reports

of the Institution of Geosciences University of Tsukuba, Section B (geological science), v.

13, p. 15–125.

Lee, Y.G., 1993, The marine diatom genus *Chaetoceros* Ehrenberg flora and some resting spores

of the Neogene Yeonil Group in the Pohang Basin, Korea: Journal of the Paleontological

Society of Korea, v. 9, p. 24–52.

Lee, Y.-N., Ichishima, H., and Choi, D.K., 2012, First record of a platanistoid cetacean from the

middle Miocene of South Korea: Journal of Vertebrate Paleontology, v. 32, p. 231–234.

https://doi.org/10.1080/02724634.2012.626005

Lim, J.-D., 2005, The first dolphin fossil from the Miocene of Korea: Current Science, v. 89, p.

939–940.

Malyshkina, T.P., Ward, D.J., Nazarkin, M.V., Nam, G.-S., Kwon, S.-H., Lee, J.-H., Kim, T.-W., Kim,

D.-K., and Baek, D.-S., 2023, Miocene Elasmobranchii from the Duho Formation, South

Korea: Historical Biology, v.35, p. 1726–1741.

https://doi.org/10.1080/08912963.2022.21 10870

Nam, G.-S., Ko, J.-Y., and Nazarkin, M.V., 2019, A new lightfish, †*Vinciguerria orientalis*, sp. Nov.

(Teleostei, Stomiiformes, Phosichthyidae), from the middle Miocene of South Korea:

Journal of Vertebrate Paleontology, v. 39, e1625911. https://doi.org/

0.1080/02724634.2019.1625911

Nam, G.-S., Nazarkin, M.V., and Bannikov, A.F., 2021, First discovery of the genus *Auxis*

(Actinopterygii: Scombridae) in the Neogene of South Korea: Bollettino della Societa

Paleontologica Italiana, v. 60, p. 61–67. https://doi.org/10.4435/BSPI.2021.05

Nam, G.-S., and Nazarkin, M.V., 2022, A new lanternfish (Myctophiformes, Myctophidae) from

the Middle Miocene Duho Formation, South Korea: Journal of Vertebrate Paleontology,

v. 42, e2121924. https://doi.org/10.1080/02724634.2022.2121924

Nazarkin, M.V., and Nam, G.-S., 2022, Miocene lightfish Vinciguerria sp. (Stomiiformes:

Phosichthyidae) from the North-West Pacific with notes on the recent evolution of the

genus: Historical Biology, v. 34, p. 102–107.

https://doi.org/10.1080/08912963.2021.1900839

Nelson, J.S., 2006, Fishes of the World (fourth edition): Hoboken, John Wiley and Sons, 601 p.

Nicol, S., Menkes, C., Jurado-Molina, J., Lehodey, P., Usu, T., Kumasi, B., Muller, B., Bell, J.,

Tremblay-Boyer, L., and Briand, K., 2014, Oceanographic characterisation of the Pacific

Ocean and the potential impact of climate variability on tuna stocks and tuna fisheries:

SPC Fisheries Newsletter, v. 145, p. 37–48.

Quintino, V., Sangiorgio, F., Ricardo, F., Mamede, R., Pires, A., Freitas, R., Rodrigues, A.M., and

Basset, A., 2009, In situ experimental study of reed leaf decomposition along a full

salinity gradient: Estuarine, Coastal and Shelf Science, v. 85, p. 497–506.

https://doi.org/10.1016/j.ecss.2009.09.016

Rafinesque, C.S., 1815, Analyse de la nature, ou tableau de l'univers et des corps organisés:

Palermo, 224 p. https://doi.org/10.5962/bhl.title.106607

Risso, A., 1810, Ichthyologie de Nice ou histoire naturelle des poissons du département des

Alpes Maritimes: Paris, F. Schoell, 388 p. https://doi.org/10.5962/bhl.title.7052

Romeo, J., and Mansueti, A.J., 1962, Little tuna, *Euthynnus alletteratus*, in northern Chesapeake

Bay, Maryland, with an illustration of its skeleton: Chesapeake Science, v. 3, p. 257–263.

https://doi.org/10.2307/1350633

Seong, M.N., Kong, D.-Y., Lee, B.J., and Lee, S.-J., 2009, Cenozoic brittle stars (Ophiuroidea) from

the Hagjeon Formation and the Duho Formation, Pohang Basin, Korea: Journal of

Korean Society of Economic and Environmental Geology, v. 42, p. 367–376 [in Korean

with English abstract].

Shaw, J.O., Briggs, D.E.G., and Hull, P.M., 2020, Fossilization potential of marine assemblages

and environments: Geology, v. 49, p. 258–262. https://doi.org/10.1130/g47907.1

Sohn, Y.K., Rhee, C.W., and Shon, H., 2001, Revised stratigraphy and reinterpretation of the

Miocene Pohang basinfill, SE Korea: sequence development in response to tectonism

and eustasy in a back-arc basin margin: Sedimentary Geology, v. 143, p. 265–285.

https://doi.org/10.1016/s0037-0738(01)00100-2

Starks, E.C., 1910, The osteology and mutual relationships of the fishes belonging to the family

Scombridae: Journal of Morphology, v. 21, p. 77–99.

https://doi.org/10.1002/jmor.1050210103

Uchida, R.N., 1981, Synopsis of biological data on frigate tuna, *Auxis thazard*, and bullet tuna, *A.*

rochei: Silver Spring, US Department of Commerce, National Oceanic and Atmospheric

Administration, National Marine Fisheries Service, 63 p.

https://doi.org/10.5962/bhl.title.63109

Vieira, J.M., Costa, P.A., Braga, A.C., São-Clemente, R.R., Ferreira, C.E., and Silva, J.P., 2022, Age,

growth and maturity of frigate tuna (*Auxis thazard* Lacepède, 1800) in the Southeastern

Brazilian coast: Aquatic Living Resources, v. 35, p. 11.

https://doi.org/10.1051/alr/2022010

Wang, Y.-C., and Lee, M.-A., 2019, Composition and distribution of fish larvae surrounding the

upwelling zone in the waters of northeastern Taiwan in summer: Journal of Marine

Science and Technology, v. 27, p. 8.

Woo, K.S., and Khim, B.-K., 2006, Stable oxygen and carbon isotopes of carbonate concretions

of the Miocene Yeonil Group in the Pohang Basin, Korea: Types of concretions and

formation condition: Sedimentary Geology, v. 183, p. 15–30.

https://doi.org/10.1016/j.sedgeo.2005.09.005

Woodward, A.S., 1901, Catalogue of the Fossil Fishes in the British Museum (Natural History)

Part IV: London, British Museum of Natural History, 636 p.

https://doi.org/10.5962/bhl.title.162341

Yoon, S., 1975, Geology and paleontology of the Tertiary Pohang Basin, Pohang District, Korea

Part 1, Geology: Journal of the Geological Society of Korea, v. 11, p. 187–214.

Yoon, S., 1976, Geology and paleontology of the Tertiary Pohang Basin, Pohang District, Korea

Part 2, Paleontology (Mollusca): Journal of the Geological Society of Korea, v. 12, p. 1–22.

Yoshida, H.O., and Nakamura, E.L., 1965, Notes on schooling behavior, spawning, and

morphology of Hawaiian frigate mackerels, *Auxis thazard* and *Auxis rochei*: Copeia, v. 1,

p. 111–114. https://doi.org/10.2307/1441254

Yoshida, H.O., 1979, Synopsis of biological data on tunas of the genus *Euthynnus*: Silver Spring,

US Department of Commerce, National Oceanic and Atmospheric Administration,

National Marine Fisheries Service, 9 p.

Yun, H., 1985, Some fossil Squillidae (Stomatopoda) from the Pohang Tetiary Basin, Korea:

Journal of the Paleontological Society of Korea, v. 1, p. 19–31.

Yun, H., 1986, Emended stratigraphy of the Miocene formations in the Pohang Basin Part I:

Journal of the Paleontological Society of Korea, v. 2, p. 54–69.