Enhanced detection for antibiotic resistance genes in wastewater samples using a CRISPR-enriched metagenomic method

Yuqing Mao[1,2], Joanna L. Shisler[3], Thanh H. Nguyen[1,2,4]

Affiliation

[1] The Grainger College of Engineering, Department of Civil and Environmental Engineering, University of Illinois Urbana-Champaign, IL, USA

[2] Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, IL, USA

[3] Department of Microbiology, University of Illinois Urbana-Champaign, IL, USA

[4] Carle Illinois College of Medicine, University of Illinois Urbana-Champaign, IL, USA

Abstract

The spread of antibiotic resistance genes (ARGs) in the environment is a global public health concern. To date, over 5,000 genes have been identified to express resistance to antibiotics. ARGs are usually low in abundance for wastewater samples, making them difficult to detect. Metagenomic sequencing and quantitative polymerase chain reaction (qPCR), two conventional ARG detection methods, have low sensitivity and low throughput limitations, respectively. We developed a CRISPR-Cas9-modified next-generation sequencing (NGS) method to enrich the targeted ARGs during library preparation. The false negative and false positive of this method were determined based on a mixture of bacterial isolates with known whole-genome sequences. Low values of both false negative (2/1,208) and false positive (1/1,208) proved the method's reliability. We compared the results obtained by this CRISPR-NGS and the conventional NGS method for six untreated wastewater samples. As compared to the ARGs detected in the same samples using the regular NGS method, the CRISPR-NGS method found up to 1,189 more ARGs and up to 61 more ARG families in low abundances, including the clinically important KPC beta-lactamase genes in the six wastewater samples collected from different sources. Compared to the regular NGS method, the CRISPR-NGS method lowered the detection limit of ARGs from the magnitude of 10^{-4} to 10^{-5} as quantified by qPCR relative abundance. The CRISPR-NGS method is promising for ARG detection in wastewater. A similar workflow can also be applied to detect other targets that are in low abundance but of high diversity.

    ## 1 Introduction

Antibiotic resistance (AR) is an increasing threat to global public health. It is
estimated that in the next decade, if without any intervention, AR will cause an annual loss of
3.4 trillion US dollars in gross domestic products and cause 24 million more people to live in
extreme poverty [1]. In addition to surveillance in clinical settings, AR surveillance based on
wastewater has been conducted [2–9]. Culture-independent methods are commonly used for
wastewater surveillance of AR [10], because genes that encode AR are included in bacterial
genomes. By extracting and analyzing the DNA extracted from wastewater samples, the AR
profiles of the entire sample can be obtained. There are two challenges to detect antibiotic
resistance genes (ARGs) in wastewater DNA samples. First, the concentration of many
ARGs can be very low because ARGs account for less than 1% of the DNA in bacterial
genomes [11]. Second, there is a wide range of ARG targets. To date, more than 5,000
reference sequences of ARGs are recorded in The Comprehensive Antibiotic Resistance
Database (CARD) [12]. Metagenomics and quantitative polymerase chain reaction (qPCR) are
two culture-independent methods that are widely used to detect ARGs [10]. These two
methods have their pros and cons in ARG detection. qPCR has a high sensitivity, so it can
capture the presence of ARGs that are in extremely low abundance in a sample [13]. A
previous study found that the detection limit of qPCR genes in manure and compost samples
was 2-8 copies/mg, while that for metagenomics was around 3×10^4 copies/mg [14]. Another
study also found that qPCR could stably detect ARGs in water samples, while metagenomics
could not constantly detect those in low abundances [15]. Besides, qPCR can absolutely
quantify the number of gene copies in a sample [16]. However, a major drawback of qPCR is
that it only returns fluorescent signals when short DNA sequence of 70-300 bp are detected
[17]. These results can neither indicate the mutations in ARGs nor tell the existence of the
remaining sequences outside of this range. This disadvantage makes qPCR easy to
generate false positive results. Another drawback of qPCR is the limited number of targeted
genes. Multiplexed qPCR and chip-based high-throughput qPCR microarrays are applied to
increase the number of targets in one qPCR run. Multiplexed qPCR can target up to 20
genes in one reaction [18], but the interaction among different primers and probes may
become complex when the number of targets increases and make the results inaccurate [19].
Chip-based high-throughput qPCR microarrays can run thousands of qPCR reactions in the
meantime by separating different assays into droplets, allowing it to target dozens to
thousands of genes [20]. This method can avoid the disadvantage of complex interactions of
different primers and probes in conventional multiplexed qPCR while largely increasing the
number of targets. However, to ensure good amplification efficiencies, the design and
validation of specific primers for each individual ARG target is required. When the number of
targets increases, the cost of primer design and validation will also increase. High-
throughput qPCR thus has a hidden cost in the large number of primer validation. In
addition, not all chip-based high-throughput qPCR machines can adjust the numbers of
samples and targets flexibly [21]. Some chip-based high-throughput qPCR machines have
upper limits of 96 or 192 genes to be targeted in one run [22]. In contrast, metagenomics does
not require primer design for specific targets, so that it can detect a wide range of ARGs
without pre-estimation [23]. Next-generation sequencing (NGS) is increasingly accessible
nowadays with the development of cost-friendly benchtop sequencers. However, because of
the low percentage of ARGs in wastewater DNA samples, most of the reads obtained in
conventional metagenomics are not relevant to ARG detection. The sensitivity of
metagenomics is lower than qPCR, making the detection of low-abundance ARGs difficult by
metagenomics [10].

To increase the sensitivity for ARG detection using metagenomics, different methods
have been developed. Hybridization capture is one of the commercialized multiplexed target-

enrichment methods for NGS. This method adds an additional step after indexed library
preparation using streptavidin magnetic beads and specifically designed biotinylated DNA
probes to capture DNA fragments with matched sequences [24]. Thousands to millions of
probes can be used to target various DNA sequences [25]. However, this method is expensive,
complex, and time-consuming because of the synthesis of highly multiplexed biotinylated
DNA/RNA probes [26,27]. Recently, CRISPR (clustered regularly interspaced short palindromic
repeats)-Cas9-based target enrichment NGS methods were developed for clinical diagnosis
[28,29]. Such methods use specifically designed guide RNA (gRNA) to guide Cas9 nuclease
specifically cleave the target genes to generate 5' phosphate group for adapter ligation. The
libraries generated by this method can be applied to both the Illumina NGS platform and the
Nanopore 3rd-generation sequencing platform [28,29]. Unlike qPCR where multiplexed primers
and probes may have complex and unexpected interactions [19], the interaction of Cas9-gRNA
complexes to DNA samples is simpler, making the Cas9 cleavage reactions possible to be
highly multiplexed [30]. Unlike biotinylated DNA probes in hybridization capture (which do not
contain conserved regions), the gRNA used for Cas9 cleavage contains conserved regions
[31], making the synthesis of multiplexed assays of CRISPR-Cas9 enriched methods more
flexible and economical.

NGS is increasingly accessible to laboratories because of its reduced cost [32]. In this
study, we developed a novel CRISPR-NGS method for ARG detection in wastewater
samples. The detailed step-by-step protocol for the entire CRISPR-NGS workflow can be
found at dx.doi.org/10.17504/protocols.io.8epv5xdnjg1b/v2. This CRISPR-NGS method
includes six major steps (**Figure 1**). Step 1 is to prepare a highly multiplexed gRNA pool
(i.e., 6,010 different sequences) targeting different sites in ARGs specifically. We chose to
synthesize trans-activating CRISPR RNA (tracrRNA) and multiplexed CRISPR RNA (crRNA)
from DNA templates to reduce the cost because purchasing gRNA individually will be
extremely costly [31]. Step 2 is to prepare DNA samples for sequencing. This step includes
three sub-steps of wastewater DNA extraction, DNA blocking by 5' phosphate group
removal, and DNA positive control spike. In NGS library preparation, adapter ligation is a
necessary step which requires DNA fragments have 5' phosphate group. To avoid adapters
ligating to DNA fragments irrelevant to ARG detection, we used alkaline phosphatase to
remove 5' phosphate groups from all DNA fragments before library preparation. After
blocking, a 250-bp DNA fragment with synthesized 5' phosphate group obtained from
Pseudogulbenkiania sp. strain NH8B which should be absent in most DNA samples was
spiked in the DNA sample as the positive control for sequencing [33]. Step 3 is to specifically
cleave the DNA samples at the sites in the ARGs using Cas9 nuclease to generate new 5'
phosphate groups for adapter ligation. The newly generated 5' phosphorylated ARG
fragments were further modified by a dA-tailing step for optimized adapter ligation efficiency.
Step 4 is to ligate the adapters to the specifically cleaved ARG fragments. Step 5 is to
amplify the adapter-ligated fragments by polymerase chain reaction (PCR) using the primers
targeting the conserved region on the adapters to enrich the ARG fragments in the DNA
sample. Step 6 is to check the fragment lengths and the DNA concentrations of the library
and load the libraries for NGS. As described here, the CRISPR-NGS method modifies the
DNA fragmentation step in NGS library preparation, so that the sensitivity of NGS in ARG
detection can be largely increased. With the same total number of reads, hundreds more
ARGs in low abundances in sewage samples can be uncovered, and a lower detection limit
can be expanded from the magnitude of 10^{-4} to 10^{-5} as determined by qPCR relative
abundance.

RT: room temperature

Figure 1. The workflow of the CRISPR-NGS method.

2 Methods

2.1 Wastewater sample collection, pre-treatment, and DNA extraction

The wastewater samples used in this study included five influents from the wastewater treatment plants in Chicago, Illinois, and one raw sewage composite samples directly collected from a manhole. More information on sample collection and processing is in **Supplementary Methods S1**.

2.2 Mock community DNA sample preparation

We estimated the false positive and false negative detection of ARGs using the mixture of DNA extracts at equal concentrations from 10 bacteria isolates (three *E. coli* from wastewater, three *E. coli* from clinical sources, one *Shigella* spp. from a clinical source, two *Salmonella* spp. from clinical sources, and one *Pseudomonas aeruginosa* PA14 (see **Table S1**)[34]. This mock community was subjected to both whole-genome sequencing and CRISPR-NGS to determine the false positive and false negative rates of the CRISPR-NGS method. Details on DNA extraction, preparation, and subsequent whole-genome sequencing are in **Supplementary Methods S2**. The whole-genome sequences are available in BioProject PRJNA1147862 (to be released upon publication)[35].

2.3 Design and synthesis of crRNA and tracrRNA

Multiplexed gRNA for ARG detection was designed by FLASHit[28]. Four thousand six hundred and forty-one ARG nucleotide sequences for gRNA design input were downloaded from the CARD protein homolog model on September 12, 2022[36]. The default exclusion of off-targets in the *E. coli* BL21 genome in FLASHit was removed, because the Cas9 protein used in this study was commercially purchased. In total 6,010 different 20-nt spacer sequences were generated by FLASHit to cleave target ARGs into approximately 200-bp fragments. More details of the synthesis of crRNA and tracrRNA are in **Supplementary Methods S3**.

2.4 CRISPR-NGS and regular NGS library preparation and sequencing

We used Cas9 nuclease to specifically cleave targeted sites for DNA fragmentation in CRISPR-NGS library preparation. For regular NGS library preparation, we used non-specific enzymes for DNA fragmentation. A synthesized 250-bp DNA fragment obtained from *Pseudogulbenkiania* sp. strain NH8B with 5' phosphate group was spiked to all libraries as the positive control [33]. In the adapter ligation step, xGen™ UDI-UMI Adapters (Integrated DNA Technologies) was used to reduce the PCR bias. Detailed information for CRISPR-NGS library preparation, regular NGS library preparation, library quality control, and Illumina sequencing is in **Supplementary Methods S4-S5**.

2.5 Sequencing read mapping

After paired-end sequencing, UMI-tools, PriceSeqFilter, Bowtie2, and SAMtools were used for read deduplication, quality control, and ARG mapping [37–41]. The same read mapping methods were used for both CRISPR-NGS and regular NGS libraries. All sequencing raw data, including the separated UMI index files, can be found in BioProject PRJNA1148079 (to be released upon publication) [42]. The reference file for read mapping includes the ARG sequences downloaded from CARD (see 2.3) and the sequence of the 250-bp NH8B DNA positive control. Any ARGs with coverages higher than 75% were considered present in the sample, according to previous studies [43–45]. The identity threshold was approximately 90%, according to the end-to-end alignment algorithm of Bowtie2 [39]. Detailed information for read mapping is in **Supplementary Methods S6**.

3 Results and Discussions

3.1 Method validation

We validated that alkaline phosphatase blocking inhibited the T4 DNA ligase from connecting different DNA fragments together. The detailed method for the validation of 5' phosphate group removal can be found in **Supplementary Methods S7**. After phosphatase blocking, the 5' phosphate group of DNA fragments is removed, and T4 ligase could not anneal DNA fragments to form new fragments of other sizes (**Figure S1**). We used a synthetic 731-bp DNA template with 5' phosphorylation to test the blocking efficiency of alkaline phosphatase. After phosphatase blocking, the DNA was incubated with T4 ligase. As shown in **Figure S1**, while the DNA sample without blocking showed multiple bands, the blocked DNA remained its original 731-bp size, suggesting the 5' phosphate groups of blocked DNA fragments were removed, and T4 DNA ligase could not ligate these DNA fragments.

We validated the Cas9 cleavage using an in-silico analysis that compares the breakpoints in raw sequencing reads with the intended Cas9 cleavage sites. In read mapping, paired 150-bp raw sequencing reads were aligned to the reference ARG sequence obtained from CARD. The paired reads, which started from different ends of the same DNA fragment, should have an overlapping region if the DNA fragment is smaller than 300 bp. If the reads are from different fragments, there would be no overlap. Because most DNA fragments were designed to be around 200 bp using a protocol developed previously [28] and the maximum read length was 150 bp in NGS, the entire DNA fragment would not be covered by one single read but could be covered by two paired reads from different directions. Therefore, in the read mapping plot of a cleaved 200-bp DNA fragment, we expected to see the middle part (usually around 100 bp) overlapped by the paired 150-bp reads from different directions and the remaining left and right part covered by the remaining lengths of the paired reads. Because the DNA fragments were generated by Cas9 cleavage

at a few designed sites, these DNA fragments should not overlap with each other in the read
mapping of the ARG. The locations where two DNA fragments meet each other should be
the same as the designed Cas9 cleavage sites. The example of overlapping regions on
different DNA fragments in the same ARG was marked and shown in **Figure S2,** which
shows the read mapping plot for CRISPR-NGS and regular NGS for the same gene OXA-36
(ARO: 3001430) [12]. The four cleavage sites on OXA-36 broke the read mapping histogram of
CRISPR-NGS into three overlapping regions while the regular NGS read mapping histogram
was smooth. We checked the read mapping of 79 ARGs from different ARG families, 78 of
the raw reads were broken at the designed cleavage sites except for *rpoB* which contained a
few off-target cleavage sites. This finding suggested most of the ARGs detected by CRISPR-
NGS truly existed in the samples.

221
222

3.2 CRISPR-NGS method showed high accuracy in ARG detection in the mock
community sample
We conducted whole-genome sequencing of 10 bacteria isolates (listed in **Table S1**)
and these genomes served as a basis to determine false positives and false negatives in
ARG detection by the CRISPR-NGS method. Because some ARGs share highly similar
sequences, we divided the 4,641 targeted ARGs into 1,208 clusters based on their
similarities in terms of ANI to avoid overcounting these ARGs (**Figure S3; Spreadsheet S1**).
The details of ARG clustering method are in **Supplementary Methods S8**. We first used
annotation for the 10 whole-genome sequences to find 100 ARGs in 99 clusters existing in
the mock community. The CRISPR-NGS method detected 1,032 ARGs in 124 clusters in the
same sample (**Spreadsheet S2**). Ninety-seven clusters were found by both CRISPR-NGS
and annotation of the whole genomes. Because the algorithm of whole-genome annotation
may miss some ARGs [46], we used BLAST to search the ARG sequences listed in CARD
within the 10 whole bacterial genomes in the mock community [47]. We found 25 additional
clusters of ARGs that were detected by the CRISPR-NGS method in the mock community.
Finally, we used Sanger sequencing to confirm the presence of two clusters of ARGs that
were not found by either annotation or BLAST in the 10 whole-genome sequences, one
containing quinolone resistance protein (*qnr*) genes and the other containing one single ARG
aph(3')-Ia. The primer sequences of these two genes are listed in **Table S2**. The amplicon of
the *qnrB5* gene was aligned to the CARD reference sequence in 99.65% identity, suggesting
this gene was missed by long-read PacBio sequencing but could be detected by short-read
NGS. In total, as summarized in **Table S3**, we confirmed the presence of 123 ARGs cluster
detected by the CRISPR-NGS method (97 from annotation of **Spreadsheet S3**, 25 from the
BLAST method and one from Sanger sequencing). These clusters are the true positives.
There were two false negatives, which were found by whole-genome sequencing but not by
CRISPR-NGS. The maximum identities of these two ARG clusters in all mock community
genomes were only 86-87% when comparing with the reference sequences, while the
maximum identities of any other true positive ARGs were equal to or greater than 93%. The
NGS libraries used Bowtie2 for ARG identification which has an identity threshold of
approximately 90% under our current settings [39]. In contrast, ARGs were identified from the
whole-genome sequences by the Genome Annotation tool on Bacterial and Viral
Bioinformatics Resource Center (BV-BRC) with an identity threshold of 80% [36,48,49]. The
difference in identity thresholds made the two ARG clusters of 86-87% identities missed by
NGS.
We also compared the ARG detection results for the mock community sample to the
regular NGS method using the similar methods as described above. The regular NGS library
detected two more ARG clusters, *dfrA14* and *aadA13*, than the CRISPR-NGS library
(**Spreadsheet S4**). However, the identities of these two ARGs were 89% in BLAST against
the reference sequence. The coverage of *aadA13* (ARO: 3002613, 799 bp) in the CRISPR-
NGS library was 65.91%, slightly below the 75% threshold. It is worth noting that regular

NGS didn't have any reads mapped to the reference sequence after approximately 700 bp, suggesting potential accumulation of mutations there (**Figure S4**). The cleavage site of CRISPR-NGS at 772-795 bp might be affected due to the mutations and resulted in the low coverage. The failure in *dfrA14* (ARO: 3002859, 475 bp) detection might be due to the unsuccessful cleavage at 15-38 bp (**Figure S4**). To conclude, though CRISPR-NGS may miss some ARGs with accumulated mutations that make the identities of the ARGs below 90% with reference sequences, its overall accuracy in ARG detection is high.

3.3 High reproducibility in ARG detection was observed in the CRISPR-NGS libraries made from the same DNA sample

The reproducibility of the CRISPR-NGS method was tested by comparing the results of two sewage samples collected from Calumet on 8/16/2023 and 10/24/2023. The detailed method is in **Supplementary Methods S9**. The DNA extracted from these samples were made into duplicated libraries. Nine hundred and sixty ARGs were detected by both libraries for Calumet (8/16/2023), with 1,020 and 1,045 ARGs detected in each library (94.1% and 91.9% were detected in both). As for Calumet (10/24/2023), 1,597 and 1,281 ARGs were detected in each duplicate, while 1,263 ARGs were shared in common (79.1% and 98.6% were detected in both).

The read depths for the ARGs detected in only one duplicated library ranged from 1.03 to 291.68, with a median of 7.70, for Calumet (8/16/2023) and ranged from 1.08 to 1041.34, with a median of 20.72, for Calumet (10/24/2023). These ARGs were missed because their coverages were below the 75% threshold, instead of no read alignment. For example, the 987-bp *AAC6_30_AAC6_Ib* gene (ARO: 3002599) detected only in one of the two libraries of Calumet (10/24/2023) with read depth of 1041.34 and coverage of 82.07%, while the other library only got a coverage of 60.59%. As a comparison, we also made duplicated regular NGS libraries for Calumet (10/24/2023). One hundred and fifty-six ARGs were detected by both libraries, while 399 and 190 ARGs were detected by each individual library (39.1% and 82.1%). The read depths of the ARGs detected only by one library ranged from 0.91 to 6.07, with a median of 1.36. These findings suggest that though having 100% same ARG detection by metagenomics is difficult, it is still possible to repeatedly detect most of the ARGs in the same DNA sample by CRISPR-NGS.

Then we compared the read depths of the ARGs detected in both duplicated libraries. The read depths of shared ARGs were fitted linearly in log-10 scale with slopes of 0.92 and 1.05, intercepts of 0.017 and -0.377 (**Figure 2**). The R^2 values were 0.90 and 0.98, respectively, suggesting low fitting variations. The slopes of both fittings were close to 1, and the intercepts were close to 0, suggesting the read depths of the ARGs detected in the duplicated libraries were reproducible.

Figure 2. The ARG read depths in replicated libraries made from the same DNA samples (a: Calumet, 8/16/2023; b: Calumet, 10/24/2023) fitted linearly in log-10 scale with R^2 values of 0.90 and 0.98, respectively.

3.4 More ARGs were detected by the CRISPR-NGS method than regular NGS method

The improved sensitivity of the CRISPR-NGS method in ARG detection was evaluated by comparing the number of ARGs and ARG families detected in the same sewage sample by the CRISPR-NGS method and regular NGS method, respectively. **Figure 3** shows the number of ARGs and ARG families that were mapped with over 75% coverage by only the CRISPR-NGS method, only the regular NGS method, and both methods. The sewage samples had 288-433 ARGs detected by regular NGS libraries, and these numbers increased to 542-1,615 for CRISPR-NGS libraries. Only from seven to 43 ARGs were uniquely detected by regular NGS libraries, while 278-1,189 ARGs were uniquely detected by CRISPR-NGS libraries. The ARGs detected in wastewater samples by different CRISPR-NGS and regular NGS libraries are listed in **Spreadsheet S5 and S6**. These additional ARGs detected by CRISPR-NGS libraries were mostly from ARG families that were not covered by regular NGS libraries. Seventeen to 61 more ARG families were only detected by CRISPR-NGS libraries for the sewage samples, while zero to three were only detected by regular NGS libraries. CRISPR-NGS libraries missed these ARG families mainly because the coverages of these ARGs were slightly lower than the 75% threshold. For example, CfxA beta-lactamase genes were missed by three CRISPR-NGS libraries (Calumet 10/24/2023, Calumet 12/19/2023, and Lemont 12/19/2023). The coverages of CfxA2 (ARO: 3003002) in these libraries were between 72.05% and 72.57%, less than 3% below the coverage threshold. The ARG families detected only by CRISPR-NGS libraries include KPC and CTX-M beta-lactamase genes which are clinically important. Another example is the mobilized colistin resistance (MCR) phosphoethanolamine transferase genes that encode resistance to colistin which is one of the last-resort antibiotics [50,51]. The CRISPR-NGS method detected MCR genes in a high diversity, including MCR-3, MCR-4, MCR-5, MCR-9, and MCR-10, while regular NGS only detected MCR-3 or MCR-5 in some libraries.

Figure 3. More ARGs (a) and ARG families (b) were detected by CRISPR-NGS method than regular NGS method in different sewage samples. The yellow bars show the number of ARGs and ARG families only detected by CRISPR-NGS method, the purple bars show the number of ARGs and ARG families detected by both CRISPR-NGS and regular NGS methods, and the dark blue bars show the number of ARGs and ARG families only detected by the regular NGS method.

3.5 CRISPR-NGS method could detect ARGs in higher read depths than regular NGS method

The ARGs detected by CRISPR-NGS libraries also had higher read depths than those detected by regular NGS libraries. The fold changes in read depths of the 145 ARGs detected in all sewage samples by both CRISPR and regular NGS libraries are plotted in **Figure 4**. All ARGs had significantly higher read depths in CRISPR-NGS libraries than regular NGS libraries. The fold changes in read depths varied from 2.73 to 126.45, suggesting the CRISPR-NGS method can detect ARGs in higher confidence than regular NGS method.

Figure 4. The read depths of all 145 ARGs detected in all sewage samples were significantly higher in CRISPR-NGS libraries than regular NGS libraries. The y axis shows the p-values in a reversed log-10 manner, and the x axis shows the fold changes in read depths (CRISPR-NGS versus regular NGS). The horizontal dash line means the threshold of a significant p-value equals to 0.05, and the vertical dash line means the thresholds in fold change of 1.5.

To compare the ARG detection efficiencies of CRISPR-NGS and regular NGS with deeper sequencing read depths, we subsampled the CRISPR-NGS reads and compared the ARG mapping results with the corresponding regular NGS libraries without subsampling (**Figure 5**). The subsampling protocol can be found in **Supplementary Methods S10**. Before subsampling, the total number of reads for CRISPR-NGS libraries for sewage samples ranged from 43.3 million to 58.2 million, and the total number of reads for regular NGS libraries ranged from 44.6 million to 60.7 million (**Table S4**). When subsampling 2% of the reads, three CRISPR-NGS libraries (Calumet 12/19/2023, Calumet 10/24/2023 A and B) detected more ARGs than the corresponding regular NGS libraries. When the subsampling ratio reached 10%, all CRISPR-NGS libraries except for Lemont (12/19/2023) detected more ARGs than corresponding regular NGS libraries. The CRISPR-NGS library for Lemont (12/19/2023) detected 89.6% of the number of ARGs detected by its regular NGS library when doing 10% subsampling, and the percentage increased to 99.2% at subsampling ratio of 20%, and 171.2% at subsampling ratio of 50%. At subsampling ratio of 50%, all curves reached a plateau phase (**Figure 5**). These findings suggest that CRISPR-NGS can detect similar numbers of ARGs as regular NGS with approximately 2-20% of the total number of reads. With approximately 25 million total number of reads (i.e., approximately 50% of the read depths used in this study), the number of ARGs detected by CRISPR-NGS approaches maximum. This improvement suggests that CRISPR-NGS can make it possible to detect ARGs deeply by metagenomics on many benchtop sequencers that can generate more than 25 million reads in one run, instead of ultra-deep sequencing for regular NGS libraries that usually require expensive and professional production-scale sequencers.

377
Figure 5. When the total read depth reaches approximately 2-20% of current read depth,
CRISPR-NGS libraries can detect similar numbers of ARGs as regular NGS libraries with
their original read depths. The CRISPR-NGS libraries were subsampled by ratios of 1%, 2%,
5%, 10%, 20%, 50%, and 70%. The horizontal dash line at y=1 means equal number of
ARGs detected by the subsampled CRISPR-NGS library and the regular NGS library with
original read depth.
384

3.6 ARG presence could be validated by Sanger sequencing

We used Sanger sequencing to validate the presence of selected ARGs in sewage
samples (**Table 1**). The ARG targets were amplified by PCR, and the PCR amplicons were
Sanger sequenced and compared to the reference sequences in CARD. The detailed
method for Sanger sequencing validation can be found in **Supplementary Methods S11**.
We chose Sanger sequencing for validation, because Sanger sequencing for PCR
amplicons is accurate, cost-effective, and fast when there are only a few targets. The
identities between the sequenced PCR amplicons and the reference sequence of the ARGs
in CARD were between 89% and 100%, suggesting the amplicon sequences were highly
similar to the reference sequences. Because the PCR primers designed for different ARGs
had different binding regions, the amplicons could not cover the full lengths of the ARGs.
After adjusting with the amplicon lengths, the coverage of the ARGs were between 58% and
99%. The 58% coverage of *aac(3)-IId* and 73% coverage of *aph(3')-Ia* were due to the low
quality of Sanger sequencing of the campus town manhole sample. After removing these
outliers, the coverages were between 77% and 99%. The high coverages and identities of
Sanger sequencing also matched the read mapping plots. For example, the two colistin
resistance genes, MCR-5.1 and MCR-4.1, were nearly fully mapped by CRISPR-NGS reads
(**Figure S5**), though these two genes were barely detected by regular NGS.

Table 1. The Sanger sequencing identities and adjusted coverages for the ARGs detected in
different CRISPR-NGS libraries

Gene	Identity (%)	Coverage adjusted by amplicon lengths (%)
aph(3")-Ib	99-100	95-98
sul1	97-99	95-96
CRP	89-94	95
sul2	95-99	92-96
aph(3')-Ia	95-99	73-97
dfrA1	96-97	93-95
GES-26	95-100	96-97
MCR-5.1	98	77-99
MCR-4.1	98	98
TEM-1	99	96
aac(3)-IId	96-99	58-93
tet(G)	94	88
qnrB5	100	97

Note: The coverages were adjusted by dividing the aligned length by the amplicon length, rather than the coverage provided in BLAST, which divides the aligned length by the query length (i.e., the reference ARG length in CARD).

3.7 CRISPR-NGS libraries had lower detection limit than regular NGS libraries, and the read depths in CRISPR-NGS libraries were proportional to the relative abundances determined by qPCR

We evaluated the detection limit of the CRISPR-NGS method by plotting the read depths against the relative abundance detected by qPCR for nine ARGs with various read depths detected by CRISPR-NGS or regular NGS methods in sewage samples (**Figure 6**). The detailed method for qPCR can be found in **Supplementary Methods S12**. The ARGs and corresponding qPCR primers are listed in **Table S5**. The Cq values of the 16S rRNA gene in the sewage samples are listed in **Table S6**. The read depths in both CRISPR-NGS and regular NGS libraries increased when the qPCR relative abundances increased, suggesting the read depths of CRISPR-NGS and regular NGS can be used for ARG abundance estimation. As shown in **Figure 6**, the lower detection limit of the CRISPR-NGS method for these nine ARGs was down to 2.23×10^{-5} copies ARG per copy of 16S rRNA gene, while the lower detection limit of regular NGS method was at 1.79×10^{-4} copies ARG per copy of 16S rRNA gene. Thus, the CRISPR-NGS method has a higher sensitivity than regular NGS method in ARG detection. A previous study reported varied relative abundances of ARGs in wastewater samples ranging from 1.4×10^{-5} to 0.09 [52], suggesting the improved lower detection limit of CRISPR-NGS can promote the detection of ARGs in low abundances in wastewater samples.

Figure 6. The read depths versus the qPCR relative abundances of the nine ARGs detected in different wastewater samples. The orange dots are the read depths of the ARGs detected in the CRISPR-NGS libraries. The blue dots are the read depths of the ARGs detected in the regular NGS libraries. The vertical dash lines in corresponding colors indicate the lower detection limits of CRISPR-NGS and regular NGS in terms of qPCR relative abundance. CRISPR-NGS had a lower detection limit of 2.23×10^{-5}, as labeled by the orange vertical line, while the lower detection limit of regular NGS was 1.79×10^{-4}, as labeled by the blue vertical line.

3.8 CRISPR-NGS is better at detecting low abundance ARG than regular NGS

The most abundant clusters of ARGs (i.e., ARG clusters with read depths greater than 200 in any of the CRISPR-NGS libraries) in the sewage samples were selected and plotted into heatmaps for comparison in **Figure 7**. These 41 ARG clusters were distributed in eight major drug classes, including aminoglycoside (8), beta-lactam (17), antiseptics (2), fluoroquinolone (2), macrolide (3), multi-drug (3), phenicol (2), and tetracycline (4). The ARG clusters in **Figure 7** are named based on the ARG names and families in the cluster. The multi-drug class was defined to include the ARGs that are resistant to at least two drug classes in the 61 sub-terms under the CARD "antibiotic molecule" ontology term [12]. Comparing between CRISPR-NGS and regular NGS libraries, the entire group of CMY-2 and KPC beta-lactamase genes and *qnrB8*-like quinolone resistance genes were missed by regular NGS. Comparing the wastewater treatment plant sewage samples, the campus town manhole sample had lower abundances of GES, MOX-9, and OXA-427-like beta-lactamase genes than the wastewater treatment plant (WWTP) influent samples. The abundance of the

tet(C) gene in the campus town manhole sample was greater than any of the Chicago
WWTP influent samples, but *tet(E)* in the campus town manhole sample was lower than the
WWTP influent samples. The above findings suggested different ARG distributions in
different geographical areas or different sampling scales. As for the ARGs shared in
common, *sul1*, *qacE/qacEdelta1*, *qacL*, and OXA-2-like beta-lactamase genes were
constantly more abundant than the other ARGs in all sewage samples. Since *sul1*, *qacE*,
and *qacEdelta1* are known to be located on class 1 integrons which play an important role in
the spread of ARGs, our findings suggest that class 1 integron may be actively transferring
ARGs in these wastewater samples [53].

Figure 7. The heatmaps for the most abundant ARGs (i.e., the ARGs with read depths
greater than 200 in at least one library detected by CRISPR-NGS) in different sewage
samples. a: The read depths of the most abundant ARGs detected by CRISPR-NGS
libraries; b: The read depths of the most abundant ARGs detected by regular NGS libraries.
The read depths increase from green to pink. The white color means the ARG was not
detected in the sample by the corresponding method. From top to bottom, the heatmaps
were divided into eight parts by horizontal black lines according to the drug classes of the
corresponding ARGs: aminoglycoside, beta-lactam, antiseptic, fluoroquinolone, macrolide,
multi-drug, phenicol, and tetracycline. The solid vertical lines divide samples by different
locations of collection, and the dash vertical lines divide samples from the same location by
different collection times.

When extending the scale to all ARGs with high or low abundances, the profiles
detected by CRISPR-NGS and regular NGS were more different (**Figure S6**). The entire
class of aminocoumarin antibiotics was not detected by regular NGS libraries but by
CRISPR-NGS libraries. Fewer classes of beta-lactamase genes were detected by regular
NGS libraries than CRISPR-NGS libraries. For example, KPC and CTX-M were not detected
by any regular NGS libraries but by CRISPR-NGS libraries. Similarly, regular NGS libraries
missed several clusters of multi-drug, peptide, diaminopyrimidine, and phenicol resistance
genes which were captured by CRISPR-NGS libraries. But for aminoglycoside, macrolide,
and tetracycline resistance genes, regular NGS libraries detected similar numbers as
CRISPR-NGS libraries. Such findings suggest that when the range of ARGs for surveillance
expands, CRISPR-NGS works better in capturing the ARGs in the classes that are less

abundant than the three drug classes mentioned above.

4 Discussion

Metagenomics has been applied widely in ARG surveillance in different water and
wastewater samples [54–58]. The advantage of metagenomics is that it can detect a wide range
of ARGs without knowing the targets. However, it is challenging to sequence ARGs present
in low abundances by metagenomics. We overcome this disadvantage by using CRISPR-
Cas9 with multiplexed gRNA to specifically enrich the ARGs in wastewater DNA samples
during NGS library preparation.

After the CRISPR-Cas9 treatment, compared to regular NGS libraries, up to 1,189
more ARGs were detected, and CRISPR-NGS libraries only require 2-20% total number of
reads to detect similar number of ARGs as regular NGS libraries. Based on the findings in
this study, CRISPR-NGS works the best for samples that contain highly diverse ARGs in
relatively low abundances. Various types of extended-spectrum beta-lactamase (ESBL)
genes were detected in the sewage samples in this study by CRISPR-NGS. These genes
include not only the common ones like TEM, CTX-M, and OXA, but also the rare ones, such
as GES, PER, VEB, BES, and TLA [59]. Among these genes, GES, PER, OXA, and VEB were
abundant enough to be constantly detected by regular NGS. Other genes were too diluted to
be stably detected without CRISPR treatment before NGS, including the clinically important
carbapenem resistance gene KPC and cefotaxime resistance gene CTX-M. The return is low
but extra effort is required to monitor these rare ESBLs using conventional methods because
their abundances were too low to be detected in most cases. However, studies showed that
these ESBLs were also evolving and spreading globally [60,61]. For example, GES-2 gained
resistance to carbapenem by only a single point mutation [62]. Though rarely detected in the
clinical isolates obtained until now, our findings indicated that these ESBL genes were
circulating in human communities and could potentially cause public health problems in the
future. For example, though considered as a less common ESBL, the read depths of GES in
WWTP samples were in the top 17 of all 76 detectable beta-lactamase gene clusters. In
contrast, GES was not as abundant as in WWTP samples for the campus town manhole
sample. This finding suggested that the presence and abundance of different ESBL may be
dynamic. Innovative high-throughput detection methods with high sensitivity such as
CRISPR-NGS will be helpful in the surveillance of the emergence of such unexpected ARGs.

The CRISPR-NGS method is promising for ARG detection, especially for the
surveillance of sewage samples which may have many unexpected targets. Early warning of
new threats in public health is challenging, because lots of new targets may be hidden in low
abundance in human communities and suddenly emerge to cause an outbreak [63].
Surveillance methods should be high in throughput and sensitivity simultaneously. When the
throughput is high, multiplexed or chip-based qPCR is labor-intensive in primer design and
sample preparation. Because of such reasons, NGS is more suitable for monitoring
unexpected targets than qPCR. Conventional NGS can increase its sensitivity by increasing
the total number of reads during a sequencing run [64,65]. However, this method can be costly,
because increasing the total read depth by 10-fold means the total sequencing cost for the
same sample is increased by 10-fold. Currently, the Illumina sequencing cost is around $2.2
per million reads [66]. For a library with 50 million reads, the cost solely for sequencing is
around $110. If the total number of reads increases by 10-fold to 500 million, the sequencing
cost for the same library will increase to around $1,100. Considering more than 99.9% reads
are irrelevant to ARG detection [24], increasing the proportion of ARG reads before loading the
samples to sequencing machine will be more economical. The CRISPR-NGS method could

detect the same number of ARGs read depth using 2-20% of total number of reads required
by regular NGS libraries with only an increased cost of around $30 in Cas9 nuclease and
gRNA preparation for each library. The relatively low cost makes the CRISPR-NGS method
attractive for routine ARG surveillance in the future. The reduced read depth requirement of
CRISPR-NGS make it possible to use downgraded but less expensive benchtop sequencers
like MiSeq to detect ARGs by metagenomic sequencing and generate results comparable to
the more advanced production-scale sequencers like NovaSeq.

There are also limitations of the CRISPR-NGS method for future improvement. First,
genetic information other than ARGs, for example, the pathogen genomes and mobile
genetic elements, may be missed by CRISPR-NGS method, because these sequences are
not targeted during assay design. ARGs that are currently unknown may also be missed for
the same reason. To detect these targets, future studies can include the sequences of
genetic markers of pathogens and mobile genetic elements in gRNA design. Combining
CRISPR with 3[rd] generation long-read sequencing can also help identify the host information
of ARGs, because the upstream and downstream sequences of target ARGs can be
identified accordingly [67]. Second, the read depths of ARGs may be biased because of the
high number of PCR cycles used during CRISPR-NGS library preparation if using
conventional adapters. UDI-UMI adapters with molecule-specific adapter sequences are
recommended to reduce the PCR bias, because the over-amplified DNA fragments can be
identified by their shared UMI indexes. Third, CRISPR-NGS cannot quantify ARGs in
absolute copy numbers as qPCR, the relative quantification is also not as accurate as qPCR.
We recommend using CRISPR-NGS periodically for wastewater surveillance to identify
ARGs relevant for public health and use qPCR to quantify these ARGs in a more frequent
manner. Fourth, between seven to 43 ARGs were missed by CRISPR-NGS but captured by
regular NGS in sewage samples, suggesting not all ARGs were successfully targeted by
CRISPR-NGS. This may be due to the low on-target cleavage efficiency of some gRNA or
due to "dropouts" in template DNA oligo pool synthesis. Though Cas9 nuclease can cleave
DNA at targeted sites with a few mismatches [68], the efficiency of Cas9 binding and cleavage
is reduced when mismatches occur. In addition, the current DNA oligo pool synthesis
technique cannot guarantee 100% synthesis of all oligos. Around 0.8% of the sequences in
a DNA oligo pool can be missed during synthesis. For the oligo pool used in this study with
6,010 different sequences, the number of "dropout" oligos is around 48. The corresponding
gRNA would not be transcribed and used in library preparation. Selecting a vendor with
relatively lower "dropout" ratio for oligo pool synthesis may help increase the on-target
cleavage ratio. Fifth, the CRISPR-NGS method may require a more flexible read mapping
threshold (e.g., different coverage threshold for different ARGs) to avoid false negative
results, making the bioinformatic pipeline after sequencing more complex. In this study, we
used a fixed coverage threshold of 75% to make CRISPR-NGS results comparable to
regular NGS, but future studies can develop a bioinformatic pipeline with flexible threshold
specifically for CRISPR-NGS. Because of the recognition of the protospacer adjacent motif
(PAM) regions, gRNA sequences cannot be designed exactly at the beginning and the end
of the ARGs. This disadvantage in gRNA design can result in reduced coverage in read
mapping of some ARGs, causing false negative results of these ARGs when using a fixed
threshold.

5 Conclusions

Here, we introduced a CRISPR-Cas9 modified NGS library preparation method to
increase the sensitivity in ARG detection in sewage samples. Up to 61 ARG families or up to
1,189 ARGs present in low abundance were uncovered by NGS after CRISPR-Cas9

modification. With the same total read depth for one sequencing run, CRISPR-NGS libraries lower the detection limit from magnitude of 10^{-4} to 10^{-5} as determined by qPCR relative abundance. CRISPR-NGS could detect similar number of ARGs as regular NGS with only 2-20% required total number of reads. CRISPR-NGS only requires approximately 25 million reads to detect ARGs deeply in sewage samples, making it possible to be applied to cost-friendly and more accessible benchtop sequencers.

Acknowledgement

We acknowledge funding from Water Research Foundation (5182) and USEPA (R840487). This study has not been formally reviewed by EPA. EPA does not endorse any products or commercial services mentioned in this publication. Our study was supported by Dr. Wang, Akin, and Dwivedy (CRISPR protocol), Schmidt, McAllister, Craft, Hoppenworth (sample collection and processing), Drs Whitaker, Chandrashekhar, and Suttenfield (*Pseudomonas aeruginosa* PA14). Roy J. Carver Biotechnology Center in their sequencing services, Drs Ceniceros, Frederick, and Wellman from Carle Health for providing clinical bacteria isolates, Drs. Cox, El-Naggar, Patel, and Patterson from Metropolitan Water Reclamation District of Greater Chicago for providing WWTP sewage samples.

References

1.   United Nations Environment Programme. *Bracing for Superbugs : Strengthening Environmental Action in the One Health Response to Antimicrobial Resistance.*; 2023.

2.   Dong C, Liu Z, Zhu L, et al. Dynamic migration and risk of cephalosporin antibiotic resistance genes: Move from pharmaceutical plants through wastewater treatment plants to coastal tidal flats sediments. *Water Res*. 2024;261:121983. doi:10.1016/J.WATRES.2024.121983

3.   Wang Z, Cai M, Du P, Li X. Wastewater surveillance for antibiotics and resistance genes in a river catchment: Spatiotemporal variations and the main drivers. *Water Res*. 2024;251:121090. doi:10.1016/J.WATRES.2023.121090

4.   Abia Akebe LK, Sibanda T, Selvarajan R, El-Liethy MA, Kamika I. Editorial: Environmental Reservoirs of Antibiotic Resistance Determinants: A Ticking Time Bomb for the Future Emergence of Super-Bugs of Environmental and Public Health Importance. *Front Environ Sci*. 2022;10:941847. doi:10.3389/FENVS.2022.941847/BIBTEX

5.   Larsson DGJ, Flach CF. Antibiotic resistance in the environment. *Nature Reviews Microbiology 2021 20:5*. 2021;20(5):257-269. doi:10.1038/s41579-021-00649-x

6.   Lin ZJ, Zhou ZC, Shuai XY, Shan XY, Zhou JY, Chen H. Deciphering Multidrug-Resistant Plasmids in Disinfection Residual Bacteria from a Wastewater Treatment Plant. *Environ Sci Technol*. 2024;58(15):6793-6803. doi:10.1021/ACS.EST.3C10895/ASSET/IMAGES/LARGE/ES3C10895_0005.JPEG

7.   Wang C, Yin X, Xu X, et al. Metagenomic absolute quantification of antibiotic resistance genes and virulence factor genes-carrying bacterial genomes in anaerobic digesters. *Water Res*. 2024;253:121258. doi:10.1016/J.WATRES.2024.121258

8.   Mao X, Yin X, Yang Y, et al. Standardization in global environmental antibiotic resistance genes (ARGs) surveillance. *Crit Rev Environ Sci Technol*. Published online 2024.

doi:10.1080/10643389.2024.2344453

9. Auerbach EA, Seyfried EE, McMahon KD. Tetracycline resistance genes in activated sludge wastewater treatment plants. *Water Res*. 2007;41(5):1143-1151. doi:10.1016/J.WATRES.2006.11.045

10. Liguori K, Keenum I, Davis BC, et al. Antimicrobial Resistance Monitoring of Water Environments: A Framework for Standardized Methods and Quality Control. *Environ Sci Technol*. 2022;56(13):9149-9160. doi:10.1021/ACS.EST.1C08918/ASSET/IMAGES/LARGE/ES1C08918_0005.JPEG

11. Debroas D, Siguret C. Viruses as key reservoirs of antibiotic resistance genes in the environment. *ISME J*. 2019;13(11):2856. doi:10.1038/S41396-019-0478-9

12. Alcock BP, Huynh W, Chalil R, et al. CARD 2023: expanded curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Res*. 2023;51(D1):D690-D699. doi:10.1093/NAR/GKAC920

13. Bengtsson-Palme J, Abramova A, Berendonk TU, et al. Towards monitoring of antimicrobial resistance in the environment: For what reasons, how to implement it, and what are the data needs? *Environ Int*. 2023;178:108089. doi:10.1016/J.ENVINT.2023.108089

14. Crossette E, Gumm J, Langenfeld K, Raskin L, Duhaime M, Wigginton K. Metagenomic quantification of genes with internal standards. *mBio*. 2021;12(1):1-12. doi:10.1128/MBIO.03173-20/ASSET/CE3BDE42-29BB-462F-9011-C09814CA41E0/ASSETS/GRAPHIC/MBIO.03173-20-F0003.JPEG

15. Ferreira C, Otani S, Aarestrup FM, Manaia CM. Quantitative PCR versus metagenomics for monitoring antibiotic resistance genes: balancing high sensitivity and broad coverage. *FEMS Microbes*. 2023;4. doi:10.1093/FEMSMC/XTAD008

16. Harshitha R, Arunraj DR. Real-time quantitative PCR: A tool for absolute and relative quantification. *Biochemistry and Molecular Biology Education*. 2021;49(5):800-812. doi:10.1002/BMB.21552

17. Ginzinger DG. Gene quantification using real-time quantitative PCR: An emerging technology hits the mainstream. *Exp Hematol*. 2002;30(6):503-512. doi:10.1016/S0301-472X(02)00806-8

18. Jacky L, Yurk D, Alvarado J, et al. Robust Multichannel Encoding for Highly Multiplexed Quantitative PCR. *Anal Chem*. 2021;93(9):4208-4216. doi:10.1021/ACS.ANALCHEM.0C04626/ASSET/IMAGES/LARGE/AC0C04626_0005.JPEG

19. Xie NG, Wang MX, Song P, et al. Designing highly multiplex PCR primer sets with Simulated Annealing Design using Dimer Likelihood Estimation (SADDLE). *Nature Communications 2022 13:1*. 2022;13(1):1-10. doi:10.1038/s41467-022-29500-4

20. Fassy J, Lacoux C, Leroy S, et al. Versatile and flexible microfluidic qPCR test for high-throughput SARS-CoV-2 and cellular response detection in nasopharyngeal swab samples. *PLoS One*. 2021;16(4):e0243333. doi:10.1371/JOURNAL.PONE.0243333

21. Lee SJ, Demeke T, Dusabenyagasani M, Saydak D, Perry D, Walkowiak S. Evaluation of two high-throughput genotyping systems for rapid identification of Canadian wheat varieties. *Canadian Journal of Plant Science*. 2023;103(4):422-425. doi:10.1139/CJPS-

669   2022-0192/SUPPL_FILE/CJPS-2022-0192SUPPLD.XLSX

670   22.   Olwagen CP, Adrian P V., Madhi SA. Performance of the Biomark HD real-time qPCR
671   System (Fluidigm) for the detection of nasopharyngeal bacterial pathogens and
672   Streptococcus pneumoniae typing. *Scientific Reports 2019 9:1*. 2019;9(1):1-11.
673   doi:10.1038/s41598-019-42846-y

674   23.   de Abreu VAC, Perdigão J, Almeida S. Metagenomic Approaches to Analyze
675   Antimicrobial   Resistance: An   Overview.   *Front   Genet*.   2020;11.
676   doi:10.3389/FGENE.2020.575592

677   24.   Guitor AK, Raphenya AR, Klunk J, et al. Capturing the Resistome: a Targeted Capture
678   Method To Reveal Antibiotic Resistance Determinants in Metagenomes. *Antimicrob
679   Agents Chemother*. 2020;64(1). doi:10.1128/AAC.01324-19

680   25.   Samorodnitsky E, Jewell BM, Hagopian R, et al. Evaluation of Hybridization Capture
681   Versus Amplicon-Based Methods for Whole-Exome Sequencing. *Hum Mutat*.
682   2015;36(9):903-914. doi:10.1002/HUMU.22825

683   26.   Sundararaman B, Sylvester MD, Kozyreva VK, Berrada ZL, Corbett-Detig RB, Green
684   RE. A hybridization target enrichment approach for pathogen genomics. *mBio*.
685   2023;14(5). doi:10.1128/MBIO.01889-23

686   27.   Gaudin M, Desnues C. Hybrid capture-based next generation sequencing and its
687   application to human infectious diseases. *Front Microbiol*. 2018;9(NOV):426578.
688   doi:10.3389/FMICB.2018.02924/BIBTEX

689   28.   Quan J, Langelier C, Kuchta A, et al. FLASH: a next-generation CRISPR diagnostic for
690   multiplexed detection of antimicrobial resistance sequences. *Nucleic Acids Res*.
691   2019;47(14):E83. doi:10.1093/NAR/GKZ418

692   29.   Gilpatrick T, Lee I, Graham JE, et al. Targeted nanopore sequencing with Cas9-guided
693   adapter   ligation.   *Nature   Biotechnology   2020   38:4*.   2020;38(4):433-438.
694   doi:10.1038/s41587-020-0407-5

695   30.   McCarty NS, Graham AE, Studená L, Ledesma-Amaro R. Multiplexed CRISPR
696   technologies for gene editing and transcriptional regulation. *Nature Communications
697   2020 11:1*. 2020;11(1):1-13. doi:10.1038/s41467-020-15053-x

698   31.   Liang X, Potter J, Kumar S, et al. Rapid and highly efficient mammalian cell engineering
699   via   Cas9   protein   transfection.   *J   Biotechnol*.   2015;208:44-53.
700   doi:10.1016/J.JBIOTEC.2015.04.024

701   32.   McCombie WR, McPherson JD. Future Promises and Concerns of Ubiquitous Next-
702   Generation   Sequencing.   *Cold   Spring   Harb   Perspect   Med*.   2019;9(9).
703   doi:10.1101/CSHPERSPECT.A025783

704   33.   Zhang Q, Ishii S. Improved simultaneous quantification of multiple waterborne
705   pathogens and fecal indicator bacteria with the use of a sample process control. *Water
706   Res*. 2018;137:193-200. doi:10.1016/J.WATRES.2018.03.023

707   34.   Grace A, Sahu R, Owen DR, Dennis VA. Pseudomonas aeruginosa reference strains
708   PAO1 and PA14: A genomic, phenotypic, and therapeutic review. *Front Microbiol*.
709   2022;13:1023523. doi:10.3389/FMICB.2022.1023523/BIBTEX

710   35.   BioProject: PRJNA1147862.   Accessed   August   13,   2024.

711   https://dataview.ncbi.nlm.nih.gov/object/PRJNA1147862?reviewer=2dbu2u54fitk5peh
712   oqct8omr2l

713   36.   Alcock BP, Raphenya AR, Lau TTY, et al. CARD 2020: antibiotic resistome surveillance
714   with the comprehensive antibiotic resistance database. *Nucleic Acids Res.*
715   2020;48(D1):D517-D525. doi:10.1093/NAR/GKZ935

716   37.   Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique
717   Molecular Identifiers to improve quantification accuracy. *Genome Res.* 2017;27(3):491-
718   499. doi:10.1101/GR.209601.116

719   38.   Ruby JG, Bellare P, DeRisi JL. PRICE: Software for the targeted assembly of
720   components of (Meta) genomic sequence data. *G3: Genes, Genomes, Genetics.*
721   2013;3(5):865-880. doi:10.1534/G3.113.005967/-/DC1/FILES1.FASTA

722   39.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature Methods*
723   *2012 9:4.* 2012;9(4):357-359. doi:10.1038/nmeth.1923

724   40.   Danecek P, Bonfield JK, Liddle J, et al. Twelve years of SAMtools and BCFtools.
725   *Gigascience.* 2021;10(2). doi:10.1093/GIGASCIENCE/GIAB008

726   41.   Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and
727   SAMtools.   *Bioinformatics.*   2009;25(16):2078-2079.
728   doi:10.1093/BIOINFORMATICS/BTP352

729   42.   BioProject: PRJNA1148079.   Accessed   August   13,   2024.
730   https://dataview.ncbi.nlm.nih.gov/object/PRJNA1148079?reviewer=p3cv3putr51hv68b
731   4nrbe332rk

732   43.   Chen B, Yuan K, Chen X, et al. Metagenomic Analysis Revealing Antibiotic Resistance
733   Genes (ARGs) and Their Genetic Compartments in the Tibetan Environment. *Environ*
734   *Sci   Technol.*   2016;50(13):6670-6679.
735   doi:10.1021/ACS.EST.6B00619/ASSET/IMAGES/LARGE/ES-2016-
736   006199_0006.JPEG

737   44.   Fang H, Wang H, Cai L, Yu Y. Prevalence of antibiotic resistance genes and bacterial
738   pathogens in long-term manured greenhouse soils as revealed by metagenomic survey.
739   *Environ   Sci   Technol.*   2015;49(2):1095-1104.
740   doi:10.1021/ES504157V/SUPPL_FILE/ES504157V_SI_001.PDF

741   45.   Li YM, Cao WW, Liang SL, et al. Metagenomic characterization of bacterial community
742   and antibiotic resistance genes in representative ready-to-eat food in southern China.
743   *Scientific Reports 2020 10:1.* 2020;10(1):1-14. doi:10.1038/s41598-020-72620-4

744   46.   Maboni G, Baptista R de P, Wireman J, Framst I, Summers AO, Sanchez S. Three
745   Distinct Annotation Platforms Differ in Detection of Antimicrobial Resistance Genes in
746   Long-Read, Short-Read, and Hybrid Sequences Derived from Total Genomic DNA or
747   from   Purified   Plasmid   DNA.   *Antibiotics.*   2022;11(10):1400.
748   doi:10.3390/ANTIBIOTICS11101400/S1

749   47.   Madden TL. Chapter 16: The BLAST Sequence Analysis Tool. Published online 2013.

750   48.   Kent WJ. BLAT--the BLAST-like alignment tool. *Genome Res.* 2002;12(4):656-664.
751   doi:10.1101/GR.229202

752   49.   Olson RD, Assaf R, Brettin T, et al. Introducing the Bacterial and Viral Bioinformatics

Resource Center (BV-BRC): a resource combining PATRIC, IRD and ViPR. *Nucleic Acids Res*. 2023;51(D1):D678-D689. doi:10.1093/NAR/GKAC1003

50. Han R, Shi Q, Wu S, et al. Dissemination of Carbapenemases (KPC, NDM, OXA-48, IMP, and VIM) Among Carbapenem-Resistant Enterobacteriaceae Isolated From Adult and Children Patients in China. *Front Cell Infect Microbiol*. 2020;10:535376. doi:10.3389/FCIMB.2020.00314/BIBTEX

51. McKenna M. Antibiotic resistance: The last resort. *Nature*. 2013;499(7459):394-396. doi:10.1038/499394A

52. Srathongneam T, Sresung M, Paisantham P, et al. High throughput qPCR unveils shared antibiotic resistance genes in tropical wastewater and river water. *Science of The Total Environment*. 2024;908:167867. doi:10.1016/J.SCITOTENV.2023.167867

53. Ghaly TM, Chow L, Asher AJ, Waldron LS, Gillings MR. Evolution of class 1 integrons: Mobilization and dispersal via food-borne bacteria. *PLoS One*. 2017;12(6):e0179169. doi:10.1371/JOURNAL.PONE.0179169

54. Garner E, Chen C, Xia K, et al. Metagenomic Characterization of Antibiotic Resistance Genes in Full-Scale Reclaimed Water Distribution Systems and Corresponding Potable Systems. *Environ Sci Technol*. 2018;52(11):6113-6125. doi:10.1021/ACS.EST.7B05419/SUPPL_FILE/ES7B05419_SI_004.XLSX

55. Yin X, Deng Y, Ma L, Wang Y, Chan LYL, Zhang T. Exploration of the antibiotic resistome in a wastewater treatment plant by a nine-year longitudinal metagenomic study. *Environ Int*. 2019;133:105270. doi:10.1016/J.ENVINT.2019.105270

56. Li W, Mao F, Ng C, et al. Population-based variations of a core resistome revealed by urban sewage metagenome surveillance. *Environ Int*. 2022;163:107185. doi:10.1016/J.ENVINT.2022.107185

57. Zhou Z, Xu L, Zhu L, et al. Metagenomic analysis of microbiota and antibiotic resistome in household activated carbon drinking water purifiers. *Environ Int*. 2021;148:106394. doi:10.1016/J.ENVINT.2021.106394

58. Davis BC, Brown C, Gupta S, et al. Recommendations for the use of metagenomics for routine monitoring of antibiotic resistance in wastewater and impacted aquatic environments. *Crit Rev Environ Sci Technol*. 2023;53(19):1731-1756. doi:10.1080/10643389.2023.2181620

59. Castanheira M, Simner PJ, Bradford PA. Extended-spectrum β-lactamases: an update on their characteristics, epidemiology and detection. *JAC Antimicrob Resist*. 2021;3(3). doi:10.1093/JACAMR/DLAB092

60. Pagani L, Mantengoli E, Migliavacca R, et al. Multifocal Detection of Multidrug-Resistant Pseudomonas aeruginosa Producing the PER-1 Extended-Spectrum β-Lactamase in Northern Italy. *J Clin Microbiol*. 2004;42(6):2523. doi:10.1128/JCM.42.6.2523-2529.2004

61. Gniadkowski M. Evolution of extended-spectrum beta-lactamases by mutation. *Clin Microbiol Infect*. 2008;14 Suppl 1(SUPPL. 1):11-32. doi:10.1111/J.1469-0691.2007.01854.X

62. Poirel L, Weldhagen GF, Naas T, De Champs C, Dove MG, Nordmann P. GES-2, a class A beta-lactamase from Pseudomonas aeruginosa with increased hydrolysis of

imipenem. *Antimicrob Agents Chemother*. 2001;45(9):2598-2603.
doi:10.1128/AAC.45.9.2598-2603.2001

63.   Diamond MB, Keshaviah A, Bento AI, et al. Wastewater surveillance of pathogens can
inform public health responses. *Nature Medicine 2022 28:10*. 2022;28(10):1992-1995.
doi:10.1038/s41591-022-01940-x

64.   Zaheer R, Noyes N, Ortega Polo R, et al. Impact of sequencing depth on the
characterization of the microbiome and resistome. *Scientific Reports 2018 8:1*.
2018;8(1):1-11. doi:10.1038/s41598-018-24280-8

65.   Gweon HS, Shaw LP, Swann J, et al. The impact of sequencing depth on the inferred
taxonomic composition and AMR gene content of metagenomic samples.
Environmental Microbiomes. 2019;14(1):1-15. doi:10.1186/S40793-019-0347-
1/FIGURES/7

66.   Pricing | Biotech. Accessed November 20, 2024.
https://biotech.illinois.edu/htdna/pricing

67.   Fuhrmeister ER, Kim S, Mairal SA, et al. Context-Seq: CRISPR-Cas9 Targeted
Nanopore Sequencing for Transmission Dynamics of Antimicrobial Resistance. *bioRxiv*.
Published online September 12, 2024. doi:10.1101/2024.09.12.612745

68.   Boyle EA, Becker WR, Bai HB, Chen JS, Doudna JA, Greenleaf WJ. Quantification of
Cas9 binding and cleavage across diverse guide sequences maps landscapes of target
engagement. *Sci Adv*. 2021;7(8).
doi:10.1126/SCIADV.ABE5496/SUPPL_FILE/ABE5496_TABLES_S1_TO_S19.XLSX

Enhanced detection for antibiotic resistance genes in wastewater samples using a CRISPR-enriched metagenomic method

Yuqing Mao[1,2], Joanna L. Shisler[3], Thanh H. Nguyen[1,2,4]

Affiliation

[1] The Grainger College of Engineering, Department of Civil and Environmental Engineering, University of Illinois Urbana-Champaign, IL, USA

[2] Carl R. Woese Institute for Genomic Biology, University of Illinois Urbana-Champaign, IL, USA

[3] Department of Microbiology, University of Illinois Urbana-Champaign, IL, USA

[4] Carle Illinois College of Medicine, University of Illinois Urbana-Champaign, IL, USA

Supplementary Information

Supplementary Methods:

S1. Wastewater sample collection, pre-treatment, and DNA extraction

The wastewater treatment plant influents were taken as 24-hour composites on weekdays. The manhole sewage sample was collected by an autosampler programmed to collect four-day composite sewage samples at four-hour intervals as described in a previous study (Oh et al., 2022). After collection, the water samples were sealed in sterile sample bags and kept in coolers filled with ice packs during transportation and stored at 4 ℃ for fewer than three days before filtering. Detailed information of the samples is listed in **Table S7**. Wastewater samples were shaken well and filtered through 47 mm mixed cellulose esters (MCE) membrane filters. After filtering, the filters were cut into quarters and stored at -80 ℃ before DNA extraction. DNA was extracted from each filter using the FastDNA SPIN Kit for Soil (MP Biomedicals), using the protocol provided by the manufacturer. The centrifugation step after bead beating was increased to 15 minutes to ensure a thorough separation of liquid and solid phases. The DNA extracts were eluted by 100 μL DES solution provided in the kit. The DNA samples were purified by OneStep PCR Inhibitor Removal Kit (Zymo Research). The DNA yield was determined by Qubit™ 1X dsDNA High Sensitivity (HS) Assay Kit (Invitrogen). Then, the DNA samples were stored at -20 ℃ before library preparation.

S2. Mock community DNA sample preparation and whole-genome sequencing

ZymoBIOMICS DNA Miniprep Kit (Zymo Research) was used for DNA extraction. Before DNA extraction, each bacterial strain was purified by streaking onto culture plates from their frozen stocks and incubated at 37 ℃ for 24 hours. Then, single colonies were picked from the plates and spiked into 900 μL of LB broth and incubated at 37 ℃ overnight for bacteria enrichment. The overnight cultures were centrifuged at 600x g for 5 minutes to separate bacteria cells from the liquid medium. The pellet was re-suspended by 750 μL of ZymoBIOMICS Lysis Solution and directly transferred to ZR BashingBead™ Lysis Tubes. Further DNA extraction occurred by following the manufacturer's protocol. The DNA was

eluted in 100 μL of ZymoBIOMICS™ DNase/RNase Free Water. The DNA yield was
determined by Qubit™ 1X dsDNA High Sensitivity (HS) Assay Kit (Invitrogen) immediately
after extraction. After mixing the DNA extracts of the 10 isolates, the mock community was
stored at -20 ℃ before library preparation.

GenFind V3 (Beckman Coulter) was used to extract high molecular weight DNA from
each isolate listed on **Table S1**. These DNA extracts were subjected to PacBio HiFi whole-
genome sequencing. Similar to above, a portion of frozen bacterial stock was streaked for
purification and incubated in liquid medium for enrichment using the same method as
mentioned in section 2.2. After the overnight incubation at 37 ℃ of the liquid culture, an
additional enrichment step was performed by transferring 100 μL of the overnight culture into
900 μL fresh LB broth and incubating at 37 ℃ for 2-3 hours to ensure the integrity of the
genome DNA. Each DNA sample was eluted by 200 μL 10 mM Tris-HCl (pH=8). The
concentrations and the lengths of the DNA extracts were determined by Qubit™ 1X dsDNA
High Sensitivity (HS) Assay Kit (Invitrogen) and 1% agarose gel electrophoresis. The DNA
extracts were stored at -80 ℃ before further analysis.

Before PacBio HiFi library preparation, all DNA extracts were sheared by Megarupter
3 (Diagenode) at speed 40 to obtain mean fragment lengths at 10 kbp for each sample. After
shearing, the sheared DNA samples were purified by OneStep PCR Inhibitor Removal Kit
(Zymo Research) followed by 1:1 beads cleaning using SMRTbell® cleanup beads (PacBio).
The fragment lengths of the sheared DNA samples were determined by 1% agarose gel
electrophoresis. Libraries were made using the SMRTbell prep kit 3.0 (PacBio). Qubit™ 1X
dsDNA High Sensitivity (HS) Assay Kit (Invitrogen) was used to determine the
concentrations of the sheared DNA samples and the libraries. The fragment lengths of the
libraries were determined by AATI Fragment Analyzer by Roy J. Carver Biotechnology
Center before sequencing. Sequencing was also conducted by Roy J. Carver Biotechnology
Center on SMRT Revio cell (PacBio) with 30-hour movie time.

The HiFi reads of the bacteria isolates were assembled into whole genomes by
Trycycler [1] using Canu, Flye, and Raven as the assemblers [2–4]. Then, the fasta files
containing all assembled contigs were uploaded to Bacterial and Viral Bioinformatics
Resource Center (BV-BRC) and annotated by the Genome Annotation service [5,6]. The
records of the ARGs identified in these genomes from the CARD source were downloaded
and labeled by their corresponding AROs using a customized Python code [7]. Due to the
difference in the IDs of the ARGs used in BV-BRC Genome Annotation and CARD
downloadable records, the remaining ARGs that were not automatically matched by the
customized code were BLASTed manually to compare the sequences and labeled by
corresponding AROs if the identity and coverage of the BLAST result were both greater than
95% [8]. The AROs of the ARGs that were not used for crRNA design were excluded. This list
of AROs was used as a reference to determine the true/false positive/negative in ARG
detection for CRISPR-NGS method.

S3. Design and synthesis of crRNA and tracrRNA

Multiplexed gRNA for ARG detection was designed by FLASHit [9]. Four thousand six
hundred and forty-one ARG nucleotide sequences for gRNA design input were downloaded
from the CARD protein homolog model on September 12, 2022 [7]. The default exclusion of
off-targets in the *E. coli* BL21 genome in FLASHit was removed, because the Cas9 protein
used in this study was commercially purchased. In total 6,010 different 20-nt spacer
sequences were generated by FLASHit to cleave target ARGs into approximately 200-bp
fragments.

For financial reasons, multiplexed gRNA was assembled using crRNA and tracrRNA
that were transcribed separately from corresponding DNA templates. The template for
tracrRNA was a double-stranded synthesized DNA fragment (5'-
AGGCGAATCAGATAATCGTTATGTCCAGACTGTATTAATACGACTCACTATAGGACAGC
ATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCGAGTCGGT
GCTTTTT-3') [9], and the template for multiplex crRNA was a single-stranded DNA oligo pool
of the following format (The 20 Ns correspond to the 20-nt spacer sequences generated by
FLASHit):

5'-
TAATACGACTCACTATAGNNNNNNNNNNNNNNNNNNNNGTTTTAGAGCTATGCTGTTTT
G-3'. The 20-nt spacer sequences in crRNA used in this study and their starting and ending
positions in all ARGs are listed in **Spreadsheet S7**.

A PCR step was performed to enrich both templates before transcription. Each PCR
reaction included 25 μL of Phusion® High-Fidelity PCR Master Mix with HF Buffer (New
England Biolabs), 1.25 μL of 10 mM forward (cr/tracr: 5'-TAATACGACTCACTATAG-3') and
reverse (cr: 5'-CAAAACAGCATAGCTCTAAAAC-3'; tracr: 5'-
AAAAGCACCGACTCGGTGCCAC-3') primers respectively, 4 μL of 10 ng/μL multiplexed
crRNA template or 1 ng/μL tracrRNA template, and filled up to 50 μL total volume using
molecular biology grade water [10]. The thermal cycle included an initial denaturation step at
98 ℃ for 10 seconds, 5 PCR cycles for crRNA template or 12 PCR cycles for tracrRNA
template with 98 ℃ denaturation for 5 seconds and annealing at 55 ℃ for 15 seconds, and a
final extension step at 72 ℃ for 1 minute. The PCR product was directly added to
TranscriptAid T7 High Yield Transcription Kit (Thermo Fisher) reaction as the DNA template
for RNA transcription following the manufacturer's instruction, followed by incubating at 37 ℃
for 4-6 hours until significant amount of white mist formed. After transcription, crRNA and
tracrRNA were purified by RNA Clean & Concentrator-5 with DNase I Set (Zymo Research
Corporation) following the total RNA clean-up protocol with DNase I treatment before RNA
clean-up. The DNase I treatment was increased to 30 minutes to ensure thorough removal
of residual DNA templates. The 100% ethanol was adjusted to 1.5 volume to increase short-
length RNA yield. The RNA product was eluted by 15 μL of DNase/RNase-free water. The
purified crRNA and tracrRNA were quantified by Qubit RNA Broad Range Assay Kit (Thermo
Fisher) using 100-fold diluted RNA products. The crRNA and tracrRNA were aliquoted and
stored separately at -80 ℃ before use [9].

S4. CRISPR-NGS library preparation

Instead of using non-specific enzymes for DNA fragmentation in most NGS library
preparation workflows, in the CRISPR-NGS library preparation workflow, we used Cas9
nuclease to specifically cleave targeted sites for DNA fragmentation. Right before CRISPR-
NGS library preparation, crRNA and tracrRNA were mixed equimolarly with the final
concentration adjusted by Duplex Buffer (Integrated DNA Technology) to 2,500 ng/μL. The
mixture was heated up to 94 ℃ for 2 minutes and slowly cooled down to room temperature
for gRNA assembly [11]. Then the duplexed gRNA was mixed with pre-diluted 25 μM Alt-R™
S.p. HiFi Cas9 Nuclease V3 (Integrated DNA Technologies) in a 10:3 volume ratio and
incubated at room temperature for 2.5 hours to form Cas9-gRNA complex.

During Cas9-gRNA complex formation, rAPid Alkaline Phosphatase (Roche) was
used to remove the 5' phosphate groups of the DNA samples. A 10 μL phosphatase reaction
included 100-300 ng of DNA samples, 1 μL of rAPid Alkaline Phosphatase Buffer (10x
concentrated), and 0.5 μL of rAPid Alkaline Phosphatase (1 U/μL). The mixture was

incubated at 37 ℃ for 10 minutes for 5' phosphate group removal followed by 75 ℃ for 2
minutes for phosphatase inactivation.

The Cas9 cleavage reaction included 3 µL of NEBuffer™ r3.1 (New England
Biolabs), 10.4 µL of the Cas9-gRNA complex, 1 µL of 0.01 ng/µL 5' phosphorylated
synthesized external 250-bp DNA spike with no cleavage site in its sequence (5'-
ACCCATACAAGGAACCCGGCCAGCACTACGCTCACTACGGCCGGTGGTACGGTGGGC
ACTCCGGTGAAATGCACGTGCTTGGCATGCCGTCAGGCCGTGAAGTCAAGCGCACCC
CGGTGTTCAACATGGACAGCAACAAGATGACCATCCACATCGCCTCGCCGGCGCCGG
CATACAGTCTGGGGGGAATTCAAGATGGAGAAGGGCGACGAGGTAATGGCGATCCTG
ACCTCGACAAGTGGAAGACCTG-3'), 6-8 µL of phosphatase treated DNA sample, and the
volume was filled up to 30 µL by molecular biology grade water. The mixture was pipette
mixed and incubated at 37 ℃ for 16 hours for thorough cleavage.

After the 16-hr incubation, 2.5 µL of RNase A (10 mg/mL) was directly added to the
mixture and incubated at room temperature for 15 minutes. Then, 2.5 µL of Thermolabile
Proteinase K (New England Biolabs) was added to each sample, and samples were
incubated at 37 ℃ for 15 minutes for protein digestion, and at 55 ℃ for 10 minutes for
Thermolabile Proteinase K inactivation. The DNA fragments were dA-tailed by Taq
polymerase. A master mix was premade by mixing 2 µL of dATP (100 mM), 5 µL of Taq
DNA polymerase (5,000 U/mL), 80 µL of Standard Taq Reaction Buffer (New England
Biolabs), and filled the volume up to 100 µL by molecular biology grade water. Five
microliters of the dA-tailing master mix were added to each reaction and incubated at 72 ℃
for 20 minutes. After dA-tailing, the cleaved DNA samples were ready for adapter ligation.

We used xGen™ UDI-UMI Adapters (Integrated DNA Technologies) in the adapter
ligation step to reduce the bias generated by the PCR step afterward. The adapter ligation
reaction included 35 µL of the dA-tailed DNA sample, 2 µL of 100-fold diluted adapter, 30 µL
of NEBNext® Ultra™ II Ligation Master Mix (New England Biolabs), and 1 µL of NEBNext®
Ligation Enhancer. The reaction was incubated at room temperature for 15 minutes. After
adapter ligation, 57 µL of AMPure XP Reagent (Beckman Coulter) was added to the reaction
for sample clean up. The adapter-ligated DNA samples were eluted by 15 µL 0.1x TE buffer
for PCR amplification.

The 50-µL PCR reaction included 15 µL of adapter-ligated DNA sample, 5 µL of
xGen™ Library Amplification Primer Mix (Integrated DNA Technologies), 25 µL of
NEBNext® Ultra™ II Q5® Master Mix (New England Biolabs), and 5 µL of molecular biology
grade water. The thermal cycle of PCR included an initial denaturation step at 98 ℃ for 30
seconds, 22 cycles of PCR step (98 ℃ for 10 seconds and 65 ℃ for 75 seconds), and a final
extension step at 65 ℃ for 5 minutes. The PCR product was cleaned by adding 45 µL of
AMPure XP Reagent (Beckman Coulter) into the 50 µL reaction. The cleaned PCR product
was eluted by 30 µL 0.1x TE buffer as the CRISPR-NGS library. The libraries were stored at
-20 ℃ before sequencing.

S5. Regular NGS library preparation, quality control, and Illumina sequencing

The regular NGS libraries were prepared using NEBNext® Ultra™ II FS DNA Library
Prep Kit for Illumina (New England Biolabs). The same amount of DNA samples as the input
of the CRISPR-NGS libraries were used to make the regular NGS libraries. One microliter of
0.01 ng/µL 5' phosphorylated synthesized external 250-bp DNA spike as mentioned in the
above section was added [12]. The library preparation steps followed manufacturer's
instructions with minor adjustments. In the fragmentation step, the samples were incubated
at 37 ℃ for 30 minutes. xGen™ UDI-UMI Adapters (Integrated DNA Technologies) was used
for adapter ligation, so that the USER® Enzyme incubation step was skipped. The adapter

dilution ratios were adjusted by the DNA input for each library (**Table S4**). Five cycles of
PCR were used in the PCR enrichment step. The final library was eluted in 30 μL of 0.1x TE
buffer and stored at -20 ℃ before sequencing.

The concentrations of the libraries were determined by Qubit™ 1X dsDNA High
Sensitivity (HS) Assay Kit (Invitrogen) right after library preparation. After concentration
check, the libraries were sent to Roy J. Carver Biotechnology Center for following quality
check steps and sequencing. DNA fragment lengths were determined by AATI Fragment
Analyzer. An additional beads purification was conducted before sequencing to remove
excess adapter dimers according to the fragment analysis results. A small portion of the
libraries were pooled and pre-sequenced on Illumina MiSeq V3 to determine the optimal
pooling ratio. Then, all libraries were pooled according to the MiSeq read depths and
sequenced on Illumina NovaSeq X Plus 10B paired-reads.

S6. Sequencing read mapping

After paired-end sequencing, the UMI indexes were extracted into a separated fastq
file. For each library, three fastq files were generated to store read 1, read 2, and the UMI
indexes. All sequencing raw data, including the separated UMI index files, can be found in
BioProject PRJNA1148079 (to be released upon publication) [13]. All fastq files were sorted by
the command "cat file.fastq | paste - - - - | sort -k1,1 -t " " | tr "\t" "\n" > file_sorted.fastq"
before processing. UMI-Tools was used to add UMI indexes from the separated UMI index
file to corresponding identifiers in read 1 and read 2 files using the command "umi_tools
extract --extract-method=string --bc-pattern=NNNNNNNNN -I [UMI_index_fastq_file] -S
[UMI_index_fastq_file_out_nothing_left] --read2-in=[read_fastq_file_to_add_UMI] --read2-
out=[output_read_fastq_file_with_UMI_added]" [14]. After adding UMI indexes, the sequencing
reads in read 1 and read 2 files were screened by PriceSeqFilter using the criteria that 85%
of nucleotides in a read must be high quality, minimum allowed probability of a nucleotide
being correct is 0.98, and 90% of nucleotides in a read that must be called [15]. The output
fastq files from PriceSeqFilter were used for read mapping in Bowtie2 with very sensitive
end-to-end mapping and reporting all alignments [16]. A fasta file including the sequences of
all ARGs downloaded from CARD for gRNA design (see 2.3) and the sequence of the 250-
bp external DNA spike was used as the reference. The output sam files were converted to
bam files by SAMtools using -bSF4 command to remove unaligned reads[17,18]. The bam files
were sorted and indexed by SAMtools before being deduplicated by UMI-Tools according to
the UMI indexes added to the read identifiers earlier [14]. The coverage and average read
depths of the mapping for all ARGs and the external DNA spike were exported by the
coverage function in SAMtools using the deduplicated bam file as the input with "--ff 0"
setting to remove the restriction on secondary alignment [17,18]. Any ARGs with coverages
higher than 75% were considered present in the sample, according to previous studies [19–21].
The identity threshold was approximately 90%, according to the end-to-end alignment
algorithm of Bowtie2 [16].

After the above steps, a validation step was conducted to check the reliability of read
mapping. In all ARG families that were detected by sewage CRISPR-NGS libraries, one
ARG from each family was randomly selected and viewed for these CRISPR libraries in
Integrative Genomics Viewer (IGV) [22]. Since DNA in CRISPR-NGS libraries were fragmented
by cleaving at designed sites and DNA in regular NGS libraries were fragmented randomly,
sharp boundaries of reads were expected at cleavage sites for CRISPR-NGS libraries while
the read mapping histograms were smooth for regular NGS libraries (**Figure S1**). Read
mapping was checked by comparing the cleavage sites shown in IGV to the bed file

generated after gRNA design indicating all possible cleavage sites for all ARGs.

S7. Validation for 5' phosphate group removal

A synthesized 731-bp DNA fragment with 5' phosphorylation was used for the
validation of 5' phosphate group removal by alkaline phosphatase. The synthesized DNA
was dissolved to a concentration of 10 ng/μL by molecular grade water to avoid inhibition to
the enzymes by EDTA. The 5' phosphate group removal reaction (10 μL) included 8.5 μL of
synthesized DNA, 1 μL of rAPid Alkaline Phosphatase Buffer (10x concentrated), and 0.5 μL
of rAPid Alkaline Phosphatase (1 U/μL). The mixture was incubated at 37 ℃ for 10 minutes
for 5' phosphate group removal followed by 75 ℃ for 2 minutes for phosphatase inactivation.
A negative control sample was made by replacing the alkaline phosphatase by the same
volume of molecular biology grade water. Then, both samples were used as the DNA input of
T4 ligation. The ligation reaction included 8.5 μL of either positive or negative DNA sample, 1
μL of T4 DNA Ligase Buffer (10X), and 0.5 μL of T4 DNA Ligase (New England Biolabs). The
mixture was incubated at room temperature for two hours for ligation. Then, the ligated DNA
samples were loaded to agarose gel for ligation results visualization. Phosphatase-treated
DNA did not have 5' phosphate groups, so that the DNA fragments would not ligate to form
new fragments. This sample would show a single band in gel. In contrast, the DNA sample
that was not treated by alkaline phosphatase should still have 5' phosphate group for
ligation. This sample would show multiple bands in gel.

S8. ARG clustering for quantification

Among the 4,641 ARGs used for gRNA design, some have highly similar sequences
with only a few mutations. One read may be mapped to multiple ARGs that share highly
similar sequences. To avoid the bias in ARG quantification due to the overlaps in read
mapping, we divided the 4,641 ARGs into 1,208 clusters based on their average nucleotide
identities (ANI). The pairwise ANIs for all ARGs were calculated by FastANI using fragment
length of 100 bp and minimum fraction of a sequence that must be shared to be 100% [23].
The FastANI output was screened by a customized python code to keep only the ANIs
above 90%. A network was built for the screened ANIs in Cytoscape [24]. The ARGs were
plotted as nodes. Two nodes were connected if the ANI between them was greater than
90%. The connected nodes were defined to be in the same cluster. If a node did not connect
to any other nodes, it was in an individual cluster. There were 228 clusters containing more
than one ARGs, and 980 clusters containing only one single ARG. The read depth of a
cluster was calculated by the average read depths of all ARGs in this cluster.

S9. Reproducibility test for CRISPR-NGS libraries

To test the reproducibility of CRISPR-NGS in ARG detection and quantification, the
libraries for Calumet (8/16/2023) and Calumet (10/24/2023) were made in duplicates. After
sequencing and read mapping, scatter plots were made in log-10 scale using the read
depths of the ARGs detected in both duplicated libraries as the x and y coordinates. Then,
the scatter plots were fitted linearly to get the correlation of the same ARGs detected in both
libraries. OriginPro 2024 was used to make the scatter plots and do the fitting.

S10. CRISPR-NGS read subsampling

To compare the ARG detection efficiency between CRISPR-NGS and regular NGS
with higher read depth, we subsampled the CRISPR-NGS fastq files by 1%, 2%, 5%, 10%,
20%, 50%, and 70% using seqtk (https://github.com/lh3/seqtk/tree/master) with the same
seed -s100 after reads sorting, UMI index labeling, and reads quality filtering as described in
section 2.5 in the main manuscript. After subsampling, all subsampled fastq files were
mapped to ARGs following the same protocol as described in section 2.5. After getting the
ARG mapping results of each individual subsampled CRISPR-NGS library, we calculated the

ratio between the number of ARGs detected by subsampled CRISPR-NGS libraries and the
corresponding regular NGS libraries. Then, these ARG detection ratios were plotted in
OriginPro 2024 against the subsampling ratio.

S11. Sanger sequencing for ARG presence validation

Sanger sequencing was conducted to validate the presence of the ARGs detected in
the DNA samples. The ARGs targeted in this validation step were randomly selected from
the ARG mapping results. The sequences of the selected ARGs were downloaded from
CARD [25]. Then, these sequences were used as the input for Primer BLAST on NCBI to
design primers for long-range PCR [26]. The minimum PCR product size was modified to be
less than 100 bp smaller than the size of the input ARG to ensure covering the largest
possible segment of the gene. The database used in Primer BLAST was nr, and the
organism was set as bacteria (taxid: 2). The long-range PCR primers are listed in **Table S2**.

For long-range PCR, we added 12.5 µL of Phusion® High-Fidelity PCR Master Mix
(New England Biolabs) or NEBNext® Ultra™ II Q5® Master Mix (New England Biolabs),
1.25 µL of 10 µM forward and reverse primers respectively, 1 µL of the DNA sample, and
filled the volume up to 25 µL by nuclease-free water for each reaction. The PCR thermal
cycle includes an initial denaturation step of 98 ℃ for 30 seconds, 34 cycles of the PCR step
with 10 seconds of denaturation at 98 ℃, 30 seconds of annealing at 55 ℃, and 90 seconds
of elongation at 72 ℃, and a final extension step at 72 ℃ for 5 minutes. The amplicon
lengths and DNA concentrations were validated by DNA electrophoresis and Qubit™ 1X
dsDNA High Sensitivity (HS) Assay Kit (Thermo Fisher) before Sanger sequencing. Sanger
sequencing was conducted by UIUC Core Sequencing Facility. Each DNA sequence was
aligned to the corresponding reference gene sequence in CARD by NCBI nucleotide BLAST
to test their similarity [8].

S12. qPCR for ARG quantification validation

Nine ARGs with various read depths detected by CRISPR-NGS or regular NGS
methods were selected for qPCR validation. The primers for these ARGs and their
corresponding amplification efficiencies are listed in **Table S5**. A 15-µL qPCR reaction
included 7.5 µL of PowerUp SYBR Master Mix (Applied Biosystems), 600 nM of forward and
reverse primers, 2 µL of 10-fold diluted sewage DNA extract, and 4.3 µL of molecular biology
grade water. The samples were loaded in triplicates in a qPCR run. Reactions targeting the
16S rRNA gene for the same samples were also loaded in triplicates on every qPCR plate
for relative abundance quantification. qPCR was performed on QuantStudio 7 Pro (Thermo
Fisher) using fast cycle followed by a melt curve stage for the quality control of non-specific
amplification. The qPCR thermal cycle included an initial hold stage at 95 ℃ for 20 seconds
and 40 PCR cycles with 95 ℃ for 1 second and 60 ℃ for 20 seconds. After the qPCR run,
the relative abundances of the ARGs were determined by ΔCq method using the following
equation [27]:

$$Relative\ abundance = 2^{Cq_{16S}-Cq_{ARG}}$$

In the equation, Cq_{16S} means the average Cq value of the 16S rRNA gene in the
sample, and Cq_{ARG} means the average Cq value of the ARG in the same sample. After
obtaining the qPCR results for the nine ARGs in all sewage samples, the read depths in both
CRISPR-NGS and regular NGS against the relative abundances in qPCR were plotted and
fitted as linear regression curves in log10 scale in Origin Pro 2024 to evaluate the ARG
quantification potential for the CRISPR-NGS method.

 Supplementary Tables:

 Table S1. Mock community composition

Isolate	Species/Genus
Environmental E. coli 1	*E. coli*
Environmental E. coli 2	*E. coli*
Environmental E. coli 3	*E. coli*
Clinical E. coli 1	*E. coli*
Clinical E. coli 2	*E. coli*
Clinical E. coli 3	*E. coli*
Clinical Shigella 1	*Shigella* spp.
Clinical Salmonella 1	*Salmonella* spp.
Clinical Salmonella 2	*Salmonella* spp.
PA14	*Pseudomonas aeruginosa*

    Table S2. The PCR primers designed for Sanger sequencing validation

#	Gene	Gene Length	Amplicon Length	Forward	Reverse
1	MCR-4.1	1626	1401	CACTGCGTTGTTTTAT GTTGCC	GCCAGCATTGGTAC GCTAGT
2	TEM-1	861	827	TTCGTGTCGCCCTTA TTCCC	TGAGGCACCTATCTC AGCGA
3	aac(3)-IId	861	810	TGCATACGCGGAAGG CAATA	TCGGAGTGGTTCCG AAATGC
4	tet(G)	1176	1101	ATCATTGCCCTGCTG ATCGT	GCCGACAATCCAAA CCCAAC
5	qnrB5	681	605	ACTCTGGCATTAGTT GGCGA	AACGATGCCTGGTA GCTGTC
6	GES-26	864	829	GCGCTTCATTCACGC ACTATT	AACTTGACCGACAGA GGCAA
7	aph(3")-Ib	804	780	GGTGAATCGCATTCT GACTGGTT	TCAACCCCAAGTCAG AGGGT
8	MCR-5.1	1644	1609	CGGTTGTCTGCATTT ATCACTTTCT	GAAGGTCCAACTCT GGCGT
9	sul1	840	833	GTGACGGTGTTCGGC ATTCT	GCATGATCTAACCCT CGGTCT
10	CRP	633	607	GACCCGACTCTCGAA TGGTT	AACGAGTGCCGTAA ACGACG
11	sul2	816	800	TCGCTCATCATTTTC GGCATC	ATTCTTGCGGTTTCT TTCAGCG
12	aph(3')-Ia	816	806	ATGAGCCATATTCAA CGGGAAAC	TCATCGAGCATCAAA TGAAACTGC
13	dfrA1	474	454	TGGTAGCTATATCGA AGAATGGAGT	CCCTTTTGCCAGATT TGGTAACTAT
14	MCR-5.1_latter_p art	1644	778	GACGGATACGGCTAC ATCCC	TCTGCATGTTGCCAG AAGGT
15	MCR-4.1_latter_p art	1626	504	TGGCAGTAGCACCAA GTCAA	GGCTCTGATAGACG GTGGTT

334 **Table S3.** Summary of positive and negative results in ARG detection using CRISPR-NGS
335 method for the mock community sample

ARG cluster	True	False
Positive	123 (97 from whole-genome sequencing, 25 from BLAST, 1 from Sanger sequencing)	1 (*aph(3')-Ia*)
Negative	NA	2 (identity < 90% in whole-genome sequencing annotation)

337 **Table S4.** Information for library preparation and sequencing

Method	Library	Adapter dilution ratio	DNA input (ng)	NH8B spike amount (ng)	Total number of reads
CRISPR-NGS	Mock community	1:100	123.8	8.66E-03	68,633,041
	Calumet 8/16/2023	1:100	108.0	5.40E-02	58,184,677
	Calumet 8/16/2023	1:100	115.2	8.06E-03	47,361,080
	Calumet 10/24/2023	1:100	111.6	4.40E-03	49,296,820
	Calumet 10/24/2023	1:100	111.6	4.40E-03	43,305,745
	Calumet 12/19/2023	1:100	110.4	6.86E-03	52,492,274
	Lemont 8/16/2023	1:100	121.2	6.86E-03	51,487,769
	Lemont 12/19/2023	1:100	136.8	6.86E-03	50,975,555
	Campus Town 11/30/2021	1:100	65.8	8.06E-03	54,097,785
Regular NGS	Mock community	1:1	122.9	7.60E-03	58,886,378
	Calumet 8/16/2023	1:1	108.0	1.11E-01	49,880,963
	Calumet 10/24/2023	1:1	178.5	8.40E-03	57,076,379
	Calumet 10/24/2023	1:1	178.5	8.40E-03	44,601,769
	Calumet 12/19/2023	1:1	166.4	7.60E-03	57,647,030
	Lemont 8/16/2023	1:1	161.2	7.60E-03	60,683,061

	Lemont 12/19/2023	1:1	200.6	7.60E-03	51,761,671
	Campus Town 11/30/2021	1:10	100.1	7.60E-03	46,173,438

Table S5. Information for primers used for qPCR

Gene	Forward	Reverse	Length	Efficiency (%)	Reference
dfrA	ACGGATCCTGGCTGTTGGTTGGACGC	CGGAATTCACCTTCCGGCTCGATGTC	237	89.4	[28]
sul1	CACCGTTGGCCTTCCTGTAA	TCTAACCCTCGGTCTCTGGC	180	98.4	[28]
sul2	GCGCTCAAGGCAGATGGCATT	GCGTTTGATACCGGCACCCGT	293	94.3	[28]
GES-26	CTTGAGAAGCTAGAGCGCGA	GCAGGAGACCATTCGACGAT	224	93	This study
MCR-5.1	CTTGCCCGTACCATTGACCT	ACAGGCTTGGTCGGCATAAA	198	101.3	This study
aph(3')-Ia	ACCGAGGCAGTTCCATAGGA	ACCTTATTTTTGACGAGGGGAAA	104	99.1	[29]
KPC	CACTGTGCAGCTCATTCAAGG	CGCCGATAGAGCGCATGAA	270	100.8	[30]
OXA-48	CTTAAACGGGCGAACCAAGC	GTTCATCCTTAACCACGCCC	94	101.9	[30]
CTX-M	CGGGCRATGGCGCARAC	TGCRCCGGTSGTATTGCC	105	87.3	[31]
16S	CCTACGGGAGGCAGCAG	ATTACCGCGGCTGCTGG	194	97.5	[32]

Table S6. The Cq values of the 16S rRNA gene in the six sewage samples

Sample	Cq
Calumet 8/16/2023	10.3
Calumet 10/24/2023	9.4
Calumet 12/19/2023	9.3
Lemont 8/16/2023	9.8
Lemont 12/19/2023	9.3
Campus Town 11/30/2021	12.1

344 **Table S7.** All sewage samples used in this study

Sample	Time	Location	Description	Volume (mL)
Calumet WWTP 1	8/16/2023	Chicago	Wastewater influent	50
Calumet WWTP 2	10/24/2023	Chicago	Wastewater influent	50
Calumet WWTP 3	12/19/2023	Chicago	Wastewater influent	50
Lemont WWTP 1	8/16/2023	Chicago	Wastewater influent	75
Lemont WWTP 2	12/19/2023	Chicago	Wastewater influent	50
Campus Town manhole	11/30/2021	Central Illinois	Raw sewage from a manhole	20

347 **Supplementary Figures:**

349 **Figure S1.** Validation of alkaline phosphatase blocking to remove 5' phosphate. a: The
350 concept figure of 5' phosphate removal validation using T4 ligase. b: The gel image of
351 blocked and unblocked samples after T4 ligase treatment.

Figure S2. The difference in read mapping between the CRISPR-NGS library (top) and the regular NGS library (bottom) using the example of OXA-36 (ARO: 3001430). The number axis at the top of the figure shows the scale of OXA-36 in base pairs (bp). The DNA fragments were cleaved at the four gRNA binding sites (85-108 bp, 296-319 bp, 517-540 bp, and 706-729 bp, labeled by orange dash lines) for the CRISPR-NGS library.

Figure S3. ARG clusters containing more than one ARGs.

Figure S4. The comparison of read mapping of *aadA13* (ARO: 3002613, 799 bp) (top) and *dfrA14* (ARO: 3002859, 475 bp) (bottom) in the mock community sample for the CRISPR-NGS library and the regular NGS library.

Figure S5. The read mapping of MCR-5.1 (ARO: 3004332) (a) and MCR-4.1 (ARO: 3004325)
(b) in the CRISPR-NGS library (yellow) and the regular NGS library (blue) of Lemont
(8/16/2023) sample.

Figure S6. The heatmap for the average read depths of all 270 ARG clusters in all sewage samples measured by CRISPR-NGS method and regular NGS method. The read depths are colored in log-10 scale. The read depths increase from green to pink. The ARG clusters are sorted by different drug classes. The drug classes are labeled by different colors on the left.

Supplementary Spreadsheets:

Spreadsheet S1. The clustering of different ARGs targeted in this study.

Spreadsheet S2. The ARGs detected in the mock community sample by CRISPR-NGS.

Spreadsheet S3. The ARG annotation results for the 10 PacBio HiFi whole-genome
sequences used in mock community.

Spreadsheet S4. The ARGs detected in the mock community sample by regular NGS.

Spreadsheet S5. The ARGs detected in all sewage samples by CRISPR-NGS.

Spreadsheet S6. The ARGs detected in all sewage samples by regular NGS.

Spreadsheet S7. The gRNA spacer sequences designed in this study.

References

1. Wick RR, Judd LM, Cerdeira LT, et al. Trycycler: consensus long-read assemblies for bacterial genomes. *Genome Biol*. 2021;22(1):1-17. doi:10.1186/S13059-021-02483-Z/FIGURES/4

2. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: Scalable and accurate long-read assembly via adaptive κ-mer weighting and repeat separation. *Genome Res*. 2017;27(5):722-736. doi:10.1101/GR.215087.116/-/DC1

3. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using repeat graphs. *Nature Biotechnology 2019 37:5*. 2019;37(5):540-546. doi:10.1038/s41587-019-0072-8

4. Vaser R, Šikić M. Time- and memory-efficient genome assembly with Raven. *Nature Computational Science 2021 1:5*. 2021;1(5):332-336. doi:10.1038/s43588-021-00073-4

5. Olson RD, Assaf R, Brettin T, et al. Introducing the Bacterial and Viral Bioinformatics Resource Center (BV-BRC): a resource combining PATRIC, IRD and ViPR. *Nucleic Acids Res*. 2023;51(D1):D678-D689. doi:10.1093/NAR/GKAC1003

6. Aziz RK, Bartels D, Best AA, et al. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics*. 2008;9(1):75. doi:10.1186/1471-2164-9-75

7. Alcock BP, Raphenya AR, Lau TTY, et al. CARD 2020: antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Res*. 2020;48(D1):D517-D525. doi:10.1093/NAR/GKZ935

8. McGinnis S, Madden TL. BLAST: at the core of a powerful and diverse set of sequence analysis tools. *Nucleic Acids Res*. 2004;32(Web Server issue):W20. doi:10.1093/NAR/GKH435

9. Quan J, Langelier C, Kuchta A, et al. FLASH: a next-generation CRISPR diagnostic for multiplexed detection of antimicrobial resistance sequences. *Nucleic Acids Res*. 2019;47(14):E83. doi:10.1093/NAR/GKZ418

10. Liang X, Potter J, Kumar S, et al. Rapid and highly efficient mammalian cell engineering via Cas9 protein transfection. *J Biotechnol*. 2015;208:44-53. doi:10.1016/J.JBIOTEC.2015.04.024

11. Miura H, Quadros RM, Gurumurthy CB, Ohtsuka M. Easi-CRISPR protocol for creating knock-in and conditional knockout mouse models using long ssDNA donors. *Nat Protoc*. 2018;13(1):195. doi:10.1038/NPROT.2017.153

12. Zhang Q, Ishii S. Improved simultaneous quantification of multiple waterborne pathogens and fecal indicator bacteria with the use of a sample process control. *Water Res*. 2018;137:193-200. doi:10.1016/J.WATRES.2018.03.023

13. BioProject: PRJNA1148079. Accessed August 13, 2024. https://dataview.ncbi.nlm.nih.gov/object/PRJNA1148079?reviewer=p3cv3putr51hv68b4nrbe332rk

14. Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique Molecular Identifiers to improve quantification accuracy. *Genome Res*. 2017;27(3):491-499. doi:10.1101/GR.209601.116

15.   Ruby JG, Bellare P, DeRisi JL. PRICE: Software for the targeted assembly of
components of (Meta) genomic sequence data. *G3: Genes, Genomes, Genetics*.
2013;3(5):865-880. doi:10.1534/G3.113.005967/-/DC1/FILES1.FASTA

16.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nature Methods*
2012 9:4. 2012;9(4):357-359. doi:10.1038/nmeth.1923

17.   Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and
SAMtools.   *Bioinformatics*.   2009;25(16):2078-2079.
doi:10.1093/BIOINFORMATICS/BTP352

18.   Danecek P, Bonfield JK, Liddle J, et al. Twelve years of SAMtools and BCFtools.
Gigascience. 2021;10(2). doi:10.1093/GIGASCIENCE/GIAB008

19.   Chen B, Yuan K, Chen X, et al. Metagenomic Analysis Revealing Antibiotic Resistance
Genes (ARGs) and Their Genetic Compartments in the Tibetan Environment. *Environ*
Sci   *Technol*.   2016;50(13):6670-6679.
doi:10.1021/ACS.EST.6B00619/ASSET/IMAGES/LARGE/ES-2016-
006199_0006.JPEG

20.   Fang H, Wang H, Cai L, Yu Y. Prevalence of antibiotic resistance genes and bacterial
pathogens in long-term manured greenhouse soils as revealed by metagenomic survey.
Environ   *Sci*   *Technol*.   2015;49(2):1095-1104.
doi:10.1021/ES504157V/SUPPL_FILE/ES504157V_SI_001.PDF

21.   Li YM, Cao WW, Liang SL, et al. Metagenomic characterization of bacterial community
and antibiotic resistance genes in representative ready-to-eat food in southern China.
Scientific Reports 2020 10:1. 2020;10(1):1-14. doi:10.1038/s41598-020-72620-4

22.   Robinson JT, Thorvaldsdóttir H, Winckler W, et al. Integrative Genomics Viewer. *Nat*
Biotechnol. 2011;29(1):24. doi:10.1038/NBT.1754

23.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nature*
Communications 2018 9:1. 2018;9(1):1-8. doi:10.1038/s41467-018-07641-9

24.   Shannon P, Markiel A, Ozier O, et al. Cytoscape: A Software Environment for Integrated
Models of Biomolecular Interaction Networks. *Genome Res*. 2003;13(11):2498.
doi:10.1101/GR.1239303

25.   Alcock BP, Huynh W, Chalil R, et al. CARD 2023: expanded curation, support for
machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance
Database. *Nucleic Acids Res*. 2023;51(D1):D690-D699. doi:10.1093/NAR/GKAC920

26.   Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL. Primer-BLAST: a
tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics*.
2012;13:134. doi:10.1186/1471-2105-13-134

27.   Rao X, Huang X, Zhou Z, Lin X. An improvement of the 2ˆ(–delta delta CT) method for
quantitative real-time polymerase chain reaction data analysis. *Biostat Bioinforma*
Biomath. 2013;3(3):71. Accessed July 10, 2024. /pmc/articles/PMC4280562/

28.   Le TH, Ng C, Chen H, et al. Occurrences and Characterization of Antibiotic-Resistant
Bacteria and Genetic Determinants of Hospital Wastewater in a Tropical Country.
Antimicrob Agents Chemother. 2016;60(12):7449. doi:10.1128/AAC.01556-16

29.   Rochegüe T, Haenni M, Cazeau G, et al. An inventory of 44 qPCR assays using
hydrolysis probes operating with a unique amplification condition for the detection and
quantification of antibiotic resistance genes. *Diagn Microbiol Infect Dis*.
2021;100(3):115328. doi:10.1016/J.DIAGMICROBIO.2021.115328

30.   Lund M, Petersen MB, Jørgensen AL, Paulmann D, Wang M. Rapid real-time PCR for
the detection of IMP, NDM, VIM, KPC and OXA-48 carbapenemase genes in isolates
and spiked stool samples. *Diagn Microbiol Infect Dis*. 2018;92(1):8-12.
doi:10.1016/J.DIAGMICROBIO.2018.04.002

31.   Roschanski N, Fischer J, Guerra B, Roesler U. Development of a Multiplex Real-Time
PCR for the Rapid Detection of the Predominant Beta-Lactamase Genes CTX-M, SHV,
TEM and CIT-Type AmpCs in Enterobacteriaceae. *PLoS One*. 2014;9(7):e100956.
doi:10.1371/JOURNAL.PONE.0100956

32.   Muyzer G, De Waal EC, Uitterlinden AG. Profiling of complex microbial populations by
denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified
genes coding for 16S rRNA. *Appl Environ Microbiol*. 1993;59(3):695-700.
doi:10.1128/AEM.59.3.695-700.1993