Assessing the effects of warming and carbonate chemistry parameters on marine microbes in the Gulf of Mexico through basin-scale DNA metabarcoding

Sean R. Anderson[1,2*], Katherine Silliman[4], Leticia Barbero[3,4], Fabian A. Gomez[7,8], Beth A. Stauffer[5], Astrid Schnetzer[6], Christopher R. Kelble[4,†], and Luke R. Thompson[4,7*]

[1]Department of Biological Sciences, University of New Hampshire, Durham, NH, USA

[2]Marine Chemistry and Geochemistry Department, Woods Hole Oceanographic Institution, Falmouth, MA, USA

[3]Cooperative Institute for Marine and Atmospheric Studies, University of Miami, Miami, FL, USA

[4]National Oceanic and Atmospheric Administration, Atlantic Oceanographic and Meteorological Laboratory, Ocean Chemistry and Ecosystems Division, Miami, FL, USA

[5]Department of Biology, University of Louisiana at Lafayette, Lafayette, LA, USA

[6]Marine, Earth, and Atmospheric Sciences, North Carolina State University, Raleigh, NC, USA

[7]Northern Gulf Institute, Mississippi State University, Starkville, MS, USA

[8]Physical Oceanography Division, Atlantic Oceanographic and Meteorological Laboratory, Miami, FL, USA

[†]National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Office of Science and Technology

*Corresponding authors: Sean R. Anderson and Luke R. Thompson

Emails: sean.anderson1@unh.edu and luke.thompson@noaa.gov

Keywords: Metabarcoding; Ocean Acidification; Warming; Microbes; Protists; Bacteria; Archaea; Gulf of Mexico; Generalized linear models; Indicator analysis

Abstract

Ocean acidification and warming threaten marine life, yet the impact of these processes on microbes remains unclear. Here, we performed basin-scale DNA metabarcoding of prokaryotes (16S V4–V5) and protists (18S V9) in the Gulf of Mexico and applied generalized linear models to reveal group-specific environmental correlates of functionally diverse microbes. Models supported prior physiological trends for some groups, like positive temperature effects on SAR11 and SAR86, and a positive effect of pH on *Prochlorococcus* that implied a negative response to decreasing pH. New insights were revealed for protists, like Syndiniales and Sagenista (e.g., positive pH effects), which offset positive relationships with temperature and reinforced the importance of considering multiple stressors simultaneously. Indicator analysis revealed phytoplankton, like *Ostreococcus* sp. and *Emiliania huxleyi*, that were associated with more acidic waters and may reflect candidate indicators of ocean change. Our findings highlight the need for sustained microbial sampling in marine systems, with implications for carbon export, nutrient cycling, and ecosystem health.

Introduction

Our ability to predict how marine ecosystems and resources will respond to future ocean conditions will require accurate monitoring of marine biodiversity over space, time, and across natural environmental gradients (*1*). The oceans are changing rapidly, heavily impacted by rising concentrations of human-derived atmospheric carbon dioxide (CO_2) that is absorbed at the ocean's surface (*2*). Atmospheric CO_2 has increased by nearly 50% (~420 ppm at present) over the last century, leading to increased levels of dissolved inorganic carbon (DIC) in the ocean, in turn lowering seawater pH (*3*). This process of ocean acidification (OA) reduces saturation states for carbonate minerals, placing stress on organisms that require these minerals for cellular growth and other functions (*4, 5*). The effects of OA are amplified by ocean warming, particularly at low latitudes, with surface temperatures expected to increase by 1–10 °C over the next century (*6*). Changes in seawater chemistry and physics can have immense impacts, both direct and indirect, on marine life (*3*). Thus, it is imperative to understand better how diverse marine organisms respond to present-day chemical and physical conditions to inform future potential shifts in community composition.

Over the past decade, research on species sensitivity to OA has expanded greatly, particularly for multicellular organisms that rely on carbonate chemistry for their structure and function (*3, 5*). Much less research has been conducted on marine microbes (i.e., protists, Bacteria, and Archaea), despite the central role of microbes in food webs and their strong influence on biogeochemical cycles and carbon export (*7, 8*). Microbes also respond quickly to shifts in their surrounding environment, making them potentially important indicators of changing ocean conditions (*9, 10*). In general, global ecosystem models predict a decline in photosynthetic biomass and a shift in composition from larger plankton (e.g., diatoms) to picophytoplankton

(0.2–2 µm), primarily driven by warming and enhanced stratification ([11–14]). Field and
laboratory experiments have measured direct and negative impacts of OA on plankton, notably
among calcifying haptophytes (e.g., coccolithophores), where increased partial pressure of CO_2
(pCO_2) and/or decreasing pH has led to reduced growth and calcification rates ([15–17]).
However, evidence suggests that some phytoplankton species, even coccolithophores, may be
resilient to rising pCO_2 and warming ([13], [18], [19]). In addition, heterotrophic bacteria may be
more resilient to OA compared to phytoplankton, impacted more directly by warming and
changes to phytoplankton-derived organic matter ([20], [21]). Employing DNA metabarcoding to
characterize the complex effects of OA parameters and temperature on a wide range of microbes
([22], [23]) will help guide lab-based experiments, identify indicator taxa, and inform model
predictions.
81
The Gulf of Mexico (GOM) is an ideal location to study the effects of multiple stressors on
marine microbes, as microbial communities in the GOM are affected by several major
hydrographic features that result in strong physicochemical gradients ([24]). The GOM is a semi-
enclosed subtropical basin, influenced by the Loop Current (and associated anticyclonic eddies)
and freshwater inflow from riverine systems (Mississippi-Atchafalaya) in the north ([25], [26]).
Most of the GOM is oligotrophic (and nutrient-limited), with phytoplankton biomass dominated
by picophytoplankton ([27]). Despite overall low biomass, microbial food webs in the GOM
support high biodiversity of mesozooplankton and micronekton ([28]), as well as several
economically important fisheries ([29]). At times, nutrient runoff from terrestrial sources promotes
eutrophication, resulting in coastal hypoxic zones that are more acidic ([30], [31]). Coastal
eutrophication combined with physical upwelling of new inorganic nutrients onto the shelf can
also enhance formation of harmful algal blooms (HABs), particularly along the western coast of
Florida ([32], [33]) and in other coastal regions in the southern Gulf ([34]). HABs pose a threat to
marine ecosystems in the GOM and can negatively impact local economies ([35]). While OA has
resulted in observable changes in ocean chemistry in the GOM ([24]), research on the impacts of
OA and warming on marine microbes has not been well explored. Most microbial genomics
studies have been localized to specific regions or depths ([36–39]) or focused on responses of
microbes to natural disturbances, like oil spills, in the northern Gulf ([40], [41]). This lack of spatial
biological sampling has made it difficult to characterize environmental drivers of diverse
microbes in the GOM ([24]), including OA parameters (e.g., pH, DIC, and pCO_2), and impedes
our ability to understand how microbial communities may shift in the future.
103
Here, we performed the first basin-scale DNA metabarcoding survey of protists, Bacteria, and
Archaea in the GOM as part of the fourth Gulf of Mexico Ecosystems and Carbon Cycle
(GOMECC-4) cruise that sailed from late summer to early fall of 2021. Overall, we collected
481 discrete DNA samples from 51 (out of 141) stations, encompassing 16 inshore–offshore
transects and up to three depths per site that corresponded to the surface, deep chlorophyll
maximum (DCM), and near bottom (Fig. 1A). Amplicon metabarcoding was performed to reveal
population dynamics of protists (18S SSU rRNA gene, V9 region) and prokaryotes (16S SSU

rRNA gene, V4–V5 region). We constructed generalized linear models (GLMs) for major
microbial groups in the photic zone to gain insight into group-specific environmental correlates,
including carbonate system parameters. These GLMs were applied to all GOMECC-4 sites,
including those where DNA samples were not collected, to expand spatial distributions of
microbial groups in the GOM. Finally, we performed indicator analysis based on profiles of DIC
and total alkalinity (TA) in the photic zone to identify microbes that were potential indicators of
more or less acidic waters (based on TA:DIC ratios). This study provides an important baseline
for microbial OA research in the GOM that will guide future DNA sampling efforts in this region
and contribute to our growing knowledge on the potential responses of marine microbes to
climate change.

Results and Discussion

Microbial population dynamics in the GOM

We obtained a total of 8,312 sequences on average per sample (range: 3,322–16,483) from 18S
metabarcoding, resulting in 13,632 protist amplicon sequence variants (ASVs) identified
throughout all GOM samples. In comparison, we obtained an average of 12,963 sequences per
sample (range: 5,056–28,620) for 16S metabarcoding which were assigned to 41,876 total
prokaryotic ASVs. Though significant to community composition ($P < 0.01$), factors like
transect, location on the shelf (< 200 m) vs. open ocean (> 200 m), and categorical depth had low
explanatory power on their own (PERMANOVA $R = 0.03–0.2$). As depth is a well-known driver
of global marine microbial communities (*42–46*), we performed hierarchical clustering of
microbial composition to better control for the impact of depth on subsequent spatial analyses.
This revealed separation of DNA samples into three clusters (Clusters 1–3), similar for both
marker gene regions, that reflected depth of samples in the water column on the continental shelf
and/or in open ocean GOM regions (Fig. 1B; fig. S1). For instance, Cluster 1 mainly consisted of
samples collected on the shelf at all depths and offshore at the surface layer, with all samples
located in the photic zone (2–99 m). Cluster 2 samples were mainly from the DCM (2–124 m) in
more stratified open ocean regions of the GOM, while Cluster 3 samples largely represented
meso- to bathypelagic waters (135–3,326 m) in the open ocean that were confined to the aphotic
zone (Fig. 2A–B; Fig. 3A–B). Though Clusters 1–2 were both in the photic zone (upper ~150
m), and had some overlap (Fig. 2B), they were separated into distinct clusters based on their
composition that reflected total depth in the water column and shifts in physicochemical
variables (fig. S2). In our case, clustering of DNA samples allowed us to better explore
microbes, and their relationships with environmental variables, within distinct spatial habitats
they occupy in the GOM.

Microbial communities in the GOM were more species-rich and diverse in the DCM and aphotic
zone (Fig. 2C; Fig. 3C), consistent with vertical profiles from other oceanic regions (*43, 46, 47*).
Higher richness and diversity with depth may be the result of microbes utilizing a broad

spectrum of sinking organic matter, exerting alternative metabolic strategies (redox reactions),
and/or forming diverse trophic relationships with other organisms to exploit such habitats (*48*).
Alpha diversity was stable along sampling transects in the photic zone for 16S (fig. S3) and 18S
samples (fig. S4), with higher variability in the DCM and aphotic zone. For example, microbial
diversity in the aphotic zone steadily decreased from coastal Florida (27°N line) to regions near
the Mississippi River outflow, increasing thereafter from Brownsville to Cancun (fig. S3).

Shifts in taxonomy between clusters were in line with depth-related microbial dynamics seen
previously in the GOM (*37, 38, 49*) and on a global scale (*42, 43, 48*). Among prokaryotes, the
photic zone and DCM were dominated by common heterotrophic bacteria, such as SAR11,
SAR86, and Flavobacteriales (Fig. 2D; fig. S5). Autotrophic cyanobacteria within the order
Synechococcales also had high relative abundance in the photic zone (Fig. 2D; fig. S5),
particularly *Prochlorococcus* and *Synechococcus* (fig. S6), both genera known to dominate
primary production in the GOM (*27*). Prokaryotic communities shifted dramatically in the
aphotic zone, with higher relative abundance of metabolically diverse taxa that are endemic to
deeper waters (*48–50*), including nitrous oxide-reducing Marinimicrobia (SAR406), ammonia-
oxidizing Nitrosopumilales, and sulfur-oxidizing Thiomicrospirales (Fig. 2D; fig. S5). These
microbes use redox reactions to acquire energy in less oxygenated waters (*48*), such as those
found in the mesopelagic zone (~200–800 m) in the GOM (fig. S2), and likely contributed to
increased richness of prokaryotic communities observed with depth (Fig. 2C). Certain 16S
groups varied at more resolved taxonomic levels between clusters. For example,
Prochlorococcus became more relatively abundant in the DCM, while SAR11 clade II increased
in the aphotic zone relative to other SAR11 clades (fig. S7). Similar patterns have been observed
elsewhere for *Prochlorococcus* (*51*) and SAR11 (*48, 52*), and reflect potential environmental
niche partitioning through the water column. High abundance of SAR11 clade II in the
mesopelagic has recently been observed in the Pacific Ocean (*48*), which may indicate particle
association among SAR11 that may be more common than previously thought.

Protist biodiversity was dominated by Dinophyceae, Syndiniales, Prymnesiophyceae, and
Sagenista in the photic zone and DCM, transitioning to Radiolaria (Polycystinea and RAD-B)
and Diplonemea in the aphotic zone (Fig. 3D). Dinophyceae and Prymnesiophyceae are common
in pelagic waters, including in the GOM (*36, 53*), and occupy important functional roles as
grazers (and mixotrophs) in microbial food webs (fig. S5). Sagenista was also abundant in the
photic zone (Fig. 3D), a group of common, yet still uncultured heterotrophic protists that have
important ecological roles (*54*). Other class level protist groups that were common in the GOM
in summer–fall, like Mamiellophyceae (Chlorophyta) and Mediophyceae (Stramenopiles), varied
more greatly across transects in the photic zone and DCM (Fig. 3D). Radiolarians dominated
relative abundance in mesopelagic samples (Fig. 3D). While these organisms remain largely
uncultivated and hard to study, they are key members of deep ocean food webs, forming
endosymbiotic relationships with other microorganisms (fig. S5) and contributing to the export
of carbon and biogenic silica (*55*).

193
DNA metabarcoding also reinforced the importance of obligate parasites within the group
Syndiniales at all depths in the water column (*42*, *56*, *57*), including at the basin scale in the
GOM (Fig. 3D; fig. S5). The prevalence of Syndiniales may be attributed to their wide host
range, active (and passive) export on sinking particles, and depth related niche partitioning (*56*,
58). We observed vertical shifts within Syndiniales at the clade level in our samples that aligned
with prior observations (*45*, *56*). For instance, there was a shift from Syndiniales Group-I Clades
1 and 4 in the photic zone to other clades, like Group-II Clade 7 and Group-I Clade 2 in the
aphotic zone (fig. S8). Radiolaria also varied between clusters, with certain members of
Polycystinea (e.g., *Heliosphaera* and *Pterocorys*) increasing in relative abundance from the
DCM to the aphotic zone (fig. S8). Diplonemea were also dominant in the GOM aphotic zone
(Fig. 3D). Though enigmatic, Diplonemea have been found globally in mesopelagic waters (*57*)
and likely represent important consumers of picoplankton and bacteria in these environments
(*59*).
207
208   **Generalized linear models reveal group-specific environmental correlates**
209
We used generalized linear models (GLMs) with either Poisson or negative-binomial error
distributions to identify potential explanatory variables of major 16S and 18S taxonomic groups
in the GOM. GLMs account for multiple predictor variables (factors) and have been applied to
ecological count (and proportional) data of higher trophic level marine organisms (*60*, *61*). Here,
we applied GLMs to microbial metabarcoding data, allowing us to observe predictor variables
and their relation to group-specific relative abundance measured spatially in the photic zone (Fig.
2D; Fig. 3D). We focused our models on the photic zone (Cluster 1), primarily because most
factors were collinear in the DCM and aphotic zone (table S1). Collinearity among variables can
result in models being less statistically reliable and confound model interpretation (*62*). Eight of
fifteen environmental variables were initially selected for models and included temperature,
salinity, dissolved oxygen (O_2), nitrate (NO_3), ammonium (NH_4), phosphate (PO_4), dissolved
inorganic carbon (DIC), and total pH recalculated to in situ temperatures (Table 1). Many
parameters related to OA that were measured or derived (e.g., total alkalinity, $p\text{CO}_2$, carbonate
ion concentration, and aragonite saturation) were strongly collinear to each other and
temperature (Spearman $r_s > 0.7$ or < -0.7), and thus were excluded from initial models (table
S1). Environmental conditions in the GOM surface were typical for this time of year (*63*, *64*).
For instance, offshore waters were warm (> 28 °C) and nutrient-limited (e.g., $NO_3 < 0.1$ μmol
kg^{-1}), while coastal regions had higher nutrient concentrations, including near the Mississippi
River outflow (Table 1; fig. S2). DIC was highest in the southern GOM and onto the Campeche
Bank (> 2050 μmol kg^{-1}), while pH often increased from the shelf to open ocean regions of the
Gulf (Table 1; fig. S2).
231
Microbial groups differed in the type and number of variables that significantly contributed to
the final models (Table 2). Pseudo R^2 values produced from GLMs ranged from 0.26–0.80

(Table 2), though several other methods confirmed appropriate model fit. First, model
simulations fit the data well (Fig. 4C; Fig. 5C) and the standardized residuals were normally
distributed for all groups (Kolmogorov–Smirnov, $P > 0.05$), except for Flavobacteriales. This
was further supported by significant and often strong positive correlations (Pearson $R = 0.31$–
0.90; $P < 0.01$) between test and model-trained relative abundance data for all groups (fig. S9;
fig. S10), with example plots shown for SAR11 (Fig. 4D) and Syndiniales (Fig. 5D).
Explanatory variables like temperature, DIC, and pH had individual, and often significant ($P <$
0.05) effects on relative abundance that varied among major 16S (Fig. 4A–B) and 18S groups
(Fig. 5A–B). Through this approach, we examined individual model terms, focusing primarily on
those related to ocean change, and explored their relationships with group-specific relative
abundances in the GOM.
245
Our model findings often supported prior physiological responses for certain microbial groups
that have been revealed in field and culture experiments. While changes in relative abundance
data with any given environmental factor does not necessarily translate to physiology, applying
DNA metabarcoding to OA research can help to verify existing trends and produce new
hypotheses for future testing on a wide range of microorganisms (*22, 23*). We found that
temperature had a positive effect on the relative abundance of SAR11 and SAR86 in our models
(Fig. 4A–B). Experimental evidence suggests that warmer conditions may favor increased
biomass of small, oligotrophic bacteria, like SAR11 and SAR86, that have low nucleic acid
content (*65*). In general, warming is thought to promote increased bacterial production, biomass,
and respiration, while also lowering growth gross efficiency (*20*). DIC had a positive effect on
SAR11, SAR86, and Flavobacteriales in group models (Fig. 4A–B), which together with
temperature effects, may imply a favorable response among these taxa to continued OA and
warming in this region. It is important to note that heterotrophic bacteria will also be influenced
by indirect changes in plankton composition, dissolved organic matter (DOM) availability and
quality, and trophic interactions (*21, 66*). These factors may outweigh direct OA effects in
natural communities and will be important to incorporate into future climate model predictions
of bacterial diversity and composition.
263
We observed contrasting effects of pH on the relative abundance of *Prochlorococcus* vs.
Synechococcus in the photic zone (Fig. 4A–B). For example, in situ pH had a strong and positive
effect on *Prochlorococcus*, implying a negative response to lower pH (more acidic) conditions.
An opposite trend was observed for pH in the *Synechococcus* model (Fig. 4A–B); however, DIC
also had a negative effect on *Synechococcus*, confounding model inference. Fu et al. (2007)
noted that combined effects of high $p\text{CO}_2$ and temperature significantly increased growth rates,
photosynthetic capacity, and cellular pigment levels of *Synechococcus* but not *Prochlorococcus*.
Mesocosm work in the subtropical North Atlantic also indicated a positive response of
Synechococcus to high $p\text{CO}_2$ (*67*), though others have noted small or insignificant physiological
shifts to changing conditions (*68*). On a global scale, ecological niche models predict increased
Prochlorococcus and *Synechococcus* biomass to ocean warming (via flow cytometry),

particularly in low latitude regions like the GOM, where these taxa already dominate plankton
biomass (*69*). Such niche models have not considered pH (or DIC), which we show may be
significant predictors. Though not resolved at the ecotype level in our amplicon dataset,
individual cyanobacterial strains or ecotypes will likely respond differently to future conditions,
as well as be influenced by indirect changes in top-down pressure (grazing or viral lysis),
nutrients, or sunlight (*70, 71*). Multiple ecotypes have already been discovered for
Prochlorococcus in the ocean, with evidence of different nutrient uptake rates, light preferences,
and thermal optima that shape population dynamics (*51, 72*). Additional field and laboratory
work is needed to identify responses among microbes at the species or ecotype level to support
accurate model predictions (*13*) and reveal underlying physiological mechanisms.
285
Future OA and warming is predicted to favor small phytoplankton, like picoeukaryotes, that can
more efficiently exploit oligotrophic and nutrient-limited waters (*11, 12, 14*), primarily due to
their larger cell surface to volume ratios that promote resource acquisition. Though for many
protists, the effects of OA and warming are less clear. This is especially true for Syndiniales and
Sagenista, enigmatic protist parasites and grazers that have seldom been considered with respect
to climate change. In our models, temperature had a significant and positive effect on the relative
abundance of Syndiniales and Sagenista (Fig. 5A–B). Temperature is often thought to enhance
physiological rates (*73*), which may include microzooplankton grazing and parasitism; however,
temperature relationships are hard to predict and can often be confounded by other factors, like
host or prey composition, that can dictate mortality rates. We found that pH had a positive effect
on Syndiniales and Sagenista, while DIC had a negative impact on Syndiniales relative
abundance (Fig. 5A–B). This implied a negative response among these groups to more acidic
conditions in the GOM at this time. Therefore, models that include only temperature or pH may
result in different outcomes for certain plankton groups (*74*), potentially misleading how we
interpret (and predict) their responses to climate change.
301
Dinophyceae were also prevalent in the photic zone on GOMECC-4 (Fig. 3D). It is well
understood that dinoflagellates are central to the microbial loop in oligotrophic regions, often
exhibiting mixotrophy and representing a key link between primary production and higher
trophic levels (*29, 53*). We found that temperature and DIC had significant and negative effects
on Dinophyceae relative abundance (Fig. 5A–B), implying a negative response to warmer and/or
more acidic conditions in the GOM. Similar findings on dinoflagellates have been observed in a
mesocosm study (*67*), though others have found dinoflagellates to benefit from or be less
sensitive to warming or increased pCO$_2$ concentrations (*23, 71, 74, 75*). Dinoflagellates often
exhibit mixotrophy, and so favorable responses to OA among this group may be indicative of
increased consumption of common prey (picoeukaryotes and cyanobacteria) that tend to grow
faster under such conditions (*75*). As is the case with many protists, dinoflagellates are extremely
diverse, not only phylogenetically but also in terms of their size, physiology, and trophic modes
(*76*). Therefore, it will likely be challenging to define a unified response for Dinophyceae to
changing ocean conditions. Future work that merges DNA metabarcoding with more targeted

approaches, like single-cell genomics or qPCR, will help to shed light on species sensitivity, interactions, and drivers that would otherwise be overlooked.

In addition to temperature and carbonate chemistry parameters, other factors like nutrients, salinity, and oxygen had significant effects on prominent 16S and 18S groups in the GOM (Fig. 4A; Fig. 5A). As an example, limiting nutrients like NO_3 and PO_4 had negative effects on the relative abundance of *Synechococcus* and *Prochlorococcus*, respectively (Fig. 4A). This may be related to the ability of cyanobacteria to uptake nutrients at low concentrations in surface waters (*72*). *Synechococcus* are thought to exploit low NO_3 concentrations in the GOM by maintaining a shallower distribution in the water column (*64*), relying on regenerated sources of NO_3 via nitrification (*29*). Salinity was also an important variable in our models, with positive effects on the relative abundance of *Prochlorococcus*, Syndiniales, Dinophyceae, and Prymnesiophyceae, as well as a negative effect on Flavobacteriales (Fig. 4A; Fig. 5A). Salinity is a known driver of bacterial and plankton distribution and diversity in the GOM (*37, 39, 77*). This is particularly evident in the northern GOM, where plankton biomass and composition are often driven by salinity-induced stratification (and nutrient availability) that result from riverine discharge via the Mississippi–Atchafalaya system, as well as by climatic processes, like the El Niño–Southern Oscillation (*77, 78*). Here, interpreting the role of salinity or nutrients in driving specific microbial groups was difficult, mainly because our sampling strategy and analysis focused on large-scale spatial patterns in microbial communities that did not allow us to explore regional trends (e.g., in the northern GOM). Even so, our results emphasize the importance of including such variables to resolve microbial composition and distribution at the basin scale in the GOM.

There are several caveats to consider with our model analysis. Models constructed from amplicon data on GOMECC-4 reflected only a specific time of the year (summer–fall) and did not integrate seasonal sampling. Temperature and carbonate chemistry parameters vary seasonally in the GOM (*79*), as does the intensity and position of the Loop Current (and eddies) and nutrient input from coastal runoff, all of which will impact microbial communities (*80, 81*). Consistent temporal sampling will be essential to better resolve microbes and their drivers over seasonal and interannual time scales (*24*). Such sustained sampling will also allow for more accurate predictions of microbial dynamics that integrate new OA data beyond the limits of GOMECC-4 measurements. We also considered GLMs for major taxonomic groups that were present in our samples (i.e., highest relative abundance), mainly to avoid issues with zero-inflation and overdispersion in the models. As a result, several groups thought to be sensitive to ocean change, like diatoms and diazotrophic cyanobacteria (*8*), were not considered here due to lower relative abundance at the basin scale. Similarly, this constrained our ability to predict model effects below the order to class level, with the exception being highly abundant cyanobacterial genera. We examined linear trends with GLMs as a simple and conservative approach to model relative abundance in the GOM. Future work may consider applying generalized additive models (GAMs) that allow for nonlinear dynamics (*82*), especially as more amplicon data is collected.

357
Though we tested for model fit, it is important to note that amplicon data is compositional, with
relative abundance of any single group being dependent on the proportion of others (*83*).
Comparing relative abundance among eukaryotic groups is also tenuous, as 18S rRNA gene copy
numbers can vary greatly (2–166 copies per cell) among protists (*84*). This is especially true for
alveolates (Syndiniales and Dinophyceae) and can lead to overestimation of read counts and
relative abundance (*42*). However, such concerns would not necessarily discount our modeling
approach that focused on groups separately and explored their relation to environmental factors.
Lastly, models did not account for trophic interactions (e.g., changes in prey or host) that may
vary along with changing conditions (*19*) or potential evolutionary adaptations among organisms
(*85*). Nevertheless, applying GLMs to amplicon data in this study offered a first step to define
multiple environmental drivers of diverse marine microbes, many of which are not easily
discerned with traditional observational methods like microscopy or cytometry. Our findings are
also timely for marine regions like the GOM that have lacked basin-scale sampling.
371
372 **GLMs expand microbial distributions in GOM surface waters**
373
Final models were used to predict the relative abundance of major 16S and 18S groups at 135
surface sites on GOMECC-4 (Fig. 1A), including 84 sites where DNA was not collected. This
allowed us to increase the spatial resolution of microbial sampling in the GOM at this time of
year. Groups like SAR11 (Fig. 6A) and Syndiniales (Fig. 6F) were well distributed throughout
the GOM, with highest relative abundance predicted offshore of Brownsville (Texas), in the Bay
of Campeche, and regions on the Campeche Bank. Cyanobacteria genera were largely partitioned
in the GOM based on their expected ecological niches (*86*). *Prochlorococcus* was most relatively
abundant offshore in stratified and nutrient-limited waters (Fig. 6B), particularly in parts of the
southern GOM. *Synechococcus* was present throughout the GOM at the surface, but relative
abundance was often highest in nutrient-rich coastal regions and in a localized area in the central
Gulf (Fig. 6C). Other groups like SAR86 (Fig. 6D), Prymnesiophyceae (Fig. 6H), and Sagenista
(Fig. 6I) were most relatively abundant offshore in the southern GOM and onto the East Mexico
Shelf, likely driven by higher temperature and DIC concentrations in these areas (fig. S2). This
was supported in the model output for these taxa, where temperature and/or DIC had positive
effects on relative abundance (Fig. 4A; Fig. 5A). Flavobacteriales was highest near the
Mississippi River outflow (Fig. 6E), in line with strong negative effects of salinity in the model
output for this group (Fig. 4A). The Mississippi River is the dominant source of freshwater into
the GOM, providing nutrients and organic matter into the system (fig. S2) that can stimulate
phytoplankton blooms (*81*). Though not widespread in the GOM, diatoms were most relatively
abundant in the photic zone near the Mississippi River (Fig. 3D), which may have contributed to
higher relative abundance of copiotrophs like Flavobacteriales that often associate with blooms
and can rapidly consume DOM (*87*).
396

Current predictions also revealed insights into the biogeography of a HAB species in the GOM.
Though prevalent through most of the GOM, including in the open ocean, relative abundance of
Dinophyceae was predicted to be highest directly off the coast of Tampa, Florida (Fig. 6G). This
was caused by a likely bloom event of the mixotrophic dinoflagellate, *Karenia brevis*, captured
in our DNA samples (fig. S11) and confirmed to be highly abundant through manual counts
(10^5–10^6 cells l^{-1}) estimated around the same time and location via the Florida Fish and Wildlife
Conservation Commission (https://myfwc.com/research/redtide/monitoring). HABs formed by
K. brevis are common in the GOM in the summer–fall, particularly along the West Florida Shelf,
and can negatively impact marine ecosystems and local economies (*32, 33, 35*). There is
evidence that warming may increase toxin production, growth rates, bloom frequency, and range
expansion of some HAB species (*88*). Temperature had negative effects on Dinophyceae in our
models (Fig. 5A–B), but responses were not explored to genus level. In culture, *K. brevis* has
shown increased growth rates with increasing pCO$_2$, though changes in toxin production were
not recorded (*89*). It remains important to monitor HABs and their drivers (*90*), combining
traditional monitoring and molecular methods to better predict pervasive blooms in the GOM
and elsewhere.
413
414 **Indicator analysis reveals candidate microbial indicator taxa of OA**
415
It is also important to determine specific microorganisms below the order to class level that may
be indicative of different OA conditions in natural waters (*9, 10, 23*). To this end, we grouped
samples in the photic zone (Cluster 1) based on TA:DIC ratios and examined microbial indicator
taxa at the ASV level. The TA:DIC ratio is a well-used proxy for carbonate chemistry in the
ocean, determining the buffering capacity against acidification (*79, 91*). In general, lower
TA:DIC ratios indicate poorly buffered waters, and so in our case, microbes that were more
prevalent in lower TA:DIC samples may be candidate indicators of more acidic conditions in the
GOM. TA:DIC ratios ranged from 1.1–1.2 in the photic zone, were not influenced by sampling
transect, and were positively correlated with pH (Pearson $R = 0.71$; $P < 0.01$) in surface waters
(Fig. 7A–B). TA:DIC ratios were manually grouped into low (< 1.16) vs. high (> 1.16)
categories to explore microbial indicators (Fig. 7A–B).
427
Overall, we found that 146 and 117 protist ASVs were significant indicators ($P < 0.001$) of low
or high TA:DIC ratios, respectively (table S2). Protist indicators spanned a range of taxonomic
groups, though several ASVs stood out (Fig. 7C). For instance, protists with the highest indicator
values (> 0.35) in samples with low TA:DIC ratios included *Ostreococcus* sp., which was the
most relatively abundant indicator ASV on average in the photic zone (~6%), as well as other
ASVs assigned to *Emiliania huxleyi* (now *Gephyrocapsa huxleyi*), Cryptomonadales,
Euduboscquella (Syndiniales), and Dino-Group I Clade 1 (Syndiniales). In comparison, ASVs
with high indicator values (> 0.55) in samples with high TA:DIC ratios consisted of ASVs
assigned to heterotrophic flagellates like MAST 3-B (and 4-B) and Kathablepharidida, as well as
parasites in Dino-Group III and V (Fig. 7D). For 16S samples, a total of 228 and 136 ASVs were

significant indicators ($P < 0.001$) of low vs. high TA:DIC ratios (table S2), dominated by
Proteobacteria (fig. S12). 16S ASVs with the highest indicator values (> 0.45) in low TA:DIC
samples included an ASV assigned to SAR11 clade Ia, which accounted for 10% of reads on
average in Cluster 1, as well as other ASVs assigned to SAR11 (clade I and Ia), SAR406, and
AEGEAN-169 (fig. S12). The 16S ASVs that were most indicative of high TA:DIC were
assigned to SAR116, SAR86, AEGEAN-169, and Rickettsiales (family S25-593; fig. S12).
444
Pico- and nanoeukaryotes dominate warm and oligotrophic regions like the open GOM (*27, 53,*
64) and are sensitive to changing ocean conditions (*15, 23*). Two of arguably the most well
studied taxa in the field of phytoplankton OA research, *Ostreococcus* sp. and *Emiliania huxleyi*,
were associated with less buffered (and more acidic) waters in the GOM (Fig. 7C). Both species
are widespread and impact global biogeochemical cycles (*92*), with *E. huxleyi* being a major
calcifier and contributor to $CaCO_3$ flux (*93*). In a prior 18S rRNA metabarcoding survey in the
southern GOM, *Ostreococcus* was the only genus with significantly different relative abundance
between upwelling and downwelling conditions in the DCM and when comparing the DCM to
mixed layer (*38*), which authors suggest may make this species an indicator of vertical nitrate
flux. Our findings imply *Ostreococcus* may also be a candidate indicator of acidic conditions in
GOM surface waters. Calcifying plankton, like *E. huxleyi*, are thought to be strongly impacted by
OA, with increased pCO_2 and/or lower pH having detrimental effects on growth and calcification
rates (*15, 16*). However, contrasting effects have been observed and may reflect considerable
strain and ecotype variability (*18, 23*). Indeed, several culture-based studies with *E. huxleyi* (and
Ostreococcus) have revealed adaptive mechanisms of cells to elevated pCO_2 over hundreds of
generations (*94, 95*). Though *E. huxleyi* was not prevalent overall in our samples (fig. S8), this
species has been measured in high concentrations ($\sim10^4$ cells l^{-1}) in the southern GOM in spring
(*96*). Together with model results at the class level (positive temperature effects on
Prymnesiophyceae), our findings highlight the potential sensitivity of haptophytes to changing
conditions in the GOM that should be further explored.
465
Indicator analysis also revealed SAR11, specifically ASVs assigned to clades 1 and 1a, as being
possible indicators of less buffered waters in the GOM in summer–fall (fig. S12). SAR11 is the
most abundant bacterial group in the oceans, playing an important role in global carbon cycling
(*97*). Though diverse, members of the SAR11 clade Ia ecotype tend to be most prevalent in
surface oceans (*52*), adapted to nutrient-poor conditions via small cell sizes and streamlined
genomes (*98*). Though direct effects of OA on SAR11 remain unclear and are likely to be less
important compared to shifts in DOM (*21*), SAR11 exhibits known seasonality in the surface
oceans and is sensitive to temperature (*48, 65*). Such temperature sensitivity was supported in
our model analysis (Fig. 4A–B). Given the ubiquity of SAR11 and its role in global carbon
cycles, it remains critical to confirm and further investigate the potential of this group as an
indicator of ocean change.
477

Future sampling to characterize microbes in changing oceans

Efforts to characterize microbial communities over natural physicochemical gradients are essential to inform how these communities may shift in the face of changing ocean conditions (*10*). In the GOM, there is evidence of increased pCO$_2$ in many parts of the open ocean that are on par with rates of change in other oligotrophic regions (*99*), like those measured in the Pacific Ocean via the Hawaiian Ocean Time-series (1.72 µatm yr^{-1}) and in the Atlantic Ocean via the Bermuda Atlantic Time-series Study (1.69 µatm yr^{-1}). Yet, knowledge on the effects of OA and warming on biological organisms is limited in the GOM, particularly for microbes. Here, we performed the first basin-scale DNA metabarcoding survey in the GOM and paired this with extensive hydrographic, nutrient, and carbonate chemistry measurements to investigate diverse prokaryotes and protists and their specific environmental drivers (Fig. 1). In line with prior physiological and modeling-based observations, our GOM model analyses suggest that more acidic and warmer conditions in the GOM may favor heterotrophic bacteria (SAR11 and SAR86) and smaller phytoplankton (e.g., Prymnesiophyceae), with groups like Dinophyceae potentially being less favored in future conditions (Fig. 4; Fig. 5). Warming and OA in the GOM may have contrasting effects on major plankton parasites (Syndiniales) and grazers (Sagenista) that are seldom considered with respect to climate change and underscores the importance to measure multiple stressors simultaneously (Fig. 5). We also defined microbial indicator taxa at the ASV level (Fig. 7), which resulted in several ubiquitous (and environmentally sensitive) microbes, like *Ostreococcus* sp., *Emiliania huxleyi*, and SAR11 clade Ia, being associated with more acidic waters in the GOM. Model inference and the utility of identified ASVs to act as indicator species of OA will need to be further tested, including at different times of the year to reflect seasonal turnover of the microbial community.

Though still unclear, empirical and predictive work suggests that changes in our ocean systems will likely have profound impacts on microbial composition, biogeography, and physiology (*8, 100*), with consequences for trophic transfer, nutrient cycling, and carbon export. Global models and experimental evidence predict increased stratification with warming, shifting communities to smaller picophytoplankton that can better exploit nutrients and other resources (*12, 14*). Warming-induced stratification may also result in an overall net reduction in carbon export that may threaten to decrease the amount of organic carbon that reaches the seafloor (*101*). Yet, predicted shifts in carbon export in global ecosystem models remains uncertain, ranging from a 41% decrease to 8% increase in carbon export flux in future oceans (*102*). Strong selection imposed by climate change may also drive rapid adaptation, competition, or the emergence of new species (e.g., with higher thermal tolerance), all restructuring microbial communities (*8, 13, 85*). In culture, some microbes demonstrate the ability to adapt to warmer or more acidic conditions (*94, 95*), though this does not necessarily mean they will remain competitive and it remains an open question on how this will apply to natural systems with mixed assemblages (*19, 85*). Further, current models do not fully account for trophic interactions, like grazing or parasitism, the rates of which will likely vary in future oceans and offset direct physiological

519 effects of OA or warming on certain microbes. It remains important to measure microbial
520 interactions, plankton mortality rates, and carbon export rates over time and in space (*10, 102*),
521 which will support a more mechanistic approach to model predictions.
522
Our findings provide an important baseline for microbial OA research in the GOM; however,
sampling on GOMECC-4 only reflected a single time of the year and did not consider known
seasonal variability in carbonate chemistry parameters or hydrography (*79, 80*), which are likely
to influence microbes and their drivers (*36*). In response, there is a need for sustained biological
measurements in the GOM, either by establishing long-term monitoring programs or continuing
to leverage existing oceanographic surveys, like GOMECC. Long term microbial sampling in the
GOM will be essential to accurately predict future changes in microbial groups that may be
expected with continued OA or warming. For example, increased DNA collection would support
ecosystem modeling of microbes in the GOM, integrating climate model scenarios (e.g., via the
Coupled Model Intercomparison Project) to predict shifts in microbial abundance by the end of
the century. Ultimately, our ability to predict the response of marine microbes to climate change
will depend on sustained and coordinated sampling efforts across a range of dynamic marine
ecosystems.
536
537 **Materials and Methods**
538
Seawater collection, DNA filtration, and environmental metadata
539
Seawater was collected on board the NOAA Ship *Ronald H. Brown* as part of GOMECC-4 from
September 13 to October 21, 2021. Sampling for GOMECC-4 occurred along 16 inshore–
offshore transects across the entire GOM and an additional line at 27°N latitude in the Atlantic
Ocean (Fig. 1A). Sampling started at the 27°N line and continued counterclockwise across the
GOM, ending at Florida Straits and Cape Coral. We also collected DNA samples near Padre
Island National Seashore (U.S. National Parks Service), a barrier island located off the coast of
south Texas (Fig. 1A). Vertical CTD sampling was employed at each site to measure discrete
physical, chemical, and biological properties. Water sampling for DNA filtration was conducted
at 51 out of 141 total sites and three depths per site, representing the surface, deep chlorophyll
maximum (DCM), and near bottom (fig. S1).
550
At each respective site and depth, seawater was collected from pre-designated Niskin bottles on a
CTD rosette. To ensure adequate amounts of water were filtered for DNA analysis, samples for
chemical parameters were taken at the same depths but with different discrete Niskin bottles.
Following a CTD cast, which varied in duration from 30 min to 3 h depending on water depth,
whole seawater was transferred from Niskin bottles to triplicate Whirl-Pak bags (3 depths x 3
replicates = 9 bags per site). Within an hour, whole seawater (~2 L per replicate) was filtered
through 0.22-μm Sterivex filters (Millipore; CAT# SVGP01050) via a peristaltic pump (100–150
rpm) and run dry. Filters were capped and outlets were sealed with parafilm. Filters were stored

at –80 °C on the ship and kept at the same temperature in the laboratory for longer-term storage. Filter lines were sterilized with 2% bleach and rinsed with Milli-Q after every site. Milli-Q blanks were also filtered randomly throughout the duration of the cruise. Accounting for blanks and replication, a total of 481 Sterivex filters were collected on GOMECC-4.

Discrete samples for water column hydrography and chemistry were taken at each site and depth during GOMECC-4, including sites sampled for DNA. Temperature, salinity, pressure, and chlorophyll fluorescence were obtained from the CTD. Vertical CTD profiles on the downcast were used to estimate the position of the DCM at each site. Blanks and quality control samples were considered for each discrete chemical parameter. Dissolved oxygen concentration was estimated from water samples (125 ml) using an automated oxygen titrator with amperometric end-point detection (*103*). Nutrient samples were collected from Niskin bottles into 50-ml acid washed bottles. Dissolved nutrients (NO_3, NO_2, NH_4, PO_4, and SiO_4) were measured on board using an automated continuous flow analytical system with colorimetric detection ((*104*); SEAL Analytical). Samples for DIC were collected from Niskin bottles into 294-ml borosilicate glass bottles, sealed with glass stoppers, and stored for 12 h at room temperature. DIC samples were analyzed on the ship using two analytical systems, each consisting of a coulometer (CM5017, UIC Inc.) coupled with a Dissolved Inorganic Carbon Extractor (*105*).

Samples for total alkalinity (TA) were collected from Niskin bottles into 500-ml collection bottles, preserved with a mercuric chloride solution, and kept in a water bath at 22 ºC for 1 h prior to analysis. TA measurements were made using a two-titration system, consisting of a Metrohm 765 or 665 Dosimat Titrator and Orion 720A or 2-Star pH meter (*106*). Samples for pCO$_2$ were drawn from Niskin bottles into 500-ml glass bottles, preserved with mercuric chloride, and stored at room temperature for 8 h before analysis. Details on the system used to measure pCO$_2$ are described in (*107*) and include equilibrating each sample with a constantly circulating gas phase. Lastly, for pH analysis, samples were collected from Niskin bottles into 10-cm (~30 ml) glass cylindrical optical cells and analyzed on an Agilent 8453 spectrophotometer with a custom-made temperature-controlled cell holder (*108*). Aragonite saturation state was calculated at each site and depth based on temperature, salinity, pressure, DIC, and TA using the CO2SYS program for CO_2 System Calculations (*109*). Measurements of pCO$_2$ (20 ºC) and pH (25 ºC) were re-calculated to in situ conditions using pressure, temperature, salinity, DIC, and TA in CO2SYS (*109*). Environmental metadata associated with DNA samples are provided in table S3.

DNA extractions, PCRs, and library preparations

Sterivex filters were extracted in-house at NOAA's Atlantic Oceanographic and Meteorological Laboratory (AOML) using the ZymoBIOMICS 96 DNA/RNA MagBead kit (Zymo; CAT# D4308), with modifications for in-cartridge bead beating as described in (*110*). Filters were thawed, the inlet caps were removed, and excess water was dried from the inlet using kimwipes

to allow for dispensing of beads into the cartridge. Premade mixtures of 0.1 mm and 0.5 mm
beads were directly added into the filters to ensure adequate lysis and recovery of hard-to-lyse
phytoplankton groups (*110*). This was followed by the addition of a lysis buffer (1 ml). Sterivex
filters were vortexed for 40 min on a Vortex-Genie at maximum speed (~3200 rpm). DNA
lysates were transferred to 2-ml LoBind tubes (Eppendorf) via syringe and centrifuged for 1 min
at 10,000 g. Supernatant (750 µl per sample) was split across three KingFisher 96-well plates
(250 µl per plate). Zymo MagBinding buffer (600 µl) and magnetic beads (25 µl) were added to
each well in each of the three plates. With this setup, 96 samples were extracted at the same time
on the automated KingFisher Flex (Thermo Fisher). Each run included three wash plates with
500–900 µl per well of MagWash and an elution plate with 150 µl per well of molecular-grade
water. DNA was eluted into a single well from the same discrete sample across replicate plates.
Concentrations of eluted DNA were measured using a Varioskan LUX plate reader and the
Quant-IT dsDNA Assay (Thermo Fisher) and corrected per replicate sample based on volume of
seawater filtered (ng l^{-1}; fig. S13). Filters were processed randomly. Extraction blanks (clean
Sterivex filters) were also included and processed similarly. A bacterial mock community
(Zymo) was included as a positive control.
616
Metabarcoding libraries were initially prepared at AOML, amplifying DNA of target organisms
with universal primers, including 16S (Bacteria and Archaea) and 18S rRNA (protists). Primers
from (*111*) were used to target the 16S V4–V5 region: forward (515f; 5'-
GTGYCAGCMGCCGCGGTAA-3') and reverse (926r; 5'-CCGYCAATTYMTTTRAGTTT-
3'). Primers from (*112*) and the Earth Microbiome Project
(http://www.earthmicrobiome.org/emp-standard-protocols/18s/) targeted the 18S V9 region:
forward (1391f; 5'-GTACACACCGCCCGTC-3') and reverse (EukBr; 5'-
TGATCCTTCTGCAGGTTCACCTAC-3'). Primers were constructed with Fluidigm common
oligos CS1 forward (CS1-TS-F: 5'-ACACTGACGACATGGTTCTACA-3') and CS2 reverse
(CS2-TS-R: 5'-TACGGTAGCAGAGACTTGGTCT-3') fused to their 5' ends, to enable two-
step library preparation at the Michigan State University Research Technology Support Facility
(RTSF).
629
PCR reactions were run in triplicate (12.5 µl per sample), with 1 µl of DNA per sample. 16S
PCR reactions consisted of 5 µl of AmpliTaq Gold, 6.25 µl of water, and 0.375 µl of each primer
(10 µM); PCR conditions included denaturation at 95 °C for 2 min, 25 cycles of 95 °C for 45 s,
50 °C for 45 s, and 68 °C for 90 s, followed by a final elongation step of 68 °C for 5 min (*111*).
18S PCR reactions consisted of 5 µl of AmpliTaq Gold, 6.5 µl of water, and 0.25 µl of each
primer (10 µM); PCR reactions involved denaturation at 94 °C for 3 min, 35 cycles of 94 °C for
45 s, 65 °C for 15 s, 57 °C for 30 s, and 72 °C for 90 s, followed by a final elongation step of 72
°C for 10 min (*112*). PCR products were pooled and run on a 2% agarose gel to confirm
amplification of target bands. Sample plates were submitted to the Michigan State University
RTSF Genomics Core for secondary PCR and sequencing.
640

Secondary PCR used dual-indexed, Illumina-compatible primers, targeting the Fluidigm
CS1/CS2 oligomers at the ends of the PCR products. PCR conditions for the secondary run
included an initial denaturation step at 95 °C for 3 min, 11 cycles of 95 °C for 15 s, 60 °C for 30
s, and 72 °C for 60 s, followed by elongation at 72 °C for 3 min. Amplicons were batch
normalized using Invitrogen SequalPrep DNA Normalization plates and the recovered product
was pooled. The pool was QC'd and quantified using a combination of Qubit dsDNA HS,
Agilent 4200 TapeStation HS DNA1000, and Invitrogen Collibri Library Quantification qPCR
assays. The RTSF Core included a sequencing blank for each sample plate. Separate sequencing
runs were performed using an Illumina MiSeq (2 × 250 bp) for 18S and 16S samples. Custom
sequencing and index primers complementary to the Fluidigm CS1 and CS2 oligomers were
added to appropriate wells of the reagent cartridge. Base calling was done by Illumina Real Time
Analysis (RTA) v1.18.54 and output of RTA was demultiplexed and converted to FASTQ
format with Illumina Bcl2fastq v2.20.0.
654
655   **Bioinformatics and functional assignments**
656
Primers were removed from demultiplexed FASTQ sequences using Cutadapt (*113*). Trimmed
reads were processed in Tourmaline, which implements QIIME 2 (and DADA2 plugins) in a
Snakemake workflow (*114*). Paired-end DADA2 was used to infer 16S and 18S amplicon
sequence variants or ASVs (*115*). Taxonomic assignments were also performed in Tourmaline
using reference files from SILVA (Version 138.1; (*116*)) and the Protistan Ribosomal Reference
or PR2 (Version 5.0.1; (*117*)) databases for 16S and 18S ASVs, respectively. In both cases,
taxonomy was assigned using a Naïve Bayes classifier trained to the respective databases and
trimmed to the primer regions (*118*). Output files (taxonomy, count, and metadata) were
imported separately into R (Version 4.3.1) using qiime2R (Version 0.99.6;
https://github.com/jbisanz/qiime2R) and merged with phyloseq (Version 1.44.0; (*119*)). Several
groups were removed from the 18S dataset: Metazoa, Streptophyta, Rhodophyta, and unassigned
reads at the Subdivision level. 18S reads assigned to non-marine taxa, e.g., Insecta, Archosauria,
and Ascomycota were also filtered out. For 16S, reads assigned to Chloroplast, Mitochondria,
and Eukaryota were removed. Samples with less than 3,000 reads counts were filtered out for
18S (5,000 reads for 16S), along with ASVs only observed once in each respective dataset.
Species accumulation curves were generated for 18S and 16S samples using the R package
ranacapa (Version 0.1.0; (*120*)). The number of reads vs. ASVs was saturated with respect to
categorial depth and position of samples on the shelf vs. open GOM, indicating that an
appropriate sequencing depth was reached (fig. S14). Samples were rarefied to the minimum
read count to normalize for differences in library size.
677
Protist ASVs were manually assigned to functional groups based on 18S V9 functional
annotations (https://doi.org/10.5281/zenodo.3768950) that were previously applied to Tara
Ocean communities (*57*). Additional databases (e.g., World Register of Marine Species) and
literature searches were also used. The following functional groups were included for 18S

protists: autotrophic protists, heterotrophic protists, mixotrophic protists, parasites,
photosymbionts, and other protists. Mixotrophic protists were further categorized as being
constitutive mixotrophs (CM) that inherently have chloroplasts and endosymbiotic specialist
non-constitutive mixotrophs (eSNCM) that harbor endosymbionts to support growth (*121*). We
recognize that many protists likely exhibit mixotrophy in some capacity, and so, our functional
annotation of this group may be underrepresented. Other protists represented higher level
taxonomic groups (domain or supergroup) that were unassigned at lower levels. Bacteria and
Archaea were categorized functionally as being heterotrophic or autotrophic.
690
691 **Statistical analyses**
692
Prior to ordination, ASV count tables were transformed to Aitchison distances, which is
estimated by transforming read counts via centered log-ratio normalization and computing
Euclidean distances. The resulting Aitchison distance matrices were used to observe microbial
composition and aimed to minimize compositional bias inherent with amplicon data (*83*).
Principal coordinate analysis (PCoA) of Aitchison distances was used to visualize 16S and 18S
community composition. Permutational multivariate analysis of variance (PERMANOVA) tests
were employed with the adonis2 function in vegan (9999 permutations) to estimate the impact of
spatial factors on community composition. This included categorical depth (surface, DCM, and
near bottom), sampling transect, and location of samples on the continental shelf vs. in open
ocean regions of the GOM designated by the 200 m isobath (Fig. 1A).
703
Samples were also grouped into clusters via hierarchical clustering (Ward's method) based on
Aitchison distances using the hclust function in vegan (Version 2.6-6.1; (*122*)). The optimal
number of clusters was determined based on average silhouette widths using the factoextra
package (Version 1.0.7; (*123*)). Silhouette widths offer an estimate on the quality of sample
clustering, with higher width coefficients indicating optimal clustering (*124*). Three clusters were
found to be optimal for both 16S and 18S (fig. S15), which largely reflected depth in the water
column (Fig. 1B; fig. S1). Cluster 1 consisted of samples collected at all depths on the shelf and
offshore in the surface layer, all confined to the photic zone (2–99 m). Cluster 2 consisted of
samples mainly from offshore and more stratified waters in the DCM (2–124 m), while Cluster 3
represented samples collected offshore in meso- to bathypelagic waters (135–3,326 m). The
photic zone extends to 200 m in many deeper regions of the GOM, and so, samples in Cluster 2
(and a handful in Cluster 3) were also technically collected within the photic zone. However, we
distinguish communities in Clusters 2–3 from Cluster 1 based on the large proportion of samples
confined to the open ocean DCM (Cluster 2; 80%) and mesopelagic (Cluster 3; 98%) that reflect
disparate habitats in the GOM.
719
Mean Shannon diversity index and richness (# of ASVs) were determined for each cluster using
the estimate_richness function in phyloseq (*119*) and compared against other clusters with
Wilcoxon tests ($P < 0.05$). Mean diversity and richness were also estimated along transects,

applying local regression (loess) curves to visualize trends using the geom_smooth function in
ggplot2 (Version 3.5.1; (*125*)). Stacked bar plots displaying mean relative abundance were
observed at the class level for 18S and order level for 16S for each sampling transect and cluster
using the microeco package in R (Version 1.7.1; (*126*)). Taxonomic profiles were also observed
using the treemap package in R (Version 2.4-4; (*127*)), a tiered approach to visualize relative
abundance across multiple taxonomic levels.
729
Indicator taxa that were more abundant and representative of high (or low) TA:DIC ratios were
statistically inferred using the indicspecies package in R (Version 1.7.14; (*128*)). The TA:DIC
ratio was chosen because it is a good proxy to determine the ocean's capacity to absorb
anthropogenic CO_2 by influencing its buffering capacity (*91*). Higher ratios indicate strong
buffering capacity (i.e., the capacity of seawater to buffer against acidification). Based on
histograms of TA:DIC, samples were grouped a priori into high (> 1.16) or low categories (<
1.16) that reflect different OA conditions (Fig. 7A–B). We focused on DNA samples collected
from the photic zone (Cluster 1) to mitigate natural depth effects and to provide additional
context to models (see next section). Indicator analysis was run separately on rarefied 16S or 18S
samples that were agglomerated to the species level using the function multipatt with 999
permutations (*128*). Significant ASVs ($P < 0.001$) were retained and summarized for high (or
low) TA:DIC and plotted against their mean relative abundance in the photic zone.
742
743 **Generalized linear models**
744
Generalized linear models (GLMs) were used to examine relationships between environmental
factors (predictor variables) and the relative abundance of major microbial groups (response
variables). GLMs focused on DNA samples collected in the photic zone (Cluster 1), in large part
to mitigate collinearity of factors that was prevalent in Clusters 2–3 (table S1). Separate GLMs
were performed for the top four most relatively abundant order level 16S and class level 18S
groups (Table 2). Separate models were constructed for *Synechococcus* and *Prochlorococcus* to
resolve differences between major cyanobacteria genera. Only variables that met requirements of
low collinearity (Spearman $r_s < 0.7$ or $> –0.7$) and a variance inflation factor (VIF) < 10 were
considered (*129*). Zurr et al. (2010) suggest using a more stringent VIF cutoff (< 3). However,
we aimed to retain as many variables in the dataset as possible, which meant a few variables
(e.g., DIC and salinity) approached VIF = 10. To select the best model for each 18S or 16S
group, variables were further selected in a stepwise manner based on Akaike Information Criteria
(AIC) values using the stepAIC function in the MASS package in R (Version 7.3-60; (*130*)).
Only significant variables ($P < 0.05$) were used in the final model.
759
Final models were constructed with either Poisson or negative binomial error distributions.
Initial model type was chosen by comparing standardized residuals and other model indices (e.g.,
AIC) with the compare_performance function in the performance package in R (Version 0.11.0;
(*131*)). GLMs were implemented with the glm.nb function for negative binomial models in the

MASS package or the glm function for Poisson models (family = Poisson) in the stats package in
base R (Version 4.3.1). We observed overdispersion in relative abundance data for several
Poisson models (Syndiniales, Dinophyceae, *Synechococcus*, and *Prochlorococcus*), in which
case negative binomial models were applied (Table 2). Model quality and fit was estimated for
each group using the check_model function in the performance package (*131*), which included
plots of posterior predictive checks (model simulations), standardized residuals (Q–Q plots),
homogeneity of variance, and collinearity of selected predictor variables. The goodness of fit
was assessed with a pseudo R^2 (Nagelkerke's), though standardized residual checks of the final
models were also carried out (*62*) to assess model fit and uniformity of the residuals
(Kolmogorov–Smirnov, $P > 0.05$). As an additional validation, relative abundance data for each
group was randomly split and trained with respective models using 80% of the data to predict a
test set that was left out (20%). Pearson correlations were performed between model-trained and
test data.
777
Model coefficients were scaled and visualized for each 18S and 16S group using the multiplot
function in the coefplot package in R (Version 1.2.8; (*132*)). Individual factors were plotted
against predicted relative abundance using the plot_model function in the sjPlot package in R
(Version 2.8.16; (*133*)). We focused predictive plots on temperature and carbonate chemistry
parameters (DIC and pH). Group-specific GLMs were used to predict relative abundance at all
GOMECC-4 sites where surface layer (< 10 m) variables were collected (135 out of 141 sites).
Six stations did not have representative CTD data available at the surface layer and were
excluded. Predicted relative abundance for all surface GOMECC-4 sites were observed in Ocean
Data View using Data-Interpolating Variational Analysis (DIVA) interpolation (*134*).
787

References

1. S. Díaz, J. Fargione, F. S. Chapin 3rd, D. Tilman, Biodiversity loss threatens human well-
being. *PLoS Biol.* **4**, e277 (2006).

2. N. Gruber, D. Clement, B. R. Carter, R. A. Feely, S. van Heuven, M. Hoppema, M. Ishii,
R. M. Key, A. Kozyr, S. K. Lauvset, C. Lo Monaco, J. T. Mathis, A. Murata, A. Olsen, F.
F. Perez, C. L. Sabine, T. Tanhua, R. Wanninkhof, The oceanic sink for anthropogenic
CO2 from 1994 to 2007. *Science* **363**, 1193–1199 (2019).

3. S. C. Doney, D. S. Busch, S. R. Cooley, K. J. Kroeker, The Impacts of Ocean
Acidification on Marine Ecosystems and Reliant Human Communities. *Annu. Rev.*
Environ. Resour. **45**, 1–30 (2020).

4. J. C. Orr, V. J. Fabry, O. Aumont, L. Bopp, S. C. Doney, R. A. Feely, A. Gnanadesikan,
N. Gruber, A. Ishida, F. Joos, R. M. Key, K. Lindsay, E. Maier-Reimer, R. Matear, P.
Monfray, A. Mouchet, R. G. Najjar, G.-K. Plattner, K. B. Rodgers, C. L. Sabine, J. L.
Sarmiento, R. Schlitzer, R. D. Slater, I. J. Totterdell, M.-F. Weirig, Y. Yamanaka, A.
Yool, Anthropogenic ocean acidification over the twenty-first century and its impact on
calcifying organisms. *Nature* **437**, 681–686 (2005).

5. M. M. White, D. C. McCorkle, L. S. Mullineaux, A. L. Cohen, Early exposure of bay
scallops (*Argopecten irradians*) to high CO_2 causes a decrease in larval shell growth.
PLoS One **8**, e61065 (2013).

6. R. K. Pachauri, M. R. Allen, V. R. Barros, J. Broome, W. Cramer, R. Christ, J. A.
Church, L. Clarke, Q. Dahe, P. Dasgupta, N. K. Dubash, Climate Change 2014: Synthesis
Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of
the Intergovernmental Panel on Climate Change (IPCC, 2014).

7. A. Z. Worden, M. J. Follows, S. J. Giovannoni, S. Wilken, A. E. Zimmerman, P. J.
Keeling, Rethinking the marine carbon cycle: Factoring in the multifarious lifestyles of
microbes. *Science* **347**, 127594 (2015).

8. D. A. Hutchins, F. Fu, Microorganisms and ocean global change. *Nat Microbiol* **2**, 17058
(2017).

9. P. W. Boyd, D. A. Hutchins, Understanding the responses of ocean biota to a complex
matrix of cumulative anthropogenic change. *Mar. Ecol. Prog. Ser.* **470**, 125–135 (2012).

10. D. A. Caron, D. A. Hutchins, The effects of changing climate on microzooplankton
grazing and community structure: drivers, predictions and knowledge gaps. *J. Plankton*
Res. **35**, 235–252 (2012).

11. D. G. Boyce, M. R. Lewis, B. Worm, Global phytoplankton decline over the past century.
Nature **466**, 591– 596 (07 2010).

12. L. Bopp, L. Resplandy, J. C. Orr, S. C. Doney, J. P. Dunne, M. Gehlen, P. Halloran, C.
Heinze, T. Ilyina, R. Séférian, J. Tjiputra, M. Vichi, Multiple stressors of ocean

ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences* **10**,
6225–6245 (2013).

13.   S. Dutkiewicz, J. J. Morris, M. J. Follows, J. Scott, O. Levitan, S. T. Dyhrman, I.
Berman-Frank, Impact of ocean acidification on the structure of future phytoplankton
communities. *Nat. Clim. Chang.* **5**, 1002–1006 (2015).

14.   S. A. Henson, B. B. Cael, S. R. Allen, S. Dutkiewicz, Future phytoplankton diversity in a
changing climate. *Nat. Commun.* **12**, 5372 (2021).

15.   K. R. M. Mackey, J. J. Morris, F. M. M. Morel, S. A. Kranz, Response of Photosynthesis
to Ocean Acidification. *Oceanography* **28**, 74–91 (2015).

16.   M. Segovia, M. R. Lorenzo, C. Iñiguez, C. García-Gómez, Physiological stress response
associated with elevated CO2 and dissolved iron in a phytoplankton community
dominated by the coccolithophore *Emiliania huxleyi*. *Mar. Ecol. Prog. Ser.* **586**, 73–89
(2018).

17.   V. Vázquez, P. León, F. J. L. Gordillo, C. Jiménez, I. Concepción, K. Mackenzie, E.
Bresnan, M. Segovia, High-CO2 Levels Rather than Acidification Restrict *Emiliania*
huxleyi Growth and Performance. *Microb. Ecol.* **86**, 127–143 (2023).

18.   M. D. Iglesias-Rodriguez, P. R. Halloran, R. E. M. Rickaby, I. R. Hall, E. Colmenero-
Hidalgo, J. R. Gittins, D. R. H. Green, T. Tyrrell, S. J. Gibbs, P. von Dassow, E. Rehm,
E. V. Armbrust, K. P. Boessenkool, Phytoplankton calcification in a high-CO2 world.
Science **320**, 336–340 (2008).

19.   A. C. Martiny, G. I. Hagstrom, T. DeVries, R. T. Letscher, G. L. Britten, C. A. Garcia, E.
Galbraith, D. Karl, S. A. Levin, M. W. Lomas, A. R. Moreno, D. Talmy, W. Wang, K.
Matsumoto, Marine phytoplankton resilience may moderate oligotrophic ecosystem
responses and biogeochemical feedbacks to climate change. *Limnol. Oceanogr.* **67**,
S378–S389 (2022).

20.   H. Sarmento, J. M. Montoya, E. Vázquez-Domínguez, D. Vaqué, J. M. Gasol, Warming
effects on marine microbial food web processes: how far can we go when it comes to
predictions? *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **365**, 2137–2149 (2010).

21.   X. A. G. Morán, F. Baltar, C. Carreira, C. Lønborg, Responses of physiological groups of
tropical heterotrophic bacteria to temperature and dissolved organic matter additions:
food matters more than warming. *Environ. Microbiol.* **22**, 1930–1943 (2020).

22.   T. E. Berry, B. J. Saunders, M. L. Coghlan, M. Stat, S. Jarman, A. J. Richardson, C. H.
Davies, O. Berry, E. S. Harvey, M. Bunce, Marine environmental DNA biomonitoring
reveals seasonal patterns in biodiversity and identifies ecosystem responses to anomalous
climatic events. *PLoS Genet.* **15**, e1007943 (2019).

23.   R. Gallego, E. Jacobs-Palmer, K. Cribari, R. P. Kelly, Environmental DNA
metabarcoding reveals winners and losers of global change in coastal waters.
Proceedings of the Royal Society B **287**, 20202424 (2020).

24.   E. Osborne, X. Hu, E. R. Hall, K. Yates, J. Vreeland-Dawson, K. Shamberger, L.
864   Barbero, J. Martin Hernandez-Ayon, F. A. Gomez, T. Hicks, Y.-Y. Xu, M. R.
865   McCutcheon, M. Acquafredda, C. Chapa-Balcorta, O. Norzagaray, D. Pierrot, A. Munoz-
866   Caravaca, K. L. Dobson, N. Williams, N. Rabalais, P. Dash, Ocean acidification in the
867   Gulf of Mexico: Drivers, impacts, and unknowns. *Prog. Oceanogr.* **209**, 102882 (2022).

868   25.   N. N. Rabalais, R. E. Turner, W. J. Wiseman Jr, Hypoxia in the Gulf of Mexico. *J.*
869   *Environ. Qual.* **30**, 320–329 (2001).

870   26.   W. Sturges, R. Leben, Frequency of Ring Separations from the Loop Current in the Gulf
871   of Mexico: A Revised Estimate. *J. Phys. Oceanogr.* **30**, 1814–1819 (2000).

872   27.   L. Linacre, R. Durazo, V. F. Camacho-Ibar, K. E. Selph, J. R. Lara-Lara, U. Mirabal-
873   Gómez, C. Bazán-Guzmán, A. Lago-Lestón, E. M. Fernández-Martín, K. Sidón-Ceseña,
874   Picoplankton carbon biomass assessments and distribution of *Prochlorococcus* ecotypes
875   linked to loop current eddies during summer in the southern gulf of Mexico. *J. Geophys.*
876   *Res. C: Oceans* **124**, 8342–8359 (2019).

877   28.   T. T. Sutton, M. R. Clark, D. C. Dunn, P. N. Halpin, A. D. Rogers, J. Guinotte, S. J.
878   Bograd, M. V. Angel, J. A. A. Perez, K. Wishner, R. L. Haedrich, D. J. Lindsay, J. C.
879   Drazen, A. Vereshchaka, U. Piatkowski, T. Morato, K. Błachowiak-Samołyk, B. H.
880   Robison, K. M. Gjerde, A. Pierrot-Bults, P. Bernal, G. Reygondeau, M. Heino, A global
881   biogeographic classification of the mesopelagic zone. *Deep Sea Res. Part 1 Oceanogr.*
882   *Res. Pap.* **126**, 85–102 (2017).

883   29.   M. R. Stukel, T. B. Kelly, M. R. Landry, K. E. Selph, R. Swalethorp, Sinking carbon,
884   nitrogen, and pigment flux within and beneath the euphotic zone in the oligotrophic,
885   open-ocean Gulf of Mexico. *J. Plankton Res.* **44**, 711–727 (2022).

886   30.   W.-J. Cai, X. Hu, W.-J. Huang, M. C. Murrell, J. C. Lehrter, S. E. Lohrenz, W.-C. Chou,
887   W. Zhai, J. T. Hollibaugh, Y. Wang, P. Zhao, X. Guo, K. Gundersen, M. Dai, G.-C.
888   Gong, Acidification of subsurface coastal waters enhanced by eutrophication. *Nat.*
889   *Geosci.* **4**, 766–770 (2011).

890   31.   R. Wanninkhof, L. Barbero, R. Byrne, W.-J. Cai, W.-J. Huang, J.-Z. Zhang, M. Baringer,
891   C. Langdon, Ocean acidification along the Gulf Coast and East Coast of the USA. *Cont.*
892   *Shelf Res.* **98**, 54–71 (2015).

893   32.   R. H. Weisberg, L. Zheng, Y. Liu, West Florida shelf upwelling: Origins and pathways.
894   *J. Geophys. Res. C: Oceans* **121**, 5672–5681 (2016).

895   33.   E. R. Hall, K. K. Yates, K. A. Hubbard, M. J. Garrett, J. D. Frankle, Nutrient and
896   carbonate chemistry patterns associated with *Karenia brevis* blooms in three West
897   Florida Shelf estuaries 2020-2023. *Frontiers in Marine Science* **11** (2024).

898   34.   M. J. Ulloa, P. Álvarez-Torres, K. P. Horak-Romo, R. Ortega-Izaguirre, Harmful algal
899   blooms and eutrophication along the mexican coast of the Gulf of Mexico large marine
900   ecosystem. *Environmental Development* **22**, 120–128 (2017).

901   35.   C. A. Heil, A. L. Muni-Morgan, Florida's Harmful Algal Bloom (HAB) Problem:
902   Escalating Risks to Human, Environmental and Economic Health With Climate Change.
903   *Frontiers in Ecology and Evolution* **9** (2021).

904   36.   P. M. Brannock, A. C. Ortmann, A. G. Moss, K. M. Halanych, Metabarcoding reveals
905   environmental factors influencing spatio-temporal variation in pelagic micro-eukaryotes.
906   *Mol. Ecol.* **25**, 3593–3604 (2016).

907   37.   C. G. Easson, J. V. Lopez, Depth-Dependent Environmental Drivers of Microbial
908   Plankton Community Structure in the Northern Gulf of Mexico. *Front. Microbiol.* **9**,
909   3175 (2018).

910   38.   K. Sidón-Ceseña, M. A. Martínez-Mercado, J. Chong-Robles, Y. Ortega-Saad, V. F.
911   Camacho-Ibar, L. Linacre, A. Lago-Lestón, Response of microbial eukaryote community
912   to the oligotrophic waters of the Gulf of Mexico: a plausible scenario for warm and
913   stratified oceans, *bioRxiv* (2023)

914   39.   M. W. Henson, J. C. Thrash, Microbial ecology of northern Gulf of Mexico estuarine
915   waters. *mSystems*, e0131823 (2024).

916   40.   B. B. Tolar, G. M. King, J. T. Hollibaugh, An analysis of thaumarchaeota populations
917   from the northern Gulf of Mexico. *Front. Microbiol.* **4**, 72 (2013).

918   41.   Thrash J. Cameron, Seitz Kiley W., Baker Brett J., Temperton Ben, Gillies Lauren E.,
919   Rabalais Nancy N., Henrissat Bernard, Mason Olivia U., Metabolic Roles of
920   Uncultivated Bacterioplankton Lineages in the Northern Gulf of Mexico "Dead Zone."
921   *MBio* **8**, 10.1128/mbio.01017–17 (2017).

922   42.   C. de Vargas, S. Audic, N. Henry, J. Decelle, F. Mahe, R. Logares, E. Lara, C. Berney,
923   N. L. Bescot, I. Probert, M. Carmichael, J. Poulain, S. Romac, S. Colin, J.-M. Aury, L.
924   Bittner, S. Chaffron, M. Dunthorn, S. Engelen, O. Flegontova, L. Guidi, A. Horak, O.
925   Jaillon, G. Lima-Mendez, J. Luke, S. Malviya, R. Morard, M. Mulot, E. Scalco, R. Siano,
926   F. Vincent, A. Zingone, C. Dimier, M. Picheral, S. Searson, S. Kandels-Lewis, T. O.
927   Coordinators, S. G. Acinas, P. Bork, C. Bowler, G. Gorsky, N. Grimsley, P. Hingamp, D.
928   Iudicone, F. Not, H. Ogata, S. Pesant, J. Raes, M. E. Sieracki, S. Speich, L. Stemmann, S.
929   Sunagawa, J. Weissenbach, P. Wincker, E. Karsenti, E. Boss, M. Follows, L. Karp-Boss,
930   U. Krzic, E. G. Reynaud, C. Sardet, M. B. Sullivan, D. Velayoudon, Eukaryotic plankton
931   diversity in the sunlit ocean. *Science* **348**, 1261605–1261605 (2015).

932   43.   S. Sunagawa, L. P. Coelho, S. Chaffron, J. R. Kultima, K. Labadie, G. Salazar, B.
933   Djahanschiri, G. Zeller, D. R. Mende, A. Alberti, F. M. Cornejo-Castillo, P. I. Costea, C.
934   Cruaud, F. d'Ovidio, S. Engelen, I. Ferrera, J. M. Gasol, L. Guidi, F. Hildebrand, F.
935   Kokoszka, C. Lepoivre, G. Lima-Mendez, J. Poulain, B. T. Poulos, M. Royo-Llonch, H.
936   Sarmento, S. Vieira-Silva, C. Dimier, M. Picheral, S. Searson, S. Kandels-Lewis, T. O.
937   Coordinators, C. Bowler, C. de Vargas, G. Gorsky, N. Grimsley, P. Hingamp, D.
938   Iudicone, O. Jaillon, F. Not, H. Ogata, S. Pesant, S. Speich, L. Stemmann, M. B.
939   Sullivan, J. Weissenbach, P. Wincker, E. Karsenti, J. Raes, S. G. Acinas, P. Bork,
940   Structure and function of the global ocean microbiome. *Science* **348**, 1261359– 1261359

941   (05 2015).

44. A. Djurhuus, P. H. Boersch-Supan, S.-O. Mikalsen, A. D. Rogers, Microbe biogeography tracks water masses in a dynamic oceanic frontal system. *R Soc Open Sci* **4**, 170033 (2017).

45. J. K. Pearman, B. Jones, M. Qashqari, S. Carvalho, Eukaryotic diversity patterns in the Red Sea: from the surface to the deep. *Frontiers in Marine Science* **10** (2024).

46. E. A. Walsh, J. B. Kirkpatrick, S. D. Rutherford, D. C. Smith, M. Sogin, S. D'Hondt, Bacterial diversity and community composition from seasurface to subseafloor. *ISME J.* **10**, 979–989 (2016).

47. T. Pommier, P. R. Neal, J. M. Gasol, M. Coll, S. G. Acinas, C. Pedrós-Alió, Spatial patterns of bacterial richness and evenness in the NW Mediterranean Sea explored by pyrosequencing of the 16S rRNA. *Aquat. Microb. Ecol.* **61**, 221–233 (2010).

48. Y.-C. Yeh, J. A. Fuhrman, Contrasting diversity patterns of prokaryotes and protists over time and depth at the San-Pedro Ocean Time series. *ISME Commun* **2**, 36 (2022).

49. N. V. Patin, Z. A. Dietrich, A. Stancil, M. Quinan, J. S. Beckler, E. R. Hall, J. Culter, C. G. Smith, M. Taillefert, F. J. Stewart, Gulf of Mexico blue hole harbors high levels of novel microbial lineages. *ISME J.* **15**, 2206–2232 (2021).

50. A. E. Santoro, C. Buchwald, M. R. McIlvin, K. L. Casciotti, Isotopic signature of N(2)O produced by marine ammonia-oxidizing archaea. *Science* **333**, 1282–1285 (2011).

51. Z. I. Johnson, E. R. Zinser, A. Coe, N. P. McNulty, E. M. S. Woodward, S. W. Chisholm, Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. *Science* **311**, 1737– 1740 (03 2006).

52. D. Tsementzi, J. Wu, S. Deutsch, S. Nath, L. M. Rodriguez-R, A. S. Burns, P. Ranjan, N. Sarode, R. R. Malmstrom, C. C. Padilla, B. K. Stone, L. A. Bristow, M. Larsen, J. B. Glass, B. Thamdrup, T. Woyke, K. T. Konstantinidis, F. J. Stewart, SAR11 bacteria linked to ocean anoxia and nitrogen loss. *Nature* **536**, 179– 183 (08 2016).

53. K. E. Selph, R. Swalethorp, M. R. Stukel, T. B. Kelly, A. N. Knapp, K. Fleming, T. Hernandez, M. R. Landry, Phytoplankton community composition and biomass in the oligotrophic Gulf of Mexico. *J. Plankton Res.* **44**, 618–637 (2022).

54. R. Massana, J. del Campo, M. E. Sieracki, S. Audic, R. Logares, Exploring the uncultured microeukaryote majority in the oceans: reevaluation of ribogroups within stramenopiles. *ISME J.* **8**, 854–866 (2014).

55. T. Biard, Diversity and ecology of Radiolaria in modern oceans. *Environ. Microbiol.* **24**, 2179–2200 (2022).

56. L. Guillou, M. Viprey, A. Chambouvet, R. M. Welsh, A. R. Kirkham, R. Massana, D. J. Scanlan, A. Z. Worden, Widespread occurrence and genetic diversity of marine

parasitoids belonging to Syndiniales (Alveolata). *Environ. Microbiol.* **10**, 3349–3365
(2008).

57.   T. Cordier, I. B. Angeles, N. Henry, F. Lejzerowicz, C. Berney, R. Morard, A. Brandt,
M.-A. Cambon-Bonavita, L. Guidi, F. Lombard, P. M. Arbizu, R. Massana, C. Orejas, J.
Poulain, C. R. Smith, P. Wincker, S. Arnaud-Haond, A. J. Gooday, C. de Vargas, J.
Pawlowski, Patterns of eukaryotic diversity from the surface to the deep-ocean sediment.
Science Advances **8**, eabj9309 (2022).

58.   C. A. Durkin, I. Cetinić, M. Estapa, Z. Ljubešić, M. Mucko, A. Neeley, M. Omand,
Tracing the path of carbon export in the ocean through DNA sequencing of individual
sinking particles. *ISME J.*, 1–11 (2022).

59.   O. Flegontova, P. Flegontov, S. Malviya, S. Audic, P. Wincker, C. de Vargas, C. Bowler,
J. Lukeš, A. Horák, Extreme Diversity of Diplonemid Eukaryotes in the Ocean. *Curr.*
Biol. **26**, 3060–3065 (2016).

60.   J. Saltzman, E. R. White, Determining the role of environmental covariates on
planktivorous elasmobranch population trends within an isolated marine protected area.
Mar. Ecol. Prog. Ser. **722**, 107–123 (2023).

61.   E. Bravo-Ormaza, R. Arauz, S. Bessudo, A. Hearn, A. P. Klimley, F. Ladino-Archila, J.
López-Macías, T. Steiner, C. Peñaherrera-Palma, Scalloped hammerhead shark *Sphyrna*
lewini relative abundance comparison in three offshore marine protected areas of the
Eastern Tropical Pacific. *Environ. Biol. Fishes*, doi: 10.1007/s10641-023-01454-6
(2023).

62.   A. F. Zuur, E. N. Ieno, C. S. Elphick, A protocol for data exploration to avoid common
statistical problems. *Methods Ecol. Evol.* **1**, 3–14 (2010).

63.   F. E. Muller-Karger, J. P. Smith, S. Werner, R. Chen, M. Roffer, Y. Liu, B. Muhling, D.
Lindo-Atichati, J. Lamkin, S. Cerdeira-Estrada, D. B. Enfield, Natural variability of
surface oceanographic conditions in the offshore Gulf of Mexico. *Prog. Oceanogr.* **134**,
54–76 (2015).

64.   N. Yingling, T. B. Kelly, T. A. Shropshire, M. R. Landry, K. E. Selph, A. N. Knapp, S.
A. Kranz, M. R. Stukel, Taxon-specific phytoplankton growth, nutrient utilization and
light limitation in the oligotrophic Gulf of Mexico. *J. Plankton Res.* **44**, 656–676 (2022).

65.   X. A. G. Morán, L. Alonso-Sáez, E. Nogueira, H. W. Ducklow, N. González, Á. López-
Urrutia, L. Díaz-Pérez, A. Calvo-Díaz, N. Arandia-Gorostidi, T. M. Huete-Stauffer,
More, smaller bacteria in response to ocean's warming? *Proc. Biol. Sci.* **282** (2015).

66.   N. Arandia-Gorostidi, P. K. Weber, L. Alonso-Sáez, X. A. G. Morán, X. Mayali,
Elevated temperature increases carbon and nitrogen fluxes between phytoplankton and
heterotrophic bacteria through physical attachment. *ISME J.* **11**, 641–650 (2017).

67.   J. Taucher, J. Arístegui, L. T. Bach, W. Guan, M. F. Montero, A. Nauendorf, E. P.
Achterberg, U. Riebesell, Response of Subtropical Phytoplankton Communities to Ocean

Acidification Under Oligotrophic Conditions and During Nutrient Fertilization. *Frontiers in Marine Science* **5** (2018).

68.   M. W. Lomas, B. M. Hopkinson, J. L. Losh, D. E. Ryan, D. L. Shi, Y. Xu, F. M. M. Morel, Effect of ocean acidification on cyanobacteria in the subtropical North Atlantic. *Vol.* **66**, 211–222 (2012).

69.   P. Flombaum, W.-L. Wang, F. W. Primeau, A. C. Martiny, Global picophytoplankton niche partitioning predicts overall positive response to ocean warming. *Nat. Geosci.* **13**, 116–120 (2020).

70.   F.-X. Fu, M. E. Warner, Y. Zhang, Y. Feng, D. A. Hutchins, Effects of increased temperature and CO2 on photosynthesis, growth, and elemental ratios in marine *Synechococcus* and *Prochlorococcus* (cyanobacteria). *J. Phycol.* **43**, 485–496 (2007).

71.   L. T. Bach, S. Alvarez-Fernandez, T. Hornick, A. Stuhr, U. Riebesell, Simulated ocean acidification reveals winners and losers in coastal phytoplankton. *PLoS One* **12**, e0188198 (2017).

72.   A. Martiny, S. Kathuria, P. Berube, Widespread metabolic potential for nitrite and nitrate assimilation among *Prochlorococcus* ecotypes. *Proc. Natl. Acad. Sci. U. S. A.*, doi: 10.1073/pnas.0902532106 (06 2009).

73.   J. H. Brown, J. F. Gillooly, A. P. Allen, V. M. Savage, G. B. West, Toward a metabolic theory of ecology. *Ecology* **85**, 1771–1789 (2004).

74.   Y. Feng, F. Chai, M. L. Wells, Y. Liao, P. Li, T. Cai, T. Zhao, F. Fu, D. A. Hutchins, The Combined Effects of Increased pCO2 and Warming on a Coastal Phytoplankton Assemblage: From Species Composition to Sinking Rate. *Frontiers in Marine Science* **8** (2021).

75.   H. G. Horn, N. Sander, A. Stuhr, M. Algueró-Muñiz, L. T. Bach, M. G. J. Löder, M. Boersma, U. Riebesell, N. Aberle, Low CO2 Sensitivity of Microzooplankton Communities in the Gullmar Fjord, Skagerrak: Evidence from a Long-Term Mesocosm Study. *PLoS One* **11**, e0165800 (2016).

76.   D. K. Stoecker, P. J. Hansen, D. A. Caron, A. Mitra, Mixotrophy in the Marine Plankton. *Ann. Rev. Mar. Sci.* **9**, 311–335 (2017).

77.   O. U. Mason, E. J. Canter, L. E. Gillies, T. K. Paisie, B. J. Roberts, Mississippi River Plume Enriches Microbial Diversity in the Northern Gulf of Mexico. *Front. Microbiol.* **7**, 1048 (2016).

78.   F. A. Gomez, S.-K. Lee, F. J. Hernandez Jr, L. M. Chiaverano, F. E. Muller-Karger, Y. Liu, J. T. Lamkin, ENSO-induced co-variability of Salinity, Plankton Biomass and Coastal Currents in the Northern Gulf of Mexico. *Sci. Rep.* **9**, 178 (2019).

79.   F. A. Gomez, R. Wanninkhof, L. Barbero, S.-K. Lee, F. J. Hernandez Jr, Seasonal patterns of surface inorganic carbon system variables in the Gulf of Mexico inferred from

a regional high-resolution ocean biogeochemical model. *Biogeosciences* **17**, 1685–1700
(2020).

80.   A. K. Williams, A. S. McInnes, J. R. Rooker, A. Quigg, Changes in Microbial Plankton
Assemblages Induced by Mesoscale Oceanographic Features in the Northern Gulf of
Mexico. *PLoS One* **10**, e0138230 (2015).

81.   S. Anglès, A. Jordi, D. W. Henrichs, L. Campbell, Influence of coastal upwelling and
river discharge on the phytoplankton community composition in the northwestern Gulf of
Mexico. *Prog. Oceanogr.* **173**, 26–36 (2019).

82.   S. Wood, Generalized Additive Models: An Introduction with R (Chapman and
Hall/CRC, 2017).

83.   G. B. Gloor, J. M. Macklaim, V. Pawlowsky-Glahn, J. J. Egozcue, Microbiome Datasets
Are Compositional: And This Is Not Optional. *Front. Microbiol.* **8**, 2224 (2017).

84.   W. Gong, A. Marchetti, Estimation of 18S Gene Copy Number in Marine Eukaryotic
Plankton Using a Next-Generation Sequencing Approach. *Frontiers in Marine Science* **6**
(2019).

85.   S. Collins, P. W. Boyd, M. A. Doblin, Evolution, Microbes, and Changing Ocean
Conditions. *Ann. Rev. Mar. Sci.* **12**, 181–208 (2020).

86.   F. Partensky, J. Blanchot, D. Vaulot, Differential distribution and ecology of
Prochlorococcus and *Synechococcus* in oceanic waters: a review. *Bulletin de l'Institut*
océanographique. Monaco. n° spécial, 457–475 (1999).

87.   A. Buchan, G. R. LeCleir, C. A. Gulvik, J. M. González, Master recyclers: features and
functions of bacteria associated with phytoplankton blooms. *Nat. Rev. Microbiol.* **12**,
686–698 (2014).

88.   C. J. Gobler, Climate Change and Harmful Algal Blooms: Insights and perspective.
Harmful Algae **91**, 101731 (2020).

89.   R. M. Errera, S. Yvon-Lewis, J. D. Kessler, L. Campbell, Responses of the dinoflagellate
Karenia brevis to climate change: pCO2 and sea surface temperatures. *Harmful Algae* **37**,
110–116 (2014).

90.   B. A. Stauffer, H. A. Bowers, E. Buckley, T. W. Davis, T. H. Johengen, R. Kudela, M. A.
McManus, H. Purcell, G. J. Smith, A. Vander Woude, M. N. Tamburri, Considerations in
Harmful Algal Bloom Research and Monitoring: Perspectives From a Consensus-
Building Workshop and Technology Testing. *Frontiers in Marine Science* **6** (2019).

91.   E. S. Egleston, C. L. Sabine, F. M. M. Morel, Revelle revisited: Buffer factors that
quantify the response of ocean chemistry to changes in DIC and alkalinity. *Global*
Biogeochem. Cycles (2010).

92.   A. Z. Worden, J. K. Nolan, B. Palenik, Assessing the dynamics and ecology of marine

picophytoplankton: The importance of the eukaryotic component. *Limnol. Oceanogr.* **49**,
168–179 (2004).

93.   P. Ziveri, B. de Bernardi, K.-H. Baumann, H. M. Stoll, P. G. Mortyn, Sinking of
coccolith carbonate and potential contribution to organic carbon ballasting in the deep
ocean. *Deep Sea Res. Part 2 Top. Stud. Oceanogr.* **54**, 659–675 (2007).

94.   C.-E. Schaum, B. Rost, S. Collins, Environmental stability affects phenotypic evolution
in a globally distributed marine picoplankton. *ISME J.* **10**, 75–84 (2016).

95.   K. T. Lohbeck, U. Riebesell, T. B. H. Reusch, Adaptive evolution of a key phytoplankton
species to ocean acidification. *Nat. Geosci.* **5**, 346–351 (2012).

96.   K.-H. Baumann, B. Boeckel, Spatial distribution of living coccolithophores in the
southwestern Gulf of Mexico. *J. Micropalaeontol.* **32**, 123–133 (2013).

97.   S. J. Giovannoni, SAR11 Bacteria: The Most Abundant Plankton in the Oceans. *Ann.*
Rev. Mar. Sci. **9**, 231–255 (2017).

98.   M. S. Rappé, S. A. Connon, K. L. Vergin, S. J. Giovannoni, Cultivation of the ubiquitous
SAR11 marine bacterioplankton clade. *Nature* **418**, 630– 633 (08 2002).

99.   N. R. Bates, Y. M. Astor, M. J. Church, K. Currie, J. E. Dore, M. González-Dávila, L.
Lorenzoni, F. Muller-Karger, J. Olafsson, J. M. Santana-Casiano, A Time-Series View of
Changing Surface Ocean Chemistry Due to Ocean Uptake of Anthropogenic CO_2 and
Ocean Acidification. *Oceanography* **27**, 126–141 (2014).

100.   S. Dutkiewicz, P. Cermeno, O. Jahn, M. J. Follows, A. E. Hickman, D. A. A. Taniguchi,
B. A. Ward, Dimensions of marine phytoplankton diversity. *Biogeosciences* **17**, 609–634
(2020).

101.   E. L. Cavan, S. A. Henson, P. W. Boyd, The Sensitivity of Subsurface Microbes to Ocean
Warming Accentuates Future Declines in Particulate Carbon Export. *Frontiers in*
Ecology and Evolution **6** (2019).

102.   S. A. Henson, C. Laufkötter, S. Leung, S. L. C. Giering, H. I. Palevsky, E. L. Cavan,
Uncertain response of ocean biological carbon export in a changing world. *Nat. Geosci.*
15, 248–254 (2022).

103.   C. Langdon, Determination of dissolved oxygen in seawater by Winkler titration using
the amperometric technique. (2010).

104.   J.-Z. Zhang, G. A. Berberian, Determination of dissolved silicate in estuarine and coastal
waters by gas segmented continuous flow colorimetric analysis. *Methods for the*
determination of chemical substances in marine and estuarine environmental matrices,
366–360 (1997).

105.   K. M. Johnson, A. E. King, J. M. Sieburth, Coulometric TCO2 analyses for marine
studies; an introduction. *Mar. Chem.* **16**, 61–82 (1985).

106. A. G. Dickson, C. L. Sabine, J. R. Christian, "Guide to best practices for ocean CO2
measurements" (Report, North Pacific Marine Science Organization, 2007).

107. R. Wanninkhof, K. Thoning, Measurement of fugacity of CO2 in surface water using
continuous and discrete sampling methods. *Mar. Chem.* **44**, 189–204 (1993).

108. R. H. Byrne, L. R. Kump, K. J. Cantrell, The influence of temperature and pH on trace
metal speciation in seawater. *Mar. Chem.* **25**, 163–181 (1988).

109. S. M. A. C. van Heuven, M. Hoppema, O. Huhn, H. A. Slagter, H. J. W. de Baar, Direct
observation of increasing CO2 in the Weddell Gyre along the Prime Meridian during
1973–2008. *Deep Sea Res. Part 2 Top. Stud. Oceanogr.* **58**, 2613–2635 (2011).

110. S. R. Anderson, L. R. Thompson, Optimizing an enclosed bead beating extraction method
for microbial and fish environmental DNA. *Environmental DNA*, doi: 10.1002/edn3.251
(2021).

111. A. E. Parada, D. M. Needham, J. A. Fuhrman, Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. *Environ. Microbiol.* **18**, 1403– 1414 (05 2016).

112. L. A. Amaral-Zettler, E. A. McCliment, H. W. Ducklow, S. M. Huse, A method for
studying protistan diversity using massively parallel sequencing of V9 hypervariable
regions of small-subunit ribosomal RNA Genes. *PLoS One* **4**, e6372 (07 2009).

113. M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**, 10–12 (2011).

114. L. R. Thompson, S. R. Anderson, P. A. Den Uyl, N. V. Patin, S. J. Lim, G. Sanderson, K.
D. Goodwin, Tourmaline: A containerized workflow for rapid and iterable amplicon
sequence analysis using QIIME 2 and Snakemake. *Gigascience* **11** (2022).

115. B. J. Callahan, P. J. Mcmurdie, S. P. Holmes, Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. *ISME J.* **11**, 2639–2643
(2016).

116. E. Pruesse, C. Quast, K. Knittel, B. M. Fuchs, W. Ludwig, J. Peplies, F. O. Glöckner,
SILVA: a comprehensive online resource for quality checked and aligned ribosomal
RNA sequence data compatible with ARB. *Nucleic Acids Res.* **35**, 7188–7196 (2007).

117. L. Guillou, D. Bachar, S. Audic, D. Bass, C. Berney, L. Bittner, C. Boutte, G. Burgaud,
C. De Vargas, J. Decelle, J. Del Campo, J. R. Dolan, M. Dunthorn, B. Edvardsen, M.
Holzmann, W. H. C. F. Kooistra, E. Lara, N. Le Bescot, R. Logares, F. Mahé, R.
Massana, M. Montresor, R. Morard, F. Not, J. Pawlowski, I. Probert, A. L. Sauvadet, R.
Siano, T. Stoeck, D. Vaulot, P. Zimmermann, R. Christen, The Protist Ribosomal
Reference database (PR2): A catalog of unicellular eukaryote Small Sub-Unit rRNA
sequences with curated taxonomy. *Nucleic Acids Res.* **41** (2013).

118. N. A. Bokulich, B. D. Kaehler, J. R. Rideout, M. Dillon, E. Bolyen, R. Knight, G. A.

Huttley, J. Gregory Caporaso, Optimizing taxonomic classification of marker-gene
amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome* **6**, 90
(2018).

119. P. J. McMurdie, S. Holmes, Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS One* **8** (2013).

120. G. S. Kandlikar, Z. J. Gold, M. C. Cowen, R. S. Meyer, A. C. Freise, N. J. B. Kraft, J.
Moberg-Parker, J. Sprague, D. J. Kushner, E. E. Curd, ranacapa: An R package and
Shiny web app to explore environmental DNA data with exploratory statistics and
interactive visualizations. *F1000Res.* **7**, 1734 (2018).

121. A. Mitra, D. A. Caron, E. Faure, K. J. Flynn, S. G. Leles, P. J. Hansen, G. B. McManus,
F. Not, H. do Rosario Gomes, L. F. Santoferrara, D. K. Stoecker, U. Tillmann, The
Mixoplankton Database (MDB): Diversity of photo-phago-trophic plankton in form,
function, and distribution across the global ocean. *J. Eukaryot. Microbiol.* **70**, e12972
(2023).

122. J. Oksanen, F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R.
Minchin, R. B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, H.
Wagne, Vegan: Community ecology package. R package version 2.6-6.1 (2024).

123. A. Kassambara, Practical Guide to Cluster Analysis in R: Unsupervised Machine
Learning Vol. 1 (Sthda, 2017).

124. L. Kaufman, P. J. Rousseeuw, Finding Groups in Data: An Introduction to Cluster
Analysis (John Wiley & Sons, 2009).

125. H. Wickham, M. Averick, J. Bryan, W. Chang, L. McGowan, R. François, G.
Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. Pedersen, E. Miller, S. Bache, K.
Müller, J. Ooms, D. Robinson, D. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke,
K. Woo, H. Yutani, Welcome to the tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

126. C. Liu, Y. Cui, X. Li, M. Yao, microeco: an R package for data mining in microbial
community ecology. *FEMS Microbiol. Ecol.* **97**, fiaa255 (2021).

127. B. Bederson, B. Shneiderman, M. Wattenberg, Ordered and quantum treemaps: Making
effective use of 2D space to display hierarchies. *ACM Trans. Graph.* **21**, 833–854 (2002).

128. M. De Cáceres, P. Legendre, M. Moretti, Improving indicator species analysis by
combining groups of sites. *Oikos* **119**, 1674–1684 (2010).

129. G. James, D. Witten, T. Hastie, R. Tibshirani, An Introduction to Statistical Learning
(Springer, 2013), vol. 112.

130. B. Ripley, B. Venables, D. M. Bates, K. Hornik, A. Gebhardt, D. Firth, M. B. Ripley,
Package "mass." *Cran r* **538**, 113–120 (2013).

131. D. Lüdecke, M. Ben-Shachar, I. Patil, P. Waggoner, D. Makowski, Performance: An R

package for assessment, comparison and testing of statistical models. *J. Open Source Softw.* **6**, 3139 (2021).

132.   J. P. Lander, coefplot: Plots coefficients from fitted models. R package version 1.2.8 (2022).

133.   D. Lüdecke, sjPlot: Data visualization for statistics in social science. R package version 2.8.16 (2024).

134.   R. Schlitzer, Ocean Data View. https://odv.awi.de (2023).

Acknowledgements

We thank the captain and crew of the NOAA Ship *Ronald H. Brown* for logistical support for sample collection in the Gulf. We acknowledge the efforts of many scientists at NOAA and partnering institutions for collecting and processing oceanographic data on GOMECC-4. We thank Easton White and Elizabeth Harvey for their helpful conversations, and Sean McAllister for carefully reviewing the manuscript. We acknowledge the genomics services performed in the RTSF Genomics Core at Michigan State University.

Funding: This work was funded in part through the NOAA Ocean Acidification Program (OAP) ROR #02bfn4816 under project numbers 21392 (Thompson) and 20708 (Barbero) and by awards NA16OAR4320199 and NA21OAR4320190 to the Northern Gulf Institute from NOAA's Office of Oceanic and Atmospheric Research, U.S. Department of Commerce. This research was carried out in part under the auspices of the Cooperative Institute for Marine and Atmospheric Studies (CIMAS) and NOAA, cooperative agreement NA20OAR4320472. This work was also supported by NSF award OCE-2019589 for the Center for Chemical Currencies of a Microbial Planet (C-CoMP). This is C-CoMP publication #046.

Author contributions: LRT, SRA, LB, and CRK conceived the study. SRA collected DNA samples. LB led collection of carbonate chemistry parameters. SRA and LRT processed DNA samples. SRA performed bioinformatics and data analysis. LRT, KS, and FAG contributed to improve data analysis. BAS and AS provided guidance on taxonomy and functional assignments. SRA and LRT led the writing of the manuscript. All authors contributed to revising the manuscript and approved the final version.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: Code and associated files needed to reproduce results and figures for this study are available on GitHub (https://github.com/aomlomics/gomecc) and have been archived on Zenodo (https://zenodo.org/records/13102580). All 18S and 16S sequence data generated from this study have been published at the National Center for Biotechnology Information (NCBI)'s Sequence Read Archive and BioSample database and are available with BioProject accession number PRJNA887898. Species count data generated from this study have been published on the Ocean Biodiversity Information System (OBIS) and the Global Biodiversity Information Facility (GBIF) at https://doi.org/10.15468/sm6fpz. Biological data has also been submitted to the National Centers for Environmental Information (NCEI) at https://www.ncei.noaa.gov/archive/accession/0250940/data/0-data/noaa-aoml-gomecc. Environmental measurements from the Niskin bottles and CTD profiles are also available at NCEI at https://doi.org/10.25921/4twf-pp50 and https://doi.org/10.25921/04h7-gv36, respectively. A cruise report detailing all the sampling and analyzing procedures during GOMECC-4 is available at https://doi.org/10.25923/rwx5-s713.

1266   **Figures and Tables**

1267

1268

Figure 1 (legend on next page)

Figure 1: Vertical and horizontal DNA sampling across the GOM. (A) Map of all sites
sampled on GOMECC-4. Sites are colored by transect and indicate instances where DNA was
(filled) or was not (empty) sampled. Samples at all sites were collected in triplicate.
Environmental metadata was collected from all stations. Contour lines indicate depth in the
GOM and correspond to the right y-axis in panel **B**. Transects are also labeled to match the color
of stations along a given transect. Samples were collected counterclockwise in the Gulf starting
at the 27°N line. **(B)** Map displaying depth-related position (log scale) of samples across the
GOM. Stations are colored by transect at the surface, matching transect colors in panel **A**.
Samples with depth are colored by their clusters (Clusters 1–3) that were determined via
hierarchical clustering of Aitchison distances and largely reflected depth in the water column.
Cluster 1 generally corresponded to shelf waters at all depths and in the open GOM at the surface
(photic zone; n = 235), Cluster 2 represented sites in the DCM in the open GOM (DCM; n =
137), and Cluster 3 was confined to open ocean sites in deep waters (aphotic zone; n = 89).
Clustering was similar between 18S (shown) and 16S samples (fig. S16).
1284

Figure 2: Bacterial and archaeal community dynamics in the GOM from 16S metabarcoding. (A) Ridgeline plots showing the depth distribution of samples within each cluster (Clusters 1–3). Clusters were determined via hierarchical clustering of Aitchison distances: Cluster 1 (photic zone), Cluster 2 (DCM), and Cluster 3 (aphotic zone). **(B)** Principal coordinates analysis of Aitchison distances, with samples colored by their respective clusters. **(C)** Mean observed richness (# of ASVs) and Shannon diversity index for Clusters 1–3, with points representing individual samples. Significant differences between clusters were determined with Wilcoxon tests (** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). **(D)** Stacked bar plots of mean relative abundance (%) at the order level in each sampling transect and faceted by cluster. Transects are ordered on the x-axis based on the order of sampling (counterclockwise) on GOMECC-4, except for FL straits and Cape Coral that were sampled last but grouped spatially with other FL lines. Bar plots display the top 12 most relatively abundant groups over all samples ("others" in gray). Taxonomy was assigned via the SILVA database. Generalized linear models focused on the top four most relatively abundant groups in Cluster 1 (red asterisks). Models for Synechococcales were constructed at the genus level to discriminate between *Prochlorococcus* and *Synechococcus*. LA = Louisiana and PAIS = Padre Island National Seashore. Transects have the same labels in all subsequent plots.

1306
1307   **Figure 3 (legend on next page)**

Figure 3: Protist community dynamics in the GOM from 18S metabarcoding. (A) Ridgeline plots showing the depth distribution of samples within each cluster (Clusters 1–3). Clusters reflected depth along the shelf and open GOM and closely resembled clustering of 16S samples. **(B)** Principal coordinates analysis of Aitchison distances, with 18S samples colored by cluster. **(C)** Mean observed richness (# of ASVs) and Shannon diversity index for Clusters 1–3, with points representing individual samples. Significant differences between clusters were determined with Wilcoxon tests (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$). **(D)** Stacked bar plots of mean relative abundance (%) at the class level in each sampling transect and faceted by cluster. Transects are ordered the same as in Fig. 2. Bar plots display the top 12 most relatively abundant groups over all samples ("others" in gray). Protist taxonomy was assigned via the PR2 database. Generalized linear models focused on the top four most relatively abundant groups in Cluster 1 (red asterisks).

1321
1322   **Figure 4 (legend on next page)**

Figure 4: Generalized linear models of major 16S taxa reveal group-specific environmental drivers in the photic zone. (A) Scaled model coefficients (± 2 standard deviations) of predictor variables (environmental factors) that were significant to the final model (based on AIC values). Models were constructed with group-specific relative abundance as the response variable. The most relatively abundant 16S groups were modeled, which included heterotrophs and autotrophs. Models were generated at the order level, except for cyanobacteria (Synechococcales), where separate models were run for *Prochlorococcus* and *Synechococcus*. Only covariates that were statistically significant to a given model were plotted. **(B)** Predicted response estimates (relative abundance) and 95% confidence intervals (CIs) of major 16S groups to temperature, DIC, and in situ pH. NS = not significant. **(C)** An example of a posterior predictive plot, highlighting the fit of observed vs. model-predicted relative abundance for the final SAR11 model. The model-predicted data was simulated with 50 bootstraps and followed a similar trend as the observed data. **(D)** Pearson correlation (with 95% CI) between SAR11 test and modeled relative abundance to estimate model fit. Predicted abundance was derived from the final SAR11 model using a subset of the data (80%; 219 samples) and correlated to test data that was left out (20%; 48 samples). Model fit of other major 16S groups is shown in fig. S9.

1340
1341   **Figure 5 (legend on next page)**

Figure 5: GLMs of major 18S taxa reveal group-specific drivers in the photic zone. (A)
Scaled model coefficients (± 2 standard deviations) of predictor variables (environmental
factors) that were significant to the final model (based on AIC values). Models were constructed
with group-specific relative abundance as the response variable. The top four most relatively
abundant 18S groups were modeled separately, spanning constitutive mixotrophs (CM),
parasites, autotrophs, and heterotrophs. Covariates that were not statistically significant to a
given model are not shown. **(B)** Predicted response estimates (relative abundance) and 95%
confidence intervals (CIs) of major 18S groups to temperature, DIC, and pH. NS = not
significant. **(C)** An example of a posterior predictive plot, highlighting the fit of observed vs.
model-predicted relative abundance for the final Syndiniales model. The model-predicted data
was simulated with 50 bootstraps and followed a similar trend as the observed data. **(D)** Pearson
correlation (with 95% CI) between Syndiniales test and modeled relative abundance. Predicted
abundance was derived from the final Syndiniales model using a subset of the data (80%; 187
samples) and correlated to test values that were left out (20%; 43 samples). Model fit of other
major 18S groups is shown in fig. S10.

1357

Figure 6: Expanding current microbial distributions in the GOM. Predicted relative abundance (%) of major 16S **(A–E)** and 18S groups **(F–I)** at 135 GOMECC-4 sites modeled with each respective GLM (from Table 2). Model results have been interpolated using DIVA interpolation in Ocean Data View. Isobaths are shown for 200 m, 2,000 m, and 3,000 m. Scales for predicted relative abundance vary by taxonomic group (on the right of each panel) but display low to high relative abundance.

Figure 7: Protist indicator taxa based on TA:DIC ratios in the photic zone. (A) Histogram showing the density distribution of 18S samples in the photic zone (Cluster 1) based on TA:DIC ratios. **(B)** Values of in situ pH vs. TA:DIC in the photic zone, with samples colored by transect. Pearson correlation between variables is shown, with 95% confidence interval. The dotted line in panels **A–B** indicate the manual cutoff used for indicator analysis: low TA:DIC < 1.16 vs. high TA:DIC > 1.16. **(C–D)** Indicator values vs. average relative abundance (%) for protist ASVs in the photic zone that were significant to the analysis ($P < 0.001$) in samples with either low TA:DIC **(C)** or high TA:DIC **(D)**. Protist ASVs are colored by division and the top five ASVs with the highest indicator values are labeled in each panel, identified to their lowest possible taxonomic assignment (via the PR2 database). DG = Dino-Group. See table S2 for a full list of 18S (and 16S) indicator ASVs. Similar plots for 16S ASVs are shown in fig. S12.

Table 1: Environmental factors used in microbial models. Factors were grouped into
parameter type and chosen for initial GLMs based on Spearman correlations (table S1) and low
variance inflation factors (VIF < 10) to mitigate collinearity among predictor variables. VIFs
varied slightly between 16S and 18S (in parentheses) due to differences in sample size (n = 274
for 16S; n = 235 for 18S) following clustering analysis. Triplicate samples were included in
models. Datasets clustered similarly, as evidenced by a similar range in the predictor values.
Initial factors were used to construct group-specific models.

1383

Parameter type	Factor	Values	VIF 16S (18S)
Hydrography	Temperature	20.83–30.12 (°C)	2.8 (2.4)
	Salinity	25.16–36.61 (psu)	7 (7.5)
	Oxygen	105.46–232.33 (μmol kg^{-1})	3.1 (5)
Nutrients	Nitrate	0–6.16 (μmol kg^{-1})	4.6 (4.2)
	Phosphate	0–0.85 (μmol kg^{-1})	5.7 (6.2)
	Ammonium	0.12–2.37 (μmol kg^{-1})	1.6 (1.7)
Carbonate chemistry	DIC	1891.67–2186.16 (μmol kg^{-1})	9.1 (8.9)
	pH	7.88–8.16	5.2 (6.1)

Table 2: Final models for major microbial groups in the photic zone. Protists were examined
at the class level and prokaryotes at the order level. GLMs were constructed for *Prochlorococcus*
and *Synechococcus*. Models were run either with negative binomial (neg bin) or Poisson
distributions. Variables that were significant to the final model ($P < 0.05$) are shown for each
group and reflect stepwise selection based on Akaike Information Criterion (AIC). See Table 1
for the full list of variables considered. Pseudo R^2 values are shown as a proxy for model fit,
though standardized residuals and validation tests confirmed model fit. Temp = temperature
(°C); Sal = salinity; Oxy = oxygen (μmol kg^{-1}); PO$_4$ = phosphate (μmol kg^{-1}); NO$_3$ = nitrate
(μmol kg^{-1}); NH$_4$ = ammonium (μmol kg^{-1}); DIC = dissolved inorganic carbon (μmol kg^{-1}).

Group	Taxonomy	GLM	Type	R^2
Protists	Dinophyceae	Temp + Sal + Oxy + DIC	Neg Bin	0.5
	Syndiniales	Temp + Sal + PO$_4$ + NH$_4$ + pH + DIC	Neg Bin	0.57
	Sagenista	Temp + Oxy + PO$_4$ + pH	Poisson	0.28
	Prymnesiophyceae	Temp + Sal + PO$_4$	Poisson	0.6
Prokaryotes	SAR11	Temp + Oxy + NH$_4$ + DIC	Poisson	0.26
	Synechococcus	Oxy + NO$_3$ + pH + DIC	Neg Bin	0.38
	Prochlorococcus	Sal + PO$_4$ + NO$_3$ + pH	Neg Bin	0.8
	SAR86	Temp + Oxy + NH$_4$ + pH + DIC	Poisson	0.33
	Flavobacteriales	Sal + DIC	Poisson	0.64

1413
1414