Dynamics of antimicrobial resistance and virulence of *Salmonella enterica* serovar

Dublin

Cheryll M Sia[1,2*], Rebecca L Ambrose[3], Mary Valcanis[2], Patiyan Andersson[2], Susan A

Ballard[2], Benjamin P Howden[1,2], Deborah A Williamson[4#], Jaclyn S Pearson[3,5,6*#], Danielle J

Ingle [1*#]

[1] Department of Microbiology and Immunology, The University of Melbourne at The Peter

Doherty Institute for Infection and Immunity, Melbourne, 3000, Victoria, Australia.

[2] Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology &

Immunology, The University of Melbourne at The Peter Doherty Institute for Infection and

Immunity, Melbourne, 3000, Victoria, Australia.

[3] Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical Research,

Clayton, 3168, Victoria, Australia.

[4] Department of Infectious Diseases, The University of Melbourne at the Peter Doherty Institute

for Infection and Immunity, Melbourne, 3000, Victoria, Australia.

[5] Department of Microbiology, Monash University, Clayton, 3168, Victoria, Australia.

[6]School of Medicine, University of St Andrews, St Andrews, KY16 9TF, Fife, United Kingdom.

 # Contributed equally

* Corresponding author

CMS: cmsia@unimelb.edu.au

JSP: jp287@st-andrews.ac.uk

DJI: danielle.ingle@unimelb.edu.au

ABSTRACT

Salmonella Dublin is a host-adapted, invasive non-typhoidal *Salmonella* (iNTS) serovar that

causes bloodstream infections in humans and demonstrates increasing prevalence of

antimicrobial resistance (AMR). Here, we characterised a global population of 1,303 *S.* Dublin

isolates for resistance and virulence mechanisms using a combined approach of *in silico* and *in*

vitro analyses. We observed distinct populations of Vi antigen-negative *S.* Dublin circulating

in different geographical regions, with the emergence of a North American cluster ~60 years

ago. Further, we identified two distinct lineages in Australia, within which we discovered a

novel hybrid plasmid encoding both AMR and mercuric resistance, highlighting the potential

for emerging virulent and resistant lineages. These differences, in addition to the lack of Vi

flagella represent adaptation mechanisms that may enhance survival and pathogenicity, and in

turn affect prevention and control strategies for future *S.* Dublin infections. Collectively, our

study embodies one of the most comprehensive investigations of *S.* Dublin and highlights the

potential impact of pathogen adaptation on global health and economic burdens experienced

by different sectors including livestock and human health.

Funding

DAW was supported by a National Health and Medical Research Council (NHMRC)

Investigator Grant (GNT1174555) and a Medical Research Future Fund (MRFF) Grant

(FSPGN000045). BPH is supported by NHMRC Investigator Grant (GNT1196103). DJI is

supported by an NHMRC Investigator Grant (GNT1195210). JSP was supported by a Sylvia

and Charles Viertel Senior Medical Fellowship. MDU PHL is funded by the Victorian

Government,   Australia.   No   conflict   of   interest   declared.

Keywords

Invasive nontyphoidal *Salmonella*, antimicrobial resistance, heavy metal resistance, virulence;

One Health

INTRODUCTION

Invasive nontyphoidal *Salmonella* (iNTS) infections are a major cause of global morbidity and

mortality (1). Of the >2,500 NTS serovars, iNTS infections are associated with a relatively

limited number of serovars. These include *Salmonella enterica* serovar Dublin (*S.* Dublin), *S.*

Panama, *S.* Virchow, *S.* Choleraesuis and *S.* Typhimurium ST313 (2–4). *S.* Dublin, a bovine-

adapted pathogen, is one of the leading iNTS serovars both in Australia and globally (5–7), and

is associated with bacteraemia in a high proportion of infections (2,5–7). Symptomatic cattle

may develop septicaemia and transplacental transmission (7,8), while asymptomatic infections

in cattle enable persistence within the herd population, facilitating disease spread (7,9). *S.*

Dublin causes zoonotic infections predominately through the consumption of contaminated

dairy and meat produce (10–12), and as such is considered a 'One Health' pathogen. (10–12).

Antimicrobial resistance (AMR) in NTS has increased over the past several decades, leading

to the reduction of therapeutic options (6,13). Although *S.* Dublin isolates are largely

susceptible to therapeutics, AMR has become increasingly prevalent in *S.* Dublin (7,12,14–16).

For example, a multidrug-resistant (MDR; defined as resistance to ≥ 3 antimicrobial classes

(17)) lineage has recently been described in North America, characterised by varying profiles

of AMR to ampicillin, streptomycin, chloramphenicol, sulphonamides, tetracyclines and third

generation cephalosporins. These MDR *S.* Dublin isolates all type as sequence type 10 (ST10),

and the AMR determinants have been demonstrated to be carried on an IncC plasmid that has

recombined with a virulence plasmid encoding the *spvRABCD* operon with IncX1 and IncFII

replicons, to produce a 329kb megaplasmid (12,16).

S. Dublin is closely related to *S.* Enteritidis and *S.* Gallinarum (18), with differences in accessory genome content reflective of adaptation to different ecological niches (18). The majority of the STs reported for *S.* Dublin are ST10 or single-locus variants of this ST, although there is a second lineage, ST74, that is distinct from the main *S.* Dublin population (19). This uncommon *S.* Dublin lineage, ST74, may be variably serotyped as *S.* Dublin or *S.* Enteritidis dependent upon the strains used to generate typing sera in different laboratories (19), which may have also contributed to it not being identified. The antigenic formula of *S.* Dublin (1,9,12:g,p:-) is highly similar to *S.* Enteritidis (1,9,12:g,m:-), with three non-synonymous point mutations in *fliC* (Ala220Val, Thr315Ilc and Thr318Ala) differentiating the two serovars. Although the Vi flagella antigen (encoded by the *viaB* locus) is part of the antigenic definition for *S.* Dublin, it has been shown that the Vi antigen is absent in many *S.* Dublin isolates (19,20). Other putative virulence determinants in *S.* Dublin include *Salmonella* pathogenicity islands (SPIs) 6 and 19, which both encode type VI secretion system (T6SS) machineries (21,22). The presence of two T6SSs in *S.* Dublin contrasts with other NTS serovars, which encode a single T6SS (21,22). T6SSs are molecular nanomachines that deliver effector proteins into eukaryotic and/or prokaryotic cells to promote pathogen survival within multispecies and complex immune environments such as the gut. Recent studies have demonstrated a role for SPI-6 and SPI-19 in interbacterial competition by specific *S.* Dublin strains (23,24), however their effect on host immune processes remains largely unknown.

We previously undertook a genomic investigation of another emerging NTS lineage in Australia, *Salmonella enterica* serovar 4,[5],12:i:-, and identified several bacterial traits that may have contributed to the successful spread of this lineage (25–29). In particular, we hypothesised that flagellin deletion and MDR status may have contributed to host immune

evasion and positive selection, respectively, in *Salmonella enterica* serovar 4,[5],12:i:- (30).

Here, we sought to understand the contribution of virulence and resistance to the recent

emergence of MDR *S.* Dublin. We interrogated a global dataset of 1,303 isolates collected

from 13 countries on five continents and identified distinct geographical lineages associated

with specific AMR and virulence profiles, including a lineage unique with increased capability

for intracellular survival and immune evasion in human macrophages.

RESULTS

Distinct S. Dublin populations circulate globally

A total of 1,303 *S.* Dublin genomes were included in our analysis, spanning four decades

(Figure 1A, Supplementary Table 1-2). This dataset included 53 genomes from humans in

Australia, and 1,250 genomes from Africa, Europe, North and South America collected from

humans (*n*=462), animals (*n*=551), food (*n*=223) and other sources that were not stated (*n*=67)

(Figure 1B, Supplementary Table 1). In total, 1,235 genomes (99.4%) were ST10 and 58 were

a single locus variant (SLV) of ST10, with two double locus variant genomes (e.g. ST4100)

restricted to isolates from South America. The exceptions were ST74 and SLV ST1545 that

were only detected in Australia (Figure 1C-E, Supplementary Table 3). All isolates, including

the ST74 lineage, were serotyped *in silico* as *S.* Dublin. However, the ST74 lineage and a single

ST10 isolate had codon profiles in *fliC* of Gly60, Leu138, Ala220, Thr315 and Ala318, in

contrast to the remaining ST10 and SLV, which both had a profile of Gly60, Leu138, Val220,

Ile315 and Ala318 (the previously described 'Du2' and 'Du1' electrophoretic profiles (20),

respectively) (Supplementary Table 4).

A maximum likelihood (ML) phylogeny was inferred from the 1,303 genomes, revealing two

major lineages of *S.* Dublin, ST74 and ST10 (Figure 1D, Supplementary Figure 1). Within the

ST10 lineage, five clades were identified using Bayesian Analysis of Population Structure (BAPS), inferred from the alignment of 2,957 core single nucleotide polymorphisms (SNPs). Four clades were broadly associated with geographical region (Supplementary Figure 1, Supplementary Table 1). The Australian ST10 isolates (n = 39) clustered in clade 1, while isolates from North America were associated with clade 5 (Supplementary Figure 1). The BAPS clade 2 were the isolates that didn't have sufficient evidence to be split into further clades, hence are spread across the tree spanning a range of geographical regions.

Emergence of S. Dublin lineages in the 20th century

To investigate the emergence of *S*. Dublin clades, Bayesian phylodynamic analysis was undertaken on a representative subset of 660 genomes, selected for phylogenetic, geographical and AMR diversity (see methods). We inferred a maximum clade credibility (MCC) phylogeny from an alignment of 3,971 SNPs using the highest supported model of a relaxed lognormal molecular clock and coalescent exponential tree prior. The estimated substitution rate was 1.05 x 10^{-7} (95% highest posterior density (HPD): 1.16 x 10^{-7}– 9.41 x 10^{-8}), slightly higher than previous estimated mean substitution rates for host-restricted *Salmonella* serovars (*S*. Typhi and *S*. Paratyphi A) but lower than rates reported for host-generalist NTS serovars *S*. Kentucky and *S*. Agona (31).

The topology of the MCC tree was consistent with the ML tree, with geographically associated sub-clades were observed in the MCC tree (Figure 2, interactive tree is available in Microreact under https://microreact.org/project/sdublinpaper). The most recent common ancestor (MRCA) for the *S*. Dublin global population was inferred to be 1901 (HPD 1878-1924), with the MRCA of the different clades and sub-clades largely falling between 1950 and 2000. For

example, the most recent common ancestor (MRCA) of clade 5 MDR isolates from North

America was estimated to be ~1964 (HPD 1952 -1974) (Figure 2). The MRCA of the Australian

sub-clade was estimated to be ~1967 (HPD 1953-1980); with the Australian and North

American isolates inferred to have a MRCA ~1931 (HPD 1914-1947). Moreover, three clades

(1, 3, and 4) had European sub-clades associated with Sweden, the UK, and Denmark,

respectively that were all largely susceptible to antimicrobials. Some regions, such as Europe

and South America, were associated with several sub-clades of *S.* Dublin, potentially

suggesting multiple importations and expansions, while other regions, such as Australia were

comprised of a single sub-clade within clade 1.

Resistance determinants are associated with geographical regions

Different plasmid types co-occurred with different AMR and HMR determinants in the *S.*

Dublin population, and these were also associated with differential geographical distribution

(Figure 2, Supplementary Figure 3-4, Supplementary Table 1). Overall, 51.5% (667/1,295) of

S. Dublin genomes had at least one AMR determinant, with 632/667 (94.8%) isolates classified

as MDR. MDR *S.* Dublin was predominantly identified from North American isolates, although

some AMR determinants were detected in *S.* Dublin from other geographic regions (Figure 3,

Supplementary Figures 2-3). AMR *S.* Dublin collected from animals was common

(*n*=305/667; 45.7%). AMR in *S.* Dublin from food sources and humans was similar in

proportion (*n*=174/667; 26.1% and *n*=173/667; 25.9% respectively). Although these profiles

are likely impacted by the sampling frame of different studies, the source of collection was not

considered as a selection criterion for this study, and as such, may broadly represent the AMR

profiles of *S.* Dublin from different sources (Figure 3A-B, Supplementary Figure 1A-C).

The most prevalent AMR determinants were *strA-B* (*n*=620), *floR* (*n*=556), *tet*(A) (*n*=608) and

sul2 (*n*=596) (Supplementary Figure 2). AMR mechanisms to ciprofloxacin (triple point

mutations in quinolone resistance determining regions *gyrA*[83:S-F]*, gyrA*[87:D-N]*,*

parC[80:S-I])*,* macrolides (*mph*(A)), and colistin (*mcr-9.1*) were each detected in single

genomes from *S.* Dublin clade 5 at a low frequency of 0.08% (*n*=1/1,303) in the dataset. The

only third generation cephalosporin (3GC) resistance mechanism detected was *bla*CMY-2,

present in 467/667 (70%) of AMR *S.* Dublin genomes. Although *strA-B*, *tet*(A) and *bla*TEM-1

were observed in multiple clades, *floR*, *sul2* and *bla*CMY-2 were restricted to North American

isolates that fell in clade 5 (Supplementary Figure 2-3). This MDR profile, in addition to *strA-*

strB and *tet*(A), was mediated by IncC plasmids in 593 genomes (*n*=509 from the United States

of America [USA], *n*=82 and *n*=2 from the UK) (Supplementary Figure 3, 4C). Nearly all clade

3 isolates linked to the UK had no AMR determinants, except for four isolates collected

between 2010-2015 (Supplementary Figure 1-3, Figure 3, Supplementary Table 1).

In addition to AMR determinants, we characterised biocide and heavy metal resistance (HMR)

profiles of the *S.* Dublin isolates. Although gold resistance determinants *golS* and *golT*, were

ubiquitous in *S.* Dublin (*n*=1,302/1,303; 99.9%), biocide and HMR determinants generally co-

occurred with AMR determinants (Figure 3C, Supplementary Figures 2-3). The biocide

quaternary ammonium compound resistance determinant, *qacEdelta1*, was primarily identified

in MDR genomes from clade 5 isolates (*n*=76/614; 12.4%), associated with North America.

Further, HMR determinants *ter* (tellurium metal), *pcoS* (copper metal) and *arsC* (arsenate

metal) were detected in a single genome, SRR5905515 from North American animal isolate

collected in 2016, with AMR mechanisms to seven different antimicrobial classes, including

colistin (*mcr-9.1*) (Supplementary Figure 3, Supplementary Table 1). Mercury resistance genes

(*merABDEPRT*) were found in 341/614 (55.5%) MDR clade 5 genomes, while *merCPRT* genes

were detected in single MDR genomes from Australia and South America (Supplementary Table 1). The *mer* operon co-occurred in 90.1% (*n* = 534) of the isolates mediated by the IncC MDR plasmid, suggesting an association of this operon with resistance determinants (Supplementary Figure 3, Supplementary Table 1).

Characterisation of S. Dublin plasmids and identification of a novel hybrid plasmid

Both IncX1 (*n*=1269/1303; 97.4%) and IncFII(S) (*n*=1213/1303; 93.1%) plasmid replicons were found at high frequency within the dataset, in line with previous findings (12). Three plasmid replicons, namely IncN, IncX1 and IncFII(S), and a MOBF/MOBP relaxase, were detected in one of the Australian *S.* Dublin ST10 genomes from clade 1 (AUSMDU00035676). Comparative analysis revealed the AUSMDU00035676 plasmid lacked the *spv* operon and had low homology to reference IncN and IncX1/IncFII(S) plasmids previously associated with *Salmonella* (Supplementary Figure 5, Supplementary Table 5). The loss of the *spv* operon may be associated with a less invasive phenotype (32). This finding suggests the detection of a potentially novel hybrid plasmid with IncN/IncX1/IncFII(S) markers that encodes AMR and HMR determinants flanked by IS26 elements, which may aid gene mobilisation (33). In contrast, the AUSMDU00056868 plasmid had IncX1/IncFII(S) plasmid replicon markers and was found to not encode any mechanisms for resistance. This plasmid was >99% homologous to known virulence reference *S.* Dublin plasmids pOU1115 and pSDU2_USMARC_69807 and encoded the *spv* operon (Supplementary Table 5, Supplementary Figure 6). This suggests that the *S.* Dublin virulence plasmid is conserved, while the *spv* operon that has been shown to enhance disease severity (14,16,34–36) is maintained on the plasmid.

Diversity in virulence determinants across S. Dublin lineages

To investigate the potential for invasiveness within *S.* Dublin, we applied a previously

described bioinformatic tool to our dataset of 1,303 *S.* Dublin genomes (37). In brief, this tool

infers the likelihood of invasiveness by assigning an index score between 0 to 1 that reflects

the level of genome degradation, whereby a higher score suggests genomic features that may

be more associated with a narrower host range and great likelihood of invasiveness (37). We

performed initial validation work using 36 complete publicly available reference genomes,

covering known invasive and non-invasive phenotypes (Supplementary Table 8). Using this

validation set, invasive and largely host-restricted serovars, including *S.* Choleraesuis, *S.*

Paratyphi and *S.* Typhi, scored >0.5, suggestive of an invasive genomic profile (Figure 4A,

Supplementary Table 8). Conversely, non-invasive NTS serovars with broader ecological

niches, including *S.* Anatum, ST19 *S.* Typhimurium and *S.* Derby, scored <0.5 (Figure 4A,

Supplementary Table 8). When applied to our *S.* Dublin dataset, only eleven genomes

(*n*=11/1,303; 0.84%) had an index score < 0.5. These included three ST10 genomes and all

eight ST74 genomes (Figure 4B, Supplementary Table 6). No association was observed

between index scores and region or source attribution (Figure 4B).

Putative virulence determinants identified in *S.* Dublin genomes included SPIs, the *spv* operon

(16), prophages and other individual genes (Supplementary Table 6). Some putative virulence

factors, such as *pagN* [cell adhesion and invasion (38,39)] and *pgtE,* [virulence (26,40)], were

found in 1,303 (100%, *n*=1,303/1,303) and 1,302 (99.9%, *n*=1,302/1,303) genomes

respectively, and no SNPs in the promoter region of *pgtE* were detected, a mutation previously

reported in invasive *S.* Typhimurium ST313 (26). In contrast, ST313-gi of unknown function

(26,40), the Gifsy-2 like prophage [unknown function (26,40)] and *ggt* gene thought to be

involved in gut colonisation (34), were detected in the major ST10 *S.* Dublin lineages, and

absent in ST74 *S.* Dublin. The *viaB* operon encoding the Vi flagella antigen was only detected

in SLV ST73, a historical isolate from France that displayed the Du3 electrophoretic type that

was consistent with a previous study (20). This suggests that the Vi antigen is not conserved in

the *S.* Dublin population or associated with virulence. Further, there was no association

between the number of virulence determinants and the AMR and HMR profile of isolates,

suggesting a potential lack of fitness cost associated with resistance mechanisms and virulence

(Supplementary Table 6).

The presence of SPI-6 and SPI-19 differed across *S.* Dublin isolates. Both SPIs encode T6SSs

that provide a means for injecting effector proteins into adjacent competing bacteria or host

cells for immune evasion (22,41). SPI-19 was absent in the ST74 *S.* Dublin lineage but present

in all ST10 *S.* Dublin (Supplementary Figure 7). The genetic differences between ST74 and

ST10 were reflected in the disparity of their index scores of <0.5 and >0.5, respectively. The

SPI-6 region was detected in all isolates, however <90% sequence coverage was detected in

422/1,303 genomes (32.4%). Long read sequencing on three genomes of the ST74 lineage and

three genomes from the ST10 clade 1 lineage corroborated these findings (Figure 4C-E),

demonstrating truncation of SPI-6 with the T6SS machinery (Figure 4D). Long-read

sequencing also revealed an absence of SPI-19 in the three ST74 isolates and presence of SPI-

19 in the three ST10 isolates (Figure 4C, E). The absence of SPI-19 in ST74 suggests that this

deletion event occurred during its divergence from the primary *S.* Dublin lineage, ST10.

Infection of human macrophages reveals distinct intracellular replication patterns between

ST10 and ST74 lineages

Given the distinct genomic profiles of ST10 and ST74 lineages, we compared the phenotypic

profile of a selection of Australian isolates from each lineage *in vitro*. The immortalised human

macrophage cell line, THP-1, was used to assess invasion, intracellular replication and host

cytotoxicity induced by a subset of clinical isolates from the two lineages. Specifically, we assayed an ST10 population comprising eleven isolates, including an isolate from the highly related ST4293 lineage, and an ST74 population comprising seven isolates including an isolate from the highly related ST1545 lineage (Supplementary Table 10). Although predicted invasive index scores were higher for the ST10 population (Figure 4B), each isolate failed to replicate more than 2-fold over 24 hours in THP-1 cells (Figure 5A), whereas isolates within the ST74 population displayed a sustained increase in fold change of intracellular replication, up to 8-fold, over 24 hours (Figure 5B). Comparing bacterial growth (CFU/ml) across both sequence types, both lineages were comparable for initial bacterial uptake (Figure 5C) however, the ST74 population replicated to significantly higher levels in THP-1 macrophages at 9- and 24-hours post-infection (hpi) compared to the ST10 population (Figure 5D-F). Given that NTS also replicate in epithelial cells, we utilised an immortalised human intestinal epithelial cell line, HT-29, to assess *S.* Dublin replication. In contrast to the results obtained in macrophages, we found that 6 out of 11 isolates within the ST10 population replicated more efficiently (up to 10-fold) in HT-29 cells, whereas all seven isolates within the ST74 population showed minimal replication (up to 2-fold) over 24 hours (Supplementary Figure 8A-C).

Increased replication of ST74 isolates in human macrophages does not correlate with increased host cell death

It is well documented that protype *S.* Typhimurium (ST19) potently induce macrophage cytotoxicity both *in vitro* and *in vivo* (42). Here, we utilised a cellular lactate dehydrogenase (LDH) release assay to assess THP-1 cytotoxicity in response to infection with *S.* Dublin ST74 and ST10 populations. Surprisingly, the substantial intracellular replication of the ST74 population in THP-1 cells induced significantly less host cell cytotoxicity at 3 and 9 hpi compared with the ST10 population suggesting potential immune evasion by ST74 (Figure 6A,

B). By 24 hpi however, cytotoxicity induced by both ST74 and ST10 populations was relatively high, with no significant difference observed across the sequence types (Figure 6C). In contrast, there was minimal cytotoxicity observed in infected HT-29 cells across the board (Supplementary Figure 8D), suggesting a less potent activation of host cell death in infected epithelial cells compared with macrophages.

To understand how the ST74 population may be supressing macrophage death at earlier stages of infection, we measured protein secretion of IL-1ß via cytokine release and immunoblot assays at 9 hpi, respectively. IL-1ß is a potent pro-inflammatory cytokine that is released upon activation of the NLRC4 inflammasome by components of the *Salmonella* type III secretion apparatus (43), resulting in a lytic cell death process known as pyroptosis (44). Here, we observed significantly less IL-1ß release by THP-1 cells infected with isolates from the ST74 population at 9 hpi compared with those from the ST10 population (Figure 6D), directly reflecting the LDH cytotoxicity data represented in Figure 6B. To further support our observations, we selected single representative isolates of ST74 and ST10 to perform immunoblot analyses of cleaved IL-1ß (activated form) within THP-1 cells. Here, ST74 induced less IL-1ß cleavage at 9 hpi compared to the ST10 (Figure 6E). We included the ubiquitous laboratory NTS reference strain, *S.* Typhimurium SL1344 in the immunoblot analysis as this is known to potently induce pyroptosis in human macrophages (45). As expected, SL1344 induced significant IL-1ß cleavage during infection of THP-1 macrophages (Figure 6E). To assess the correlation of cellular cytotoxicity as measured by LDH, IL-1ß release and intracellular replication, we generated a PCA plot of all isolates within the ST74 and ST10 populations. Here we can clearly demonstrate that isolates from the ST74 population have increased replicative ability, induce less cytotoxicity, and lower levels of IL-1ß release (Figure 6F) compared with those within the ST10 population. There were however, some ST10

isolates that induced similarly low levels of host cell death, presumably due to lower levels of intracellular replication (Figure 6F and Supplementary Figure 9A, B). These isolates generally clustered away from the ST10 isolates that induced more cell death (Supplementary Figure 9C).

A pangenome analysis of the isolates from both ST10 and ST74 lineages revealed that out of a total of 5,792 genes, there were 473 and 1,143 genes specific to the ST10 and ST74 lineage, respectively (Supplementary Figure 10, Supplementary Table 11). Further, out of 1,143 genes, there were 362 core genes present in all isolates from the ST74 lineage (Supplementary Table 11). While these findings require further investigation, an increased gene content and genome size were consistently observed in all isolates of the ST74 lineage (Supplementary Table 9 and 11). These results suggest an association with unique genome content of the ST74 lineage and may account for altered host responses and replicative ability (Supplementary Table 9).

DISCUSSION

Our data provide comprehensive insights into the emergence and spread of distinct lineages of *S*. Dublin, with striking geographic associations with AMR and virulence, consistent with previous work (12). Analysing the largest global dataset of *S*. Dublin genomes to date, we show that highly resistant *S*. Dublin is almost entirely confined to North America, with emergence in the late 20th century and ongoing local microevolution. North American *S*. Dublin isolates in clade 5 were characterised by the presence of an IncC plasmid, often with co-location of AMR and heavy metal resistance determinants. Heavy metals and biocides are both used in the agri-food industry and may contribute to the persistence of AMR via co-selection (46). The co-occurrence of AMR, HMR and biocide resistance determinants in animal, food and human *S*. Dublin genomes highlights the potential for selection pressure in one ecological niche (e.g. the

use of biocides in farming or heavy metals in animal feeds) leading to AMR in another (e.g.

AMR in humans).

The phylogeographic separation of *S*. Dublin lineages is consistent with other studies (7,12,47),

and is likely to be influenced by several factors, including local AMR use and patterns of

livestock movement. Given the livestock reservoir of *S*. Dublin is relatively host-restricted to

cattle, it is conceivable that both historic and contemporary global cattle movement may have

contributed to region-specific differences in *S*. Dublin. For example, Australian *S*. Dublin

genomes in clade 1 shared a common ancestor with North American isolates in clade 5. During

the early 1930's, Brahman cattle were imported from the U.S into Australia (48), supporting

our phylodynamic finding that the common ancestral link between *S*. Dublin isolates from

North America and Australia emerged in the early-mid 20[th] century. Following region-specific

introductions, it is likely that local antimicrobial and husbandry practices may then contribute

to the local evolution of *S*. Dublin. In contrast to North American MDR isolates, ST10 *S*.

Dublin isolates from Australia were predominantly fully susceptible to antimicrobials.

Selective pressure, misdirected treatment, the misuse of antimicrobials in cattle and MDR

plasmids in agricultural settings have been associated with the increase of MDR *S*. Dublin cases

in North America (49). Conversely, the use of antimicrobials has been extensively minimised

in the Australian dairy industry, hence cattle infections are rarely associated with AMR (50).

Consistent with regional differences in AMR profiles were geographic differences in plasmid

profiles. IncC plasmids are large (70-250kb), can encode up to 10 AMR determinants and are

widespread in North America (51–53), while IncN plasmids are smaller (30-70kb), encode up

to seven AMR determinants and are widespread in Europe (51,53,54).

Of note, amongst *S.* Dublin isolates from Australia, genomic analysis (cf. serotyping) revealed

a smaller population of ST74 isolates. *S.* Dublin ST74 was previously identified as *S.*

Enteritidis ST74, and sits as an intermediate between *S.* Dublin and *S.* Enteritidis (19), and has

been understudied to date in studies of *S.* Dublin. The difference between the antigenic formula

of *S.* Dublin and *S.* Enteritidis is in the H1 flagella region, which is typed as "*g,p*" for *S.* Dublin

and "*g,m*" for *S.* Enteritidis. ST74 *S.* Dublin genomes were characterised by: (i) a lack of any

AMR or HMR determinants; (ii) lack of complete SPI-6 regions, specifically the T6SS, and

(iii) a lack of putative virulence determinants including ST313-gi, Gifsy-2 like prophage and

SPI-19 (which encodes a T6SS mechanism for intestinal colonisation and survival that interacts

with the host by mediating macrophage cytotoxicity (55)). Our data demonstrated that while

machine learning predicted the ST74 population to have a lower potential for invasiveness

compared to ST10, the genetic differences in ST74 may contribute to increased replication in

human macrophages. Collectively, these findings highlight the co-circulation in Australia of

two *S.* Dublin populations, an ST10 and ST74 population, which may be phenotypically

characterised as *S.* Dublin.

The lack of the Vi antigen in 1,302/1,303 (99.9%) *S.* Dublin genomes has several implications.

First, the antigenic formula of *S.* Dublin, 1,9,12[Vi] may need to be reviewed if most *S.* Dublin

genomes lack the machinery to express the Vi flagella. These limitations with serotyping have

been highlighted in a previous study, suggesting a transition to a novel genotyping scheme that

would prevent these phenotypic discrepancies (56). Second, the Vi antigen is a vaccine target

for *S.* Typhi (57), and the Vi antigen has been explored as a target for *S.* Dublin (58). While

previous studies have noted that some *S.* Dublin genomes may be capable of producing the Vi

antigen (58), the genomic data from this study shows that these genomes are in the minority of

the *S.* Dublin population and may not be stable targets for *S.* Dublin specific vaccines.

Our phenotypic data demonstrated a striking difference in replication dynamics between ST10

and ST74 populations in human macrophages, whereby isolates within the ST74 population

replicated significantly over 24 hours compared with the rapid clearance of those within the

ST10 population after 9 hours of infection. The ST74 population generally induced

significantly less host cell death during the early-mid stages of macrophage infection,

supported by a limited processing and release of the pro-inflammatory cytokine IL-1ß at 9 hpi.

Research to date suggest that *Salmonellae* are potent activators of inflammasome activation

(59), however most of the data that support this theory have been generated using laboratory

adapted strains of *S.* Typhimurium. Our data, however, suggest that ST74 (and highly related)

isolates may utilise specific immune evasion mechanisms to avoid host recognition and the

activation of host cell death signalling pathways in phagocytes in the early stages of infection.

This observation is not unique to NTS. In 2015, Carden *et al.* (60) demonstrated that *S.*

Typhimurium ST313, a major cause of bacteraemia in humans in sub-Saharan Africa, induced

significantly less inflammasome activation compared with *S.* Typhimurium ST19 isolates

during infection of murine bone marrow-derived macrophages. Such events could potentially

facilitate increased pathogen replication and dissemination in the host at later stages of

infection. Indeed, we observed similarly high levels of host cell cytotoxicity in response to

ST10, ST74 and related isolates at 24 hpi, indicating that ST74 likely induce cell death via

alternative mechanisms once intracellular bacterial levels become unsustainable within the

macrophage. Further investigation is required to identify and validate specific genomic factors

that may contribute to the mechanisms of enhanced host cell survival, in particular inhibition

of inflammasome activation and subsequent IL-1ß release during infection with ST74 isolates.

Indeed, a pangenome analysis of the ST10 and ST74 lineages included in phenotypic assays

have revealed multiple SPI and non-SPI-related genes that are unique to the ST74 lineage

(Supplementary Figure 11), >900 of which remain uncharacterised. A systematic analysis of

these unique genes, including comparison with other invasive NTS serovars, protein function

prediction and targeted mutagenesis may reveal genes of interest that contribute to pathogen

survival in the hostile immune environment of a macrophage.

To date, SPI-19 has been identified in NTS serovars Dublin, Weltevreden, Agona, Gallinarum

and Enteritidis (22), however the lack of SPI-19 in the *S.* Dublin ST74 lineage (Figure 4C and

E, Supplementary Figure 7, Supplementary Table 6) is an interesting observation and may

suggest a specific adaptation that supports increased replicative ability in phagocytes. In

previous studies, *S.* Dublin SPI-19 has been associated with colonisation and manipulation of

intestinal microbiome in poultry (21,55), as well as contributing to oral (but not systemic)

virulence in mice (55). Collectively, these outcomes suggest a role for SPI-19 at the mucosal-

luminal interface of the intestinal tract, possibly contributing to colonisation of intestinal

epithelial cells. Given that ST74 isolates replicate so efficiently in macrophages, it may be that

they simply do not require the added benefit of SPI-19-mediated mucosal colonisation and rely

solely on their ability to access phagocytes via M cell or dendritic cell uptake at the apical

surface of the intestine. The fact that ST74 isolates have a much larger pangenome than ST10

isolates, may also contribute to successful colonisation, replication and survival within the host

and thus compensate for the lack of SPI-19 function. Regardless, this work highlights how little

we know about the specific host-pathogen interactions of NTS and what specifically drives the

success of potential emerging lineages of iNTS.

This study has both strengths and limitations. It represents the most comprehensive genomic

study of *S.* Dublin to date, systematically characterising antimicrobial, heavy metal and biocide

resistance determinants, in addition to exploring the virulome in the global population of *S.*

Dublin. Importantly, our phenotypic data reveals the caution required when making generalised

statements about infection outcomes and virulence mechanisms of *Salmonella* serovars. Our

data clearly shows that within the Dublin serovar, there are distinct phenotypic differences

observed across sequence types and host cell types, all of which could impact significantly on

future development of effective therapeutics or vaccine strategies for NTS infections. The key

limitation of this dataset is the geographical distribution bias with an overrepresentation of

North American genomes and the lack of data from other regions, such as Asia. However, this

is a common limitation of many genomic studies and often results in a bias towards high

income countries that have greater resources for whole genome sequencing. This situation may

change in the future as many countries build their public health genomics capacity.

In conclusion, we provide genomic insights into the evolutionary, resistance and virulence

dynamics in *S.* Dublin circulating globally. The increasing incidence of iNTS caused by *S.*

Dublin may warrant this pathogen becoming a notifiable disease, similar to typhoid and

paratyphoid fever. The distinct virulence and AMR profiles of *S.* Dublin globally warrant

revisitation of the current classification scheme for NTS, in keeping with the rapid and ongoing

evolution of these significant pathogens. Importantly, our data further highlight the need for a

One Health approach to genomic surveillance.

METHODS

Setting and dataset

For Australian isolates, whole genome sequencing (WGS) was performed on all viable *S.*

Dublin isolates from human clinical samples received at the Microbiological Diagnostic Unit

Public Health Laboratory (MDU PHL), the primary bacterial reference laboratory for the State

of Victoria in Australia. In total, 53 *S.* Dublin genomes between 2000-2019 were included. For

publicly available data, we included *S.* Dublin sequences that had associated metadata

(geographical origin and date of sample collection), providing a total of 1,250 publicly

available sequences. Data were collected in accordance with the Victorian Public Health and

Wellbeing Act 2008. Ethical approval was received from the University of Melbourne Human

Research Ethics Committee (study number 1954615.3).

Genome quality control, assembly and serovar prediction

Genomes were assembled and quality controlled using the Nullarbor pipeline v2.0.20191013

(https://github.com/tseemann/nullarbor). Briefly, Skesa v2.4.0 (61) was used to assemble the

genomes before determining the sequence type (ST) using the multilocus sequence typing

(MLST) scheme '*senterica*' within mlst v2.19.0 (https://github.com/tseemann/mlst). Kraken

v1.1.1 (62) was then used to confirm the taxonomic classification. Snippy v4.6.0

(https://github.com/tseemann/snippy) with parameters set to a 0.90 variant rate proportion

against a minimum coverage of 10X mapped the reads against a well-studied reference

genome, *S.* Dublin CT_02021853 (NC_011205.1). To be included in subsequent analysis,

genomes had to pass the following quality control parameters: mapped to $\geq$ 80% of the

reference genome; had a $\geq$ 80% taxonomic sequence similarity to *S. enterica*; N50 >25000,

and <300 contigs. To verify that each genome was *S.* Dublin, all sequences were typed using

Salmonella in silico Typing Resource (SISTR) v1.1.1 (63). The electrophoretic type of each

isolate was determined using Antimicrobial Resistance Identification by Assembly (ARIBA)

v2.14.5 (64) against the reference Du 2 (Genbank accession: M84972.1).

Core genome phylogenomic analysis

Variant calls from snippy were used to construct a core genome single nucleotide

polymorphism (SNP) alignment using snippy-core v4.6.0

(https://github.com/tseemann/snippy). Phaster v2020-12-22 (65) was used to create a BED file

containing prophage regions. The option '—mask' in snippy-core was used to exclude the

identified prophage regions from the BED file. Constant sites from the reference genome were

obtained using the script iqtree-calc_const_sites.sh (https://github.com/MDU-PHL/mdu-

tools/blob/master/bin/iqtree-calc_const_sites.sh). A maximum-likelihood (ML) phylogeny was

inferred using IQ-TREE v2.1.0 (66) with parameters set to a GTR+G4 nucleotide substitution

model with constant sites and ultrafast bootstrapping (67) to 1000 replicates. The resulting ML

tree was then visualised using *ggtree* v2.3.3 R package (68). Population clades were defined

using rhierBAPS v1.1.3 to one level on an alignment of 2,957 core SNPS of 1,295 isolates in

the ST10 *S.* Dublin lineage (69)

In silico prediction of resistance determinants and plasmid replicons

To screen for quinolone point mutations and acquired AMR determinants, ARIBA and

AbriTAMR v0.2.2 (https://github.com/MDU-PHL/abritamr) were used, respectively. The

AMR database used for this study was the NCBI AMRFinderPlus database v2020-01-22.1.

Each isolate's susceptibility profile was classified as either susceptible; resistant to 1-2

antimicrobials, or multidrug resistant (MDR). Heavy metal and biocide determinants were

identified by screening the AMRFinder database v2020-07-16.2 with AMRFinder v3.8.4 (70).

Replicon types were determined using ABRicate v1.0.1 (https://github.com/tseemann/abricate)

at a threshold of 80% nucleotide identity and 95% coverage, and screening against the

Plasmidfinder database (71). A strict threshold was applied to ensure the detection of only

unambiguous replicon types. To establish the most common resistant profiles, co-occurrence

matrices of resistant determinants were constructed in R using *igraph* v1.2.4.1 (72).

Virulome analysis

Virulence genes were identified using ABRicate at 90% nucleotide identity/coverage against the virulence factor database (vfdb) v2018-03-18 (73). Databases from previous studies (16,26,34,74–78) were also interrogated and collated. This included a database containing additional virulent determinants and *Salmonella* pathogenicity islands (SPIs) -1, -2, -6 and -19 (Supplementary Table 7). Using ABRicate, the presence of SPIs -1, -2, -6 and -19 were identified at 80% ID and 10% COV. ABRicate was also used to screen the manually created SPI determinant database for genes relevant to SPI-1, -2, -6 and -19 with a threshold of 90% identity/coverage (Supplementary Table 7). To quantify invasiveness, index scores were calculated *in silico* for each isolate using Wheeler *et al.*'s tool developed in 2018 for inferring the likelihood of invasiveness (http://www.github.com/UCanCompBio/invasive_*Salmonella*), with a higher score for an isolate deemed more invasive (37). Prior to running the tool, a set of reference genomes consisting of invasive and non-invasive *Salmonella* serovars were used as a validation set (Supplementary Table 8). The validation results were then visualised using *ggtree*.

Bayesian phylodynamic analysis

A subset of 660 *S*. Dublin genomes from the main cluster was used for BEAST v1.10.4 (79) analysis. This subset comprised all Australian genomes and a proportion of international *S*. Dublin genomes. The sampling strategy for selecting representative international genomes involved capturing all geographical regions and sources harbouring distinct susceptibility profiles. When two genomes demonstrated the same AMR profile and were from the same region and source, the oldest genome was preferred. Recombination sites were filtered using Gubbins v2.4.1 (80). A core genome SNP alignment of these genomes was created using

snippy-core v4.6.0 and then a ML phylogeny was inferred as above. Temporal signal was

assessed using TempEst v1.5.3 (81) by assessing the best-fitting root using the heuristic

residual mean squared method. The core alignment was then used to create an xml file in

BEAUti v1.10.4 using the GTR+G4 substitution model. To identify the most appropriate model

for our dataset, a combination of using either a "relaxed lognormal" or a "strict" molecular

clock with either a "coalescent constant" or a "coalescent exponential" population prior was

tested. For all models, the Markov chain Monte Carlo chain length was set to 300,000,000 and

sampling was set to every 20,000 trees with the marginal likelihood estimation set to

"generalised stepping stone (GSS)" to provide statistical support. The analysis was repeated in

triplicates with and without tip dates to ensure there was no temporal bias. Bayesian evaluation

of temporal signal (BETS) was used to determine whether a temporal signal was present (82).

To optimise computational resource and time, BEAST v1.10.4 was used with BEAGLE v3.0.2

(83). A Maximum Clade Credibility (MCC) tree from the highest supported model using GSS

was extracted using TreeAnnotator v1.10.4 and median heights. The resulting MCC tree was

then visualised using *ggtree*.

Long read sequencing

The Oxford Nanopore Technologies MinION platform was used to obtain complete genomes

of six Australian *S*. Dublin genomes (Supplementary Table 9). These genomes, comprised of

ST10 lineage clade 1 (AUSMDU00056868 and AUSMDU00035676,) and ST74

(AUSMDU00016676 and AUSMDU00021231), were selected based on their AMR profile,

virulome and index scores. To prepare new DNA libraries for long read sequencing, genomes

were freshly re-cultured. DNA was extracted using the Virus/Pathogen DSP midi kit on the

QiaSymphony SP instrument. Sequencing libraries were prepared using a Ligation sequencing

kit (SQK-LSK109) with native barcode expansion and sequenced for 48 hours on a GridION

Mk2 instrument using a R9.4 flowcell. WGS was performed using the Illumina NextSeq

500/550 platform (Illumina, San Diego, CA). Demultiplexed long reads and corresponding

WGS reads were subjected to QC to exclude unreliable reads using fastp v0.20.1 (84) and

Filtlong v0.2.1 (https://github.com/rrwick/Filtlong), respectively. Long reads were assembled

using Trycycler v0.4.1 (85). The final sequence was polished using Medaka v1.4.3

(https://github.com/nanoporetech/medaka). To ensure small plasmids were not excluded during

the initial QC step, hybrid assemblies using both long and WGS reads were generated

Unicycler v0.4.4 (86). Trycycler assemblies were checked against Unicycler assemblies with

Bandage v0.8.1 (87). The final genome assemblies were confirmed as *S*. Dublin using SISTR

and annotated using Prokka v1.14.6 (88) (Supplementary Table 9).

Chromosomal assemblies were aligned to complete genomes of known *S*. Dublin references

CT_02021853 (CP001144.1, without Vi flagella) and CFSAN000518 (CP074226.1, with Vi

flagella) and visualised using Blast Ring Image Generator (BRIG) v0.95 (89). Plasmid

assemblies were screened using PlasmidFinder (90) and mob-typer v3.1.0 (91) to identify

replicon and relaxase types, respectively. Plasmid profiles were subjected to BLASTn against

previously identified IncX1/IncFII(S) virulence plasmids reported in *S*. Dublin and IncN

plasmids associated with serovars of *Salmonella* (Supplementary Table 5). Genome

comparisons of plasmids demonstrating the top four highest BLASTn max scores were

visualised using BRIG. Prokka was used to annotate the regions of interest, while Easyfig

v2.2.2 (92) was used to compare regions of the genome. Regions of the genome were

visualised using the R package *gggenomes* v0.9.12.9000

(https://github.com/thackl/gggenomes).

Human cell line infections

Human macrophage (THP-1) (ATCC® TIB-202™) and colonic epithelial (HT-29) (ATCC®

HTB-38™) cell lines were maintained in Roswell Park Memorial Institute (RPMI) 1640 media

+ 200 mM GlutaMAX (Life Technologies) supplemented with 10% v/v Fetal Bovine Serum

(FBS; Bovogen) and grown in a humidified 5% CO2 37ºC incubator. THP-1 monocytes were

differentiated into macrophages with 50 ng/mL phorbol 12-myristiate-12 acetate (PMA,

Sigma-Aldrich) for 72 hours prior to infection.

Single colonies of each *S*. Dublin isolate were inoculated into 10 mL Luria Bertani (LB) broth

and grown overnight at 37ºC at 200 rpm, then sub-cultured for 3 hours in the same conditions

in 5 mL LB. Colony forming units/mL were measured by absorbance at 600 nM (OD_{600}). THP-

1 and HT-29 cells were infected at a multiplicity of infection (MOI) of 10 in RPMI in triplicate

wells of a 96-well plate and centrifuged at 525 x g for 5 minutes to synchronize bacterial

uptake. Infected cells were incubated at 37ºC for 30 minutes, then infective media removed

and replaced with 100 µg/mL gentamicin in 10% v/v FBS/RPMI to inhibit extracellular

bacterial growth. 0 hours post infection (hpi) samples were collected at this time to measure

initial bacterial uptake. Cells were then incubated in high dose gentamicin for 1 hour, then

replaced with 10 µg/mL gentamicin in 10% v/v FBS/RPMI for the remainder of the infection.

For THP-1 cells, samples were collected at time 0-, 3-, 9- and 24-hours post-infection. For HT-

29 cells, samples were collected at 0- and 24-hours post-infection.

Cytotoxicity of infection was measured by lactate dehydrogenase (LDH) release by infected

cells before lysis for intracellular bacterial enumeration. Briefly, 35 uL of infected (and

uninfected) cell culture supernatants were collected from each well and LDH quantified as per

manufacturer's instructions (Promega), with percent cytotoxicity calculated as a proportion of

lysed control cells and blank media. Following this, infected cells were washed three times

with warm phosphate-buffered saline (PBS) to remove any lingering extracellular bacteria,

then lysed in 0.1% Triton X-100/PBS and plated out onto LB agar after serial dilution. Bacterial

enumeration and cell viability assays were performed over 3 biological replicates, each

performed independently, with a new passage of host cells, on a separate day. Within each

biological replicate, technical triplicates were performed for each isolate. All statistical

analyses were performed using Prism software (GraphPad Software v9.0) and determined by

nested students t-test, which allowed for inclusion of individual isolates in sub-groups of each

sequence type (ST10 vs ST74).

Measurement of cellular responses

THP-1 cells were infected at an MOI of 10 with a representative isolate from each of the ST10

and ST74 lineages as described above, then lysed in radio-immunoprecipitation (RIPA) buffer

(20 mM Tris pH 8.0, 0.5 mM EDTA, 150 mM NaCl, 1% v/v Triton X-100, 0.5% w/v sodium

deoxycholate, 0.1% w/v SDS) supplemented with Complete protease and phosphatase inhibitor

cocktails (Roche) at 0-, 3-, 8- and 24 hpi . Total protein concentration was quantified by

Bicinchoninic acid (BCA) assay and 15 μg separated on 4-12% Bolt Bis-Tris polyacrylamide

gels (Thermo Scientific), transferred to nitrocellulose (iBlot, Thermo Scientific) and blotted

for cleaved IL-1β (clone D3A3Z, Cell Signalling). Cell culture supernatants were also collected

from 96-well plate infections at 9 hpi and secreted IL-1β measured by DuoSet© Human IL-

1β/IL-1F2 ELISA as per manufacturer's instructions (R&D Systems). A core genome SNP ML

phylogeny was constructed on the Australian ST10 lineage to demonstrate the genetic

relationship of representative ST10 isolates that underwent *in vitro* phenotypic assays.

Pangenome analysis

The pangenomes of the isolates used in the macrophages and epithelial infections were

analysed using Panaroo v1.3.4 (93) set to default threshold settings, under strict mode and with

the removal of invalid genes on the draft genome assemblies (Supplementary Table 11).

DATA AVAILABILITY

Paired end reads of public data were obtained from either the European Nucleotide Archive

(ENA) or National Center for Biotechnology Information (NCBI). The Australian short read

data and complete genomes for the four Australian *S.* Dublin isolates are available at the NCBI

Sequence Read Archive under BioProject PRJNA319593. Details of all 1,303 genomes

included in the study are available in Supplementary Tables 1-11. These supplementary tables

also includes *i*) the resistance, virulence, Vi and *fliC* profiles of all 1,303 genomes; *ii*) the

genbank accession numbers of all reference genomes, virulence determinants and IncN

plasmids used in this study, and *iii*) all raw CFU, cytotoxicity and ELISA values from the

phenotypic analyses. The interactive tree is available in Microreact under

https://microreact.org/project/sdublinpaper).

CODE AVAILABILITY

The machine learning tool developed by Wheeler *et al.* for predicting the likelihood of

invasiveness is available in a public Github depository

http://www.github.com/UCanCompBio/invasive_*Salmonella*.

REFERENCES

1. Stanaway JD, Parisi A, Sarkar K, Blacker BF, Reiner RC, Hay SI, et al. The Global
Burden of Non-typhoidal *Salmonella* Invasive Disease: A Systematic Analysis for the
Global Burden of Disease Study 2017. The Lancet Infectious Diseases.
2019;19(12):1312–24.

2. Parisi A, Crump JA, Stafford R, Glass K, Howden BP, Kirk MD. Increasing incidence of
invasive nontyphoidal *Salmonella* infections in Queensland, Australia, 2007-2016. PLOS
Neglected Tropical Diseases. 2019 Mar 18;13(3):e0007187.

3. Mughini-Gras L, Pijnacker R, Duijster J, Heck M, Wit B, Veldman K, et al. Changing
Epidemiology of Invasive Non-typhoid *Salmonella* Infection: A Nationwide Population-
based Registry Study. Clinical Microbiology and Infection. 2020;26(7):941.e9-941.e14.

4. Kingsley RA, Msefula CL, Thomson NR, Kariuki S, Holt KE, Gordon MA, et al.
Epidemic Multiple Drug Resistant *Salmonella* Typhimurium Causing Invasive Disease in
Sub-Saharan Africa have a Distinct Genotype. Genome Res. 2009;19(12):2279–87.

5. Cheng RA, Eade CR, Wiedmann M. Embracing Diversity: Differences in Virulence
Mechanisms, Disease Severity, and Host Adaptations Contribute to the Success of
Nontyphoidal *Salmonella* as a Foodborne Pathogen. Front Microbiol. 2019 Jun
26;10:1368.

6. Williamson DA, Lane CR, Easton M, Valcanis M, Strachan J, Veitch MG, et al.
Increasing Antimicrobial Resistance in Nontyphoidal *Salmonella* Isolates in Australia
from 1979 to 2015. Antimicrob Agents Chemother. 2018 Jan 25;62(2). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5786757/

7. Mangat CS, Bekal S, Avery BP, Côté G, Daignault D, Doualla-Bell F, et al. Genomic
Investigation of the Emergence of Invasive Multidrug-Resistant *Salmonella enterica*
Serovar Dublin in Humans and Animals in Canada. Antimicrobial Agents and
Chemotherapy. 2019 Jun 1;63(6). Available from:
https://aac.asm.org/content/63/6/e00108-19

8. Reichel MP, Wahl LC, Hill FI. Review of Diagnostic Procedures and Approaches to
Infectious Causes of Reproductive Failures of Cattle in Australia and New Zealand. Front
Vet Sci. 2018 Oct 2;5.

9. Kirchner M, McLaren I, Clifton-Hadley FA, Liebana E, Wales AD, Davies RH. A
Comparison Between Longitudinal Shedding Patterns of *Salmonella* Typhimurium and
Salmonella Dublin on Dairy Farms. Veterinary Record. 2012 Aug 25;171(8):194–194.

10. McDonough PL, Fogelman D, Shin SJ, Brunner MA, Lein DH. *Salmonella enterica*
Serotype Dublin Infection: an Emerging Infectious Disease for the Northeastern United
States. J Clin Microbiol. 1999 Aug;37(8):2418–27.

11. Davidson KE, Byrne BA, Pires AFA, Magdesian KG, Pereira RV. Antimicrobial
resistance trends in fecal *Salmonella* isolates from northern California dairy cattle
admitted to a veterinary teaching hospital, 2002-2016. PLoS One. 2018 Jun 28;13(6).
Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6023112/

12. Fenske GJ, Thachil A, McDonough PL, Glaser A, Scaria J. Geography Shapes the
Population Genomics of *Salmonella enterica* Dublin. Genome Biol Evol. 2019 Aug
1;11(8):2220–31.

13. Sia CM, Baines SL, Valcanis M, Lee DYJ, Gonçalves da Silva A, Ballard SA, et al.
Genomic Diversity of Antimicrobial Resistance in Non-typhoidal *Salmonella* in Victoria,
Australia. Microbial Genomics. 2021;7(12):000725.

14. Paudyal N, Pan H, Elbediwi M, Zhou X, Peng X, Li X, et al. Characterization of
Salmonella Dublin isolated from bovine and human hosts. BMC Microbiology. 2019 Oct
16;19(1):226.

15. Harvey RR, Friedman CR, Crim SM, Judd M, Barrett KA, Tolar B, et al. Epidemiology of *Salmonella enterica* Serotype Dublin Infections among Humans, United States, 1968–2013 - Volume 23, Number 9—September 2017 - Emerging Infectious Diseases journal - CDC. [cited 2020 Nov 24]; Available from: https://wwwnc.cdc.gov/eid/article/23/9/17-0136_article

16. Hsu CH, Li C, Hoffmann M, McDermott P, Abbott J, Ayers S, et al. Comparative Genomic Analysis of Virulence, Antimicrobial Resistance, and Plasmid Profiles of *Salmonella* Dublin Isolated from Sick Cattle, Retail Beef, and Humans in the United States. Microbial Drug Resistance. 2019 May 31;25(8):1238–49.

17. Magiorakos AP, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, et al. Multidrug-Resistant, Extensively Drug-resistant and Pandrug-resistant Bacteria: An International Expert Proposal for Interim Standard Definitions for Acquired Resistance. Clinical Microbiology and Infection. 2012;18(3):268–81.

18. Langridge GC, Fookes M, Connor TR, Feltwell T, Feasey N, Parsons BN, et al. Patterns of Genome Evolution that have Accompanied Host Adaptation in *Salmonella*. Proc Natl Acad Sci USA. 2015;112(3):863–8.

19. Achtman M, Wain J, Weill FX, Nair S, Zhou Z, Sangal V, et al. Multilocus Sequence Typing as a Replacement for Serotyping in *Salmonella enterica*. PLoS Pathogens. 2012;8(6):e1002776.

20. Selander RK, Smith NH, Li J, Beltran P, Ferris KE, Kopecko DJ, et al. Molecular Evolutionary Genetics of the Cattle-Adapted Serovar *Salmonella* Dublin. J Bacteriol. 1992;174(11):3587–92.

21. Pezoa D, Blondel CJ, Silva CA, Yang HJ, Andrews-Polymenis H, Santiviago CA, et al. Only One of the Two Type VI Secretion Systems Encoded in the *Salmonella enterica* Serotype Dublin Genome is Involved in Colonization of the Avian and Murine Hosts. Veterinary Research. 2014;45(1):2.

22. Blondel CJ, Jiménez JC, Contreras I, Santiviago CA. Comparative Genomic Analysis Uncovers 3 Novel Loci Encoding Type Six Secretion Systems Differentially Distributed in *Salmonella* Serotypes. BMC Genomics. 2009 Aug 4;10:354.

23. Blondel CJ, Amaya FA, Bustamante P, Santiviago CA, Pezoa D. Identification and distribution of new candidate T6SS effectors encoded in *Salmonella* Pathogenicity Island 6. Front Microbiol. 2023 Aug 17;14. Available from: https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2023.1252344/full

24. Amaya FA, Blondel CJ, Barros-Infante MF, Rivera D, Moreno-Switt AI, Santiviago CA, et al. Identification of Type VI Secretion Systems Effector Proteins That Contribute to Interbacterial Competition in *Salmonella* Dublin. Frontiers in Microbiology. 2022;13. Available from: https://www.frontiersin.org/article/10.3389/fmicb.2022.811932

25. Vilela FP, Gomes CN, Paziani MH, Braz VS, Rodrigues D dos P, Costa RG, et al. Virulence traits and expression of *bstA, fliC* and *sopE2* in *Salmonella* Dublin strains

isolated from humans and animals in Brazil. Infection, Genetics and Evolution. 2020;80:104193.

26. Hammarlöf DL, Kröger C, Owen SV, Canals R, Lacharme-Lora L, Wenner N, et al. Role of a Single Noncoding Nucleotide in the Evolution of an Epidemic African Clade of *Salmonella*. Proc Natl Acad Sci U S A. 2018;115(11):E2614–23.

27. Herrero-Fresno A, Espinel IC, Spiegelhauer MR, Guerra PR, Andersen KW, Olsen JE. The Homolog of the Gene *bstA* of the BTP1 Phage from *Salmonella enterica* Serovar Typhimurium ST313 Is an Antivirulence Gene in *Salmonella enterica* Serovar Dublin. Infection and Immunity. 2018;86(1).

28. Mohammed M, Cormican M. Whole Genome Sequencing Provides Insights into the Genetic Determinants of Invasiveness in *Salmonella* Dublin. Epidemiology and Infection. 2016;144(11):2430–9.

29. De Sousa Violante M, Podeur G, Michel V, Guillier L, Radomski N, Lailler R, et al. A Retrospective and Regional Approach Assessing the Genomic Diversity of *Salmonella* Dublin. NAR Genom Bioinform. 2022 Jul 9;4(3):lqac047.

30. Ingle DJ, Ambrose RL, Baines SL, Duchene S, Gonçalves da Silva A, Lee DYJ, et al. Evolutionary Dynamics of Multidrug Resistant *Salmonella enterica* serovar 4,[5],12:i:- in Australia. Nature Communications. 2021;12(1):4786.

31. Duchêne S, Holt KE, Weill FX, Le Hello S, Hawkey J, Edwards DJ, et al. Genome-scale Rates of Evolutionary Change in Bacteria. Microb Genom. 2016 Nov 30;2(11):e000094.

32. Silva C, Puente JL, Calva E. *Salmonella* Virulence Plasmid: Pathogenesis and Ecology. Pathogens and Disease. 2017;75(6).

33. Harmer CJ, Hall RM. IS26 Family Members IS257 and IS1216 Also Form Cointegrates by Copy-In and Targeted Conservative Routes. mSphere. 2020;5(1):811.

34. Mohammed M, Le Hello S, Leekitcharoenphon P, Hendriksen R. The invasome of *Salmonella* Dublin as revealed by whole genome sequencing. BMC Infect Dis. 2017 Aug 4;17. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5544996/

35. Hong SF, Chiu CH, Chu C, Feng Y, Ou JT. Complete Nucleotide Sequence of a Virulence Plasmid of *Salmonella enterica* Serovar Dublin and Its Phylogenetic Relationship to the Virulence Plasmids of Serovars Choleraesuis, Enteritidis and Typhimurium. FEMS Microbiology Letters. 2008;282(1):39–43.

36. Chu C, Feng Y, Chien AC, Hu S, Chu CH, Chiu CH. Evolution of Genes on the *Salmonella* Virulence Plasmid Phylogeny Revealed From Sequencing of the Virulence Plasmids of *S. enterica* Serotype Dublin and Comparative Analysis. Genomics. 2008;92(5):339–43.

37. Wheeler NE, Gardner PP, Barquist L. Machine Learning Identifies Signatures of Host Adaptation in the Bacterial Pathogen *Salmonella enterica*. PLoS Genetics. 2018;14(5):e1007333.

38. Lambert MA, Smith SG. The PagN protein of *Salmonella enterica* Serovar Typhimurium
is an Adhesin and Invasin. BMC Microbiology. 2008;8(1):142.

39. Folkesson A, Advani A, Sukupolvi S, Pfeifer JD, Normark S, Löfdahl S. Multiple
Insertions of Fimbrial Operons Correlate with the Evolution of *Salmonella* Serovars
Responsible for Human Disease. Molecular Microbiology. 1999;33(3):612–22.

40. Herrero-Fresno A, Wallrodt I, Leekitcharoenphon P, Olsen JE, Aarestrup FM, Hendriksen
RS. The Role of the *st313-td* Gene in Virulence of *Salmonella* Typhimurium ST313.
PLOS ONE. 2014 Jan 3;9(1):e84566.

41. Sabbagh SC, Forest CG, Lepage C, Leclerc JM, Daigle F. So Similar, Yet So Different:
Uncovering Distinctive Features in the Genomes of *Salmonella enterica* Serovars
Typhimurium and Typhi. FEMS Microbiology Letters. 2010;305(1):1–13.

42. Chen LM, Kaniga K, Galán JE. *Salmonella* spp. are cytotoxic for cultured macrophages.
Molecular Microbiology. 1996;21(5):1101–15.

43. Miao EA, Mao DP, Yudkovsky N, Bonneau R, Lorang CG, Warren SE, et al. Innate
immune detection of the type III secretion apparatus through the NLRC4 inflammasome.
Proceedings of the National Academy of Sciences. 2010 Feb 16;107(7):3076–80.

44. Bergsbaken T, Fink SL, Cookson BT. Pyroptosis: host cell death and inflammation. Nat
Rev Microbiol. 2009 Feb;7(2):99–109.

45. Doerflinger M, Deng Y, Whitney P, Salvamoser R, Engel S, Kueh AJ, et al. Flexible
Usage and Interconnectivity of Diverse Cell Death Pathways Protect against Intracellular
Infection. Immunity. 2020 Sep;53(3):533-547.e7.

46. Cheng G, Ning J, Ahmed S, Huang J, Ullah R, An B, et al. Selection and Dissemination
of Antimicrobial Resistance in Agri-food Production. Antimicrob Resist Infect Control.
2019;8:158.

47. Srednik ME, Lantz K, Hicks JA, Morningstar-Shaw BR, Mackie TA, Schlater LK.
Antimicrobial Resistance and Genomic Characterization of *Salmonella* Dublin Isolates in
Cattle from the United States. PLOS ONE. 2021 Sep 21;16(9):e0249617.

48. O'Connell J. Australian Food Timeline. 1930 [cited 2022 Nov 11]. Australian food
history timeline - Brahman cattle imported. Available from:
https://australianfoodtimeline.com.au/brahman-cattle-imported/

49. Fritz HM, Pereira RV, Toohey-Kurth K, Marshall E, Tucker J, Clothier KA. *Salmonella
enterica* Serovar Dublin from Cattle in California from 1993–2019: Antimicrobial
Resistance Trends of Clinical Relevance. Antibiotics (Basel). 2022;11(8):1110.

50. Izzo M, Mohler V, House J. Antimicrobial Susceptibility of *Salmonella* Isolates
Recovered From Calves with Diarrhoea in Australia. Australian Veterinary Journal.
2011;89(10):402–8.

51. McMillan EA, Jackson CR, Frye JG. Transferable Plasmids of *Salmonella enterica*
Associated With Antibiotic Resistance Genes. Frontiers in Microbiology. 2020;11.
Available from: https://www.frontiersin.org/article/10.3389/fmicb.2020.562181

52. Zhang Y, Lei CW, Chen X, Yao TG, Yu JW, Hu WL, et al. Characterization of IncC
Plasmids in Enterobacterales of Food-Producing Animals Originating From China. Front
Microbiol. 2020;11:580960.

53. Carattoli A. Resistance Plasmid Families in Enterobacteriaceae. Antimicrob Agents
Chemother. 2009;53(6):2227–38.

54. Rozwandowicz M, Brouwer MSM, Fischer J, Wagenaar JA, Gonzalez-Zorn B, Guerra B,
et al. Plasmids Carrying Antimicrobial Resistance Genes in Enterobacteriaceae. Journal
of Antimicrobial Chemotherapy. 2018;73(5):1121–37.

55. Schroll C, Huang K, Ahmed S, Kristensen BM, Pors SE, Jelsbak L, et al. The SPI-19
encoded type-six secretion-systems (T6SS) of *Salmonella enterica* serovars Gallinarum
and Dublin play different roles during infection. Veterinary Microbiology. 2019;230:23–
31.

56. Chattaway MA, Langridge GC, Wain J. *Salmonella* Nomenclature in the Genomic Era: A
Time for Change. Sci Rep. 2021;11(1):7494.

57. Hu X, Chen Z, Xiong K, Wang J, Rao X, Cong Y. Vi Capsular Polysaccharide: Synthesis,
Virulence, and Application. Critical Reviews in Microbiology. 2017;43(4):440–52.

58. Janis C, Grant AJ, McKinley TJ, Morgan FJE, John VF, Houghton J, et al. *In Vivo*
Regulation of the Vi Antigen in *Salmonella* and Induction of Immune Responses with an
in vivo-inducible Promoter. Infect Immun. 2011;79(6):2481–8.

59. Clare B. Inflammasome activation by *Salmonella*. Current Opinion in Microbiology.
2021 Dec 1;64:27–32.

60. Carden S, Okoro C, Dougan G, Monack D. Non-typhoidal *Salmonella* Typhimurium
ST313 Isolates that Cause Bacteremia in Humans Stimulate Less Inflammasome
Activation than ST19 Isolates Associated with Gastroenteritis. Pathogens and Disease.
2015;73(4):ftu023.

61. Souvorov A, Agarwala R, Lipman DJ. SKESA: Strategic K-mer Extension for
Scrupulous Assemblies. Genome Biology. 2018;19(1):153.

62. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using
exact alignments. Genome Biology. 2014 Mar 3;15(3):R46.

63. Yoshida CE, Kruczkiewicz P, Laing CR, Lingohr EJ, Gannon VPJ, Nash JHE, et al. The
Salmonella In Silico Typing Resource (SISTR): An Open Web-accessible Tool for
Rapidly Typing and Subtyping Draft *Salmonella* Genome Assemblies. PLoS One.
2016;11(1):e0147101.

64. Hunt M, Mather AE, Sánchez-Busó L, Page AJ, Parkhill J, Keane JA, et al. ARIBA:
Rapid Antimicrobial Resistance Genotyping Directly from Sequencing Reads. Microb
Genom. 2017;3(10). Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5695208/

65. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster
version of the PHAST phage search tool. Nucleic Acids Res. 2016 Jul 8;44(Web Server
issue):W16–21.

66. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: A Fast and Effective
Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol.
2015 Jan 1;32(1):268–74.

67. Minh BQ, Nguyen MAT, von Haeseler A. Ultrafast Approximation for Phylogenetic
Bootstrap. Mol Biol Evol. 2013 May 1;30(5):1188–95.

68. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. *ggtree*: An R Package for Visualization and
Annotation of Phylogenetic Trees with their Covariates and other Associated Data.
Methods in Ecology and Evolution. 2017;8(1):28–36.

69. Tonkin-Hill G, Lees JA, Bentley SD, Frost SDW, Corander J. RhierBAPS: An R
Implementation of the Population Clustering Algorithm hierBAPS. Wellcome Open Res.
2018;3:93.

70. Feldgarden M, Brover V, Haft DH, Prasad AB, Slotta DJ, Tolstoy I, et al. Using the NCBI
AMRFinder Tool to Determine Antimicrobial Resistance Genotype-Phenotype
Correlations Within a Collection of NARMS Isolates. bioRxiv. 2019;550707.

71. Carattoli A, Zankari E, García-Fernández A, Larsen MV, Lund O, Villa L, et al. In Silico
Detection and Typing of Plasmids using PlasmidFinder and Plasmid Multilocus Sequence
Typing. Antimicrobial Agents and Chemotherapy. 2014 Jul 1;58(7):3895–903.

72. Csardi G, Nepusz T. The igraph software package for complex network research.
InterJournal. 2006; Complex Systems:1695.

73. Chen L, Zheng D, Liu B, Yang J, Jin Q. VFDB 2016: Hierarchical and Refined Dataset
for Big Data Analysis—10 Years On. Nucleic Acids Res. 2016;44:D694–7.

74. Mansour MN, Yaghi J, El Khoury A, Felten A, Mistou MY, Atoui A, et al. Prediction of
Salmonella Serovars Isolated from Clinical and Food Matrices in Lebanon and Genomic-
based Investigation Focusing on Enteritidis Serovar. International Journal of Food
Microbiology. 2020;333:108831.

75. Suez J, Porwollik S, Dagan A, Marzel A, Schorr YI, Desai PT, et al. Virulence Gene
Profiling and Pathogenicity Characterization of Non-Typhoidal *Salmonella* Accounted for
Invasive Disease in Humans. PLoS One. 2013 Mar 7;8(3):e58449.

76. Sévellec Y, Vignaud ML, Granier SA, Lailler R, Feurer C, Le Hello S, et al. Polyphyletic
Nature of *Salmonella enterica* Serotype Derby and Lineage-Specific Host-Association
Revealed by Genome-Wide Analysis. Frontiers in Microbiology. 2018;9:891.

77. Hayward MR, Jansen VA, Woodward MJ. Comparative Genomics of *Salmonella enterica*
serovars Derby and Mbandaka, Two Prevalent Serovars Associated with Different
Livestock Species in the UK. BMC Genomics. 2013;14:365.

78. Kérouanton A, Hirchaud E, Rose V, Esnault E, Naquin D, Denis M. First Complete
Genome Sequence of a *Salmonella enterica* subsp. *enterica* Serovar Derby Strain
Associated with Pork in France. Genome Announc. 2015;3(4):e00853-15.

79. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian
Phylogenetic and Phylodynamic Data Integration Using BEAST 1.10. Virus Evolution.
2018;4(1). Available from: https://doi.org/10.1093/ve/vey016

80. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid
Phylogenetic Analysis of Large Samples of Recombinant Bacterial Whole Genome
Sequences Using Gubbins. Nucleic Acids Research. 2015;43(3):e15.

81. Rambaut A, Lam TT, Max Carvalho L, Pybus OG. Exploring the Temporal Structure of
Heterochronous Sequences using TempEst (formerly Path-O-Gen). Virus Evol.
2016;2(1):vew007.

82. Duchene S, Lemey P, Stadler T, Ho SYW, Duchene DA, Dhanasekaran V, et al. Bayesian
Evaluation of Temporal Signal in Measurably Evolving Populations. Molecular Biology
and Evolution. 2020;37(11):3363–79.

83. Ayres DL, Darling A, Zwickl DJ, Beerli P, Holder MT, Lewis PO, et al. BEAGLE: An
Application Programming Interface and High-Performance Computing Library for
Statistical Phylogenetics. Systematic Biology. 2012 Jan 1;61(1):170–3.

84. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018 Sep 1;34(17):i884–90.

85. Wick RR, Judd LM, Cerdeira LT, Hawkey J, Méric G, Vezina B, et al. Trycycler:
Consensus Long-read Assemblies for Bacterial Genomes. Genome Biology. 2021 Sep
14;22(1):266.

86. Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: Resolving Bacterial Genome
Assemblies From Short and Long Sequencing Reads. PLOS Computational Biology.
2017;13(6):e1005595.

87. Wick RR, Schultz MB, Zobel J, Holt KE. Bandage: interactive visualization of de novo
genome assemblies. Bioinformatics. 2015 Oct 15;31(20):3350–2.

88. Seemann T. Prokka: Rapid Prokaryotic Genome Annotation. Bioinformatics (Oxford,
England). 2014;30(14):2068–9.

89. Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. BLAST Ring Image Generator
(BRIG): Simple Prokaryote Genome Comparisons. BMC Genomics. 2011;12(1):402.

90. Carattoli A, Hasman H. PlasmidFinder and *In Silico* pMLST: Identification and Typing of
Plasmid Replicons in Whole-Genome Sequencing (WGS). Methods in Molecular
Biology (Clifton, NJ). 2020;2075:285.

91. Robertson J, Nash JHEY. MOB-suite: Software Tools for Clustering, Reconstruction and
Typing of Plasmids from Draft Assemblies. Microbial Genomics. 2018;4(8):e000206.

92. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer.
Bioinformatics. 2011 Apr 1;27(7):1009–10.

93. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing
Polished Prokaryotic Pangenomes with the Panaroo Pipeline. Genome Biology.
2020;21(1):180.

ACKNOWLEDGEMENTS

We thank all staff members of the Enterics and MDDI section at the Microbiological Diagnostic

Unit Public Health Laboratory (MDU PHL) for their technical assistance. We would also like

to thank Jake Lacey for his assistance uploading the reads to NCBI. The BEAST analysis for

this paper was undertaken using the LIEF HPC-GPGPU Facility hosted at the University of

Melbourne. This Facility was established with the assistance of LIEF Grant LE170100200.

AUTHOR INFORMATION

These authors contributed equally: Deborah A Williamson, Jaclyn S Pearson, Danielle J Ingle

Authors and Affiliations

Department of Microbiology and Immunology, The University of Melbourne at The Peter

Doherty Institute for Infection and Immunity, Melbourne, Victoria, Australia.

Cheryll M Sia, Benjamin P Howden, Danielle J Ingle

Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical

Research, Clayton, Melbourne, VIC, Australia.

Rebecca Ambrose, Jaclyn S Pearson

Department of Infectious Diseases, The University of Melbourne at the Peter Doherty

Institute for Infection and Immunity, Melbourne, VIC, Australia.

Deborah A Williamson

Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology

& Immunology, The University of Melbourne at The Peter Doherty Institute for Infection

and Immunity, Melbourne, Victoria, Australia.

Mary Valcanis, Patiyan Andersson, Susan A Ballard, Benjamin P Howden

Department of Microbiology, Monash University, Clayton, Victoria, Australia.

Jaclyn S Pearson

School of Medicine, University of St Andrews, St Andrews, Fife, Scotland.

Jaclyn S Pearson

Contributions

Study design and oversight: CMS, RA, DAW, JSP, DJI. Isolate collection and metadata: MV,

PA, SAB, BJP. Performed wet lab experiments: RA, JSP. Data analysis: CMS, RA, JSP, DJI.

Manuscript writing: CMS, RA, DAW, JSP, DJI. All authors contributed to manuscript editing.

ETHICS DECLARATIONS

Competing interests

The authors declare no competing financial interests.

 Figures and Tables

 Figure 1. Overview of *S.* Dublin genomes in study

Geographical distribution and clonal relationship of 1,303 *S.* Dublin genomes in the study. A) Number of *S.* Dublin genomes according to geographical region and time interval of collection. B) Distribution of human, animal and food *S.* Dublin genomes collected over time. C) Distribution of *S.* Dublin sequence types (ST) from each region. Each coloured circle correlates to an ST. Circles in green indicate no ST was identified. D) Unrooted phylogenetic tree of 1,303 *S.* Dublin genomes. Two main lineages of ST74 (*n*=8) and ST10 (*n*=1,295) shown in blue and pink respectively. E) Relatedness of known sequence types (STs) to ST10. The nodes are labelled with the ST and the branch lengths are indicative of the number of dissimilar locus variants to ST10.

 Figure 2. Estimation of divergence time

 Maximum clade credibility (MCC) tree of the 667 *S.* Dublin ST10 genomes. The time in years

 is shown on the x-axis. From left to right, the coloured heatmaps refer to the region of origin,

 the susceptibility profile, the BAPs clade, and the presence of IncC and IncN plasmid types.

1017

1018

1019

1020

1021

 Figure 3. Resistance profiles of *S*. Dublin genomes

Observed resistance profiles of *S*. Dublin genomes spanning three-time intervals for each region. A) Frequency of susceptible and resistant *S*. Dublin genomes from each time period B) Antimicrobial resistance profiles observed spanning three time periods. The colour of the circle corresponds to an antimicrobial class, while the height is relative to the number of genomes. C) Heavy metal and biocide resistance profiles observed. The colour of the circle corresponds to its respective heavy metal/biocide class, while the height is relative to the number of genomes.

Figure 4. Invasive factors associated with *S.* Dublin genomes

Correlation of *Salmonella* pathogenicity islands (SPIs) to *S.* Dublin's invasiveness. A)

Validation of the likelihood of invasiveness prediction tool using a variety of publicly available

genomes consisting of non-invasive and invasive *Salmonella* serovars (Supplementary Table

8). B) The correlation of index scores to the source and its sequence type based on the region.

* indicates a statistically significant p-value between the index scores of two sources. C)

Chromosomal assemblies of ST10 (blue) and ST74 (pink) genomes aligned to *S.* Dublin

chromosomal reference CT_02021853 (CP001144.1) with a lower limit of 80% identity D)

Aligned sequence of SPI-6 region of an ST10, AUSMDU00056868, and an ST74,

AUSMDU00016676, in comparison to SPI-6 region of reference CT_02021853. E) Aligned

sequence of SPI-19 region of an ST10, AUSMDU00056868, and an ST74,

AUSMDU00016676 in comparison to SPI-19 region of reference CT_02021853.

SPI, *Salmonella* pathogenicity island.

1045

1046

Figure 5. *S.* Dublin ST74 isolates demonstrate increased intracellular replication in

human macrophages over ST10 isolates

Differentiated THP-1 (human monocyte) cells were infected at an MOI:10 with selected ST10

and ST74 *S.* Dublin isolates. **A, B)** THP-1 cells were lysed, and intracellular bacteria

enumerated as fold change in replication and **C-F)** CFU/well at time 0, 3, 9 and 24 hpi. For **A)**,

each measurement at times 3, 9 and 24 hpi represents the fold change in CFU/well compared

to 0 hpi, the start of infection, recorded as a biological replicate (performed in technical

triplicate), with error bars indicating ± 1 standard deviation of $n = 3$ biological replicates. For

C-D), each dot represents CFU/well of a biological replicate (performed in technical triplicate),

with error bars indicating ± 1 standard deviation of $n = 3$ biological replicates. Statistical

significance was determined by nested student's t-test. MOI, multiplicity of infection, hpi,

hours post-infection, CFU, colony-forming units, FC, fold-change.

1060

1061

Figure 6. *S.* **Dublin ST74 isolates induce comparable cytotoxicity to ST10 isolates in human macrophages, despite increased intracellular growth**

A-C) Cell supernatants were assayed for LDH as a measure of cytopathic effects of infection, and % cytotoxicity was calculated by comparison with 100% lysed uninfected control cells. Samples were collected from THP-1 cells at 3, 9 and 24 hpi. Each point represents the % cytotoxicity of a biological replicate (performed in technical triplicate), with error bars indicating ± 1 standard deviation of $n = 3$ biological replicates. Statistical significance was determined by nested student's t-test. **D)** IL-1ß secretion from THP-1 cells infected for 9 hours with *S.* Dublin isolates. Error bars indicate ± 1 standard deviation of $n = 3$ biological replicates

and statistical significance determined by student's t-test **E)** Immunoblot of cleaved IL-1ß

(Asp166) in THP-1 cell lysates infected with *S.* Typhimurium reference strain SL1344,

S. Dublin ST10 (AUSMDU00016673), or *S.* Dublin ST74 (AUSMDU00016682) over 0, 3, 8

and 24 hours. **F)** PCA plot comparing replication rate, % cytotoxicity and IL-1β secretion from

THP-1 cells infected with *S.*Dublin isolates at 9 hpi. Each dot represents each biological

replicate of three (averaged from three technical replicates). .MOI, multiplicity of infection,

hpi, hours post-infection, CFU, colony-forming units, LDH, lactate dehydrogenase.

1078