Title: Charting Taxa in *Amanita* Section *Amidella* (*Basidiomycota* : *Amanitaceae*)

Authors:

1. Paulo Oliveira, ORCID 0000-0001-7979-6520
2. Ricardo Arraiano-Castilho, ORCID 0000-0001-8465-5909

Affiliations:

1. University of Évora, Biology Department, Évora, Portugal
1. Mediterranean Institute for Agriculture, Environment and Development (MED), Évora, Portugal
1. Research Centre in Biodiversity and Genetic Resources (CIBIO), Vairão, Portugal
2. University of Lausanne, Department of Ecology and Evolution, Lausanne, Switzerland

Corresponding author: Paulo Oliveira, oliveira@uevora.pt

Abstract

Species determination in the *Amidella* clade is notoriously difficult, because of the relative dearth of diagnostic characters and the rather common occurrence of homoplasies. This results in a substantial number of misnamed and unnamed collections, a misapprehension of the geographic range of known species, and a gross underestimation of the number of species it contains. To assess the diversity that should be considered as part of *Amidella*, DNA sequences available for this group were retrieved in public nucleotide databases, using a combination of approaches to achieve a comprehensive representation. Phylogenetic analysis based on the aligned ITS sequences, consistently with the results from other molecular markers (ncLSU, *RPB2*, *TEF1*, *BTUB*), suggests five major clades: one containing the type species *Amanita volvata*, another for *Amanita ponderosa* and allies, a third one (roughly half of all species) with *Amanita lepiotoides*, and two others without valid species yet. At species level, around 81 clades were delimited, of which only 16–17 can be assigned a valid name, with a few more corresponding to provisional taxa listed in the amanitaceae.org website. Up to three further species without assigned sequences might correspond to the proposed clades. The current evidence suggests a rather narrow geographic range for most of these clades. This study provides a phylogeny-arranged outlook of the worldwide distribution of *Amidella* species, and an infrasectional framework for optimizing taxonomic sampling and designing clade-specific molecular markers to assist in identification.

Keywords
Amanita, *Amidella*, taxonomy, molecular phylogeny, biogeography.

Acknowledgements
No funding was received for conducting this study.

Introduction

The creation of *Amidella* as a taxonomic group dates from Edouard-Jean Gilbert's initial proposal as a separate genus, within family *Amanitaceae* created by the same author, and subsequently as a subgenus in *Amanita* Pers. (Neville & Poumarat 2004). The name underscores the importance attached to the basidiospore amyloid reaction with iodine which the name refers to (Konrad and Maublanc 1948).

The current taxonomic arrangement within *Amanita* recognises *Amidella* at the section level in subgenus *Amanitina*, bracketed by sections *Roanokenses* and *Arenariae* (Cui et al. 2018). This is the culmination of a process whereby several groups with amyloid basidiospores segregated from the original concept of *Amidella* (Wolfe et al. 2012; Cui et al. 2018; Riccioni et al. 2019), leaving the name *Amidella* solely for a morphologically distinctive, relatively small group of species (Cui et al. 2018; Tulloss and Yang 2024).

Its demarcation from other congeneric groups is such that identification in the field as 'Amidella' can be straightforward, but in contrast reaching a determination at species level is very challenging (Tulloss and Yang 2024). It may be achieved at regional level, as done recently for Europe with prominent recourse to non-morphological characteristics such as soil, vegetation and phenology (Arraiano-Castilho et al. 2022). This notwithstanding, too many specimens have been assigned equivocally with species epithets such as *volvata*, *avellaneosquamosa* or *curtipes*. The problems arising from this are very difficult to tackle, and the only compelling argument against misidentifications have been the evident phylogenetic disparities (Moreno et al. 2008; Kim et al. 2013; Cho et al. 2015; Cui et al. 2018).

The ingrained notion of *Amidella* as a small group of species, in spite of some evidence against it (Cui et al. 2018; Tulloss and Yang 2024), favours the continued misapplication of known epithets and does not encourage the exploration of any meaningful taxonomic subdivisions. Partial phylogenetic sampling, not to mention mere blast searches, does little to address these issues; a comprehensive overview of the diversity of *Amidella* is definitely needed.

This study attempts such comprehensiveness based on publicly available molecular databases. The approach cannot cover a few known species lacking molecular representation, but it provides a novel picture of the group, establishing its boundaries, recognising more than 80 clades at species level, most of them undescribed, and demonstrating a phylogenetic substructure of great value for separating sympatric species. Knowledge gaps, such as the need for new morphological criteria and the underrepresentation of vast geographical regions, are clearly defined under the new taxonomic view provided here.

Materials and Methods

Adopted identifications

There are concerns about the correct application of some names. Following known authorities of genus *Amanita*, a tentative set of sequenced specimens are adopted as reference (Table 1).

{Table 1}

Sources and search strategies

Named sequences in Amanitaceae.org (Tulloss and Yang 2024)

The GenBank accession numbers for all sequences in Section *Amidella* listed in the details tab of each species were collected and used to retrieve from GenBank the full FastA records using the NCBI Batch Entrez interface (https://www.ncbi.nlm.nih.gov/sites/batchentrez last accessed December 2, 2024).

5.8S phylogeny

On February 25th 2023 a total of 7,045 sequences under the genus name *Amanita*, containing at least one of the nuclear ribosomal internal transcribed spacers (ITS), were downloaded from the GenBank database, and processed by ITSx (Bengtsson-Palme et al. 2013) in the PlutoF platform (Abarenkov et al. 2010), producing a total of 6,445 5.8S sequences that were aligned in UGene (Okonechnikov et al. 2012) and further processed with MEGA (Tamura et al. 2021), using the Maximum Likelihood tree construction method (see Outline of phylogenetic analysis, below). The resulting dendrogram was then searched for species epithets known to belong to Section *Amidella*. Three major clades were identified (Table 2), all containing many accessions without known epithets, and the GenBank accessions contained in them were collected, and used to retrieve from GenBank the full FastA records using the NCBI Batch Entrez interface as mentioned above.

{Table 2}

A preliminary phylogenetic analysis of the retrieved ITS sequences was made after this, to constitute an outgroup made only from sequences that branched out closer to the known *Amidella* sequences. Those that ended up in the outgroup were all from clade I (Table 2), all from Section *Roanokenses*.

Affinity with collected sequences

Two strategies were followed: first, using the voucher identifiers, an in-house wrapper script was developed to parse the output of esearch | efetch | xtract from Entrez Direct (Kans 2013) and used to retrieve the corresponding non-ITS records from GenBank (i.e. for the nuclear ribosomal 28S large subunit gene ncLSU, for the second largest protein subunit of the RNA polymerase II *RPB2*, for the translation elongation factor EF-1 alpha *TEF1*, and for the tubulin beta chain *BTUB*) that matched the specimens already available; second, using the BLAST homology algorithm, either in the PlutoF platform for the ITS sequences (including environmental samples), or in the NCBI interface for ncLSU, *RPB2*, *TEF1* and *BTUB*.

Additional sequences, summary and database checks

Several other sequences were spotted as they were referred to in specimen observation online platforms such as MushroomObserver.org, iNaturalist.org or MycoMap.com.

The final search took place on December 2, 2024. A total of 237 ITS, 132 ncLSU, 30 *RPB2*, 24 *TEF1* and 11 *BTUB* nucleotide accessions belonging to Section *Amidella* are included in this study (Online Resource 1). Excluded GenBank accessions: MK782535 (it contains errors and was suppressed at the submitter's request), OQ754998–9 (two partial reads of the same specimen that fit badly in the ncLSU phylogeny), duplicates KT779088, OP681731 and OP681759, as well as those of other genes with very little coverage (the 'etc' column in Online Resource 1). Redundant environmental sequences in the UNITE database version 10FU (Abarenkov et al. 2024) were not included either. The ncLSU sequences that appeared to be unique ("singletons") were entered (using the GenBank accession number) in the search page on ARB-SILVA (Quast et al. 2012) to check for their presence in the ncLSU r138.2 database and retrieve quality parameters dependent on ambiguous, repetitive or chimeric features. Except for accession KP012973, all sequences deposited till August 2018 were thus validated; the exception has a long string of 'N' that lies upstream from the ncLSU alignment, and it was decided to include it in the present analyses. Later singleton sequences in our ncLSU dataset were absent from the EUKARYOME 'blacklist'(Table S1 in Tedersoo et al. 2024). A Species Hypothesis code at 1.5% distance level (Kõljalg et al. 2013) was obtained for most of the ITS sequences.

Geographic references for each sequence were obtained primarily from the GenBank records, complemented by further information that could be retrieved from other sources (scientific publications, the reference website amanitaceae.org (Tulloss and Yang 2024) and online platforms).

Outline of the phylogenetic analyses

Most ITS sequences were processed in the ITSx programme (Nilsson et al. 2010; Bengtsson-Palme et al. 2013) in the PlutoF platform (Abarenkov et al. 2010), marking the 'Concat' option to obtain the ITS1:ITS2 concatenate. Those that, for being incomplete, were left unprocessed by ITSx, were aligned to the concatenated sequences, placing a gap in the latter that marked the intercurrent 5.8S gene sequence, thus allowing its safe removal.

The decision to use the ITS1:ITS2 concatenate fell out from the surprising heterogeneity of the 5.8S sequences within *Amidella* (Table 2). This hinted at the possibility that the 5.8S gene might distort the phylogenetic analyses based on the ITS region, and although it was later realised that, as expected from the fact that most of the phylogenetic signal is from the spacer regions, the differences of including the 5.8S gene were minimal (not shown), we had two important reasons to prefer the ITS1:ITS2 concatenate: first, to avoid potential inconsistency in the more critical clades; second, to have a phylogeny based purely on intronic sequences, which are in principle not subject to evolutionary constraints that act on the exonic ribosomal sequences, thus enhancing the contrast with the phylogenetic analyses based on the ncLSU.

All applications used for offline analyses were run in Windows 10. Alignments were made in the UGene platform versions 49–51 (Okonechnikov et al. 2012) using the built-in MUSCLE tool version 5 (Edgar 2022), complemented by visual corrections. The alignments were finally trimmed of

terminal tracts with low coverage. They can be retrieved in a figshare repository, https://doi.org/10.6084/m9.figshare.26352388.

The main phylogenetic reconstructions were based on the Bayesian Maximum Likelihood approach as implemented in the BEAST package version 1.10.4 (Suchard et al. 2018) supported by the BEAGLE acceleration libraries version 4.0.0 (Suchard and Rambaut 2009; Ayres et al. 2019). Alignments were converted in BEAUTi to XML input files using the default parameters, except that the substitution model was GTR+G+I with 5 gamma categories. Each BEAST output tree file was then analysed with TreeAnnotator using the default parameters to obtain the maximum likelihood tree. A few phylogenetic reconstructions were made using MEGA version 12.0.8 (Kumar et al. 2024). Splitstree CE version 6 (Huson and Bryant 2006) was used for Pairwise Homoplasy Index calculations with the default setting (Bruen et al. 2006).

To evaluate which sequences presumably belong to the same species/Operational Taxonomic Units (OTUs), and to allow the application of p-distance thresholds for tentative species, dendrograms were built with MEGA using the Minimal Evolution method with pairwise deletion, p-distances without corrections and uniform rates among sites (no Gamma correction). In the resulting 'tree sessions', using the 'Collapse Nodes' menu, option 'By Sequence Difference', the branches within a p-distance below a given (adjustable) threshold were condensed. The threshold was estimated for the ncLSU dataset using the barcode gap discovery approach (Puillandre et al. 2012), with the option of including phylogenetic trees in the output. The USEARCH software, version 11, was used as an alternative approach for delimiting centroid-based sequence clusters (Edgar 2010), the centroid-representative sequences aligned in UGene and the alignments used for a phylogenetic reconstruction in MEGA.

Except for the MEGA trees, which were drawn by the integrated tree session module, all trees were rendered in the iTOL website (Letunic and Bork 2024) from input files in text format produced by BEAST and ABGD.

Results

Clades at subsection and stirp level

The phylogenetic reconstruction based on the ITS concatenate subset suggests that section *Amidella* is subdivided into five main subclades (Figure 1a) that are named 'Volvatae', 'Sagittariae', 'Lepiotidae', 'Ponderosae' and 'Baakani'. This structure is also obtained for the ncLSU subset, except that Volvatae and Sagittariae are nested in a clade (with 57% posterior probability) that contains all Lepiotidae sequences (Figure 1b). The results also indicate further subdivisions with high support: Volvata, Peckiana and Fulvisquamea (in the Volvatae clade), Lepiotoides, Avellaneosquamosa, Subviscosa and Pseudomutabilis (Lepiotidae clade), Ponderosa, Claristriata and Pinophila (Ponderosae clade) (Table 3). However, two clades containing specimens classified in Lepiotoides and Subviscosa for the ITS concatenate dataset form separate branches in the phylogeny based on the ncLSU dataset.

{Figure 1}

{Table 3}

Further analyses of other markers (*RPB2*, *TEF1*, *BTUB*) as well as a concatenated ncLSU:*RPB2*:*TEF1* dataset, though limited by the reduced taxa coverage, support this structure (Online Resource 2), the only exception being the low posterior probability for Lepiotidae in the *RPB2* dataset (Table 3). Some of the 100% support values are based on a very limited taxon sampling. Pairwise Homoplasy Index calculations showed no evidence of recombination, with $\Phi_w > 0.05$ in all alignments (Bruen et al. 2006).

Clades at species level

A phylogenetic reconstruction based on the ITS concatenates was made without distance corrections (i.e., based on the percentage divergences calculated pairwise, and without gamma corrections for between-site heterogeneities) to allow the application of p-distance thresholds for tentative species. By adjusting the minimum distance threshold to separate species at different levels, the 1.25% level was chosen based on the observation that none of the previously recognised species were merged and very few (*Amanita rufobrunnescens*, *A. pseudovalens* and *A. hiltonii*) were split. As discussed in the respective Taxonomy sections below, the *subviscosa* s.l. branches are considered to be three species, while the sp. 'Naletali' branch merges two SH codes. At this level, a total of 67 branches were obtained within *Amidella* (Online Resource 3a). For the ncLSU dataset the choice of a 0.8% threshold was determined by the Automatic Barcode Gap Discovery method (Puillandre et al. 2012) and applied as in the ITS concatenate dataset (Online Resource 4). Although in this case a few mergers occur (for example, between *Amanita lanigera* and *A. brunneomaculata*), ad-hoc attempts at lower levels were judged less satisfactory because more taxa were split. The 0.8% threshold value was also applied for the UCLUST clustering algorithm in USEARCH (Edgar 2010) to provide an alternative phylogeny-independent delimitation at species level. Both ABGD and UCLUST provided data for phylogenetic reconstructions that differed only slightly from the one in Online Resource 3b, actually sorting out some of the merged clades in the latter (Online Resource 4).

The ncLSU dataset allowed the identification of the branches that likely correspond to *Amanita volvata*, *A. peckiana*, *A. curtipes*, *A. avellaneosquamosa* and *A. clarisquamosa* (Table 1) and suggested further species not represented in the ITS1:ITS2 dataset. Thus, after studying the possibility of assigning such branches to species already delimited in the latter dataset (see Discussion), a complete list of 81 tentative species was obtained (Table 4). The potential splitting of the two sp. 'Naletali' sequences and the potential merging of aff. volvata8 with the sp. S19/aff. volvata9 clade may cause a slight alteration of the total number (see Taxonomy section). Overall, 49 clades at species level (60% of the total of 81) are represented by a unique specimen collection or environmental sequence, and designated "singletons".

{Table 4}

Biogeography of *Amidella*

Though limited by the geographically uneven representation, the present species outline (Table 4 and Online Resources 5–6) brings out noteworthy aspects. The species are distributed over temperate, subtropical and tropical regions only, comprising 12 of the biogeographic regions defined by Loidi and Vynokurov (2024); the highest diversity is in Asia (40–41 species, half of the total), followed by North America (21 species), Africa (13–14 species), Europe (5 species) and Oceania (4 species). No sequences are known from South America, and 2 species are known (as basidiomes) to be shared between Southern Europe and North Africa (Neville and Poumarat 2004). All Baakani species are from Sub-Saharan Africa and Madagascar.

Furthermore, most species have a limited geographic range. Examples with a relatively wider distribution are *Amanita fulvisquamea*, *A. volvata*, *A. ponderosa* and *A. whetstoneae*, but at least in two of these the ranges might outsize that of the basidiomes: the presence of *A. ponderosa* in oak ectomycorrhizas from South England, for example, is far to the North from the known northernmost occurrence of this species as basidiomes (Neville and Poumarat 2004), and *A. fulvisquamea* basidiomes are known only from Northern Thailand (Liu et al. 2023). Currently, only *A. ponderosa* and *A. pseudovalens* overlap biogeographical regions.

Finally, Tulloss and Yang (2024) list 14 species (some with provisional names) without available sequences (Online Resource 6): seven from Africa (*Amanita floccosolivida* Beeli, *A. fulvosquamulosa* Beeli, *A. goossensiae* Beeli, *A. irreperta* nom. prov. E.-J. Gilbert, *A. olivacea* Beeli, *A. subviscosa* Beeli, *A. subviscosa* sensu Pegler & Shah-Sm.), one from Asia (*A. duplex* Corner & Bas), four from North America (*A. dolichopus* nom. prov. Tulloss, *A. fallax* nom. prov. Tulloss & G. Wright, *A. occidentalis* O. K. Mill. & D. J. Lodge, *A. pseudovolvata* nom. prov. Tulloss) and three from Oceania (*A. curta* (Cooke & Massee) E.-J. Gilbert, *A. grisea* Massee & Rodway, *A. pallidogrisea* A. E. Wood). Given that *A. floccosolivida* and *A. pallidogrisea* may not belong to section *Amidella* (Tulloss and Yang 2024), it would be less likely to have them captured in our search; apart from that, a relation to specimens or samples for which any of the classified sequences are known seems to be possible only for *A. subviscosa* sensu Pegler & Shah-Sm. (noting in particular the Zambian *subviscosa* s.l. species clade), *A. irreperta* (to the sp. 'Madagascar' clade) and *A. fallax* (most likely the sp. 'California' clade).

Taxonomy

We first propose an update of the description of section *Amidella* and later outline the species

262 composition of the clades within *Amidella* (Table 4).

Updated description for Section *Amidella*

In their taxonomic scheme for European species of *Amanita*, Neville and Poumarat (2004) adopted
subsection *Ovoideinae* Singer, corresponding to the original genus *Amidella* E.-J. Gilbert, and within
it they made a split between series *Ovoidea* and series *Amidella*, proposing for the first time to our
knowledge a separation between the current concept of *Amidella* and species that have been replaced
in section *Roanokenses* (Wolfe et al. 2012; Cui et al. 2018; Riccioni et al. 2019). The keys to their
Amidella series result in the following diagnosis (our translation from the original in French):
Margin of the pileus smooth and sometimes appendiculate, saccate volva under the stipe, annulus
rudimentary or absent, quick discolouration of the context to reddish tones either by surface rubbing
or exposure to air, later to brownish tones, amyloid spores (Neville and Poumarat 2004).
According to Neville and Poumarat (2004), the separation of the two series is for "reasons of
convenience" based on the discolourations. However, for non-European species the above diagnosis
and the discolouration criterium require further clarification: besides the fact that in many specimens
the pileus margin is striated and not smooth, the requirement for discolouration is left to some
questioning, with an "unchanging trama" commonly reported, while several species in the current
concept of *Roanokenses* have discolourations considered typical of *Amidella*, for example *Amanita*
daucipes (Mont.) Lloyd, *A. fulvopulverulenta* Beeli, *A. luteofolia* Yang-Yang Cui, Qing Cai & Zhu L.
Yang or *A. mutabilis* Beardslee (Tulloss 1984; Cui et al. 2018; Tulloss and Yang 2024). It is only in
combination with other characters, namely the saccate volva and lack of a persistent membranous
ring, that the discolourations can help define *Amidella*.
Currently, at section level, *Amidella* is the taxon introduced by Konrad and Maublanc (1948), who
sweepingly transferred it from the one defined by Gilbert at subgenus level. Although the taxon is not
listed in the Indexfungorum (Royal Botanic Gardens Kew et al. 2024) and Mycobank (Robert et al.
2005) nomenclatural databases, it is considered valid by different authorities (Neville and Poumarat
2004; Cui et al. 2018; Tulloss and Yang 2024). However, contrary to Gilbert's nuanced original
description of the protolog for genus *Amidella* (Neville and Poumarat 2004), the concept of section
Amidella needs revision because it excludes striation of the pileus margin, which is sometimes
mentioned for the section (Yang et al. 2001; Sanmee et al. 2008) and specifies a bulbous stipe base
(Konrad and Maublanc 1948). Furthermore, we find that several European species show a faint grey
reaction with iron sulphate on the exposed context, as suggested first by the analysis of *Amanita*
ponderosa (Tulloss and Yang 2024), and in contradiction to a rule put forward for the whole genus
(Neville and Poumarat 2004). We believe that this reaction should be systematically tested for the
whole of *Amanita* in order to determine its phylogenetic extent. Therefore, we propose the following
update:
Section *Amidella* Konrad & Maublanc mut. char. (emend. P. Oliveira & R. Arraiano-Castilho)
Type species: *Amanita volvata* (Peck) Lloyd
Agaricoid basidiomata, all parts initially whitish and becoming ochre to tan with aging or drying.
Usually, the surface of fresh specimens quickly reddens where rubbed or bruised, in some cases with

a brownish or greyish discolouration. Universal veil persisting as a saccate membranous volva around
the stipe base and, sometimes, as membranous patches on the pileus, with remnants of the inner friable
layer on the lower part of the stipe and/or on the pileus standing out as earlier and more intensely
coloured markings, forming zebroid patterns on the stipe or squamules or patches on the pileus
(lepiotoid appearance). Remains of the partial veil adherent to the upper part of the stipe, sometimes
contrasting with the more intensely discoloured friable remains of the universal veil on the lower part,
forming in some immature specimens an evanescent creamy annulus and/or tiny appendiculate rags
on the pileus margin. The pileus margin can be striated (*Vaginatae* appearance). Lamellulae truncate
to nearly truncate, lamellar edge normally concolorous. Stipe base usually not bulbous in cross section
(discounting the added thickening by the volva). Context (fresh) frequently showing a quick
reddening on exposure, and/or (fresh as well as dried) a faint grey reaction with iron sulphate solution.
Spores amyloid, usually elongate to cylindric (typically average Q > 1.6), septa without clamps in all
parts.
315
Clades at subsection level
317
Volvatae
The proposed name assumes that *Amanita volvata*, the type species for section *Amidella*, is
represented by accession AF024485 (Table 1). At least one close ncLSU sequence (GenBank
accession OQ311333) is from a specimen also represented in the ITS concatenate dataset, thus
enabling the merging of various ITS sequences within this species and forming the reference species
clade for section *Amidella*.
Neville and Poumarat (2004) have used *Volvatae* at section level, following previous authors to
encompass (for Europe) current sections *Amidella*, *Phalloideae*, *Validae* and part of *Roanokenses*,
thus it is an altogether different taxon.
Three resolved and generally well-supported subclades at stirp level are considered:
1. Volvata
Comprises the type species *Amanita volvata* (Peck) Lloyd, *A. clarisquamosa* (S. Imai) S. Imai, *A.*
curtipes E.-J. Gilbert (all based on the assignments in Table 1), *A. pseudovalens* (Neville & Poumarat)
Arraiano-Castilho et al., and 3 unnamed or provisional species (Table 4). According to our analyses,
a variety of provisional names from the Eastern United States (sp. T52 Tulloss & S. D. Russell, IN57,
IN36) are all *A. volvata*.
2. Peckiana
Comprises *Amanita peckiana* Kauffman in Peck (comprising sequences from specimens clustering
around accession HQ539720, Table 1), *A. brunneomaculata* Yang-Yang Cui, Qing Cai & Zhu L. Yang,
and *A. lanigera* Yang-Yang Cui, Qing Cai & Zhu L. Yang (Table 4). According to our analyses, *A.*
floridella Tulloss & S. D. Russell nom. prov. is *A. peckiana*.
3. Fulvisquamea
Comprises *Amanita fulvisquamea* Yuan S. Liu & S. Lumyong and *A.* sp. 'Sarawak1', the latter
represented by ncLSU sequences clustering with those of *A. fulvisquamea* with moderate support
(Table 3)..

Sagittariae

This clade is composed of 8 singletons, and named after *Amanita sagittaria* Tulloss & K. W. Hughes nom. prov.. Two sequences are named *A. clarisquamosa*, but the specimen assumed to represent this species (Table 1) has an ncLSU sequence in the Volvatae clade, unrelated to Sagittariae specimens (Figure 1b); therefore, under the assumption of Table 1 the *clarisquamosa* epithet may not be applicable in clade Sagittariae.

Lepiotidae

The proposed name is based on the oldest accepted epithet present, *Amanita lepiotoides*. This group of species is by far the most diverse (about half of Section *Amidella*). Notwithstanding the strong support in the ITS concatenate phylogeny for this clade and its main subclades (Figure 1a), the ncLSU phylogeny shows the corresponding group in a clade with low support that includes Volvatae and Sagittariae (Figure 1b, Table 3), and with an even lower support in the *RPB2* phylogeny but separate from Volvatae (Table 3, Online Resource 2a). Therefore, the supposed natural groups in the ITS phylogeny should be viewed with caution.

Moreover, there is a potential difficulty with the names to be used. The available sequences named *Amanita lepiotoides* are of the forma *subcylindrospora* (Neville and Poumarat 2004; Arraiano-Castilho et al. 2022) and one cannot be certain whether the autonym form would be phylogenetically close or not. Furthermore, as pointed out in the footnote to Table 1, the original *A. avellaneosquamosa* could be different from our assumption, noted here as 'cf. avellaneosquamosa1' (Table 4, Online Resources 1 and 3). Hence, if any of these two taxa turn out to belong outside the clades under consideration, the respective naming should be revised.

Based on the ITS concatenate phylogeny, four subclades are considered at stirp level:

1. Lepiotoides

Comprises *Amanita lepiotoides* f. *subcylindrospora* Neville & Poumarat, *A. rufobrunnescens* W. Q. Deng, T. H. Li & Hosen, *A. parvirufobrunnescens* A. Kumar, Y. P. Sharma & Mehmood, *A. insolens* I. Safonov & L. V. Kudzma nom. prov., and up to 4 other species (Table 4). One of the latter might be *A. texidella* Tulloss, Pastorino & K. W. Hughes nom. prov., since its ncLSU sequence is close to *A.* sp. JPN01/sp. OMS-015 (80% posterior probability), but the branch containing this species pair is separate from the remainder of the Lepiotoides sequences.

2. Avellaneosquamosa

Comprises 12 clades, including *Amanita parvicurta* Yang-Yang Cui, Qing Cai & Zhu L. Yang and the specimen presumed to represent the original *A. avellaneosquamosa* (S. Imai) S. Imai in Ito, included here in clade *A.* cf. *avellaneosquamosa1* (see note on Table 1), as well as 10 unnamed species. The epithet *rosea*, given with a cf. to one ncLSU sequence (accession OQ628446), is of a species belonging in Section *Roanokenses*, and for this reason *A.* aff. *rosea* is used to name the corresponding clade.

3. Subviscosa

This branch bears the name of *Amanita subviscosa* Beeli but this species might not be represented in the current dataset: a group of three closely related species clades, each corresponding to

geographically segregate SH codes, was named *A. subviscosa* s.l. instead, for it is uncertain whether any of these correspond to the described *A. subviscosa* (Table 4, see also Discussion). The Subviscosa group comprises up to 14 species from warm climates, including almost all from Africa that are not Baakani (Table 4). Although the ITS phylogeny places *A.* sp. Thai04 in this group, this species forms a separate branch in the ncLSU phylogeny, together with the additional *A.* sp. 'Sarawak2', thus the inclusion of both in Subviscosa is rather tentative. The ncLSU sequence named *A.* aff. goossensiae KM 12 has a position in this subclade (close to *A. subviscosa* s.l.) that appears not to be the one expected for the clade Baakani (see below), where other specimens from the same Cameroon location are also named in approximation of the same epithet.

4. Pseudomutabilis

This clade is named in connection with one of the ncLSU sequences (accession MH620228) identified as *Amanita mutabilis*, which is expected to be wrong since this is a species with a membranous ring belonging in Section *Roanokenses* (Tulloss & Yang 2024). The species clade for that accession is named *A.* aff. *mutabilis*. Further species clades include *A. pseudorufobrunnescens* K.C. Semwal, K. Das, R.P. Bhatt, Mehmood & V.K. Bhatt, and three others, one of them possibly *A. fallax* nom. prov. as noted in the Biogeography section above.

5. Unclassified

There are three Lepiotidae species that form single branches and thus are kept unclassified: *Amanita* sp. 19 GMB-2014 and *A.* sp. canadensis01/sp. IN28 are placed next to Avellaneosquamosa (Figure 1a-b); and *A.* aff. volvata1, represented by a single ncLSU sequence.

Ponderosae

The proposed name is based on the oldest accepted species in this clade, *Amanita ponderosa*. Three well-resolved and well-supported subclades at stirp level are considered:

1. Ponderosa

Comprises *Amanita ponderosa* Malençon & R. Heim and *A. whetstoneae* Tulloss, Goldman and Kudzma nom. prov..

2. Claristriata

Comprises *Amanita claristriata* Yuan S. Liu & S. Lumyong, *A. neocaesariensis* Tulloss, I. Safonov & K. W. Hughes nom. prov. (including specimens labelled IN26), and around 6 other unnamed species. The uncertainty in the number of species is due to the limited species-level correspondence between some ITS clades (sp. OMS-010/aff. volvata6, aff. volvata7, aff. volvata8, sp. S19/aff. volvata9) and their presumed ncLSU counterparts (see Discussion).

3. Pinophila

Comprises *Amanita pinophila* Yang-Yang Cui, Qing Cai & Zhu L. Yang and an unidentified specimen from China labelled *A.* aff. *albidostipes*, an epithet that applies to a species in Section *Vaginatae* (Cui et al. 2018).

Baakani

This clade is named with reference to the Ba'Aka nation in Cameroon, who live in the provenance of six undescribed specimens named for the apparent affinity to *Amanita gossensiae* Beeli, a species

classified in Section *Amidella* (Tulloss and Yang 2024). These specimens belong in 2 species, and another 2–3 species from Zimbabwe and 1 from Madagascar also belong in this clade. Furthermore, an unnamed species from Zambia (*A*. sp. Arora-01-560) represented by a single ncLSU sequence has a position in the phylogeny congruent with belonging in Baakani. We note that the position of *A. goossensiae* still needs to be confirmed, and a separate collection named *A.* aff. *goossensiae* KM 12, from the same Cameroon provenance, might be an alternative candidate, but its ncLSU sequence is placed within Lepiotidae (see Subviscosa, above; Figure 1b).

Hesleri

Sequences belonging to either *Amanita hesleri* Bas or *A. zangii* Zhu L. Yang, T.H. Li & X.L. Wu, which belong to Section *Phalloideae* (Cai et al. 2014; Cui et al. 2018), frequently appeared in the BLAST results, and early phylogenetic reconstructions of *Amidella* suggested them to form a clade intermediate between the outgroup and *Amidella*, even forming a clade with the Baakani sequences in one case (not shown), thus raising questions about the actual limits between sections. Their morphologies are atypical for *Phalloideae* (Cui et al. 2018) and certainly do not conform to the established concept of Section *Amidella*, namely for the lack of saccate volva and lack of discolourations of the basidiome, as well as for the microscopy of the volva remnants on the pileus (Yang et al. 2001; Tulloss and Yang 2024), so their placement had to be examined, both for the ITS1:ITS2 concatenate and the ncLSU. The results confirmed their position within *Phalloideae*, neatly separate from *Amidella* (Online Resource 7). The short branch separating Hesleri from the *Phalloideae* node in most phylogenies may explain why these taxa were retrieved in our BLAST-based searches.

Discussion

The present work collects for the first time a comprehensive coverage of the sequence data available in Section *Amidella*, establishing its limits with regard to neighbouring sections, delimiting clades at species level and thereby tripling the total count, uncovering their arrangement into subsectional clades and opening a new discussion on their biogeography.

Evidence for subsectional clades and undescribed diversity

The present work presents for the first time, as far as the authors are aware, a nearly exhaustive sampling of the DNA sequence diversity available for section *Amidella*. Our analyses provide a boundary separating the section from neighbouring sections *Roanokenses*, *Arenariae* and *Phalloideae*, to include unsuspected phylogenetic branches even in the face of misleading epithets such as vaginata. Furthermore, a previously unknown structure is revealed, with five clades at subsection level, with further subdivision at stirp level in three of them. This structure adds at least one level of precision in the currently problematic task of species determination, and can be very useful in collecting sequences for phylogenetic analyses more effectively, both by overriding the need for very large datasets in studies of closely related groups (Dissanayake et al. 2024) and by helping form a balanced representation of the groups to be studied. Although well-supported in general (the main exception is a partial disruption of Lepiotidae in the ncLSU phylogeny, Figure 1b, Table 3), the topology between the five clades is rather constant, with Baakani as the early-branching clade, while Sagittariae and Volvatae appear as the most derived. It remains to be seen whether it will hold up with the publication of new taxa, for example the five vouchers from China (Cai et al. 2024) placed within *Amidella* thanks to the *TEF1* sequences released so far (HKAS57709/69526/84822/92026/129313, Online Resource 8).

The complexity within *Amidella* is also expressed by a total of about 81 clades at species level (Table 4, Online Resource 3), of which only 16–17 have accepted names (Table 4). Adding 9–12 described species without sequences (Online Resource 6), the current total for section *Amidella* reaches circa 90 species, of which only one third have accepted names. The delimitation of clades at species level was done with a branch clustering option in MEGA, using consensus cutoffs for the pairwise uncorrected p-distances that, in this group, were set at 1.25% for the ITS concatenate and 0.8% for the ncLSU. This approach is tree-based and is not limited by the high prevalence of singletons, which is an obstacle for statistical approaches that rely on intraspecies diversity such as sGMYC or mPTP (Fujisawa and Barraclough 2013; Leavitt et al. 2015; Sánchez-Ramírez et al. 2017; Maharachchikumbura et al. 2021). However, the cutoff used for a given phylogeny is likely heterogeneous in different branches, as suggested by the slight differences between different approaches to the ncLSU dataset (Online Resources 3b and 4). Notwithstanding the suggestion that only the ITS2 region has an effective barcoding gap for *Amanita* (Badotti et al. 2017), the high prevalence of species not represented by ITS sequences argues for the use of all information available. At any rate, the evidence of a high number of novel species, mainly in clades Baakani, Sagittariae, Lepiotidae (subsection level) and Claristriata (stirp level), is compelling.

It is worth mentioning that, because of nrDNA cistron heterogeneity, a single sequence might not be representative of the collection itself — as so well exemplified in Online Resource 3a with the

Amanita hiltonii ITS sequences in the outgroup (Davison and Giustiniano 2020). Genomic analyses have demonstrated that the same can happen in the ncLSU (Paloi et al. 2022; Cedeño-Sanchez et al. 2024). Nevertheless, a recent survey has obtained a >99% pairwise identity between ITS cistrons in *Amanita* (Bradshaw et al. 2023), therefore we feel that nrDNA cistron heterogeneity would affect, if anything, the separations at interspecies level rather than higher-level relationships.

Unfortunately, DNA sequences are the only support for these subclades. Even with the very detailed descriptions available (Deng et al. 2016; Cui et al. 2018; Semwal et al. 2020; Liu et al. 2023; Tulloss and Yang 2024), attempts to derive morphology-based infrasectional simplesiomorphies were unsuccessful (not shown). This, and the possibility of minor changes as more species are discovered, deters their formal proposal as infrasectional taxa. On the other hand, the available alignments suggest efficient means to design taxon-specific PCR primers to retrieve enough sequence information from type specimens and thus resolve the hitherto unclear identification of taxa such as *Amanita avellaneosquamosa* (Table 1; Online Resource 9).

Biogeographical implications

Strikingly, practically all species have a relatively narrow geographical range (Table 4, Online Resource 5), as reported already for other taxa (Looney et al. 2016, 2020; Peay and Matheny 2016; Bazzicalupo et al. 2019; Noffsinger et al. 2024). This could help avoid using the same name for geographically distinct collection origins — as the clade naming used here at species level demonstrates, the *volvata*, *peckiana*, *curtipes* and *avellaneosquamosa* epithets have been used very confusingly (Figure 1, Table 4, Online Resource 1). Four contrasting examples serve to highlight the application of this geographic premise, as well as difficulties that may arise:

i. The clade *Amanita subviscosa* s.l. spans an unusually vast distribution in Africa, from Benin to Zambia and Zimbabwe (Table 4). We named it so for two reasons: first, the SH codes at 1.5% are unique for each country, thus suggesting that three species of narrower range are to be considered; second, *A. subviscosa* has been described from a fourth country (Online Resource 6) and is regarded as different from an unsequenced collection from Zambia (Tulloss and Yang 2024), therefore one cannot be sure which of the sequenced specimens, if any, represents *A. subviscosa*.

ii. There are three North American ITS clades in Claristriata (Ponderosae) with contiguous geographical distributions: *Amanita* sp. S19/aff. volvata9 ranging from Massachusetts to Tennessee, *A.* sp. neocaesariensis/sp. IN26 from New York to Wisconsin, and *A.* sp. T51 in Mississippi and Eastern Texas. The latter two collapse into a single lineage at 1.4% divergence, raising the possibility that they could be a single species, but their disjunct distributions suggest vicariant species. The three clades merge at 1.55% divergence, strengthening the hypothesis of a recent speciation. They might be useful for a case study of a speciation process in ectomycorrhizal fungi.

iii. Still with the *Amanita* sp. S19/aff. volvata9 group, there is an ncLSU sequence from a South Korean collection that is very close phylogenetically. An ITS sequence for this collection would help understand its relationship with the North American group, but none is available. Tentatively, we consider the South Korean collection autochthonous and assign a separate epithet (volvata8),

but it could turn out to be the same species as the North American collections, and one possible explanation for such a wide distribution could be the accidental introduction with an ectomycorrhizal host originating in the United States (Kim and Zsuffa 1994; Vellinga et al. 2009).

iv. The provenances of Eastern Asia (specifically, Mainland China, Taiwan, Japan and South Korea) have only a minority of shared species (Online Resource 10), most of them between Japan and South Korea, while Mainland China shares only 2 out of 16 with Japan (one being *Amanita* aff. avellaneosquamosa3/OMS-012, but the Chinese occurrence is in a botanical garden in Jianxi; the other is *A. fulvisquamea*, a rather wide-ranging species when soil and root tip samples are considered, however it occurs as fruitbodies only in Thailand until now). One should expect these separations to blur as more findings are reported, but currently, for any species in this group, it is worth considering with caution a shared geographical distribution between Mainland China and Japan, as we note on Table 1.

Uniqueness of species defined with different datasets

A considerable part of the species deduced from clades in the present work (twelve, to be precise) are represented by ncLSU sequences without an ITS counterpart. This raises two questions: i) is any of these species the same as one of those defined by ITS sequences without ncLSU counterpart, such that both should be merged, and ii) is the species threshold used for ncLSU (0.8% p-distance, Online Resource 3b) anywhere less applicable?

To answer the first question, the cases to be considered for merging narrow down to a few possible pairs classified in the same subclade and possibly not too far geographically: in Claristriata, there are *Amanita* aff. avellaneosquamosa6 (ITS; China, Yunnan) and *A.* aff. volvata8 (LSU; South Korea), but the latter is phylogenetically very close to *A.* sp. S19/aff. volvata9 (Eastern United States) and the former is not; in Avellaneosquamosa, *Amanita* aff. rosea (LSU) is Australian like *Amanita* sp. 19 GMB-2014 (ITS) and both are phylogenetically close to *Amanita parvicurta*, but the first is from New South Wales and the second from Northern Territory; in Lepiotidae, *Amanita* aff. volvata1 (LSU; China, Yunnan) remains unclassified and the species that are nearest geographically (in Thailand and South China) are in unrelated Subviscosa branches; in Baakani, the *Amanita* sp. Arora-01-560 specimen (LSU; Zambia) could be of the same species as one of those in the clade represented by ITS sequences, especially those in Zimbabwe, at least 1000 Km away nevertheless. Hence, each of these four pairs seems unlikely to be merged. Finally, the *Amanita* sp. S17/sp. IN41 clade has a single publicly available ncLSU accession (HQ539665) for the reference specimen RET 444-9 (Tulloss and Yang 2024), but there are several others in the ITS concatenate dataset identified as *Amanita* sp. S17, in a cluster that includes specimens designated *A.* sp. IN41, *A.* "volvata" or *A.* "peckiana". This extension of *Amanita* sp. S17 seems reasonable since all specimens are from Eastern United States, with similar ecology and phenology and, like accession HQ539665, they cluster with the same group of clades at species level (*Amanita* aff. volvata5, *A.* aff. avellaneosquamosa4/aff. volvata3, *A.* cf. avellaneosquamosa1).

The second question highlights a contrast between the criterium used here for the ITS (1.25%, Online Resource 3a) and the one used for the ncLSU phylogeny (0.8%, Online Resource 3b). While the

former was chosen by trial and error, guided by branch collapsing within accepted species, for the
latter there was no consensus value that could mimic the results with the species also present in the
ITS concatenate dataset. Thus, the ABGD method (automatic barcoding gap discovery) was used to
obtain the 0.8% estimate for the p-distance to be used for ncLSU, to settle on a common threshold
(Online Resource 3b) and choose the similarity level of 99.2% for UCLUST, used as a phylogeny-
independent approach to delimiting sequence clusters. From both approaches it was possible to obtain
alternative ncLSU groupings (Online Resource 4) not unlike the one in Online Resource 3b,
separating adequately some of the merged clades in the latter, for example the one containing *Amanita*
lanigera and *A. brunneomaculata*. We take these results as independent support for the proposed
species list (Table 4), but it remains admissible that part of those twelve ncLSU-only clades might be
later considered for merging with others, especially as the respective ITS sequences become known.

Conclusion and Outlook

The present study exposes the unsuspected dimension of some knowledge gaps regarding section
Amidella: a large majority of undescribed species, a sizeable number of described species without
molecular information, the need for molecular studies on types to clarify the identity of species such
as *Amanita avellaneosquamosa* and *A. clarisquamosa*, the narrow geographical range as a rule, and
an essential need for novel discriminant characters, the utility of which is framed from now on by the
infrasectional taxonomic structure here revealed.

One final point. In naming the clades, we had no choice but to use three epithets (*fulvisquamea*,
claristriata, *pinophila*) that have been accepted only recently (Cui et al. 2018; Liu et al. 2023),
sagittaria that is still provisional (Tulloss and Yang 2024), and devise a novel name for the Baakani
clade. Section *Amidella* is still a largely untrodden field of research.

References

Abarenkov K, Nilsson RH, Larsson K-H, et al (2024) The UNITE database for molecular identification and taxonomic communication of fungi and other eukaryotes: sequences, taxa and classifications reconsidered. Nucleic Acids Res 52:D791–D797. https://doi.org/10.1093/nar/gkad1039

Abarenkov K, Tedersoo L, Nilsson RH, et al (2010) PlutoF—a Web Based Workbench for Ecological and Taxonomic Research, with an Online Implementation for Fungal ITS Sequences. Evol Bioinforma 6:189–196. https://doi.org/10.4137/EBO.S6271

Arraiano-Castilho R, Silva AC, Vila-Viçosa C, et al (2022) The Amidella Clade in Europe (Basidiomycota: Amanitaceae): Clarification of the Contentious Amanita valens (E.-J.Gilbert) Bertault and the Importance of Taxon-Specific PCR Primers for Identification. Cryptogam Mycol 43:139–157. https://doi.org/10.5252/cryptogamie-mycologie2022v43a6

Ayres DL, Cummings MP, Baele G, et al (2019) BEAGLE 3: Improved Performance, Scaling, and Usability for a High-Performance Computing Library for Statistical Phylogenetics. Syst Biol 68:1052–1061. https://doi.org/10.1093/sysbio/syz020

Badotti F, de Oliveira FS, Garcia CF, et al (2017) Effectiveness of ITS and sub-regions as DNA barcode markers for the identification of *Basidiomycota* (*Fungi*). BMC Microbiol 17:42. https://doi.org/10.1186/s12866-017-0958-x

Bazzicalupo AL, Whitton J, Berbee ML (2019) Over the hills, but how far away? Estimates of mushroom geographic range extents. J Biogeogr 46:1547–1557. https://doi.org/10.1111/jbi.13617

Bengtsson-Palme J, Ryberg M, Hartmann M, et al (2013) Improved software detection and extraction of ITS1 and ITS 2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. Methods Ecol Evol 4:914–919. https://doi.org/10.1111/2041-210X.12073

Bradshaw MJ, Aime MC, Rokas A, et al (2023) Extensive intragenomic variation in the internal transcribed spacer region of fungi. iScience 26:107317. https://doi.org/10.1016/j.isci.2023.107317

Bruen TC, Philippe H, Bryant D (2006) A Simple and Robust Statistical Test for Detecting the Presence of Recombination. Genetics 172:2665–2681. https://doi.org/10.1534/genetics.105.048975

Cai Q, Codjia JEI, Buyck B, et al (2024) The evolution of ectomycorrhizal symbiosis and host-plant switches are the main drivers for diversification of Amanitaceae (Agaricales, Basidiomycota). BMC Biol 22:230. https://doi.org/10.1186/s12915-024-02031-8

Cai Q, Tulloss RE, Tang LP, et al (2014) Multi-locus phylogeny of lethal amanitas: Implications for species diversity and historical biogeography. BMC Evol Biol 14:143. https://doi.org/10.1186/1471-2148-14-143

Cedeño-Sanchez M, Cheng T, Lambert C, et al (2024) Unraveling intragenomic polymorphisms in the high-quality genome of Hypoxylaceae: a comprehensive study of the rDNA cistron. Mycol Prog 23:5. https://doi.org/10.1007/s11557-023-01940-2

Cho HJ, Park MS, Lee H, et al (2015) Four New Species of *Amanita* in Inje County, Korea. Mycobiology 43:408–414. https://doi.org/10.5941/MYCO.2015.43.4.408

Cui Y-Y, Cai Q, Tang L-P, et al (2018) The family *Amanitaceae*: molecular phylogeny, higher-rank taxonomy and the species in China. Fungal Divers 91:5–230. https://doi.org/10.1007/s13225-018-0405-9

Davison EM, Giustiniano D (2020) Amanita hiltonii (Amanitaceae), a common but frequently misidentified mushroom in southwestern Australia, and reconsideration of A. albifimbriata and A. brunneibulbosa. Muelleria Aust J Bot 39:59–73. https://doi.org/10.5962/p.340570

Deng W, Li T, Hosen MdI (2016) Amanita rufobrunnescens, a new species of Amanita section Amidella from South China. Phytotaxa 243:147–154. https://doi.org/10.11646/phytotaxa.243.2.4

Dissanayake AJ, Zhu J-T, Chen Y-Y, et al (2024) A re-evaluation of Diaporthe: refining the boundaries of species and species complexes. Fungal Divers. https://doi.org/10.1007/s13225-024-00538-7

Edgar RC (2022) Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. Nat Commun 13:6968. https://doi.org/10.1038/s41467-022-34630-w

Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. Bioinformatics 26:2460–2461. https://doi.org/10.1093/bioinformatics/btq461

Fujisawa T, Barraclough TG (2013) Delimiting Species Using Single-Locus Data and the Generalized Mixed Yule Coalescent Approach: A Revised Method and Evaluation on Simulated Data Sets. Syst Biol 62:707–724. https://doi.org/10.1093/sysbio/syt033

Huson DH, Bryant D (2006) Application of Phylogenetic Networks in Evolutionary Studies. Mol Biol Evol 23:254–267. https://doi.org/10.1093/molbev/msj030

Kans J (2013) Entrez Direct: E-utilities on the Unix Command Line.

Kim CS, Jo JW, Kwag Y-N, et al (2013) Taxonomic Study of *Amanita* Subgenus *Lepidella* and Three Unrecorded *Amanita* Species in Korea. Mycobiology 41:183–190. https://doi.org/10.5941/MYCO.2013.41.4.183

Kim KH, Zsuffa L (1994) Reforestation of South Korea: The history and analysis of a unique case in forest tree improvement and forestry. For Chron 70:58–64. https://doi.org/10.5558/tfc70058-1

Kõljalg U, Nilsson RH, Abarenkov K, et al (2013) Towards a unified paradigm for sequence-based identification of fungi. Mol Ecol 22:5271–5277. https://doi.org/10.1111/mec.12481

Konrad P, Maublanc A (1948) Tribu des Amaniteae Genre 1. Amanita (Fries ex Person) Quélet 1872. In: Les Agaricales. Classification - Revision des espèces - Iconographie - Comestibilité. Tome 1. Agaricaceae. Paul Lechevalier, pp 56–68

Kumar S, Stecher G, Suleski M, et al (2024) MEGA12: Molecular Evolutionary Genetic Analysis Version 12 for Adaptive and Green Computing. Mol Biol Evol 41:msae263. https://doi.org/10.1093/molbev/msae263

Leavitt SD, Moreau CS, Thorsten Lumbsch H (2015) The Dynamic Discipline of Species Delimitation: Progress Toward Effectively Recognizing Species Boundaries in Natural Populations. In: Upreti DK, Divakar PK, Shukla V, Bajpai R (eds) Recent Advances in

672   Lichenology. Springer India, New Delhi, pp 11–44

673   Letunic I, Bork P (2024) Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree
674   display and annotation tool. Nucleic Acids Res gkae268.
675   https://doi.org/10.1093/nar/gkae268

676   Liu YS, Liu J-K, Kumla J, et al (2023) Taxonomic Novelties and New Records of Amanita
677   Subgenus Amanitina from Thailand. J Fungi 9:601. https://doi.org/10.3390/jof9060601

678   Loidi J, Vynokurov D (2024) The biogeographical kingdoms and regions of the world. Mediterr Bot
679   45:e92333. https://doi.org/10.5209/mbot.92333

680   Looney BP, Adamčík S, Matheny PB (2020) Coalescent-based delimitation and species-tree
681   estimations reveal Appalachian origin and Neogene diversification in Russula subsection
682   Roseinae. Mol Phylogenet Evol 147:106787. https://doi.org/10.1016/j.ympev.2020.106787

683   Looney BP, Ryberg M, Hampe F, et al (2016) Into and out of the tropics: global diversification
684   patterns in a hyperdiverse clade of ectomycorrhizal fungi. Mol Ecol 25:630–647.
685   https://doi.org/10.1111/mec.13506

686   Maharachchikumbura SSN, Chen Y, Ariyawansa HA, et al (2021) Integrative approaches for
687   species delimitation in Ascomycota. Fungal Divers 109:155–179.
688   https://doi.org/10.1007/s13225-021-00486-6

689   Moreno G, Platas G, Peláez F, et al (2008) Molecular phylogenetic analysis shows that *Amanita*
690   *ponderosa* and *A. curtipes* are distinct species. Mycol Prog 7:41–47.
691   https://doi.org/10.1007/s11557-007-0551-z

692   Neville P, Poumarat S (2004) *Amidella*. In: *Amaniteae*: *Amanita*, *Limacella* et *Torrendia*. Candusso,
693   Alassio, pp 645–699

694   Nilsson RH, Veldre V, Hartmann M, et al (2010) An open source software package for automated
695   extraction of ITS1 and ITS2 from fungal ITS sequences for use in high-throughput
696   community assays and molecular ecology. Fungal Ecol 3:284–287.
697   https://doi.org/10.1016/j.funeco.2010.05.002

698   Noffsinger CR, Adamčíková K, Eberhardt U, et al (2024) Three new species in *Russula* subsection
699   *Xerampelinae* supported by genealogical and phenotypic coherence. Mycologia 116:322–
700   349. https://doi.org/10.1080/00275514.2023.2295957

701   Okonechnikov K, Golosova O, Fursov M (2012) Unipro UGENE: a unified bioinformatics toolkit.
702   Bioinformatics 28:1166–1167. https://doi.org/10.1093/bioinformatics/bts091

703   Paloi S, Luangsa-ard JJ, Mhuantong W, et al (2022) Intragenomic variation in nuclear ribosomal
704   markers and its implication in species delimitation, identification and barcoding in fungi.
705   Fungal Biol Rev 42:1–33. https://doi.org/10.1016/j.fbr.2022.04.002

706   Peay KG, Matheny PB (2016) The biogeography of ectomycorrhizal fungi – a history of life in the
707   subterranean. In: Martin F (ed) Molecular Mycorrhizal Symbiosis, 1st edn. Wiley, pp 341–
708   361

709   Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery
710   for primary species delimitation. Mol Ecol 21:1864–1877. https://doi.org/10.1111/j.1365-
711   294X.2011.05239.x

Quast C, Pruesse E, Yilmaz P, et al (2012) The SILVA ribosomal RNA gene database project:
improved data processing and web-based tools. Nucleic Acids Res 41:D590–D596.
https://doi.org/10.1093/nar/gks1219

Riccioni C, Paolocci F, Tulloss RE, Perini C (2019) Molecular phylogenetic analyses show that
Amanita ovoidea and *Amanita proxima* are distinct species and suggest their assignment to
Roanokenses section. Mycol Prog 18:1275–1283. https://doi.org/10.1007/s11557-019-
01521-2

Robert V, Stegehuis G, Stalpers J (2005) The MycoBank engine and related databases

Royal Botanic Gardens Kew MS, Landcare Research-NZ MG, Chinese Academy of Science I of M
(2024) Index Fungorum

Sánchez-Ramírez S, Wilson AW, Ryberg M (2017) Overview of Phylogenetic Approaches to
Mycorrhizal Biogeography, Diversity and Evolution. In: Tedersoo L (ed) Biogeography of
Mycorrhizal Symbiosis. Springer International Publishing, Cham, pp 1–37

Sanmee R, Tulloss RE, Lumyong P, et al (2008) Studies on Amanita (Basidiomycetes:
Amanitaceae) in Northern Thailand. Fungal Divers 32:97–123

Semwal KC, Das K, Bhatt RP, et al (2020) Amanita pseudorufobrunnescens, a new species of
Amanita section Amidella from Indian Himalaya. Kew Bull 75:22.
https://doi.org/10.1007/s12225-020-9874-6

Suchard MA, Lemey P, Baele G, et al (2018) Bayesian phylogenetic and phylodynamic data
integration using BEAST 1.10. Virus Evol 4:. https://doi.org/10.1093/ve/vey016

Suchard MA, Rambaut A (2009) Many-core algorithms for statistical phylogenetics. Bioinformatics
25:1370–1376. https://doi.org/10.1093/bioinformatics/btp244

Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. Mol Biol Evol 38:3022–3027. https://doi.org/10.1093/molbev/msab120

Tedersoo L, Hosseyni Moghaddam MS, Mikryukov V, et al (2024) EUKARYOME: the rRNA gene
reference database for identification of all eukaryotes. Database 2024:baae043.
https://doi.org/10.1093/database/baae043

Tulloss RE (1984) Distribution and Taxonomic Notes on *Amanita mutabilis*. Mycologia 76:555–
558. https://doi.org/10.1080/00275514.1984.12023879

Tulloss RE, Yang Z (2024) Section Amidella. http://www.amanitaceae.org?section+Amidella.
Accessed 2024-06-30.

Vellinga EC, Wolfe BE, Pringle A (2009) Global patterns of ectomycorrhizal introductions. New
Phytol 181:960–973. https://doi.org/10.1111/j.1469-8137.2008.02728.x

Wolfe BE, Tulloss RE, Pringle A (2012) The Irreversible Loss of a Decomposition Pathway Marks
the Single Origin of an Ectomycorrhizal Symbiosis. PLoS ONE 7:e39597.
https://doi.org/10.1371/journal.pone.0039597

Yang Z, Li T, Wu X (2001) Revision of Amanita collections made from Hainan, Southern China.
Fungal Divers 6:149–165

Statements and Declarations

No funding was received for conducting this study.

The authors have no relevant financial or non-financial interests to disclose.

Both authors made substantial contributions to the conception of the work, and to the acquisition, analysis, and interpretation of data; both authors drafted the work and revised it critically for important intellectual content; both authors approved the version to be published; and both authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Tables

Table 1. Outline of the nucleotide accessions representing names that are frequently misapplied.

Current name (basionym)	GenBank accessions	Authority
Amanita volvata (*Agaricus volvatus* Peck 1872)	AF024485	Tulloss, R. E. http://www.amanitaceae.org/?Amanita%20volvata
Amanita peckiana (*Amanita peckiana* Kauffman 1913)	HQ539720	Tulloss, R. E. http://www.amanitaceae.org/?Amanita%20peckiana
Amanita curtipes (*Amanita baccata* f. *minor Bres.* 1928)	EF653963 EF653960	Moreno et al. (2008)
Amanita avellaneosquamosa (*Amanitopsis avellaneosquamosa* S. Imai 1933)[a]	MH508259 AF024441 MH486380 MH508682	Tulloss, R. E. http://www.amanitaceae.org/?Amanita%20avellane osquamosa and Cui et al. (2018)
Amanita clarisquamosa (*Amanitopsis clarisquamosa* S. Imai 1933)[a]	AF024448	Yang, Z. L. & Tulloss, R. E. http://www.amanitaceae.org/?Amanita%20clarisqua mosa and Cui et al. (2018)

[a] The choice for *Amanita avellaneosquamosa* is particularly difficult. Of the 6 specimens included by Cui et al. (2018) under this name, 5 separate clades were found in the present study. Furthermore, the sequence from specimen HKAS 38300, highlighted in http://www.amanitaceae.org/?Amanita%20avellaneosquamosa (Tulloss and Yang 2024), is quite distant from all other *A. avellaneosquamosa*, as already noted in Cui et al. (2018). The accession AF024441 also considered by Tulloss & Yang (2024) is more consensual, but its origin (Yunnan, China) is rather different from that of the type (Japan; incidentally, the same happens with the ncLSU sequence chosen here to represent *A. clarisquamosa*). Thus, the clade hereby chosen as reference for *A. avellaneosquamosa* is named 'cf. avellaneosquamosa1', and more generally we caution that the reference sequences listed in this Table 1 are subject to revision.

771 Table 2. Major clades derived from the 5.8S phylogeny with known *Amidella* species epithets. Only
772 the epithets of accepted species are in italic, and those in double quotes are probably misapplied.

Clade	Number of accessions	known species epithets (*Amidella*)	nearest species epithets (examples)
I	53	*lanigera*, *brunneomaculata*, floridella, *curtipes*, *pseudovalens*, sagittaria, "clarisquamosa", *rufobrunnescens*	*goossensfontanae*, *proxima*, *cinereoconia*, *fulvopulverulenta*, *yenii*, *hiltonii*
II	81	*ponderosa*, *pinophila*, whetstoneae, neocaesariensis, "volvata", "avellaneosquamosa", "goossensiae"	*quenda*, *daucipes*, *walpolei*, *ovoidea*
III	30	"avellaneosquamosa", *parvicurta*, "volvata", "peckiana"	*neo-ovoidea*, *preissii*

Table 3. Bayesian posterior probabilities (percentage) for the clades[a]

Marker	ITS1:ITS2	ncLSU	*RPB2*	*TEF1*	*BTUB*	ncLSU:*RPB2*:*TEF1*
Volvatae	100	100	100	80	100	r
Volvata	100	99	s	—	—	—
Peckiana	99	100	100	100	100	100
Fulvisquamea	m	85	m	m	s	—
Sagittariae	91	95	—	—	—	—
Lepiotidae[b]	99	p!	44	r	r	r
Lepiotoides	91	p!	m	—	—	—
Avellaneosquamosa	100	100	96	100	100	100
Subviscosa	98	p!	—	—	—	—
Pseudomutabilis	99	100	—	—	—	—
Ponderosae	99	88	100	100	100	100
Ponderosa	99	100	m	—	—	—
Claristriata	99	100	m	m	m	m
Pinophila	100	m	m	m	m	m
Baakani	100	s	—	—	—	—
[outgroup]	100	100	100	100	99	100

[a] m: one species, monophyletic; p!: paraphyletic; s: single sequence; r: redundant; — no data.
[b] includes three additional single-species branches.

Table 4. List of the clades at species level, arranged according to the adopted taxonomy (subsection and stirp levels) and alphabetically ordered by geographic location name.

Subsection	Strip	Epithet	SH (1.5%)	Geography	Ecology
Volvatae	Volvata	*clarisquamosa*		Asia, China, Yunnan	
		aff. curtipes1	SH0932681.10FU	Asia, South Korea	
		pseudovalens	SH0932676.10FU	Europe, Portugal/France	
		curtipes	SH0932674.10FU	Europe, Portugal/Spain	Mediterranean, sand
		sp. QU06	SH0932677.10FU	NA, Canada, Québec	
		sp. whartonensis	SH0932675.10FU	NA, USA, New Jersey	
		volvata	SH0932680.10FU	NA, USA, Texas/Louisiana/Missis-sipi/Indiana/New Jersey/New York	conifers/ hardwoods, sand
	Peckiana	*brunneomaculata*	SH0932678.10FU	Asia, China, Yunnan	*Pinus*
		lanigera	SH0932679.10FU	Asia, China/South Korea	*Pinus*/Fagaceae
		peckiana	SH0932672.10FU	NA, USA, Florida/Georgia/West Virginia/New York	*Pinus*, sand
	Fulvisquamea	*fulvisquamea*	SH0932673.10FU	Asia, Thailand/China/Taiwan/Japan	Dipterocarps/ *Pinus*
		sp. 'Sarawak1'		Asia, Malaysia, Sarawak	Dipterocarps
Sagittariae		aff. clarisquamosa2	SH0884338.10FU	Asia, China, Hunan	
		aff. clarisquamosa1	SH0884339.10FU	Asia, China, Yunnan	
		sp. OMS-013	SH0884342.10FU	Asia, Japan, Tokyo	
		sp. OMS-017	SH0884337.10FU	Asia, Japan, Tokyo	
		sp. montelongense		Europe, Portugal, Minho	*Quercus*
		aff. vaginata	SH0884340.10FU	NA, USA, North Carolina	
		sp. sagittaria	SH1158201.09FU	NA, USA, Florida	*Pinus*/*Quercus*
		sp. 17 GMB-2014	SH1031636.10FU	Oceania, Australia, Northern Territory	

Subsection	Strip	Epithet	SH (1.5%)	Geography	Ecology
Lepiotidae	Lepiotoides	*rufobrunnescens*	SH0858857.10FU	Asia, China, Guangdong	*Castanopsis*
		parvirufobrunnescens		Asia, India, Jammu & Kashmir	Pinaceae
		sp. JPN01/sp. OMS-015	SH0891149.10FU	Asia, Japan/South Korea	*Pinus*
		lepiotoides f. *subcylindrospora*	SH0891145.10FU	Europe, Spain/France	Fagaceae
		sp. QU05	SH0891151.10FU	NA, Canada, Québec	conifers/ hardwoods, sand
		sp. IN25	SH0891156.10FU	NA, USA, Indiana	*Quercus*, sand
		sp. IN33	SH0858858.10FU	NA, USA, Indiana	
		sp. insolens	SH0858859.10FU	NA, USA, Indiana/New Jersey	
		sp. texidella		NA, USA, Texas	mixed woods
	(unclassified)	sp. canadensis01/sp. IN28	SH0891154.10FU	NA, USA, Connecticut/Indiana	*Quercus*/*Carya*/ *Pinus*
	(unclassified)	sp. 19 GMB-2014	SH1029735.10FU	Oceania, Australia, Northern Territory	*Eucalyptus*
	Avellaneosquamosa	aff. avellaneosquamosa2	SH0891153.10FU	Asia, China, Anhui	mixed woods
		parvicurta	SH0891146.10FU	Asia, China, Fujian/Yunnan	Fagaceae
		cf. *avellaneosquamosa*1	SH0891147.10FU	Asia, China, Yunnan	*Pinus*
		aff. avellaneosquamosa3/OMS-012	SH0891155.10FU	Asia, China/Japan	conifers?
		aff. avellaneosquamosa4/aff. volvata3	SH0891142.10FU	Asia, Japan/South Korea	Fagaceae
		aff. volvata4	SH0891148.10FU	Asia, South Korea	
		aff. volvata5	SH0891144.10FU	Asia, South Korea, Gyeonggi	Hardwoods
		aff. avellaneosquamosa5	SH0891141.10FU	Asia, Taiwan, Nantou County	*Quercus*/*Pinus*

Subsection	Strip	Epithet	SH (1.5%)	Geography	Ecology
(Lepiotidae)	(Avella neosqu amosa)	sp. IN31	SH0891143.10FU	NA, USA, North Carolina/New York/Indiana	
		sp. S17/sp. IN41	SH0891150.10FU	NA, USA, Virginia/Pennsylva-nia/New Jersey/New York/New Hampshire/Massachu setts/Indiana/Wiscons in	Hardwoods
		aff. rosea		Oceania, Australia, New South Wales	*Eucalyptus*
		sp. 12 GMB-2014	SH0891152.10FU	Oceania, Australia, Northern Territory	
	Subvis cosa	aff. subviscosa	SH1045708.10FU	Africa, Benin	
		subviscosa s.l.	SH0846553.10FU	Africa, Benin	
		subviscosa s.l.	SH0846552.10FU	Africa, Zambia, Mufumbwe district	
		subviscosa s.l.	SH0846554.10FU	Africa, Zimbabwe	Fabaceae
		aff. goossensiae KM 12		Africa, Cameroon, Dja Biosphere Reserve	*Gilbertiodendro n*/*Uapaca*
		sp. TANZ01		Africa, Tanzania, Iringa	Fabaceae (Miom-bo), red soil
		sp. soil subtropical China 32		Asia, China	
		sp. soil subtropical China 33		Asia, China	
		sp. clone d28 3Ama	SH1013321.10FU	Asia, China, Eastern Himalaya	*Polygonum*
		sp. TUF116503	SH1013317.10FU	Asia, Thailand, Chiang Mai	
		sp. Thai04	SH1013318.10FU	Asia, Thailand, Isan	Dipterocarps
		sp. clone Ama3	SH1013320.10FU	Asia, Thailand, Kamphaeng Phet	Dipterocarpus
		sp. 'Sarawak2'		Asia, Malaysia, Sarawak	Dipterocarps
		sp. 'Jalisco'	SH1013319.10FU	NA, Mexico, Jalisco	*Quercus*/*Pinus*

Subsection (Lepiotidae)	Strip	Epithet	SH (1.5%)	Geography	Ecology
	Pseudo mutabilis	sp. 'Lupane'	SH0937059.10FU	Africa, Zimbabwe, Lupane	
		pseudorufobrunnescens		Asia, India, Uttarakhand	*Quercus*
		aff. volvata2		Asia, Japan, Yakushima Island	Fagaceae
		sp. 'California'	SH0944450.10FU	NA, USA, California	*Quercus*
		aff. mutabilis	SH0944451.10FU	NA, USA, Florida	*Quercus*?
	(unclassified)	aff. volvata1		Asia, China, Yunnan	
Ponderosae	Ponderosa	*ponderosa*	SH0916963.10FU	Europe, Portugal/Spain/France/ Italy/UK	*Quercus*/*Cistus*
		sp. whetstoneae	SH0916957.10FU	NA, USA, Mississipi/Missouri/ Tennessee/North Carolina/West Virginia/District of Columbia/Pennsylvania/New Jersey/New York/Indiana/Ohio	
	Claristriata	aff. avellaneosquamosa6	SH0916962.10FU	Asia, China, Yunnan	
		sp. OMS-010/aff. volvata6	SH0916961.10FU	Asia, Japan/South Korea	
		aff. volvata7	SH0916952.10FU	Asia, South Korea, Gangwon	
		aff. volvata8		Asia, South Korea	
		claristriata	SH0916955.10FU	Asia, Thailand, Chiang Mai	Dipterocarps
		sp. neocaesariensis/sp. IN26	SH0916954.10FU	NA, USA, New Jersey/New York/ Massachusetts/Indiana/Wisconsin	*Pinus*
		sp. S19/aff. volvata9	SH0916956.10FU	NA, USA, Tennessee/North Carolina/Virginia/New Jersey/Massachusetts	*Quercus*/*Carya*/*Tsuga*
		sp. T51	SH0916960.10FU	NA, USA, Texas/Mississippi	mixed woods

Subsection	Strip	Epithet	SH (1.5%)	Geography	Ecology
(Ponderosae)	Pinoph ila	aff. albidostipes	SH0916959.10FU	Asia, China, Jiangxi	
		pinophila	SH0916953.10FU	Asia, China, Yunnan	*Pinus*
Baakani		aff. goossensiae	SH0952146.10FU	Africa, Cameroon, Dja forest	
		cf. *goossensiae*	SH0883755.10FU	Africa, Cameroon, Dja forest	
		sp. 'Madagascar'	SH1029443.10FU	Africa, Madagascar, Mandena Conservation Zone	
		sp. Arora-01-560		Africa, Zambia, Copper Belt	
		sp. Sharp2816	SH0883752.10FU	Africa, Zimbabwe	Caesalpiniaceae
		sp. 'Naletali' s.l.	SH0883753.10FU SH0883754.10FU	Africa, Zimbabwe, Shangangwe	

Figure legends

Fig. 1 Phylogenetic trees obtained with BEAST (Suchard et al. 2018), with posterior probabilities above 50% shown to the left of each node. The sequence labels have the GenBank accession numbers followed by unique identifiers, and the proposed clades are highlighted with background colours. The scale bars (top left) are for substitutions per site. a) ITS1:ITS2 concatenate dataset, tree log likelihood −14,030.9, with the names for major clades (subsection level, corresponding to a different background colour) and, in smaller type, the minor (stirp level) clades. b) ncLSU dataset, tree log likelihood − 7311.5, background colours matching those of a)

Figure 1