Episomal virus maintenance enables bacterial population recovery from infection and virus-bacterial coexistence

Rodrigo Sanchez-Martinez[1]*, Akash Arani[2]*, Mart Krupovic[3], Joshua S. Weitz[2,4,5]**, Fernando Santos[1,7], Josefa Anton[1,6,7]**

Affiliations

[1] Department of Physiology, Genetics and Microbiology, University of Alicante, San Vicent del Raspeig, Alicante, Spain

[2] Department of Biology, University of Maryland, College Park, MD, USA

[3] Institut Pasteur, Université Paris Cité, Archaeal Virology Unit, Paris, France

[4] Department of Physics, University of Maryland, College Park, MD, USA

[5] Institut de Biologie, École Normale Supérieure, Paris, France

[6] Multidisciplinary Institute of Environmental Studies Ramon Margalef, Alicante, Spain

[7] Institute of Health and Biomedical Research of Alicante (ISABIAL), Alicante, Spain

* These authors contributed equally

** Corresponding authors: jsweitz@umd.edu; anton@ua.es

ABSTRACT

Hypersaline environments harbor the highest concentrations of virus-like particles (VLPs) reported for aquatic ecosystems. The substantial densities of both microbial populations and VLPs challenge traditional explanations of top-down control exerted by viruses. At close to saturation salinities, prokaryotic populations are dominated by *Archaea* and the bacterial clade *Salinibacter*. In this work we examine the episomal maintenance of a virus within a *Salinibacter ruber* host. We found that infected cultures of *Sal. ruber* M1 developed a population-level resistance and underwent systematic and reproducible recovery post infection that was counter-intuitively dependent on the multiplicity of infection (MOI), where higher MOI led to better host outcomes. Furthermore, we developed a nonlinear population dynamics model that successfully reproduced the qualitative features of the recovery. This suggests that the maintenance of the virus episomally, often referred to as pseudolysogeny, and lysis inhibition allow for host-virus co-existence under high MOI infections. Our results emphasize the ecological importance of exploring a spectrum of viral infection strategies beyond the conventional binary of lysis or lysogeny.

INTRODUCTION

Diverse microorganisms adapted to life at high salt concentrations are found in globally distributed hypersaline environments that comprise ~50% of continental waters (1). The abundance of prokaryotes increases with salinity, reaching values even higher than 10^8 cells per milliliter at salinities above 25% (2, 3, 4). The microbial communities in these environments, especially at extreme salinities, are dominated by members of the domain *Archaea* along with *Bacteria* predominantly from the *Salinibacter* clade (5, 6). Amongst bacterial clades, *Salinibacter ruber* was the first bacterium confirmed to grow actively in extreme salinities (7). Global-scale biogeographic studies have revealed that *Sal. ruber* is one of the most prevalent and dispersed bacterial lineages in hypersaline environments. Moreover, it is also characterized by a high intraspecific diversity, which can be caused as a response to an intense viral predation (8, 9, 10). At saturated salt conditions, hypersaline environments are also characterized by the presence of viral-like particles (VLP) that can reach up to 10^{10} VLP/ml (11), more than three orders of magnitude higher than estimates from surface marine communities (12, 13). Measured ratios of

viruses to microbes increase with salinity (from seawater to highly salt-saturated ponds) with

values between 10 and 100 VLPs per cell, and can even reach 300 VLPs per cell (14). Hypersaline

environments at close-to-saturation salinities are typically inhospitable to bacterivorous organisms

found in other aquatic habitats, including protists and heterotrophic nanoflagellates (15, 16, 17).

Unusually high viral abundances and virus-to-cell ratios along with reduced bacterivory in

hypersaline environments suggest that viral infections and the subsequent modulation of microbial

cell fate have a significant role in shaping prokaryotic population dynamics (18).

At the cellular scale, viral infections can lead to a continuum of outcomes spanning pathways that

lead to rapid lysis as well as long-term persistence via lysogeny (19). In the lytic pathway, the

virus redirects host metabolism to replicate the viral genome, produce virions, and lyse the host

cell, thereby releasing viral progeny to the environment. In the lysogenic pathway, the infecting

viral genome replicates along with the host chromosome. The provirus (i.e., the integrated viral

genome) can reactivate stochastically or in response to changes in host state, re-initiating the lytic

pathway leading to the release of virus particles (20, 21). Viral life cycles span the continuum

between these archetypes, including life cycles such as chronic infections, in which there is a

progressive release of virions into the environment without cell lysis (22). Another alternative viral

life cycle is pseudolysogeny, in which genomes of non-temperate viruses persist inside host cells

without integration as episomes, do not replicate synchronously with their host cell, and are

transmitted asymmetrically from mother to daughter cells (23, 24).

The presence of high microbial densities, low bacterivory, and unusually high VLP levels in

saturated brines presents a challenge to conventional explanations of top-down control. If

extremely high viral densities are achieved through efficient infection and lysis, one might expect

strong selection for viral-resistance via Kill-the-Winner like mechanisms (25, 26) or even the local

collapse of archaeal and bacterial communities (and their viruses) until recolonized via dispersal.

Alternatively, prokaryotes and viruses may persist at high levels in hypersaline environments

because of sustained production from infected cells or a high stability of non-infectious viruses.

Here, we utilize a *Sal. ruber*-virus model system to explore the mechanistic basis for the

persistence and coexistence of high-density populations of viruses and bacterial hosts in

hypersaline environments. Initial experiments revealed that virus infections led to a transient

bacterial population crash followed by a recovery. However, contrary to our expectations and

leveraging a combination of experiments, sequencing, and mathematical modeling, we conclude

that bacterial population recovery was not enabled by the emergence of virus-resistant bacterial

mutants or lysogeny. Instead, bacterial population recovery was enabled via the initiation of a

persistent infection characterized by the episomal maintenance of viruses that protected infected

cells from subsequent infection and lysis. As we show, this persistent infection enables lytic

viruses to persist at high abundances without necessarily eliminating their hosts, providing new

challenges to paradigms of virus-microbe coexistence in extreme environments.

RESULTS

Systematic and reproducible recovery of *Sal. ruber* populations after viral-induced lysis

We infected *Sal. ruber* strain M1 with the virulent M1EM-1 virus (27) (named henceforth EM1

for convenience) at a MOI=0.01 in triplicate at time=0 h (Fig. 1a). In all three replicates, we

observed a rapid stabilization in the host optical density (OD) following infection due to lysis,

followed by a population-level recovery at 80 h such that infected cultures eventually recovered

to the levels comparable to those of the virus-free control, reaching the stationary state ODs (see

Methods for further details). This systematic recovery of the population post infection was also

observed in other *Sal. ruber*-virus pairs (see Extended Data Fig. 1). This finding was unexpected

as the EM1 virus does not encode lysogenic markers (i.e., excisionases and integrases) which

would allow its integration into the host chromosome, facilitating superinfection exclusion-

mediated population level recovery as seen for some temperate viruses (28). We further examined

this recovery phenomenon by infecting *Sal. ruber* M1 with the EM1 virus at ten-fold higher viral

levels (MOI=0.1) and using more replicates (10 replicates for both control and infection treatment;

Fig. 1b). In this case, we observed a sharp decrease in the OD levels of the infected cultures,

consistent with the 10x increase in viral pressure (MOI=0.1 vs. MOI=0.01). However, we once

again observed recovery dynamics comparable in all ten replicates.

Sal. ruber recovery is not consistent with the proliferation of virus-resistant mutants or the integration of viral genomes into bacterial genomes

113
114

We hypothesized that reproducible recovery of infected cell cultures is not caused by spontaneous
mutations and the emergence and proliferation of virus-resistant bacterial populations which
typically involves jackpot like variability that is incompatible with the observed reproducibility in
recovery times (29). Nonetheless, to assess the link between recovery and virus-resistance, all
replicates (both the infected cultures and the controls) from the recovered cultures shown in Fig.
1a (named C_F) were sequenced. The initial *Sal. ruber* M1 used as a host (named C_0) was also
sequenced as a control. We then analyzed and compared the sequenced genomes to evaluate the
evidence for selective sweeps of virus-resistance genes in infected replicates.

Only 6 mutations showed a frequency close to 100% (Supplementary Table 1) in all the infected
replicates. Of these 6 mutations, 3 involved non-synonymous changes, affecting an ATP-
dependent helicase/deoxyribonuclease, an orotate phosphoribosyltransferase and a
sodium/glucose cotransporter. The first two are related to helicase/exonuclease activities and to
the synthesis of pyrimidines, respectively, so they did not seem to have a direct relationship with
virus resistance. The third mutation, affecting the stop codon in the synthesis of the sodium/glucose
cotransporter, could be involved in the resistance, as it affects a transmembrane protein. However,
upon further examination, we found that both resistant and sensitive cells had the same mutation,
confirming that it was unrelated to resistance. In addition, it was subsequently found that cells that
regained susceptibility to the virus retained all the mutations mentioned above (see below and
Mutations Supplementary Information for further details). Therefore, no mutations could be
unequivocally tied to the observed population recovery phenomenon in the infected cultures.

According to previous studies (30), these dynamics after viral infection could be explained by the
integration of the viral genome into the chromosome of its host, yielding lysogens resistant to
superinfection with the same virus. Although EM1 lacked any lysogeny-associated traits in its
genome, including excisionase and integrases (Supplementary Table 2), the reads of the infected
C_F cultures were reexamined to search for hybrid host-virus reads, which would signify integration
into the chromosome (see Methods). No hybrid reads were found, and therefore recovery cannot
be attributed to virus integration. However, we observed a remarkably high sequencing depth of
the virus genome within infected C_F cultures, averaging 146 nt/position, while the cellular
chromosome exhibited 116 nt/position (Supplementary Table 3). This similar level of sequencing

depths suggests that viral genomes are maintained intracellularly in the recovered cultures, albeit
not via integration as a prophage.

Infected cells with episomally maintained viruses comprise part of the recovered *Sal. ruber* population

The lack of evidence for mutations related to virus resistance or integrated prophage within
recovered *Sal. ruber* populations led us to hypothesize that recovered bacterial populations
infected by viruses might maintain viral genomes intracellularly in an episomal form. To explore
this alternative, we grew colonies from time point C_0 (initial *Sal. ruber* M1) and C_F (final recovered
cells). Twenty individual colonies from both C_0 and C_F were grown in liquid medium and PCR-
tested for the presence of the EM1 virus. As expected, all C_0 colonies tested positive with the
specific primers for *Sal. ruber* M1 but tested negative for the virus. However, all the final colonies
from the infected cultures tested positive for the EM1 virus. Three randomly selected cultures from
the C_F, colonies that tested PCR positive for the virus (which we refer to as 1R, 2R and 3R
respectively), were stained with SYBR^{TM}Gold and observed under the epifluorescence microscope
(Fig. 2a). After observation of more than thirty fields per culture, extracellular virions were not
observed in any case. Hence, although EM1 virus was present in the culture, its virions were not
detected extracellularly. The intracellular presence of viral genomes was confirmed by pulsed field
gel electrophoresis (PFGE) with DNA from the final colony 1R, but not from the control C_0 colony
(wild type *Sal. ruber* M1). The PFGE showed a band with the size of the EM1 virus genome (35
kb) in 1R, which was confirmed by Southern blot hybridization with an EM1-specific probe (Fig.
2b). These results, in conjunction with the sequencing data that revealed no signs of integration
(no hybrid host-virus reads were found), the lack of integrases and/or recombinases in the genome
of EM1 (Supplementary Table 2), supports the hypothesis that the EM1 virus was maintained in
an episomal form within *Sal. ruber* M1 recovered populations. In addition, the terminal
redundancy found in sequencing may indicate that it is maintained within cells as a circular
episome.

Episomal maintenance of viruses protects recovered cells from subsequent infection and viral-induced lysis

To test whether the acquisition and episomal maintenance of the virus had an influence on the
recovered cell growth, we conducted growth experiments on 1R and compared the population
dynamics to that of the wild type *Sal. ruber* M1 (Fig. 3a). We found no statistical differences in
the reached OD or growth dynamics between 1R and *Sal. ruber* M1 (*Modeling Supplementary*
Information: Sal. ruber M1 wild-type and 1R (pseudolysogen) growth comparison). Hence, there
is no evidence that viral acquisition provides a direct cost or benefit to host growth. However, the
benefits of intracellular maintenance of viral genomes may be contextual.

During typical lysogeny, the prophages can protect hosts from viral infection and lysis via
superinfection exclusion (31, 32, 33). We hypothesize that similar principles may apply to
recovered bacterial populations. To test for superinfection exclusion in recovered populations, we
conducted spot test assays with 1R and the wild type *Sal. ruber* M1. In both cases, we plated cells
and then overlaid agar plates with spots of EM1 virus serially diluted from 10^{10} PFU/ml to 10^4
PFU/ml (see Methods). The wild type strain was susceptible to EM1 infection, whereas 1R
exhibited resistance (Fig. 3b). We interpret the results of the spot assay experiment to imply that
episomal maintenance of the EM1 virus confers protection against subsequent EM1 infection and
lysis. Furthermore, by conducting an adsorption assay with 1R and the wild type *Sal. ruber* M1,
we found that protection against subsequent EM1 infection occurs at the surface level, i.e.,
preventing effective adsorption (Extended Data Fig. 2).

Finally, to test how stable is the intracellular maintenance of EM1, we plated the 1R strain and
picked colonies at random to test for the presence of the virus by PCR. The results showed that 13
of 17 colonies retained the virus (*Mutations Supplementary Information*). A spot test with EM1 on
the cultures derived from the 17 colonies revealed that only those strains that had lost the virus
genome were sensitive to EM1 infection, confirming that the resistance was due to the virus
acquisition and episomal maintenance. Subsequent genomic sequencing of 3 of these sensitive and
3 resistant colonies (all derived from 1R) showed that the mutations observed in recovered cultures
following the infection (see above) had not been lost and therefore were not involved in the
development of resistance to EM1 infection (*Mutations Supplementary Information*). The
reappearance of sensitive strains within the 1R recovered population suggests that the viral genome
is not replicating synchronously or faithfully segregated with the host chromosome. Consistently,
no marker was found in the virus genome that would indicate cell-synchronous replication and

partition, such as the Par system of bacteriophage P1 (34) (Supplementary Table 2). Therefore, we
hypothesize the viral persistence inside recovered cells is enabled through asymmetric episomal
virus transmission during cell division consistent with a pseudolysogenic state (see (24)). For the
remainder of the manuscript, we refer to bacterial populations post-infection as 'recovered' and
use the term resistant pseudolysogens to refer to the subset of bacterial populations that have
episomally maintained the virus genome and are resistant to subsequent infection and lysis by the
same virus.

Bacterial population recovery depends non-monotonically on MOI

The spot assay experiment yielded a counter-intuitive result. When high virus concentrations were
added on the plates of wild type *Sal. ruber* M1 at levels between MOI of 10-100, we observed
growth of cells in the center of spots. This suggests that there may be a MOI dependence to the
initiation of pseudolysogeny, akin to that found in classic studies of lysogeny in which the
probability of lysogeny increases with increasing MOI (21, 35, 36). Hence, cultures with cells
infected by multiple viruses may be more likely to yield pseudolysogens which can grow even in
the presence of viruses. To explore the MOI-dependence of recovery further, *Sal. ruber* M1 was
infected with EM1 at MOIs ranging from 0.01 to 10 viruses per host cell. The M1 population was
infected during its exponential growth phase (66 h). The population dynamics (Fig. 3c) exhibit dip
and recovery dynamics for MOI 1, 5 and 10 in a similar fashion to the initial infection curve
experiment. In contrast, population dynamics for MOI 0.01 and 0.1 solely exhibit growth, albeit
slightly slower than in the control population. Notably, the final cell densities in the experiments
with MOI 1, 5 and 10 were non-monotonically related to MOI, i.e., the OD values were 0.22 +/-
0.01, 0.18 +/- 0.02 and 0.27 +/- 0.01 for MOI 1, 5, and 10, respectively. The OD value at time 230
hours for MOI 10 was statistically significantly different from the OD value for MOI 1 (t-test,
p=0.041) and MOI 5 (t-test, p=0.038). Note that we could not reject the null hypothesis that the
OD values at time 230 hours for MOI 1 vs 5 were statistically different (t-test, p=0.206). These
results raise the question: if EM1 generates resistant pseudolysogens (as was confirmed in the spot
assay experiment for 1R (see Fig. 3b)), then why would recovery dynamics depend non-
monotonically on MOI?

Typically, superinfection exclusion requires certain cellular rearrangement post-infection (31, 32,

37). As such, we hypothesize that multiple EM1 viruses can jointly infect the same *Sal. ruber* M1

cell, insofar as multiple infections precede the development of resistance (38). For example, given

a MOI of 10 and assuming that all virions have adsorbed to the cell and injected their genomes,

the infection would yield an average of 10 viruses per cell with a standard deviation of ~3.

Therefore, to test whether superinfection was possible in this system, we performed adsorption

assays with the wild type *Sal. ruber* M1 and EM1 at MOI of 0.01 and 10 (Fig. 3d). In the case of

MOI 0.01, the decrease of free viruses was equivalent to that expected in theory for complete

adsorption of virions onto cells (i.e., approximately 1% of cells should be infected). In the case of

MOI 10, the exponential decrease of free viruses plateaus at ~20%, implying that 80% of virions

adsorbed to cells, consistent both with superinfection (i.e., the average cellular-level MOI would

be 8) and interference that limits viral superinfection at high MOI. Hence, we conclude that

superinfection of *Sal. ruber* M1 is possible by the EM1 virus given sufficiently short intervals

between infection events even if the development of a superinfection exclusion state prevents

subsequent infection over the long term in resistant pseudolysogens.

Scaling-up the dynamical effects of pseudolysogeny from cells to populations

The contextual impact of superinfection raises questions on how pseudolysogeny modulates

recovery dynamics, in light of three key empirical observations (Fig. 3c). First, the *Sal. ruber* M1

populations infected at MOI 0.1 and 0.01 do not exhibit an observable population decline. Second,

the M1 host population recovers to a higher population densities when infected at MOI 10

compared to MOI 5 and MOI 1, highlighting a non-monotonic relationship between viral levels

and population-level lysis. Third, when infected at MOI 10, the *Sal. ruber* M1 population does not

crash immediately upon infection and maintains persistent transient growth post infection. To

connect mechanisms occurring at cellular-scales with these population-level dynamics, we

developed a nonlinear ordinary differential equation (ODE) model of the EM1 virus interacting

with *Sal. ruber* M1 hosts (Fig. 4). The model includes dynamics of resources (R), sensitive cells

(S), pseudolysogens (P) and virus particles (V) (39). We separate pseudolysogens into "early" (P_e)

and "fully developed" pseudolysogens (P_f), where viruses can infect sensitive cells and early

pseudolysogens (consistent with findings of Fig. 3d) which then transition to become resistant, P_f

types (consistent with findings of Fig. 3b). We account for cellular-level multiplicity of infection

of the pseudolysogens through a superscript [k] where k denotes the number of viral genomes per

cell e.g., $P_e^{[2]}$ denotes the number of early pseudolysogens with 2 viral genome copies. The number

of viruses able to infect a single cell is limited, consistent with adsorption results from Fig. 3d

(details in Extended Data Fig. 3). Cell division for both sensitive cells and pseudolysogens is

modeled as a function of resource consumption. Full details of the model specification and

parameters are provided in the Supplement (*Modeling Supplementary Information: Main Model*).

Simulated population dynamics (Fig. 5a) recapitulate the key empirical observations (as shown in

Fig. 3c). The match between theory and experiment is driven by the following mechanisms. First,

the model assumes that host lysis decreases with decreasing resources (40, 41). In Fig. 5b, we

show that decoupling host lysis from resource concentrations leads to a rapid crash in *Sal. ruber*

dynamics amongst the low MOI populations that is inconsistent with experimental observations.

Second, the model assumes that cellular lysis rates decrease as a function of increasing cellular

MOI (42, 43). In Fig. 5c, we show that if lysis rates are independent of cellular MOI, then

population crashes are most apparent at the highest MOI and decrease with decreasing MOI,

inconsistent with experimental observations. Third, we assume that pseudolysogens divide and

that their daughter cells can inherit viral genomes, albeit randomly, from the parent cell (akin to

random segregation of episomal elements (44)). In Fig. 5d, we show that if pseudolysogens do not

divide then the *Sal. ruber* population at MOI 10 (yellow curve) immediately dips upon infection

at 66 hours (blue dotted line) and there is no persistent transient growth, thereby inconsistent with

the experimental dynamics. In *Modeling Supplementary Information*: *Subsets of main model with*

corresponding dynamics, we explore all potential combinations of the presence/absence of each

mechanism and find that the model only recapitulates observed dynamics when all three

mechanisms operate jointly. Hence, we contend that each of these elements (resource-dependent

lysis, latent period delays in multiply infected cells, and pseudolysogen cellular division) are

necessary to explain experimentally observed dynamics of the EM1 virus and its host across 3

orders of magnitude difference in initial ratios of virus to host. The details of how all the model

features are implemented is highlighted under *Modeling Supplementary Information: Key Model*

Features and *Modeling Supplementary Information: Viral passage in superinfected*

pseudolysogens. Finally, we note that the results are robust to variation in the variability associated

with the passage of episomally maintained viruses to daughter cells during replication (*Modeling*

Supplementary Information: Episomal viral replication).

Episomal maintenance of viruses and the population-scale maintenance of virus-microbe coexistence

Finally, we set out to explore the extent to which episomal maintenance of viruses inside recovered bacterial populations could enable virus-microbe coexistence over longer time scales. The persistence of the virus within the pseudolysogen population was tested by transferring pseudolysogen 1R every 9-10 generations to fresh medium. We made four such transfers, resulting in ~ 40 generations of pseudolysogen growth and cell division (Extended Data Fig. 4a). At the end of the exponential phase of each transfer, we conducted a PCR test and found that the virus was not lost over time (Extended Data Fig. 4b). Upon qPCR analysis, we found that the ratio of virus genome/chromosome was close to 1 in the first transfer and upon further transfers, the ratio increased to 2-3 virus genomes per chromosome of M1 and was maintained at this level (Extended Data Fig. 4c). To determine whether viral resistance was maintained within the population, cells from each of the aforementioned transfers were exposed to EM1 through a spot test. We found that the cells from every transfer exhibited resistance to additional infection (Extended Data Fig. 5). Next, to observe how the pseudolysogens might ecologically interact with other *Sal. ruber* strains, we conducted spot tests in which eight strains of *Sal. ruber* used routinely in our laboratory were individually exposed to pseudolysogen 1R. We observed that *Sal. ruber* M1, the only strain sensitive to the EM1 virus, was killed while the rest of the seven strains were maintained (Extended Data Fig. 6). Similar dynamics have been observed with temperate viruses, where lysogens induce and release free virions as a weapon to kill sensitive cells in shared environments (45). We hypothesize that a subset of pseudolysogens of *Sal. ruber* reactivate the lytic pathway, releasing virions that infect sensitive competitors, generating a mix of pseudolysogens and increasing the level of extracellular virions.

DISCUSSION

Here we explored the dynamics arising from interactions between a virulent virus (EM1) and one strain of a prevalent and globally dispersed bacterial host found in hypersaline environments (*Sal. ruber* M1). We found that *Sal. ruber* M1 populations reproducibly recover from infections by EM1 viruses across three orders of magnitude variation in MOI. Notably, M1 bacterial populations

recover to higher densities when infected with more viruses (MOI=10) than with less viruses (MOI

1 or 5) while recovering to densities comparable to those of non-infected controls for MOI levels

of 0.1 or 0.01. Both sequence and experimental evidence suggest that recovered populations have

a significant fraction of cells with episomally maintained virus genomes that pass asymmetrically

from mother to daughter cells in the absence of synchronized replication with their cellular hosts.

Subsequent infection experiments revealed that these cells with episomally maintained virus

genomes (i.e., pseudolysogens) are resistant to infection and lysis by the original virus, providing

an opportunity to explore mechanisms by which infection recovery could stabilize bacteria-virus

feedback. We developed and analyzed a nonlinear population dynamics model that recapitulated

key features of the MOI dependent recovery dynamics that depend on the potential joint infections

of cells, lysis inhibition via superinfection immunity, and lysis inhibition due to resource

limitation. Altogether, the initiation and maintenance of pseudolysogeny stabilizes short-term

population dynamics while providing a mechanism for long-term coexistence of viruses and their

bacterial hosts in extreme environments.

The current study connects cellular-level interactions to feedback mechanisms that stabilize virus-

host population dynamics that could otherwise lead to population collapse, especially at high MOI

values. While previous studies have shown similar recovery dynamics in different virus-host

systems (46, 47, 48, 49), there has been limited effort to explain the mechanisms responsible.

Studies observing recovery dynamics phenomenon with temperate viruses have pointed to

emergence of resistant lysogens which are protected against further infections (28, 30). However,

bacterial population recovery has also been observed in viruses without any lysogeny markers as

well (50, 51), in which the emergence of resistance has been overlooked or simply attributed to

spontaneous mutations of the host (52, 53). Here, recovery dynamics could not be attributed to

lysogeny given the use of the *Sal. ruber* M1-EM1 system where EM1 is strictly virulent. The

recovery dynamics observed here was also highly reproducible, inconsistent with recovery due to

the emergence and expansion of spontaneous resistant mutants that grow from low to high

frequency where substantial variability is a hallmark feature of dynamics (29). Furthermore, our

analysis revealed no clear resistance-associated mutations in the recovered M1 populations.

Similarly, we also identified lysis inhibition at high MOI and/or when resources are limited as an

essential part of explaining the MOI-dependence of recovery dynamics. Lysis inhibition has

commonly been noted in infection of *Escherichia coli* by T4 (40, 42, 53, 54), with some studies

suggesting that it plays a key role in the maintenance and development of *E. coli*-T4
pseudolysogens (24, 54). We hypothesize that lysis inhibition could similarly be a key feature of
pseudolysogeny in the *Sal. ruber* M1-EM1 system, allowing for persistent infection in starved
conditions and/or under high MOI.

Our combined study comes with caveats. First, we focused primarily on contextual benefits of
pseudolysogeny, noting that identifying direct impacts of episomal maintenance of viruses on
cellular fitness has limited support (55). While we found that there were no statistical differences
between the growth rate of pseudolysogens and wild type host cells, we found that pseudolysogeny
provided surface-level protection against further infection. Furthermore, we found that loss of
episomally maintained viruses renders pseudolysogens sensitive to further infection. Hence, viral
acquisition comes with a cost (the risk of lytic infection) but is key to enabling surface-level
resistance and population level recovery. More work is needed to understand when pseudolysogens
become resistant and the cellular mechanisms underlying transient protection to infection. We also
found that *Sal. ruber* strains that were sensitive to the EM1 virus were killed in the presence of
pseudolysogens. Similar dynamics have been observed with temperate viruses of *Roseobacter*,
where lysogens induce and release virions to kill competing sensitive strains in head-to-head
experiments (45). We hypothesize that *Sal. ruber* M1 pseudolysogens may switch from a
maintenance to an actively infected state, e.g., when episomal numbers are low and/or when
resources become available. Again, future work will be needed to elucidate the cellular drivers of
lytic initiation and the extent to which intrinsic or extrinsic heterogeneity plays a role in the lysis
of pseudolysogens that release EM1 viruses that may then infect and kill competing sensitive
strains and potentially initiate new pseudolysogens.

In closing, this study reveals how feedback mechanisms at the cellular scale can transform
population dynamics leading to non-monotonic relationships between virus infection pressure and
bacterial impacts. Put simply: for strictly lytic viruses, more infection does not always result in
more killing. Within the *Sal. ruber* M1-EM1 bacteria-virus model system, we have identified
feedback mechanisms that limit lysis, whether because of multiple infections, resource limitation,
or the establishment of an episomally maintained virus (i.e., a pseudolysogen). Each of these
mechanisms stabilizes virus-microbe population dynamics. We recognize that it is challenging to
move from experimental settings to extreme field conditions, hence we are forced - for now - to

speculate that pseudolysogeny may be an evolutionarily adaptive strategy for obligately lytic
viruses to preserve hosts until environmental conditions improve or host density increases (23, 24,
56). Depending on the durability of viruses extracellularly, such a balance between infection and
production may help to explain virus:host ratios ranging from 10 to as high as 300 in the face of
fluctuating conditions (14). More broadly, it may be of some value to explore other, related cellular
mechanisms that stabilize virus-host population dynamics, enabling reproducible recovery
dynamics seen in ecosystems outside hypersaline environments, such as the bloom-bust dynamics
observed in oceanic *Emiliania huxleyi* populations after infection by EHV at high MOI (46, 57) or
the long-term persistence of crAss-like phage crAss001 and its host (58). Overall, the results from
this work highlight the ecological relevance of exploring a continuum of viral infection strategies
beyond the dichotomous options of lysis or lysogeny.

METHODS

Growth conditions and DNA extraction

Sal. ruber strain M1 was grown aerobically with gentle shaking (60 rpm) at 37 ºC in 25% SW (sea
water) with 0.2% yeast extract (59). *Sal. ruber* M1 was infected with the lytic virus EM1 (27),
using a multiplicity of infection (MOI) of 0.01 plaque forming units (PFUs) per cell at time 0 h.
Briefly, viruses were mixed with cells at exponential phase (optical density, OD, at 600 nm=0.3)
and incubated during 30 minutes at room temperature without shaking to facilitate the adsorption.
After the adsorption, the abovementioned culture medium was added up to 25 ml and incubated
as previously described. Infections were monitored by OD at 600 nm. Non-infected *Sal. ruber* M1
was grown as the control. All experiments were conducted in triplicate. Another curve was made
with the same methodology but using a MOI of 0.1 and 10 replicates in both control and infection.

Bacterial pellets of the initial (C_0, before virus addition) and final cells (C_F) of the curves performed
using a MOI of 0.01 were obtained by centrifugation at 17.000 xg during 10 min. The cells were
washed three times with sterile medium, the supernatant removed and the cell pellets stored at -80
ºC. C_0 and C_F DNAs were extracted with the DNeasy Blood & Tissue Kit (Qiagen, Hilden,

Germany) following the manufacturer's protocol, and nucleic acids eluted in 70 µl of milli-Q water

and quantified using Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, USA).

DNA sequencing and bioinformatic analysis

DNA was sequenced on an Illumina HiSeq (2 x 150 bp) at Novogene (Novogene, Seur, UK).

Primers and adapters were removed from sequences, and reads were filtered based on quality

scores using Trimmomatic v0.36.0 (60). Trimmed reads were assembled using SPAdes v3.13.1

with the trusted option using the reference genome of *Sal. ruber* M1 as scaffold (61). The mean

sequencing depth was calculated by a BLASTN-recruitment analysis where the assembled contigs

were used as reference.

For the search for hybrid host-virus reads, a BLASTN of the trimmed reads was performed, using

the genome of *Sal. suber* M1 of NCBI as a database (GenBank accession number:

NZ_CP030364.1). The reads with >95% ID were filtered by best hit and extracted. No horizontal

coverage filter was used. Another BLASTN analysis was performed with these reads using the

genome of the EM1 virus of NCBI as a database (GenBank accession number: NC_042348.1) with

the same parameters. No hybrid host-virus reads were found.

Mutations generated during the infective process were identified comparing reads from C_F with

the reference contigs assembled from C_0 using Geneious software 6.1.8 (Biomatters, Auckland,

New Zealand) with a minimum variant frequency of 0.01% and a coverage >25% to the mean

coverage.

PCR and epifluorescence microscopy

Cells from C_0 and C_F were diluted and plated on SW 25% with 0.2% yeast extract agar plates

(concentration 2%) and incubated at 37 °C for 30-45 days until colonies visualization. Twenty

colonies from every point (C_0 and C_F) were grown in 2 ml of culture medium at 37 °C and 60 rpm.

Once the cultures grew to late exponential phase, 200 µl of each were washed three times by

centrifugation (see above). DNA was extracted from washed pellets by boiling at 100 ºC for 10

min after resuspending the cells in 80 µl of milli-Q water. Two parallel PCR reactions were

performed with each DNA, using specific primers for *Sal. ruber* M1 and the virus EM1

(Supplementary Table 4). PCR reactions were carried out in a final volume of 25 µl, containing:

0.75 µl of 1.5 mM MgCl₂, 2.5 µl of 10X reaction buffer, 0.5 µl of 10 mM dNTPs, 0.1 µl of Taq

polymerase (5 U/µl, Invitrogen, Waltham, USA), 0.5 µl of 10 µM primers, 1 µl of extracted DNA

and milli-Q water to complete the final volume. PCR conditions are described in Supplementary

Table 5. PCR products were electrophoresed and UV-visualized after staining with ethidium

bromide (100 µg/ml).

Three random cultures from C_F colonies that were PCR positive for both *Sal*. *ruber* M1 and the

EM1 virus (named 1R, 2R and 3R) were selected for the visualization of extracellular virions by

SYBR™Gold staining (62). Briefly, 100 µl of the culture were fixed with formaldehyde (4% final

concentration) for 1 h at 4 ºC. Fixation was stopped by adding phosphate-buffered saline (PBS)

1X up to 1 ml. Twenty microliters (equivalent to 2 µl of sample) were filtered through 0.02 µm

Anodisc 25 filters (Whatman Int. Ltd, Maidstone, UK) to retain cells and viruses. Filters were

stained with SYBR™Gold (25X) for 15 min, washed twice for 1 min with milli-Q water and

visualized with an epifluorescence microscope (Leica, type DM4000B).

Pulsed-field gel electrophoresis (PFGE) and Southern Blot

One ml of liquid cultures of C_0, PCR positive for M1, and C_F, PCR positive for M1 and EM1, were

centrifuged and the cell pellets mixed with agarose at 0.8% (final concentration) to obtain plugs,

as detailed in Santos et al., 2007 (63). Plugs were incubated overnight at 50 ºC in ESP (0.5 M

EDTA, pH 9.0, 1% N-laurylsarcosine, 1 mg ml⁻¹ proteinase K) for cell lysis and virion

disruption and stored at 4 ºC. DNA was separated by PFGE on a 1% LE agarose gel in 0.5X TBE

buffer, using a Bio-Rad CHEF DR-III System (Bio-Rad, Richmond, USA), under the following

conditions: 8–12 s pulse ramp, 6 V/cm, at 14 °C for 30 hrs. After electrophoresis, gels were stained

with ethidium bromide (100 µg/ml) for 10 min, washed with distilled water for 30 min, and UV-

visualized with a transilluminator.

Agarose gels were transferred to a positively charged nylon membrane (GE Healthcare,

Buckinghamshire, UK) as previously described (64). In parallel, DNA from the EM1 virus was

labelled to be hybridized against the membrane using the DIG High Prime DNA Labelling and

Detection Starter Kit II (Roche, Mannheim, Germany), following manufacturer's instructions.

Growth curves, spot tests and adsorption assays

Sal. ruber M1 wild type (as a control) and pseudolysogen 1R were grown in liquid, transferred to

fresh media (1% v/v) and incubated at 37 ℃ with gentle shaking at 60 rpm. Cellular density was

measured by OD at 600 nm. The growth curves of *Sal. ruber* M1 wild type and pseudolysogen 1R

were compared for statistical differences using 99% confidence intervals obtained through the

Delta method (*Modeling Supplementary Information: Sal. ruber M1 wild-type and 1R*

(pseudolysogen) growth comparison).

To test the resistance/susceptibility of *Sal. ruber* M1 wild type and pseudolysogen 1R to the EM1

virus, both hosts were exposed to the virus by a spot test. Four ml of molten 0.7% top agar of 25%

SW with 0.2% yeast extract were mixed with 500 μl of bacterial cultures at exponential phase and

plated on solid medium. Once solidified, 3 μl of the EM1 virus, tittered at 10^{10} PFUs/ml and diluted

10^2, 10^4 and 10^6 times, were added to the lawn. The spotted plates were left to dry and incubated

at 37 ℃ for 10 days. The detection of a clearance zone was interpreted as evidence of viral lysis

while the areas exhibiting growth were attributed to resistance against EM1.

An adsorption assay was performed with the EM1 virus and *Sal. ruber* M1 wild type. Cultures in

exponential phase were diluted with fresh medium to an optical density of 0.3 and $2x10^8$ cells were

mixed with $2x10^6$ PFU (MOI=0.01) or $2x10^9$ PFU (MOI=10) of the EM1 virus. This was

considered as the initial time. It was incubated at 37 ℃ and 60 rpm. Aliquots of 150 μl were taken

along 2 h (at 15, 30, 45, 60, 90 and 120 min), centrifuged at 17.000 xg for 8 min, and 120 μl of the

cells-free supernatant were stored in ice. The number of infective free viruses was measured by

plaque assay, diluting the supernatants in sterile SW 25% and mixing 100 μl of each with 500 μl

of *Sal. ruber* M1 wild type in exponential phase. Four ml of top agar were added and this was

plated on solid plaques. After solidifying, the plates were incubated at 37 ℃ for 10 days until

plaques were visible. The number of free infective viruses (PFU/ml) were counted and represented.

A control of the viral decay of the EM1 virus was added, in which the virus was incubated without

host. All the experiments were conducted in triplicate.

To get more information about the resistance mechanism, an adsorption assay was performed with

the EM1 virus and *Sal. ruber* M1 wild type and pseudolysogen 1R with a MOI of 0.01 as described

above.

To calculate the adsorption rate, the graph of the free infective viruses (PFU/mL) was plotted with
respect to time and fit to an exponential function. As the assay was conducted within a 120 min
time period and the latent period of the EM1 virus is around 21-22 h, we can assume that there is
negligible killing and the susceptible host population stays constant at $S = 2x10^8$ cells/ml. We fit
the adsorption data to the function $V = V_0 e^{-\phi St}$ and get an adsorption rate $\phi \approx 4x10^{-10}$ ml/h
(Fig. S11).

Infection curves at varying MOIs and revival of lysis in pseudolysogens

Sal. ruber M1 wild type was grown at 37 ºC and 60 rpm. The EM1 virus was added at 67 h at
middle exponential phase. Cultures were infected with a MOI of 0.01, 0.1, 1, 5 and 10 plaque
forming units (PFUs) per cell in separate experiments. Infections were monitored by OD at 600
nm. *Sal. ruber* M1 was grown without viruses as a control. All experiments were conducted in
triplicate.

A total of $2x10^8$ cells from the end of each infection curve were transferred to 4 ml of fresh medium
(SW 25% with 0.2% yeast extract). This caused all cultures to be diluted to an OD of
approximately 0.1. Cultures were monitored by OD at 600 nm.

Numerical Simulations

All the numerical simulations were performed on Python 3.11.5 through the Jupyter IDE. To solve
the model ODEs and obtain the simulated results, we used the numerical integration module *odeint*
from the *SciPy* package. To convert the experimental OD measurements to concentrations, we fit
a calibration curve using the *curve_fit* module from the *SciPy* package. All the code that can
reproduce the simulated results seen in this paper are available on the Github:
https://github.com/aranilah/SalruberM1_EM1_recovery/tree/main

Transfers of pseudolysogen 1R to fresh media, PCR and qPCR

In order to assess if the EM1 virus was with generations, pseudolysogen 1R was transferred four
times to fresh media. Briefly, pseudolysogen 1R was transferred in triplicates to fresh media (1 %
v/v) and incubated at 37 ºC with shaking at 60 rpm, measuring the growth by OD at 600 nm. At
the late exponential phase cultures were transferred again to fresh media until a total of 4 transfers.

A spot test of each transfer was also carried out as previously mentioned to check if the different

transfers of pseudolysogen 1R were still resistant to the virus. Moreover, 100 µl of each transfer

were fixed with formaldehyde (0.5% final concentration to avoid PCR inhibition) for 1 h at 4 ℃.

Fixation was stopped by adding phosphate-buffered saline (PBS) 1X up to 1 ml.

To eliminate free viruses, assuming that viruses could have been spontaneously released during

the transfer periods, the fixed samples were centrifuged at 17.000 xg during 10 min and washed

with PBS five times. PCR reactions for *Sal. ruber* M1 and the EM1 virus were carried out as

mentioned above, with 1 µl of the fixed and washed cells as DNA template (first, we checked that

the PCR initial thermal shock was sufficient for cell lysis; data not shown).

Once the presence of EM1 was detected in all the transfers, a qPCR was performed to monitor

virus concentration along the subsequent transfers. The qPCR assay was conducted using TaqMan

hydrolysis probes (Supplementary Table 4) labeled with hexachlorofluorescein (for *Sal. ruber* M1)

and fluorescein (for EM1). The experiment was carried out using the standard run in a

StepOnePlus™ PCR System (Life Technologies, Carlsbad, USA) in a 10 µl reaction mixture with

PrimeTime™ Gene Expression Master Mix (Integrated DNA Technologies, Coralville, USA). The

reaction contained: 5 µl of 2X Master Mix, 0.2 µl of each 10 µM primer, 0.2 µl of 10 µM TaqMan

probe, 1 µl of fixed sample and milli-Q water to complete volume. Conditions are detailed in

Supplementary Table 6. The results were analyzed with the Applied Biosystems StepOne™

Instrument program. All samples were run in triplicate (including the standards and negative

controls).

Plating of pseudolysogen 1R, mutation screening and spot test

Pseudolysogen 1R at late exponential phase was plated on solid medium (SW 25% with 0.2%

yeast extract and 2% agar) to obtain individual colonies. After two months of incubation, 50

colonies developed, which were picked and transferred to 1 ml of liquid media each. The presence

of the EM1 virus and the resistance/susceptibility was checked by PCR and spot test as previously

described.

To check if the colonies had the mutation affecting the sodium/glucose cotransporter, primers were

designed to amplify the specific region (Supplementary Table 7). DNAs of the 6 selected colonies

picked in 1 ml of medium and *Sal. ruber* M1 wild type (as a control of the mutation) were extracted
by boiling as previously described and PCR reactions done as described previously. PCR
conditions are described in Supplementary Table 8. PCR products were visualized under
ultraviolet light after staining with ethidium bromide (100 μg/ml). Products were sequenced with
Sanger method at Stab Vida (Stab Vida, Caparica, Portugal) and analyzed with Geneious software
6.1.8 (Biomatters, Auckland, New Zealand).

Eight strains of *Sal. ruber* (M1, M8, M31, P13, P18, SP38, SP73 and RM158) were exposed to
pseudolysogen 1R through spot test as previously described, adding spots of 3 μl of 1R at late
exponential phase and diluted 10^2, 10^4 and 10^6 times on the double layer formed by the strains.

One step growth curve

Burst size was determined after carrying out a one-step growth experiment as detailed in
Kropinski, 2018 (65) (Supplementary Table 9).

REFERENCES

1.   Shiklomanov, I. A. World water resources: a new appraisal and assessment for the 21st century (UNESCO-IHP, 1998).

2.   Mutlu, M. B. et al. Prokaryotic diversity in Tuz Lake, a hypersaline environment in Inland Turkey. FEMS Microbiol. Ecol. 65, 474–483 (2008).

3.   Ramos-Barbero, M. D. et al. Prokaryotic and viral community structure in the singular chaotropic salt lake Salar de Uyuni. Environ. Microbiol. 21, 2029–2042 (2019).

4.   Boujelben, I. et al. Virioplankton Community Structure in Tunisian Solar Salterns. Appl. Environ. Microbiol. 78, 7429–7437 (2012).

5.   Ventosa, A. *Unusual micro-organisms from unusual habitats: hypersaline environments* (Cambridge University Press, 2006).

6.   Gomariz, M. *et al.* From community approaches to single-cell genomics: the discovery of ubiquitous hyperhalophilic Bacteroidetes generalists. *ISME J.* 9, 16–31 (2015).

7.   Antón, J. *et al. Salinibacter ruber* gen. nov., sp. nov., a novel, extremely halophilic member of the Bacteria from saltern crystallizer ponds. *Int. J. Syst. Evol. Microbiol.* 52, 485–491 (2002).

8.   Viver, T. *et al.* Diversity of extremely halophilic cultivable prokaryotes in Mediterranean, Atlantic and Pacific solar salterns: Evidence that unexplored sites constitute sources of cultivable novelty. *Syst. Appl. Microbiol.* 38, 266–275 (2015).

9.   Mora-Ruiz, M. del R. *et al.* Biogeographical patterns of bacterial and archaeal communities from distant hypersaline environments. *Syst. Appl. Microbiol.* 41, 139–150 (2018).

10.   Viver, T. *et al.* Description of two cultivated and two uncultivated new *Salinibacter* species, one named following the rules of the bacteriological code: *Salinibacter grassmerensis* sp. nov.; and three named following the rules of the SeqCode: *Salinibacter pepae* sp. nov., *Salinibacter abyssi* sp. nov., and *Salinibacter pampae* sp. nov. *Syst. Appl. Microbiol.* 46, 126416 (2023).

11.   Santos, F. *et al.* Culture-Independent Approaches for Studying Viruses from Hypersaline
Environments. *Appl. Environ. Microbiol.* 78, 1635–1643 (2012).

12.   Wigington, C. H. *et al.* Re-examination of the relationship between marine virus and
microbial cell abundances. *Nat. Microbiol.* 1, 1–9 (2016).

13.   López-García, P. *et al.* Metagenome-derived virus-microbe ratios across ecosystems. *ISME*
J. 17, 1552–1563 (2023).

14.   Di Meglio, L. *et al.* Seasonal dynamics of extremely halophilic microbial communities in
three Argentinian salterns. *FEMS Microbiol. Ecol.* 92, fiw184 (2016).

15.   Javor, B. J. Planktonic standing crop and nutrients in a saltern ecosystem1. *Limnol.*
Oceanogr. 28, 153–159 (1983).

16.   Ventosa, A., Fernández, A. B., León, M. J., Sánchez-Porro, C. & Rodriguez-Valera, F. The
Santa Pola saltern as a model for studying the microbiota of hypersaline environments.
Extremophiles 18, 811–824 (2014).

17.   Harding, T. & Simpson, A. G. B. Recent Advances in Halophilic Protozoa Research. *J.*
Eukaryot. Microbiol. 65, 556–570 (2018).

18.   Santos, F., Yarza, P., Parro, V., Briones, C. & Antón, J. The metavirome of a hypersaline
environment. *Environ. Microbiol.* 12, 2965–2976 (2010).

19.   Correa, A. M. S. *et al.* Revisiting the rules of life for viruses of microorganisms. *Nat. Rev.*
Microbiol. 19, 501–513 (2021).

20.   Ptashne, M. *A genetic switch: Gene control and phage λ* (Cell Press & Blackwell Scientific
Publications, 1986).

21.   Zeng, L. *et al.* Decision making at a subcellular level determines the outcome of
bacteriophage infection. *Cell* 141, 682–691 (2010).

22.   Hay, I. D. & Lithgow, T. Filamentous phages: masters of a microbial sharing economy.
EMBO Rep. 20, e47427 (2019).

23.   Łoś, M. & Węgrzyn, G. Pseudolysogeny. *Adv. Virus Res.* 82, 339–349 (2012).

24.   Mäntynen, S., Laanto, E., Oksanen, H. M., Poranen, M. M. & Díaz-Muñoz, S. L. Black

box of phage–bacterium interactions: exploring alternative phage infection strategies. *Open Biol.*

11, 210188 (2021).

25.   Thingstad, T. F. Elements of a theory for the mechanisms controlling abundance, diversity,

and biogeochemical role of lytic bacterial viruses in aquatic systems. *Limnol. Oceanogr.* 45, 1320–

1328 (2000).

26.   Weitz, J. S., Beckett, S. J., Brum, J. R., Cael, B. B. & Dushoff, J. Lysis, lysogeny and

virus–microbe ratios. *Nature* 549, E1–E3 (2017).

27.   Villamor, J. *et al.* Characterization of ecologically diverse viruses infecting co-occurring

strains of cosmopolitan hyperhalophilic Bacteroidetes. *ISME J.* 12, 424–437 (2018).

28.   Erez, Z. *et al.* Communication between viruses guides lysis–lysogeny decisions. *Nature*

541, 488–493 (2017).

29.   Luria, S. E. & Delbrück, M. Mutations of bacteria from virus sensitivity to virus resistance.

Genetics 28, 491–511 (1943).

30.   Maynard, N. D. *et al.* A Forward-Genetic Screen and Dynamic Analysis of Lambda Phage

Host-Dependencies Reveals an Extensive Interaction Network and a New Anti-Viral Strategy.

PLOS Genet. 6, e1001017 (2010).

31.   Shi, K. *et al.* Structural basis of superinfection exclusion by bacteriophage T4 Spackle.

Commun. Biol. 3, 691 (2020).

32.   van den Berg, B. *et al.* Structural basis for host recognition and superinfection exclusion

by bacteriophage T5. *Proc. Natl. Acad. Sci.* 119, e2211672119 (2022).

33.   Labrie, S. J., Samson, J. E. & Moineau, S. Bacteriophage resistance mechanisms. *Nat. Rev.*

Microbiol. 8, 317–327 (2010).

34.   Treptow, N., Rosenfeld, R. & Yarmolinsky, M. Partition of nonreplicating DNA by the par
system of bacteriophage P1. *J. Bacteriol.* 176, 1782–1786 (1994).

35.   Kourilsky, M. Learning Through Advocacy: An Experimental Evaluation of an Adversary
Instructional Model. *J. Econ. Educ.* 3, 86–93 (1972).

36.   Kourilsky, P. Lysogenization by bacteriophage lambda. *Mol. Gen. Genet. MGG* 122, 183–
195 (1973).

37.   Cornett, J. B. Spackle and Immunity Functions of Bacteriophage T4. *J. Virol.* 13, 312–321
(1974).

38.   Shabram, P. & Aguilar-Cordova, E. Multiplicity of Infection/Multiplicity of Confusion.
Mol. Ther. 2, 420–421 (2000).

39.   Weitz, J. S. *Quantitative Viral Ecology: Dynamics of Viruses and Their Microbial Hosts*.
(Princeton University Press, 2016).

40.   Łoś, M. *et al.* Effective inhibition of lytic development of bacteriophages λ, P1 and T4 by
starvation of their host, *Escherichia coli*. *BMC Biotechnol.* 7, 13 (2007).

41.   Bryan, D., El-Shibiny, A., Hobbs, Z., Porter, J. & Kutter, E. M. Bacteriophage T4 Infection
of Stationary Phase *E. coli*: Life after Log from a Phage Perspective. *Front. Microbiol.* 7, (2016).

42.   Abedon, S. T. Lysis of lysis-inhibited bacteriophage T4-infected cells. *J. Bacteriol.* 174,
8073–8080 (1992).

43.   Dressman, H. K. & Drake, J. W. Lysis and Lysis Inhibition in Bacteriophage T4: *rV*
Mutations Reside in the Holin *t* Gene. *J. Bacteriol.* 181, 4391–4396 (1999).

44.   Million-Weaver, S. & Camps, M. Mechanisms of plasmid segregation: have multicopy
plasmids been overlooked? *Plasmid* 0, 27–36 (2014).

45.   Basso, J. T. R. *et al.* Genetically similar temperate phages form coalitions with their shared
host that lead to niche-specific fitness effects. *ISME J.* 14, 1688–1700 (2020).

46.   Thyrhaug, R., Larsen, A., Thingstad, T. & Bratbak, G. Stable coexistence in marine algal
host-virus systems. *Mar. Ecol. Prog. Ser.* 254, 27–35 (2003).

47.   Huang, C. *et al.* Isolation, characterization, and application of a novel specific *Salmonella*
bacteriophage in different food matrices. *Food Res. Int.* 111, 631–641 (2018).

48.   Sant, D. G., Woods, L. C., Barr, J. J. & McDonald, M. J. Host diversity slows bacteriophage
adaptation by selecting generalists over specialists. *Nat. Ecol. Evol.* 5, 350–359 (2021).

49.   Han, K. *et al.* Characterization and genome analysis of a novel phage Kayfunavirus TM1.
Virus Genes 59, 302–311 (2023).

50.   Tan, D. *et al.* Characterization of *Klebsiella pneumoniae* ST11 Isolates and Their
Interactions with Lytic Phages. *Viruses* 11, 1080 (2019).

51.   Li, Z. *et al.* Characterization and Application of a Lytic Phage D10 against Multidrug-
Resistant *Salmonella*. *Viruses* 13, 1626 (2021).

52.   Yang, Y. *et al.* Development of a Bacteriophage Cocktail to Constrain the Emergence of
Phage-Resistant *Pseudomonas aeruginosa*. *Front. Microbiol.* 11, (2020).

53.   Alexyuk, P. *et al.* Isolation and Characterization of Lytic Bacteriophages Active against
Clinical Strains of *E. coli* and Development of a Phage Antimicrobial Cocktail. *Viruses* 14, 2381
(2022).

54.   Los, M., Wegrzyn, G. & Neubauer, P. A role for bacteriophage T4 *rI* gene function in the
control of phage development during pseudolysogeny and in slowly growing host cells. *Res.*
Microbiol. 154, 547–552 (2003).

55.   Latino, L., Midoux, C., Vergnaud, G. & Pourcel, C. Investigation of *Pseudomonas*
aeruginosa strain PcyII-10 variants resisting infection by N4-like phage Ab09 in search for genes
involved in phage adsorption. *PLOS ONE* 14, e0215456 (2019).

56.   Ripp, S. & Miller, R. V. The role of pseudolysogeny in bacteriophage-host interactions in
a natural freshwater environment. *Microbiology* 143, 2065–2070 (1997).

57.   Bratbak, G., Egge, J. & Heldal, M. Viral mortality of the marine alga *Emiliania huxleyi*

(Haptophyceae) and termination of algal blooms. *Mar. Ecol. Prog. Ser.* 93, 39–48 (1993).

58.   Shkoporov, A. N. *et al.* Long-term persistence of crAss-like phage crAss001 is associated

with phase variation in *Bacteroides intestinalis*. *BMC Biol.* 19, 163 (2021).

59.   Rodriguez-Valera, F., Ventosa, A., Juez, G. & Imhoff, J. F. Variation of Environmental

Features and Microbial Populations with Salt Concentrations in a Multi-Pond Saltern. *Microb.*

Ecol. 11, 107–115 (1985).

60.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* 30, 2114–2120 (2014).

61.   Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its Applications

to Single-Cell Sequencing. *J. Comput. Biol.* 19, 455–477 (2012).

62.   Noble, R. T. & Fuhrman, J. A. Use of SYBR Green I for rapid epifluorescence counts of

marine viruses and bacteria. *Aquat. Microb. Ecol.* 14, 113–118 (1998).

63.   Santos, F. *et al.* Metagenomic approach to the study of halophages: the environmental

halophage 1. *Environ. Microbiol.* 9, 1711–1723 (2007).

64.   Smith, C. L., Klco, S. R. & Cantor, C. R. Pulsed-field gel electrophoresis and the

technology of large DNA molecules. *Genome Anal. Pract. Approach* 41–72 (1988).

65.   Kropinski, A. M. *Bacteriophages: Methods and Protocols, Volume 3* (Springer, 2018).

ACKNOWLEDGEMENTS

We warmly thank to Julia Esclapez and Laura Matarredona for their help with the Southern Blot. We thank Cristina López for providing the initial infection curves. This research was supported by the projects "Virhost" CIPROM/2021/006 (PROMETEO 2022, Generalitat Valenciana) and METACIRCLE PID2021-126114NB-C41 (Spanish Ministry of Science and Innovation). R.S.M. received funding for his doctoral thesis from the Spanish Ministry of Science and Innovation PRE2019-087998. R.S.M., F.S. and J.A. are members of the National Excellence Network FAGOMA (RED2022-134837-T). This project was supported, in part, by Simons Foundation Grant 722153 (to J.S.W.) and by the Chaires Blaise Pascal program of the Île-de-France Region (to J.S.W.).

AUTHOR CONTRIBUTIONS

R.S.M., J.S.W., F.S. and J.A. conceived and designed the study. R.S.M. performed the experiments and analyzed the sequences under the inputs and supervision of M.K., F.S., and J.A.. A.A. and J.S.W. designed and implemented the mathematical model. A.A. wrote the code and performed the simulations and statistical analysis. R.S.M. and A.A. drafted the original manuscript. All authors contributed to manuscript final writing and approved the final paper.

DATA AVAILABILITY

The raw files used for mutation analysis were deposited in the NCBI database with BioProject accession ID PRJNA1136859. Raw sequences of the Sanger sequencing of the mutation affecting the sodium/glucose cotransporter can be found in the *Mutations Supplementary Information*

CODE AVAILABILITY

The code used in the study is available via GitHub at:

https://github.com/aranilah/SalruberM1_EM1_recovery/tree/main

ETHICS DECLARATIONS

The authors declare no competing interests.

 FIGURES

Fig. 1: **Infection experiments show a reproducible recovery after viral lysis**. Infection experiments of *Sal. ruber* strain M1 with the EM1 virus were performed using two different MOIs. Black lines represent uninfected control and red (**a**, MOI=0.01) and purple (**b**, MOI=0.1) lines represent virus-infected cultures. Each of the individual replicates used are shown as dotted lines. The dots show the mean and the vertical bars the standard error. The experiments were performed with 3 replicates for MOI 0.01 and 10 replicates for MOI 0.1 for both the control and infected cultures. Viruses were mixed with the host at time=0 h.

Fig. 2: The EM1 virus establishes a pseudolysogenic state in *Sal. ruber* M1. a, Three randomly selected cultures of the 20 colonies of C_F, PCR positive for the EM1 virus, that we named 1R, 2R and 3R, were stained with SYBR™Gold and observed under epifluorescence microscope to test the presence of extracellular viruses. No extracellular viruses were seen in any of them. 30 fields were examined per sample. The photos shown here are a representation of each sample. Scale bar: 10 μm. **b**, A PFGE was performed (left) comparing initial cells (C_0) with final colony 1R (lines 3 and 4 of the gel, respectively). The DNA of the virus was also loaded as a control (line 5). The PFGE showed a band in 1R with the size of EM1 (35kb) which was not found in C_0. A southern blot (right) with a labeled probe targeting the genome of EM1 confirmed that this band belonged to the virus.

Fig. 3: Viral acquisition protects Sal. ruber M1 from further infection and has MOI
dependent properties. a, Growth curves (performed in triplicate) of *Sal. ruber* M1 wild type
(black dotted line) and pseudolysogen 1R (blue line) were performed and their growth dynamics
were found to have no statistical differences (*Modeling Supplementary Information: Sal. ruber M1*
wild-type and 1R (pseudolysogen) growth comparison). **b**, Susceptibility to the virus was assessed
using a spot test. *Sal. ruber* M1 wild type (left) and pseudolysogen 1R (right) were exposed to 3
µl of the EM1 virus tittered at 10^{10} PFUs/ml (left part of each plate) and diluted 10^2, 10^4 and 10^6
(left to right) times. **c**, Infection curves of *Sal. ruber* M1 and the EM1 virus were performed at
different MOI. The M1 cultures were infected at 66 hours, during the exponential growth phase.
The black dotted line represents the non-infected control and the different colors indicate the MOI,
as is indicated in the legend. The experiment results exhibit three key features (*see text*). **d**, An
adsorption assay of the EM1 virus to *Sal. ruber* M1 wild type at MOI 0.01 (red line) and 10 (yellow
line), where the number of extracellular viruses (PFU/ml) was measured over 120 minutes. The
virus stock used was also set as a decay control (grey line). All the experiments were performed
by triplicate. Error bars represent the standard error.

800
Fig. 4: Schematic of nonlinear dynamics population model a, Schematic representation of the
ODE model. R denotes the resource level and V denotes the free virus concentration. The host
population is split into three classes: sensitive (S), early pseudolysogen (P_e^k), and fully developed
pseudolysogen (P_f^k), where k is the number of viral genome copies within the host cell. **b,**
Dynamics of pseudolysogen growth and superinfection assuming primary infection has already
occurred (primary infection: $S \rightarrow P_e^1$). Early pseudolysogens get superinfected causing $P_e^k \rightarrow$
P_e^{k+1} transitions, while fully developed pseudolysogens are resistant to superinfection at the
surface level (denoted by *blocked infection* arrow). Pseudolysogen cell division for both early and
fully developed pseudolysogens randomly splits the viral genome of the parent, causing $P^k \rightarrow$
$P^{k'}$ transitions where k' is a randomly chosen positive integer that is $\leq k$ (P^0 is virus-free and is
therefore equivalent to S). Early stage pseudolysogens eventually transition into fully developed
pseudolysogens that still cell divide but are resistant to superinfection.

Fig. 5: Simulated population dynamics recapitulate experimental features through
implementation of lysis inhibition and pseudolysogenic cell division. a, Simulated host
population dynamics ($S + \sum_{i=1}^{k} P_e^{[k]} + \sum_{i=1}^{k} P_f^{[k]}$) using the ODE model proposed *(Modeling*
Supplementary Information: Main Model) to successfully recapitulate the experimental features
highlighted in Fig 3c. Subfigures b,c,d graphically show how decoupling features from the model
prevents simulations from capturing experimental features. **b,** Simulated host population dynamics
with lysis rates independent of resource levels. **c**, Simulated host population dynamics with lysis
rates independent of cellular MOI. **d**, Simulated host population dynamics without pseudolysogen
cell division.

   EXTENDED DATA

Extended Data Fig. 1 Infection curves of different *Sal. ruber* strains and its viruses. The black dotted line represents the control without viruses and the pink dotted lines represent each replicate of the infected culture. In the infected cultures, the mean is represented by a pink dot and the bars represent the standard error. These curves were made with different strains of *Sal. ruber* and different viruses (**a**, strain M1 and EM1 virus; **b**, strain M8 and CR2 virus; **c**, strain M31 and CC1 virus; **d**, strain M1 and CR41-2 virus). The control has only one replicate, while the infection has 3 replicates.

842
Extended Data Fig. 2 Pseudolysogens develop resistance to EM1 at external surface level. An
adsorption assay of the EM1 virus to *Sal. ruber* M1 wild type (black dotted line) and
pseudolysogen 1R (blue line) was carried out with a MOI of 0.01, quantifying the quantity of
extracellular viruses through PFU/ml during 120 minutes. The experiment was also carried out
with the virus alone as a decay control (grey line). The experiment was performed by triplicate.
Error bars represent the standard error.

Extended Data Fig. 3 Expected adsorption results with superinfection. The simulated
adsorption dynamics when only a single virus can infect a cell (*k=1, green)*, when eight viruses at
maximum can infect a cell *k= 8* viruses (*k=8, red*), and when viruses can continuously infect a cell
with no limit (*black*). The dynamics when *k=8* seem to match the experimental adsorption
dynamics observed in Figure 3d. In reality, the reasoning behind the limit to the number of viruses
per cell could be due to competition for receptor sites, or perhaps explicit spatial limitations.

860
Extended Data Fig. 4 The EM1 virus is maintained at the population level across time. a,
Growth curves of the different transfers of pseduolysogen 1R. The points taken for new transfers,
spot tests, PCR and qPCR are marked with a grey oval. It was transferred by triplicate. Error bars
represent the standard error of the replicates. **b**, PCR of the three replicates of the transfers of
pseudolysogen 1R shown above. The PCR was performed with primers for *Sal. ruber* M1 (upper
part) and the EM1 virus (bottom part). **c**, qPCRs were performed on each of the 4 transfers to
quantify the ratio of EM1 virus genome/chromosome of *Sal. ruber* M1 in the culture. Biological
replicates are represented with dots. The mean is represented as a horizontal line.

870
Extended Data Fig. 5 Resistance to the EM1 virus is also maintained over time in the pseudolysogens. Spot tests of the three replicates of the transfers of pseudolysogen 1R to assess the susceptibility to the EM1 virus. All the transfers of pseudolysogen 1R were plated in solid medium and exposed to spots of the EM1 virus tittered at 10^{10} PFUs/ml and diluted 10^2, 10^4 and 10^6 times. A control of *Sal. ruber* M1 wild type (upper part) was included in each transfer to test the infectivity of the virus.

878 **Extended Data Fig. 6 Pseudolysogens kill other virus-susceptible strains.** Spot test of eight
879 wild type strains of *Sal. ruber* against the pseudolysogen 1R. All the cultures were plated in solid
880 medium and exposed to spots of pseudolysogen 1R diluted 10^0, 10^2, 10^4 and 10^6 times. Spots of
881 *Sal. ruber* M1 wild type and the EM1 virus were also added as a control.