1 An effective model for community-based conservation around authorized

2 fishing settlements inside a devolved Wildlife Management Area in

3 southern Tanzania

4 Abstract

Wildlife Management Areas (WMAs) represent a relatively new institutional model for

devolved, locally-led conservation in Tanzania, in which local villages set aside part of their

land for wildlife conservation and manage that resource collectively, so that their stakeholder

communities can collectively leverage economic and social benefits from income-generating

activities like tourism. This study examines the relationship between community-defined land

use plans and *de facto* land use practices, and the influence of the latter on the relative

abundance and distribution of large wild mammals in a across the Ifakara-Lupiro-Mangula

(ILUMA) WMA, which acts as a key buffer zone between Nyerere National Park (NNP) to

the east and adjacent stakeholder villages to the north and west. All observed signs of wildlife

and human activity were recorded across 32 locations inside ILUMA and in the permanent

settlements and national park that respectively border it to the west and east. Across much of

ILUMA WMA, in areas where agreed land use plans were not adhered to, rampant cattle

herding and land clearing for agriculture were associated with reductions in wildlife richness

and biodiversity, as well as overall ecosystem integrity. Although human settlement was also

generally associated with reduced natural ecosystem integrity, some important exceptions to

this rule illustrate how sustainable livelihoods for local people that are based on well-

managed natural resource harvesting practices may actually enhance conservation

effectiveness: Three authorised human settlements within the WMA, where fishing was the

primary permitted livelihood and local communities collaborated with the WMA

management, were surrounded by pristine land cover with thriving terrestrial wildlife

populations. Correspondingly, the best conserved parts of the WMA not only included those

closest to the boundary with the national park to the east, but also these fishing villages along

the riverbank to the north, where compliance with agreed land use plans was most rigorous.

Overall, this study documents a useful example of how a devolved conservation area may

conditionally host resident local communities undertaking selective natural resource

extraction activities and collaborate with them to achieve effective *de facto* conservation

practices.

37 1 | INTRODUCTION.

Locally devolved community-based conservation (CBC) is considered a promising

alternative to traditional fortress-style conservation, many different forms of which have been

developed in dozens of countries around the world, with varying degrees of success and/or

failure. Two outcomes that are considered successes for CBC are the conservation of natural

ecosystems and the many species that live in them, and participation by the local

communities, not only in the governance and management of the area, but also in the derived

economic and social benefits (Songorwa, 1999). By providing tangible economic benefits to

local communities, through a participatory approach to land use planning and management

(Gibson and Marks, 1995). The overall idea of CBC is that protected landscapes and natural

resources are managed in a devolved manner by empowered local communities, to secure

their future on a sustainable basis (Gibson and Marks, 1995). The aim of CBC is to shift the

focus of conservation from exclusively centralized, state-controlled protection models, like

national parks and game reserves, to management through bottom-up institutional structures

that ensure community participation and inclusion (Goldman, 2003). In principle, local

communities can benefit from CBC via consumptive tourism, such as regulated trophy

hunting (Lindsey et al., 2006) and non-consumptive eco-tourism (Kiss, 2004), both of which

aim at strengthening conservation efforts by keeping financial rewards within the local

economy, including sustainable livelihoods for community members.

Having said that, many CBC initiatives have failed achieve these objectives, and this has

often been attributed to a lack of business, tourism and marketing skills among the local

stakeholder community (Spenceley, 2012). Furthermore, efforts to develop CBC practices

across Africa during the 1990s have been criticised for sidelining local communities from the

processes of developing management plans, while nevertheless including the community in

implementing externally-imposed conservation policies (Agrawal and Ribot, 1999, Goldman,

2003). In essence, while relevant communities were, in theory, included in wildlife

conservation policies, meaningful devolution of power was not implemented in practice,

leading to a lack of governance stakeholdership and management authority at a local level.

This prevented relevant communities from playing an active role in conservation or accrue

the derived socio-economic benefits that are so important to sustain enthusiasm for such

initiatives (Agrawal and Ribot, 1999).

It is therefore unsurprising that various CBC initiatives across Africa have yielded a mixed

picture of success and failure (Salerno *et al.*, 2016). With pressure on land use only expected

to increase in the coming decades, with a predicted 9.7 billion global citizens by 2050

(Population Division of the United Nations Department of Economic and Social Affairs,

2022). Africa in particular is expected to see declines in wild mammal habitat of up to 25%

by 2050, the primary driver of this being land-use change (Baisero et al., 2020). There is

therefore an urgent need to establish more effective mechanisms for sustained CBC that

benefit both people and wildlife as scalable models of best practice.

The establishment of Wildlife Management Areas (WMAs), as an institutional mechanism for

CBC in Tanzania, is correspondingly intended to satisfactorily devolve power and rectify

these management and governance issues. Although mentioned in the Wildlife Policy of

1998, WMAs were only established formally through the *Establishment and Management of*

Wildlife Management Areas and Benefit Sharing Government Notice in The Wildlife

Conservation Act of 2005, which was then revised in 2022 (United Republic of Tanzania

Ministry for Natural Resources and Tourism, 2022). It is the intention of this institutional

basis for CBC is to "empower local communities to be more involved and to hold more

authority over the management of wildlife to promote sustainable biodiversity conservation

and rural economic development" (USAID, 2018). In accordance with this act, WMAs

should establish administrative mechanisms whereby wildlife conservation is promoted,

economic development of the stakeholder villages is enhanced, rural poverty is reduced, and

the distribution of benefits and costs is equitable (United Republic of Tanzania Ministry for

Natural Resources and Tourism, 2022).

Each WMA has its own individual set of by-laws and is independently managed. Once

formally established and authorized, the community-based management body of the WMA

has the right to negotiate and sign agreements with customers and investors, under the

condition that representatives of the district council are present and involved in the

negotiation, with the aim of advancing both community development and wildlife

conservation objectives. The WMA is monitored for unauthorised human activities by

Village Game Scouts (VGS), who are appointed by the WMA management as its equivalent

of wildlife rangers. VGS are also responsible for both protecting the natural resources of the

area and also protecting people's lives and property from wildlife. The VGS and WMA

management also collaborate with other legal bodies, such as the Tanzanian National Parks

Authority (TANAPA), The Wildlife Division, the police and the district government in

operations to prevent various forms of encroachment upon protected land. Currently 19

WMAs are operational in Tanzania, accounting for 7% of the country's total land area, and a

further 19 were in the planning stages as of 2018 (Lee and Bond, 2018).

In May 2015, the Ifakara-Lupiro-Mangula (ILUMA) WMA was issued user rights and

declared the 19th WMA in Tanzania (United Republic of Tanzania Ministry of Natural

Resources and Tourism, 2015). Fourteen villages from both the Ulanga and Kilombero

districts, with a total of more than 80,000 residents, are included as stakeholder communities

in this WMA (United Republic of Tanzania Ministry of Natural Resources and Tourism,

2015). Three authorised local cooperative fishing communities are established and located

inside the wetland zone of ILUMA; *Mikeregembe*, *Mdalangwila* and *Funga*. In the

management plan these are referred to as *Beach Management Units* (BMUs). Monitored

fishing is permitted in these areas under licence and the ILUMA by-laws include regulations

on how these settlements are maintained. For example, construction of a permanent residence

in these locations is not permitted, so all structures must be temporary in nature. Specifically,

this means that they are built using only natural materials like mud, sticks and thatch, rather

than bricks and iron sheets. The residents of these areas are allowed to collect dry wood for

such purposes as cooking and drying fish but are not permitted to harvest any live trees.

Charcoal burning is also not permitted.

At face value, these guidelines and management mechanisms appear ideal for protecting both

the livelihoods of the local people from the 14 villages and conserving wildlife. However,

anecdotal evidence and informal experiences of some of the authors circa 2018 suggested that

the WMA was struggling, and that the land use plans formally agreed upon were not being

adhered to. The aim of this study was therefore to evaluate how well the formally agreed land

use plans compared with the *de facto* land use practices ongoing across the area of ILUMA

WMA, assess how well the WMA was functioning from a wildlife conservation viewpoint,

and glean useful insights to inform improved WMA management going forward.

128   2 | MATERIALS AND METHODS.

129   2.1 | Study Area

The study area is located both inside ILUMA Wildlife Management Area (ILUMA WMA),

and outside it in the villages immediately surrounding its borders to the west, and in NNP to

the east. ILUMA WMA is located in the Kilombero Valley [08°40'S 36°10'E], spanning both

Kilombero and Ulanga districts in the Morogoro region of Southern Tanzania (Figure 1).

Covering an area of over 7, 000km^2, the Kilombero Valley is the largest lowland freshwater

wetland in East Africa and an established RAMSAR site. ILUMA WMA is nested within this

valley, covering an area of 509km^2 immediately to the west of NNP (United Republic of

Tanzania Ministry for Natural Resources and Tourism, 2023). Located in south-eastern

Tanzania, NNP is Africa's largest national park, encompassing 30,893 km^2 ("Nyerere

National Park," 2023). It was designated national park status in 2019, having previously

formed part of the Selous game reserve. ILUMA's northern boundary is demarcated by the

Kilombero river, while areas to the south, west and north contain expanses of agricultural

land and villages.

143   INSERT FIGURE 1 HERE

Figure 1. Map displaying the survey area in a national and local context in southern

Tanzania. The insert at the top left is a map of Tanzania, while the enlarged area shows the

part of the Ifakara-Lupiro-Mangula Wildlife Management Area (ILUMA WMA) to the south

of the Kilombero River where most of the surveys were carried out in the context of the

nearby Nyerere National Park (NNP) to the east, the extensive wetlands of the Kilombero

Valley inland delta upstream to the west and Ifakara town to the north-west.

ILUMA acts as a buffer zone for NNP and includes zones for tourist hunting and local

hunting, in addition to a wetland conservation zone. Establishment began in 2011 and user

rights were declared in 2015, making ILUMA the 19[th] WMA in Tanzania (United Republic of

Tanzania Ministry of Natural Resources and Tourism, 2015). Land from 14 villages is

encompassed by ILUMA (Daconto et al., 2018), and its cover consists of miombo woodlands,

dense groundwater forest, riparian forest, open grasslands and wetlands. The climate of the

region is typical of moist tropical savannahs with distinct seasons. The main rainy season of

reasonably consistent heavy precipitation occurs between March and May each year, with the

dry season from June through to October generally seeing little to no rainfall, and then

followed by a shorter, less predictable season of generally more sporadic rainfall between

November and February. Annual average rainfall in the Kilombero valley typically varies

between 1,200 and 1,4000mm (Mombo et al., 2011) and temperatures are highest from

December to March.

The survey area consisted of 32 distinct locations, referred to as *camps*, each assigned an

identity numbers from 1 to 32 (Figure 2 and supplementary file 1). These camp survey

locations were distributed throughout the study area: 22 inside the borders of the ILUMA

conservation area, 6 outside it in the domesticated areas close to nearby villages and 4 camps

inside adjacent parts of NNP. It should be noted that this study was nested within a larger

project related to malaria vector mosquito ecology (Kavishe *et al.,* 2024, Walsh, 2024), the

objectives of which were the primary determinant of survey location selection. The logistical

requirements of this study therefore had to be adapted to fit within the data collection

requirements of that larger multi-disciplinary project. The most important requirement

considered when choosing survey sites was year-round availability of surface water, where

malaria vector mosquitoes could breed and the survey team could obtain sufficient water for

drinking, cooking and bathing (Kavishe *et al*., forthcoming and Walsh *et al.,* forthcoming).

176   INSERT FIGURE 2 HERE

Figure 2. Map displaying the distribution of suitable camping locations used as the

sampling frame for all the surveys described in this report. Each of the 32 camps detailed

in supplementary file 1 are illustrated in the geographic context of the boundaries of the

Ifakara-Lupiro-Mangula Wildlife Management Area (ILUMA WMA) and Nyerere National

Park (NNP).

2.2 | Study design and sampling frame

As explained in detail elsewhere (Duggan, 2024) this study was carried out using a repeated

rolling cross-sectional design with three rounds of surveys over the course of one calendar

year, specifically 2022. Round 1 ran from 21st January to 16th March, while round 2 ran from

22nd March to 25th May, and round 3 then ran from 25th August to 29th November. Rounds

one and two occurred during the wet season and round 3 during the dry season. Each round

encompassed most of these 32 camps, but not all of them (Supplementary file 1) for various

practical reasons, including physical inaccessibility during the rains or lack of water in the

dry season. These camps were assigned to one of five circuits, with each one surveyed over a

period of approximately two weeks, staying at each camp for two nights before departing for

the next camp the following morning. The radial surveys of various detected human,

livestock and wild mammal activities around surface water bodies within a 2km radius of

each camp were carried out on the mornings after arriving at each camp (Duggan, 2024,

Duggan et al. (2024), forthcoming, Duggan et al., forthcoming).

2.3 | Formal quantitative radial surveys of human, livestock and wild mammal activities

around surface water bodies

As explained in detail elsewhere (Duggan, 2024, Duggan et al. (2024), Duggan et al.,

forthcoming), radial surveys for various signs of activity by humans, livestock or wild

animals (Supplementary file 3) were carried out along the fringes of water bodies within a

2km radius of each camp, These water bodies were first surveyed for mosquito larvae by the

accompanying investigators focusing on malaria vector ecology (Walsh, 2024), so the exact

direction taken and waterbodies surveyed was primarily determined by the requirements of

those entomological surveys. The waterbodies surveyed included seasonal streambeds and

the pools of water within them, puddles and pools outside of streambeds, waterholes, pools of

water in rice fields and rivers and any other surface water body that might attract mosquitoes,

people or animals, even if they were clearly ephemeral in nature. All direct visual

observations and indirect signs of activity by humans, livestock or wild mammals

(Supplementary file 3) that were encountered along the routes taken between and around the

fringes of surveyed water bodies were recorded using the standardized tools provided in

supplementary files 4, 5, 6 and 7). The fully explicit data collected through all these formal

quantitative observations are presented in supplementary file 8.

2.4 | Numerical estimation of conventional indices of wild animal community or whole

ecosystem integrity based on formal quantitative data from the radial activity surveys

Estimates for each of the following indices of the integrity of wild animal communities or the

whole ecosystem at each camp were calculated as follows based on the data obtained through

radial activity surveys (Section 2.3). All the following analyses were carried out using the $R^{®}$

open-source software package through the *RStudio*$^{®}$ version 2023.03.0+386 environment.

While only data relating to activities by wild mammals was used to calculate scores for the

first three indices of wild animal community integrity for each camp, the fourth and final

index described below, intended as an indicator of whole ecosystem integrity, included all of

the variables available in supplementary file 8. Assigned scores for the each of the four

following indices at each camp are detailed in supplementary file 2.

Species Richness Index (SRI) for each camp was calculated as the absolute number of species

in a given area (Whittaker et al., 2001), based on wildlife activity data only, using the *plyr*

package and *richness* command. SRI was calculated as: $\sum_{i=1}^{x} n$, where n = the number of

species detected (Moore, 2013), with higher values indicating higher biodiversity.

Simpson's Index of Diversity (SID) were calculated as 1- D, where $D = \sum_{i=0}^{x} iPi^{\ 2}$, where P_i (P_i

$= \frac{xi}{N}$) is the proportion of observations accounted for by each species i, x_i is the total number

of each species recorded and N is the total number of individuals recorded.

Objective Wild Animal Community Integrity Index (OWACII) was estimated by using the

factoextra package to carry out a principal component analysis (PCA) of the values for all the

various types of detections of all the wildlife species for each camp in the processed datasets

provided in supplementary file 8. OWACII represents a synthetic statistical summary

parameter, based on this data reduction analysis of all the variables that relate to recorded

activities of wild mammals. To make these values easier to interpret, they were normalized by

scaling as their Z-score means.

Objective Natural Ecosystem Integrity Index (ONEII) was also estimated by PCA using the

factoextra package (James and McCulloch, 1990; Ortega *et al.,* 2004; Castela *et al.,* 2008;

Reza and Abdullah, 2011; Janžekovič and Novak, 2012; Caniani *et al.,* 2016). This synthetic

statistical summary parameter represents more comprehensive index of overall ecosystem

integrity that OWACII because it is not only based on all the detections of wildlife, livestock

and humans but also the recorded summary indicators of land cover at each camp. This index

therefore combines all the recorded indicators of wildness with those of encroachment to

provide a composite score for comparing the ecological status of different camps. Again, to

make these values easier to interpret, they were scaled as their Z-score means.

2.5 | Cluster analysis to classify survey locations into statistically distinctive categories

with common characteristics

Cluster analysis using the k-medoids method was undertaken using the *cluster, dplyr* and

Rtsne packages, plus *ggplot2* and *scales* for graphing. The silhouette method was used to

determine the optimum number of clusters using the *cluster* package. The output suggested

two clusters as the optimum, followed by four and then eight (Figure 3). Two clusters were

not just the best fit to the data (Figure 3), but also the most practically useful, because the two

clusters could be readily and intuitively interpreted as representing degraded versus pristine

classes (Figure 4). Plots of Gower dissimilarities were created using *Rtsne* and *ggplot2*

packages to reveal clear patterns of clustering.

258   INSERT FIGURE 3 HERE

Figure 3. Plot of silhouette widths as output from k-medoids cluster analysis described

above. Higher values for width suggest a more accurate number of clusters.

2.6 | Consensus perception-based estimation of a *Subjective Natural Ecosystem Integrity*

***Index* (SNEII)**

Following the three rounds of data collection and prior to any analyses, a *subjective natural*

ecosystem integrity index (SNEII) was devised, as an alternative means of assessing the

ecological intactness of the area immediately around each camp and the level of degradation,

if any, exhibited over the course of the study. This index was scaled as a percentage score,

with a score of zero representing a fully domesticated landscape with well-established human

settlements and comprehensive land conversion for agricultural purposes, whereas a score of

100% represents a fully pristine, natural ecosystem, with no evidence of human exploitation

of any kind.

As described in detail elsewhere (Duggan, 2024, Duggan et al. (2024), Duggan et al.,

forthcoming) individual scores were carefully assigned based on a consensus reached through

discussions among the three investigators on the field team (LD, KW and LT), based on their

subjective recollections. The primary and most important criterion for assigning an ecosystem

integrity score was the subjective impressions of the intensity of land degradation caused by

the conversion of land for agriculture, deforestation for the purposes of charcoal production,

timber harvesting or human settlement and for livestock herding. The intensity of

degradation of the land surrounding a 2km radius of the camp was determined by estimating

the proportion of the natural ecosystem remaining more-or-less intact, based on informal

personal observations that were made while present in the camp and during camp radial

surveys. This subjectively estimated proportion of intact land was then used as a reference

point for initiating the discussions that concluded with the assignment of the score. Camps

that evidently belonged on the extreme ends of the scale were assigned first, to provide clear

reference points, before those that were less clear and required a more in depth-discussion.

The investigators recollections regarding the abundance of wildlife and humans as indicators

of ecosystem integrity featured less in conversation when assigning scores but were

nevertheless useful as secondary factors to be considered following the discussion of the

intensity of land degradation.

These SNEII scores were initially assigned, then discussed in depth, refined and finalized

based on consensus (Supplementary file 2) in December 2022. Note that this consensus-based

score assignation was completed before the statistical analyses of the numerical summaries of

formal, quantitative survey data described above were initiated, to avoid biasing the former

towards the latter.

2.7 | General linear modelling (GLM) analyses

In order to determine which human activities had the most severe impact on natural

ecosystems, a series of separate general linear modelling (GLM) analyses were undertaken, in

which each of the individual indices described above were treated as an alternative dependent

variable, thus acting as one of several potential indicators of natural ecosystem integrity. One

exception is that GLM analysis was not carried out on ONEII as variables relating to human

activity were used in creating this index, so testing for their association with this derived

index would be confounded by circular logic. Each human activity was firstly scaled using Z-

score means, using the same calculation method described above. GLM analysis was then

carried out in *RStudio®* using the *lme4* package.

A series of univariate GLMs were fitted defining the index as the dependent variable and one

given human activity as the sole independent variable. Multiple distributions and link

functions were specified until the best fit model with the lowest AIC score was identified.

The selected best-fit distribution for SRI was Gaussian family with an inverse link. The

selected best-fit distribution for SID was Gamma family with a log link function. The

Gaussian family without any link function was chosen as the best fit for both SNEII and

OWACII. Multiple univariate GLM analyses were run using each human activity as the

independent variable. Then multivariate analyses were then carried out, using a forward

stepwise selection procedure to build the most informative and parsimonious explanatory

model possible, beginning with the human activity variable with the lowest P value in

univariate analysis, which was livestock herding in all cases. Human activities were added

and removed from the model based on objective criteria specified *a priori*, until the best fit

model based on AIC score was produced: each retained variable was required to at least

approach significance in the model it was included in and significantly improve the goodness

of fit relative to the otherwise equivalent model from which it was excluded.

321   3 | RESULTS

322   3.1 | Geographic variations in wildlife activity and ecosystem integrity across the study
area

Overall, the study area exhibited a consistent gradient of landscape characteristics, ranging

from fully domesticated lands with minimal wild fauna in and around human settlements in

the west and south, all the way through to completely intact natural ecosystems with their full

complement of wildlife to the north and east (Figure 4 and 5). The overall pattern was readily

explained in terms of the geographic distribution of various human activities, even including

the establishment of substantial human settlements, which extended deep into the ILUMA

conservation area from the villages immediately outside it to the east and south

(Supplementary file 9). However, much of the northeast of the WMA was well conserved and

even surpassed the rigorously protected survey locations inside NNP to the east in terms of

mammalian species diversity. The transition zone between these two extremes inside the

WMA exhibited a mosaic of mixed land cover, where wildlife, humans and livestock

intermingled.

336   INSERT FIGURE 4 HERE

337   **Figure 4:** Maps illustrating the distribution of scores for three different indicators of the

338   integrity of the wild animal community across the study area, estimated based on formal

339   quantitative surveys (Sections 2.3 and 2.4) at each of the camps (Section 2.2 and

340 supplementary files 1 and 2): **A**; Species Richness Index (SRI), **B**; Simpson's Index of

341 Diversity (SID), and **C**; Objective Wild Animal Community Integrity Index (OWACII).

342   INSERT FIGURE 5 HERE

Figure 5: Maps illustrating the distribution of scores for three different indicators of the

integrity of the wild animal community across the study area, respectively estimated based on

either formal quantitative surveys (Sections 2.3 and 2.4) or subjective appraisal based on

consensus investigator perspective (Section 2.5) at each of the camps (Section 2.2 and

supplementary files 1 and 2): **A**; Objective Natural Ecosystem Integrity Index (ONEII),

B; Subjective Natural Ecosystem Integrity Index (SNEII).

The number of wild mammalian species recorded at each camp varied considerably, ranging

from 2 (camp number 20 – *Makingi*) to 22 (camp number 12 – *Bwawa la Maya* and camp

number 15 – *Mdalangwila*) (Supplementary file 2). Interestingly, all of the five highest

ranked camps for SRI occurred in the WMA rather than in the adjacent national park (Figure

4A and Supplementary file 2). This is likely due to the natural cover of the WMA consisting

largely of miombo woodland and groundwater forest habitats which appears to host to a

larger diversity of species than the acacia savanna habitat that dominates most of the camps

sampled in NNP.

The range of SID values calculated for each of these camps for this index varied widely

(Figure 4B ad supplementary file 2). The most diverse camp *Bwawa la Nyati* had a score of

0.96, while the least diverse camp, *Mavimba Porini* scored 0.16. Exactly half of the camps

have a SID score >0.90, which confirms that the better conserved areas of the WMA are

highly species diverse. Of the top five camps all are located inside ILUMA in locations with

Miombo woodland land cover. While the camps located inside NNP did yield high SID

scores (all > 0.90), none of them were ranked in the top 5. This observation illustrates how

even a degraded miombo woodland ecosystem can support a greater diversity of species than

a well-protected, non-encroached acacia savanna.

However, the top ranked camps in terms of OWACII were the four NNP camps (Figure 4C

and supplementary file 2). Meat poaching was only ever recorded at one of these locations

inside NNP and neither cattle herding nor any of the other observed human activities were

recorded at any of them (Supplementary file 9). The five worst-ranked camps in terms of

OWACII all contained large areas of human settlement where extensive human activities

were ongoing. Similarly, the top ten scoring camps in terms of ONEII were all located either

inside NNP (all ranked in the top 4) or in the northeastern section of ILUMA, very close to

the border of the national park, while those with the lowest scores were all either in the

neighbouring villages to the west or around heavily settled areas inside the eastern and

southern boundary of the WMA. Overall, the SNEII estimates were closely correlated with

both of these objective alternatives, ONEII in particular (Duggan, 2024, Duggan et al. (A),

forthcoming, Duggan et al. (B), forthcoming), so the cartographic picture obtained with this

subjective approach was very similar (Figure 5), even though it was based on consensus

investigator perceptions rather than statistical synthesis of formal quantitative activity

surveys.

3.2 | Influences of specific human activities on wildlife and ecosystem integrity

For all three of the objective indices of wild animal community diversity (SRI, SID) and

integrity (OWACII), single independent variable GLM analyses indicated that a number of

human activities were associated with variation of the index in question, but only a small

subset proved significant and were retained in the subsequent final multiple independent

explanatory variable analysis (Supplementary file 10). Livestock herding was the only human

activity to significantly impact species richness in multivariate analyses of SR. In

multivariate analysis of SID, however, livestock herding and unauthorized fishing were the

only two significant human activities. Both of these variables were associated with lower SID

scores, but the effect size for the former was more than twice the latter. Livestock herding and

meat poaching both appeared associated with lower ONACII scores in multivariate analysis,

each having an apparently a similar magnitude of impact.

While similar GLM analyses could not be applied to assess the influence of these activities

upon ONEII, because this dependent variable was estimated using these same activities that

would also be treated as the independent variables (Section 2.6), multiple independent

explanatory variables analysis revealed that six human activities were predictive of the SNEII

(Table 1). It therefore appears that the perception based SNEII represents a far more sensitive,

statistically powerful dependent variable with which to elucidate the associations of

ecosystem conservation successes and failures with various human activities than any of the

more objective alternatives based on formal quantitative surveys of wild animal activity

(Duggan, 2024, Duggan et al. (B), forthcoming). Livestock herding, charcoal burning, timber

harvesting, human settlement and both rice and other tillage agriculture all proved significant

predictors of SNEII in the multivariate analysis presented in table 1, so these relationships are

represented cartographically in supplementary file 11.

Table 1. Results of single and multiple explanatory variables generalized linear modelling

analyses of scaled human activity indicators as determinants of the estimated Subjective

Natural Ecosystem Integrity Index (SNEII) for each camp surveyed, with statistically

significant associations highlighted in bold. The best fit models identified for this SNEII

outcome all assumed a Gaussian distribution and identity link function for the dependent

variable.

Activity	Single Independent Variable		Multiple Independent Variables	

	β ± SEM	T	P	β ± SEM	t	P
Livestock Herding	**-29.3 ± 3.1**	**-9.51**	**<<0.0001**	**-17.0 ± 2.9**	**-5.85**	**<<0.0001[a]**
Charcoal Burning	**-14.5 ± 5.6**	**-2.59**	**0.0146**	**6.8 ± 2.7**	**3.01**	**0.0058[a]**
Timber Harvesting	-7.7 ± 6.0	-1.27	0.213	**4.4 ± 2.0**	**2.25**	**0.0333[a]**
Fishing	1.1 ± 6.2	0.18	0.862	3.6 ± 2.4	1.50	0.1454[b]
Hunting	-4.9 ± 6.1	-0.80	0.429	-0.88 ± 2.2	-0.40	0.6917[b]
Human Settlement	**-23.3 ± 4.5**	**-5.19**	**<<0.0001**	**-6.5 ± 2.7**	**-2.38**	**0.0256[a]**
Well	**-21.4 ± 4.8**	**-4.49**	**0.0001**	3.3 ± 3.0	1.10	0.2814[b]
Meat Poaching	-8.2 ± 6.0	-1.37	0.180	-1.2 ± 2.0	-0.62	0.5388[b]
Human Presence	**-21.6 ± 4.7**	**-4.6**	**0.0001**	5.4 ± 3.2	1.70	0.1011[b]
Rice Farming	**-27.9 ± 3.5**	**-8.00**	**<<0.0001**	**-9.4 ± 2.9**	**-3.25**	**0.0033[a]**
Other Tillage Farming	**-29.4 ± 3.0**	**-9.69**	**<<0.0001**	**-9.7 ± 3.7**	**-2.59**	**0.0157[a]**

SEM: Standard error of the mean for the β coefficient estimate.

[a] As estimated from the final, most parsimonious best-fit GLM, in which this was included on

the basis of either being a significant (P≤0.05) determinant of SNEII or approached

significance (P≤0.10) as a determinant of SNEII and significantly (P≤0.05) improved the

goodness of fit based on the estimated Akaike Information Criterion (AIC).

[b] As estimated from the last GLM in which this variable was still included before being

removed on the basis that it was not a significant determinant of the SNEII (P>0.05) or

approached significance (P≤0.10) but did not significantly improve the goodness of fit based

on the estimated AIC (P>0.05).

While charcoal burning and timber harvesting led to an increased SNEII score in multiple

independent variables, this is most likely due to reverse causality: These activities obviously

do not improve ecosystem integrity but rather tended to take place in areas inside the WMA

with moderate to high SNEII scores, where the prerequisite natural resources remained

available (Supplementary file 11, figures S11.2 and S11.3).

Interestingly, cattle grazing was recorded just as frequently at the degraded camps inside

ILUMA as at those in fully domesticated areas outside the WMA (Supplementary file 11,

figure S11.1), where this activity is not restricted but must be traded off against agricultural

land use options during the growing season for obvious reasons. It is also notable that cattle

herding was encountered remarkably frequently at some relatively pristine locations with

moderate to high SNEII scores inside the WMA (Supplementary file 11, figures S11.1),

suggesting that grazing *per se* may not, in itself, have as great an impact as that suggested by

table 1. Having said that, one evident trend was that livestock herding had a consistently

significant negative impact on all the estimated indices of wild animal community

(Supplementary files 10 and 11) and overall ecosystem integrity (Table 1) and was the single

most quantitatively important variable in all four multivariate models presented in table 1.

Furthermore, no other human activity was consistently associated with all of these indices in

multivariate analyses (Table 1, supplementary files 10 and 11).

In contrast, human settlement appeared the least influential of the four activities found to be

associated with lower SNEII scores in the multivariate analysis detailed in table 1: Once the

effects of the following other activities was accounted for, the effect size for human

settlement appeared similar to or less than that for farming of rice or other tillage crops and

far less than that for cattle herding (Table 1). Furthermore, it was not significantly associated

with reductions of any of the indicators of a healthy wild animal community detailed in

supplementary file 11. One obvious reason for this surprisingly modest and inconsistent

relationship between ecosystem integrity and human settlements was that the 3 northernmost

camps inside the WMA, namely camps 14, 25 and 26 along the banks of the Kilombero River

counterintuitively had both substantial human settlements and remarkably intact natural land

cover (Figure 6).

450   INSERT FIGURE 6 HERE

Figure 6: Map of ILUMA WMA showing each camp location's Subjective Natural

Ecosystem Integrity Index (SNEII) score, ranging from the lowest (red) to the highest (green)

scores, and layered over the extent of human settlement estimated for each camp, because the

latter was identified as a significant predictor of the former in table 1.

3.3 | Cluster analysis: General rules and informative exceptions

Experience-informed interpretation of a cluster analysis of survey location characteristics

revealed that these three planned, carefully managed human settlements within the WMA

actually appeared beneficial in conservation terms. Cluster analysis split the 32 camp

locations into two equally sized, clearly distinguished groups composed of 16 camps in each

(Supplementary file 2), which were clearly and intuitively related to their geographic

distribution (Figure 7) and level of environmental conservation or degradation (Figures 4, 5,

6, and 8 plus supplementary files 2, 9, 10, 11 and 12).

464   INSERT FIGURE 7 HERE

Figure 7. Map of ILUMA WMA displaying the 32 camp locations sampled and the cluster

they were assigned to by k-medoids analysis as described in section 2.5. Shape represents

location, either inside ILUMA WMA, outside in the villages along its western boundary or

inside NNP to the east, while colour represents assignment to either the *pristine* (green) or

degraded (red) cluster, the interpretations of which based on figures 4, 5, 6 and 7, as well as

supplementary files 2, 9, 10, 11 and 12).

471   INSERT FIGURE 8 HERE

472 **Figure 8.** Dot and violin plots of the estimated values for all of the various indices of wild

473   animal community and overall ecosystem integrity (Supplementary file 2) for each camp

474   classified by location and cluster assignment as illustrated in figure 4: **A**; Species richness

475   index (SRI), **B**; Simpson's index of biodiversity (SID), **C**; Objective wild animal community

476   integrity index (OWACII), **D**; Objective natural ecosystem integrity index (ONEII) and **E**;

477   Subjective natural ecosystem integrity index (SNEII).

The first cluster contained all of the village-based camps outside the western border of

ILUMA that were fully domesticated, as well as highly encroached camps inside ILUMA to

the west and to the south with similarly *degraded* land cover. In stark contrast, all the camps

statistically attributed to the second cluster were better conserved, so this cluster was

interpreted as reflecting a relatively intact natural landscape with natural vegetation cover and

wild fauna. These *pristine* camps occur where ILUMA borders NNP on the eastern side of

ILUMA. where there is less encroachment by humans and livestock, and to the north where

the Kilombero river acts as a physical boundary against encroachment. This second cluster

also included the NNP camps to the east.

Generally speaking, the geographic patterns identified by cluster analysis were clear and

consistent with those apparent from maps of the various estimated indices of the integrity of

the wild animal community or the overall ecosystem (Figures 4, 5, 6 and 7, as well as

supplementary files 2, 9, 10, 11 and 12). There were no pristine camps outside the western

boundary of ILUMA and there are no degraded camps inside NNP. ILUMA contains a mix of

both degraded and pristine, but with more degraded camps located to the west and south of

the area and more pristine camps located to the east and north. Broadly speaking, this overall

pattern mirrored the geographic distribution of all the various human activities observed, notably the

establishment of settlements that were typically surrounded by a range of other unauthorized activities

(Figure 9A).

498   INSERT FIGURE 9 HERE

Figure 9: Photographic illustrations of how human settlements inside the ILUMA WMA

were usually observed to be surrounded by comprehensive degradation of the natural

ecosystem (Panel **A**) but were also noted to have negligible impact upon the surrounding

natural terrestrial ecosystem in the case of some authorized fishing camps established along

the Kilombero River (Panel B). Panel **A** depicts a highly encroached area visited during a

radial survey around camp 4, with obvious evidence of land conversion from forest to tillage

agriculture and human settlement. The image in panel **B** was taken at camp 14, depicting how

this authorised fishing settlement on the banks of the Kilombero river had negligible impact

upon the groundwater forest immediately behind it. Note the contrast in level of land

clearance between these two settlements.

However, consistent with figure 6 some surveyed locations within the ILUMA conservation

area that had authorised fishing camps (camps 14. 15 and 26) as part of the agreed

management plan for the WMA were assigned to the pristine cluster by the objective

statistical cluster analysis (Figure 7). These three authorized fishing villages established

within the WMA conservation area therefore represented informative exceptions to the

general rule that human settlements, and all the various human activities that are otherwise

closely associated with them, appear inevitably associated with habitat degradation (Figures 6

and 9A). These fishing camps were well managed settlements that were confined to a very

small geographical area, with an authorised and carefully regulated sustainable livelihood

(fishing) that has negligible impact on the adjacent terrestrial habitat (Figures 6 and 9B).

Consequently, the surrounding forest and floodplain were observed to be essentially intact

with no sprawl of dwellings beyond the designated settlement limits, and experienced

minimal encroachment due to being monitored and patrolled frequently (Figures 6 and 9B).

4 | DISCUSSION.

Data detected, gathered, and analysed throughout the course of this study indicate that there

was both adherence to, and at times strong deviation from, the land use plans that had been

negotiated and agreed with the stakeholder communities for ILUMA WMA. Unauthorised

human activities were recorded most frequently close to the western and southern boundaries

of ILUMA, where the WMA conservation area is bordered by formally established villages.

As a general trend, the southwest experienced the highest levels of unauthorised human

activity, which lessened in frequency approaching the northeast of the WMA and the eastern

boundary with NNP.

Unauthorised human settlement marks a clear breech of the agreed land use plans, and the

extractive and ancillary activities that are associated with human settlement such as land

clearing or deforestation for building materials and agriculture, all alter the landscape and

ecosystem functioning. Most unauthorised human settlements within ILUMA's boundaries

were clearly associated with high levels of land clearing and extractive activities. On the

African continent, the area south of the equator accounts for 13% of the land globally that is

facing degradation due to conversion of natural lands for agriculture or encroachment into

protected areas, a phenomenon which is increasing in both extent and severity across the

earth (Bai et al., 2008). This degradation of natural ecosystems on a global scale is a crucial

factor in why it is imperative for small-scale community-based conservation programmes,

such as the one in place here, to be well maintained and in doing so could prove vital in

preventing land degradation.

Rice farming took place on a much large scale throughout ILUMA. Rice agriculture is

primarily a cash crop in the Kilombero valley, grown largely for sale rather than subsistence

(Nkuba et al., 2016), so it is correspondingly grown wherever suitable soil allows. The

practice of rice farming in the Kilombero has long been established due to the largely flat

topography of this low-lying valley that seasonally floods and fertile *fluvisol* soil type that

can retain water for long periods. Due to the nature and practice of rice paddy growth, and the

large amount of water it requires, in many areas where rice farming was recorded in ILUMA

degradation of a waterbody nearby was evident. The majority of rice cultivation in ILUMA

involved some level of water course diversion for irrigation, whether from ephemeral

waterholes in the dry season or streams. The diversion of water from its original path caused

waterholes to dry out quicker and provided less resources for the wildlife that depend on

them. A study on the Kilombero wetlands, part of which was included in this study area,

found that the flow of many rivers has been reduced and that a number of swamp areas have

been reduced in size or dried up completely as a result of intensive agriculture and

deforestation in the Kilombero wetlands (Mombo et al., 2011).

Overall, while livestock herding has appeared to be the most statistically obvious factor

directly associated with to species richness, diversity, and ecological integrity in ILUMA

WMA over the course of this study, it was also the most frequently detected unauthorised

human activity. Indeed, the most frequently detected unauthorised human activity in ILUMA

was undoubtedly livestock grazing, with over 1400 instances cattle herding recorded. It

should therefore be cautioned that other unauthorised human activities, like cultivation of rice

and other tillage crops, may have at least comparable impacts that might not necessarily be as

readily attributable to them in statistical terms.

Cattle grazing was encountered just as often around the degraded camps inside ILUMA as

around those outside it, where herding is not restricted but has to compete with agriculture as

a land use option. Although it is clear in the land use agreement (District Authorities, 2016)

that cattle grazing is not a permitted activity inside of ILUMA's boundaries, the evidence

reported herein illustrates just how extensively this aspect of the WMA plan has been

breached. Indeed, the informal observations and anecdotal evidence accumulated by the

investigators indicate that grazing pressure on lands around the ILUMA WMA was being

alleviated by unauthorised cattle herding within the supposedly protected area, especially

close to its the western and southern boundaries.

ILUMA's habitat type is predominantly miombo woodland of different densities, and cattle

herding in Tanzanian miombo woodland, in an area with similar abiotic and biotic

characteristics to that assessed in this study, revealed mixed ecosystem responses to cattle

grazing (Mtimbanjayo and Sangeda, 2018). High intensity cattle grazing caused a significant

negative effect on both the density and diversity of miombo tree species, while less intense

herding, especially during the dry season, was shown to have no adverse effect on the

ecosystem (Mtimbanjayo and Sangeda, 2018). Given the somewhat flexible nature of land

use agreements for WMAs in Tanzania, regulated seasonal grazing is permitted in some

WMAs, such as the Enduimet WMA in the north of the country (University of Copenhagen

Department of Food and Resource Economics, 2015). It may therefore be useful to consider

whether rigorously regulated seasonal grazing in ILUMA might help to control the level of

unregulated herding, by permitting it at certain times and in certain areas only, and in doing

so protecting the areas that are highly valuable for wildlife. The village lands outside ILUMA

are largely used for the production of agricultural crops during the wet season, thus

displacing cattle into the nearby WMA. Seasonally allowing limited cattle grazing inside the

WMA during the rains might not only allow more controlled regulation of this important

local livelihood, it might also enhance productivity of this protected area by preventing the

overgrowth of perennial grasses and encouraging shorter more-nutritious grasses to grow,

thus benefiting both livestock and wildlife (Fynn et al., 2016). Indeed, it is notable that cattle

grazing was common at some locations at the interface between relatively intact and heavily

degraded lands with intermediate to high scores for SRI, SID, OWACII (Supplementary file

10), ONEII and SNEII (Supplementary file 11, figures S11.1), suggesting that this very

common land use activity may not necessarily have as large a negative impact upon wildlife

populations and ecosystem integrity as face value interpretation of table 1 might otherwise

indicate.

Limiting regulated grazing within the ILUMA WMA to the wet season may also be

remarkably compatible with wildlife migration patterns and complementary non-extractive

income-generating activities. Several important wild mammalian species, notably zebra, sable

and hartebeest are known by the VGS and park rangers to migrate westwards into the open

acacia savannah areas of NNP during the rains to access short green grass with high protein

content (Estes, 2012) so the grazing of cattle within the WMA during this season, when taller

grasses suitable for bulk grazing is most abundant, should minimize competition with wildlife

over resources. Conversely, these same wildlife species tend to migrate back into the ILUMA

WMA during the dry season, to exploit its extensive miombo woodland, floodplain

grasslands and groundwater forest habitats. The dry season, when livestock may be returned

to fallow agricultural land, typically runs from June through October, thus overlapping with

the summer holiday season of tourists from Europe and North America. The dry season is

therefore an ideal time for hosting tourists, at a time when it is most convenient to exclude

cattle grazing. However, based on the results presented herein, it is evident that

approximately two thirds of the WMA is already too degraded for tourism and that current

rates of livestock encroachment into the WMA are unsustainable. Without any interventions

like a rigorously regulated wet season grazing plan, to both regulate this important local

livelihood and disincentivise potentially more destructive ones like land conversion for

agriculture, continued encroachment may well result in this WMA becoming non-viable as a

conservation area.

Fortunately, evidence of success with respect to the established practice of another natural

resource utilization activity that conducted in an authorized, carefully regulated manner

within the WMA conservation area is encouraging, suggesting that there may be scope for

participatory development of such compromise solutions. In stark contrast to all the

aforementioned breaches of the ILUMA land use plan across the southern and western parts

of the supposedly protected area of the WMA, one section of the management plan that was

effectively adhered to over the course of this study was the authorised fishing camps in the

north of ILUMA, along the banks of the Kilombero river. Very few unauthorised activities

were detected at these locations despite the presence of substantive long-established fishing

villages, and the areas immediately outside of these settlements appeared ecologically intact

with very little evidence of degradation (Figures 6, 7 and 9B). Crucially, not only do the

residents of these locations have permission to extract fish from the Kilombero river and sell

it as a means of making a living, they also can and do call upon the WMA management and

VGS to enforce fishing regulations and apprehend unlicensed competitors from outside the

area.

This vital collaborative relationship between the WMA authority and one of the stakeholder

communities it serves was reflected in many obvious ways during surveys conducted in and

around these villages. Fishing is carried out with nets or by hook and line, from dugout

canoes hollowed from tree trunks and is traditional in its forms of practice. Based on both

formal data collected and informal observations made while conducting the field work,

notably conversations with the VGS about their interactions with these villages, the residents

of these fishing settlements, generally speaking, agreed with and adhered to the WMA the

land use plan. All three villages, plus two others along the same stretch of river that were not

included in the sampling frame, all maintained human settlement patterns that were confined

within a small area. These settlements are also much smaller in size than the formally

established villages outside of ILUMA, and no large-scale agriculture was recorded in the

vicinity of the three relevant camps. All of these fishing villages were surrounded by fully

intact, dense groundwater forest within meters from the centres of these settlements.

The presence of authorised residents may actually be helpful in deterring unlawful activity in

the areas surrounding these settlements. Due to the mutually beneficial nature of the fishing

camps for both the residents, who practice sustainable livelihoods provided by the intact

natural ecosystem that surrounds them, residents are quick to report suspicions of illegal

activities within the WMA to the VGS. This evidence clearly demonstrates that well-managed

human settlements within a protected area can, not only have minimal direct impact upon on

the ecological intactness of an ecosystem, they can actually bolster its conservation through

resident community vigilance and reporting on a year-round basis. Such passive but

continuous surveillance by resident communities living within a protected area contrasts with,

but also complements, more active but intermittent forms of surveillance by community-

based professionals like VGS conducting patrols.

Overall, this success story illustrates how sustainable livelihoods for local people and

effective ecological conservation may be achieved concurrently if adherence to formally

agreed land use plans can be adhered to through compromise and collaboration between

participatory conservation authorities and the resident stakeholder communities that live in

and around such such protected areas. Learning from this example, an analogous approach to

negotiating seasonal grazing agreements with villages adjacent to the WMA, or even those

already established within it, might involve conditionally allowing limited use of protected

rangelands within the WMA, in exchange for de facto participation of beneficiary

communities in collaborative regulation of not only grazing itself, but also potentially more

destructive activities like agriculture and extractive deforestation.

Such compromise solutions may prove crucially important to delivering on the roles that

WMAs, and other sundry institutional models for community-based management of fringe

conservation areas, are meant to play as buffer zones around more vertically managed,

rigorously protected parks and reserves. Despite the extensive encroachment seen in the west

and south of the ILUMA WMA, the gradient of wildlife activity across the WMA clearly

increases moving eastward and northward as the conservation status of the land improves and

evidence of illegal human encroachment dissipates, with some parts of the WMA having

similar ecosystem integrity to the adjoining national park, not to mention higher mammalian

biodiversity. The ILUMA WMA clearly does indeed act as an important buffer zone for NNP

by providing a protected area with extensive woodland, forest and floodplain habitats that are

useful for many species, including many from inside the national park that use it on a

seasonal or opportunistic basis.

The remaining pristine areas of ILUMA offers invaluable habitat supporting a wide diversity

of wild mammals, further emphasising the need for creative solutions to achieve

collaborative, community-based regulation of human encroachment into the WMA.

Beyond merely halting ongoing ecosystem degradation in such compromised fringe

conservation area across the tropics, it has never been more important to restore them to their

intact natural state (Strassburg, 2020, Wilkinson, 2019) so the ongoing evolution of effective

institutional models for their devolved governance and management is of paramount

importance. Indeed, more effective schemes for managing vulnerable forested areas in the

face of growing human population pressure have a particularly important role to play in

simultaneously combating climate change (Strassburg, 2020) and the emergence of novel

pathogens capable of causing pandemics (Wilkinson, 2019). The natural land cover of

ILUMA is predominantly miombo woodland - an ecosystem type with deep soils, abundant

hardwood trees and very high carbon sequestration potential when it recovers: The wood of

regrowing miombo trees sequesters 0.5 to 0.9 tons of carbon ha^{-1}per year, while soils in a

regrowing miombo woodland sequester >100 tons of carbon ha^{-1}per year (Williams et al.,

2008). Furthermore, the biodiversity of mammals (and presumably the pathogens they carry)

was highest in parts of the study area with miombo woodland vegetation cover, notably in

where it overlapped with various human activities locations at the scrubby interface between

degraded and pristine land (Supplementary files 1, 2, 10).

702 5 | CONCLUSION.

Across much of ILUMA WMA, in areas where agreed land use plans were not adhered to,

rampant cattle herding and land clearing for agriculture were associated with reductions in

wildlife richness and biodiversity, as well as overall ecosystem integrity. Although human

settlement was also generally associated with reduced natural ecosystem integrity, some

notable exceptions illustrate how sustainable livelihoods for local people that are based on

well-managed natural resource harvesting practices may actually enhance conservation

effectiveness: Several authorised fishing villages within the WMA, where fishing was

permitted and local communities collaborated with the WMA management to enforce

conservation regulations, were surrounded by pristine land cover with thriving terrestrial

wildlife populations. Correspondingly, the best conserved parts of the WMA not only

included those closest to the boundary with the national park to the east, but also these fishing

villages along the riverbank to the north, where compliance with agreed land use plans was

most rigorous. Applying lessons learned from this success story more broadly across the

WMA, the offer to negotiate conditional grazing rights within the ILUMA during the rainy

season, when cattle need to be moved off neighbouring agricultural land, might motivate

beneficiary communities to participate in more effective collaborative interventions against

unsustainable levels of ongoing encroachment. Overall, this study illustrates how well-

managed WMAs may host resident local communities undertaking selective natural resource

extraction activities and collaborate with them to achieve effective *de facto* conservation

practices, thus maintaining robust wildlife populations and acting as effective buffer zones

along the fringes of larger core conservation areas like national parks and game reserves.

COMPETING INTERESTS STATEMENT

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

Supplementary file 1; Table S1: The number, name, location, coordinates, and ecological

characteristics of each camp location, together with the quadrant circuit to which it was

assigned and the number of times it was surveyed https://doi.org/10.5281/zenodo.11099575

Supplementary file 2; Table S2. Scores for various indices of the integrity of the wild animal

community or ecosystem as a whole, estimated for each surveyed camp location (Section 2.2)

based on either a consensus approach to synthesizing investigator-perception (SNEII, Section

2.6) or numerical syntheses (SRI, SID, OWACII and ONEII, Section 2.4) or cluster analysis

(Section 2.5) of extensive data from formal, quantitative surveys of observable human,

livestock and wildlife activities, as well as land cover characteristics (Section 2.3 and

Supplementary file 8) https://doi.org/10.5281/zenodo.11099588

Supplementary File 3; Figures S3.1 to S3.4: Photographs illustrating different ways in

which signs of livestock, humans or wild mammal activity, as well as land cover attributes,

were observed and classified https://doi.org/10.5281/zenodo.11099593

Supplementary File 4; Complete list of all wild herbivores, wild carnivores, wild

primates   and   prosimians   and   wild   rodents   sampled   in   this   study

https://doi.org/10.5281/zenodo.11099600

Supplementary File 5; Data collection key devised and refined during a preliminary pilot

study between October and November 2021 and used throughout the course of the field study

to inform accurate and consistent recording of data https://doi.org/10.5281/zenodo.11099602

Supplementary   File   6;   Criteria   defining   terms   used   in   data   dictionary   key

https://doi.org/10.5281/zenodo.11099616

Supplementary File 7; Data collection sheet developed during preliminary investigation in

October and November 2021 and used for all data collection in both transect segment surveys

and camp radial surveys throughout the course of the study

https://doi.org/10.5281/zenodo.11099618

Supplementary File 8; The full data set derived from formal quantitative surveys of land

cover types and activities of humans, livestock and wildlife (Section 2.3) that were used for

the analyses described in sections 2.4, 2.5 and 2.7 https://doi.org/10.5281/zenodo.11099625

Supplementary File 9; Figures S9.1 to S9.10: Maps of the ILUMA WMA, respectively

illustrating the geographic distribution of all the various human activities surveyed as

described in section 2.3 https://doi.org/10.5281/zenodo.11099630

Supplementary File 10; Tables S10.1 to S10.3 detailing the results of univariate and

multivariate generalized linear modelling analyses of scaled human activity indicators as

respective determinants of the estimated Species Richness Index (SRI), Simpson's Index of

Diversity (SID) and Objective Wild Animal Community Integrity Index (OWACII) for each

camp surveyed. Integrity https://doi.org/10.5281/zenodo.11099636

Supplementary File 11; Supplementary figures S11.1 to S11.5; Maps illustrating how the

distributions of the values for all the significant independent variables that were included in

the final multivariate model detailed in table 1, except for the extent of human settlement

(See figure 8), compared with those for the Subjective Natural Ecosystem Integrity Index

(SNEII) scores that were used as the dependent variable in that model

https://doi.org/10.5281/zenodo.11099639

Supplementary File 12; Supplementary figures S12.1 to S12.5; Dot plots and violin plots

respectively illustrating the absolute frequency and probability density distributions of all the

various recorded signs of activity by wild animals, livestock, work animals and humans

during the surveys described in section 2.3 https://doi.org/10.5281/zenodo.11099645

Reference List

776   Agrawal, A., Ribot, J., 1999. Making decentralization accountable: A framework for analysis
777   and empirical studies from South Asia and West Africa. J. Dev. Areas 33, 473–490.
778

779   Bai, Z.G., Dent, D.L., Olsson, L., Schaepman, M.E., 2008. Proxy global assessment of land
780   degradation. Soil Use and Management 24, 223–234. https://doi.org/10.1111/j.1475-
781   2743.2008.00169.x
782

783   Baisero, D., Visconti, P., Pacifici, M., Cimatti, M., Rondinini, C., 2020. Projected global loss
784   of mammal habitat due to land-use and climate change. One Earth 2, 578–585.
785

Daconto, G., Games, I., Lukumbuzya, K., Raijmakers, F., 2018. Integrated Management Plan for the Kilombero Valley Ramsar Site. Ministry of Natural Resources and Tourism: Dar es Salaam, Tanzania.

District Authorities, 2016. ILUMA's wildlife association by-laws for the villages of Ulanga and Ifakara in the KIlombero District 2016. Local government act, chapter 287, composed under article 163.

Duggan, L. (2024). The influence of community-defined land use plans and *de facto* land use practices on the relative abundance and distribution of large wild mammals in a community-based Wildlife Management Area in Southern Tanzania [MSc Dissertation, University College Cork, Ireland].

Duggan, L.M, Tarimo, L.J., Walsh, K.A., Kavishe, D.R., Crego, R.D., Elisa, M., Mombo, F., Butler, F., and Killeen, G.F. (Forthcoming A). Comparative assessment between radial and transect surveys to document wild mammal activity across diverse habitat types in southern Tanzania.

Duggan, L.M, Tarimo, L.J., Walsh, K.A., Kavishe, D.R., Crego, R.D., Elisa, M., Mombo, F., Butler, F., and Killeen, G.F. (Forthcoming B). A subjective and intuitive approach to rapid, holistic assessment of natural ecosystem integrity across a community-managed conservation area in southern Tanzania.

Estes, R. D. (2012). The behaviour guide to African mammals: Including hoofed mammals, carnivores, primates. Univ of California Press.

Fynn, R.W., Augustine, D.J., Peel, M.J., de Garine☐Wichatitsky, M., 2016. Strategic management of livestock to improve biodiversity conservation in A frican savannahs: a conceptual basis for wildlife–livestock coexistence. Journal of Applied Ecology 53, 388–397.

Gibson, C.C., Marks, S.A., 1995. Transforming rural hunters into conservationists: an assessment of community-based wildlife management programs in Africa. World development 23, 941–957.

Goldman, M., 2003. Partitioned nature, privileged knowledge: community☐based conservation in Tanzania. Development and change 34, 833–862.

Kavishe, D.R., Msoffe R.V., Malika, G.Z., Walsh, K.A., Duggan, L.M., Tarimo, L.J., Butler, F., Kaindoa, E.W.W., Ngowo, H.S., & Killeen, G.F. (2024) 'A self-cooling self-humidifying mosquito carrier backpack for transporting live adult mosquitoes on foot over long distances

under   challenging   field   conditions'.   BioRχive
https://biorxiv.org/cgi/content/short/2024.04.13.588955v1

Kiss, A., 2004. Is community-based ecotourism a good use of biodiversity conservation
        funds? Trends in ecology & evolution 19, 232–237.

Lee, D.E., Bond, M.L., 2018. Quantifying the ecological success of a community-based
        wildlife conservation area in Tanzania. Journal of Mammalogy 99, 459–464.

Lindsey, P.A., Alexander, R., Frank, L.G., Mathieson, A., Romanach, S., 2006. Potential of
        trophy hunting to create incentives for wildlife conservation in Africa where
        alternative wildlife☐based land uses may not be viable. Animal conservation 9, 283–
        291.

Mombo, F.M., Speelman, S., Van Huylenbroeck, G., Hella, J., Pantaleo, M., Moe, S., 2011.
        Ratification of the Ramsar convention and sustainable wetlands management:
        Situation analysis of the Kilombero Valley wetlands in Tanzania. Journal of
        Agricultural Extension and Rural Development 3, 153–164.

Moore, J.C., 2013. Diversity, Taxonomic versus Functional, in: Levin, S.A. (Ed.),
        Encyclopedia of Biodiversity (Second Edition). Academic Press, Waltham, pp. 648–
        656. https://doi.org/10.1016/B978-0-12-384719-5.00036-8

Mtimbanjayo, J., Sangeda, A., 2018. Ecological effects of cattle grazing on Miombo tree
        species regeneration and diversity in Central-Eastern Tanzania. Journal of
        Environmental Research 2, 1–7.

Nkuba, J., Ndunguru, A., Madulu, R., Lwezaura, D., Kajiru, G., Babu, A., Chalamila, B., Ley,
        G., 2016. Rice value chain analysis in Tanzania: identification of constraints,
        opportunities and upgrading strategies. African Crop Science Journal 24, 73–87.
Nyerere National Park, 2023.

Population Division of the United Nations Department of Economic and Social Affairs, 2022.
        World Population Prospects 2022: Summary of Results (UN DESA).

Salerno, J., Mudler, M.B., Grote, M.N., Ghiselli, M. and Packer, C. (2016) 'Household
livelihoods and conflict with wildlife in community-based conservation areas across northern
Tanzania', Oryx, 50(4), pp. 702–712.

Songorwa, A.N., 1999. Community-based wildlife management (CWM) in Tanzania: are the
        communities interested? World development 27, 2061–2079.

872 Spenceley, A., 2012. Implications of responsible tourism for conservation and development in
873   southern Africa, in: Responsible Tourism. Routledge, pp. 389–404.
874

Strassburg, B.B.N, Iribarrem, A., Beyer, H.L., Cordeiro, C.L., Crouzeilles, R., Jakovac, C.C.,
Junqueira, A.B., Lacerda, E., Latawiec, A., Balmford, A., Brooks, T.M., Butchart, S.H.M.,
Chazdon, R.L., Erb, K-H, Brancalion, P., Buchanan, G., Cooper, D., Diaz, S., Donald, P.F.,
Kapos, V., Leclere, D., Miles, L., Obersteiner, M., Plutzar, C., Scaramuzza, C.A.M., Scarano,
F.R. and Visconti, P. (2020) Global priority areas for ecosystem restoration. *Nature* **586**, 724–
729

United Republic of Tanzania Ministry for Natural Resources and Tourism, 2023. Tanzanian
Wildlife Management Authority.
884

885 United Republic of Tanzania Ministry for Natural Resources and Tourism, 2022. The Wildlife
886   Conservation Act. Revised Edition 2022. Chapter 283.
887

888 United Republic of Tanzania Ministry of Natural Resources and Tourism, 2015.
889   KILORWEMP News Letter 4/15.
890

891 University of Copenhagen Department of Food and Resource Economics, 2015. Policy briefs
892   (Copenhagen Centre for Research and Development) (Online) (Policy Briefs •
893   Copenhagen Centre For Development Research No. ISSN 2246-6800).
894

895 USAID, 2018. Wildlife Management Area (WMA) Viability Assessment: Pilot Report.
896

Walsh, K.A. (2024) Blood host preferences and competitive inter-species dynamics within an
African malaria vector species complex inferred from signs of animal activity around aquatic
larval habitats distributed across a gradient of fully domesticated to fully pristine ecosystems
in southern Tanzania. M.Sc. Dissertation. University College Cork, Ireland.

Whittaker, R.J., Willis, K.J., Field, R., 2001. Scale and species richness: towards a general,
hierarchical theory of species diversity. Journal of biogeography 28, 453–470.
904

Wilkinson, D.A., Marshall, J.C., French, N.P., Hayman, D.T.S. (2018) Habitat fragmentation,
biodiversity loss and the risk of novel infectious disease emergence. Journal of the Royal
Society Interface, 15, 20180403

Williams, M., Ryan, C., Rees, R., Sambane, E., Fernando, J., Grace, J., 2008. Carbon
sequestration and biodiversity of re-growing miombo woodlands in Mozambique.
Forest Ecology and management 254, 145–155.
912

SNEII

- ■ 0 - 17
- ■ 17 - 50.8
- ■ 50.8 - 78.8
- ■ 78.8 - 85.8
- ■ 85.8 - 100

Human Settlement (no. of structures)

- ○ 1 - 10
- ○ 11 - 20
- ○ 50 +

- NNP
- Kilombero Wetlands
- ILUMA
- Ifakara Road

0   5   10 km