Disruption of mitochondrial folate metabolism leads to mitochondrial DNA leakage and activation of apoptotic or inflammatory pathways

Sinwoo Hwang[a], Cameron Baker[a], Martha S. Field[a*]

[a]Division of Nutritional Sciences, Cornell University, Ithaca, NY, USA

Corresponding author: Martha S. Field, Cornell University, mas246@cornell.edu

Abstract

Folate-mediated one-carbon metabolism (FOCM) is required for the biosynthesis of

purines, thymidylate (dTMP), and methionine. Maintenance of adequate cellular dTMP

levels is essential to preserve the integrity of the nuclear and mitochondrial genomes.

Inadequate dTMP production leads to uracil misincorporation into mitochondrial DNA

(mtDNA) and impaired mitochondrial function. However, the mechanisms whereby uracil

in mtDNA impairs mitochondrial function are uncharacterized. The release of mtDNA

into the cytosol acts as a damage-associated molecular pattern (DAMP), resulting in

inflammation and cell death. Previously, we reported that disrupting mitochondrial dTMP

synthesis through serine hydroxymethyltransferase 2 (*Shmt2*) heterozygosity increases

uracil in mtDNA and impairs mitochondrial function in mice. This study aimed to

investigate whether impairment of mitochondrial FOCM through *Shmt2* disruption leads

to the release of mtDNA into the cytosol. $Shmt2^{+/-}$ MEF cells showed a > 2-fold increase

in cytosolic mtDNA leakage compared to $Shmt2^{+/+}$ cells (p<0.05). There was no

significant difference in total mtDNA content between $Shmt2^{+/+}$ and $Shmt2^{+/-}$ MEF cells.

MEFs with decreased *Shmt2* expression activated apoptosis by the ratio of cleaved

caspase-3 to caspase-3. In addition, SHMT2 knock-out (SHMT2 KO) haploid chronic

myeloid leukemia (HAP1) cells also exhibited increased cytosolic mtDNA content

compared to the wild-type (WT) HAP1 cells. HAP1 lacking *Shmt2* expression activated

the cGAS/STING pathway, but suppressed apoptosis, compared to WT HAP1 cells.

This study demonstrates that decreased *Shmt2* expression leads to cytosolic mtDNA

leakage and that the downstream effects of mtDNA leakage vary by cell type.

Keywords: Folate, folate-mediated one-carbon metabolism, mtDNA, apoptosis,

inflammation

Introduction

Folate, also known as vitamin B9, is an essential cofactor required for folate-mediated

one-carbon metabolism (FOCM) [1]. This metabolic pathway provides one-carbon units

for biological processes, including biosynthesis of purines, thymidylate (dTMP), and

methionine [2]. Disruptions of FOCM are associated with an increased risk of various

pathologies such as anemia, cancer, neurodegeneration, and neural tube defects.

FOCM is compartmentalized within the cell, taking place in the cytosol, nucleus, and

mitochondria [3, 4]. FOCM is essential to maintain an adequate cellular dTMP level to

preserve the integrity of the nuclear and mitochondrial genomes. Impaired folate-

dependent *de novo* synthesis of dTMP leads to uracil misincorporation into nuclear DNA

(nDNA) and mitochondrial DNA (mtDNA), in turn, causing genome instability [5]. In

HeLa cells and in mouse liver, folate deficiency elevates uracil in in mtDNA without

increasing uracil in nDNA, indicating that mtDNA is more susceptible to disruption of

FOCM than is nDNA [6, 7].

Serine hydroxymethyltransferase 2 (SHMT2) facilitates the reversible conversion of THF

and serine into 5,10-methylene-THF and glycine within the mitochondria [8]. In addition

to SHMT2, mitochondrial biosynthesis of dTMP also requires folate-dependent enzymes

synthase (TYMS) and dihydrofolate reductase (DHFR). These enzymes are responsible

for converting deoxyuridine monophosphate (dUMP) into dTMP [4, 9, 10]. Therefore,

disruption of *Shmt2* expression impairs mitochondrial dTMP synthesis. In mouse liver

and cultured cells, *Shmt2* disruption leads to uracil misincorporation in mtDNA and

impaired oxidative phosphorylation [11]. The mechanisms whereby uracil

misincorporation in mtDNA impairs mitochondrial function have not been elucidated.

Repair of uracil misincorporation by base-excision repair mechanisms can damage DNA

by generating strand-breaks and abasic sites during the repair process. MtDNA

damage can lead to mtDNA leakage into the cytosol, where mtDNA then acts as a

mitochondrial-derived damage-associated molecular pattern (DAMP). Increasing

evidence suggests that mitochondria can initiate the innate immune response and

apoptosis by releasing mitochondrial DAMPs under various stress conditions [12, 13]. In

addition to its role in innate immunity [14], mitochondrial DAMPs are involved in

apoptotic cell death through the activation of caspases [15]. Still, it is unclear how

mitochondrial DAMPs regulate the balance between immune response and apoptosis.

Given the role of SHMT2 in maintenance of mtDNA integrity, the aim of this study was to

investigate whether decreased *Shmt2* expression induces mtDNA release into the

cytosol and how cytosolic mtDNA release affects downstream processes including

apoptosis and the immune response.

Materials and Methods

Mouse models, cell lines, and cell culture conditions:

Shmt2$^{+/-}$ whole-body knockout mice (homozygous knockout is embryonic lethal) on a

C57BL/6J background were generated as previously described [11]. Mouse embryonic

fibroblasts (MEFs) were isolated from C57BL/6J female mice bred to *Shmt2$^{+/-}$* male mice

as previously described [11]. HAP1 cells (wild-type) and SHMT2 knock-out HAP1

(*SHMT2* KO) HAP1 cells were obtained from Horizon Discovery. MEF cells were

passaged in alpha-minimal essential medium (alpha-MEM; Hyclone Laboratories,

Logan, UT, USA) supplemented with 10% FBS (VWR, Radnor, PA, USA) and 1%

penicillin/streptomycin (Gibco, Grand Island, NY, USA). HAP1 cells were passaged in

Iscove's Modification of DMEM (IMDM, Corning, NY, USA) supplemented with 10% FBS

(VWR, Radnor, PA, USA) and 1% penicillin/streptomycin (Gibco, Grand Island, NY,

USA). Cells were cultured at 37 ℃ in a humidified atmosphere of 5% CO_2. To identify

the effect of folate availability, HAP1 cells were cultured in the folate deficient (0 nM)

and the folate sufficient (25 nM) modified DMEM (Hyclone Laboratories, Logan, UT,

USA) as previously described (35).

Preparation of cell extracts:

When cultured MEF and HAP1 cells reach 80% confluency in a 6-well plate, the cells

were washed with ice-cold phosphate buffered saline (PBS) three times. The cells were

collected in 1X radioimmunoprecipitation assay buffer (RIPA buffer; Cell Signaling,

Danvers, MA, USA), supplemented with 1 mM PMSF (Cell Signaling, Dancers, MA,

USA), 0.5 M EDTA Solution (Thermo Fisher Scientific, Waltham, MA, USA), and Halt

Phosphatase Inhibitor Cocktail (Thermo Fisher Scientific, Waltham, MA, USA). The cells

were incubated on ice for 20 minutes and centrifuged at 13,000 × g for 10 minutes at

4 ℃. The supernatant was transferred to a new 1.5 mL tube. Total protein was quantified

by the Pierce BCA protein assay (Thermo Fisher Scientific, Waltham, MA, USA)

following manufacturer's instruction.

Immunoblotting:

Protein lysates were denatured by heating with 4X Laemelli buffer (Bio-Rad, Hercules,

CA, USA) for 5 minutes at 95 ℃. Samples were loaded onto 4-12% Sodium dodecyl-

sulfate polyacrylamide gel electrophoresis (SDS-PAGE) gels for 60-90 minutes in SDS-

PAGE running buffer and transferred to Polyvinylidene difluoride (PVDF) membranes

(Bio-Rad, Hercules, CA, USA). Membranes were stained with Ponceau S to verify the

transfer of proteins. Membranes were blocked using 5% (w/v) non-fat dry milk in 1X

Tris-buffered saline and tween-20 (TBS-T) for 1 hour at room temperature. The blocked

membranes were incubated in the respective primary antibody at 4 ℃ overnight.

Primary antibodies that were used are as follows: CS (1:1000, ab96600, Abcam,

Cambridge, UK), cGAS (1:1000, 31689, Cell Signaling, Danvers, MA, USA), STING

(1:1000, 50494, Cell Signaling, Danvers, MA, USA), TBK1 (1:1000, 3504, Cell Signaling,

Danvers, MA, USA), Phospho-TBK1 (1:1000, 5483, Cell Signaling, Danvers, MA, USA),

Caspase-3 (1:1000, 9662, Cell Signaling, Danvers, MA, USA), Cleaved caspase-3

(1:1000, 9664, Cell Signaling, Danvers, MA, USA), and GAPDH (1:2118S, Cell

Signaling, Danvers, MA, USA). Membranes were washed 3 times with 1X TBS-T for 5

minutes each and incubated in the appropriate horseradish peroxidase-conjugated

secondary antibody for 2 hours at room temperature, followed by washing 3 times with

1X TBS-T. To detect antibody, an enhanced chemiluminescent reagent (SuperSignal

west femto maximum sensitivity substrate, Thermo Fisher Scientific, Waltham, MA, USA)

was used at a 1:10000 dilution. Blots were imaged using FluorChem E imaging system

(Protein Simple, San Jose, CA, USA). Densitometry was performed with Fiji software

using GAPDH as the control.

Sub-fractionation of cellular compartments (Mitochondria):

Cultured MEF cells (5×10^6) or HAP1 (1×10^7) were pelleted by centrifugation at $500 \times$

g for 10 minutes at 4 ℃. The pellets were washed with 0.9 % sodium chloride (NaCl)

and pelleted again by centrifugation at $500 \times$ g for 10 minutes at 4 ℃. Qproteome

mitochondria isolation assay (Qiagen, Hilden, Germany) was performed to extract high-

purity mitochondria. The pellets were resuspended in ice-cold Qproteome lysis buffer

supplemented with 100X protease inhibitor solution and incubated for 10 minutes at 4 ℃

on an end-over-end shaker. The lysates were pelleted by centrifugation at $1000 \times$ g for

10 minutes at 4 ℃ and resuspended in ice-cold Qproteome disruption buffer. The

lysates were disrupted with a blunt-ended needle and a syringe and centrifugated at

$1000 \times$ g for 10 minutes at 4 ℃. The supernatant was transferred to a fresh 1.5 mL tube,

followed by another centrifugation at $6000 \times$ g for 10 minutes at 4 ℃ to obtain

mitochondria-containing pellets. To obtain high-purity mitochondria, the pellets were

resuspended in Qproteome mitochondrial purification buffer and pipetted on top of

layers of purification buffer and disruption buffer, followed by a centrifugation at $14000 \times$

g for 15 minutes at 4 ℃. The mitochondrial pellets were washed three times with

Qproteome mitochondrial storage buffer and prepared for further experiments.

Sub-fractionation of cellular compartments (Cytosol):

Cultured cells were washed three times with ice-cold PBS and pelleted by centrifugation

at $500 \times$ g for 5 minutes at 4 ℃. Cell pellets were resuspended in cytosolic extraction

buffer (150 mM NaCl, 50 mM HEPES pH 7.4, 25 μg/ml digitonin) and homogenized

using a 2 mL Dounce grinder with a tight-fitting pestle (Kimble Chase, Vineland, NJ,

USA) for 10 minutes at 4 ℃. Lysates were centrifuged at $10000 \times$ g for 15 minutes at 4

°C. The supernatant was transferred to a fresh 1.5 mL tube, followed by another

centrifugation at 12000 × g for 15 minutes at 4 °C. The cytosolic fraction was isolated by

collecting supernatant into a fresh 1.5 mL tube. The purity of cytosolic fraction without

mitochondrial contamination was determined by immunoblotting analysis with CS

(1:1000, ab96600, Abcam, Cambridge, UK)

Measurement of mitochondrial DNA content:

Total genomic DNA was isolated from whole, mitochondrial, and cytosolic cellular

fractions respectively, using the QIAamp DNA Micro Kit (Qiagen, Hilden, Germany)

following manufacturer's instructions. DNA purity and concentration were quantified

using Nanodrop spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA). To

determine mtDNA copy number, real-time quantitative polymerase chain reaction (qPCR)

was performed with the LightCycler 480 SYBR-Green I Master (Roche Diagnostics,

Basel, Switzerland) and primers specific to the mitochondrial D-loop region that is not

inserted into nuclear DNA region (non-NUMT). Nuclear DNA encoding *Tert* primers were

used for normalization. PCR amplification was performed with a 10-minutes

denaturation step at 95°C, followed by 40 cycles of 95°C for 10 seconds, 60°C for 30

seconds, and 72°C for 10 seconds. Primer sequences are listed as follows:

Mouse D-loop forward: 5'- AATCTACCATCCTCCGTGAAACC-3'

Mouse D-loop reverse: 5'-TCAGTTTAGCTACCCCCAAGTTTAA-3'

Mouse *Tert* forward: 5'-CTAGCTCATGTGTCAAGACCCTCTT-3'

Mouse *Tert* reverse: 5'-GCCAGCACGTTTCTCTCGTT-3'

Human D-loop forward: 5'-GTTTATGTAGCTTACCTCCTC-3'

Human D-loop reverse: 5'-TTGTTTATGGGGTGATGTGAG-3'

Human *Tert* forward: 5'-TCACGGAGACCACGTTTCAAA-3'

Human *Tert* reverse: 5'-TTCAAGTGCTGTCTGATTCCAAT-3'

Statistical analysis:

Statistical analysis was performed using GraphPad Prism 8.0. The Welch's t-test was

used to determine whether there is a significant difference between the means of

Shmt2$^{+/+}$ and *Shmt2*$^{+/-}$ MEFs, and wild-type HAP1 and SHMT2 KO HAP1. A *p*-value of

0.05 or less was considered statistically significant.

Results

Reduced *Shmt2* expression leads to mtDNA leakage into the cytosol without

changing total mtDNA content.

To examine mtDNA leakage into the cytosol, cytosol was isolated from the cultured

Shmt2$^{+/+}$ and *Shmt2*$^{+/-}$ MEF cells. Using the primers specific to the mitochondrial D-loop

region that is not inserted into nuclear DNA (non-NUMT) and nuclear DNA encoding *Tert*

primers as control, cytosolic mtDNA content was measured by qPCR. Citrate synthase,

a mitochondrial matrix enzyme, is commonly used as a maker for the presence of intact

mitochondria (i.e. lack of mitochondrial contamination in cytosolic fraction). As shown in

Figure 1A and 1D, respectively, citrate synthase was only observed in the total cell

lysate and not in the cytosol of MEF (*Shmt2*$^{+/+}$ and *Shmt2*$^{+/-}$) and HAP1 (wild-type and

SHMT2 KO) cells, confirming the purity of the cytosolic fractions. There was no

significant difference in total cellular mtDNA content between $Shmt2^{+/+}$ and $Shmt2^{+/-}$

MEF cells (Figure 1B), consistent with previous results [11]. However, the $Shmt2^{+/-}$ MEF

cells showed a > 2-fold increase in cytosolic mtDNA leakage compared to the $Shmt2^{+/+}$

group (Figure 1C). Similarly, *SHMT2* KO HAP1 cells exhibited higher levels of mtDNA

content in the cytosol compared to the wild-type HAP1 cells (Figure 1F). As observed in

MEF cells (Figure 1B), there was no significant difference in total mtDNA between wild-

type and *SHMT2* KO HAP1 cells (Figure 1E). In addition, wild-type HAP1 cells were

cultured in the folate-deficient modified cell culture medium to determine if the folate

availability affects the cytosolic mtDNA leakage. No significant change was observed in

the cytosolic mtDNA content between the control and folate-deficient media groups

(Figure 1G), suggesting that exposure to folate-deficient culture medium does not

increase mtDNA leakage to the cytosol.

$Shmt2$ heterozygosity-induced mtDNA leakage increases apoptosis in MEFs but

suppresses apoptosis in HAP1 cells.

Heterozygous disruption of $Shmt2$ expression in MEF cells lead to an increase in

cytosolic mtDNA leakage (Figure 1). Since cytosolic mtDNA leakage is linked to both

and inflammation, we investigated whether reduced expression of $Shmt2$ increases pro-

apoptotic activity by measuring the ratio of cleaved caspase-3 to caspase-3 in whole-

cell lysates of $Shmt2^{+/+}$ and $Shmt2^{+/-}$ MEF cells. $Shmt2^{+/-}$ MEF cells had a significantly

higher level of the cleaved caspase-3 compared to $Shmt2^{+/+}$ MEF cells. Furthermore,

the ratio of cleaved caspase-3 to pro-caspase-3 was more than 10-fold higher in

$Shmt2^{+/-}$ MEF cells compared to $Shmt2^{+/+}$ MEF cells (Figure 2A-C). In contrast, *SHMT2*

KO-induced cytosolic mtDNA in HAP1 cells was associated with decreased apoptotic

pathway activation compared to WT HAP1 cells (Figure 2D-F). Taken together, these

results suggest that *Shmt2* heterozygosity-induced mtDNA leakage into the cytosol is

associated with apoptosis in MEF, but not HAP1 cells.

Cytosolic mtDNA is associated with activation of the cGAS/STING pathway in

HAP1 cells lacking *SHMT2* but not with *Shmt2* heterozygosity in MEF cells.

Previous studies have demonstrated that apoptotic caspases attenuate cGAS/STING-

mediated inflammatory signals triggered by mtDNA [16]. In light of our discovery that the

release of mtDNA into the cytosol due to *Shmt2* heterozygosity is linked to apoptosis in

MEF cells (Figure 2), we hypothesized that the activity of the cGAS/STING pathway

would be decreased in *Shmt2*$^{+/-}$ MEF cells. The protein levels of cGAS and STING were

evaluated in *Shmt2*$^{+/+}$ and *Shmt2*$^{+/-}$ MEF cells using western blotting. No notable

differences were observed between two groups (Figure 3A-C). Interestingly, SHMT2 KO

HAP1 cells exhibited significant higher STING expression and an increase in the ratio of

pTBK-1/TBK-1, compared to the wild-type HAP1 cells (Figure 3D-3F). This suggests

that the *Shmt2* heterozygosity-induced mtDNA release into the cytosol does not activate

cGAS/STING pathway in MEF cells; conversely, HAP1 cells lacking SHMT2 expression

was associated with cGAS/STING activation.

Conclusions and discussion

Mitochondria are essential organelles within eukaryotic cells that play a crucial role in

energy production and cellular respiration [17-19]. SHMT2 is a folate-dependent

enzyme that is integral to mitochondrial FOCM. Mitochondrial FOCM is essential for

various cellular processes, including the biosynthesis of formate for cytosolic/nuclear

FOCM, mitochondrial dTMP production, and intermediates required for mitochondrial

protein translation [4]. Recent studies have identified biallelic human *SHMT2* variants

that result in decreased in SHMT2 expression in human fibroblasts with age [20, 21].

This has led to a growing interest in understanding the cellular implications of perturbed

mitochondrial FOCM.

Moreover, research on the *Shmt2* gene and its association with mtDNA stability and

mitochondrial function has attracted significant attention. Previously, we demonstrated

that *Shmt2* heterozygosity results in uracil misincorporation into mtDNA and impaired

mitochondrial function without affecting the total mtDNA mass [11]. However, the impact

of *Shmt2* heterozygosity on other aspects of mitochondrial biology remains unclear. In

this study, we aimed to investigate whether *Shmt2* heterozygosity could lead to mtDNA

leakage, activation of apoptosis, increased mtDNA common deletions, and accumulated

deleted mtDNA within the mitochondrial compartment.

Here, we demonstrated that reduced expression of *Shmt2* led to mtDNA leakage into

the cytosol in MEF cells. Interestingly, there was no significant change in total mtDNA

content comparing wild-type and *Shmt2*$^{+/-}$ MEF cells (Figure 1C), suggesting that

impaired mitochondrial FOCM induced by decreased *Shmt2* may lead to leakage of

mtDNA into the cytosol by altering mitochondrial membrane integrity. These findings

corroborate the previous research of Fiddler *et al.* (2021), demonstrating that *Shmt2*

heterozygous MEF cells cultured in low-folate media had decreased mitochondrial

membrane potential [11].

As mentioned above, reduced *Shmt2* expression leads to mitochondrial FOCM

disruption, resulting in the accumulation of uracil misincorporation in mtDNA of mouse

liver [11]. Repairing uracil in nuclear DNA is known to induce genome instability through

single- and double-strand breaks [22]. However, the consequences of uracil

misincorporation into mtDNA remain uncharacterized. Nonetheless, mtDNA breaks –

expected during uracil excision repair in mitochondria – are known to cause mtDNA

deletions [23]. The findings presented here suggest that uracil-induced mtDNA damage

may lead to mtDNA release into the cytosol.

MtDNA is increasingly recognized as a regulator of immune responses and apoptotic

cell death [14, 24]. Mitochondrial DAMPs are endogenous molecules that can alert the

immune system to potential threats, such as tissue damage or infection [25]. In

particular, mtDNA leakage is a crucial component of this alarm system, triggering the

activation of PRRs to initiate a cascade of immunological events to repair or eliminate

threats to the host organism [14]. Our hypothesis posited that decreased *Shmt2*

expression would lead to mtDNA leakage into the cytosol and trigger an innate immune

response. Contrary to this initial hypothesis, no significant difference was observed in

the expression levels of cGAS/STING between wild-type and *Shmt2*$^{+/-}$ MEF cells (Figure

3). The cGAS/STING signaling pathway is one type of the innate immune system that

detects the presence of cytosolic DNA. Specifically, cGAS recognizes and binds to

double-stranded DNA (dsDNA), catalyzing the synthesis of cyclic guanosine

monophosphate-adenosine monophosphate (cGAMP) from ATP and GTP [26, 27]. The

cGAMP, in turn, activates the STING protein as a second messenger [28]. Subsequently,

STING activates TANK-binding kinase 1 (TBK1) through its C-terminal PLPLRT/SD

motif [29]. TBK1 phosphorylates the interferon regulatory factor 3 (IRF3) [30, 31].

Dimerized IRF3 translocate into the nucleus and induces the expression of type I IFNs

[30].

It may be possible that *Shmt2* heterozygosity-induced mtDNA leakage is not sufficient to

activate the cGAS/STING pathway. Alternatively, there may be other cellular

mechanisms that can compensate for the increased mtDNA leakage in *Shmt2*

heterozygous MEF cells and maintain a balanced immune response to prevent

cGAS/STING activation. Interestingly, we found that *Shmt2* heterozygosity-induced

mtDNA leakage can promote apoptosis through a cascade of caspase, instead of

activating cGAS/STING-mediated inflammation (Figure 2). This finding is consistent with

previous research that apoptotic caspases suppress type I IFN production via the

cleavage of cGAS [32]. In contrast, *SHMT2* KO HAP1 cells also exhibit increased

mtDNA leakage to the cytosol, but in this cell type the cGAS/STING pathway is

activated as a result of *SHMT2* loss (Figure 2).

The findings presented in this study offer novel insights into the mechanisms underlying

SHMT2 disruption-induced impairments in FOCM, which can lead to mitochondrial

dysfunction and immune response activation and/or cell death. It is important to note

that mtDNA release as a result of impaired mitochondrial FOCM is observed only in

response to reduced *SHMT2* expression, and not as a result of exposure to folate-

deficient culture medium. Both folate deficiency and *Shmt2* disruption were shown to

increase uracil accumulation in mtDNA and impair mitochondrial function in previous

studies [11, 33] , suggesting that decreased SHMT2 may cause cytosolic mtDNA

leakage through mechanisms other than just impairing FOCM such as impaired

mitochondrial translation or impaired redox balance [4]. *SHMT2* expression decreases

with aging in human tissues [21], highlighting the need to better understand the

mechanisms linking decreased *SHMT2* expression, mtDNA leakage, and immune

activation in the context of aging. Future studies should focus on determining whether

cytosolic mtDNA can initiate the innate immune response through TLR9 or

inflammasome-mediated pathways in addition to the cGAS/STING pathway. In addition,

it remains unclear which membrane pore is responsible for the mtDNA release under as

a result of reduced *Shmt2* expression. Therefore, future work should focus on

investigating the role of SHMT2 in cell-free mtDNA-mediated cell death across a range

of tissues and determine the molecular mechanisms that cause mtDNA leakage.

Acknowledgements

The authors thank Luisa Castillo and Katarina Heyden for technical assistance.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Author contributions

SH and MSF designed research. SH and CB conducted research. SH, CB and MSF

analysed data. SH prepared the figures. SH and MSF prepared the manuscript. MSF

has primary responsibility for the final content. All authors reviewed and approved the

final manuscript.

Funding

This work was funded by institutional funds provided to MSF. Cameron Baker was a

student in the Molecular Biology and Genetics Research Experience for Undergraduate

(MBG-REU) program, which was supported by the NSF (DBI1950247), the Department

of Molecular Biology and Genetics, the Weill Institute of Cell and Molecular Biology, and

the Division of Nutritional Sciences at Cornell University.

References

1.   P. J. Stover, Polymorphisms in 1-carbon metabolism, epigenetics and folate-
related pathologies, Lifestyle Genomics, 4 (5) (2012) 293-305.
2.   P. J. Stover and M. S. Field, Trafficking of intracellular folates, Advances in
nutrition, 2 (4) (2011) 325-331.
3.   A. S. Tibbetts and D. R. Appling, Compartmentalization of Mammalian folate-
mediated one-carbon metabolism, Annual review of nutrition, 30 (2010) 57-81.
4.   Y. Xiu and M. S. Field, The roles of mitochondrial folate metabolism in supporting
mitochondrial DNA synthesis, oxidative phosphorylation, and cellular function,
Current Developments in Nutrition, 4 (10) (2020) nzaa153.
5.   B. C. Blount, et al., Folate deficiency causes uracil misincorporation into human
DNA and chromosome breakage: implications for cancer and neuronal damage,
Proceedings of the National Academy of Sciences, 94 (7) (1997) 3290-3295.
6.   L. Martiniova, et al., Maternal dietary uridine causes, and deoxyuridine prevents,
neural tube closure defects in a mouse model of folate-responsive neural tube
defects, The American journal of clinical nutrition, 101 (4) (2015) 860-869.
7.   J. R. Alonzo, et al., The mitochondrial inner membrane protein MPV17 prevents
uracil accumulation in mitochondrial DNA, Journal of Biological Chemistry, 293
(52) (2018) 20285-20294.
8.   D. D. Anderson, C. M. Quintero, and P. J. Stover, Identification of a de novo
thymidylate biosynthesis pathway in mammalian mitochondria, Proceedings of
the National Academy of Sciences, 108 (37) (2011) 15163-15168.
9.   S. Spizzichino, et al., Cytosolic localization and in vitro assembly of human de
novo thymidylate synthesis complex, The FEBS Journal, 289 (6) (2022) 1625-
1649.
10.   D. Watkins, et al., Novel inborn error of folate metabolism: identification by
exome capture and sequencing of mutations in the MTHFD1 gene in a single
proband, Journal of medical genetics, 48 (9) (2011) 590-592.
11.   J. L. Fiddler, et al., Reduced Shmt2 expression impairs mitochondrial folate
accumulation and respiration, and leads to uracil accumulation in mouse
mitochondrial DNA, The Journal of Nutrition, 151 (10) (2021) 2882-2893.
12.   R. Sandhir, A. Halder, and A. Sunkaria, Mitochondria as a centrally positioned
hub in the innate immune response, Biochimica et Biophysica Acta (BBA)-
Molecular Basis of Disease, 1863 (5) (2017) 1090-1097.

13. Y. Song, Y. Zhou, and X. Zhou, The role of mitophagy in innate immune
responses triggered by mitochondrial stress, Cell Communication and signaling,
18 (2020) 1-14.
14. A. P. West and G. S. Shadel, Mitochondrial DNA in innate immune responses
and inflammatory pathology, Nature Reviews Immunology, 17 (6) (2017) 363-375.
15. W. Zheng, et al., How the innate immune DNA sensing cGAS-STING pathway is
involved in apoptosis, International Journal of Molecular Sciences, 24 (3) (2023)
3029.
16. M. J. White, et al., Apoptotic caspases suppress mtDNA-induced STING-
mediated type I IFN production, Cell, 159 (7) (2014) 1549-1562.
17. H. M. McBride, M. Neuspiel, and S. Wasiak, Mitochondria: more than just a
powerhouse, Current biology, 16 (14) (2006) R551-R560.
18. D. R. Green and J. C. Reed, Mitochondria and apoptosis, science, 281 (5381)
(1998) 1309-1312.
19. P. F. Chinnery and E. A. Schon, Mitochondria, Journal of Neurology,
Neurosurgery & Psychiatry, 74 (9) (2003) 1188-1199.
20. À. García-Cazorla, et al., Impairment of the mitochondrial one-carbon
metabolism enzyme SHMT2 causes a novel brain and heart developmental
syndrome, Acta neuropathologica, 140 (2020) 971-975.
21. O. Hashizume, et al., Corrigendum: Epigenetic regulation of the nuclear-coded
GCAT and SHMT2 genes confers human age-associated mitochondrial
respiration defects, Scientific Reports, 5 (2015).
22. J. Chakraborty and P. J. Stover, Deoxyuracil in DNA in health and disease, Curr
Opin Clin Nutr Metab Care, 23 (4) (2020) 247-252.
23. S. Srivastava and C. T. Moraes, Double-strand breaks of mouse muscle mtDNA
promote large deletions similar to multiple mtDNA deletions in humans, Human
molecular genetics, 14 (7) (2005) 893-902.
24. K. Shimada, et al., Oxidized mitochondrial DNA activates the NLRP3
inflammasome during apoptosis, Immunity, 36 (3) (2012) 401-414.
25. K. Thurairajah, G. D. Briggs, and Z. J. Balogh, The source of cell-free
mitochondrial DNA in trauma and potential therapeutic strategies, European
Journal of Trauma and Emergency Surgery, 44 (2018) 325-334.
26. C. Shu, X. Li, and P. Li, The mechanism of double-stranded DNA sensing through
the cGAS-STING pathway, Cytokine & growth factor reviews, 25 (6) (2014) 641-
648.
27. B. Zhao, et al., The molecular basis of tight nuclear tethering and inactivation of
cGAS, Nature, 587 (7835) (2020) 673-677.
28. J. Wu, et al., Cyclic GMP-AMP is an endogenous second messenger in innate
immune signaling by cytosolic DNA, Science, 339 (6121) (2013) 826-830.
29. B. Zhao, et al., A conserved PLPLRT/SD motif of STING mediates the
recruitment and activation of TBK1, Nature, 569 (7758) (2019) 718-722.
30. M. Yoneyama, W. Suhara, and T. Fujita, Control of IRF-3 activation by
phosphorylation, Journal of interferon & cytokine research, 22 (1) (2002) 73-76.
31. A. I. Dragan, et al., Mechanisms of activation of interferon regulator factor 3: the
role of C-terminal domain phosphorylation in IRF-3 dimerization and DNA binding,
Nucleic acids research, 35 (11) (2007) 3525-3534.

32.   X. Ning, et al., Apoptotic caspases suppress type I interferon production via the
395   cleavage of cGAS, MAVS, and IRF3, Molecular cell, 74 (1) (2019) 19-31. e7.
396   33.   J. L. Fiddler, et al., Impairments in SHMT2 expression or cellular folate
397   availability reduce oxidative phosphorylation and pyruvate kinase activity, Genes
398   & Nutrition, 18 (1) (2023) 5.

400   Figures

Figure 1. Decreased *Shmt2* expression leads to mitochondrial DNA leakage into

the cytosol without changing cellular mtDNA content in MEFs and HAP1 cells. (A)

Representative western blots of CS protein expression in subcellular fractionation of

cytosol and whole cell lysates. (B) Total mtDNA content and (C) cytosolic mtDNA

content in *Shmt2* MEF cells. (D) Representative western blots of CS protein expression

in subcellular fractionation of cytosol and whole cell lysates. (E) Total mtDNA content

and (F) cytosolic mtDNA content of Shmt2 wild-type and knock-out HAP1 cells cultured

in IMDM. (G) Cytosolic mtDNA content of wild-type HAP1 cells in folate modified media.

Densitometry was performed with Fiji software using GAPDH as a control. A Welch's t

test was performed to determine the statistical significance. AU, arbitrary units; CS,

citrate synthase; D-loop, displacement loop; GAPDH, glyceraldehyde-3-phosphate

dehydrogenase; Tert, telomerase reverse transcriptase. ns denotes statistically not

significant, * denotes $P < 0.05$, and ** denotes $P < 0.01$.

Figure 2. *Shmt2* heterozygosity-induced mtDNA leakage increases apoptosis in MEF cells but suppresses apoptosis in SHMT2 KO HAP1 cells. (A) Representative western blot of caspase-3 and cleaved caspase-3 and quantification graphs of (B) cleaved caspase-3 and (C) the ratio of cleaved caspase-3 to caspase-3 in *Shmt2*$^{+/+}$ and *Shmt2*$^{+/-}$ MEF cells. (D) Representative western blot of caspase-3 and cleaved caspase-3 and quantification graphs of (E) cleaved caspase-3 and (F) the ratio of cleaved caspase-3 to caspase-3 in WT or SHMT2 KO HAP1 cells. Densitometry was performed with Fiji software using GAPDH as a control. A Welch's t test was performed to determine the statistical significance. ns denotes statistically not significant and * denotes P < 0.05.

Figure 3. *Shmt2* loss activates cGAS/STING-mediated inflammation in *SHMT2* KO HAP1 cells. (A) Representative western blot of cGAS and STING and the quantification graphs of (B) cGAS and (C) STING in *Shmt2*$^{+/+}$ and *Shmt2*$^{+/-}$ MEF cells. As the samples are the same as those shown in Figure 2A, the GAPDH western blot image was reused in Figure 3A as a control. (D) Representative western blot of TBK1, p-TBK1, and STING and the quantification graphs of (E) STING and (F) the ratio of p-TBK1/TBK1 in wild-type and SHMT2 KO HAP1 cells. As the samples are the same as those shown in Figure 2D, the GAPDH western blot image was reused in Figure 3D as a control. Densitometry was performed with Fiji software using GAPDH as a control. A Welch's t test was performed to determine the statistical significance. cGAS, cyclic GMP-AMP synthase; STING, stimulator of interferon gene; TBK1, tank binding kinase 1; p-TBK1, phosphorylated tank binding kinase 1. ns denotes statistically not significant and * denotes P < 0.05.