1 **A rapid CAT transformation protocol and nuclear transgene expression tools for**

2 **metabolic engineering in *Cyanidioschyzon merolae* 10D**

3 Melany Villegas-Valencia[1], Martha R. Stark[2], Mark Seger[3], Gordon B. Wellman[1],
4 Sebastian Overmans[1], Peter J. Lammers[3], Stephen D. Rader[2], Kyle J. Lauersen[1,3]
5
6 [1]Bioengineering Program, Biological and Environmental Sciences and Engineering
7 Division, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-
8 6900, Kingdom of Saudi Arabia.
9
10 [2]Department of Chemistry and Biochemistry, University of Northern British Columbia,
11 Prince George, Canada.
12
13 [3]Arizona Center for Algae Technology and Innovation, Arizona State University, Mesa,
14 United States.
15
16 Corresponding author:
17 Kyle.lauersen@kaust.edu.sa
18
19

**Abstract**

The eukaryotic red alga *Cyanidioschyzon merolae* 10D is an emerging algal host for

synthetic biology and metabolic engineering. Its small nuclear genome (16.5 Mb; 4775

genes), low intron content (38), stable transgene expression, and capacity for

homologous recombination into its nuclear genome make it ideal for genetic and

metabolic engineering endeavors. Here, we present an optimized transformation and

selection protocol, which yields single chloramphenicol-resistant transformants in under

two weeks. Transformation dynamics and a synthetic modular plasmid toolkit are

reported, including several new fluorescent reporters. Techniques for fluorescence

reporter imaging and analysis at different scales are presented to facilitate high-

throughput screening of *C. merolae* transformants. We use this plasmid toolkit to

overexpress the *Ipomoea batatas* isoprene synthase and demonstrate the dynamics of

engineered volatile isoprene production during different light regimes using multi-port

headspace analysis coupled to parallel photobioreactors. This work seeks to promote *C.*

*merolae* as an algal system for metabolic engineering and future sustainable

biotechnological production.

**Keywords:** *Cyanidioschyzon merola*e 10D, polyextremophile, genetic engineering,

fluorescent reporters, transformation, isoprene.

**Abstract**

**Abbreviations**

BCA: Bicinchoninic Acid

CAT: Chloramphenicol transferase

CBB: Coomassie Brilliant Blue

cfu: Colony-forming units

CTP: Chloroplast targeting peptide

HR: Homologous recombination

IspS: Isoprene synthase

NTC: No template controls

PEG: Polyethylene glycol

qPCR: Quantitative PCR

SDS-PAGE: Sodium dodecyl sulfate-polyacrylamide gel electrophoresis

TSP: Total soluble protein

**Introduction**

The unicellular red alga *Cyanidioschyzon merolae* 10D was isolated from the volcanic Phlegrean fields near Naples, Italy as a haploid clone without a cell wall (Toda et al., 1995). This alga is considered to represent the simplest free-living eukaryote; composed of a nucleus, a single mitochondrion, one plastid, a simple ER, one Golgi body with two cisternae, few lysosomes, and one peroxisome (Miyagishima & Tanaka, 2021a). *C. merolae* contains a small nuclear genome and had one of the first telomere-telomere complete genome sequences of any model species, with ~16 Mbp; 4775 protein-coding genes, only 38 introns (Schärfen et al., 2022), and low genetic redundancy (Nozaki et al., 2007). Current genome-wide and multi-omics analysis in *C. merolae* has allowed investigation of the molecular mechanisms related to cell proliferation (Miyagishima et al., 2014; Suzuki et al., 1994), organelle development (Imoto & Yoshida, 2017; Misumi et al., 2005), and various metabolic pathways (Imamura et al., 2010, 2015; Mori et al., 2016; Moriyama, Sakurai, et al., 2014). Genetic manipulation of *C. merolae* 10D has been reported, including precise integration of transgenes in both nuclear (Fujiwara et al., 2019; Minoda et al., 2004; Sumiya et al., 2015) and chloroplast (Zienkiewicz et al., 2017) genomes via homologous recombination (HR) using 200–500 bp targeting sequences (Fujiwara et al., 2017; Takemura, Imamura, et al., 2019). In addition, transgenes are stably expressed, and there appears to be no endogenous silencing of transgenes (Watanabe et al., 2014; Zienkiewicz et al., 2019).

The biomass of *C. merolae* 10D is protein-rich (Villegas-Valencia et al., 2023) and contains valuable bioproducts like heat-stable phycocyanin, carotenoids, and β-glucan (Lang et al., 2022; Rahman et al., 2017; Villegas-Valencia et al., 2023). Because the cells lack a rigid cell wall and are easily disrupted, the introduction of DNA and extraction of intracellular contents is straightforward (Miyagishima & Tanaka, 2021b). The *C. merolae* 10D strain can be cultivated in acidified medium (pH 1–3) and elevated temperatures (42-46 ⁰C) in freshwater and seawater to high cell densities in both lab-scale and outdoor conditions with minimal contamination (Hirooka et al., 2020; Villegas-Valencia et al., 2023). These features suggest that *C. merolae* could be a valuable and scalable platform for metabolic engineering or other recombinant bioproduct accumulation.

Further advances in metabolically engineering this alga are of interest to add value to the
algal biomass. There is a growing number of engineered traits in *C. merolae*, including
the enhanced generation of triacylglycerol (TAG) without growth inhibition (Sumiya et al.,
2015), the incorporation of a plasma membrane sugar transporter from its relative
*Galdieria sulphuraria* to enable heterotrophic growth in the presence of exogenous
glucose in the dark (Fujiwara et al., 2019), and recently, our demonstration of the
production of the non-native ketocarotenoids canthaxanthin and astaxanthin (Seger et
al., 2023). Although various genetic tools have been developed in *C. merolae* 10D,
current methods for its transformation and standard screening of transformants can take
several weeks (Fujiwara et al., 2017; Fujiwara & Ohnuma, 2017; Zienkiewicz et al., 2019).
Metabolic engineering attempts in this host are still in their infancy and the development
of techniques to achieve mature genetic engineering concepts are still needed. This work
seeks to create a user-friendly and standardized transformation approach and we present
new modular genetic tools for *C. merolae* nuclear genome transgene expression for
metabolic engineering and recombinant product accumulation.
Here, we report an optimized transformation and selection protocol for *C. merolae* 10D,
which yields single chloramphenicol-resistant colonies in under two weeks and enables
phenotypic screening within three. We present an *in silico* designed and synthetically
constructed modular plasmid toolkit which includes several fluorescent reporters. We
show its transformation, the metrics of successful homologous recombination, and
techniques for fluorescence imaging for the high-throughput screening of transformants.
The transgene expression dynamics from the nuclear genome including total soluble
recombinant protein accumulation and a first demonstration of metabolic engineering
volatile isoprene production from the alga are also presented. This work will serve as a
foundation of open-source molecular tools for the *C. merolae* research community and
indicates it could be a reliable chassis for further engineering concepts.
126
127

**Materials and Methods**

**2.1. Algae culture**

*C. merolae* 10D strain (wild-type; NIES-3377) was obtained from the National Institute of Environmental Studies' microbial culture collection in Japan. *C. merolae* culture and its transgenic lines were routinely maintained in MA2G medium, a modified version of MA2 (Minoda et al., 2004). This medium is a version of Modified Allen's medium (MA, (Minoda et al., 2004)) wherein several components have doubled concentrations. MA2G consists of 40 mM $(NH_4)_2SO_4$, 8.0 mM $KH_2PO_4$, 4.0 mM $MgSO_4$, 1.0 mM $CaCl_2.2H_2O$, 100 µM $FeCl_3.6H_2O$, 80 µM $Na_2EDTA.2H_2O$, 92.2 µM $H_3BO_3$, 18.2 µM $MnCl_2-4H_2O$, 1.54 µM $ZnCl_2$, 3.2 µM $Na_2MoO_4.2H_2O$, 0.34 µM $CoCl_2.6H_2O$, 0.64 µM $CuCl_2-2H_2O$, and 50 mM glycerol. $H_2SO_4$ was used to adjust the pH to 2.5. Glycerol was added as it encourages slightly faster growth of the Cyanidiophyceae and has been shown to encourage respiration (Moriyama et al., 2015). Working stocks of cultures were maintained on corn starch beds on MA2G Gellan gum plates, and 50 mL liquid cultures were shaken at 100 rpm in 250 mL Erlenmeyer flasks with 0.22 µm filter vented caps to allow gas exchange. Cultures were maintained under continuous illumination (90–130 µE) at 42 ºC in a Percival incubator (AL-30, Percival Scientific) supplemented with 2– 4% $CO_2$. Wild-type and transgenic cultures were cryopreserved in 8% DMSO and stored at -80 ºC for long-term preservation.

**2.2. Plasmid construction**

Plasmids were designed to integrate the selectable marker chloramphenicol acetyltransferase (CAT), a fluorophore (one of mTagBFP2 (Subach et al., 2011), mTFP1 (Ai et al., 2006), Clover (Lam et al., 2012), mVenus (Kremers et al., 2006), LSSmOrange (Tsutsui et al., 2008), mKOk (Shcherbakova et al., 2012) , mScarlet (Bindels et al., 2017)), and isoprene synthase (IspS) into the intergenic region between the nuclear glycogen phosphorylase (CMD184C) and TATA-box binding protein-associated factor 13 (CMD185C) genes on *C. merolae* chromosome 4 ("HR-L" and "HR-R" arms in Figures 1a, 2b, 3a and 4a) via homologous recombination (HR (Fujiwara et al., 2013)). Gene elements used in our *in silico* genetic designs are listed in Supplemental File S1 and all plasmids can be obtained from the authors or with permission from Genscript.

Endogenous elements such as promoters, terminators, and homology arms were taken
from the reference genome of *C. merolae* 10D (Fujiwara et al., 2013, 2017, 2019;
Moriyama, Tajima, et al., 2014). The *Staphylococcus aureus* chloramphenicol
acetyltransferase (CAT) (NCBI: M58516.1 (Schwarz & Cardoso, 1991)) was used as a
selection marker and *Ipomoea batatas* isoprene synthase (*Ib*IspS, NCBI: AZW07551.1
(Ilmén et al., 2015)) for isoprene production. Fluorophore sequences were taken from the
sources indicated in Figure 3b. Codon optimization after back translation of amino acid
sequences was conducted for the most frequent codon usage for the algal nuclear
genome using *C. merolae's* codon usage table found in the Kazusa database
(https://www.kazusa.or.jp/codon/cgi-bin/showcodon.cgi?species=280699). For all genes,
native targeting peptides were removed from the amino acid sequence before
optimization. Gene synthesis and subcloning were carried out by Genscript (Piscataway,
NJ, USA), and plasmids were delivered as lyophilized DNA, transformed, and preserved
in *Escherichia coli* DH5α with ampicillin as the selection agent in lysogeny broth (LB). Full
annotated sequences of all plasmid elements can be found in Supplemental File S2.
174

**2.3. *C. merolae* 10D transformation**
The transformation procedure was similar to the ones reported before (Fujiwara et al.,
2017; Ohnuma et al., 2008, 2014) with slight modifications, as follows: 1 pmol of linear
DNA was used in transformations, which was prepared by PCR amplification using
primers targeting the homologous arms with a high-fidelity polymerase (Figure 2A, primer
set 1). PCR products (50 µL reactions) were purified using a PCR clean-up kit (ZR-96
DNA Clean & Concentrator) and re-suspended in DNase and RNase-free water. DNA
concentration and purity were measured on a NanoDrop One spectrophotometer
(Thermo Fisher Scientific, UK).
To prepare for transformation, cell cultures were maintained in MA2G (MA2 + 50 mM
glycerol) medium under continuous light at 90–130 µE, 42 ºC and constantly
supplemented with 2–4% $CO_2$. The doubling time under these conditions is ~9–10 h for
wild-type cells. Wild-type cell cultures were diluted 2–3 days before transformation so that
a culture of actively dividing cells with an $OD_{740nm}$ lower or equal to 3.0 was ready prior to
transformation ("transformation stock"). One day before transformation, the stock culture

was diluted to an $OD_{740nm}$ of ~0.2 in 50 mL MA2G and grown under the same conditions

as above until the $OD_{740nm}$ of the culture reached ~0.8–1.0. On the day of transformation,

polyethylene glycol (PEG)-4000 was prepared (60% w/v) by combining 0.9 g PEG4000

(A16151.30, ThermoFisher Scientific) and 750 μL MA-I (note: same concentrations of

MA2 except 20 mM $(NH_4)_2SO_4$, 2 mM $MgSO_4$ + metals, pH adjusted to 2.5) in a 2 mL

Eppendorf tube and dissolved at 42 ℃ with occasional inversion. Next, 50 mL of

transformation culture was centrifuged (10 min at 2000 x g), the supernatant filter

sterilized to make conditioned media ('MA2G-C'), and cells washed with 1 mL of MA-I

medium kept at 42 ℃, and then transferred to a 1.5 mL tube, centrifuged again (1 min at

2000 x $g$) and resuspended in ~100-150 μL warm MA-I to a total final volume of 200 μL

(250x concentrated). In a 1.5 mL tube, ~1 pmol of linear- (PCR product) or circular-DNA

was diluted in water to a total volume of 84 μL and combined with 10 μL 10XMA medium

(to bring the transformation mixture to 1XMA-I concentration), and 6 μL 10 mg/mL

UltraPure Salmon Sperm DNA Solution (Invitrogen, USA). Salmon sperm DNA was

denatured by heating to 98 ℃ for 5 min and then rapidly cooled on ice before adding.

Next, 25 μL of concentrated cells were added to the DNA mixture. A no-DNA control

reaction was also prepared as a negative control. Then, one sample at a time, 125 μL of

PEG4000 solution was added to the reaction and mixed quickly by flicking the wrist 8-10

times so that the PEG and the cells/DNA were completely mixed. 1 mL of warm MA2G

was immediately added to the tube, which was then poured into 50 mL of warm MA2G in

a 250 mL vented flask and allowed to recover while shaking at 100 rpm for 48 h under

the same cultivation conditions as described above.

After recovery, transformed cells were collected by centrifugation and resuspended in 2.3

mL MA2G-C, a medium in which cells have previously been grown to an $OD_{740nm}$ 0.8-1.0,

and has been filter sterilized. MA2G-C was used to dilute cornstarch and cells for plating

after transformations. This medium was prepared by growing wild-type *C. merolae* cells

in MA2G to a culture $OD_{740nm}$ ~1.0, centrifuging the cultures, and using the 0.2 μm filtered

supernatant. We have observed this medium from *C. merolae* cells in the exponential

growth phase accelerates the formation of single colonies on starch beds. Cell

suspensions were then serially diluted (1:9, 1:27, 1:81, and 1:243) in a 96-well plate and

10 μL pipetted onto freshly prepared starch beds on 0.77X MA2G Gellan gum plates

(0.46% Gellan gum; 120mm x 120mm x 17 mm square petri dishes) containing 250 µg/mL

chloramphenicol. Based on our observations, colonies tend to come up faster on 0.77X

MA2G Gellan gum plates than on 1X. This dilution series should result in at least one

series of starch beds that contains uncrowded colonies appropriate for isolation (~1-

10/spot) across a wide range of transformation efficiencies. The plates are prepared the

day before use with 20% (v/v) cornstarch slurry beds (Kobayashi et al., 2010), as shown

in Figure 1B. Before spotting the starch, MA2G Gellan gum plates were left open in the

incubator for 20 min so that spots that do not run into each other are formed. Briefly, 20%

cornstarch slurry was prepared from cornstarch that was washed 3X with sterile $H_2O$,

resuspended to 50% (v/v) in 75% EtOH, and kept at 4 ºC until further processing. For

plating, 20% cornstarch was prepared by taking 10 mL from the 50% cornstarch/EtOH

stock, centrifuging briefly, washing the pellet 3X with MA2G-C medium, then

resuspending to a final volume of 25 mL in MA2G-C. This was poured into a Universal

Reagent Reservoir and manually agitated to avoid settling of the starch while pipetting 15

µL spots onto the plates using a multichannel pipette. Chloramphenicol was only added

to the Gellan gum + MA2G solid medium, and the cornstarch was antibiotic-free.

Plates with approximately 144 cornstarch spots were inoculated with serially diluted

transformants (10 µL per spot), along with 6 spots of nurse cells (Kobayashi et al., 2010)

throughout the plate (chloramphenicol resistant and actively dividing cells that may

encourage neighboring colony growth; Figure 1B). Nurse cells are spotted on top of a few

transformants for ease of plating with the multichannel pipette. Spots were allowed to dry

and plates were incubated in $CO_2$-supplied Percival incubators under previously

described conditions until colony formation appeared ~6–10 d after plating. Colonies were

then picked and resuspended in 15 µL MA2G, and this liquid was used to re-inoculate

cornstarch beds for long-term storage and maintenance. Picked liquid samples can also

be grown for ~3–4 d (Figure 1b) and used in further analysis. Isolates were then scaled

up in 1 mL of liquid MA2G medium in 24-well plates until dense ($OD_{740nm}$ ~1.0; after 3 d),

and used in plate-level fluorescence, in-gel fluorescence, and flow cytometry assays to

confirm the expression of fluorescent reporters.

## 2.4. Transformation efficiencies

Constructs containing mVenus and CAT were transformed into *C. merolae* to determine transformation efficiencies of integrated linear DNA and circular plasmids. The DNA concentration of each was normalized to 1 pmol. Transformations were carried out in triplicate as previously described. 10 days after plating, colony-forming units (cfu) were counted using images of the plates and a software for biological-image analysis (Fiji; Schindelin et al., 2012). Transformation efficiencies were calculated based on cfu from the dilution series on each plate (Supplemental File S3), using the following formula:

$$Efficiency = \frac{Number\ of\ transformants}{pmol\ of\ DNA} \times \frac{Final\ vol.\ at\ recovery\ (mL)}{vol.\ plated\ (mL)}$$

## 2.5. DNA extraction and molecular screening

*C. merolae* strains were harvested at mid-log phase by centrifugation (10 min at 2000 x *g*) and total genomic DNA was extracted from algal pellets using a Quick-DNA Fungal/Bacterial Miniprep Kit (Zymo Research, USA) and a FastPrep-24 5 g bead beating grinder and lysis system (MP Biomedicals, USA) according to the manufacturer's protocol. DNA extracts were quantified using a NanoDrop One spectrophotometer (Thermo Fisher Scientific, UK).

Q5 High-Fidelity DNA Polymerase (New England BioLabs, UK) and Phusion Green Hot Start II High-Fidelity PCR Master Mix (Thermo Fisher Scientific, Lithuania) were used for PCR according to the manufacturer's protocols (Table 1). All primers used in this study were synthesized by IDT (Integrated DNA Technologies Inc., Belgium). Primer set 1 was used to linearize target constructs from plasmids before transformation, whereas primer sets 2 and 3 were used to screen transformants for the presence of the insert at the target neutral site and to verify the positive integration of the selectable marker (CAT), respectively.

Quantitative PCR (qPCR) was carried out using a CFX96 Real-Time PCR Detection System to determine the copy number of the integrated mVenus. The copy number of heterologous mVenus in each strain was normalized with that of native 60S rDNA (NCBI: 16997147) as the reference gene. To determine the optimal annealing temperature of the

primers, a thermal gradient across 7 different temperatures ranging from 59 ⁰C to 69 ⁰C
was used for each primer set in a T100 Thermal Cycler (Bio-Rad). 5 ng of DNA was added
to each well in 20 µL reactions and a no template control (NTC) consisting of $H_2O$ was
also included. SsoAdvanced Universal SYBR Green Supermix (Bio-Rad) was used for
qPCR according to the manufacturer's protocol. Standard curves were constructed using
serially diluted DNA (1/5, 1/25, 1/125, 1/625, and 1/3125; Supplemental Figure 3) isolated
from a mVenus-expressing *C. merolae* strain and the relevant set of primers. For each
primer pair, technical triplicates were done for each dilution factor. An NTC was also
included for each primer set. PCR and qPCR conditions are shown in Supplemental File
S4.
In addition, a semi-quantitative assessment of mVenus accumulation was carried out in
selected transformants. The total soluble protein (TSP) fraction of the algal cells was
quantified by the bicinchoninic acid (BCA) assay using bovine serum albumin (BSA) as a
standard of known concentrations. Samples of TSP extracted from mVenus-expressing
transformants, along with a dilution series of *E. coli* produced and StrepTrap (Cytiva
StrepTrap XT, Sigma-Aldrich, Germany) column chromatography purified mVenus were
analyzed under non-denaturing sodium dodecyl sulfate-polyacrylamide gel
electrophoresis (SDS-PAGE) conditions (5% stacking, 15% resolving), followed by
fluorescence imaging using excitation wavelengths for mVenus (Figure 2E).
301

**2.6. Fluorescence imaging**
*C. merolae* transformants cultivated in 24-well plates in 1 mL MA2G were prepared for
screening analysis during the mid-log phase. The presence of fluorescent proteins in
selected transformants was observed using a ChemStudio PLUS (Analytik Jena, USA)
gel documentation system with an eLite xenon lamp and filter wheel extension as
previously described (Gutiérrez et al., 2022). Different filters with specific bandpass
ranges that allowed the selective excitation and emission of different fluorophores were
employed. Plate-level fluorescence was carried out by spotting 10 µL of selected
transformants onto a Gellan gum plate with amido black (150 mg/L). The latter was
employed to reduce background fluorescence (Wichmann et al., 2018) and it was added

to the Gellan gum-MA2 mix by dissolving into the medium before autoclaving. The
excitation/emission filters used for each fluorophore are stated in Figure 3D.
For in-gel fluorescence analysis, 1 mL of each sample at mid-log phase was centrifuged
(10 min at 2000 x $g$), the supernatant discarded, and the pellet was either snap-frozen in
liquid nitrogen and stored at -80 ⁰C for future analysis, or resuspended in ~150–200 µL
of sample buffer (0.2 M SDS, 0.3 M Tris, 30% glycerol, 0.02% bromophenol blue),
vortexed and then incubated at 40 ⁰C for 5 min on a heat block and quickly placed on ice
until loaded on an SDS-PAGE gel. 15 µL of each sample was loaded along with 4 µL of
PageRuler Plus Prestained Protein Ladder (Thermo Fisher Scientific, UK). After
electrophoresis separation, gels were visualized in the ChemStudio PLUS using the filter
combinations listed in Figure 3E. 504/10 nm excitation and no emission filter, was used
to visualize the pre-stained protein ladder. Lastly, Coomassie Brilliant Blue (CBB) or
Ponceau S staining of the protein gels was done to compare sample loading. All filter
settings for the fluorescent screening of transformants on the Analytik Jena are provided
in Supplemental File S5.
327

**2.7. Flow cytometry**
Flow cytometry of wild-type and fluorescent protein-expressing cells was performed using
an Invitrogen Attune NxT flow cytometer (Thermo Fisher Scientific, UK) equipped with a
488 nm blue excitation laser for forward scatter (FSC), used to measure size, and with
lasers for excitation at wavelengths 405 nm (Violet, VL), 488 nm (Blue, BL), 561 nm
(Yellow, YL), and 638 nm (Red, RL), along with corresponding sets of emission filters
supplied by the manufacturer. For dual population analysis, transformants and wild-type
cells were grown in a 24-well plate as shown in Fig. 1b until they reached the mid-log
phase. Each sample was diluted 1:100 with 0.9% (w/v) NaCl solution and 500 µL of each
transformant and 500 µL of wild-type cells were combined in a 2 mL Eppendorf and
measured through the sample injection port (100 µL sample/min). The first 15 µL of the
combined sample was not recorded to ensure a stable cell flow rate during measurement.
The Attune NxT Software v3.2.1 (Life Technologies, USA) was used to group cell
populations and conduct further post-acquisition analyses.
342

**2.8. Quantification of heterologous isoprene biosynthesis**

To investigate the overexpression of the *I. batatas* isoprene synthase in *C. merolae* and the consequent production of isoprene, a transformant line was cultivated in 400 mL Algem photobioreactors (Algenuity, UK) along with the wild-type 10D and an mVenus-expressing strain as negative controls. Strains were cultivated under different light regimes at 42 °C and 120 rpm agitation. Preculturing before growth and isoprene analysis was performed by inoculation of cells to Algem flasks with MA2G medium for 96h, then, each strain was resuspended in the same medium to an $OD_{740nm}$ ~0.5 at the start of the experiment. Cell lines were grown in biological duplicates under constant illumination at 1200 μE or 750 μE, or under 12 h:12 h light: dark light cycling with the same light intensities.

Volatile isoprene was monitored during growth using a real-time headspace gas analysis system equipped with a triple filter mass spectrometer and multi-port inlet (Hiden Analytical HPR-20 R&D, UK). All cultures were continuously sparged with a 3% $CO_2$ in air mix (25 mL/min), and the off-gas of each reactor was redirected through separate gas lines, first to a 250 mL bottle containing 80 g of $CaCl_2$ used as a desiccant. Then, the dried gas was further directed to a 20-port inlet of the online headspace gas analysis system. The gas composition of each flask's headspace was analyzed with a rotation to the next gas stream occurring every ~3.5 mins. Isoprene was quantified by monitoring atomic mass units (amu) 67, 68, 53, 39, 40, 41, and 27; any overlapping amu of other gases present ($O_2$, $CO_2$, $N_2$, Ar) were automatically deconvoluted by the Hiden Analytical QGA Software (version 2.0.8).

The amount of isoprene detected in the headspace of each culture was integrated over time, and the isoprene output (in ppm) reported by the instrument was corrected by comparison against a standard curve (Supplemental File S6) generated with flasks that were also kept in the Algems but contained different concentrations of loaded pure isoprene standard. In total, 10 control flasks were set up to cover five concentrations (10, 5, 2.5, 1.25, and 0.625 mg of isoprene) in duplicate. The temperature of the standard flasks was gradually increased from 20 °C to 38 °C over 1 h and then held for 10 h, during which time the isoprene standard in the headspace was quantified. The integrated isoprene amounts were plotted against the amounts added to each flask to generate the

374   standard curve. The Hiden Analytical QGA Software was employed for real-time gas
375   analysis until the end of algal cultivations (Figure 4E).
376

**2.9. Statistical analysis**
Data analysis was conducted using both biological and technical replicates. Student $t$-test
was carried out to determine if mean transformation efficiencies were significantly
different ($p < 0.05$). Mean values and standard error (SEM) of replicates were calculated
using the software JIMP Pro 16.2 (SAS Institute Inc., Cary, NC), and are illustrated in
relevant graphs.
383

**Results & Discussion**

*C. merolae* 10D exhibits interesting properties like a small genome, ease of cell lysis, homologous recombination, and stable transgene expression that warrant its consideration as an emerging alga for biotechnology concepts. Here, we sought to develop a protocol that can more rapidly generate transgenic strains, characterize the behavior of integrated DNA, and present a useful molecular toolset for engineering its nuclear genome. We demonstrate the functionality of these tools in the proof-of-concept engineering of *C. merolae* 10D to produce volatile isoprene for the first time and characterize the dynamic nature of this production.

**3.1.    Development of a rapid chloramphenicol transformation protocol**

A rapid transformation protocol for *C. merolae* 10D was developed, which generates single chloramphenicol-resistant colonies within 10 days (Figure 1). We employed a two-expression-cassette system which we designed *de novo* for this study (Figure 1A). The *APCC* and *CPCC* promoters were used to drive transgene expression as they produce high transcriptional activity under illuminated conditions  (Fujiwara et al., 2019). *CPCC* was chosen for the gene of interest cassette as this promoter exhibited higher recombinant protein accumulation than the *APCC* in a previous study (Fujiwara et al., 2019). The *APCC* promoter was used to express the selectable marker CAT. Homologous recombination for transgene integration was directed by 550 bp sequences (Figure 2A,B) targeting the CMD184C–185C locus found on *C. merolae* 10D chromosome 4 (Fujiwara et al., 2017).

The gene expression cassettes were amplified by PCR and used in the PEG-mediated transformation of *C. merolae* 10D (Figure 1A-B)*.* The alga can take up foreign DNA without a physical agitation agent like the glass beads used for cell wall deficient *Chlamydomonas reinhardtii* (Kindle, 1990). Salmon sperm DNA was used in the transformation mix along with 1 pmol of linearized transgene cassette DNA. Salmon sperm DNA is denatured before use and is typically employed as a DNA carrier to improve transformation efficiency in yeast (Longmuir et al., 2019) and marine microalgae (Zhang & Hu, 2014). The recovery of transformants as single colonies requires selection on 20% cornstarch slurry spots on freshly prepared chloramphenicol-containing MA2-G plates.

Compared to the top-starch method (Takemura, Kobayashi, et al., 2019), this is a more

labor-intensive protocol, as the plates must be prepared fresh; cornstarch slurry spotted,

and cells serially diluted before inoculation. However, the use of a 3X serial dilution of

cells before plating allows for balancing the culture density to enable single colonies to

appear without overcrowding. Once colonies are visible, they can be directly picked into

liquid MA2G medium to later inoculate a maintenance starch spot for long-term storage,

and the liquid culture can be used directly for further scale-up and screening analysis

(Figure 1B). Following this new experimental pipeline, the time required to generate

transformants is reasonable for conducting iterative engineering cycles required for most

biotechnology concepts.

**Figure 1 (next page).** Plasmid design, PEG4000-mediated *C. merolae* 10D transformation and
transformation efficiencies. **A –** Synthetic plasmids were designed *in silico* and constructed *de*
*novo* for integration into the 184C-185C neutral site on *C. merolae* chromosome 4. The
expression cassette was designed to be modular, with a separate gene of interest and a
selectable marker. Each element was separated by unique restriction endonuclease sites as
illustrated. pCPCC – phycocyanin-associated rod linker protein promoter, CTP CMH166C – DNA
Gyrase B chloroplast targeting peptide, mVenus – yellow fluorescent protein reporter, StrepII –
C-terminal peptide tag with stop codon, NOS – nopaline synthase terminator, pAPCC –
allophycocyanin-associated rod linker protein promoter, CTP CMO250C – allophycocyanin-
associated rod linker protein chloroplast targeting peptide, CAT – chloramphenicol
acetyltransferase, FLAG – peptide tag with stop codon, β-tub – *C. merolae* β-tubulin terminator
CMN263C. **B** – Workflow of the optimized transformation protocol in *C. merolae* 10D to obtain
positive transformants ~1.5 weeks after selective agent plating. **C+D** – Circular vs. linear DNA
transformation. Representative transformation plates ~ 10 days after inoculating transformants in
cornstarch beds. A dilution series of cells transformed with circular (C) and linearized (D) DNA
were spotted along with untransformed wild-type cells (WT; last row) and nurse cells (see
methods). **E** – Transformation efficiencies compared between linear to circular DNA. *** indicates
results of Student's *t*-test were significantly different ($p < 0.0001$) when comparing the mean
number of transformants between linear and episomal transformation.

**A**

| Primer set | Target | Primer sequence (5' to 3') | Product size (bp) | Tm (℃) | Type |
|---|---|---|---|---|---|
| 1 | D184C-D185C | F: CTCATCATCGGCACTATGGACGG R: ATACGCCAAGCTCGCCGTG | 4550 | 69 | Plasmid linearization |
| 2 | D184C-D185C | F:GTTGCCTCAAGAAGTTGACC R: TGGCAGCATCAGCACTATTA | 4397 | 68 | Genotyping PCR |
| 3 | CAT | F:GGCGTTCATCTTCCTGGTGA R: TCGAGTTCATGAACAGGCCC | 418 | 68 | Genotyping PCR |
| 4 | mVenus | F:GGACGACGGCAACTACAAGA R:GCCGTGATGTACACGTTGTG | 119 | 60 | qPCR |
| - | 60S | F:AAGTTTCGCTGTACGCTTGG R:TAACCAGGACCATATCGCCG | 155 | 60 | qPCR |

**B**

**C**

**D**

**Figure 2.** Molecular screening of mVenus-expressing transformants. **A** – Primer sets used in this study to linearize all plasmids before transformation (1), screen for integration at the desired locus and presence of selectable marker (2 and 3, respectively), and estimate the copy number of the gene of interest (4) by quantitative PCR (qPCR). **B** – Schematic diagram of mVenus and CAT insertion into the neutral site between CMD184C and CMD185C by homologous recombination. Introduced linear DNA vector (top) and the genomic structure of the parental (P) wild-type (WT) strain (bottom). Arrowheads indicate the position of the primers used to screen for total insert (~ 5.0 kb; orange) and the insertion of the CAT gene (~ 0.5 kb; blue), as shown in C. **C** – Polymerase chain reaction (PCR) confirmation of linearized plasmid integration at the D184-185 neutral locus (insert) and presence of transgene (CAT) in 24 transformants. The predicted size of the parental strain (or off-target insertion) and correctly integrated PCR products is 0.9 kb and 4.4 kb, respectively. The asterisks (*) above the lane numbers show the selected transformants used in further qPCR and TSP analysis. PCR was followed by plate-level fluorescence analysis of the mVenus transformants on Gellan gum plates stained with amido black. Chlorophyll *a* emission (red) and mVenus fluorescent signal (yellow), as well as in-gel fluorescence, were measured to assess the expression level of mVenus in 24 transformants. The expected molecular weight of mVenus is ~26 kDa. CBB: Coomassie brilliant blue stain is included as a loading control. L: DNA / protein ladder, P: parental strain. Full-length images can be found in Supplemental Figure 1. **D** – Quantitative-PCR analysis of selected transformants to estimate the copy number of the mVenus gene. Transformants 4, 5, 23, and 24 exhibit medium-mVenus-expression with correct integration at locus (MedExp-Int); 20 and 21: high-mVenus-expression, correct integration at locus (HighExp-Int); 7 and 8: high-mVenus-expression and integration outside of target locus (HighExp-NInt). The value of each transformant was normalized against the value from the 60S rDNA gene reference gene to estimate their copy number (n=3). **E** – Percent of total soluble protein (%TSP) analysis of recombinant mVenus expression in transformants measured by in-gel fluorescence. The cell number of the transformants was normalized before extraction and gel loading. A dilution series of purified mVenus from *E. coli* was included for semiquantitative assessment. CBB is shown as a loading control.

Both linear and circular plasmid constructs containing mVenus were transformed into *C. merolae* 10D to determine differences in transformation efficiencies and potential transgene expression differences. It has been recently shown in the red alga *Porphyridium purpureum* that the bacterial origin of replication could confer autonomous replication of circular, episomal, plasmids within cells and consequent higher recombinant protein titers (Hammel et al., 2024; Li & Bock, 2018). The efficiency of transforming PCR linearized constructs in *C. merolae* 10D was 18-fold higher than in the episomal transformation ($p < 0.0001$; Figure 1E). Episomal transformation was carried out with the same workflow as linear, but the plasmid was not linearized before the transformation. The number of colony-forming units in the linear and circular transformation plates was counted 10 days after plating. Only a few colonies were recovered when the cells were transformed with a circular plasmid.

Efficient gene targeting and stable transgene expression represent two advantages of *C.*

*merolae* 10D as a host for synthetic biology and metabolic engineering. The intergenic

region within the CMD184C-CMD185C loci was selected as the target site, as it has been

previously used successfully (Fujiwara et al., 2013, 2017; Fujiwara & Ohnuma, 2017).

The occurrence of HR at this locus was examined by PCR. Integration of the target binary

cassette carrying mVenus and CAT was observed in 22 out of 24 transformants (92%;

Figure 2C). Previously, targeted integration of the CAT selection marker was observed in

24 out of 33 transformants (73%), when the transgene was flanked with 500-bp homology

arms and the cells were selected in a liquid medium with 200 µg/mL chloramphenicol for

10 days and then plated for another 2 weeks (Fujiwara et al., 2017). In our efforts to obtain

transformants more rapidly, direct plating on 250 µg/mL chloramphenicol after 48 h

recovery was employed. This modification was able to speed the CAT transformation up

by two weeks. Two transformants (7 and 8) presented the same amplicon size as the

wild-type (~ 1 kb) indicating integration outside the intended region. Although not directly

integrated at the target locus, these transformants exhibited mVenus fluorescence (Figure

2C). Both at the agar plate and in-gel fluorescence levels, the mVenus reporter was

detected in all 24 transformants. Across the clones, a consistent mVenus signal was

observed, except for those with non-targeted integration (7 and 8) and clones 20 and 21

(Figure 2C). These transformants, as well as properly-integrated and moderate mVenus-

expressing colonies 4, 5, 23, and 24, were selected for assessment of gene copy number

by qPCR. All clones with higher fluorescence than the average exhibited 2 or more copies

of the transgene construct. Colonies 8 and 20, with the highest mVenus fluorescence,

appear to have 30 and 20 copies of the transgene in their genomes respectively. This is

similar to a previous report of multicopy insertion wherein tandemly repeated insertion of

the introduced fragments was observed (Fujiwara et al., 2013). A semi-quantitative

assessment of mVenus accumulation was also carried out in these eight transformants.

The TSP fraction of the transformants was quantified by the BCA test and compared

against a dilution series of purified mVenus using an SDS-PAGE gel and fluorescence

imaging. Recombinant protein levels accumulated to between 0.3-0.6% of total soluble

protein (Figure 2E). The copy number of gene insertion likely influences the levels of

recombinant protein within the cell but does not match the levels reported for *P.*

*purpureum* episomal transgene expression (i.e. up to 5% of the total soluble protein (Li &
Bock, 2018)).

**3.2    A suite of new fluorescent reporters for *C. merolae***

New fluorescent reporters applied in *C. merolae* can enable combinations for protein
interaction studies, or as fusion partners to target recombinant proteins. We designed
codon-optimized transgenes for the expression of fluorescent reporters mTagBFP2,
mTFP1, Clover, mVenus, mKOk, LSSmOrange, and mScarlet (Figure 3A-B). mVenus
and mScarlet have been previously used in this host (Fujiwara et al., 2021; Seger et al.,
2023), however, the addition of broader spectral tools can expand applications in this
host. Here, stable expression of these fluorescent reporters was demonstrated using the
same transformation and recovery protocols as outlined above (Figure 3C-E). Cell
populations for each reporter could be distinguished from that of the wild-type cells in flow
cytometry analyses and represent a sensitive analytic technique where fluorophores
could be multiplexed (Figure 3C). Additionally, visualization of transformants and their
expression at the agar plate level and in protein gels can aid gene expression assessment
and transformant status. We compared *C. merolae* with our previous efforts in practical
fluorescence imaging developed for the green alga *C. reinhardtii* (Figure 3D-E, as
reported in (Gutiérrez et al., 2022)). We applied different bandpass filters in a modified
gel doc system to visualize pre-selected transformants either in agar plates or protein gels
and show that their signal can be visualized separately from chlorophyll *a* and
phycocyanin autofluorescence (Gutiérrez et al., 2022). The emissions of mTagBFP2,
mVenus, mKOk, and LSSmOrange were visible without spectral overlap, and to a lesser
extent mTFP1, mVenus, and mScarlet could be combined in this analysis (Figure 3D-E
and Supplemental File S5). The ability to conduct targeted HR into the *C. merolae*
genome means fewer transformants must be screened compared to efforts in green algae
(Abdallah et al., 2022; Gutiérrez et al., 2024) and the ability to rapidly detect reporter
expression is a practical asset for analyzing trangene expression.

**A**

**B**

| Fluorophore | MW (kDa) | MW + CTP | Source | FPbase ID | Ex λ | Em λ | Reference |
|---|---|---|---|---|---|---|---|
| mTagBFP2 (B) | 26.7 | 36.0 | *Entacmaea quadricolor* | ZO7NN | 399 | 454 | Subach *et al.*, 2011 |
| mTFP1 (T) | 26.9 | 36.3 | *Clavularia* sp. | ASSY9 | 462 | 492 | Ai *et al.,* 2006 |
| Clover | 26.8 | 34.9 | *Aequorea victoria* | 4Z641 | 505 | 515 | Lam *et al.*, 2012 |
| mVenus (V) | 26.9 | 36.2 | *Aequorea victoria* | WCSN6 | 515 | 527 | Kremers *et al.*, 2006 |
| mKOk (KO) | 24.5 | 33.8 | *Verrillofungia concinna* | HMK8R | 551 | 563 | Tsutsui *et al.*, 2008 |
| LSSmOrange (LSS) | 26.7 | 36.1 | *Discosoma* sp. | OVJRJ | 437 | 572 | Shcherbakova *et al.*, 2012 |
| mScarlet (S) | 26.4 | 35.7 | Synthetic | FVS3D | 569 | 594 | Bindels *et al.*, 2017 |

**C**

**D**

**E**

**Figure 3.** Fluorescent screening of *C. merolae* strains expressing various fluorescent reporters. **A** – Transgenes were codon optimized for *C. merolae* nuclear genome expression and subcloned into a two-cassette system as illustrated. **B** – Fluorescent proteins highlighted in this work, molecular weight of the protein alone or fused to a chloroplast targeting peptide, biological source or synthetic status, Fluorescent Protein Database ID (https://www.fpbase.org), optical properties and sequence source. **C** – Flow cytometry of cells expressing fluorescent reporters that can be separated from the parental strain. mTagBFP2 and mTFP1 signals compared to wild-type (WT) with violet laser (405 nm) excitation and emission from the VL1 and VL2 filter plotted against forward scatter (FSC). Clover and mVenus signals compared to WT were obtained using blue laser excitation at 488 nm and measuring the emission through the BL1 filter, with the populations plotted against FSC. In addition, mKOk and mScarlet signals compared to WT were acquired by using a 561 nm laser for excitation and then visualized with the YL1 emission filter plotted against FSC. **D** – Transformant screening by plate-level fluorescence analysis on Gellan gum MA2 plates stained with amido-black. 10 µL of three representative transformants from each fluorescent strain were spotted horizontally. Signals from each row after emission imaging are shown for each strain along with their respective excitation (Ex) and emission (Em) filter combination sets. Chlorophyll *a* emission (left side). A composite of all images is shown for comparison. **E** – In-gel fluorescence screening of reporter protein signals. One representative transformant from each strain was subjected to SDS-PAGE and imaged using the ChemStudio Plus with the same filter sets as above. The signal from each reporter appears as a single band, which were then combined to create a composite image. Colors in D and E have been incorporated digitally using the VisionWorks ver. 9.0 software. Full-length images can be found in Supplemental Figure 2. MW – molecular weight, MW + CTP – total molecular weight with chloroplast targeting peptide, L – ladder, B – mTagBFP2, T – mTFP1, V – mVenus, KO – mKOk, LSS – LSSmOrange, S – mScarlet, PSS-Ponceau S staining.

## 3.3 Overexpression of *Ib*IspS results in volatile isoprene production

We wanted to test the heterologous expression of isoprene synthase in *C. merolae* using the transformation and some of the screening tools described before. Terpenoids, also known as isoprenoids, constitute a diverse class of natural compounds many of which are essential metabolites found in all organisms where they play roles in regulating electron transfer, participate in photosynthesis, and a host of other cellular functions (Masyita et al., 2022). Isoprenoids are generated from either the 2-C-methyl-D-erythritol 4-phosphate (MEP) or mevalonate (MVA) pathways to produce the 5-carbon prenylated precursors of terpenoids, isopentyl- or its isomer dimethylallyl- diphosphate (IPP and DMAPP, respectively) (Lichtenthaler, 1999; Perez-Gil et al., 2024; Pu et al., 2021). DMAPP is used by many plants to reduce reactive oxygen species stress through the cleavage of the diphosphate group and release of the 5-carbon volatile molecule isoprene through the action of isoprene synthases. Isoprene production has been engineered into several microbes, both as a means by which to determine how much carbon flux can be diverted to this product and because it represents a bulk platform chemical for use in the

production of rubber or jet fuels (Aldridge et al., 2021; Chaves & Melis, 2018; Diner et al.,
2018; Gomaa et al., 2017; Rana et al., 2022; Yahya et al., 2023). Based on in *silico*
studies, only the MEP pathway is found in the chloroplast of *C. merolae,* where it
generates all required precursors for cellular terpenoids (Grauvogel & Petersen, 2007;
Lohr et al., 2012). *C. merolae* does not naturally produce volatile isoprene and we found
no evidence of isoprene synthase in its genome. We sought to determine whether we
could engineer the production of isoprene from *C. merolae* using our modular plasmid
toolkit. We chose this volatile product as the alga can grow to high densities on $CO_2$ and
its cultivation temperature (42 ˚C) is higher than the boiling point of isoprene (34 ˚C) which
should facilitate its production.

Many isoprene synthases (IspSs) have been heterologously expressed in microbial hosts,
including IspS from the poplar tree (*Populus alba*; (Gao et al., 2016)), the Kudzu vine
(*Pueraria montana; (Gomaa et al., 2017)*), and eucalyptus (*Eucalyptus globulus*; (Englund
et al., 2018; Gao et al., 2016)). The IspS gene from *I. batatas* (sweet potato; *Ib*IspS) was
chosen for heterologous expression in *C. merolae* as it exhibited robust isoprene
production in our previous tests in the green alga *C. reinhardtii* (Yahya et al., 2023). The
amino acid sequence for *Ib*IspS was synthetically optimized for expression from the *C.*
*merolae* nuclear genome after removal of its native plastid targeting peptide and
expressed as an N-terminal fusion with mVenus (Figure 4A). This was done both to
facilitate the identification of robustly expressing clones and to aid protein stability
(Grauvogel & Petersen, 2007; Lohr et al., 2012; Yahya et al., 2023). The native plastid
transit peptide from DNA Gyrase B was used to target the heterologous enzyme to the
algal plastid. We also chose this enzyme as an *in vitro* study had shown the optimum
catalytic activity for this enzyme was 42 ºC, the same as the alga's optimum cultivation
temperature (Li et al., 2019), adding to our interest in its use in *C. merolae*.

After transformation of the linearized plasmid, in-gel fluorescence screening was used to
identify the fusion protein expression in transformants selected for mVenus signal. The
correct molecular mass of the fusion protein was observed (Figure 4B). The transformant
showing the highest expression was then chosen and grown in 400 mL Algem
photobioreactors (Algenuity, UK) to test growth and isoprene accumulation over 6 days
using different light regimes in parallel photobioreactors using a multi-port online

headspace analyzer (Figure 4C–H). The parental *C. merolae* 10D, as well as an mVenus expressing control, were also grown in parallel and their headspaces were compared to the transformant. The transformant was subjected to either day-night cycling or 24-hour continuous illumination using two different settings.

All strains grew in their respective conditions as observed by live monitoring of optical density throughout the cultivation (Figure 4D). No isoprene was detected in either parental or mVenus alone controls, however, isoprene was continuously detected from the transformant expressing the *Ib*IspS (Figure 4E). The isoprene detected was dependent on illumination conditions, with both continuous illumination cultures reaching the same maximum output by day 3 and exhibiting a reduction in the later stages of cultivation (Figure 4E). This did not coincide with the optical density measurements, which continued to increase until day 5 (Figure 4D). The cultures with 12:12 illumination and dark cycles exhibited daily isoprene production which followed the increasing and lowering of the light programs (Figure 4F). This may indicate that the *CPCC* promoter is dynamically controlled by light as very low levels of isoprene were observed in non-illuminated conditions daily. Isoprene production was lower in constant 1200 µE illumination than 750 µE, which may indicate other levels of pathway flux regulation if the promoter is light-controlled. The daily accumulated isoprene production was measured (Figure 4G) and totaled (Figure 4H), with the overall highest amount in the continuous illumination at 750 µE light yielding 80 mg isoprene/L culture in 6 days. This is already higher than ~55 mg/L achieved by expression of the same gene alone in the green alga *C. reinhardtii* (Yahya et al., 2023). The productivity was highest for both continuous cultivations on days 3-4 (Figure 4G), which may also mean growth stage and culture densities play a role in transgene expression. Further systematic investigations of promoter expression rates during cultivation and influences of cultivation conditions will be the subject of follow-up studies.

**Figure 4.** Characterization of isoprene production in an *Ib*IspS expressing *C. merolae* transformant. **A** – Isoprene synthase (IspS) expression plasmid (top) and negative control plasmid expressing mVenus (bottom). The *Ib*IspS gene was directly fused to mVenus and has a predicted molecular mass of 98.6 kDa in contrast to mVenus alone (36.2 kDa). **B** – Molecular mass of the direct fusion between *Ib*IspS-mVenus vs. mVenus alone confirmed by in-gel fluorescence analysis. Six *Ib*IspS-mVenus expressing transformants were compared and the transformant with the strongest expression (arrow) was selected for further experiments. **C** – from left to right: Algem photobioreactors (Algenuity, UK), Hiden Analytical HPR-20 R&D headspace gas analysis system (UK), and 20-port gas inlet. **D** – Optical densities at 740 nm ($OD_{740nm}$) measured every 10 min during cultivation in Algem photobioreactors at different light intensities, and either constant light or under a 12 h light:12 h dark cycle. **E** – Volatile isoprene concentration in headspace under light regimes is shown in F. **F** – Light profiles used for each biological duplicate. For constant illumination, the light intensity increased linearly from 350 µE to the final experimental intensities (i.e., 720 and 1200 µE) after 120 h. **G** – Daily cumulative isoprene recorded in each condition. **H** – Cumulative isoprene produced during the entire cultivation period of 7 days.

**Conclusions**

In this work, we optimized the CAT transformation and selection protocol. This reduces the time from transformation to colony selection from about four weeks to 10 days. This shortened time frame will be beneficial for the often necessary iterative transformations involved in engineering strains. *C. merolae* 10D is a promising alternative photosynthetic cell chassis to green algae because of its unique cultivation conditions, possibilities of transgene HR integration, and stable transgene expression. In this work, we used *in-silico* design and *de novo* construction of transgene expression constructs to demonstrate the speed with which engineering concepts can be implemented in this host. Through this optimized transformation and selection protocol, colonies can be recovered within a reasonable time frame to enable iterative engineering activities. We demonstrated the production of engineered volatile isoprene as a proof of concept, but many other targets that require the introduction of multiple gene pathways are now more feasible to investigate. Given its stable cultivation and requirement for photoautotrophic growth, *C. merolae* is an interesting algal host for scaled cultivation and bioproduction concepts. Further understanding of the behavior of promoters and terminators under different conditions will be required to determine how expressed genes will behave in different cultivation conditions. *C. merolae* represents an interesting emerging model alga that could be used for many biotechnological investigations. Future work should broaden its molecular toolkit with antibiotic-based, counter-selectable markers and further examples of metabolic engineering for which the work presented here will be a useful foundation.

**Acknowledgments**

The authors acknowledge the labs of Shin-Ya Miyagashima and Kan Tanaka for foundational studies of *C. merolae*. KJL acknowledges baseline research funding from KAUST. MVV and MLPM are funded by the KAUST opportunity fund grant #5576 awarded to KJL. PJL acknowledges funding from Xylem, Inc. and ASU Lightworks. SDR and MRS acknowledge funding from the NSERC Discovery Grant program and UNBC's Office of Research and Innovation.

**Conflict of interest**

The authors declare that they have no conflict of interest.

**Supplementary figures legends**

**Suppl. Figure 1.** Full-length images of the molecular screening of mVenus-expressing transformants. **A** – Polymerase chain reaction (PCR) to confirm integration of the linearized plasmid at the D184-185 neutral locus (insert) and presence of the selectable marker (CAT) in 24 transformants. **B** – Plate-level fluorescence analysis of the same transformants on Gellan gum plates stained with amido black. 10 µL of liquid culture were spotted per transformant along with the parental strain (top) to compare for Chlorophyll *a* vs mVenus fluorescent emission, measured at the indicated wavelengths using ChemStudio Plus. **C** – In-gel fluorescence also used to assess the expression level of mVenus in the 24 transformants. The expected molecular weight of mVenus is ~26 kDa. CBB: Coomassie brilliant blue stain included as a loading control.

**Suppl. Figure 2. A-F** – Raw in-gel fluorescence images captured using different filter sets. Filters and exposure times for different fluorophores are indicated. Images were captured in ChemStudio plus. **G** – Ponceau S staining is also included as a loading control.

**Supp. Figure 3.** Quantitative PCR (qPCR) optimization  **A** – Electrophoresis gel showing the results from a thermal gradient PCR to determine the optimal annealing temperature of the primers. A no-template control (NTC; DNA replaced with water) was included. Arrows indicate the best-predicted temperature (~60 ºC ) for both primer sets targeting regions within mVenus and 60S rDNA (reference gene). **B** – Standard curve to measure the efficiency of the amplicon doubling. 91.0% and 109.6% were obtained for 60S rDNA and mVenus, respectively. **C-D** – Template amplification for both target genes when 5 concentrations of genomic DNA were tested (1/5, 1/25, 1/125, 1/625, and 1/3125). **E-F** – Meltcurve generated by the CFX Maestro software. One peak confirms one product is being generated without the presence of non-specific binding. **G** – Cq values of the amplification of 60S rDNA and mVenus for 8 different transformants. Cq values were measured in technical triplicates for each primer set and the standard error is expressed with the error bars.

**References**

Ai, H., Henderson, J. N., Remington, S. J., & Campbell, R. E. (2006). Directed evolution of a monomeric, bright and photostable version of *Clavularia* cyan fluorescent protein: Structural characterization and applications in fluorescence imaging. *Biochemical Journal*, *400*(3), 531–540. https://doi.org/10.1042/BJ20060874

Bindels, D. S., Haarbosch, L., Van Weeren, L., Postma, M., Wiese, K. E., Mastop, M., Aumonier, S., Gotthard, G., Royant, A., Hink, M. A., & Gadella, T. W. J. (2017). mScarlet: A bright monomeric red fluorescent protein for cellular imaging. *Nature Methods*, *14*(1), 53–56. https://doi.org/10.1038/nmeth.4074

Fujiwara, T., Hirooka, S., Mukai, M., Ohbayashi, R., Kanesaki, Y., Watanabe, S., & Miyagishima, S. (2019). Integration of a *Galdieria* plasma membrane sugar transporter enables heterotrophic growth of the obligate photoautotrophic red alga *Cynanidioschyzon merolae*. *Plant Direct*, *3*(4), e00134. https://doi.org/10.1002/pld3.134

Fujiwara, T., & Ohnuma, M. (2017). Procedures for Transformation and Their Applications in Cyanidioschyzon merolae. In T. Kuroiwa, S. Miyagishima, S. Matsunaga, N. Sato, H. Nozaki, K. Tanaka, & O. Misumi (Eds.), *Cyanidioschyzon merolae* (pp. 87–103). Springer Singapore. https://doi.org/10.1007/978-981-10-6101-1_7

Fujiwara, T., Ohnuma, M., Kuroiwa, T., Ohbayashi, R., Hirooka, S., & Miyagishima, S.-Y. (2017). Development of a Double Nuclear Gene-Targeting Method by Two-Step Transformation Based on a Newly Established Chloramphenicol-Selection System in the Red Alga Cyanidioschyzon merolae. *Frontiers in Plant Science*, *8*. https://doi.org/10.3389/fpls.2017.00343

Gutiérrez, S., Wellman, G. B., & Lauersen, K. J. (2022). Teaching an old 'doc' new tricks for algal biotechnology: Strategic filter use enables multi-scale fluorescent protein signal detection. *Frontiers in Bioengineering and Biotechnology*, *10*, 979607. https://doi.org/10.3389/fbioe.2022.979607

Hammel, A., Cucos, L.-M., Caras, I., Ionescu, I., Tucureanu, C., Tofan, V., Costache, A., Onu, A., Hoepfner, L., Hippler, M., Neupert, J., Popescu, C.-I., Stavaru, C., Branza-Nichita, N., & Bock, R. (2024). The red alga *Porphyridium* as a host for molecular farming: Efficient production of immunologically active hepatitis C virus glycoprotein. *Proceedings of the National Academy of Sciences*, *121*(24), e2400145121. https://doi.org/10.1073/pnas.2400145121

Hirooka, S., Tomita, R., Fujiwara, T., Ohnuma, M., Kuroiwa, H., Kuroiwa, T., & Miyagishima, S. (2020). Efficient open cultivation of cyanidialean red algae in acidified seawater. *Scientific Reports*, *10*(1), 13794. https://doi.org/10.1038/s41598-020-70398-z

Imamura, S., Kawase, Y., Kobayashi, I., Sone, T., Era, A., Miyagishima, S., Shimojima, M., Ohta, H., & Tanaka, K. (2015). Target of rapamycin (TOR) plays a critical role in triacylglycerol accumulation in microalgae. *Plant Molecular Biology*, *89*(3), 309–318. https://doi.org/10.1007/s11103-015-0370-6

Imamura, S., Terashita, M., Ohnuma, M., Maruyama, S., Minoda, A., Weber, A. P. M., Inouye, T., Sekine, Y., Fujita, Y., Omata, T., & Tanaka, K. (2010). Nitrate Assimilatory Genes and Their Transcriptional Regulation in a Unicellular Red Alga Cyanidioschyzon merolae: Genetic Evidence for Nitrite Reduction by a Sulfite Reductase-Like Enzyme. *Plant and Cell Physiology*, *51*(5), 707–717. https://doi.org/10.1093/pcp/pcq043

Imoto, Y., & Yoshida, Y. (2017). Cellular Structure of Cyanidioschyzon merolae: A Minimum Set of Organelles. In T. Kuroiwa, S. Miyagishima, S. Matsunaga, N. Sato, H. Nozaki, K. Tanaka, & O. Misumi (Eds.), *Cyanidioschyzon merolae* (pp. 17–27). Springer Singapore. https://doi.org/10.1007/978-981-10-6101-1_2

Kobayashi, Ohnuma, Mio, Kuroiwa, Tsuneyoshi, Tanaka, Kan, & Hanaoka, Mitsumasa. (2010). The basics of cultivation and molecular genetic analysis of the unicellular red alga Cyanidioschyzon merolae. *Endocytobiosis & Cell Research*, *20*.

Kremers, G.-J., Goedhart, J., Van Munster, E. B., & Gadella, T. W. J. (2006). Cyan and Yellow Super Fluorescent Proteins with Improved Brightness, Protein Folding, and FRET Förster Radius ,. *Biochemistry*, *45*(21), 6570–6580. https://doi.org/10.1021/bi0516273

Lam, A. J., St-Pierre, F., Gong, Y., Marshall, J. D., Cranfill, P. J., Baird, M. A., McKeown, M. R., Wiedenmann, J., Davidson, M. W., Schnitzer, M. J., Tsien, R. Y., & Lin, M. Z. (2012). Improving FRET dynamic range with bright green and red fluorescent proteins. *Nature Methods*, *9*(10), 1005–1012. https://doi.org/10.1038/nmeth.2171

Lang, I., Bashir, S., Lorenz, M., Rader, S., & Weber, G. (2022). Exploiting the potential of Cyanidiales as a valuable resource for biotechnological applications. *Applied Phycology*, *3*(1), 199–210. https://doi.org/10.1080/26388081.2020.1765702

Li, Z., & Bock, R. (2018). Replication of bacterial plasmids in the nucleus of the red alga Porphyridium purpureum. *Nature Communications*, *9*(1), 3451. https://doi.org/10.1038/s41467-018-05651-1

Matsuzaki, M., Misumi, O., Shin-i, T., Maruyama, S., Takahara, M., Miyagishima, S., Mori, T., Nishida, K., Yagisawa, F., Nishida, K., Yoshida, Y., Nishimura, Y., Nakao, S., Kobayashi, T.,

Momoyama, Y., Higashiyama, T., Minoda, A., Sano, M., Nomoto, H., … Kuroiwa, T. (2004).
Genome sequence of the ultrasmall unicellular red alga Cyanidioschyzon merolae 10D.
*Nature*, *428*(6983), 653–657. https://doi.org/10.1038/nature02398

Minoda, A., Sakagami, R., Yagisawa, F., Kuroiwa, T., & Tanaka, K. (2004). Improvement of
Culture Conditions and Evidence for Nuclear Transformation by Homologous Recombination
in a Red Alga, Cyanidioschyzon merolae 10D. *Plant and Cell Physiology*, *45*(6), 667–671.
https://doi.org/10.1093/pcp/pch087

Misumi, O., Matsuzaki, M., Nozaki, H., Miyagishima, S., Mori, T., Nishida, K., Yagisawa, F.,
Yoshida, Y., Kuroiwa, H., & Kuroiwa, T. (2005). *Cyanidioschyzon merolae* Genome. A Tool
for Facilitating Comparable Studies on Organelle Biogenesis in Photosynthetic Eukaryotes.
*Plant Physiology*, *137*(2), 567–585. https://doi.org/10.1104/pp.104.053991

Miyagishima, Fujiwara, T., Sumiya, N., Hirooka, S., Nakano, A., Kabeya, Y., & Nakamura, M.
(2014). Translation-independent circadian control of the cell cycle in a unicellular
photosynthetic eukaryote. *Nature Communications*, *5*(1), 3807.
https://doi.org/10.1038/ncomms4807

Miyagishima, & Tanaka, K. (2021). The Unicellular Red Alga *Cyanidioschyzon merolae*— The
Simplest Model of a Photosynthetic Eukaryote. *Plant and Cell Physiology*, *62*(6), 926–941.
https://doi.org/10.1093/pcp/pcab052

Mori, N., Moriyama, T., Toyoshima, M., & Sato, N. (2016). Construction of Global Acyl Lipid
Metabolic Map by Comparative Genomics and Subcellular Localization Analysis in the Red
Alga Cyanidioschyzon merolae. *Frontiers in Plant Science*, *7*.
https://doi.org/10.3389/fpls.2016.00958

Moriyama, T., Mori, N., & Sato, N. (2015). Activation of oxidative carbon metabolism by nutritional
enrichment by photosynthesis and exogenous organic compounds in the red alga
Cyanidioschyzon merolae: Evidence for heterotrophic growth. *SpringerPlus*, *4*(1), 559.
https://doi.org/10.1186/s40064-015-1365-0

Moriyama, T., Sakurai, K., Sekine, K., & Sato, N. (2014). Subcellular distribution of central
carbohydrate metabolism pathways in the red alga Cyanidioschyzon merolae. *Planta*,
*240*(3), 585–598. https://doi.org/10.1007/s00425-014-2108-0

Moriyama, T., Tajima, N., Sekine, K., & Sato, N. (2014). Localization and Phylogenetic Analysis
of Enzymes Related to Organellar Genome Replication in the Unicellular Rhodophyte
Cyanidioschyzon merolae. *Genome Biology and Evolution*, *6*(1), 228–237.
https://doi.org/10.1093/gbe/evu009

Nozaki, H., Takano, H., Misumi, O., Terasawa, K., Matsuzaki, M., Maruyama, S., Nishida, K., Yagisawa, F., Yoshida, Y., Fujiwara, T., Takio, S., Tamura, K., Chung, S. J., Nakamura, S., Kuroiwa, H., Tanaka, K., Sato, N., & Kuroiwa, T. (2007). A 100%-complete sequence reveals unusually simple genomic features in the hot-spring red alga Cyanidioschyzon merolae. *BMC Biology*, *5*(1), 28. https://doi.org/10.1186/1741-7007-5-28

Ohnuma, M., Yokoyama, T., Inouye, T., Sekine, Y., Kuroiwa, T., & Tanaka, K. (2014). Optimization of polyethylene glycol (PEG)-mediated DNA introduction conditions for transient gene expression in the unicellular red alga Cyanidioschyzon merolae. *The Journal of General and Applied Microbiology*, *60*(4), 156–159. https://doi.org/10.2323/jgam.60.156

Ohnuma, M., Yokoyama, T., Inouye, T., Sekine, Y., & Tanaka, K. (2008). Polyethylene Glycol (PEG)-Mediated Transient Gene Expression in a Red Alga, Cyanidioschyzon merolae 10D. *Plant and Cell Physiology*, *49*(1), 117–120. https://doi.org/10.1093/pcp/pcm157

Rahman, D. Y., Sarian, F. D., Van Wijk, A., Martinez-Garcia, M., & Van Der Maarel, M. J. E. C. (2017). Thermostable phycocyanin from the red microalga Cyanidioschyzon merolae, a new natural blue food colorant. *Journal of Applied Phycology*, *29*(3), 1233–1239. https://doi.org/10.1007/s10811-016-1007-0

Schärfen, L., Zigackova, D., Reimer, K. A., Stark, M. R., Slat, V. A., Francoeur, N. J., Wells, M. L., Zhou, L., Blackshear, P. J., Neugebauer, K. M., & Rader, S. D. (2022). Identification of Alternative Polyadenylation in Cyanidioschyzon merolae Through Long-Read Sequencing of mRNA. *Frontiers in Genetics*, *12*, 818697. https://doi.org/10.3389/fgene.2021.818697

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676–682. https://doi.org/10.1038/nmeth.2019

Seger, M., Mammadova, F., Villegas-Valencia, M., Bastos De Freitas, B., Chang, C., Isachsen, I., Hemstreet, H., Abualsaud, F., Boring, M., Lammers, P. J., & Lauersen, K. J. (2023). Engineered ketocarotenoid biosynthesis in the polyextremophilic red microalga Cyanidioschyzon merolae 10D. *Metabolic Engineering Communications*, *17*, e00226. https://doi.org/10.1016/j.mec.2023.e00226

Shcherbakova, D. M., Hink, M. A., Joosen, L., Gadella, T. W. J., & Verkhusha, V. V. (2012). An Orange Fluorescent Protein with a Large Stokes Shift for Single-Excitation Multicolor FCCS and FRET Imaging. *Journal of the American Chemical Society*, *134*(18), 7913–7923. https://doi.org/10.1021/ja3018972

899     Subach, O. M., Cranfill, P. J., Davidson, M. W., & Verkhusha, V. V. (2011). An Enhanced
900           Monomeric Blue Fluorescent Protein with the High Chemical Stability of the Chromophore.
901           *PLoS ONE*, *6*(12), e28674. https://doi.org/10.1371/journal.pone.0028674

902     Sumiya, N., Kawase, Y., Hayakawa, J., Matsuda, M., Nakamura, M., Era, A., Tanaka, K., Kondo,
903           A., Hasunuma, T., Imamura, S., & Miyagishima, S. (2015). Expression of Cyanobacterial
904           Acyl-ACP Reductase Elevates the Triacylglycerol Level in the Red Alga *Cyanidioschyzon*
905           *merolae*. *Plant and Cell Physiology*, *56*(10), 1962–1980. https://doi.org/10.1093/pcp/pcv120

906     Suzuki, K., Ehara, T., Osafune, T., Kuroiwa, H., Kawano, S., & Kuroiwa, T. (1994). Behavior of
907           mitochondria, chloroplasts and their nuclei during the mitotic cycle in the ultramicroalga
908           Cyanidioschyzon merolae. *European Journal of Cell Biology*, *63*(2), 280–288.

909     Takemura, T., Imamura, S., & Tanaka, K. (2019). Identification of a chloroplast fatty acid exporter
910           protein, CmFAX1, and triacylglycerol accumulation by its overexpression in the unicellular
911           red alga Cyanidioschyzon merolae. *Algal Research*, *38*, 101396.
912           https://doi.org/10.1016/j.algal.2018.101396

913     Takemura, T., Kobayashi, Y., Imamura, S., & Tanaka, K. (2019). Top Starch Plating Method for
914           the Efficient Cultivation of Unicellular Red Alga Cyanidioschyzon merolae. *BIO-PROTOCOL*,
915           *9*(4). https://doi.org/10.21769/BioProtoc.3172

916     Tsutsui, H., Karasawa, S., Okamura, Y., & Miyawaki, A. (2008). Improving membrane voltage
917           measurements using FRET with new fluorescent proteins. *Nature Methods*, *5*(8), 683–685.
918           https://doi.org/10.1038/nmeth.1235

919     Villegas-Valencia, M., González-Portela, R. E., De Freitas, B. B., Al Jahdali, A., Romero-Villegas,
920           G. I., Malibari, R., Kapoore, R. V., Fuentes-Grünewald, C., & Lauersen, K. J. (2023).
921           Cultivation of the polyextremophile Cyanidioschyzon merolae 10D during summer conditions
922           on the coast of the Red Sea and its adaptation to hypersaline sea water. *Frontiers in*
923           *Microbiology*, *14*, 1157151. https://doi.org/10.3389/fmicb.2023.1157151

924     Watanabe, S., Sato, J., Imamura, S., Ohnuma, M., Ohoba, Y., Chibazakura, T., Tanaka, K., &
925           Yoshikawa, H. (2014). Stable expression of a GFP-reporter gene in the red alga
926           *Cyanidioschyzon merolae. Bioscience, Biotechnology, and Biochemistry*, *78*(1), 175–177.
927           https://doi.org/10.1080/09168451.2014.877823

928     Wichmann, J., Baier, T., Wentnagel, E., Lauersen, K. J., & Kruse, O. (2018). Tailored carbon
929           partitioning for phototrophic production of (E)-α-bisabolene from the green microalga
930           Chlamydomonas reinhardtii. *Metabolic Engineering*, *45*, 211–222.
931           https://doi.org/10.1016/j.ymben.2017.12.010

Zienkiewicz, M., Krupnik, T., Drożak, A., Golke, A., & Romanowska, E. (2017). Transformation of
the Cyanidioschyzon merolae chloroplast genome: Prospects for understanding chloroplast
function in extreme environments. *Plant Molecular Biology*, *93*(1–2), 171–183.
https://doi.org/10.1007/s11103-016-0554-8

Zienkiewicz, M., Krupnik, T., Drożak, A., & Kania, K. (2019). PEG-mediated, Stable, Nuclear and
Chloroplast Transformation of Cyanidioschizon merolae. *BIO-PROTOCOL*, *9*(17).
https://doi.org/10.21769/BioProtoc.3355