1

MLCK/MLCP regulates mammalian axon regeneration and redistributes the growth cone F-actin

Saijilafu[1, 2#], Wei-Hua Wang[2#], Jin-Jin Ma[2], Yin Yin[3], and Yan-Xia Ma[2]

1 Key Laboratory of Novel Targets and Drug Study for Neural Repair of Zhejiang Province, School of

Medicine, Hangzhou City University, Hangzhou, Zhejiang 310015, China

2 Department of Orthopaedics, the First Affiliated Hospital, Orthopaedic Institute, Soochow University,

Suzhou, China, 215007

3 Laboratory Animal Center, Suzhou Medical College of Soochow University, Suzhou, China, 215123

*Correspondence: Saijilafu, Ph.D.

Phone: 86-15190093375

Email: saijlf@hzcu.edu.cn

Key words: axon regeneration, myosin light chain, MLCK, MLCP, optic nerve injury, spinal cord injury

Abstract

Axon regrowth is a key determinant of the restoration of the biological function of the nervous system

after trauma. However, mature mammalian neurons have limited capacity for axon regeneration. We have

previously demonstrated that neuronal axon growth both in the central and the peripheral nervous systems is

markedly enhanced when non-muscle myosin II (NMII) is inhibited with blebbistatin. The activity of NMII is

primarily regulated by MLCK and MLCP *via* the phosphorylation and dephosphorylation of its light chain,

respectively; however, the functional roles of MLCK and MLCP in mammalian axonal regeneration remain

unknown. In the present study, we provide strong evidence that the inhibition of MLCK activity significantly

blocks axon regeneration in mice. Conversely, the inhibition of MLCP promotes axon regrowth of both the

peripheral and central nervous system. Our findings further indicate that the MLCK/MLCP regulates axon

regeneration and redistributes the growth cone F-actin, and this result suggests that direct regulation of the

growth cone machinery is a potential strategy to promote axon regeneration.

Introduction

The coordinated regulation of growth cone cytoskeleton components is essential for axon growth. Additionally, cytoskeletal proteins in the growth cone are converging targets of extracellular inhibitory molecules. It is now recognized that extracellular inhibitory molecules, such as myelin-based inhibitors and chondroitin sulfate proteoglycans (CSPGs) in glial scars are a major hurdle for successful axon regeneration. These extracellular inhibitory molecules typically trigger the activation of the RhoA/ROCK pathway, leading to phosphorylation and, consequently, the deactivation of cofilin, a protein that severs actin filaments and promotes their depolymerization. Cofilin deactivation results in the stabilization of the growth cone actin cytoskeleton protein, and, subsequently, growth cone collapses. Accordingly, in order to overcome those inhibitory substrates and promote axonal outgrowth, several recent studies have focused on manipulating growth cone cytoskeleton dynamics, and have obtained promising results[1–4]. Here, we hypothesized that the appropriate spatiotemporal regulation of the local cytoskeletal machinery in the growth cone can provide an alternative and effective approach for enhancing axon regeneration.

We have previously shown that the inhibition of non-muscle myosin II (NMII) activity can promotes axonal growth over inhibitory substrates, such as CSPGs and myelin, in both the central and peripheral nervous systems (CNS and PNS, respectively)[1]. Furthermore, this growth-promoting effect is achieved *via* the reorganization of microtubules (MTs) in the growth cone[1]. Myosin proteins constitute a superfamily of ATP-dependent motor proteins known primarily for their role in muscle contraction. In the nervous system, however, myosin II usually drives actin retrograde flow in the growth cone, while the actin cytoskeleton

powers directional growth cone motility[5,6]. An increasing number of studies have shown that the actin

cytoskeleton and its regulators also play a critical role in axon elongation and guidance. Myosin II is typically

activated *via* the phosphorylation of the N-terminus of its light chain by myosin light chain kinase (MLCK)

and deactivated through dephosphorylation by myosin light chain phosphatase (MLCP). An early study

reported that the expression of MLCK, a calcium/calmodulin-dependent kinase, is increased during axonal

outgrowth in goldfish retinal ganglion cells[7]. The overexpression of constitutively active MLCK in Drosophila

CNS neurons increases myosin II activity and leads to impaired axonal outgrowth[8]. Zhang *et al.* demonstrated

that MLCK inhibition attenuates 5-HT-dependent neurite outgrowth in bag cell neurons of the mollusk

Aplysia[9]. Additionally, the activation of MLCP/MLC signaling in PC12 cells was also shown to promote

neurite outgrowth[10]. Together, these reports suggested that the MLCK/MLCP regulates axon growth in

non-mammalian organisms. However, to our knowledge, the functional roles of MLCK and MLCP during

mammalian axon regeneration remain unknown.

In the present study, we found that sciatic nerve injury elevated MLCK protein expression, leading to

increased levels of MLC phosphorylation in dorsal root ganglion (DRG) neurons. Furthermore, the

pharmacological inhibition or genetic silencing of MLCK blocked axon regeneration in the PNS and CNS both

in vitro and *in vivo*, whereas the pharmacological inhibition or transcriptional silencing of MLCP dramatically

promotes axon regeneration. We further found that the MLCK/MLCP activity also affects F-actin redistribution

in the growth cone. Taken together, our results indicated that MLCK and MLCP may regulate axon

regeneration and redistribute the growth cone F-actin, and the direct manipulation of cytoskeleton proteins in

the growth cone may be a promising strategy for enhancing mammalian axon regeneration.

Materials and Methods

Animals and reagents

Adult ICR (8–10 weeks old, 20–30 g) were used. BDA (D1956) and CTB labeled with Alexa Fluor 488

(CTB-488, C34775) were obtained from Life Technologies. ML-7 hydrochloride (110448-33-4) was purchased

from Santa Cruz Biotechnology. PDBu (P1269) was acquired from Sigma-Aldrich. Blebbistatin (S7099) was

obtained from Selleck Chemicals. Antibodies against βIII-tubulin (TUJ1, T8578), MLCK (SAB1300116), and

GFAP (G3893) were purchased from Sigma-Aldrich. Antibodies against β-actin (#4970) and p-MLC (#3675)

were acquired from Cell Signaling Technology. Actin-stain 555 phalloidin (Cat[#] PHDH1) was purchased from

Cytoskeleton, Inc. siRNAs designed to target mouse MLCK (siMLCK) (5′-GGCAAATACACCTGTGAAG-3′,

5′-TGGTCAAAGAAGGGCAGAT-3′,   5′-AGCCAAAGGGAGTCAACAT-3′,   and

5′-TCACGACGGGAATGAGATT-3′) and mouse MYPT1 (siMYPT1) (5′-CTGTGGATATCTCGATATTGC-3′)

were obtained from GenePharma (Shanghai, China). The scrambled siRNA was used as control.

Cell culture

Adult DRG neurons were dissociated from 8 to 10-week-old ICR mice and cultured in Minimal Essential

Medium (MEM) containing 5% (*v/v*) fetal bovine serum (FBS), 5-fluoro-2-deoxyuridine/uridine (20 μM), and

penicillin/streptomycin. Embryonic cortical neurons (from E14.5 embryos) and hippocampal neurons (from

E18.5 embryos) were cultured in neurobasal medium supplemented with penicillin/streptomycin, GlutaMAX,

and B27 supplement. To investigate axonal growth on inhibitory substrates, glass coverslips were first coated

with poly-D-lysine (100 μg/mL) for 2 h, and then coated again with myelin or 10 μg/mL CSPGs for 2 h, as

previously described[1,11].

For the cell replating experiment, DRG neurons were cultured for 3 days, resuspended in culture medium,

replated on new coverslips, and finally re-cultured for a further 18 h to allow new axons to grow, as previously

described[12,13]. Briefly, in treatments denoted as "before replating", neurons were first grown in the presence of

10 μM ML-7 during the initial 3-day culture period, and, after washing out the ML-7, they were replated and

grown in the absence of drug(s) for 18 h. Treatments denoted as "after replating" refer to procedures in which

neurons were first grown in the absence of a drug or drugs during the initial 3-day culture period, after which

10 μM ML-7 was added immediately after replating, followed by 18 h of culture in the presence of drug(s).

Sciatic nerve axotomy

The sciatic nerve was transected at the sciatic notch using ophthalmic scissors. After surgery, the wound was

closed, and the mice were allowed to recover. The mice were then euthanized and L4–L5 DRGs were isolated

for cell culture, quantitative reverse transcription-PCR (qRT-PCR), and western blotting.

siRNA transfection

Cells were transfected with siRNA by electroporation in accordance with the manufacturer's instructions.

Briefly, dissociated neurons were suspended in 80-μL solutions of Amaxa electroporation buffer containing

siRNAs (2–3 nmol) and/or an EGFP-expressing plasmid (5 mg). The cells were then transferred to a 2-mm

cuvette and electroporated using an Amaxa Nucleofector apparatus. After electroporation, the cells were

immediately transferred to 500 µL of prewarmed culture medium and cultured on glass coverslips coated with

100 mg/mL poly-D-lysine. After 4 h, when the neurons were fully attached to the coverslips, the remaining

electroporation buffer was discarded, and fresh culture medium (500 mL) was added. We used the pCMV–

EGFP–N3 as control, and the pCMV– EGFP–N3 plasmid was from Clontech, Inc.

qRT-PCR and western blotting

Total RNA was isolated using Trizol Reagent and assessed for integrity and concentration in a NanoDrop 1000

spectrophotometer. Each sample was then reverse transcribed using Maxima H Minus Reverse Transcriptase.

Real-time qPCR was performed using SYBR Premix ExTaq II in a CFX96 Real-Time qPCR Detection System

(Bio-Rad). Glyceraldehyde 3-phosphate dehydrogenase (*Gapdh*) was used as internal control. The sequences

of the forward and reverse primers were MLCK: 5′-AGAAGTCAAGGAGGTAAAGAATGATGT-3′ and

5′-CGGGTCGCTTTTCATTGC-3′, respectively; and GAPDH: 5′-AGGTCGGTGTGAACGGATTTG-3′ and

5′-TGTAGACCATGTAGTTGAGGTCA-3′, respectively.

For western blotting, samples were homogenized in RIPA buffer. After boiling for 5 min, equal amounts of

extracted protein (20 µg/sample) were separated using 10% sodium dodecyl sulfate–polyacrylamide gel

electrophoresis, and electro-transferred to polyvinylidene fluoride membranes (Immobilon-P; Millipore). After

blocking with 0.01 M PBS containing 5% (*w/v*) skimmed milk overnight, the membranes were incubated first

with primary antibodies and then with the appropriate secondary peroxidase-conjugated antibodies. Protein

bands were developed using ECL Prime Western Blotting Detection Reagent and subsequently quantified

using Image Lab software. An anti-β-actin antibody was used as a loading control.

Immunohistostaining

Mice were deeply anesthetized and transcardiacally perfused with 4% paraformaldehyde (PFA). L4–L5 DRG

tissues were isolated, further fixed in 4% PFA, cryosectioned into 12-μm slices, and blocked with 5% FBS

containing 0.3% Triton X-100. The sections were then incubated with the indicated primary antibody

(monoclonal anti-βIII-tubulin antibody and/or polyclonal anti-MLCK antibody). Immunoreactivity was

visualized using the appropriate fluorescently labeled secondary antibody. Alternatively, cultured DRG neurons

were fixed in 4% PFA, blocked with 2% BSA and 0.1% Triton X-100, and then stained as described above. To

visualize F-actin in the growth cones, *in vitro*-cultured neurons were stained with actin-stain 555 phalloidin

following the manufacturer's instructions.

Axonal growth and growth cone quantification

All images were acquired at 1388 × 1040-pixel resolution using a Zeiss fluorescence microscope with

AxioVision 4.7 software (Carl Zeiss Micro Imaging, Inc.) equipped with a 10× or a 20× objective. To quantify

axon length, neurons with axonal processes exceedingly twice the diameter of their cell body were selected,

and the longest axon from 100 neurons for each condition was measured using the curve tool in AxioVision 4.7.

Meanwhile, growth cone size and F-actin area were measured using the outline tool in AxioVision 4.7. The

size of the growth cone was ascertained from the neck of the microtubule to the edge of the axon tip and the

F-actin area was determined from the hillock to the most distant edge of the growth cone.

In vivo **electroporation of DRGs and sciatic nerve crush**

In vivo electroporation of adult mouse DRGs was performed as previously described[14]. Briefly, each mouse

was anesthetized *via* an intraperitoneal injection of a ketamine (100 mg/kg) and xylazine (10 mg/kg) solution.

The transverse process of L4 and L5 was carefully removed to expose the DRG tissue. A 1 μl solution

containing indicated siRNAs together with the plasmid encoding EGFP (pCMV–EGFP–N3) was then

microinjected into the L4–L5 DRG tissue using a Picospritzer II Microcellular Injection Unit followed by

electroporation using a BTX ECM830 Electro Square Porator (five 15-ms pulses, 35 V, 900-ms interval).

Following this, the muscle and skin were carefully sutured using 5-0 nylon sutures. The mouse was placed on a

heated blanket (35°C) until it had completely recovered from anesthesia. Two days after electroporation, the

sciatic nerves were crushed with forceps #5 and the crush sites were marked with an 11-0 nylon epineuria

suture. After 3 more days, the mice were euthanized by perfusion of 4% PFA, and the whole sciatic nerves

were dissected out and postfixed overnight in 4% PFA at 4°C. Prior to the whole-mount flattening process, it

was essential to verify that the location of the epineural suture corresponded with the site of injury. Only

experiments where the crush site was distinctly recognizable were considered for inclusion in the subsequent

analysis. To quantify axon regeneration, all discernible axons labeled with EGFP within the sciatic nerve were

manually traced and measured from the crush site to the distal tip.

Optic nerve injury

The optic nerves of adult C57BL/6 mice were carefully exposed and crushed with a jeweler's forceps #5 just

behind the eyeball for 1 s. A small piece of gelatin sponge fully soaked with PBS or 50 μM PDBu was placed

at the lesion site; alternatively, 2 μL of PDBu (50 μM) was intravitreally injected into the vitreous body.

Simultaneously, 2 μL of CTB-488 was injected into the vitreous body of each animal after the optic nerve had

been crushed. All microinjections were performed using the Picospritzer II Microcellular Injection Unit (Paker

Ins.; pressure: 15 psi, duration: 6 ms). After 3 days, the optic nerves were dissected out, fixed in 4% PFA

overnight at 4°C, and cut into a series of 12-μm longitudinal sections. The numbers of CTB-labeled axon fibers

were counted at distances of 50 and 100 μm from the injury site. The length of the longest CTB-labeled axon

fiber of each nerve was also measured.

Spinal cord crush injury and PDBu injection

The T8 spinal cord was exposed under a microscope, the whole spinal cord was crushed for 2 s with a

modified jeweler's forceps #5 as previously described[15], and the muscle and skin were separately sutured. A

100-μL volume of PDBu (50 μM) or DMSO was administered every 4 days by intrathecal injection. After 14

days, to trace and visualize the corticospinal tract axon, 1.6 μL of 10% BDA was injected into 4 sites of the

sensorimotor cortex (1.0 mm lateral; 0.5 mm deep into the cortex; 1.0, 0.5, - 0.5, and -1.0 posterior to bregma)

with a 5 μL Hamilton syringe[16]. Fourteen days after BDA injection, mice were transcardically perfused with

4% PFA. The whole spinal cord was isolated from each mouse and post-fixed in 4% PFA for 6 h at 4°C. After

dehydration using an increasing gradient of sucrose concentrations, the spinal cord was cut into 25-μm-thick

sagittal sections, which were then stained with Cy3-streptavidin for 3 h following the manufacturer's protocol.

The lengths of regenerating axons were determined by measuring the distance from the lesion site to the tip of

the BDA-labeled axon tip. The final axon length was the average calculated from 5 sections for each spinal

cord. To quantify the retraction bulb, both maximum width of the enlarged distal tip of the axon and the width

of its immediately adjacent axon shaft were measured. Then, the ratio of these two widths was then calculated.

An axonal tip was considered as a retraction bulb if its tip/shaft ratio exceeded 4. The average number of

retraction bulbs were calculated from 3 sections in every mouse for each group. Five mice were used for each

condition.

BBB locomotor scale score assessments

BBB score assessments were performed as previously described[17]. Briefly, mice were individually placed in an

open, quiet space (45×30 cm) for 5 min while two individuals blinded to the experiments scored the hindlimb

motor function of each animal. Assessments were made on day -1 (the day before injury) and on days 1, 3, 7,

14, 21, 28, 35, and 42 post-injuries for both the DMSO and PDBu treatment groups.

Results

The MLCK expression increased in adult sensory neurons following peripheral axotomy.

Following injury, the expression of axon regeneration-related genes is usually upregulated. Thus, we first

measured MLCK protein levels in DRG neurons following sciatic nerve transection. Compared with naive

neurons, MLCK protein expression was significantly increased in DRG neurons 3 days after sciatic nerve

transection (Figure 1A, B). The mRNA levels of MLCK in DRG neurons were also increased on days 1, 3, and

7 after axotomy, with the highest expression being observed on day 3 (Figure 1C). Given that MLCK

phosphorylates MLC, we subsequently assessed the level of MLC phosphorylation and found that it was

higher in DRG neurons following nerve injury than in control DRG neurons, likely due to the increase in

MLCK expression (Figure 1A, D). Immunohistochemical staining further confirmed that both MLCK

expression and the levels of phosphorylated MLC (p-MLC) increased in DRG neurons following sciatic nerve

axotomy (Figure 1E, F). Taken together, these results indicated that MLCK is activated in DRG sensory

neurons following peripheral axotomy, which leads to an increase in MLC phosphorylation.

The inhibition of MLCK impaired axon regeneration in peripheral sensory DRG neurons.

Next, to explore the functional role of MLCK during peripheral axon regeneration, ML-7 (10 μM), a specific

pharmacological inhibitor of MLCK, was added to DRG neuron cultures for 3 days. The results showed that

ML-7 administration led to a significant reduction in MLC phosphorylation levels (Figure 2A, B, C) and

impaired axonal growth in sensory neurons (Figure 2D, E), and these effects were dose-dependent

(Supplementary Figure S1). The proportion of neurons with axons also decreased after ML-7 treatment (Figure

2E). Furthermore, the transfection of a small interfering RNA (siRNA) targeting MLCK (siMLCK) into

dissociated DRG neurons significantly inhibited MLCK expression and reduced p-MLC levels (Figure 2F, G),

while concomitantly blocking axon growth in adult DRG sensory neurons (Figure 2H, I). Next, using our

previously described *in vivo* DRG electroporation technique[14], we further explored the functional role of

MLCK in sciatic nerve axon regeneration *in vivo*. We co-transfected siMLCK and an EGFP-expressing

plasmid into lumbar vertebrae 4 and 5 (L4 and L5) DRGs of adult mice by electroporation. After 2 days, mice

were subjected to sciatic nerve crush injury, and, after another 3 days, the lengths of regenerating axons were

measured in whole-mount sciatic nerves. Our data showed that the mean length of *vivo* regenerating axons was

significantly reduced in siMLCK-treated mice compared with that in control animals (Figure 2J, K). These

findings suggested that MLCK activity is required for mammalian peripheral axon regeneration both *in vitro*

and *in vivo*.

The inhibition of MLCP promoted peripheral axon growth over inhibitory substrates.

It is well known that the opposite roles of MLCK and MLCP in regulating the MLC phosphorylation status.

Consequently, we also investigated the functional role of MLCP during axon regeneration. It has been reported

that phorbol 12,13-dibutyrate (PDBu) can block MLCP activity, which leads to an increase in MLC

phosphorylation[18]. Here, the results demonstrated that PDBu treatment (20 µM) significantly increased the

p-MLC level in cultured DRG neurons (Supplementary Figure S2A, B). Importantly, we found that this

occurred in parallel with greatly enhanced axonal growth in DRG neurons cultured on permissive substrates,

such as poly-D-lysine (Supplementary Figure S2C, D). A key reason why CNS neurons cannot spontaneously

regenerate axons after injury is due to the inhibitory local microenvironment. Myelin and CSPGs are

well-characterized CNS-based inhibitory substrates that block axon regeneration. Therefore, we next examined

whether inhibition of MLCP activity with PDBu promoted axon regeneration in the presence of these potential

CNS-based inhibitory substrates. When dissociated adult DRG neurons were cultured on myelin extracts or

CSPGs, sensory neuronal axon growth was significantly suppressed (Supplementary Figure S2C, D).

Interestingly, however, when PDBu was added to the culture medium, axon growth was significantly enhanced

on either substrate (Supplementary Figure S2C, D). This result indicated that PDBu treatment relieved the

inhibitory effect of myelin and CSPGs on axonal growth in DRG neurons. MLCP is a phosphatase complex

comprising PP1δ, MYPT1, and M20[19]. In this study, a siRNA specifically targeting MYPT1 (siMYPY1) was

found to markedly increase MLC phosphorylation in cultured DRG neurons (Figure 3A, B). In line with the

results obtained with the pharmacological inhibitor, siMYPT1 also significantly stimulated axonal growth in

DRG sensory neurons cultured on myelin or CSPGs (Figure 3C, D). Furthermore, the transfection of siMYPT1

into L4–L5 DRGs of adult mice by electroporation also significantly promoted sciatic nerve axon regeneration

in vivo (Figure 3E, F). Combined, these data indicated that the inhibition of MLCP activity enables mammalian

axon regeneration in vitro and in vivo.

MLCK/MLCP activity regulated axon growth in the embryonic CNS in vitro.

Compared with neurons of the PNS, those of the CNS have intrinsically poor axonal growth capability, which

also helps explain why CNS neurons cannot regenerate axons spontaneously after injury. Therefore, we next

examined the role of MLCK/MLCP on CNS axon regeneration using in vitro cultured cortical and

hippocampal neurons from embryonic day (E)14.5 and E18.5 mice, respectively. To block MLCK activity, the

neurons were treated with 10 µM ML-7 for 3 days. The results showed that the MLC phosphorylation levels

were markedly inhibited with ML-7 treatment (Figure 4A); simultaneously, the mean axon length of

ML-7-treated E14.5 cortical neurons was significantly reduced compared with that of the control neurons

(Figure 4C, D). Furthermore, consistent with the pharmacological results, the siRNA-mediated suppression of

MLCK expression in E14.5 cortical neurons (Figure 4B) significantly inhibited their axonal growth (Figure 4E,

F). Conversely, the blockade of MLCP activity with PDBu (20 µM; Figure 4A) notably augmented cortical

neuronal axonal growth (Figure 4C, D). The siRNA-mediated transcriptional silencing of MYPT1 expression

also upregulated p-MLC levels in cortical neurons (Figure 4B), and, accordingly, also enhanced axonal growth

(Figure 4E, F). Similar results were obtained with E18.5 hippocampal neurons (Figure 4G-J). Overall, these

findings demonstrated that MLCK/MLCP activity also regulates axon growth in the developing CNS.

The local inhibition of MLCP activity at an injury site promoted optic nerve regeneration.

Given the promising axon growth-promoting effects observed *in vitro*, we wondered whether inhibiting MLCP

activity could stimulate *in vivo* CNS axon regeneration. The optic nerve forms part of the CNS and, like other

neurons of the CNS, also have weak intrinsic axon-regenerating ability (Supplementary Figure S3). To test this,

the right optic nerve of adult mice was crushed, and PBS or PDBu (50 μM) was applied locally to the lesion

site using a gelatin sponge; alternatively, 2 μL of PDBu (50 μM) was microinjected into the vitreous body of

another mice. Subsequently, 2 μL of Alexa Fluor 488-conjugated cholera toxin subunit B (CTB) was injected

into the vitreous body to anterogradely label regenerating axons. After 3 days, the whole optic nerve was

harvested, and CTB-labeled axon regeneration was assessed. The local administration of PDBu using a gelatin

sponge resulted in numerous CTB-labeled axon fibers growing over the injury site, the longest being

approximately 400 μm long (Figure 5A-C). In contrast, local PBS administration at the injury site or

intravitreal PDBu injection did not significantly enhance axon regeneration beyond the injury site (Figure 5

A-C). This observation suggests that the only treatment employed in the injury site (the inhibition of MLCP

activity within the growth cone) effective promote axonal growth.

The local inhibition of MLCP activity prevented axonal retraction in the injured spinal cord.

After spinal cord injury, corticospinal tract (CST) axons also did not regenerate spontaneously, principally

because of the inhibitory environment (Supplementary Figure S4). To test whether local inhibition of MLCP

activity can also promote spinal cord axon regeneration, mice were subjected to spinal cord crush injury at the

level of the eighth thoracic vertebra (T8)[15]. Briefly, the T8 spinal cord was exposed and crushed for 1 second

with jeweler's forceps #5. A 100 μL PDBu (50 μM) or its vehicle, DMSO, was then administrated once every 4

days by intrathecal injection to block MLCP activity. After 2 weeks, 1.6 μL of 10 % biotinylated dextran amine

(BDA) was microinjected into the sensorimotor cortex to label regenerating axons[16]. After a further 2 weeks,

BDA -labeled regenerating axon was evaluated in the spinal cord section. Intrathecal PDBu injection

significantly boosted p-MLC levels (Figure 5D, E) and prevented injury-induced axon retraction (Figure 5F,

G), while also greatly reducing the number of retraction bulbs (Supplementary Figure S5). However, we did

not observe any BDA-labeled axon fibers beyond the lesion site, and Basso–Beattie–Bresnahan (BBB) score

evaluation following PDBu treatment did not differ from that of the control group (Figure 5H). Together, these

results indicate that the local inhibition of MLCP activity reduces injury-induced retraction bulb formation and

axon retraction in the spinal cord.

MLCK may regulate axon regeneration independent of myosin II activity.

We further examined whether the regulatory effect of MLCK/MLCP on axonal growth was mediated through

myosin II activity. For this, dissociated DRG neurons were co-treated with blebbistatin (a pharmacological

myosin II inhibitor) and ML-7. Consistent with our previous study [1], the inhibition of myosin II activity with

25 μM blebbistatin markedly promoted axonal growth (Figure 6A, B). Surprisingly, however,

ML-7/blebbistatin co-treatment did not block the inhibitory effect of ML-7 on sensory axon regeneration

(Figure 6A, B). Additionally, blebbistatin administration did not influence MLC phosphorylation levels in

cultured DRG neurons (Figure 6C, D). Collectively, these results interestingly suggested that MLCK regulates

axon regeneration may be independent of myosin II activity.

Local F-actin distribution in the growth cone is regulated by MLCK and MLCP.

Axonal regeneration requires both regulation of gene expression in the soma and the cytoskeletal assembly at

the growth cone. We previously established a culture-and-replating model to investigate whether axonal

regeneration was soma gene transcription-dependent or local growth cone cytoskeleton

assembly-dependent[12,13]. Interestingly, only ML-7 treatment after replating was found to significantly inhibit

axonal growth in peripheral sensory neurons. However, no effect was observed on axonal growth when ML-7

was administrated before cell replating (Figure 7A, B, C). Several studies have reported that peripheral

axotomy triggers local cytoskeleton dynamics-dependent axon growth in DRG neurons [13,16,20]. Consistent with

the culture-and-replating results, axotomy-induced, local cytoskeleton assembly-dependent axon growth was

also markedly inhibited by ML-7 treatment (Figure 7D, E). These results suggested that MLCK primarily

influences local cytoskeleton dynamics during axonal growth. Accordingly, we further examined how

MLCK/MLCP activity regulates growth cone cytoskeletal distribution during axonal growth. E14.5 cortical

neurons were treated with ML-7 or PDBu for 3 days, following which microtubule and F-actin distribution in

the growth cone was visualized using Tuj1 and phalloidin staining. The results indicated that ML-7 treatment

increased growth cone size and F-actin content, whereas PDBu treatment exerted the opposite effect (Figure

8A-C). These findings may indicate that MLCK and MLCP activity regulates F-actin redistribution.

Discussion

In the present study, we demonstrated that peripheral axotomy increases MLCK and MLC phosphorylation levels in adult DRG neurons subsequently. Thus, when MLCK activity was suppressed in mature DRG neurons *via* ML-7- or specific siRNA, axonal growth was blocked in treated neurons both *in vitro* and *in vivo*. Conversely, an inhibit in MLCP activity *via* the knockdown of MYPT1 significantly promoted axonal growth in both PNS and CNS neurons. We further found that the MLCK and MLCP primarily regulate F-actin redistribution in the growth cone during the axon regeneration process.

Axon injury leads to disconnection between neurons and their targets, thereby destroying functional circuits and behavioral outputs. This implies that for the successful restoration of impaired nervous system function, injured axons must regenerate to their original targets and reconstruct functional circuits. Interestingly, it has been reported that the coordinated regulation of the neuronal cytoskeleton dynamics is essential for axon shaft extension [21]. Among them, the microtubule and actin are two major components of neuronal cytoskeleton that involves axonal extension. For example, Athamneh et al. found that the speed of axon elongation is highly correlated with the bulk forward translocation rate of microtubules [22]. In addition, it also has been reported that the axon extension depends on actin assembly and microtubule-actin interactions[23]. Unsurprisingly given these observations, increasing attention has been paid to promoting axon regeneration *via* the modulation of local cytoskeletal dynamics in the growth cone. Myosin II is a key cytoskeleton protein, which activated through the MLCK-mediated phosphorylation of its light chain and deactivated *via* the dephosphorylation of the N-terminus of its light chain by MLCP. Here, we found that the MLCK expression

and phosphorylation level of MLC was increased in adult DRG neurons after sciatic nerve injury, while the

inhibition of MLCK blocked axon regeneration. These results indicate that activated MLCK is required for

mammalian axon regeneration. If thus, an increase of MLCK expression will activate myosin II during axon

regeneration. However, in our previous study, we found that pharmacological blockade of myosin II activity

with blebbistatin stimulated axon growth [1], which suggested that myosin II plays an inhibitory role in this

process. This apparent contradiction might be attributed to blebbistatin being a specific inhibitor of non-muscle

myosin II ATPase activity and may not affect MLCK activity or the phosphorylation status of MLC[24]. Indeed,

blebbistatin treatment did not influence MLC phosphorylation levels in our study, in line with a previous report

in which it was found that blebbistatin does not alter the phosphorylation status of MLC in sensory neurons

and trabecular meshwork cells[24,35]. The reported mechanism of blebbistatin is not through competition with the

ATP binding site of myosin. Instead, it selectively binds to the ATPase intermediate state associated with ADP

and inorganic phosphate, which decelerates the phosphate release. Importantly, blebbistatin does not impede

myosin's interaction with actin or the ATP-triggered disassociation of actomyosin[36]. It rather inhibits the

myosin head when it forms a product complex with a reduced affinity for actin. This indicates that blebbistatin

functions by stabilizing a particular myosin intermediate state that is independent of the phosphorylation status

of MLC. Additionally, our results demonstrated that blebbistatin treatment could not rescue the inhibitory

effect of ML-7 on axon regeneration. If MLCK regulates axon growth through the activation of myosin, the

inhibitory effect of ML-7 on axon growth might be influenced by blebbistatin, a NMII inhibitor. However, our

findings reveal that the combination of blebbistatin and ML-7 does not alter the rate of axon outgrowth

compared to ML-7 alone. This suggests that the roles of ML-7 and blebbistatin in axon growth are independent.

It means MLCK may regulate axon growth independent of NMII activity. Interestingly, it has been reported

that MLCK regulates cell migration not by MLC phosphorylation, but possible MLCK may serve as an

F-actin-binding protein stabilizing the F-actin/myosin II network of the membrane cytoskeleton[25]. Additionally,

recent study also identified multiple noncanonical targets of MLCK through

CRISPR-Cas9/phosphoproteomics method[26], thus it is also possible that MLCK regulates axon growth its

noncanonical targets or unknown mechanism. They also found that MLCK plays an important role in

vasopressin-induced actin depolymerization[26]. Studies have shown that the combination of MLCK with F-actin

is critical to its function. Short-chain MLCK binds to F-actin via three DFRXXL motifs at its N-terminal,

while long-chain MLCK has two additional DFRXXL motifs and six IG-like modules, which may give it

unique binding properties in cell localization[25]. During axon growth, the dynamic changes of F-actin are

regulated by a variety of proteins, such as ADF/cofilin, Arp2/3, Eps8, Profilin, myosin II, myosin V, etc. These

proteins regulate axon growth and orientation by influencing the polymerization and depolymerization of

F-actin. For example, cofilin increases the availability of actin by promoting the depolymerization of F-actin,

which promotes axon growth[37]. Furthermore，MLCP inhibition also increase actin arc movement in growth

cone[5]. This means that MLCK/MLCP may also regulate axon growth via actin dynamics. In addition, other

studies also have shown that the absence of MLCK not only impairs the synthesis of transmembrane

complexes, such as integrin, and fibronectin, but also disrupts the balance of membrane tension and protrusion,

thereby causing the membrane cytoskeleton instability[38, 39]. We think these may affect axon growth.

Meanwhile, adult sensory neurons express different isoforms of myosin II, which also exert distinct effects on

axonal regeneration[27]. For example, the knockdown of myosin IIA enhances axonal growth, while the

knockdown of myosin IIB increases F-action retrograde flow and inhibits axon growth[1,28–31]. A study in

Neuro-2A cells also revealed that myosin IIA is involved in axon retraction, whereas myosin IIB was reported

to contribute to axon extension[28,29]. Interesting，in neurons, although expression of both NMIIA and NMIIB

has been reported, the NMIIB being the predominant one[40]. Under physiological conditions, MLCP is

activated by PKC, and it has been reported that the PKC-mediated activation of myosin II is restricted to the T

zone of growth cones; the results of the present study suggest that growth cones require low-to-moderate levels

of myosin II activity and F-actin for best growth rates. This observation emphasizes the complexity of the role

of myosin II in axonal growth, the precise molecular mechanisms of which require further in-depth

investigation.

The inability of neurons in the CNS to regenerate their axons after injury is attributable, at least in part, to

the presence of inhibitory molecules at the injury site, including CSPGs and myelin. Here, we found that the

inhibition of MLCP activity with PDBu or the siRNA-mediated knockdown of MYPT1 promoted axonal

growth in adult sensory neurons over both CSPG and myelin substrates. Our findings also further indicated

that the local inhibition of MLCP by PDBu not only induces axon regeneration in peripheral sensory neurons

but also promotes axon regeneration in the optic nerve. Studies have shown that MLCK possesses several

actin-binding domains to which F-actin can bind and form bundles[32,33] and that directional growth cone

motility during axon growth is controlled by actin-based machinery. Growth cone movement during axon

growth is achieved through protrusion towards attractive guidance cues and retraction from repulsive ones in

an actin-dependent manner. In the classic molecular clutch model, strong mechanical coupling between the

F-actin cytoskeleton and the substrate, mediated by cell adhesion complexes, transmits the forces generated by

the cytoskeleton into rearward traction forces, enabling the forward movement of the growth cone. Our results

indicated that treating DRG neurons with ML-7 induces the increases F-actin content in the growth cone.

Conversely, the inhibition of MLCP following PDBu administration resulted in decreased F-actin content. Our

findings are consistent with the results of a previous study on Helisoma, in which treatment with ML-7

increased F-actin content at the C/P-domain interface, thus inhibiting axonal regeneration[34]. This suggests that

MLCK or MLCP may control mammalian axonal regeneration by regulating growth cone F-actin distribution.

Together, our findings indicate that directly targeting growth cone cytoskeleton components may be a

promising strategy for promoting mammalian axon regeneration.

Statements & Declarations

Funding

This work was supported by the National Natural Science Foundation of China (Nos. 81571189 to S), A

Priority Academic Program Development of Jiangsu Higher Education Institutions, and Innovation and

Entrepreneurship Program of Jiangsu Province.

Author Contribution

Saijilafu, W.H. W and J.J. M designed the experiment. W.H. W and J.J. M performed the experiments and

analyzed the data. W.H. W, Y.Y, Y.X. M and Saijilafu co-wrote the paper with all authors' input.

Ethics approval

The animal experiments were conducted in accordance with the National Institutes of Health Guide for the

Care and Use of Laboratory Animals (NIH Publications No. 8023, revised 1978) and were approved by the

Ethics Committee of the First Affiliated Hospital of Soochow University.

Consent to publish

All the authors consent for publication.

Conflict of interests

The authors have no relevant financial interests to disclose.

Data availability

All data generated or analyzed during this study are available from the corresponding author upon reasonable

request.

References

1.   Hur, E.-M. *et al.* Engineering neuronal growth cones to promote axon regeneration over inhibitory

molecules. *Proceedings of the National Academy of Sciences* 108, 5057–5062 (2011).

2.   Sengottuvel, V., Leibinger, M., Pfreimer, M., Andreadaki, A. & Fischer, D. Taxol Facilitates Axon

Regeneration   in   the   Mature   CNS.   *The   Journal   of   Neuroscience*   2688–2699   (2011)

doi:10.1523/jneurosci.4885-10.2011.

3.   Hellal, F. *et al.* Microtubule Stabilization Reduces Scarring and Causes Axon Regeneration After Spinal

Cord Injury. *Science* 331, 928–931 (2011).

4.   Rivieccio, M. A. *et al.* HDAC6 is a target for protection and regeneration following injury in the nervous

system. *Proceedings of the National Academy of Sciences* 106, 19599–19604 (2009).

5.   Zhang, X.-F., Schaefer, A. W., Burnette, D. T., Schoonderwoert, V. T. & Forscher, P. Rho-Dependent

Contractile Responses in the Neuronal Growth Cone Are Independent of Classical Peripheral Retrograde

Actin Flow. *Neuron* 40, 931–944 (2003).

6.   Schaefer, A. W., Kabir, N. & Forscher, P. Filopodia and actin arcs guide the assembly and transport of

two populations of microtubules with unique dynamic parameters in neuronal growth cones. *The Journal*

of Cell Biology 158, 139–152 (2002).

7.   Jian, X., Szaro, B. G. & Schmidt, J. T. Myosin light chain kinase: expression in neurons and upregulation

during axon regeneration. *Journal of Neurobiology* 31, 379–391 (1996).

8.   Kim, Y.-S., Fritz, J. L., Seneviratne, A. K. & VanBerkum, M. F. A. Constitutively Active Myosin Light

Chain Kinase Alters Axon Guidance Decisions in Drosophila Embryos. *Developmental Biology* 249,

367–381 (2002).

9.   Zhang, X.-F., Hyland, C., Van Goor, D. & Forscher, P. Calcineurin-dependent cofilin activation and

increased retrograde actin flow drive 5-HT–dependent neurite outgrowth in *Aplysia* bag cell neurons.

Molecular Biology of the Cell 23, 4833–4848 (2012).

10.   Tao, T. *et al.* Minocycline Promotes Neurite Outgrowth of PC12 Cells Exposed to Oxygen-Glucose

Deprivation and Reoxygenation Through Regulation of MLCP/MLC Signaling Pathways. *Cellular and*

Molecular Neurobiology 37, 417–426 (2017).

11. Minor, K. *et al.* Decorin promotes robust axon growth on inhibitory CSPGs and myelin via a direct effect

on neurons. *Neurobiology of Disease* 32, 88–95 (2008).

12.   Saijilafu & Zhou, F. Q. Genetic Study of Axon Regeneration with Cultured Adult Dorsal Root Ganglion

Neurons. *Journal of Visualized Experiments* (2012) doi:10.3791/4141.

13.   Saijilafu *et al.* PI3K–GSK3 signalling regulates mammalian axon regeneration by inducing the

expression of Smad1. *Nature Communications* 4, (2013).

14.   Saijilafu, Hur, E. M. & Zhou, F. Q. Genetic dissection of axon regeneration via in vivo electroporation of

adult mouse sensory neurons. *Nature Communications* (2011) doi:10.1038/ncomms1568.

15. Liu, K. *et al.* PTEN deletion enhances the regenerative ability of adult corticospinal neurons. *Nature*

Neuroscience 1075–1081 (2010) doi:10.1038/nn.2603.

16. Xi, F. *et al.* Calcium/calmodulin-dependent protein kinase II regulates mammalian axon growth by

affecting F-actin length in growth cone. *Journal of Cellular Physiology* 234, 23053–23065 (2019).

17. BASSO, D. M., BEATTIE, M. S. & BRESNAHAN, J. C. A Sensitive and Reliable Locomotor Rating

Scale for Open Field Testing in Rats. *Journal of Neurotrauma* 12, 1–21 (1995).

18. El-Yazbi, A. F., Abd-Elrahman, K. S. & Moreno-Dominguez, A. PKC-mediated cerebral vasoconstriction:

Role of myosin light chain phosphorylation versus actin cytoskeleton reorganization. *Biochemical*

Pharmacology 95, 263–278 (2015).

19. Shirazi, A. *et al.* Purification and characterization of the mammalian myosin light chain phosphatase

holoenzyme. The differential effects of the holoenzyme and its subunits on smooth muscle. *Journal of*

Biological Chemistry 269, 31598–31606 (1994).

20. Smith, D. S. & Pate Skene, J. H. A Transcription-Dependent Switch Controls Competence of Adult

Neurons for Distinct Modes of Axon Growth. *The Journal of Neuroscience* 17, 646–658 (1997).

21. Poulain, F. E. & Sobel, A. The microtubule network and neuronal morphogenesis: Dynamic and

coordinated orchestration through multiple players. *Molecular and Cellular Neuroscience* 43, 15–32

(2010).

22. Athamneh, A. I. M. *et al.* Neurite elongation is highly correlated with bulk forward translocation of

microtubules. *Scientific Reports* 7, (2017).

23. Miller, K. E. & Suter, D. M. An Integrated Cytoskeletal Model of Neurite Outgrowth. *Frontiers in*

Cellular Neuroscience 12, (2018).

24. Yu, P., Santiago, L. Y., Katagiri, Y. & Geller, H. M. Myosin II activity regulates neurite outgrowth and

guidance in response to chondroitin sulfate proteoglycans. *Journal of Neurochemistry* 120, 1117–1128

(2012).

25. Chen, C. *et al.* Myosin Light Chain Kinase (MLCK) Regulates Cell Migration in a Myosin Regulatory

Light Chain Phosphorylation-independent Mechanism. *Journal of Biological Chemistry* 289,

28478–28488 (2014).

26. Isobe, K. *et al.* CRISPR-Cas9/phosphoproteomics identifies multiple noncanonical targets of myosin light

chain   kinase.   *American   Journal   of   Physiology-Renal   Physiology*   F600–F616   (2020)

doi:10.1152/ajprenal.00431.2019.

27. Bridgman, P.C & Elkin, L.L. Axonal myosins. *Journal of Neurocytology* 29: 831-41 (2000).

28. Wylie, S. R. & Chantler, P. D. Myosin IIA Drives Neurite Retraction. *Molecular Biology of the Cell* 14,

4654–4666 (2003).

29. Wylie, S. R., Wu, P., Patel, H. & Chantler, P. D. A conventional myosin motor drives neurite outgrowth.

Proceedings of the National Academy of Sciences 95, 12967–12972 (1998).

30. Turney, S. G. & Bridgman, P. C. Laminin stimulates and guides axonal outgrowth via growth cone

myosin II activity. *Nature Neuroscience* 8, 717–719 (2005).

31. Bridgman, P. C., Dave, S., Asnes, C. F., Tullio, A. N. & Adelstein, R. S. Myosin IIB Is Required for

Growth Cone Motility. *The Journal of Neuroscience* 21, 6159–6169 (2001).

32. Hayakawa, K., Okagaki, T., Higashifujime, S. & Kohama, K. Bundling of Actin Filaments by Myosin

Light Chain Kinase from Smooth Muscle. *Biochemical and Biophysical Research Communications* 199,

786–791 (1994).

33. Sellers, J. R. & Pato, M. D. The binding of smooth muscle myosin light chain kinase and phosphatases to

actin and myosin. *Journal of Biological Chemistry* 259, 7740–7746 (1984).

34. Torreano, P. J., Waterman-Storer, C. M. & Cohan, C. S. The effects of collapsing factors on F-actin

content and microtubule distribution of *Helisoma* growth cones. *Cell Motility and the Cytoskeleton* 60,

166–179 (2005).

35. Zhang M. Rao PV. Blebbistatin, a novel inhibitor of myosin II ATPase activity, increases oqueous humor

outflow facility in perfused enucleated porcine eyes. Investigative ophthalmology& visual science, 2005,

46:4130-4138

36. Mihály Kovács, Judit Tóth, Csaba Hetényi, András Málnási-Csizmadia, James R Sellers. Mechanism of

blebbistatin inhibition of myosin II. J Biol Chem 279(34) (2004):35557-35563.

37. Andrea Tedeschi, Sebastian Dupraz, Michele Curcio, Claudia J Laskowski, Barbara Schaffran, Kevin C

Flynn, Telma E Santos, Sina Stern, Brett J Hilton, Molly J E Larson, Christine B Gurniak, Walter Witke,

Frank Bradke. ADF/Cofilin-Mediated Actin Turnover Promotes Axon Regeneration in the Adult CNS,

Neuron 103(6) (2019):1073-1085.

38. C.X. Yang, H.Q. Chen, C. Chen, W.P. Yu, W.C. Zhang, Y.J. Peng, W.Q. He, D.M. Wei, X. Gao, M.S. Zhu,

Microfilament-binding properties of n-terminal extension of the isoform of smooth muscle long myosin

light chain kinase, Cell Res 16 (4) (2006) 367–376.

39. Y. Gao, L.H. Ye, H. Kishi, T. Okagaki, K. Samizo, A. Nakamura, K. Kohama, Myosin light chain kinase as

a multifunctional regulatory protein of smooth muscle contraction, Iubmb Life 51 (6) (2001) 337–344.

40. Ana Rita Costa, Monica M Sousa. Non-Muscle Myosin II in Axonal Cell Biology: From the Growth Cone

to the Axon Initial Segment. Cells. 2020 Aug 26;9(9):1961.

 Figure Legends

Figure 1. MLCK and phosphor-MLC expression increased in sensory neurons after peripheral axotomy.

(A-D) Sciatic nerve axotomy was performed 3 days before and L4-L5 DRG tissues were collected for Western

blotting or qPCR to examine the expression level of MLCK and phosphor-MLC (p-MLC). Representative

images (A) and quantification of Western blots (B, D) demonstrate that expression levels of MLCK and

p-MLC increase significantly in adult DRGs following sciatic nerve transection. (N=3, ** $p < 0.01$; *** $p <$

0.001).

(C) L4-L5 DRGs were collected for qRT-PCR, 1, 3 and 7 days after axotomy, respectively. The results reveal

that mRNA expression of MLCK in the DRG neurons also increased after axotomy after 1, 3 and 7 days,

although the difference at 3 days was the most significant. (N=3, * $p < 0.05$; ** $p < 0.01$).

(E-F) Sections of L4-L5 DRGs from naive and sciatic nerve axotomized mice were stained with antibodies

against MLCK, Tuj-1 or p-MLC. Immunohistochemistry images indicate that MLCK (E) and p-MLC (F)

expression were elevated in the sensory neurons following sciatic nerve axotomy. Scale bar: 100μm.

Figure 2. Suppression of MLCK activity inhibits axon growth in adult DRG neurons.

(A-E) DRG neurons from adult mice were cultured and treated with DMSO or 10.0 μM ML-7. After 3 days,

neurons were lysed for Western blotting or staining with Tuj1 antibody. Western blot images (A) and

quantification of p-MLC (B) indicate that ML-7 treatment successfully inhibits MLCK activity. Quantification

of axon length indicated that inhibition of MLCK activity with ML-7 dramatically blocks adult sensory

neuronal axon growth *in vitro* (D, E), and reduces the numbers of neurons with axons (E). Scale bar: 100μm

(N=3, *$p < 0.05$; *** $p < 0.001$).

(F-I) Adult DRG neurons were isolated and electroporated with scramble siRNA (Ctrl) or MLCK siRNA

(siMLCK) mixed with GFP. Tuj1 (Red) antibody was used to stain neurons 3 days after transfection. A

representative image (F) and quantification of Western blots for MLCK (G) indicate that siMLCK significantly

suppressed MLCK and p-MLC expression (N=3, *$p < 0.05$). Quantification of mean axon length (H, I)

indicates that transfection of siMLCK inhibits adult DRG neuronal axon growth. Scale bar: 100μm. (N=3, ***

$p < 0.001$).

(J, K) L4/L5 DRGs of adult mice were electroporated with scramble siRNA (Ctrl) or siMLCK combined with

GFP, respectively. Two days later, sciatic nerves were crushed and then regenerated axons traced after a further

three days. Representative images (J) and measurement of regenerating axonal length (K) indicated that

decreased MLCK expression with siRNA inhibits sensory axon regeneration *in vivo*. Crush sites are marked

using red arrows, the marking suture was placed immediately after the crush injury. Regenerating axon tips are

marked with yellow arrow (N=5, * $p < 0.05$). Scale bar: 1mm.

Figure 3. Knockdown of MLCP protein promotes axon growth from adult peripheral sensory neurons.

(A-D) Adult DRG neurons were isolated and electroporated with scrambled siRNA (Ctrl) or MYPT1 siRNA

(siMYPT1) mixed with GFP. A representative Western blot image (A) and quantification of protein band (B)

indicate that transfection of siMYPT1 in cultured DRG neurons significantly increases p-MLC expression

(N=3, ** $p < 0.01$). Quantification of axon length (C, D) indicated siMYPT1 transfection in the adult sensory

neurons promotes its axon growth on the permissive substrate PDL or inhibitory substrates, such as CSPGs and

myelin. (N=3, ***$p < 0.001$).

(E, F) L4/L5 DRGs of adult mice were electroporated with scrambled siRNA (Ctrl) or siMYPT1, respectively.

Two days later, sciatic nerves were crushed and then regenerated axons traced after a further three days.

Representative images (E) and measurement of regenerating axonal length (F) indicated that siMYPT1

promotes sensory axon regeneration *in vivo*. Crush sites are marked using red arrows, and the marking suture

was placed immediately after the crush injury. Regenerating axon tips are marked with yellow arrow (N=5, * p

< 0.05). Scale bar: 1mm.

Figure 4. Inhibition of MLCP activity promotes axon growth from developing embryonic neurons.

(A, B) Cortical neurons were isolated from embryonic day 14.5 embryos. Neurons were treated with DMSO,

ML-7 or PDBu, or transfected with ctrl siRNA, siMLCK or siMYPT1. The p-MLC expression levels were

examined using Western blot analysis after 3 days of culture.

(C-F) coritcal neurons were treated with DMSO, ML-7 or PDBu or transfected with ctrl siRNA, siMLCK or

siMYPT1. After three days, cells were stained with Tuj1 to visualize axons. Representative images (C, E) and

quantification (D, F) of axonal length demonstrate ML-7 or siMLCK inhibits embryonic cortical neuronal

axon growth. In contrast, inhibition of MLCP activity with PDBu or siMYPT1 promoted embryonic cortical

neuronal axon growth (N=3, ***$p < 0.001$). Scale bar: 100 μm.

(G-J) Hippocampal neurons were isolated from embryonic day 18.5 embryos. Neurons were treated with

DMSO, ML-7 or PDBu or transfected with scramble siRNA, siMLCK or siMYPT1. Similarly, Representative

images (G, I) and quantification (H, J) of axonal length demonstrate that ML-7 or siMLCK inhibits

hippocampal neuronal axon growth. In contrast, suppression of MLCP with PDBu or siMYPT1 promoted

hippocampal neuronal axon growth (N=3, ***$p < 0.001$). Scale bar: 100 μm.

Figure 5. Inhibition of MLCP activity promotes CNS axon regeneration in vivo.

(A-C) Local application of a gelatin sponge soaked with PBS (Ctrl) or 50.0 μM PDBu on the site of the lesion

(ON), or microinjection of PDBu (2.0 μl, 50.0 μM) into the vitreous body (ivit) were performed after optic

nerve crush injury. Three days later, CTB-labeled regenerating axons were quantified. Representative images

of optic nerve sagittal sections (A) and quantification of CTB-positive fibers (B) at the point distance 100 μm

and 200 μm from the injury site indicate that only local administration of PDBu at the site of the lesion

promotes the axon growth. (C) The longest axons per section of group were measured and quantified.

(D-H) T8 whole spinal cord crush injury was performed in adult mice then treated with PBS or 50.0 μM PDBu

by intrathecal injection once every 4 days. BDA was injected into the right sensorimotor cortex 2 weeks after

spinal cord injury to label the regenerating cortical spinal tract (CST) axons. After 2 weeks, sagittal sections of

the spinal cord around the crush site were created, and the distance between the tip of the axons and injury site

was measured. Representative image of Western blots (D) and its quantification (E) demonstrates that

intrathecal injection of PDBu increases p-MLC level, it means that PDBu sufficiently inhibits MLCP activity.

Representative image of spinal cord sagittal section (F) and the measurement of distance from BDA-positive

axon tips to the injury site demonstrate that intrathecal injections of PDBu prevent injury-induced axon

retraction. Arrow indicates BDA-positive axons, dash line indicates lesion site. (H) Mean BBB scores

evaluation indicated that there was no difference between PDBu treatment and the PBS control group. (N=5,

***$p < 0.001$).

Figure 6. MLCK regulates axon regeneration independent of myosin II activity.

(A, B) Adult L4-L5 DRG neurons were cultured *in vitro* for 3 days with the addition of the pharmacological

inhibitors indicated. Representative images (A) and quantification of axon length (B) demonstrate that,

although myosin II inhibitor blebbistatin markedly promoted sensory neuronal axon growth, however, it did

not rescue the inhibitory effect of ML-7 on sensory axon growth (N=3, ***$p < 0.001$). (C, D) The blebbistatin

administration did not influence MLC phosphorylation levels in cultured DRG neurons, representative image

of Western blots (C) and its quantification (D) (N=3, ns: no significant).

Figure 7. Inhibition of MLCK blocks transcription-independent axon growth of adult sensory neurons.

(A) Schematic figure of the culture-and-replate model. Adult sensory neurons were cultured for 3 days then

replated to allow growth of new axons for one day. During this process, ML-7 was added prior to or after

replating.

(B, C) Representative images (B) and quantification of axon length (C) indicate that treatment with ML-7 after

replating significantly inhibits adult sensory axon growth. However, treatment with ML-7 prior to replating has

no effect on adult sensory neural axon growth (N=3, ***$p < 0.001$).

(D, E) Mice underwent sciatic nerve axotomy 7 days previously. L4-L5 DRGs were harvested from naive or

axotomized mice and cultured for 1 day with ML-7 treatment. Neurons were stained with Tuj1. Representative

images (D) and quantification of axon length (E) indicate that blockade of MLCK activity with ML-7 inhibits

axotomy-induced axon growth of adult sensory neurons (N=3, ***$p < 0.001$).

Figure 8. MLCK and MLCP activity regulate F-actin distribution of the growth cone.

(A, B) Embryonic E14.5 cortical neurons were isolated and treated with DMSO, PDBu or ML-7. After 3 days

of culture, neurons were stained with Tuj1 antibody (Green) to visualize tubulin and phalloidin (Red) to

visualize F-actin. Representative images (A), quantification of the size of the growth cone (B), and F-actin

content (C) indicate that ML-7 treatment increases the growth cone size and F-actin content, and PDBu

treatment reduces the growth cone size and the F-actin content. Scale bar: 1μm (N=3, ***$p < 0.001$).

Figure S1. ML-7 treatment inhibits axon growth in a dose-dependent manner.

The DRG neurons from adult mice were cultured and treated with a series of doses of ML-7. After 3 days

culture, neurons were stained with Tuj1 antibody and axonal lengths quantified.

Figure S2. PDBu treatment promotes axon growth from adult peripheral sensory neurons.

(A, B) Representative Western blot image (A) and quantification of protein band (B) indicates that

administration of PDBu, the MLCP pharmacological inhibitor, in cultured DRG neurons increases p-MLC

expression (N=3, ** $p < 0.01$).

(C, D) DRG neurons were cultured on permissive (PDL) or inhibitory substrates (CSPGs or myelin) in the

presence or absence of PDBu, as indicated. Neurons were stained with Tuj1. Representative images (C) and

quantification of axon length (D) show that increased p-MLC expression with PDBu addition, which promotes

adult sensory neuronal axon growth on either permissive substrate PDL or inhibitory substrates, such as

CSPGs and myelin. (N=3, ***$p < 0.001$).

Figure S3. Optic nerve crush injury model

Optic nerve was crushed behind the eyeball with a custom-modified jeweler's 5[#] forceps for 1 second. Two

weeks later, 2.0 µl Alexa Fluor 488-conjugated CTB (green) was injected intravitreally to label regenerating

optic nerve axons. Immunofluorescent staining of GFAP staining (red) was also performed in the longitudinal

section of crushed optic nerves.

Figure S4. Whole spinal cord crush injury model

The whole spinal cord was crushed at T8 level with a custom-modified jeweler's 5[#] forceps for 1 second. Two

weeks later, a 1.6 µl aliquot of 10% BDA was injected into the sensorimotor cortex to label regenerating

cortical spinal tract axons, and another two weeks later, the spinal cord was dissected out. Immunofluorescent

staining of BDA (red) and GFAP staining (green) was performed in the sagittal section of spinal cord.

Figure S5. Intrathecal PDBu injection prevented axonal retraction bulb formation.

Representative images of spinal cord sagittal section (A) and quantification of retraction bulb formation (B).

White triangle indicates retraction bulb (N=3, *** $p < 0.001$).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplementary Figure S1

Supplementary Figure S2

Supplementary Figure S3

Supplementary Figure S4

Supplementary Figure S5