Cell-type-specific striatal modulation of amygdalar acetylcholine in salience assignment

Aixiao Chen[1,3*], Yunjing Li[1,4*], Hangfei Zhu[2*], Hanmei Gu[1,5], Yanhong Weng[1,4], Fuqiang Xu[3], Hanfei Deng[2#], Xiong Xiao[1,6#]

1. Institute of Neuroscience, Key Laboratory of Brain Science on Cognition and Brain-inspired Intelligence, CAS Center for Excellence in Brain Science and Intelligence Technology, Chinese Academy of Sciences, Shanghai 200031, China
2. Institutes of Brain Science, State Key Laboratory of Medical Neurobiology and MOE Frontiers Center for Brain Science, Fudan University, Shanghai 200032, China
3. Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences; Shenzhen-Hong Kong Institute of Brain Science-Shenzhen Fundamental Research Institutions, Shenzhen, China
4. Institute of Neuroscience, Fujian Key Laboratory of Molecular Neurology, Fujian Medical University, Fuzhou, China
5. University of Chinese Academy of Sciences, Shanghai 200031, China
6. Lead Contact

* These authors contributed equally

#Correspondence: hanfeideng@fudan.edu.cn (H.D.), xiao@ion.ac.cn (X.X.)

ABSTRACT

The salience assignment is pivotal for both natural and artificial intelligence. Pioneering studies established that basal forebrain cholinergic neurons process behaviorally relevant salient information. However, the neural circuit mechanism underlying salience assignment remains poorly understood. Here we show that the acetylcholine (ACh) level in the basolateral amygdala (BLA) dynamically represented behavioral salience. Distinct neuronal subpopulations in the nucleus accumbens (NAc), D1- and D2-expressing medium spiny neurons (MSNs), antagonistically and specifically promote and suppress ACh release in the BLA, but not the cortex and hippocampus. These striatal D1 and D2 MSNs regulate BLA ACh by disinhibiting and inhibiting cholinergic neurons in the basal forebrain subregion substantia innominata (SI), respectively. Optogenetic manipulations of the pathway from striatal D1 and D2 MSNs to the SI opposingly affect associative learning. Our findings uncover an unconventional role of striatal MSNs in salience assignment via regulating the salience-representing amygdalar ACh activity.

INTRODUCTION

The brain receives a constant influx of complex sensory inputs from the environment. Salient stimuli, distinguished by their physical distinctiveness, relevance, or emotional impact, are easily noticed, perceived, and remembered. Behavioral salience, the property by which certain stimuli stand out, enables the brain to efficiently allocate its limited perceptual and cognitive resources to the most behaviorally relevant stimuli [1,2]. In the ever-changing environment where our goals and internal states also fluctuate, the stimuli salience can change swiftly. Salience assignment mechanisms allow the brain to dynamically adjust attentional priorities in response to shifting environmental demands, internal states, and goals. This process optimizes the allocation of

resources to the most pertinent information for the intended behavior, such as perception,

attentional control, and decision-making [2–4]. In the realm of artificial intelligence, attention

mechanisms akin to salience assignment have propelled recent advances by enhancing the ability

of transformer models to extract relevant signals and suppress noise within vast datasets [5,6].

However, the neural mechanism underlying salience assignment in the brain remains elusive.

Previous studies have shown that acetylcholine (ACh) is a key neuromodulator critical for

conscious awareness, salience, and attentional control, and is essential for associative learning and

memory [7–15,15,16,16–19]. Cholinergic deficiencies represent a prominent feature of attention-related

impairments observed in Alzheimer's disease and Parkinson's dementia [20,21]. Selective lesion of

cholinergic neurons, pharmacological blockade of ACh receptors, or inhibition of cholinergic

neural activity all result in impairment of sensory discrimination, attention, memory, and

conditioning performance [16,22–24]. Activation of cholinergic neurons has been shown to facilitate

sensory discrimination and associative learning [16,18,25]. Cholinergic neurons in the basal forebrain

also respond to both positive and negative reinforcers [11–13,26].

The basal forebrain cholinergic neurons project predominantly to three main targets: the cerebral

cortex, hippocampus, and basolateral amygdala (BLA) [27–29]. Stimuli that evoke strong emotions,

or are associated with values are more likely to be perceived as salient. The BLA is particularly

pivotal in processing emotional and motivational information, associating cues with valenced

outcomes, and maintaining emotional salience [27,30–33]. The principal neurons in the BLA have been

shown to encode both positive and negative values of stimuli [32,34]. Recent studies have begun to

reveal the essential role of BLA ACh in cue-reward learning, fear conditioning, and reinforcement

prediction error [12,13,15,25,35]. However, the coding of various forms of behaviorally relevant salience

in the BLA ACh has not been systematically investigated. Importantly, a fundamental question is

how the behavioral salience, which is represented by the ACh level, is regulated and contributes

to associative learning.

To address how the brain solves the salience-assignment problem, we attempted to identify the

neural circuit that could tailor the ACh level in the BLA. We first recorded dynamic changes of

ACh release in the BLA during the mouse behavior. We demonstrated that BLA ACh is robustly

activated by a variety of behaviorally salient events, including novel ("unfamiliar") stimuli,

reinforcers and their predicting cues that gain salience, dependent on animals' learning experience,

and changes in task engagement and homeostatic states. We further identified that different

neuronal subpopulations in the striatal region nucleus accumbens (NAc), D1- and D2-receptor-

expressing medium spiny neurons (MSNs), opposingly regulate the release of BLA ACh and affect

associative learning, via disinhibiting and inhibiting the basal forebrain cholinergic neurons in the

substantia innominata (SI), respectively. Our findings uncover a unique role for the basal ganglia

in regulating the salience assignment, a process crucial for adaptive and efficient associative

learning.

RESULTS

BLA ACh encodes multiple forms of behavioral salience

We set out to systematically characterize the *in vivo* response properties of BLA ACh to a battery

of behaviorally salient events. For this purpose, we injected adeno-associated virus (AAV)

expressing the genetically encoded fluorescent ACh sensor (GRAB-ACh3.8m, abbreviated as

ACh3.8m) [36] into the BLA, and then used fiber photometry to record intrinsic dynamics of the

ACh signal in head-fixed behaving mice (Figure 1A). Behavioral salience can be influenced by

various factors, such as novelty, stimulus intensity, and behavioral relevance [37,38].

First, we presented water-restricted mice with white noise (novel stimulus) or water reward (Figure

1B). BLA ACh responses to the water delivery were robust and sustained across trials, whereas

the responses evoked by white noise showed a gradual decline (Figures 1C-1F and S1A-S1D).

Next, we presented mice with different magnitudes of reward (water) or punishment (air puff)

(Figure S1E). BLA ACh responses scaled proportionally with the magnitude of the reward or

punishment (Figures 1G, 1H, S1F, and S1H). Importantly, this ACh response profile did not reflect

the rate of licking or eye blinking (Figures S1F-S1O), thus differed from the cortical ACh that

tracks animals' motor-categorized arousal levels [39].

To further test whether behavioral relevance affects BLA ACh responses, we trained mice to

associate an auditory cue (conditioned stimulus, CS) with either an appetitive or aversive outcome

(unconditioned stimulus, US) in a Pavlovian conditioning task (Figure 1I). The CS responses of

BLA ACh increased significantly after learning (Figures 1J-1M). The CS responses can distinguish

"early" versus "late" phases, but not as much for "reward" versus "punishment" cues (Figures 1N

and 1O). Notably, cue responses of BLA ACh preceded the emergence of anticipatory licking

during CS (Figures S2A-S2C) and did not track licking behavior (Figures 2D and 2E), resembling

the temporal dynamics observed in dopamine signaling during associative learning [40].

Prediction error (PE), representing deviations from our expectation, is a salient signal that captures

our attention and drives adaptive learning. To test whether BLA ACh can encode PE, we recorded

BLA ACh in different settings of associative learning, pairing cues with probabilistic outcomes

(Figure S2F). We found that the uncued (or surprised) reward evoked relatively larger ACh

responses (Figures S2G, S2I, and S2J), which is a feature of PE. However, the surprised

punishment did not induce larger responses of BLA ACh (Figures S2H, S2K, and S2L). These

results indicate that BLA ACh is particularly sensitive to the salience of unexpected positive

reinforcement events.

BLA ACh encodes motivational salience in a context-dependent manner

The salience of behaviorally relevant stimuli can be shaped by animals' motivation, influenced by

both the external behavioral setting and their internal homeostatic state. To understand how

behavioral context affects BLA ACh signals, we first trained mice to perform an operant "go/no-

go" task, where mice need to lick to the "go" cue (CS_G) to receive a reward and withhold licking

to the "no-go" cue (CS_N) to avoid punishment (Figure 2A). With learning, the ACh response to

the CS_G increased significantly, whereas the response to the CS_N remained consistently low

(Figures 2B-2F). Next, we trained mice to perform an active avoidance task, where mice need to

lick in response to the "avoid" cue (CS_A, the same auditory stimuli as the "no-go" cue) to avoid

air puff delivery (Figure 2G). In the early learning period, the ACh response to the CS_A was

relatively weak (Figures 2H and 2J). Notably, the ACh response increased significantly to the CS_A

after learning (Figures 2I, 2K, and 2L-2O). In both tasks, we used the same auditory tone for "no-

go" and "avoid" trials, yet we observed strikingly different ACh responses (Figure 2P). In the "no-

go" trials, mice displayed passive "no-lick" behavior, whereas in the "avoid" trials, they exhibited

heightened behavioral engagement. These observations strongly suggest that BLA ACh encoding

of salience is contingent upon the motivational engagement of the animal.

To understand how homeostatic state affects BLA ACh signals, we examined the impact of thirsty

versus quenched states on the CS- and US-evoked ACh activity. The cue predicting the availability

of water elicited robust anticipatory licking in well-trained, thirsty mice (Figure 2Q, left). After

these mice were allowed to drink freely until quenched, the same auditory cue no longer elicited

anticipatory licking (Figure 2Q, left). We recorded BLA ACh activity in both states and observed

a significant reduction in reward CS-evoked ACh activity in quenched mice (Figures 2Q and 2R),

consistent with a decrease of salience for the water-predicting cue in quenched mice. Additionally,

we observed that BLA ACh responses to shock became significantly larger in the quenched state

than in the thirsty state (Figures 2S and 2T), indicating that shock became more salient when

homeostatic needs were met.

Taken together, these results demonstrate that BLA ACh activities represent the dynamics of

behavioral salience tracked by animals' motivational engagement across various behavioral

contexts or homeostatic states.

Opposing effects of NAc D1 and D2 MSNs on ACh release in the BLA

We next want to explore how the salience, as represented by BLA ACh activity, is modulated. As

previously acknowledged, BLA ACh originates from the basal forebrain, which includes the

nucleus basalis of Meynert, diagonal band of Broca, medial septum, and SI [25,27,28]. It has long been

known that basal forebrain cholinergic neurons receive the most prominent inputs from the NAc

[41–44], but the function of this connection has not yet been explored. We hypothesized that the NAc

might regulate BLA ACh levels via its projection onto the basal forebrain cholinergic system. To

test this, we injected AAV with Cre-inducible expression of the light-gated cation channel

channelrhodopsin-2 (ChR2) into the NAc of *D1-Cre* or *D2-Cre* mice to express ChR2 in two major

neuronal populations of the NAc, and injected AAV expressing ACh sensor (ACh3.8m) into the

BLA (Figures 3A, 3B, 3H, and 3I). Optical fibers were subsequently implanted in the NAc and

BLA for photostimulation and fiber photometry, respectively (Figure S3A). To avoid the laser

artifacts to the photometry signal, we interleaved the photometry recording and optogenetic laser

stimulation at the frequency of 20 Hz (Figure S3B; see Methods).

To assess whether NAc D1 MSNs regulate BLA ACh, we activated NAc D1 MSNs while

simultaneously recording BLA ACh release. We presented the laser stimulation with durations of

0, 0.5, 1.0 or 2.0 seconds interleaved in one session (Figures S3C-S3E). We observed that

activation of NAc D1 MSNs induced robust elevation of BLA ACh responses (Figures 3C and

S3C). The ACh responses scaled with the duration of laser stimulation, with longer durations

resulting in larger area under the curve (AUC) of ACh responses (Figures 3C-3G and S3C). As to

the control mice with GFP expression in the NAc, the laser stimulation had no effect on BLA ACh

responses (Figures S3E-S3K). These data reveal that activation of NAc D1 MSNs robustly

increases BLA ACh levels.

Next, we set out to test whether stimulation of NAc D2 MSNs affects BLA ACh levels. Employing

a similar approach to that used with NAc D1 MSNs, we optogenetically activated NAc D2 MSNs

and monitored BLA ACh signal with fiber photometry (Figures 3H and 3I). Interestingly, we found

that activation of NAc D2 MSNs evoked potent suppression of BLA ACh responses (Figures 3J

and S3D). The suppression effect scaled with the duration of laser stimulation, with longer

durations inducing a greater suppression effect (Figures 3J-3N). These results show that NAc D2

MSNs potently decreased BLA ACh levels. Together, these data demonstrate that distinct

populations of NAc MSNs have opposing effects in modulating ACh release in the BLA.

NAc MSNs modulate ACh release in the BLA via their projections to the SI

To address what circuit mechanisms account for the accumbal modulation of ACh release, we

performed anterograde tracing for NAc D1 and D2 MSNs, and observed dense axonal terminals

in the SI (Figures S3L-S3N). Although basal forebrain ChAT neurons project to many target

regions, BLA-projecting ChAT neurons are preferentially localized in the SI (Figures S4A-S4C)

and the population of BLA-projecting SI ChAT neurons predominantly targets the BLA only

(Figures S4D-S4F), suggesting potentially unique anatomical and functional characteristics. As a

first step to test whether activation of SI ChAT neurons induces ACh release in the BLA, we

expressed ChR2 in SI ChAT neurons and ACh sensor in the BLA (Figures S4G and S4H). We

found that brief stimulation of SI ChAT neurons induced robust ACh release in the BLA (Figures

S4I-S4K). The enhancement effect scaled with the duration of laser stimulation, with longer

durations resulting in a greater enhancement effect (Figures S4K-S4N).

To further examine the causal effect of NAc→SI pathway on BLA ACh release, we sought to

perform optogenetic stimulation of the NAc→SI pathway while simultaneously recording BLA

ACh release. To this end, we infected NAc D1 or D2 MSNs with ChR2, and implanted optical

fibers in the SI above the axons originating from the infected neurons, and infected BLA neurons

with ACh sensor, and implanted optical fibers in the BLA to detect ACh release (Figures 4A, 4B,

4F, and 4G). Similar to the stimulation of cell bodies of NAc D1 and D2 MSNs, we found robust

increase and decrease of BLA ACh responses induced by activation of the $NAc^{D1} \rightarrow SI$ and

$NAc^{D2} \rightarrow SI$ pathways, respectively (Figures 4C-4E and 4H-4J). These data show that NAc MSNs

modulated BLA ACh levels via the projections to the SI.

Activation of SI cholinergic neurons and $SI^{ChAT} \rightarrow BLA$ pathway does not affect valence

Further, we aim to test whether stimulating SI ChAT neurons would drive any form of preference

or aversion. To achieve this, we expressed ChR2 selectively into SI ChAT neurons, and positioned

the optic fibers above the SI region for laser delivery (Figure S5A). To assess the effects, we

subjected these *ChAT-Cre* mice to a real-time place preference test, wherein photo-activation of

ChAT neurons was contingent on their entry into a particular side of a chamber (Figure S5B). We

found that activation of SI ChAT neurons did not prompt any preference for a specific location

(Figure S5C), and it had no impact on the movement speed (Figure S5D) or distance covered by

the mice (Figure S5E). Furthermore, we explored whether optogenetic stimulation of the

$SI^{ChAT} \rightarrow BLA$ pathway would induce any observable valence-related behaviors (Figure S5F).

Consistent with prior observations [15,25], we found that activation of the $SI^{ChAT} \rightarrow BLA$ pathway did

not induce any place preference (Figures S5G and S5H), or affect the movement (Figures S5I and

S5J). In addition, activation of this pathway failed to support self-stimulation (Figures S5K-S5N).

Taken together, these observations provide evidence that BLA ACh does not actively drive

valence-related responses.

Activation of NAc→SI pathway had no effect on cortical and hippocampal ACh release

To ask whether the stimulation effect of the NAc→SI pathway is selective to ACh release in the

BLA, we assessed ACh release in other brain regions originating from the basal forebrain. In

addition to the BLA, SI ChAT neurons also project to various other target regions, including the

cerebral cortex and hippocampus (Figures 5A-5C). Specifically, we chose to optogenetically

manipulate the NAc→SI pathway and monitor the ACh signal in cortical regions, including the

medial prefrontal cortex (mPFC) and auditory cortex (AUD), which have been extensively

characterized for ACh activity, as well as in the CA1 region of the hippocampus [27]. We expressed

ChR2 in the NAc D1 or D2 MSNs, and implanted optic fibers in the SI for pathway stimulation,

and expressed ACh sensor in the mPFC, the AUD, and the CA1 for ACh monitoring (Figures 5D,

5G, and 5J). Both reward and punishment induced a robust increase of ACh release in the mPFC,

the AUD, and the CA1 (Figure S6). However, in contrast to the effect observed in the BLA, the

NAc^{D1}→SI stimulation did not evoke an immediate and potent increase of ACh signal in the mPFC,

the AUD, and the CA1 (Figures 5E, 5H, 5K, and 5M; S7A, S7C, S7E, S7G, S7I, and S7K). The

amplitude of ACh responses in the cortex and hippocampus was quite small, compared to that of

the BLA (Figure 5M). Although there were some detectable changes in the mPFC, AUD, and the

CA1, the peak time lagged the laser stimulation significantly (Figure 5N), suggesting an indirect

effect on these areas. Activation of NAc^{D2}→SI pathway did not induce any changes to ACh release

in the mPFC, the AUD, and the CA1 (Figures 5F, 5I, 5L, and 5O; S7B, S7D, S7F, S7H, S7J, and

S7L). Notably, the suppression effect is only present in the BLA for activating NAc D2 MSNs

(Figure 5O), highlighting a unique role of D2 MSNs in modulating BLA ACh levels. These results

demonstrate that the modulatory effect of NAc→SI pathway is specific to the BLA.

SI ChAT neurons form direct connection preferentially with NAc D2, but not D1 MSNs

As distinct subpopulations of NAc MSNs have opposing modulatory effects on the BLA ACh

levels, we next sought to understand whether and how NAc D1 and D2 MSNs are connected

differentially to SI ChAT neurons. Given that both NAc D1 and D2 MSNs are GABAergic and

release inhibitory neurotransmitters, it is unlikely for the majority of D1 MSNs to form direct

monosynaptic connections with SI ChAT neurons. Conversely, D2 MSNs are more likely to have

direct monosynaptic connections with these neurons. We proposed that NAc D1 MSNs could

activate SI ChAT neurons by inhibiting intermediate, tonically active GABAergic afferents to

ChAT neurons, while D2 MSNs directly inhibit ChAT neurons.

To test this prediction, we performed high-resolution volumetric imaging of NAc axon terminals

and ChAT cell bodies to directly visualize the connections between NAc MSNs and SI ChAT

neurons. By injecting an AAV expressing a membrane-targeted green fluorescent protein (mGFP)

and synaptophysin, which is found on nearly all presynaptic vesicles, fused with a red fluorescent

protein, mRuby, in a Cre-dependent manner into the NAc of *D1-Cre* or *D2-Cre* mice (Figures 6A

and 6D), we could identify the synaptic targets of NAc MSNs. This method confines mRuby

expression to pre-synaptic terminals, enabling us to differentiate between potential synaptic targets

and passing axons. We observed robust labeling of synaptic terminals originating from NAc MSNs

in the SI (Figures 6B, 6C, 6E, and 6F). To quantify the number of synaptic connections, we counted

the red spots present within 1.0 μm distance from ChAT neurons, which were immunostained with

green fluorescence (Figure 6G). We found that the number of synaptophysin-mRuby spots around

ChAT neurons for D2 MSNs was significantly higher than that for D1 MSNs (Figure 6H),

indicating that D2 MSNs predominantly target SI ChAT neurons, thereby inhibiting their ACh

release.

To confirm that BLA-projecting SI ChAT neurons indeed receive inputs from the NAc, we utilized

an intersectional monosynaptic retrograde rabies system, which was injected into the SI of ChAT-

Cre mice in a pathway- and cell-type-specific manner (Figures 6I and 6J). We found that BLA-

projecting SI ChAT neurons received substantial inputs from the NAc (Figure 6K). To further

characterize the specific cell types of NAc MSNs with monosynaptic inputs onto SI ChAT neurons,

we performed single-molecule *in situ* hybridization on retrogradely labeled NAc MSNs infected

by rabies virus (Figure 6L). We found that the vast majority of rabies virus-labeled cells expressed

the dopamine receptor gene *Drd2*, but not *Drd1* (Figure 6M). This data shows that NAc D2 MSNs

directly targeted SI ChAT neurons, which could inhibit ACh release in the BLA. In the basal

forebrain, the vast majority of neurons are cholinergic and GABAergic [45]. ChAT neurons receive

inputs from local GABAergic neurons [43] and can be inhibited by local GABAergic neurons [17]. To

further test whether NAc D1 MSNs could disinhibit SI ChAT neurons, we recorded SI ChAT

neural activities while activating the $NAc^{D1} \rightarrow SI$ pathway (Figure 6N). We observed that activation

of the $NAc^{D1} \rightarrow SI$ pathway resulted in a potent increase in SI ChAT neural activities (Figures 6O

and 6P). Our results thus support the notion that NAc D1 MSNs inhibit local SI GABAergic

neurons, resulting in disinhibition of ChAT neurons, hence increased ACh release in the BLA.

Activation of $NAc^{D1} \rightarrow SI$ facilitates and $NAc^{D2} \rightarrow SI$ delays associative learning

Learning how environmental stimuli predict the availability of food and other natural rewards is

critical for survival. Salient stimuli capture attention, facilitating associative learning. If the NAc

modulates the salience representation of BLA ACh, then activation of the NAc→SI pathway,

which either enhances or suppresses BLA ACh levels, should affect the efficiency of cue-reward

learning. To this end, we trained mice to perform a cue-reward learning task as described before

[25]. In this task, water-restricted mice were trained to perform a correct nose poke when signaled

by an auditory cue to receive a water reward on a variable intertrial interval (Figures 7A and 7B).

We first characterized the *in vivo* activity of BLA ACh in this task (Figure S8). We observed that

BLA ACh release occurred in response to various events across the learning of the task (Figures

S8A and S8B). In rewarded trials of the pre-training stage, the highest levels of ACh release

occurred immediately after the reward retrieval (entry into the reward port) (Figures S8A and S8B).

As the training began, ACh release during reward trials showed a marked shift toward the time

before nose poking reward port (Figures S8A and S8B). As animals became proficient in the task,

there was a significant increase in the magnitude and a significant decrease in the peak latency of

ACh release upon cue presentation (Figures S8C-S8F). This indicates that as animals learned the

cue-reward contingency, the cue became a more salient event. At this time point, there was still a

peak of ACh release at the time of reward, but its magnitude was diminished, and the peak occurred

earlier (Figure S8G). Consistent with previous findings [25], the ACh responses shifted as mice

gradually learned the task. This shows that BLA ACh is actively involved in the reward learning

association.

Given the observed shift of BLA ACh towards tone onset during acquisition of cue-reward

learning, we hypothesized that BLA ACh may potentiate learning the cue-reward contingency. To

test this, we optogenetically activated the NAc^{D1}→SI pathway in a closed-loop manner, contingent

on the correct nose poke and reward retrieval, thereby boosting BLA ACh release during behavior.

Activation of NAc^{D1}→SI pathway significantly facilitated animals' learning rate (Figures 7C-7G).

Conversely, activation of NAc^{D2}→SI pathway significantly delayed mice's reward learning

(Figures 7H-7L). Since reaction time tends to be shorter when stimuli are more salient [46,47], we

used reaction time (from cue onset to port entry) as an indicator of stimulus salience, measured

during periods without laser stimulation. We found that reaction time was significantly shorter in

D1-Cre mice compared to controls (Figures S9A-S9C), while it was notably prolonged in *D2-Cre*

mice (Figures S9F-S9H). This data suggests that activation of $NAc^{D1} \rightarrow SI$ pathway promotes

salience, leading to quicker responses in mice, whereas $NAc^{D2} \rightarrow SI$ pathway suppresses salience,

resulting in slower responses. Considering the inhibitory role of D2 MSNs in the dorsal striatum

on movement, we speculated whether impaired reward learning could be attributed to movement

deficits. However, our results indicated that activation of NAc D2 MSNs did not significantly

affect general poking behavior (Figures S9I and S9J). Together, our results reveal that distinct

NAc neural populations opposingly modulate the BLA ACh levels, leading to opposite changes in

the learning rate in a cue-outcome contingency setting.

DISCUSSION

Previous studies of selective attention and the salience of behaviorally relevant stimuli have

primarily focused on the frontoparietal cortex and the thalamus [1,38,48,49]. Here we have identified

the NAc, located within the ventral striatum of the basal ganglia, as a key regulator of salience

assignment. Specifically, we demonstrate that two distinct subpopulations of striatal MSNs exert

opposing control over salience signals via modulating SI ChAT neurons (Figures 3 and 4). The

NAc is uniquely positioned at the crossroads of emotional, cognitive, and motor circuits, enabling

the integration of diverse information [50,51]. While D1 and D2 MSNs in the dorsal striatum are

segregated based on their respective projection targets, NAc D1 and D2 MSNs project to several

overlapping downstream targets, including the SI (Figures 4B, 4G, and S3L-S3N). Then how do

NAc D1 and D2 MSNs exhibit opposing roles in salience assignment? Given the diverse cortical

and limbic inputs received by the NAc [52–54], future investigations are warranted to delineate the

contributions of individual inputs while examining salience control by NAc MSNs.

Salience, defined as the absolute (i.e., unsigned) value of a stimulus, is a concept often intertwined

with valence, making their neural separation challenging. Activation of NAc D1 or D2 MSNs can

induce potent valence-related responses, either rewarding or aversive [54–57]. In contrast, stimulation

of SI ChAT neurons or the SI→BLA pathway failed to evoke any valence-specific responses

(Figure S5), consistent with ACh's representation of salience rather than valence. This suggests

that the rewarding and aversive responses may be mediated by other downstream targets of the

NAc or the SI, such as the lateral hypothalamus and ventral tegmental area [58–61]. Our results

highlight the ability to disentangle salience from valence at the neural circuit level.

Associative learning, a fundamental cognitive process that allows organisms to extract

relationships among environmental cues, actions, and their outcomes, underlies the formation of

memories and prediction of future events [49,62]. To achieve this, our brains need to not only form

and store these associations but also assign importance or "salience" to relevant cues in the

environment, thus facilitating learning and memory processes. Activation of the cholinergic

system is capable of modulating synaptic plasticity by altering synaptic strength and regulating

secretion of neurotrophic factors [14,63–66]. Through these mechanisms, BLA-projecting cholinergic

neurons may boost neural plasticity and thereby contribute to associative learning. In line with this

notion, we found that enhancing and suppressing ACh release in the BLA via manipulation of NAc

D1 and D2 MSNs facilitated and impaired reward associative learning, respectively (Figure 7). It

is widely acknowledged that the NAc is involved in various forms of learning and motivational

control. In particular, our results reveal the striatal involvement, mediated by NAc→SI pathway,

in the salience assignment as an important contributing aspect of learning and motivational control.

In the striatum, the interaction among local ChAT interneurons, dopaminergic inputs, and MSNs

has been extensively explored [67–69]. In the striatum, axons of midbrain dopamine neurons express

high levels of nicotinic ACh receptors [67]. Activation of these ACh receptors can directly trigger

action potentials in striatal dopaminergic axons, resulting in dopamine release [67,70,71]. Here we

showed that the striatal neurons interact with the extra-striatum cholinergic system, the basal

forebrain cholinergic system. The NAc MSNs modulated the SI cholinergic system potently, with

NAc D1 and D2 MSNs facilitating and suppressing ACh release in the BLA, respectively (Figures

3 and 4). Interestingly, dopamine in the BLA has recently been demonstrated to encode salience

[72]. Future studies will need to determine whether dopamine signals in the BLA are triggered by

the cholinergic inputs from the basal forebrain, and whether dopamine and ACh signals both

participate in salience assignment in a coordinated manner.

ACKNOWLEDGEMENTS

We thank M.M. Poo, X. Jin, B. Li, B. Hangya, A. Laszlo, N.L. Xu, and Y.G. Sun for comments

on earlier versions of the manuscript, and members of the Xiao laboratory for helpful discussions.

We thank Yulong Li for providing us with the ACh3.8 sensor. We thank Qian Hu (the Optical

Imaging Core at the CEBSIT) for providing assistance with the analysis of volumetric imaging

data. This work was supported by grants from the National Natural Science Foundation of China

(32371060 to X.X., 32271065 to H.D.), the Lingang Laboratory (LG-QS-202203-06 to X.X., LG-

AUTHOR CONTRIBUTIONS

398 A.C., H.D., and X.X. designed the study. A.C. conducted most of the experiments with help from

399 Y.L. and H.Z. A.C. analyzed data and prepared figures. H.G. helped with fiber photometry and

400 histology. Y.W. helped with the behavioral training. F.X. provided critical consultation and

401 feedback. X.X., A.C., and H.D. wrote the paper with inputs from all authors. X.X. supervised all

402 aspects of the work.

DECLARATION OF INTERESTS

405 The authors declare no competing interests.

RESOURCE AVAILABILITY

Lead Contact

409 Further information and requests for resources and reagents should be directed to and will be

410 fulfilled by the Lead Contact, Xiong Xiao (xiao@ion.ac.cn).

Data and code availability

The authors declare that the data supporting the findings of this study are available within the paper

and its supplementary information files. The custom codes that support the findings from this study

will be made available online.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All experiments were approved by the Animal Care and Use Committee of the Institute of

Neuroscience, Chinese Academy of Sciences (Center for Excellence in Brain Science and

Intelligence Technology, Chinese Academy of Sciences). We obtained *ChAT-Cre* (Stock No:

006410) mice from Jackson Laboratory. The *D1-Cre* (MMRRC_030989-UCD) and *D2-Cre*

(MMRRC_032108-UCD) mice were provided by Dr. Ninglong Xu at the CEBSIT. Male and

female mice (2-4 months old) were used for all the experiments. Mice were housed under a 12-h

light/dark cycle (7 a.m. to 7 p.m. light) in groups of 2-6 animals, with food and water available *ad*

libitum before being used for experiments in the animal facility at the Institute of Neuroscience.

Littermates were randomly assigned to different groups before experiments.

METHOD DETAILS

Immunohistochemistry

Immunohistochemistry experiments were conducted following standard procedures [32,73,74]. Briefly,

mice were anesthetized with isoflurane and perfused with PBS, followed by 4% paraformaldehyde

(PFA). Brains were extracted and post-fixed overnight in 4% PFA at 4 °C. Coronal sections (50-

μm) were cut using a vibratome (VT1200S, Leica). Sections were first washed in PBS (5 min),

then sections were blocked in 5% normal goat serum or BSA in PBST (0.1% Triton X-100 in PBS)

for 1 h at room temperature (RT) and then incubated with primary antibodies overnight at 4 °C.

Sections were washed with PBS (3 × 15 min) and incubated with fluorescent secondary antibodies

at RT for 2 h. Next, sections were washed twice in PBS, and then incubated with DAPI (Sigma,

Cat. No. d9542; dilution, 1:5000) for 10 min. After that, sections were mounted onto slides with

Fluoromount-G (Southern Biotech). Images were captured by a 10× objective fluorescent

microscope (Olympus, VS120).

The primary antibodies used were: chicken anti-GFP (Abcam, Cat. No. ab13970; dilution, 1:1000),

and goat anti-ChAT (Millipore, Cat. No. AB144P; dilution, 1:500). Appropriate fluorophore-

conjugated secondary antibodies (Jackson ImmunoResearch Laboratories) were used depending

on the desired fluorescence colors.

Stereotaxic surgery

All surgery was performed under aseptic conditions. Standard surgical procedures were used for

stereotaxic injection and implantation, as previously described [32,73,74]. Briefly, mice were

anesthetized with isoflurane (1-2% in a mixture with oxygen), and head-fixed in a stereotaxic

injection apparatus (Rotational Digital Stereotaxic Frame, RWD Life Science, Shenzhen, China).

For virus injection, the following stereotaxic coordinates were used for the NAc: 1.40 mm anterior

from bregma, 1.50 mm lateral from the midline, 4.60 mm vertical from skull surface; for the SI,

0.00 mm from bregma, 1.65 mm lateral from the midline, 5.10 mm vertical from skull surface; for

the BLA, 1.50 mm posterior from bregma, 3.40 mm lateral from the midline, 4.80 mm vertical

from skull surface; for the mPFC, 1.80 mm anterior from bregma, 0.50 mm lateral from the midline,

2.10 mm from skull surface; for the AUD, 2.10 mm posterior from bregma, 3.80 mm from the

midline, 2.80 mm from skull surface; for the CA1, 2.00 mm posterior from bregma, 1.40 mm from

the midline, 1.50 mm from skull surface.

For fiber photometry recordings, optic fibers (core diameter, 200 mm; N.A., 0.5; Inper, Hangzhou,

China) were implanted 50 μm above the coordinates for viral injection. For the optogenetic

experiments, optic fibers (core diameter, 200 mm; N.A., 0.37; Inper, Hangzhou, China) were

implanted 200 μm above the coordinates for viral injection. Then a metal head-bar (for head-

restraint) was subsequently mounted onto the skull with dental cement. We waited for a minimum

of 3 weeks before starting the experiments with these mice.

Viral vectors

The following adeno-associated viruses (AAVs) were obtained from Shanghai Taitool Bioscience

(Shanghai, China): AAV2/8-hSyn-EGFP-WPRE-pA, AAV2/8-hEF1α-DIO-hChR2-EYFP-

WPRE-pA, AAV2/8-hEF1α-DIO-hChR2-mCherry-WPRE-pA, AAV2/2Retro- Plus-hEF1a-DIO-

EGFP-WPRE-pA, AAV2/2Retro-CAG-FLEX-FlpO-WPRE-pA. The AAV2/8-EF1α-DIO-ChR2-

mCherry, AAV2/9-EF1α-DIO-EGFP, AAV-nEF1α-fDIO-EGFP, AAV2/9-hsyn-Flex-mGFP-

T2A-Synaptophysin-mRuby and AAV2/8-hSyn-ACh3.8 were produced by BrainCase (Shenzhen,

China). The components of the rabies viral tracing system were also produced by BrainCase:

rAAV-nEF1α-fDIO-EGFP-T2A-TVA, rAAV-nEF1α-fDIO-oRVG, AAV2/9-EF1α-DIO-EGFP-

T2A-TVA, AAV2/9-EF1α-DIO-N2c-G, RV-CVS-EnVA-ΔG-tdTomato (the modified rabies

virus). All viral vectors were aliquoted and stored at −80°C until use.

Mapping monosynaptic inputs with pseudotyped rabies virus

To map monosynaptic inputs onto ChAT$^{SI→BLA}$ neurons, we implemented a pathway-specific and

cell-specific tracing strategy utilizing a refined rabies virus system, as previously detailed [32,74]. We

injected the BLA of *ChAT-Cre* mice with the AAVretro-Flex-FlpO, which allows selective

expression of FlpO within ChAT$^{SI \to BLA}$ neurons. Subsequently, in the same surgery, we injected

the SI of these mice with a mixture of AAV-fDIO-TVA-EGFP and AAV-fDIO-oG (1:2 in volume,

for a total of 0.1~0.2 µl) that express the following components in a Flp-dependent manner: a

fluorescent reporter EGFP, TVA (which is a receptor for the avian virus envelope protein EnvA),

and the rabies envelope glycoprotein (oG). Three weeks after the first injection, mice were injected

in the SI with RV-CVS-EnVA-ΔG-tdTomato (0.1 µl), a rabies virus that is pseudotyped with EnvA,

lacks the envelope glycoprotein, and expresses tdTomato. This variant of the rabies virus has been

demonstrated to exhibit enhanced retrograde trans-synaptic transfer and reduced neurotoxicity.

Histological examination of brain tissue was conducted one week post-rabies virus administration.

Fluorescent *in situ* hybridization

Single molecule fluorescent *in situ* hybridization was employed to detect the expression of *Drd1*

and *Drd2* mRNAs in the nucleus accumbens (NAc) of adult *ChAT-Cre* mice following rabies virus

injection, with tdTomato expression in NAc neurons projecting to the substantia innominata (SI).

For tissue preparation, mice were anesthetized with isoflurane and perfused with DEPC-PBS and

4% PFA. Brains were taken out and fixed overnight at 4 °C and dehydrated by 30% sucrose in

DPEC-PBS. Brains were sectioned at 20 µm and sections containing the NAc were collected and

stored at -80°C until processing. Hybridization was carried out using the RNAscope kit (ACDBio).

On the day of the experiment, sections were washed once in RNA-free H$_2$O (1 min), and

dehydrated using 100% ethanol (twice, 3 min each). Sections were then dried at RT and incubated

with Hydrogen Peroxide for 20 min. Retrieval reagents were then applied to the sections for 5-7

min, followed by a wash with 100% ethanol (twice, 3 min each) and incubation with Protease III

for 30 min at RT. Sections were washed in PBS for 5 min at RT before hybridization. Probes

against *Drd1* (Cat. No. #406491, dilution 1:50) and *Drd2* (Cat. No. #406501, dilution 1:50) were

applied to NAc sections, followed by hybridization at 40°C for 2 hours. After hybridization,

sections were washed twice in wash buffer (2 min each) at RT, then incubated with three

consecutive rounds of amplification reagents (30 min, 30 min and 15 min at 40°C). Finally,

fluorescence detection was carried out by HRP-TSA system. To visualize tdTomato-expressing

(rabies virus-infected) cells, we performed immuno-staining using primary antibody rabbit anti-

dsRed (Takara Bio, Cat. No. 632496, 1:300) and secondary antibody goat anti-rabbit, Cy3

(Jackson ImmunoResearch, Cat. No. 111-655-003, 1:2,000) after *in situ* hybridization. Finally,

sections were incubated with DAPI for 2 min, and then mounted with coverslip using mounting

medium. Images were acquired using a 40× objective confocal microscope (Nikon, C2) and

visualized and processed using ImageJ.

Measurement of synaptophysin spots

Synaptic terminals in the SI were labeled using synaptophysin-mRuby, a presynaptic marker.

Volumetric images were captured by 40× objective confocal microscopy (FV3000, Olympus).

Subsequently, we utilized Imaris 8.1 software (Bitplane) for neuronal cell body and synaptophysin

spot identification. During the analysis, we meticulously adjusted the diameter parameter to ensure

the accurate detection of ChAT neurons, filtering out those close proximity to the image boundary.

To achieve 3D segmentation of ChAT neurons, we utilized the Imaris surface and mask selection

feature. Following this step, we employed ImageJ 3D Fast Filters to expand the cellular boundary

by 1.0 µm. Synaptophysin spots were detected based on their distance from the surface of ChAT

neuron cell body. Only spots located within a maximum distance of 1.0 µm from the surface of

ChAT neurons were considered positive and included for further analysis. This procedure was

conducted in three-dimensional space to better represent the complex architecture of the synaptic

connectivity. Finally, ImageJ 3D Manager was employed to quantify the number of synapse

termini spots surrounding individual ChAT neurons.

Markerless pose tracking

We used DeepLabCut software (v.2.2.0.5) for tracking the eyes of mice during behavior. The

training data set included 120 frames from 6 videos across 6 mice. These frames were meticulously

selected to encompass a diverse range of eye sizes and positions, ensuring robust model training

and generalization. For eye tracking, we strategically placed eight tracking points evenly

distributed along the periphery of the eye. This configuration enabled comprehensive tracking of

eye movements and provided sufficient spatial resolution for capturing subtle variations in eye size

and position. The training of the model was conducted over 1.0 million iterations with default

parameters (trained on 95% of labeled frames, initialized with ResNet-50, batch size of 8).

Real-time place preference or aversion test

Freely moving mice were initially habituated to a two-sided chamber ($20 \times 42 \times 30$ cm) for 10

minutes to assess baseline preferences. In the subsequent test sessions, lasting 10 minutes each,

one side of the chamber was designated as the photo-stimulation side, while the other served as

the non-stimulation side. The order of stimulation sides was counterbalanced across mice. Upon

entry into the stimulation side, photo-stimulation was initiated using a 473-nm laser (Shanghai

Laser & Optics Century, Shanghai, China), delivering 15-ms pulses at 20 Hz and 10 mW intensity

(measured at the tip of optic fibers). Laser stimulation ceased as the mouse exited the stimulation

side. The behavior of the mice was videotaped with a CCD camera interfaced with Bonsai software

(Bonsai), which was used to control the laser and extract real-time behavioral parameters,

including position, time, distance, and velocity.

Self-stimulation test

Freely moving mice were individually placed in a custom-designed chamber equipped with two

distinct ports. One port, termed the active port, was designated to trigger photo-stimulation (15-

ms pulses, 20 Hz, 10 mW; $\lambda = 473$ nm) for 2 s upon poking. In contrast, poking into the other port,

termed the inactive port, did not elicit any photo-stimulation. Mice were allowed to freely explore

and poke both ports during two separate 1-hour testing sessions, conducted on consecutive days.

The assignment of the active port in each session was systematically counterbalanced across

experimental subjects.

Pavlovian conditioning task

Three weeks after surgery, water-deprived mice were trained on an auditory Pavlovian

conditioning task, during which the mice were head restrained using custom-made clamps and

head-bars mounted on the skull. Prior to training, each mouse underwent a one-day habituation

period to acclimate to the head-restraint setup. During training, each trial began with a conditioned

stimulus (CS), which was a 0.5-s sound (12 kHz or 3 kHz), followed by a 1-s delay and then an

unconditioned stimulus (US; the outcome). The US was either a water reward (5 µl) or an electric

shock (0.02 mA, 100 ms). The electric shock was administered towards the tail of the animals.

Once mice reached a performance level of at least 80% successful trial, mice were subjected to the

probabilistic conditioning task. The trials were categorized into three types: predicted US (70%),

US omission (15%), and unpredicted US (15%). In predicted US trials, a 0.5-s sound was followed

by a predicted reward (water) or punishment (shock). In US omission trials, a 0.5-s sound was

presented but without subsequent reward or punishment. In unpredicted US trials, a reward or

punishment was delivered without preceding CS introduction. These contingencies were carefully

calibrated to keep mice motivated for the task. Different trial types were presented in a

pseudorandomized sequence.

Water rewards were dispensed via a metal spout positioned in front of the mice's mouths, which

also functioned as part of a custom "lickometer" circuit. This circuit registered a lick event each

time a mouse completed the circuit by licking the spout. Experimental control, including the

delivery of CSs and USs, as well as the recording of licking events, was managed by a custom

software written in MATLAB (MathWorks, Massachusetts) interfaced with a Bpod State Machine

(Sanworks, NY).

Go/no-go task

Water-deprived mice were trained in an auditory go/no-go task under head restraint. The training

started with habituation, during which mice received water rewards by licking the water spout (2

μl for each lick), with no auditory stimuli presented. Once mice reliably licked the spout (2-3 days),

they were subjected to the go/no-go training, consisting of both "go" and "no-go" trials. In go trials,

a 12-kHz tone (termed the "go cue", 0.5 s in duration; CS_G) was delivered, followed by a 1-s delay

("response window"). Licking during this window resulted in a water reward (5 μl). Conversely,

in no-go trials, a different 3-kHz tone (termed the "no-go cue", 0.5 s in duration; CS_N) was

delivered, followed by a response window (1 s). Liking during this period was punished by an air

puff blowing to the face. The go and no-go trials were randomly interleaved. For analysis, trials

were sorted into go trials and no-go trials. A correct response during a go trial ("hit") occurred

when the mouse successfully licked the spout during the response window and subsequently

received the water reward. A correct response during a no-go trial ("correct rejection") occurred

when the mouse successfully refrained from licking during the response window and thus avoided

the air puff.

Active avoidance task

The task was designed to train mice to actively avoid punishment. In avoid trials, a 3-kHz tone

(termed the "avoid cue", 0.5 s in duration; CS_A, the same tone as CS_N) was presented, followed by

a 1-s decision window. If mice lick the spout during the decision window, they would avoid an

unpleasant air puff (200 ms) blowing to the face. Otherwise, mice would receive the air puff

immediately after the decision window. Animals were trained one session per day, with each

session consisting of 100 trials.

Cue-reward learning task

Freely moving water-deprived mice underwent a cue-reward learning task within a chamber

equipped with three ports: left (reward port), middle (inactive port), and right (active port). Nose

pokes were detected via infrared beam breaks. In the habituation phase of the task, mice could

receive a reward with a nose poke in either of the ports. During the Pre-training phase, the trial

starts with a 6-s auditory cue (the "response window"), mice need to poke the active port during

the 6-second window. Upon successful poking, a 4-s reward collection window was followed,

indicated by an illuminated LED light in the reward port. Only nose pokes into the reward port

during this window led to reward delivery. In the Pre-training phase, there was no punishment for

improper nose pokes, neither in the active port outside the tone (incorrect nose pokes) nor in the

inactive port (inactive nose pokes). But in the training phase of the task, the nose pokes during the

inter-trial-interval (ITI) would result in a 2-s timeout and the restart of the ITI period. For the

optogenetic experiments, the photo-stimulation (1 s, 15-ms pulses, 20 Hz, 10 mW; $\lambda = 473$ nm)

was delivered in the SI when the mice correctly poked the active port.

***In vivo* fiber photometry and data analysis**

To record the activities of acetylcholine *in vivo* in behaving animals, we used a commercial fiber

photometry system (Inper, Hangzhou, China) to measure fluorescence signals in these neurons

through an optical fiber (core diameter, 200 µm; Inper, Hangzhou, China). A patch cord (core

diameter, 200 µm; Inper, Hangzhou, China) was used to connect the photometry system with the

implanted optical fiber. The intensity of the blue light ($\lambda = 470$ nm) for excitation was adjusted to

a low level (20~40 µW) at the tip of the patch cord. Emitted fluorescence was bandpass filtered

and focused on the sensor of a CCD camera. Photometry signals and behavioral events were

aligned based on an analog TTL signal generated by the Bpod. Recorded data were exported to

MATLAB for further analysis.

To correct for photobleaching of fluorescence signals (baseline drift), a bi-exponential curve was

fit to the raw fluorescence trace and subtracted as follows:

$F_{raw\_fit} = fit(Timestamp, F_{raw}, 'exp2')$

$$F_{raw\_correction} = \frac{F_{raw} - F_{raw\_fit}}{F_{raw\_fit}}$$

After baseline drift correction, the fluorescence signals were z-scored relative to the mean and

standard deviation of signals during the baseline.

Simultaneous fiber photometry and optogenetic stimulation

To perform simultaneous fiber photometry recording and optogenetic activation experiments, we

devised a custom trigger system (based on the ThinkerTech photometry system) capable of

delivering laser stimulation outside the photometry signal acquisition time window. Our

photometry system employs two sampling channels: green fluorescence channel (blue LED, λ =

470 nm) and isosbestic channel (violet LED, λ = 410 nm). Laser stimulation is administered using

either a blue laser (470 nm for ChR2) or a red laser (593 nm for ChrimsonR). The photometry TTL

signal indicates the time window for photometry signal sampling, while the TTL trigger for laser

stimulation is gated by both the control trigger from the Bpod and the low level of the photometry

TTL. We sampled the fluorescence and isosbestic signals at a frequency of 20 Hz. During each

50-ms time window, we sequentially deliver blue LED (470 nm), violet LED (405 nm), and laser

(blue laser, 470 nm; red laser, 593 nm) stimulation to the target area. This strategy effectively

avoided any potential interference from laser artifacts on the photometry recordings.

ROC analysis

We used the Receiver Operating Characteristic curve (ROC) to evaluate BLA ACh decodability

of two trial types or outcomes. The ROC is a non-parametric statistic calculated by sweeping a

classification threshold from the smallest to the largest data value and measuring the increase of

true and false positives. This sweep yields a curve of values. To condense this information to a

single number, the area under the ROC was measured and was referred to as auROC. The statistical

significance of auROC (whether it was significantly different from the chance level of 0.5) was

determined using the permutation test by shuffling trial labels (1000 permutations).

Generalized linear model (GLM)

We fit a linear encoding model for each mouse to capture the linear contribution of behavioral

events to the overall observed ACh signal. The feature matrix includes representations of three

behavioral events: the onset of reward, the onset of punishment, and the licking. To make the

contributions of these events temporally flexible, we convolved them with a set of cubic splines.

We chose splines spanning from 1 s before to 2 s after the behavior to capture both preparatory

and reactive neural responses. Then we set the number of splines to seven based on three-fold

cross-validation accuracy optimized on a linear encoding model. With 7 splines associated with

each of the 3 behavior events, we obtained a total of 21 splines as regressors.

The encoding model takes the form $y = \beta X + \varepsilon$, where y is the photometry signal for a given mouse,

with dimension T, or the number of time points. X is the design matrix composed of behavior

events convolved with each cubic spline (plus one column of ones for learning the intercept term),

with dimensions $22 \times T$. β is the set of weights, one weight for each cubic spline associated with

each behavioral type plus an intercept term; the dimensions of β are 1×22. A percent explained

by a behavioral event was expressed as the reduction of the variance in the residual responses

compared to the original responses. Contributions of each behavioral event in the model were

measured by the reduction of the deviance compared to a reduced model excluding the behavioral

event.

QUANTIFICATION AND STATISTICAL ANALYSIS

All statistics are indicated where used. Statistical analyses were conducted using GraphPad Prism 7 Software (GraphPad Software, CA) and MATLAB statistical toolbox (MathWorks). The statistical test used for each comparison is indicated when used. Parametric tests were used whenever possible to test differences between two or more means. Non-parametric tests were used when data distributions were non-normal. Analysis of variance (ANOVA) was used to check for main effects and interactions in experiments with repeated measures and more than one factor. All comparisons were two-tailed. Statistic hypothesis testing was conducted at a significance level of 0.05.

REFERENCES

1. Fecteau, J.H., and Munoz, D.P. (2006). Salience, relevance, and firing: a priority map for target selection. Trends Cogn Sci *10*, 382–390. https://doi.org/10.1016/j.tics.2006.06.011.

2. Knudsen, E.I. (2007). Fundamental Components of Attention. Annu. Rev. Neurosci. *30*, 57–78. https://doi.org/10.1146/annurev.neuro.30.051606.094256.

3. Schimmelpfennig, J., Topczewski, J., Zajkowski, W., and Jankowiak-Siuda, K. (2023). The role of the salience network in cognitive and affective deficits. Front. Hum. Neurosci. *17*. https://doi.org/10.3389/fnhum.2023.1133367.

4. Uddin, L.Q. (2015). Salience processing and insular cortical function and dysfunction. Nat Rev Neurosci *16*, 55–61. https://doi.org/10.1038/nrn3857.

5. Vig, J. (2019). A Multiscale Visualization of Attention in the Transformer Model. arXiv. https://doi.org/10.48550/arXiv.1906.05714.

6. Vig, J., and Belinkov, Y. (2019). Analyzing the structure of attention in a transformer language model. arXiv.

7. Everitt, B.J., and Robbins, T.W. (1997). Central Cholinergic Systems and Cognition. Annual Review of Psychology *48*, 649–684. https://doi.org/10.1146/annurev.psych.48.1.649.

8. Hasselmo, M.E., and Sarter, M. (2011). Modes and Models of Forebrain Cholinergic
Neuromodulation of Cognition. Neuropsychopharmacol *36*, 52–73.
https://doi.org/10.1038/npp.2010.104.

9. Zhang, K., Chen, C.D., and Monosov, I.E. (2019). Novelty, Salience, and Surprise Timing Are
Signaled by Neurons in the Basal Forebrain. Current Biology *29*, 134-142.e3.
https://doi.org/10.1016/j.cub.2018.11.012.

10. Li, B., Ma, C., Huang, Y.-A., Ding, X., Silverman, D., Chen, C., Darmohray, D., Lu, L., Liu,
S., Montaldo, G., et al. (2023). Circuit mechanism for suppression of frontal cortical ignition
during NREM sleep. Cell *186*, 5739-5750.e17. https://doi.org/10.1016/j.cell.2023.11.012.

11. Hangya, B., Ranade, S.P., Lorenc, M., and Kepecs, A. (2015). Central Cholinergic Neurons
Are Rapidly Recruited by Reinforcement Feedback. Cell *162*, 1155–1168.
https://doi.org/10.1016/j.cell.2015.07.057.

12. Sturgill, J., Hegedus, P., Li, S., Chevy, Q., Siebels, A., Jing, M., Li, Y., Hangya, B., and Kepecs,
A. (2020). Basal forebrain-derived acetylcholine encodes valence-free reinforcement
prediction error (Neuroscience) https://doi.org/10.1101/2020.02.17.953141.

13. Hegedüs, P., Sviatkó, K., Király, B., Martínez-Bellver, S., and Hangya, B. (2023). Cholinergic
activity reflects reward expectations and predicts behavioral responses. iScience *26*.
https://doi.org/10.1016/j.isci.2022.105814.

14. Guo, W., Robert, B., and Polley, D.B. (2019). The Cholinergic Basal Forebrain Links Auditory
Stimuli with Delayed Reinforcement to Support Learning. Neuron *103*, 1164-1177.e6.
https://doi.org/10.1016/j.neuron.2019.06.024.

15. Kimchi, E.Y., Burgos-Robles, A., Matthews, G.A., Chakoma, T., Patarino, M., Weddington,
J.C., Siciliano, C., Yang, W., Foutch, S., Simons, R., et al. (2024). Reward contingency gates
selective cholinergic suppression of amygdala neurons. eLife *12*, RP89093.
https://doi.org/10.7554/eLife.89093.

16. Pinto, L., Goard, M.J., Estandian, D., Xu, M., Kwan, A.C., Lee, S.-H., Harrison, T.C., Feng,
G., and Dan, Y. (2013). Fast modulation of visual perception by basal forebrain cholinergic
neurons. Nat Neurosci *16*, 1857–1863. https://doi.org/10.1038/nn.3552.

17. Xu, M., Chung, S., Zhang, S., Zhong, P., Ma, C., Chang, W.-C., Weissbourd, B., Sakai, N.,
Luo, L., Nishino, S., et al. (2015). Basal forebrain circuit for sleep-wake control. Nat Neurosci
18, 1641–1647. https://doi.org/10.1038/nn.4143.

18. Liu, C.-H., Coleman, J.E., Davoudi, H., Zhang, K., and Hussain Shuler, M.G. (2015). Selective
Activation of a Putative Reinforcement Signal Conditions Cued Interval Timing in Primary
Visual Cortex. Current Biology *25*, 1551–1561. https://doi.org/10.1016/j.cub.2015.04.028.

19. Teles-Grilo Ruivo, L.M., Baker, K.L., Conway, M.W., Kinsley, P.J., Gilmour, G., Phillips,
K.G., Isaac, J.T.R., Lowry, J.P., and Mellor, J.R. (2017). Coordinated Acetylcholine Release

in Prefrontal Cortex and Hippocampus Is Associated with Arousal and Reward on Distinct
Timescales. Cell Reports *18*, 905–917. https://doi.org/10.1016/j.celrep.2016.12.085.

20. Whitehouse, P., Price, D., Struble, R., Clark, A., Coyle, J., and DeLong, M.R. (1982).
Alzheimer's Disease and Senile Dementia: Loss of Neurons in the Basal Forebrain. Science
215, 1237–1239. https://doi.org/10.1126/science.7058341.

21. Müller, M.L.T.M., and Bohnen, N.I. (2013). Cholinergic Dysfunction in Parkinson's Disease.
Curr Neurol Neurosci Rep *13*, 377. https://doi.org/10.1007/s11910-013-0377-9.

22. McGaughy, J., Dalley, J.W., Morrison, C.H., Everitt, B.J., and Robbins, T.W. (2002).
Selective behavioral and neurochemical effects of cholinergic lesions produced by intrabasalis
infusions of 192 IgG-saporin on attentional performance in a five-choice serial reaction time
task. J Neurosci *22*, 1905–1913. https://doi.org/10.1523/JNEUROSCI.22-05-01905.2002.

23. Luchicchi, A., Bloem, B., Viaña, J.N.M., Mansvelder, H.D., and Role, L.W. (2014).
Illuminating the role of cholinergic signaling in circuits of attention and emotionally salient
behaviors. Front. Synaptic Neurosci. *6*. https://doi.org/10.3389/fnsyn.2014.00024.

24. Rangel, L.M., Rueckemann, J.W., Riviere, P.D., Keefe, K.R., Porter, B.S., Heimbuch, I.S.,
Budlong, C.H., and Eichenbaum, H. (2016). Rhythmic coordination of hippocampal neurons
during associative memory processing. Elife *5*, e09849. https://doi.org/10.7554/eLife.09849.

25. Crouse, R.B., Kim, K., Batchelor, H.M., Girardi, E.M., Kamaletdinova, R., Chan, J.,
Rajebhosale, P., Pittenger, S.T., Role, L.W., Talmage, D.A., et al. (2020). Acetylcholine is
released in the basolateral amygdala in response to predictors of reward and enhances the
learning of cue-reward contingency. eLife *9*, e57335. https://doi.org/10.7554/eLife.57335.

26. Harrison, T.C., Pinto, L., Brock, J.R., and Dan, Y. (2016). Calcium Imaging of Basal Forebrain
Activity during Innate and Learned Behaviors. Front. Neural Circuits *10*.
https://doi.org/10.3389/fncir.2016.00036.

27. Paul, S., Jeon, W.K., Bizon, J.L., and Han, J.-S. (2015). Interaction of basal forebrain
cholinergic neurons with the glucocorticoid system in stress regulation and cognitive
impairment. Frontiers in Aging Neuroscience *7*.

28. Barabás, B., Reéb, Z., Papp, O.I., and Hájos, N. (2024). Functionally linked amygdala and
prefrontal cortical regions are innervated by both single and double projecting cholinergic
neurons. bioRxiv, 2023.12.29.573623. https://doi.org/10.1101/2023.12.29.573623.

29. Li, X., Yu, B., Sun, Q., Zhang, Y., Ren, M., Zhang, X., Li, A., Yuan, J., Madisen, L., Luo, Q.,
et al. (2018). Generation of a whole-brain atlas for the cholinergic system and mesoscopic
projectome analysis of basal forebrain cholinergic neurons. Proc Natl Acad Sci U S A *115*,
415–420. https://doi.org/10.1073/pnas.1703601115.

30. Sengupta, A., Yau, J.O.Y., Jean-Richard-Dit-Bressel, P., Liu, Y., Millan, E.Z., Power, J.M.,
and McNally, G.P. (2018). Basolateral Amygdala Neurons Maintain Aversive Emotional
Salience. J. Neurosci. *38*, 3001–3012. https://doi.org/10.1523/JNEUROSCI.2460-17.2017.

31. Cunningham, W.A., and Brosch, T. (2012). Motivational Salience: Amygdala Tuning From
Traits, Needs, Values, and Goals. Curr Dir Psychol Sci *21*, 54–59.
https://doi.org/10.1177/0963721411430832.

32. Zhang, X., Guan, W., Yang, T., Furlan, A., Xiao, X., Yu, K., An, X., Galbavy, W.,
Ramakrishnan, C., Deisseroth, K., et al. (2021). Genetically identified amygdala–striatal
circuits for valence-specific behaviors. Nat Neurosci *24*, 1586–1600.
https://doi.org/10.1038/s41593-021-00927-0.

33. Lin, S.-C., and Nicolelis, M.A.L. (2008). Neuronal ensemble bursting in the basal forebrain
encodes salience irrespective of valence. Neuron *59*, 138–149.
https://doi.org/10.1016/j.neuron.2008.04.031.

34. Kim, J., Pignatelli, M., Xu, S., Itohara, S., and Tonegawa, S. (2016). Antagonistic negative
and positive neurons of the basolateral amygdala. Nat Neurosci *19*, 1636–1646.
https://doi.org/10.1038/nn.4414.

35. Jiang, L., Kundu, S., Lederman, J.D., López-Hernández, G.Y., Ballinger, E.C., Wang, S.,
Talmage, D.A., and Role, L.W. (2016). Cholinergic Signaling Controls Conditioned Fear
Behaviors and Enhances Plasticity of Cortical-Amygdala Circuits. Neuron *90*, 1057–1070.
https://doi.org/10.1016/j.neuron.2016.04.028.

36. Jing, M., Li, Y., Zeng, J., Huang, P., Skirzewski, M., Kljakic, O., Peng, W., Qian, T., Tan, K.,
Zou, J., et al. (2020). An optimized acetylcholine sensor for monitoring in vivo cholinergic
activity. Nat Methods *17*, 1139–1146. https://doi.org/10.1038/s41592-020-0953-2.

37. Ghazizadeh, A., Griggs, W., and Hikosaka, O. (2016). Ecological Origins of Object Salience:
Reward, Uncertainty, Aversiveness, and Novelty. Frontiers in Neuroscience *10*.
https://doi.org/10.3389/fnins.2016.00378.

38. Zhu, Y., Nachtrab, G., Keyes, P.C., Allen, W.E., Luo, L., and Chen, X. (2018). Dynamic
salience processing in paraventricular thalamus gates associative learning. Science *362*, 423–
429. https://doi.org/10.1126/science.aat0481.

39. Lohani, S., Moberly, A.H., Benisty, H., Landa, B., Jing, M., Li, Y., Higley, M.J., and Cardin,
J.A. (2022). Spatiotemporally heterogeneous coordination of cholinergic and neocortical
activity. Nat Neurosci *25*, 1706–1713. https://doi.org/10.1038/s41593-022-01202-6.

40. Coddington, L.T., Lindo, S.E., and Dudman, J.T. (2023). Mesolimbic dopamine adapts the
rate of learning from action. Nature *614*, 294–302. https://doi.org/10.1038/s41586-022-05614-
z.

41. Do, J.P., Xu, M., Lee, S.-H., Chang, W.-C., Zhang, S., Chung, S., Yung, T.J., Fan, J.L.,
Miyamichi, K., Luo, L., et al. (2016). Cell type-specific long-range connections of basal
forebrain circuit. eLife *5*, e13214. https://doi.org/10.7554/eLife.13214.

42. Chen, Z., Yang, Y., Li, M., Gao, L., Qu, W., Huang, Z., and Yuan, X. (2023). Whole-brain
neural connectivity to cholinergic neurons in the nucleus basalis of Meynert. Journal of
Neurochemistry *166*, 233–247. https://doi.org/10.1111/jnc.15873.

43. Gielow, M.R., and Zaborszky, L. (2017). The Input-Output Relationship of the Cholinergic
Basal Forebrain. Cell Reports *18*, 1817–1830. https://doi.org/10.1016/j.celrep.2017.01.060.

44. Henderson, Z. (1997). The projection from the striatum to the nucleus basalis in the rat: an
electron microscopic study. Neuroscience *78*, 943–955. https://doi.org/10.1016/S0306-
4522(96)00636-7.

45. Yang, C., Thankachan, S., McCarley, R.W., and Brown, R.E. (2017). The menagerie of the
basal forebrain: how many (neural) species are there, what do they look like, how do they
behave and who talks to whom? Current Opinion in Neurobiology *44*, 159–166.
https://doi.org/10.1016/j.conb.2017.05.004.

46. Sabol, K.E., Richards, J.B., Broom, S.L., Roach, J.T., and Hausknecht, K. (2003). Effects of
stimulus salience and methamphetamine on choice reaction time in the rat: central tendency
versus distribution skew. Behav Pharmacol *14*, 489–500. https://doi.org/10.1097/00008877-
200311000-00001.

47. Bell, A.H., Meredith, M.A., Van Opstal, A.J., and Munoz, D.P. (2006). Stimulus intensity
modifies saccadic reaction time and visual response latency in the superior colliculus. Exp
Brain Res *174*, 53–59. https://doi.org/10.1007/s00221-006-0420-z.

48. Gottlieb, J.P., Kusunoki, M., and Goldberg, M.E. (1998). The representation of visual salience
in monkey parietal cortex. Nature *391*, 481–484. https://doi.org/10.1038/35135.

49. Ballinger, E.C., Ananth, M., Talmage, D.A., and Role, L.W. (2016). Basal Forebrain
Cholinergic Circuits and Signaling in Cognition and Cognitive Decline. Neuron *91*, 1199–
1218. https://doi.org/10.1016/j.neuron.2016.09.006.

50. Floresco, S.B. (2015). The Nucleus Accumbens: An Interface Between Cognition, Emotion,
and Action. Annual Review of Psychology *66*, 25–52. https://doi.org/10.1146/annurev-psych-
010213-115159.

51. Castro, D.C., and Bruchas, M.R. (2019). A Motivational and Neuropeptidergic Hub:
Anatomical and Functional Diversity within the Nucleus Accumbens Shell. Neuron *102*, 529–
552. https://doi.org/10.1016/j.neuron.2019.03.003.

52. Li, Z., Chen, Z., Fan, G., Li, A., Yuan, J., and Xu, T. (2018). Cell-Type-Specific Afferent
Innervation of the Nucleus Accumbens Core and Shell. Front. Neuroanat. *12*.
https://doi.org/10.3389/fnana.2018.00084.

53. Liu, Z., Le, Q., Lv, Y., Chen, X., Cui, J., Zhou, Y., Cheng, D., Ma, C., Su, X., Xiao, L., et al.
(2022). A distinct D1-MSN subpopulation down-regulates dopamine to promote negative
emotional state. Cell Res *32*, 139–156. https://doi.org/10.1038/s41422-021-00588-5.

54. Klawonn, A.M., and Malenka, R.C. (2018). Nucleus Accumbens Modulation in Reward and
Aversion.   Cold   Spring   Harb   Symp   Quant   Biol   *83*,   119–129.
https://doi.org/10.1101/sqb.2018.83.037457.

55. Lobo, M.K., Covington, H.E., Chaudhury, D., Friedman, A.K., Sun, H., Damez-Werno, D.,
Dietz, D.M., Zaman, S., Koo, J.W., Kennedy, P.J., et al. (2010). Cell Type–Specific Loss of
BDNF Signaling Mimics Optogenetic Control of Cocaine Reward. Science *330*, 385–390.
https://doi.org/10.1126/science.1188472.

56. Yang, H., de Jong, J.W., Tak, Y., Peck, J., Bateup, H.S., and Lammel, S. (2018). Nucleus
Accumbens Subnuclei Regulate Motivated Behavior via Direct Inhibition and Disinhibition of
VTA   Dopamine   Subpopulations.   Neuron   *97*,   434-449.e4.
https://doi.org/10.1016/j.neuron.2017.12.022.

57. Soares-Cunha, C., de Vasconcelos, N.A.P., Coimbra, B., Domingues, A.V., Silva, J.M.,
Loureiro-Campos, E., Gaspar, R., Sotiropoulos, I., Sousa, N., and Rodrigues, A.J. (2020).
Nucleus accumbens medium spiny neurons subtypes signal both reward and aversion. Mol
Psychiatry *25*, 3241–3255. https://doi.org/10.1038/s41380-019-0484-3.

58. O'Connor, E.C., Kremer, Y., Lefort, S., Harada, M., Pascoli, V., Rohner, C., and Lüscher, C.
(2015). Accumbal D1R Neurons Projecting to Lateral Hypothalamus Authorize Feeding.
Neuron *88*, 553–564. https://doi.org/10.1016/j.neuron.2015.09.038.

59. Thoeni, S., Loureiro, M., O'Connor, E.C., and Lüscher, C. (2020). Depression of Accumbal
to Lateral Hypothalamic Synapses Gates Overeating. Neuron *107*, 158-172.e4.
https://doi.org/10.1016/j.neuron.2020.03.029.

60. Bond, C.W., Trinko, R., Foscue, E., Furman, K., Groman, S.M., Taylor, J.R., and DiLeone,
R.J. (2020). Medial Nucleus Accumbens Projections to the Ventral Tegmental Area Control
Food Consumption. J Neurosci *40*, 4727–4738. https://doi.org/10.1523/JNEUROSCI.3054-
18.2020.

61. Liu, Y., Wang, Y., Zhao, Z., Xie, G., Zhang, C., Chen, R., and Zhang, Y. (2024). A molecularly
defined NAcSh D1 subtype controls feeding and energy homeostasis. bioRxiv,
2023.02.27.530275. https://doi.org/10.1101/2023.02.27.530275.

62. Rescorla, R.A., and Holland, P.C. (1982). Behavioral Studies of Associative Learning in
Animals.   Annual   Review   of   Psychology   *33*,   265–308.
https://doi.org/10.1146/annurev.ps.33.020182.001405.

63. Gu, Z., Lamb, P.W., and Yakel, J.L. (2012). Cholinergic coordination of presynaptic and
postsynaptic activity induces timing-dependent hippocampal synaptic plasticity. J Neurosci *32*,
12337–12348. https://doi.org/10.1523/JNEUROSCI.2129-12.2012.

64. Gu, Z., and Yakel, J.L. (2011). Timing-dependent septal cholinergic induction of dynamic
hippocampal   synaptic   plasticity.   Neuron   *71*,   155–165.
https://doi.org/10.1016/j.neuron.2011.04.026.

65. Palacios-Filardo, J., and Mellor, J.R. (2019). Neuromodulation of hippocampal long-term
synaptic   plasticity.   Current   Opinion   in   Neurobiology   *54*,   37–43.
https://doi.org/10.1016/j.conb.2018.08.009.

66. Dennis, S.H., Pasqui, F., Colvin, E.M., Sanger, H., Mogg, A.J., Felder, C.C., Broad, L.M.,
Fitzjohn, S.M., Isaac, J.T.R., and Mellor, J.R. (2016). Activation of Muscarinic M1
Acetylcholine Receptors Induces Long-Term Potentiation in the Hippocampus. Cerebral
Cortex *26*, 414–426. https://doi.org/10.1093/cercor/bhv227.

67. Liu, C., Cai, X., Ritzau-Jost, A., Kramer, P.F., Li, Y., Khaliq, Z.M., Hallermann, S., and
Kaeser, P.S. (2022). An action potential initiation mechanism in distal axons for the control of
dopamine release. Science *375*, 1378–1385. https://doi.org/10.1126/science.abn0532.

68. Krok, A.C., Maltese, M., Mistry, P., Miao, X., Li, Y., and Tritsch, N.X. (2023). Intrinsic
dopamine and acetylcholine dynamics in the striatum of mice. Nature *621*, 543–549.
https://doi.org/10.1038/s41586-023-05995-9.

69. Chantranupong, L., Beron, C.C., Zimmer, J.A., Wen, M.J., Wang, W., and Sabatini, B.L.
(2023). Dopamine and glutamate regulate striatal acetylcholine in decision-making. Nature
621, 577–585. https://doi.org/10.1038/s41586-023-06492-9.

70. Threlfell, S., Lalic, T., Platt, N.J., Jennings, K.A., Deisseroth, K., and Cragg, S.J. (2012).
Striatal Dopamine Release Is Triggered by Synchronized Activity in Cholinergic Interneurons.
Neuron *75*, 58–64. https://doi.org/10.1016/j.neuron.2012.04.038.

71. Cachope, R., Mateo, Y., Mathur, B.N., Irving, J., Wang, H.-L., Morales, M., Lovinger, D.M.,
and Cheer, J.F. (2012). Selective Activation of Cholinergic Interneurons Enhances Accumbal
Phasic Dopamine Release: Setting the Tone for Reward Processing. Cell Reports *2*, 33–41.
https://doi.org/10.1016/j.celrep.2012.05.011.

72. Lutas, A., Kucukdereli, H., Alturkistani, O., Carty, C., Sugden, A.U., Fernando, K., Diaz, V.,
Flores-Maldonado, V., and Andermann, M.L. (2019). State-specific gating of salient cues by
midbrain dopaminergic input to basal amygdala. Nat Neurosci *22*, 1820–1833.
https://doi.org/10.1038/s41593-019-0506-0.

73. Xiao, X., Deng, H., Furlan, A., Yang, T., Zhang, X., Hwang, G.-R., Tucciarone, J., Wu, P.,
He, M., Palaniswamy, R., et al. (2020). A Genetically Defined Compartmentalized Striatal
Direct   Pathway   for   Negative   Reinforcement.   Cell   *183*,   211-227.e20.
https://doi.org/10.1016/j.cell.2020.08.032.

74. Deng, H., Xiao, X., Yang, T., Ritola, K., Hantman, A., Li, Y., Huang, Z.J., and Li, B. (2021).
A genetically defined insula-brainstem circuit selectively controls motivational vigor. Cell *184*,
6344-6360.e18. https://doi.org/10.1016/j.cell.2021.11.019.

 FIGURES, SUPPLEMENTARY FIGURES, and LEGEND

3
Figure 1. BLA ACh encodes behavioral salience irrespective of valence
(A) Left: schematic of the approach to record BLA ACh. Right: representative coronal image showing the
expression of a fluorescent acetylcholine sensor (ACh3.8) in the BLA and the placement of optical fiber
for photometry.
(B) Schematics of the experimental setup.
(C) Trial-by-trial heatmap of BLA ACh responses to the white noise in an example session. Each session
consists of 50 trials, divided into 5 sections (10 trials per section).
(D) Trial-by-trial heatmap of BLA ACh responses to the water in an example session.
(E) Average BLA ACh responses (n = 12 mice) to white noise along sections.
(F) Average BLA ACh responses (n = 12 mice) to water along sections.
(G) Left: average BLA ACh responses (n = 12 mice) to different sizes of reward. Right: quantification of
peak BLA ACh responses ($F_{(2, 33)}$ = 18.93, p < 0.001, one-way ANOVA followed by Tukey's test).
(H) Left: average BLA ACh responses (n = 12 mice) to different sizes of punishment. Right: quantification
of peak BLA ACh responses ($F_{(2, 33)}$ = 14.4, p < 0.001, one-way ANOVA followed by Tukey's test).
(I) Schematics of the experimental setup (top) and the Pavlovian conditioning task design (bottom).

(J) Trial-by-trial (top) and trial-averaged (bottom) BLA ACh responses in reward trials at the early (left)
and late (right) stages of learning from a representative mouse in the Pavlovian task.
(K) Trial-by-trial (top) and trial-averaged (bottom) BLA ACh responses in punishment trials at the early
(left) and late (right) learning stages from a representative mouse in the Pavlovian task.
(L) Quantification of the mean (left) ($F_{(1, 16)} = 40.36$, $p < 0.001$) and peak (right) ($F_{(1, 16)} = 38.24$, $p < 0.001$)
BLA ACh responses to conditioned stimuli (CS) in the Pavlovian task. Two-way ANOVA followed by
Bonferroni's test.
(M) Quantification of the mean (left) ($F_{(1, 16)} = 3.07$, $p = 0.099$) and peak (right) ($F_{(1, 16)} = 7.42$, $p = 0.015$)
BLA ACh responses to unconditioned stimuli (US) in the Pavlovian task. Two-way ANOVA followed by
Bonferroni's test.
(N) Receiver operating characteristic (ROC) curve evaluating the discriminability of BLA ACh responses
to reward CS (left, auROC = 0.774) and punishment CS (right, auROC = 0.767) between early and late
trials from a representative mouse.
(O) Quantification of area under the ROC curve (auROC) for CS responses in reward and punishment
trials (auROC vs 0.5, $p = 0.002$ for reward, $p < 0.001$ for punishment, paired t-test; reward vs punishment,
$p = 0.66$, t-test).
35
36 **$p < 0.01$, ***$p < 0.001$; n.s., non-significant ($p > 0.05$). Data are presented as mean $\pm$ SEM. Shaded
37 areas represent SEM.
38

39
Figure 2. Modulation of salience response in BLA ACh by behavioral context and homeostatic state
(A) Schematic of the go/no-go task design. CR, correct rejection; FA, false alarm. CS_G, CS for "go"; CS_N,
CS for "no-go".
(B, C) Trial-by-trial (top) and average (bottom) BLA ACh responses of a representative mouse at the early
(B) and late (C) stages of the go/no-go task.
(D, E) Average BLA ACh responses (left) and quantification of peak CS responses (right) for different
trial outcomes at the early (D) (hit, p = 0.0039; CR, p = 0.22; FA, p = 0.071; paired t-test) and late (E) (hit,
p < 0.001; CR, p = 0.75; FA, p = 0.87; paired t-test) stages of the go/no-go task.
(F) Quantification of peak CS responses of BLA ACh in different trial outcomes at the early and late
training stages of the go/no-go task (n = 5 mice; hit, p = 0.022; CR, p = 0.34; FA, p = 0.26; paired t-test).
(G) Schematic of the active avoidance task design. CS_A, CS for "avoid".
(H, I) Trial-by-trial (top) and average (bottom) BLA ACh responses of a representative mouse at the early
(H) and late (I) stages of the active avoidance task.

(J, K) Average BLA ACh responses (left) and quantification of peak CS responses (right) for different trial outcomes at the early (J) (success, $p = 0.46$; failure, $p = 0.99$; paired t-test) and late (K) (success, $p = 0.004$; failure, $p = 0.066$; paired t-test) stages of the active avoidance task.

(L) Quantification of peak CS responses for different trial outcomes at the early and late training stages of the active avoidance task ($n = 5$ mice; success, $p = 0.033$; failure, $p = 0.046$; paired t-test).

(M) ROC curve evaluating the discriminability of BLA ACh responses to CS_A and baseline at the late stage of the active avoidance task from a representative mouse.

(N) Quantification of auROC for the early and late training stages of the active avoidance task (auROC vs 0.5, $p = 0.41$ for early, $p = 0.015$ for late; early vs late, $p = 0.044$; paired t-test).

(O) Quantification of peak BLA ACh responses for CS_A at the early and late stages of the active avoidance task ($p = 0.033$, t-test).

(P) Quantification of peak BLA ACh responses for CS_N and CS_A at the late stage of the go/no-go and the active avoidance task, respectively ($p = 0.0058$, t-test).

(Q) Average lick rate (left) and BLA ACh responses (right) of a representative mouse during reward trials in the thirsty and quenched states in the Pavlovian task.

(R) Quantification of the peak responses of BLA ACh for reward CS (left; $p = 0.0037$; paired t-test) and US (right; $p = 0.0168$; paired t-test) in the thirsty and quenched states ($n = 5$ mice).

(S) Average BLA ACh responses of a representative mouse during punishment trials in the thirsty and quenched states in the Pavlovian task.

(T) Quantification of the peak responses of BLA ACh for punishment CS (left; $p = 0.7485$; paired t-test) and US (right; $p = 0.0136$; paired t-test) in the thirsty and quenched states ($n = 5$ mice).

*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; n.s., non-significant ($p > 0.05$). Data are presented as mean ± SEM. Shaded areas represent SEM.

78
Figure 3. NAc D1 and D2 MSNs opposingly regulate the release of ACh in the BLA
(A) Schematic of the strategy to record BLA ACh while activating NAc D1 MSNs.
(B) Left: representative image of ChR2 expression in NAc D1 MSNs. Right: representative image of
ACh3.8 expression in the BLA.
(C) Trial-by-trial (top) and average (bottom) responses of BLA ACh while activating NAc D1 MSNs with
different laser durations in an example session.
(D) Average BLA ACh responses (n = 8 mice) while activating NAc D1 MSNs with different laser
durations.

(E-G) Quantification of the mean (E) ($F_{(3, 28)}$ = 7.28, p < 0.001), peak (F) ($F_{(3, 28)}$ = 11.02, p < 0.001), and area under curve (AUC) (G) ($F_{(3, 28)}$ = 7.32, p < 0.001) of BLA ACh responses while activating NAc D1 MSNs with different laser durations (n = 8 mice). One-way ANOVA.

(H) Schematic of the strategy to record BLA ACh while activating NAc D2 MSNs.

(I) Left: representative image of ChR2 expression in NAc D2 MSNs. Right: representative image of ACh3.8 expression in the BLA.

(J) Trial-by-trial (top) and average (bottom) responses of BLA ACh while activating NAc D2 MSNs with different laser durations in an example session.

(K) Average BLA ACh responses (n = 7 mice) while activating NAc D2 MSNs with different laser durations.

(L-N) Quantification of the mean (L) ($F_{(3, 24)}$ = 22.52, p < 0.001), trough (M) ($F_{(3, 24)}$ = 36.08, p < 0.001), and AUC (N) ($F_{(3, 24)}$ = 22.1, p < 0.001) of BLA ACh responses while activating NAc D2 MSNs with different laser durations (n = 7 mice). One-way ANOVA.

***p < 0.001. Data are presented as mean ± SEM. Shaded areas represent SEM.

Figure 4. Activation of NAc^{D1}→SI and NAc^{D2}→SI pathways opposingly regulates the release of ACh in the BLA

(A) Schematic of the strategy to record BLA ACh while activating NAc^{D1}→SI.

(B) Left: representative image of ChR2-expressing NAc^{D1} projections to the SI. Right: representative image of ACh3.8 expression in the BLA.

(C) Trial-by-trial (top) and average (bottom) responses of BLA ACh while activating NAc^{D1}→SI pathway with different laser durations in an example session.

(D, E) Quantification of the peak (D) ($F_{(3, 28)} = 31.54$, $p < 0.001$) and AUC (E) ($F_{(3, 28)} = 22.5$, $p < 0.001$) of BLA ACh responses while activating NAc^{D1}→SI pathway with different laser durations (n = 8 mice). One-way ANOVA.

(F) Schematic of the strategy to record BLA ACh while activating NAc^{D2}→SI.

(G) Left: representative image of ChR2-expressing NAc^{D2} projections to the SI. Right: representative image of ACh sensor expression in the BLA.

(H) Trial-by-trial (top) and average (bottom) responses of BLA ACh while activating NAc^{D2}→SI pathway with different laser durations in an example session.

(I, J) Quantification of the trough (I) ($F_{(3, 24)} = 19.05$, $p < 0.001$) and AUC (J) ($F_{(3, 24)} = 22.44$, $p < 0.001$) of BLA ACh responses while activating NAc^{D2}→SI pathway with different laser durations (n = 7 mice). One-way ANOVA.

***$p < 0.001$. Data are presented as mean ± SEM. Shaded areas represent SEM.

125
Figure 5. Effects of NAc→SI activation on the ACh release in the cortex and hippocampus
(A) Schematic of anterograde tracing of SI ChAT neurons.
(B, C) Example coronal sections showing EGFP-expressing cell bodies in the SI (B) and their labeled
axonal downstream in the BLA, mPFC, AUD, and CA1 (C).
(D) Schematic of the strategy to record mPFC ACh while activating NAc^{D1}→SI or NAc^{D2}→SI.
(E, F) Average mPFC ACh responses across mice while activating NAc^{D1}→SI (E) and NAc^{D2}→SI (F)
with different laser durations.
(G) Schematic of the strategy to record AUD ACh while activating the NAc^{D1}→SI or NAc^{D2}→SI pathway.
(H, I) Average AUD ACh responses across mice while activating the NAc^{D1}→SI (H) and NAc^{D2}→SI (I)
with different laser durations.
(J) Schematic of the strategy to record CA1 ACh while activating NAc^{D1}→SI or NAc^{D2}→SI.
(K, L) Average CA1 ACh responses across mice while activating NAc^{D1}→SI (K) and NAc^{D2}→SI (L)
with different laser durations.
(M) Quantification of the ACh release in the mPFC, AUD, and CA1 while activating NAc^{D1}→SI in
comparison to ACh release in the BLA (left, $F_{(3, 23)} = 16.58$, $p < 0.001$; middle, $F_{(3, 23)} = 22.96$, $p < 0.001$;
right, $F_{(3, 23)} = 47.73$, $p < 0.001$; one-way ANOVA followed by Tukey's test).

(N) Quantification of peak latency of ACh signal in the BLA, mPFC, AUD, and CA1 while activating
$NAc^{D1} \rightarrow SI$ (left, $F_{(3, 23)} = 5.36$, $p = 0.006$; middle, $F_{(3, 23)} = 6.08$, $p = 0.003$; right, $F_{(3, 23)} = 10.86$, $p < 0.001$;
one-way ANOVA followed by Tukey's test).
(O) Quantification of the ACh release in the BLA, mPFC, AUD, and CA1 while activating $NAc^{D2} \rightarrow SI$ in
comparison to ACh release in the BLA (left, $F_{(3, 26)} = 3.36$, $p = 0.034$; middle, $F_{(3, 26)} = 6.89$, $p = 0.0014$;
right, $F_{(3, 26)} = 50.67$, $p < 0.001$; one-way ANOVA followed by Tukey's test).
148
149 $*p < 0.05$, $**p < 0.01$, $***p < 0.001$; n.s., non-significant ($p > 0.05$). Data are presented as mean ± SEM.
150 Shaded areas represent SEM.
151
152

153
Figure 6. Synaptic and functional connections between NAc MSNs and SI ChAT neurons
154
(A, D) Schematic of the approach.
(B, E) Confocal images showing synaptic puncta (red) of NAc D1 (B) and D2 (E) MSNs and cell body of
ChAT neurons (green).
(C, F) High-magnification images showing 3D reconstructed ChAT neurons and surrounding
synaptophysin spots from a representative *D1-Cre* (C) and *D2-Cre* (F) mouse.
(G) Schematic of synaptophysin spot quantification around ChAT neurons.
(H) Quantification of synaptophysin spots surrounding ChAT neurons for D1 and D2 MSNs (p = 0.032,
t-test).
(I) Schematic of the approach.
(J) Confocal images from a mouse prepared as in (I), showing the BLA-projecting ChAT neurons in the
SI infected by the helper viruses (green) and the cells infected by the rabies virus (RV) (red). The starter
cells are yellow, as indicated by the arrowhead.
(K) Representative image of the NAc, showing input neurons labeled by the RV.
(L) Confocal images of in situ hybridization for RV-tdTomato, Drd1, and Drd2 in the NAc. Right, high-
magnification views of the boxed area on the left.
(M) Quantification of the percentage of D1 or D2 NAc MSNs among RV-tdTomato cells (n = 6; p =
0.0029, paired t-test).
(N) Schematic of the approach.

(O) Average SI ChAT neural activities while activating the $NAc^{D1} \rightarrow SI$ pathway with different laser
durations (n = 7 mice).
(P) Quantification of the mean (left; $F_{(2, 18)} = 4.35$, $p = 0.03$) and AUC (right; $F_{(2, 18)} = 4.29$, $p = 0.03$) of
SI ChAT neural activities (n = 7 mice). One-way ANOVA.
177
178 $*p < 0.05$, $**p < 0.01$. Data are presented as mean ± SEM.
179

180
Figure 7. Activation of NAc^D1→SI and NAc^D2→SI facilitates and impairs reward associative learning, respectively

(A) Schematics of the cue-reward learning task. In the pre-training phase, there was no consequence for incorrect nose pokes (left). In training, timeout punishment was triggered by incorrect nose pokes (right).
(B) Schematic of the experimental procedure.
(C) Schematic of the strategy to activate NAc^D1→SI pathway.
(D, E) Rewards earned by the ChR2 (n = 6) and GFP (n = 8) mice in each session (D) (interaction, $F_{(11, 132)}$ = 4.18, p < 0.0001; group, $F_{(1, 12)}$ = 5.75, p = 0.034; two-way ANOVA), and average across sessions (E) (p = 0.034, t-test) with closed-loop stimulation of NAc^D1→SI in the cue-reward learning task.
(F, G) Behavioral accuracy of the ChR2 (n = 6) and GFP (n = 8) mice in each session (F) (interaction, $F_{(11, 132)}$ = 2.01, p = 0.032; group, $F_{(1, 12)}$ = 18.32, p = 0.0011; two-way ANOVA), and average across sessions (G) (p = 0. 0011, t-test) with closed-loop stimulation of NAc^D1→SI in the cue-reward learning task.
(H) Schematics of the strategy to activate NAc^D2→SI pathway.
(I, J) Rewards earned by the ChR2 (n = 7) and GFP (n = 4) mice in each session (I) (interaction, $F_{(11, 99)}$ = 6.25, P < 0.001; group, $F_{(1, 9)}$ = 19.52, p = 0.0017; two-way ANOVA), and average across sessions (J) (p = 0.0017, t-test) with closed-loop stimulation of NAc^D2→SI in the cue-reward learning task.
(K, L) Behavioral accuracy of the ChR2 (n = 7) and GFP (n = 4) mice in each session (K) (interaction, $F_{(11, 99)}$ = 3.16, p = 0.0011; group, $F_{(1, 9)}$ = 18.21, p = 0.0021; two-way ANOVA), and average across sessions (L) (p = 0.0021, t-test) with closed-loop stimulation of NAc^D2→SI in the cue-reward learning task.

*p < 0.05, **p < 0.01. Data are presented as mean ± SEM.

Figure S1. BLA ACh encodes salience but not movement

(A) Section-by-section (left) and average BLA ACh responses across mice in response to white noise over sections (n = 13 mice).

(B) Quantification of BLA ACh responses following white noise delivery (0-1 s) in the first and last section (n = 13 mice; p < 0.001, paired t-test).

(C) Section-by-section (left) and average BLA ACh responses across mice in response to water over sections (n = 12 mice).

(D) Quantification of BLA ACh responses following water delivery (0-1 s) in the first and last section (n = 12 mice; p = 0.15, paired t-test).

(E) Schematic of the experimental design.

(F) Average BLA ACh responses (left) and licking rate (right) to different sizes of reward in one session.

(G) Quantification of licking rate during reward (0-1 s) from the same session as in (F) ($F_{(2, 72)}$ = 37.97, p < 0.001, one-way ANOVA followed by Tukey's test).

(H) Average BLA ACh responses (left) and licking rate (right) to different sizes of punishment in one session.

(I) Quantification of licking rate during punishment (0-1 s) from the same session as in (h) ($F_{(2, 72)}$ = 8.48, p < 0.001, one-way ANOVA followed by Tukey's test).

(J) Representative images of the eye before (left) and after (right) air puff delivery, tracked with DeepLabCut.

(K) Average (left) and quantification (right) of the eye size change in response to different sizes of punishment ($F_{(2, 72)}$ = 9.5, p < 0.001, one-way ANOVA followed by Tukey's test).

(L) Left: correlation between BLA ACh responses and licking rate in an example session (Pearson's r = -
0.06, p = 0.43). Right: distribution of Pearson's correlation coefficient (n = 16 sessions).
(M) Schematic of the generalized linear model (GLM) for the relationship between behavioral events and
BLA ACh responses.
(N) Example of observed (left) and model-predicted (right) BLA ACh responses in reward (top) and
punishment (bottom) trials.
(O) Relative contribution of single model variables (lick, reward, and punishment) to model-predicted
BLA ACh responses (n = 16 sessions; $F_{(2, 45)}$ = 26.21, p < 0.001, one-way ANOVA followed by Tukey's
test).
236
237   *p < 0.05, **p < 0.01, ***p < 0.001; n.s., non-significant (p > 0.05). Data are presented as mean ± SEM.
238   Shaded areas represent SEM.
239

Figure S2. Dynamics of BLA ACh activity during learning

(A) Schematic of the Pavlovian conditioning task design.

(B) Example traces of simultaneously measured behavioral and BLA ACh responses in a representative mouse at the early (top) and late (bottom) learning stages.

(C) Top: licking events for a representative mouse at the early (left) and late (right) learning stages in the Pavlovian task. Middle: average licking rate. Bottom: average BLA ACh responses from this mouse. Dashed lines indicate the onset of CS and US.

(D, E) Correlation between licking rate and BLA ACh responses during baseline in an example session (left) and the distribution of Pearson's correlation coefficient (right) at the early (D) (Pearson's r = -0.14, p = 0.24) and the late (E) (Pearson's r = -0.01, p = 0.90) learning stages (n = 6 mice).

(F) Schematics for the probabilistic Pavlovian conditioning task with cue and outcome probabilities indicated.

(G) Trial-by-trial (top) and average (bottom) responses of BLA ACh in different types of reward trials from an example session.

(H) Trial-by-trial (top) and average (bottom) responses of BLA ACh in different types of punishment trials from an example session.

(I) Average BLA ACh responses (n = 6 mice) in different types of reward trials.

(J) Quantification of the peak BLA ACh responses in different types of reward trials ($F_{(2, 15)}$ = 5.19, p = 0.019, one-way ANOVA followed by Tukey's test).

(K) Average BLA ACh responses (n = 8 mice) in different types of punishment trials.

261 **(L)** Quantification of the peak BLA ACh responses in different types of punishment trials ($F_{(2, 21)} = 9.73$,
262 $p = 0.001$, one-way ANOVA followed by Tukey's test).
263
264 $^*p < 0.05$, $^{**}p < 0.01$; n.s., non-significant ($p > 0.05$). Data are presented as mean ± SEM. Shaded areas
265 represent SEM.
266

Figure S3. Laser stimulation of the NAc does not affect BLA ACh in GFP mice
(A) Schematic of the approach to activate the NAc and record BLA ACh simultaneously.
(B) Diagram showing simultaneous opto-stimulation and photometry recording interleaved at the frequency of 20 Hz.

(C-E) Example trace of BLA ACh responses with laser stimulation from a representative *D1-Cre* mouse
(C), *D2-Cre* mouse (D), and wild-type (WT) mouse (E).
(F) Schematic of the strategy to record BLA ACh while delivering laser into the NAc of a WT mouse.
(G) Left: representative image of GFP expression in the NAc. Right: representative image of ACh3.8
expression in the BLA.
(H) Trial-by-trial heatmap (top) and average (bottom) responses of BLA ACh while delivering laser into
the NAc with different durations in an example session of a WT mouse.
(I) Average BLA ACh responses (n = 6) while delivering the laser into the NAc with different laser
durations in WT mice.
(J, K) Quantification of the mean (J) ($F_{(3, 20)}$ = 0.36, p = 0.78) and AUC (K) ($F_{(3, 20)}$ = 0.38, p = 0.77) of
BLA ACh signals while delivering laser into the NAc with different laser durations in WT mice (n = 6).
One-way ANOVA.
(L) Schematic of anterograde tracing of NAc D1 or D2 MSNs.
(M, N) Example confocal images of the injection site (soma; green) in the NAc (left) and the axonal
terminals (red) in the SI (right) from a representative *D1-Cre* (M) or *D2-Cre* (N) mouse.
287
288 n.s., non-significant (p > 0.05). Data are presented as mean ± SEM. Shaded areas represent SEM.
289

Figure S4. Activation of SI ChAT neurons induces potent ACh release in the BLA

(A) Schematic of retrograde labeling of ChAT neurons that project to the BLA.

(B) Representative images showing labeled ChAT neurons in the basal forebrain.

(C) Quantification of labeled ChAT neurons in the medial septum (MS), diagonal band of Broca (HDB), and SI (n = 4; $F_{(2, 9)}$ = 25408, p < 0.001). One-way ANOVA followed by Tukey's test.

(D) Schematic of anterograde tracing of BLA-projecting SI ChAT neurons.

(E, F) Representative images showing EGFP-expressing cell bodies in SI (E) and their labeled axonal downstream (F).

(G) Schematic of the strategy to record BLA ACh while activating SI ChAT neurons.

(H) Left: representative image of ChR2 expression in SI ChAT neurons. Right: representative image of Ach3.8 expression in the BLA.

(I) Trial-by-trial (top) and average (bottom) responses of BLA ACh while activating SI ChAT neurons
with different laser durations in an example session.
(J) Example trace of BLA ACh responses when activating SI ChAT neurons with different laser durations.
(K) Average BLA ACh responses (n = 4 mice) while delivering the laser into the SI with different laser
durations.
(L-N) Quantification of the mean (L) ($F_{(3, 12)}$ = 12.22, $p < 0.001$), peak (M) ($F_{(3, 12)}$ = 13.59, $p < 0.001$),
and AUC (N) ($F_{(3, 12)}$ = 12.22, $p < 0.001$) of BLA ACh signals while activating SI ChAT neurons with
different laser durations (n = 4 mice). One-way ANOVA.
310
311   ***$p < 0.001$. Data are presented as mean ± SEM. Shaded areas represent SEM.
312

314
Figure S5. Activation of SI$^{\text{ChAT}}$ neurons and SI$^{\text{ChAT}}$→BLA pathway does not induce valence or drive
reinforcement
(A) Schematic of the approach for optogenetic activation of SI$^{\text{ChAT}}$ neurons.
(B) Heatmaps for the activity of a representative mouse at baseline (top), or in a situation where entering
the left (middle) or right (bottom) side of the chamber triggered photo-stimulation in the SI.
(C-E) Quantification of the mouse activity as shown in (B), showing that activation of SI$^{\text{ChAT}}$ neurons
does not induce place preference or aversion (C) ($F_{(2, 12)} = 0.54$, p = 0.59, one-way ANOVA followed by
Tukey's test), affect movement velocity (D) ($F_{(2, 12)} = 0.74$, p = 0.50, two-way ANOVA followed by
Bonferroni's test), or affect distance traveled (E) ($F_{(2, 12)} = 0.44$, p = 0.66, two-way ANOVA followed by
Bonferroni's test).
(F) Schematic of the approach for optogenetic activation of SI$^{\text{ChAT}}$→BLA pathway.
(G) Heatmaps for the activity of a representative mouse at baseline (top), or in a situation where entering
the left (middle) or right (bottom) side of the chamber triggered photo-stimulation of SI$^{\text{ChAT}}$→BLA
pathway.
(H-J) Quantification of the mouse activity as shown in (G), showing that activation of SI$^{\text{ChAT}}$ →BLA
pathway does not induce place preference or aversion (H) ($F_{(2, 9)} = 1.73$, p = 0.23, one-way ANOVA
followed by Tukey's test), affect movement velocity (I) ($F_{(2, 9)} = 1.30$, p = 0.32, two-way ANOVA
followed by Bonferroni's test), or affect distance traveled (J) ($F_{(2, 9)} = 1.77$, p = 0.23, two-way ANOVA
followed by Bonferroni's test).
(K) Schematic of the self-stimulation test.
(L-N) Optogenetic activation of SI$^{\text{ChAT}}$→BLA does not support self-stimulation.

(L) Poking events at one port where poking triggered the photoactivation of $SI^{ChAT} \rightarrow BLA$ (active), and
the other port where poking did not trigger the photo-stimulation (inactive) from an example session of a
ChR2 mouse.
(M) Cumulative curves for the poking responses at a port where poking triggered the photo-stimulation
(active), and a port where poking did not trigger the photo-stimulation (inactive), in ChR2 mice (n = 4),
or WT mice (as the control; n = 5).
(N) Quantification of the poking responses as shown in (M) ($F_{(1,\ 7)}$ = 0.22, p = 0.66, two-way ANOVA
followed by Bonferroni's test).
344
345   n.s., non-significant (p > 0.05). Data are presented as mean ± SEM.
346

347
Figure S6. The responses of cortical and hippocampal ACh to reward and punishment
(A) Schematic of ACh recording in the mPFC.
(B) Average response of mPFC ACh to water (left) and air puff (right) from a representative mouse.
(C) Quantification of the mean (left) and peak (right) mPFC ACh responses to water and air puff ($p < 0.001$, paired t-test).
(D) Schematic of ACh recording in the AUD.
(E) Average response of AUD ACh to water (left) and air puff (right) from a representative mouse.
(F) Quantification of the mean (left) and peak (right) AUD ACh responses to water and air puff ($p < 0.001$, paired t-test).
(G) Schematic of ACh recording in the CA1.
(H) Average response of CA1 ACh to water (left) and air puff (right) from a representative mouse.
(I) Quantification of the mean (left) and peak (right) CA1 ACh responses to water and air puff ($p < 0.001$, paired t-test).

***p < 0.001. Data are presented as mean ± SEM. Shaded areas represent SEM.

Figure S7. Activation of NAc→SI pathway does not affect ACh release in the cortex and hippocampus

(A, B) Trial-by-trial heatmap (top) and average (bottom) ACh responses in the mPFC while activating NAc^{D1}→SI (A) and NAc^{D2}→SI (B) with different laser durations in an example session.

(C, D) Trial-by-trial heatmap (top) and average (bottom) ACh responses in the AUD while activating NAc^{D1}→SI (C) and NAc^{D2}→SI (D) with different laser durations in an example session.

(E, F) Trial-by-trial heatmap (top) and average (bottom) ACh responses in the CA1 while activating NAc^{D1}→SI (E) and NAc^{D2}→SI (F) with different laser durations in an example session.

(G, H) Quantification of the mean (left), peak (middle) and AUC (right) of mPFC ACh responses while activating NAc^{D1}→SI (G) (left, $F_{(3, 20)} = 2.41$, $p = 0.097$; middle, $F_{(3, 20)} = 5.34$; $p = 0.0072$; right, $F_{(3, 20)} =$

2.43, p = 0.095) and NAc^{D2}→SI (H) (left, $F_{(3, 28)}$ = 2.81, p = 0.058; middle, $F_{(3, 28)}$ = 2.14, p = 0.12; right, $F_{(3, 28)}$ = 2.81, p = 0.058) with different laser durations. One-way ANOVA.

(I, J) Quantification of the mean (left), peak (middle) and AUC (right) of AUD ACh responses while activating NAc^{D1}→SI (I) (left, $F_{(3, 20)}$ = 4.50, p = 0.014; middle, $F_{(3, 20)}$ = 6.85, p = 0.0023; right, $F_{(3, 20)}$ = 4.49, p = 0.015) and NAc^{D2}→SI (J) (left, $F_{(3, 28)}$ = 0.54, p = 0.66; middle, $F_{(3, 28)}$ = 0.44, p = 0.72; right, $F_{(3, 28)}$ = 0.52, p = 0.67) with different laser durations. One-way ANOVA.

(K, L) Quantification of the mean (left), peak (middle) and AUC (right) of CA1 ACh responses while activating NAc^{D1}→SI (K) (left, $F_{(3, 24)}$ = 1.06, p = 0.38; middle, $F_{(3, 24)}$ = 1.35, p = 0.28; right, $F_{(3, 24)}$ = 1.06, p = 0.38) and NAc^{D2}→SI (L) (left, $F_{(3, 24)}$ = 1.78, p = 0.18; middle, $F_{(3, 24)}$ = 0.24, p = 0.87; right, $F_{(3, 24)}$ = 1.81, p = 0.17) with different laser durations. One-way ANOVA.

*p < 0.05, **p < 0.01; n.s., non-significant (p > 0.05). Data are presented as mean ± SEM. Shaded areas represent SEM.

389
Figure S8. BLA ACh responses shift from US to CS during cue-reward learning
(A) Average BLA ACh responses aligned to the cue onset (left), initial poke (middle), and reward retrieval
(right) in the pre-training (top), early-training (middle), and late-training (bottom) stages of the cue-reward
learning task from a representative mouse.
(B) Trial-by-trial heatmap of BLA ACh responses aligned to the cue onset (left) and reward retrieval (right)
at the pre-training (top), early-training (middle), and late-training (bottom) stages from the same mouse
as in (A). White dashed line, the cue onset; dark green dashed line, the initial poke; green dashed line, the
reward retrieval.
(C, D) Quantification of the mean (C) ($F_{(2, 143)} = 36.74$, $p < 0.001$) and peak (D) ($F_{(2, 143)} = 16.24$, $p <$
0.001) of BLA ACh responses to the cue at different training stages. One-way ANOVA followed by
Tukey's test.
(E-G) Quantification of the peak latency of BLA ACh responses to the cue (E) ($F_{(2, 143)} = 17.98$, $p < 0.001$),
initial poke (F) ($F_{(2, 143)} = 39.83$, $p < 0.001$), and reward retrieval (G) ($F_{(2, 143)} = 107.2$, $p < 0.001$) at different
training stages. One-way ANOVA followed by Tukey's test.
404
405 $*p < 0.05$, $**p < 0.01$, $***p < 0.001$. Data are presented as mean ± SEM. Shaded areas represent SEM.
406

407
Figure S9. Activation of NAc^{D1}→SI and NAc^{D2}→SI pathways affect reaction time but do not impair
movement
(A) Schematic of the approach to activate NAc^{D1}→SI pathway.
(B, C) Reaction time of the ChR2 (n = 6) and GFP (n = 8) mice in each session (B) (interaction, $F_{(11, 132)}$
= 1.90, p = 0.044; group, $F_{(1, 12)}$ = 11.78, p = 0.0050; two-way ANOVA), and average across sessions (C)
(p = 0.0050, t-test) with closed-loop stimulation of NAc^{D1}→SI in the cue-reward learning task.
(D, E) Number of total nose pokes of the ChR2 (n = 6) and GFP (n = 8) mice in each session (D)
(interaction, $F_{(11, 132)}$ = 0.69, p = 0.75; group, $F_{(1, 12)}$ = 1.70, p = 0.22; two-way ANOVA), and average
across sessions (E) (p = 0.22, t-test) with closed-loop stimulation of NAc^{D1}→SI in the cue-reward learning
task.
(F) Schematic of the approach to activate NAc^{D2}→SI pathway.
(G, H) Reaction time of the ChR2 (n = 7) and GFP (n = 4) mice in each session (G) (interaction, $F_{(11, 99)}$
= 1.62, p = 0.104; group, $F_{(1, 9)}$ = 8.36, p = 0.018; two-way ANOVA), and average across sessions (H) (p
= 0. 018, t-test) with closed-loop stimulation of NAc^{D2}→SI in the cue-reward learning task.
(I, J) Number of total nose pokes of the ChR2 (n = 7) and GFP (n = 4) mice in each session (I) (interaction,
$F_{(11, 99)}$ = 0.40, p = 0.95; group, $F_{(1, 9)}$ = 1.15, p = 0.31; two-way ANOVA), and average across sessions (J)
(p = 0.31, t-test) with closed-loop stimulation of NAc^{D2}→SI in the cue-reward learning task.
425
426 *p < 0.05, **p < 0.01; n.s., non-significant (p > 0.05). Data are presented as mean ± SEM.
427