The Neuropeptide Sulfakinin, a peripheral regulator of insect behavioral switch between mating and foraging

Hong-Fei Li[1,2], Bao Dong[1,2], Yuan-Yuan Peng[1,2], Hao-Yue Luo[1,2], Xiao-Lan Ou[1,2], Zheng-Lin Ren[1,2], Yoonseong Park[3*], Jin-Jun Wang[1,2*], Hong-Bo Jiang[1,2*]

[1] Key Laboratory of Entomology and Pest Control Engineering, College of Plant Protection, Southwest University, Chongqing 400715, China.

[2] Key Laboratory of Agricultural Biosafety and Green Production of Upper Yangtze River (Ministry of Education), Academy of Agricultural Sciences, Southwest University, Chongqing 400715, China.

[3] Department of Entomology, Kansas State University, Manhattan, KS 66506, United States

* Corresponding author: Yoonseong Park, Jin-Jun Wang and Hong-bo Jiang

Email: ypark@ksu.edu; wangjinjun@swu.edu.cn; jhb8342@swu.edu.cn

Author Contributions: H.L., J.W., and H.J. designed research; H.L., B.D., Y.P., H.L., X.O. and Z.R. performed research; H.L., B.D., H.L., and X.O. analyzed data; H.L., Y.P., J.W., and H.J. wrote the paper, and all authors read and approved the final manuscript.

Competing Interest Statement: The authors declare no competing interest.

Keywords: neuropeptide, behavior switching, peripheral olfaction, cholecystokinin

Abstract

Behavioral strategies for foraging and reproduction in the oriental fruit fly (*Bactrocera dorsalis*) are alternative options for resource allocation and are controlled by neuropeptides. Here we show that the behavioral switch between foraging and reproduction is associated with changes in antennal sensitivity. Starved flies became more sensitive to food odors while suppressing their response to opposite-sex pheromones. The gene encoding sulfakinin receptor 1 (*SkR1*) was significantly upregulated in the antennae of starved flies, so we tested the behavioral phenotypes of null mutants for the genes encoding the receptor (*skr1$^{-/-}$*) and its ligand sulfakinin (*sk$^{-/-}$*). In both knockout lines, the antennal responses shifted to mating mode even when flies were starved. This suggests that sulfakinin signaling via SkR1 promotes foraging while suppressing mating. Further analysis of the mutant flies revealed that sets of odorant receptor (OR) genes were differentially expressed. Functional characterization of the differentially expressed ORs suggested that sulfakinin directly suppresses the expression of ORs that respond to opposite-sex hormones while enhancing the expression of ORs that detect food volatiles. We conclude that sulfakinin signaling via SkR1, modulating OR expressions and leading to altered antenna sensitivities, is an important component in starvation-dependent behavioral change.

Main Text

Introduction

Foraging and mating behaviors are alternative options for resource allocation in insects. Starved and reproductively immature individuals intensify their foraging activity, whereas satiated and reproductively mature individuals increase their mating activity to facilitate reproduction (Chen et al., 2023; Cheriyamkunnel et al., 2021; Ebrahim et al., 2023; Ren et al., 2021). Such behavioral changes involve complex neural pathways triggered by neuropeptide signaling (González Segarra et al., 2023; Lin et al., 2022; Nässel and Zandawala, 2022; Schoofs et al., 2017; Sun et al., 2023; Zhang et al., 2022). The key regulators have been identified in the fruit fly *Drosophila melanogaster*, including the neuropeptides sulfakinin (Sk) (Wang et al., 2022; Wu et al., 2019) and short neuropeptide F (sNPF) (Ko et al., 2015; Root et al., 2011). Tyramine and diuretic hormone 31 then resolve the conflict between foraging and mating behavior in the central nervous system (Cheriyamkunnel et al., 2021; Lin et al., 2022).

In mammals, cholecystokinin (CCK) transmits satiety by activating cholecystokinin receptor 1, leading to the expression of genes encoding other anorectic and orexigenic peptides and receptors in vagal neurons (Beutler et al., 2017; Burdyga et al., 2008; Rachel et al., 2017). CCK is also involved in the central integration of sensory inputs to regulate sexual behavior (Bloch et al., 1987; Micevych et al., 2004). In insects, Sk, the ortholog of CCK, and its signals via two receptors, SkR1 (CCKLR-17D3 of *D. melanogaster*) and SkR2 (CCKLR-17D1), regulate sub-behavioral steps during foraging, food consumption, and mating in *D. melanogaster* (Fedina et al., 2023; Wang et al., 2022). For example, Sk suppresses the expression of the *Gr64f* gustatory receptor gene, which inhibits feeding as a signal for the

onset of satiety (Guo et al., 2021). Sk also inhibits both male and female sexual behavior (Wu et al., 2019). However, the complexity of Sk pathways, spatial and temporal dynamics of multiple ligands and receptors, is exemplified by cell-specific signaling via SkR1 for the promotion of mating receptivity in virgin females (Wang et al., 2022), which is an opposite action of the inhibition of sexual behavior (Wu et al., 2019). Modulation of peripheral sensory inputs, including odorant perception, is involved in the control of mating and foraging behavior in addition to the central neuronal circuits.

The oriental fruit fly, *Bactrocera dorsalis,* is a destructive agricultural pest insect. Control of *B. dorsalis* includes the use of the behavioral responses to semiochemicals, such as the components in the sex pheromones and in the food smell (Jaffar et al., 2023). Chemical components in the sex pheromone of this species includes 2,3,5-trimethyl pyrazine, 2,3,5,6-tetramethylpyrazine, ethyl laurate, ethyl myristate, ethyl cis-9-Hexadecenoate, and ethyl palmitate (Chen et al., 2023; Ren et al., 2021) and in the food smells are ethyl benzoate, ethyl butyrate, and methyl eugenol, among others. (Kamala Jayanthi et al., 2014; Zhu et al., 2018). The physiology and molecular mechanisms underlying the behavioral modulation involved in the different types of chemical attractants have not yet been studied in this species of insect.

We hypothesized that neuropeptide Sk-SkR and other neuropeptidergic systems are involved in the switch between foraging success and mating behaviors in the *B. dorsalis*. We were specifically interested in the alteration in odorant perceptions in the antennae leading to the success in foraging and/or mating depending on the physiological status. Sequential approaches, starting with the unbiased test for involvement of odorant perception in the

behavioral switch, led us to further in-depth investigations of the role of neuropeptide and Sk-SkR systems. We demonstrate that sulfakinin signaling via SkR1 in the antennae alters the expression of sets of odorant receptor (OR) genes during starvation enhancing the foraging success rate. Our results thus provide insight into sulfakinin signal on peripheral olfactory system that is coordinated with the central neuronal circuits for modulation of behavior.

Results

Altered antennal sensitivity is associated with starvation-induced foraging and satiation-induced mating success. We developed an assay to investigate the behavioral switch between foraging and mating induced by the physiological state of *B. dorsalis* (Fig. 1*A*, *B*). Foraging success increased with the duration of the starvation period (Fig. 1*D*, and S1), whereas courtship and copulation rates decreased, also with the starvation length-dependent manner (Fig. 1*G*, *H*). Starved flies tended to move directionally toward the food source located in the center of the arena (Fig. 1*C*).

We therefore tested antennal sensitivities toward artificial food and opposite-sex body extracts using an electroantennogram (EAG). Flies of both sexes starved for 12 h showed a greater EAG response to food odor (Fig. 1*E*, *F*) but a weaker response to opposite-sex body extracts (Fig. 1*I*, *J*). Furthermore, starved flies also showed greater EAG responses to food volatile components and weaker responses to sex pheromone components (Fig. S2). This suggested antenna-mediated olfaction is associated with and required for successful foraging and mating.

Starvation-induced *SkR1* mRNA in the antennae is required for the switch from mating to foraging behavior. To identify signaling molecules responsible for the olfactory-dependent

behavioral switch, we compared the expression profiles of neuropeptide signaling components in the antennae of fed and starved flies. We found that the components of three neuropeptide signaling systems were significantly upregulated in the antennae of starved flies, namely *short neuropeptide F receptor* (*sNPFR)*, *SIFamideR1,* and *SkR1*, the latter showing the strongest induction (Fig. 2*A*). *Sk* did not show the expression change in the antennae, but significantly increased in the head in comparison between those in starvation and after re-feeding, suggesting that the circulatory Sk released from the central nervous system could be the source for activation of SkR1 in the antennae (Fig. S3).

We introduced a 7-bp deletion into the sulfakinin gene (*Sk*) using the CRISPR/Cas9 system, causing a frameshift that abolished the sequence region encoding two predicted mature peptides: Sk-1 and Sk-2 (Fig. 2*B*, S4*A*, *B*). We confirmed the absence of an anti-sulfakinin immunoreactive signal in the central nervous system of the $sk^{-/-}$ flies (Fig. S4*C*). Only *SkR1* was upregulated in the starved flies among two predicted Sk receptor genes *SkR1* and *SkR2* (Fig. 2*A*, and S6). We therefore generated a *SkR1* null mutant ($skr1^{-/-}$) by introducing a 187-bp deletion using CRISPR/Cas9, resulting in a frameshift that resulted in a premature stop codon (Fig. 2*B*, and S5*A–C*). We confirmed the absence of the anti-SkR1 immunoreactive signal in the antennae of the $skr1^{-/-}$ flies (Fig. S5*D*).

Starved $sk^{-/-}$ and $skr1^{-/-}$ flies showed the same locomotor activity and body size as wild-type (WT) controls when we measured their velocity and body length (Fig. 2*C*, and S7). However, both null mutants consumed more food than WT flies (Fig. 2*D*), consistent with previous results in *D. melanogaster* (Guo et al., 2021). Interestingly, $sk^{-/-}$ and $skr1^{-/-}$ male flies consumed more food at all times compared to the wild type. However, the corresponding

mutant females consumed more food only at night. This may be due to the high basal feeding rate of females during the daytime, masking the increase in feeding in the knockout of Sk signaling. As shown in Figure 2D, WT females ate more food than males, and WT females consumed more food during the daytime. Another major difference between the null mutants and WT controls was that the mutant flies were less successful at foraging even after starvation, similar to the characteristics of satiated WT flies (Fig. 3A, B). However, foraging was not completely abolished. In addition, starved mutant females (Fig. 3C) and males (Fig. 3D) showed high mating success rates, as same rates as fed WT flies. We tested the behavioral phenotypes of heterozygous mutant of Sk knockout flies. The results showed that foraging and mating behaviors of Sk heterozygous mutants were unaffected during starvation, suggesting the mutants are completely recessive (Fig. S8).

We next tested antennal sensitivity to food odors and body extracts by EAG in each genotype using both fed and starved flies in each case. The starved $sk^{-/-}$ and $skr1^{-/-}$ flies showed significantly weaker responses to food odors compared to the starved WT flies, although the response of $sk^{-/-}$ flies was slightly stronger than that of fed WT flies (Fig. 3E, F and S8), which was similar to the behavioral response for a successful foraging rate (Fig. 3A, B). The antennae of starved $sk^{-/-}$ and $skr1^{-/-}$ flies showed the same EAG response to opposite-sex body extracts as fed WT flies (Fig. 3G, H and S9), which was significantly higher than that of starved WT flies.

Sets of ORs are differentially expressed when comparing satiated/starved WT and $sk^{-/-}$ flies. To gain insight into the molecular basis of alteration of olfactory sensitivities by the Sk-SkR1 system, we extracted RNA from the head epidermal tissues (including antennae) of

fed and starved flies for sequencing (RNA-Seq). We compared starved WT and starved $sk^{-/-}$

flies with satiated WT flies, focusing on the differentially expressed OR genes (Fig. 4*A*). We

identified three OR genes (*OR7a.4*, *OR7a.8* and *OR10a*) that were upregulated in the starved

WT flies but unchanged in starved $sk^{-/-}$ flies compared to satiated WT controls, and two

(*OR49a* and *OR63a*) that were downregulated in starved WT flies but unchanged in starved

$sk^{-/-}$ flies (Fig. 4*B*). OR67d.3 and OR74a mRNA levels were similar in the satiated and starved

WT flies. The differentially expressed genes identified by RNA-Seq were further validated by

real-time reverse-transcription PCR (qRT-PCR) (Fig. 4*B*).

Sulfakinin regulates the expressions of starvation-induced ORs that detect food

volatiles and starvation-suppressed ORs that detect pheromones. The OR genes

identified in the RNA-Seq experiments were expressed in HEK 293 cells followed by the

evaluation of functional responses to odorants using calcium mobilization assays. Tests with

single high doses of volatiles representing food or sex pheromones allowed the functional

categorization of ORs based on odorant specificities (Fig. 4*C*). OR10a was activated by ethyl

benzoate (EBE), OR7a.8 was activated by ethyl butyrate (EBU) or diethyl maleate (DM), and

OR7a.4 was activated by methyl eugenol (ME). The responses to these food odor volatiles

were all dose-dependent, with OR10a showing the strongest response to EBE (Fig. 4*E*). In

contrast, OR63a and OR49a showed dose-dependent responses to sex pheromone

components but not to food odor volatiles (Fig. 4*C*, *D*). Therefore, the upregulated *OR* genes

in starved WT flies, *OR7a.4*, *OR7a.8* and *OR10a*, were activated by the pheromonal

components, while down reulated genes, *OR49a* and *OR63a,* were activated by food volatiles.

Finally, we determined whether *SkR1* was expressed in OR neurons by testing the antennae

for immunoreactivity to SkR1 and odorant receptor co-receptor (Orco) antisera. A subset of

clustered OR neurons showed immunoreactivity to SkR1 (Fig. 4*F*), indicating that sulfakinin

acts directly on SkR1-positive OR neurons.

Discussion

Dynamic changes in the olfactory sensitivity of insect antennae are associated with

behavioral transitions in insects. Sensory inputs via olfactory perception often trigger certain

behaviors, including foraging and mating (Anton and Rössler, 2021; Gadenne et al., 2016).

Fine-tuning of the antennal OR repertoire would be an efficient way to co-opt behavioral

outputs (Anton and Rössler, 2021; Xu et al., 2024) as we have shown here for the oriental fruit

fly *B. dorsalis*. Changes in odorant sensitivity during transitions between foraging and mating

have been reported in previous studies. For example, the antennae of female *D. suzukii*

become significantly more sensitive to the host fruit odor isoamyl acetate after mating (Crava

et al., 2019). Similarly, starved *Aedes aegypti* mosquitoes downregulate the sensitivity of lactic

acid receptor neurons in grooved peg sensilla, which are used during host search (Davis, 1984;

Oliveira et al., 2017), whereas satiated mosquitoes increase the sensitivity of OR neurons in

trichoid sensilla to compounds representing the oviposition site (Siju et al., 2010). In this study,

we determined the mechanisms underlying the changes in antennal sensitivity when switching

between foraging and mating behavior. Sk-SkR1 signaling was required to increase the

success of foraging and to suppress mating during a period of starvation. The activation of

SkR1 on the antennal OR neurons upregulates the expression of ORs that detect food

volatiles while suppressing those that detect pheromones.

The coordinated actions of sulfakinin on multiple downstreams, the antennae and other parts of the peripheral and central nervous system in *B. dorsalis*, are not yet fully understood. Sulfakinin is generally known as a satiety signal that reduces food intake and suppresses the digestive system in *D. melanogaster* and a number of different invertebrate species (Nässel and Wu, 2022), whereas we found that the Sk-SkR1 signal is required to promote foraging by making the antennae more sensitive to food odorants during starvation. In *D. melanogaster*, sulfakinin inhibits the sugar receptor GR64 in the proboscis and proleg tarsi, thus reducing food ingestion (Guo et al., 2021). Our data also support the role of sulfakinin as a satiety signal because feeding rate was increased in $sk^{-/-}$ and $skr1^{-/-}$ null mutants (Fig. 2*D*). Sulfakinin was shown to inhibit both male and female sexual behavior in *D. melanogaster* (Wu et al., 2019), in agreement with our results. In addition to the previous study showing the roles of central nervous system in Sk-mediated inhibition of sexual behavior in *D. melanogaster*, we described a Sk direct action on the peripheral olfactory system in *B. dorsalis.* The underlying mechanism is the secretion of the mature peptide Dsk-2 (but not Dsk-1) by Dsk-MP1 and Dsk-MP3 neurons, which then triggers downstream neurons expressing SkR1 in the central nervous system to suppress male mating behavior (11). However, there was a contradictory report showing the role of sulfakinin promoting the receptivity in mating in virgin females (Wang et al., 2022). The activity of sulfakinin peptides Sk1 and Sk2, through the receptors SkR1 and SkR2, is therefore likely to be complex, with spatially and temporally dynamic effects in multiple downstream systems.

Although successful foraging and EAG responses to food odors were strongly reduced in both null mutants, there was still a greater response than observed in fed WT controls (Fig. 3*A*,

B, *E*, *F*). This incomplete suppression suggests that other mechanisms, in addition to the main Sk-SkR1 signal, are relevant to the response. Indeed, sNPF and tachykinin signaling have both been described in the modulation of olfactory sensitivity in *D. melanogaster*, with sNPF enhancing sensitivity to food odors and tachykinin inhibiting sensitivity to aversive odors (Ignell et al., 2009; Ko et al., 2015; Root et al., 2011). Furthermore, neuropeptide F (NPF) and SIFamide also regulate foraging behavior at the level of the olfactory circuit (Jennifer and Yi, 2013; Martelli et al., 2017; Wang et al., 2013). Notably, NPF and tachykinin also regulate mating behavior at the peripheral level. Tachykinin inhibits courtship behavior by mediating the perception of an anti-aphrodisiac pheromone in male *D. melanogaster*. (Shankar et al., 2015). NPF neurons are also required to detect female sex pheromones (Gendron et al., 2014; Kuo et al., 2012). Similarly, SIFamide also regulates mating behavior, although it is not yet clear whether it acts via the olfactory circuit (Sellami and Veenstra, 2015; Terhzaz et al., 2007). We have not yet investigated the roles of other neuropeptides in *B. dorsalis* foraging and mating behavior, but we found that starvation-induced the neuropeptide receptor genes SkR1, sNPFR, and *SIFamideR1* ($P < 0.05$). In addition, tachykinin, SIFamide, and ion transport peptide (ITP)-like peptide were also induced by starvation, although the increase was not statistically significant. Therefore, other neuropeptides in addition to sulfakinin are likely to be involved in the switch between different sets of ORs in the antenna as a means to control behavior.

The evolutionary origin of CCK/sulfakinin signaling can be traced back to the common ancestor of the Bilateria (Mirabeau and Joly, 2013; Nässel and Wu, 2022; Staljanssens et al., 2011; Tinoco et al., 2021; Yu and Smagghe, 2014). Most coelomate and pseudocoelomate phyla produce CCK/sulfakinin peptides and orthologs of the corresponding receptors.

Sulfakinin is involved in foraging, satiety, locomotion, development, aggression and reproductive behavior in insects (Fedina et al., 2023; Nässel and Wu, 2022; Nassel and Zandawala, 2019) although some insect lineages have lost both the peptide and its receptors (Consortium, 2010; Hauser et al., 2010). CCK is also involved in satiety, memory, anxiety, and depression in mammals (Jens et al., 2007; Lau et al., 2023; Rana et al., 2022; Shen et al., 2019). The many functions of CCK/sulfakinin in diverse taxonomic groups frequently include signaling between the digestive system (endoderm) and the neuronal system (ectoderm) (Nässel and Wu, 2022). This suggests an ancestral function associated with coelomic development, and that additional complexity has evolved in each lineage.

In summary, we found that the neuropeptide sulfakinin mediates the transition between foraging and mating behavior by activating the receptor SkR1 and reprogramming the peripheral OR repertoire. We propose a working model to characterize this mechanism, in which Sk-SkR1 regulates the expression of different sets of ORs, leading to the behavioral switch (Fig. 5). The modulation of peripheral sensory systems by sulfakinin, coordinating with the central neural circuit also controlled by sulfakinin, is the major trigger for a successful behavioral switch. We have thus provided new insight into the control of behavior by sulfakinin in *B. dorsalis*, which is one of the most widely distributed, fecund, and invasive insect pests to threaten global agriculture (Wan and Yang, 2016; Zhao et al., 2024).

Materials and methods

Insects

The wild-type (WT) oriental fruit flies were originally collected from Hainan province, China in 2008 and were reared in the laboratory (27.5 ± 0.5 °C, 75 ± 5% relative humidity, 14-h

photoperiod). Larvae and adults were fed with artificial food as previously described (Shen et al., 2013). Artificial food for adults, consisting of yeast powder, honey, sugar, ascorbic acid and water, was also used as a food source for foraging behavior assays.

Behavioral assays

Virgin females and males, 3-day old, were placed each in groups of 25 flies and reared at 27.5 °C and 60% humidity, and 9-day old flies were subjected to behavioral assays. Food consumption rates of individual flies were measured by the amount Bromophenol Blue dye (BPB) in the fly after the food including 0.1% (BPB) was provided for 1 h for ad libitum feeding, which is long enough time to allow satiation to occur. BPB was measured from decapitated and homogenized bodies in 1 ml PBS buffer with 1% Triton X-100. Absorbance of the supernatant after centrifugation (13000 rpm) for 3 min was measured at 614 nm on a xMarkerTM Microplate Spectrophotometer (Bio-Rad). Flies were starved for 12 h prior to the experiments with only water. Foraging behavior assay was performed according to previously published methods with modifications (Li et al., 2022). In brief, single flies were introduced into chambers (100 mm diameter x 10 mm height) and acclimatized for 20 min. The foraging behavior was recorded using a camera (Sony HDR CX405) for 15 min at 30 fps. Successful foraging was counted when the fly stayed 5 s or longer within the 5 mm radius of the center where 10 µL of artificial food was placed. The experiments consisted of the assays of 3 biological replications for each experimental condition. The behavior was analyzed by EasyFlyTracker (Qu et al., 2022) with the default parameters.

Mating behavior assay was conducted in the courtship chambers (diameter: 3.5 cm; height: 1.5cm per layer). The virgin females and males were cold anesthetized at ⊡20 °C for 1

min and introduced into the courtship chamber separated by a removable transparent film. After 1 h acclimation, the film was removed to allow the paired flies to get in contact, and mating behavior was recorded by a video camera for 15 min. We measured the level of courtship in mutant males and receptivity in mutant females respectively. For courtship assay, a mutant male ($sk^{-/-}$ or $skr^{-/-}$) and a WT virgin female were introduced into the chambers. Courtship rate is the percentage of the mutant males performs attempted copulation, which was used to measure mutant courtship level. For female receptivity assay, a mutant female ($sk^{-/-}$ or $skr^{-/-}$) and a WT virgin male were introduced into the chambers. Copulation rate is the percentage of the mutant females had engaged in copulation, which is used to measure the mutant female receptivity. The number of flies that had performed attempted copulation or engaged in copulation by the end of each 1 min interval were summed within 15 min and plotted as a percentage of total flies for each time point.

Electroantennograms

Electroantennograms were used to record the stimulus-dependent changes in the electrical potential summed over the whole antenna. To test volatiles were food odor or body extract of each sex on the flies conditioned for satiated and starved. Different genotypes of the flies were wild-type, Sk mutant, and SkR1 mutants. Female body extracts were tested by male antennae and male body extracts by female antennae (Table S2). Body extracts were prepared by washing 15 9-day old virgin flies of each sex in 1 ml of hexane for 4 hours (Chen et al., 2023; Dweck et al., 2015; Ebrahim et al., 2023) at room temperature with gentle shaking.

For electroantennograms, similar to previous study (Li et al., 2022), the head (including the antennae) of fly was excised and a glass capillary filled with 0.9% NaCl was inserted into

the cut end as the reference electrode. The tip of an antenna was placed into another glass capillary filled with 0.9% NaCl as the recording electrode. The reference electrode was connected to a grounding electrode, while the recording electrode was connected to a 10× high-impedance DC amplifier (Syntech; The Netherlands). The signal was then sent to an analog/digital converter (IDAC-4, USB, Syntech) and transferred to a computer. Finally, the data were analyzed and saved by using GC-EAD software (Syntech). A filter paper strip (3 × 0.75 cm), loaded with 10 µL of the odor solution or body extract, was placed into a pipette and delivered a stimulus into a constant, humidified airstream flowing at 100 ml/min by a stimulus controller (Syntech, Hilversum, Netherlands). Two stimuli were made for 0.5s duration with a 30s interval. Average EAG response, as shown by mV changes, was used for the calculation. EAG for sex pheromones and body surface extracts were measured from 21:00□23:00, and food odor was measured from 9:00-11:00.

Quantitative real-time reverse transcription PCR

Antennal tissues from flies of different genotypes with different treatments were dissected and immediately stored at □80 °C. Total RNA was extracted with Trizol Reagent (Invitrogen) following the manufacturer's protocol. First-strand cDNA was synthesized using the PrimeScript 1st Strand cDNA Synthesis Kit (Takara, Dalian, China) and specific primers designed using the online software, Primer 3 (http://primer3.ut.ee/, Table S1). Quantitative real-time reverse transcription PCR (qRT-PCR) was performed using a CFX Connect Real-Time System (Bio-Rad, Hercules, CA) with NovoStart SYBR PCR SuperMix kit (Novoprotein, Shanghai, China). Two reference genes (Shen et al., 2010), *α-tubulin* (GenBank: GU269902) and RPS3 (GenBank: XM_011212815), were used selected to normalize the

expression levels for the calculation in qBASE (Hellemans et al., 2007). Each experiment comprised 3 independent biological replicates and two technical replicates.

CRISPR/Cas9-mediated mutants

In vitro synthesis of single guide RNA was performed as previously described (Li et al., 2022). In brief, the suitable target sites of *Sk* and *SkR1* genes were searched by CasOT (Xiao et al., 2014), and the corresponding single guide RNAs (sgRNAs) were synthesized *in vitro* and purified using the GeneArt Precision gRNA Synthesis Kit (Invitrogen, Thermo Fisher Scientific, Waltham, MA, USA). We designed and synthesized one sgRNA for the *Sk* gene (Fig. S4) and two single guide RNAs for the *SkR1* gene (Fig. S5).

The sgRNA and cas9 protein (Invitrogen) were gently pipetted and mixed in nuclease-free water to form a mixture of 400 ng/μL sgRNA and 300 ng/μL Cas9 protein. Eggs collected within 20 min of oviposition were injected within 2 h with the mixture of sgRNA and Cas9 protein using an IM300 Electric Microinjector (Narishige, Tokyo Japan). The injected eggs were incubated at 27.5 ± 0.5 °C and 75 ± 5% relative humidity for hatching.

Mutant screening and off-target analysis

Heterozygous G1 offspring were obtained using 9-day-old virgin adults (G0) crossed to wild-type opposite sex. Suitable mutant G1 ($sk^{+/-}$ and $skr1^{+/-}$) were backcrossed to WT flies for at least eight generations, then homozygous mutant flies ($sk^{-/-}$ and $skr1^{-/-}$) were obtained by self-crossing. Genomic DNA was extracted from individual flies (G0) or a single hind leg (G1 and later generations) by incubating tissue samples in 30 μL InstaGene Matrix (Bio-Rad, Hercules, CA, USA) at 56 °C for 30 min and inactivating the enzyme at 100 °C for 10 min. Fragments of 300–500 bp spanning the sgRNA target sites were amplified using the primers

(Table S1). The fragments were genotyped using the QIAxcel Advanced System (Qiagen, Hilden Germany), and potential edited products were directly sequenced or cloned into vector pGEM-T Easy (Promega, Wisconsin, Madison, USA) for Sanger sequencing to identify the mutation.

To assess off-target effects, CasOT was used to search potential off-target sites for the *Sk* and *SkR1* genes. All loci with one potential mismatch allowed in the seed region and up to three potential mismatches in the non-seed region were searched on the genome sequence (https://db.cngb.org/search/project/CNP0003192/). Two potential off-target sites were found for *Sk* gene and three for *SkR1* gene. PCR amplification and sequencing of fragments including potential off-target sites were performed to check for mutations. In examination of the potential off-target sites with > 4 bp mismatches to the sgRNA sequences, no mutations were found in the off-sites (Fig. S10).

Immunohistochemistry

Rabbit anti-BdSkR1 antibody was generated by immunization with the peptide N'-RQGLRKCRDQWP-C'. Mouse anti-BdOrco antibody was raised by immunization with two peptides, N'-GTNPNGLTRKQE-C' and N'-HWYDGSEEAKTF-C'. Peptide were synthesized and conjugated with keyhole limpet hemocyanin through additional cysteine residues at their N termini by Zoonbio Biotechnology (Nanjing, China). Rabbit anti-BdSk antibody was a gift from Dr. Chuan Zhou (Wu et al., 2020). The antisera used to recognize Sk peptide were raised in New Zealand white rabbits using the synthetic peptide N'-GGDDQFDDYGHMRFG-C.

Immunohistochemistry was performed as previously described (Li et al., 2022). In brief, whole brains of 9-day-old flies were dissected in phosphate-buffered saline (PBS), fixed

overnight in 4% paraformaldehyde (PFA) in PBS at 4 °C, and washed (3 × 15 min) in PBST

(0.5% Triton X-100 in PBS). The tissues were blocked in 5% normal goat serum for 20 min at

room temperature, then incubated in primary antibody for 48 h at 4 °C. After three washes in

PBST (3 × 15 min), the tissues were then incubated in secondary antibody overnight at 4 °C.

After washing in PBST again, the tissues were mounted on a slide with 100% glycerol for

imaging. Adult antennae were dissected and embedded in Tissue-Tek O.C.T. Compound

(Sakura, Tokyo, Japan). Sections (18 µm) were prepared using a Thermo HM525 NX

microtome at □22 °C and then fixed in 4% PFA in PBS at room temperature for 2 h. The

subsequent procedures were as described above for the brain tissues. Primary antibody used:

rabbit anti-Sk (1:1000), rabbit anti-SkR1 (1:500), mouse anti-Orco (1:200). Secondary

antibodies used: goat anti-rabbit IgG conjugated to Alexa Fluor 488 (1:1000, Cell Signaling

Technology), goat anti-mouse IgG conjugated to Alex Fluor 647 (1:1000, Cell Signaling

Technology). Samples were imaged using an LSM780 confocal microscope (Carl Zeiss, Jena,

Germany) and processed with Image J software (Schneider et al., 2012).

RNA-seq analysis

Total RNA was extracted from the head epidermal tissues, including the antennae, of fed

WT flies, starved WT flies, and starved $sk^{-/-}$ flies using TRIzol reagent (Invitrogen). Three

replicates were performed. Library construction and sequencing was performed by Novogene

with Illumina HiSeq2000 platform (Novogene Bioinformatics Technology Co.Ltd, Beijing,

China). The raw data were analyzed after filtering the low-quality sequences by fastp with the

default parameters (Chen et al., 2018). Sequences were aligned to the *B. dorsalis* genome

(https://db.cngb.org/search/project/CNP0003192/) using Hisat2 (version 2.2.1) (Kim et al.,

2015; Yang et al., 2023). Samtools (version 1.7) was used to convert sam files to bam files in the process (Li et al., 2009). Finally, to calculate the count, we use featureCounts (version 2.0.1) (Liao et al., 2014). The expression level of genes from the RNA sequencing was normalized by the FPKM method. Differential expression analysis was performed using the DESeq2 R package (1.16.1) (Anders and Huber, 2010).

Functional heterologous expression of ORs and calcium imaging

Heterologous expressions of ORs were made in Human Embryonic Kidney-293 (HEK-293) cells, which were cultured in Dulbecco's modified Eagle medium (DMEM, 11995065, Gibco, Thermo Fisher Scientific, USA) containing 10% Fatal Bovine Serum (FBS, 10091148, Gibco) and 1% penicillin-streptomycin-Amphotericin B solution (C0224-100 ml, Beyotime) with 5% CO_2 at 37 °C. Cells for transfection were seeded onto a 96-well black well plate (165305, Thermo Fisher Scientific, USA). Approximately 50%–70% confluent HEK-293 cells were transfected by 2 μg pcDNA3.1 plasmid containing respective olfactory receptors with BdOrco of *B. dorsalis* using *Trans*IT®-LT1 Transfection Reagent (MIR 2300, Mirusbio) for 24 h.

For calcium imaging experiment, the culture media was replaced to 100 μl of 2.5 μM Fluo-4 AM (F14201, Invitrogen) and incubated at 37 °C with 5% CO_2 for 1 h. Subsequently, after another 30 min of incubation at room temperature, the culture media containing Fluo-4 was removed, and the cells were washed three times with Hanks' balanced salt solution (HBSS; 14065056, Gibco, Thermo Fisher Scientific, USA) in order to fully remove the Fluo-4 residue. Then, 99 ul of HBSS was added to each well and plate was placed in the dark at RT to be tested. Finally, 1 μL of test odor was added to each well and fluorescence immediately was measured for at least 80 s. Finally, images were captured by a Zeiss Observer 7 inverted

microscope (Carl Zeiss, Jena, Germany). Test odors were dissolved in dimethylsulfoxide

(DMSO, ST038-100ml, Beyotime) and prepared five final concentrations: 10^{-7}, 10^{-6}, 10^{-5},

10^{-4} M and 10^{-3} M. the fluorescent ratio (ΔF/F) was calculated as peak intensity divided by

basal intensity (intensity before ligand perfusion). For all experiment, the response to DMSO

was used as negative control.

Statistical analysis

For qRT-PCR result analysis, relative expression levels were calculated using qBase

(Hellemans et al., 2007) based on the expression of the reference genes. Statistical analysis is

performed using GraphPad Prism and indicated in each figure legend. Data were first verified

for normal distribution using the Shapiro-Wilk normality test. One-way ANOVA, followed by

Tukey's multiple comparisons (if normal distribution) and Kruskal-Wallis and post-hoc

Mann-Whitney U-tests (if non-normal distribution), was used for comparisons among multiple

groups. Values are reported as means ± standard errors of the mean (SEM).

Acknowledgments

This research was supported by funding from the National Natural Science Foundation of China (32072491, 31772233), National Key R&D Program of China (2022YFC2601000), 111 Project (B18044), and the China Agriculture Research System of MOF and MARA. We are grateful to Dr. Chuan Zhou and Dr. Fengming Wu (Institute of Zoology, Chinese Academy of Sciences) for the DSK antibody. We thank Dr. Tao Wang (Institute of Zoology, Chinese Academy of Sciences) for her graphical assistance for this manuscript.

References

Anders S., Huber W., 2010. Differential expression analysis for sequence count data. Genome Biol. 11(10), R106.

Anton S., Rössler W., 2021. Plasticity and modulation of olfactory circuits in insects. Cell Tissue Res. 383(1), 149-164.

Beutler L.R., Chen Y., Ahn J.S., Lin Y.-C., Essner R.A., Knight Z.A., 2017. Dynamics of Gut-Brain Communication Underlying Hunger. Neuron. 96(2), 461-475.e465.

Bloch G.J., Babcock A.M., Gorski R.A., Micevych P.E., 1987. Cholecystokinin stimulates and inhibits lordosis behavior in female rats. Physiol. Behav. 39(2), 217-224.

Burdyga G., de Lartigue G., Raybould H.E., Morris R., Dimaline R., Varro A., Thompson D.G., Dockray G.J., 2008. Cholecystokinin regulates expression of Y2 receptors in vagal afferent neurons serving the stomach. J. Neurosci. 28(45), 11583-11592.

Chen S.F., Zhou Y.Q., Chen Y.R., Gu J., 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 34(17), i884-i890.

Chen Y.Y., Zhang Y.H., Ai S.P., Yi X., 2023. Female semiochemicals stimulate male courtship but dampen female sexual receptivity. Proc. Natl. Acad. Sci. U. S. A. 120(49), e2311166120.

Cheriyamkunnel S.J., Rose S., Jacob P.F., Blackburn L.A., Glasgow S., Moorse J., Winstanley M., Moynihan P.J., Waddell S., Rezaval C., 2021. A neuronal mechanism controlling the choice between feeding and sexual behaviors in Drosophila. Curr. Biol. 31(19), 4231-4245.e4234.

Consortium T.I.A.G., 2010. Genome Sequence of the Pea Aphid Acyrthosiphon pisum. PLOS Biol. 8(2), e1000313.

Crava C.M., Sassù F., Tait G., Becher P.G., Anfora G., 2019. Functional transcriptome analyses of Drosophila suzukii antennae reveal mating-dependent olfaction plasticity in females. Insect Biochem. Mol. Biol. 105, 51-59.

Davis E.E., 1984. Regulation of sensitivity in the peripheral chemoreceptor systems for host-seeking behaviour by a haemolymph-borne factor in Aedes aegypti. J. Insect Physiol. 30(2), 179-183.

Dweck H.K., Ebrahim S.A., Thoma M., Mohamed A.A., Keesey I.W., Trona F., Lavista-Llanos S., Svatos A., Sachse S., Knaden M., Hansson B.S., 2015. Pheromones mediating copulation and attraction in Drosophila. Proc. Natl. Acad. Sci. U. S. A. 112(21), E2829-2835.

Ebrahim S.A.M., Dweck H.K.M., Weiss B.L., Carlson J.R., 2023. A volatile sex attractant of tsetse flies. Science. 379(6633), eade1877.

Fedina T.Y., Cummins E.T., Promislow D.E.L., Pletcher S.D., 2023. The neuropeptide drosulfakinin enhances choosiness and protects males from the aging effects of social perception. Proc. Natl. Acad. Sci. U. S. A. 120(51), e2308305120.

Gadenne C., Barrozo R.B., Anton S., 2016. Plasticity in Insect Olfaction: To Smell or Not to Smell? Annu. Rev. Entomol. 61(1), 317-333.

Gendron C.M., Kuo T.-H., Harvanek Z.M., Chung B.Y., Yew J.Y., Dierick H.A., Pletcher S.D., 2014. Drosophila Life Span and Physiology Are Modulated by Sexual Perception and Reward. Science. 343(6170), 544-548.

González Segarra A.J., Pontes G., Jourjine N., Del Toro A., Scott K., 2023. Hunger- and thirst-sensing neurons modulate a neuroendocrine network to coordinate sugar and water ingestion. eLife. 12.

Guo D., Zhang Y.J., Zhang S., Li J., Guo C., Pan Y.F., Zhang N., Liu C.X., Jia Y.L., Li C.Y., Ma J.Y., Nassel D.R., Gao C.F., Wu S.F., 2021. Cholecystokinin-like peptide mediates satiety by inhibiting sugar attraction. PLoS Genet. 17(8), e1009724.

Hauser F., Neupert S., Williamson M., Predel R., Tanaka Y., Grimmelikhuijzen C.J.P., 2010. Genomics and Peptidomics of Neuropeptides and Protein Hormones Present in the Parasitic Wasp Nasonia vitripennis. J. Proteome Res. 9(10), 5296-5310.

Hellemans J., Mortier G., De Paepe A., Speleman F., Vandesompele J., 2007. qBase relative quantification framework and software for management and automated analysis of real-time quantitative PCR data. Genome Biol. 8(2), R19.

Ignell R., Root C.M., Birse R.T., Wang J.W., Nässel D.R., Winther Å.M.E., 2009. Presynaptic peptidergic modulation of olfactory receptor neurons in drosophila. Proc. Natl. Acad. Sci. U. S. A. 106, 13070–13075.

Jaffar S., Rizvi S.A., Lu Y. (2023). Understanding the Invasion, Ecological Adaptations, and Management Strategies of Bactrocera dorsalis in China: A Review. *Horticulturae, 9*, 1004. Retrieved from

Jennifer B., Yi Z., 2013. Graded encoding of food odor value in the Drosophila brain. J. Neurosci. 33(40), 15693.

Jens F.R., Lennart F.H., Jens P.G., Thomas V.O.H., 2007. The biology of cholecystokinin and gastrin peptides. Curr. Top. Med. Chem. 7(12), 1154-1165.

Kamala Jayanthi P.D., Kempraj V., Aurade R.M., Venkataramanappa R.K., Nandagopal B., Verghese A., Bruce T.J., 2014. Specific volatile compounds from mango elicit oviposition in gravid Bactrocera dorsalis females. J Chem Ecol. 40(3), 259-266.

Kim D., Langmead B., Salzberg S.L., 2015. HISAT: a fast spliced aligner with low memory requirements. Nat. Methods. 12(4), 357-360.

Ko K.I., Root C.M., Lindsay S.A., Zaninovich O.A., Shepherd A.K., Wasserman S.A., Kim S.M., Wang J.W., 2015. Starvation promotes concerted modulation of appetitive olfactory behavior via parallel neuromodulatory circuits. Elife. 4, e08298.

Kuo T.H., Fedina T.Y., Hansen I., Dreisewerd K., Dierick H.A., Yew J.Y., Pletcher S.D., 2012. Insulin Signaling Mediates Sexual Attractiveness in Drosophila. PLOS Genet. 8(4), e1002684.

Lau S.H., Young C.H., Zheng Y., Chen X., 2023. The potential role of the cholecystokinin system in declarative memory. Neurochem. Int. 162, 105440.

Li H., Handsaker B., Wysoker A., Fennell T., Ruan J., Homer N., Marth G., Abecasis G., Durbin R., 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 25(16), 2078-2079.

Li H.F., Huang X.Y., Yang Y.H., Chen X.F., Yang Y., Wang J.J., Jiang H.B., 2022. The short neuropeptide F receptor regulates olfaction-mediated foraging behavior in the oriental fruit fly Bactrocera dorsalis (Hendel). Insect Biochem. Mol. Biol. 140.

Liao Y., Smyth G.K., Shi W., 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 30(7), 923-930.

Lin H.H., Kuang M.C., Hossain I., Xuan Y., Beebe L., Shepherd A.K., Rolandi M., Wang J.W., 2022. A nutrient-specific gut hormone arbitrates between courtship and feeding. Nature. 602(7898), 632-638.

Martelli C., Pech U., Kobbenbring S., Pauls D., Bahl B., Sommer M.V., Pooryasin A., Barth J., Arias C.W.P., Vassiliou C., Luna A.J.F., Poppinga H., Richter F.G., Wegener C., Fiala A., Riemensperger T., 2017. SIFamide Translates Hunger Signals into Appetitive and Feeding Behavior in Drosophila. Cell Rep. 20(2), 464-478.

Micevych P., Akesson T., Elde R., 2004. Distribution of cholecystokinin-immunoreactive cell bodies in the male and female rat: II. Bed nucleus of the stria terminalis and amygdala. J. Comp. Neurol. 269(3), 381-391.

Mirabeau O., Joly J.S., 2013. Molecular evolution of peptidergic signaling systems in bilaterians. Proc. Natl. Acad. Sci. U. S. A. 110(22), E2028-E2037.

Nässel D.R., Wu S.F., 2022. Cholecystokinin/sulfakinin peptide signaling: conserved roles at the intersection between feeding, mating and aggression. Cell Mol. Life Sci. 79(3).

Nassel D.R., Zandawala M., 2019. Recent advances in neuropeptide signaling in Drosophila, from genes to physiology and behavior. Prog. Neurobiol. 179, 101607.

Nässel D.R., Zandawala M., 2022. Endocrine cybernetics: neuropeptides as molecular switches in behavioural decisions. Open Biol. 12(7).

Oliveira P.L., Christ P., Reifenrath A., Kahnt J., Hauser F., Hill S.R., Schachtner J., Ignell R., 2017. Feeding-induced changes in allatostatin-A and short neuropeptide F in the antennal lobes affect odor-mediated host seeking in the yellow fever mosquito, Aedes aegypti. Plos One. 12(11).

Qu S.S., Zhu Q.J., Zhou H., Gao Y., Wei Y., Ma Y., Wang Z., Sun X., Zhang L., Yang Q., Kong L., Zhang L., 2022. EasyFlyTracker: A simple video tracking python package for analyzing adult Drosophila locomotor and sleep activity to facilitate revealing the effect of psychiatric drugs. Front. Behav. Neurosci. 15.

Rachel A.E., Alison G.S., Adam A.J., Anna R.R., Zoe D.T., Matthew E.C., 2017. AgRP Neurons Can Increase Food Intake during Conditions of Appetite Suppression and Inhibit Anorexigenic Parabrachial Neurons. J. Neurosci. 37(36), 8678.

Rana T., Behl T., Sehgal A., Singh S., Sharma N., Abdeen A., Ibrahim S.F., Mani V., Iqbal M.S., Bhatia S., Abdel Daim M.M., Bungau S., 2022. Exploring the role of neuropeptides in depression and anxiety. Prog. Neuropsychopharmacol. Biol. Psychiatry. 114, 110478.

Ren L., Ma Y.G., Xie M.X., Lu Y.Y., Cheng D.F., 2021. Rectal bacteria produce sex pheromones in the male oriental fruit fly. Curr. Biol. 31(10), 2220-2226 e2224.

Root C.M., Ko K.I., Jafari A., Wang J.W., 2011. Presynaptic facilitation by neuropeptide signaling mediates odor-driven food search. Cell. 145(1), 133-144.

Schneider C.A., Rasband W.S., Eliceiri K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods. 9(7), 671-675.

Schoofs L., De Loof A., Van Hiel M.B., 2017. Neuropeptides as Regulators of Behavior in Insects. Annu. Rev. Entomol. 62, 35-52.

Sellami A., Veenstra J.A., 2015. SIFamide acts on fruitless neurons to modulate sexual

behavior in Drosophila melanogaster. Peptides. 74, 50-56.

Shankar S., Chua J.Y., Tan K.J., Calvert M.E.K., Weng R., Ng W.C., Mori K., Yew J.Y., 2015. The neuropeptide tachykinin is essential for pheromone detection in a gustatory neural circuit. eLife. 4.

Shen C.J., Zheng D., Li K.X., Yang J.M., Pan H.Q., Yu X.D., Fu J.Y., Zhu Y., Sun Q.X., Tang M.Y., Zhang Y., Sun P., Xie Y., Duan S.M., Hu H.L., Li X.M., 2019. Cannabinoid CB1 receptors in the amygdalar cholecystokinin glutamatergic afferents to nucleus accumbens modulate depressive-like behavior. Nat. Med. 25(2), 337-349.

Shen G.M., Dou W., Huang Y., Jiang X.Z., Smagghe G., Wang J.J., 2013. In silico cloning and annotation of genes involved in the digestion, detoxification and RNA interference mechanism in the midgut of Bactrocera dorsalis [Hendel (Diptera: Tephritidae)]. Insect Mol. Biol. 22(4), 354-365.

Shen G.M., Jiang H.B., Wang X.N., Wang J.J., 2010. Evaluation of endogenous references for gene expression profiling in different tissues of the oriental fruit fly Bactrocera dorsalis (Diptera: Tephritidae). BMC Mol. Biol. 11(76).

Siju K.P., Hill S.R., Hansson B.S., Ignell R., 2010. Influence of blood meal on the responsiveness of olfactory receptor neurons in antennal sensilla trichodea of the yellow fever mosquito, Aedes aegypti. J. Insect Physiol. 56(6), 659-665.

Staljanssens D., Azari E.K., Christiaens O., Beaufays J., Lins L., Van Camp J., Smagghe G., 2011. The CCK(-like) receptor in the animal kingdom: functions, evolution and structures. Peptides. 32(3), 607-619.

Sun M.S., Ma M.Z., Deng B.W., Li N., Peng Q.L., Pan Y.F., 2023. A neural pathway underlying hunger modulation of sexual receptivity in Drosophila females. Cell Rep. 42(10).

Terhzaz S., Rosay P., Goodwin S.F., Veenstra J.A., 2007. The neuropeptide SIFamide modulates sexual behavior in Drosophila. Biochem. Biophys. Res. Commun. 352(2), 305-310.

Tinoco A.B., Barreiro-Iglesias A., Yañez Guerra L.A., Delroisse J., Zhang Y., Gunner E.F., Zampronio C.G., Jones A.M., Egertová M., Elphick M.R., 2021. Ancient role of sulfakinin/cholecystokinin-type signalling in inhibitory regulation of feeding processes revealed in an echinoderm. eLife. 10, e65667.

Wan F.H., Yang N.W., 2016. Invasion and management of agricultural alien insects in China. Annu. Rev. Entomol. 61, 77-98.

Wang T., Jing B., Deng B., Shi K., Li J., Ma B., Wu F., Zhou C., 2022. Drosulfakinin signaling modulates female sexual receptivity in Drosophila. Elife. 11.

Wang Y., Pu Y., Shen P., 2013. Neuropeptide-Gated Perception of Appetitive Olfactory Inputs in Drosophila Larvae. Cell Rep. 3(3), 820-830.

Wu F.M., Deng B.W., Xiao N., Wang T., Li Y.N., Wang R.C., Shi K., Luo D.G., Rao Y., Zhou C., 2020. A neuropeptide regulates fighting behavior in Drosophila melanogaster. Elife. 9, e54229.

Wu S.F., Guo C., Zhao H., Sun M.S., Chen J., Han C.H., Peng Q.L., Qiao H.H., Peng P., Liu Y., Luo S.D., Pan Y., 2019. Drosulfakinin signaling in fruitless circuitry antagonizes P1 neurons to regulate sexual arousal in Drosophila. Nat. Commun. 10(1), 4770.

Xiao A., Cheng Z., Kong L., Zhu Z., Lin S., Gao G., Zhang B., 2014. CasOT: a genome-wide Cas9/gRNA off-target searching tool. Bioinformatics. 30(8), 1180-1182.

Xu L., Jiang H.B., Yu J.L., Wang J.J., 2024. Plasticity of the olfactory behaviors in Bactrocera dorsalis under various physiological states and environmental conditions. Curr. Opin. Insect Sci. 63, 101196.

Yang Y., Jiang H.-B., Liang C.-H., Ma Y.-P., Dou W., Wang J.-J., 2023. Chromosome-level genome assembly reveals potential epigenetic mechanisms of the thermal tolerance in the oriental fruit fly, Bactrocera dorsalis. Int. J. Biol. Macromol. 225, 430-441.

Yu N., Smagghe G., 2014. CCK(-like) and receptors: structure and phylogeny in a comparative perspective. Gen. Comp. Endocrinol. 209, 74-81.

Zhang L.W., Guo X., Zhang W., 2022. Nutrients and pheromones promote insulin release to inhibit courtship drive. Sci. Adv. 8(10).

Zhao Z.H., Carey J.R., Li Z.H., 2024. The Global Epidemic of Bactrocera Pests: Mixed-Species Invasions and Risk Assessment. Annu. Rev. Entomol. 69(Volume 69, 2024), 219-237.

Zhu X., Li Q., Li J., Luo J., Chen W., Li X., 2018. Comparative Study of Volatile Compounds in the Fruit of Two Banana Cultivars at Different Ripening Stages. Molecules. 23(10).

Figures

Figure 1. Starvation alters the behavior and olfactory responses of *B. dorsalis*. (*A*) Schematic illustration of the experimental design. (*B*) Schematic illustration of the mating behavior assay device. (*C*) Representative foraging trajectories in the 100 mm diameter arenas within a 15-min observation period of flies starved for different durations. (*D*) Cumulative successful foraging within a 15-min observation period of male flies starved for different durations. Data are means ± SEM, n = 60 flies for each condition. Different letters above the error bars indicate significant differences (one-way ANOVA followed by Tukey's multiple comparisons test; $P <$ 0.05). (*E*–*F*) EAG responses to artificial food in fed and starved male (*E*) and female (*F*) flies (unpaired *t*-test). (*Top*) Representative EAG recordings. (*G*) Cumulative courtship rate within a 15-min observation period of flies starved for different durations (n = 78, 76, 78, 84 and 91, respectively, from fed to 12 h). (*H*) Cumulative copulation rate within a 15-min observation

period of the flies starved for different durations (n = 78, 76, 79, 83 and 93, respectively, from fed to 12 h). (*G–H*) Kruskal-Wallis and *post hoc* Mann-Whitney U tests were applied. (*I*–*J*) EAG responses to body extracts in fed and starved male (*I*) and female (*J*) flies. (*Top*) Representative EAG recordings. Data are means ± SEM, n = 10–12 antennae per genotype (unpaired *t*-test; *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$).

Figure 2. Starvation-induced expression of neuropeptide system components and phenotypes of null mutant flies. (*A*) Changes in the expression of neuropeptide system components in the antennae of starved flies. The relative expression values are the fold-changes compared to the control. Data are means ± SEM, $n = 3$ (unpaired *t*-test; $*P < 0.05$, $**P < 0.01$). (*B*) Altered amino acid sequence and mature peptides in the null mutants $sk^{-/-}$ and $skr1^{-/-}$ compared to wild-type (WT) flies. (*C*) Mean velocity of flies representing each genotype during a 15-min observation period. Data are means ± SEM (Kruskal–Wallis test). (*D*) Knockout of sulfakinin and SkR1 increases food consumption in *B. dorsalis* at different times of day. Each experiment consisted of one fly, and at least 15 flies were contained for the assay. Different lower-case

letters indicate significant differences between treatments (one-way ANOVA followed by

Tukey's multiple comparisons test; $P < 0.05$).

Figure 3. Sk-SkR1 signaling is required for changes in behavior and olfactory sensitivity induced by starvation. (*A□B*) Cumulative successful foraging within a 15-min observation period of fed and starved flies with different genotypes. The successful foraging of females (*A*) and males (*B*) was measured separately. Data are means ± SEM, *n* = 60 flies for each condition. Different letters above the error bars indicated significant differences (one-way ANOVA followed by Tukey's multiple comparisons test; *P* < 0.05). (*C*) Cumulative copulation rates within a 15-min observation period of fed and starved flies with different genotypes (*n* = 92, 86, 88, 106, 80 and 76, respectively, from fed WT to starved *skr1*⁻/⁻). (*D*) Cumulative courtship rates within a 15-min observation period of fed and starved flies with different genotypes (*n* = 91, 85, 86, 98, 80 and 77, respectively, from fed WT to starved *skr1*⁻/⁻). (*C–D*) Kruskal-Wallis and *post hoc* Mann-Whitney U tests were applied (**P* < 0.05, ***P* < 0.01, ****P* < 0.001). (*E□F*) EAG responses to artificial food in fed and starved flies of different genotypes. The EAG responses to artificial food in females (*E*) and males (*F*) were measured separately. (*Top*) Representative EAG recordings. (*G□H*) EAG responses to body extracts in fed and starved flies of different genotypes. The EAG responses to body extracts of females (*G*) and

males (*H*) were measured separately. (*Top*) Representative EAG recordings. Data are means ± SEM, n = 10–12 antennae for each condition. Different letters above the error bars indicated significant differences (one-way ANOVA followed by Tukey's multiple comparisons test; $P <$ 0.05).

Figure 4. The transition of olfactory responses is associated with the expression of different

sets of ORs in OR neurons induced by Sk-SkR1 signaling. (*A*) Principal component analysis

(PCA) using differentially expressed genes obtained from pairwise comparisons between

different treatments. (*B*) The expression profiles of the candidate OR genes in wild-type (WT)

and $sk^{-/-}$ flies determined by qRT-PCR. The OR names with red fonts represent the set of ORs

for food odor and blue fronts represent the set of ORs for sex pheromone. Data are mean

relative expression levels ± SEM. R1–R3 represent biological replicates. Different lower-case

letters indicate significant differences between treatments (one-way ANOVA followed by

Tukey's multiple comparisons test; $P < 0.05$). (*C*) Quantification of calcium levels following the

response of candidate ORs to sex pheromones and food odors at a concentration of 10^{-4} M (*n*

= 3⎵6). Different lower-case letters indicate significant differences between treatments

(one-way ANOVA followed by Tukey's multiple comparisons test; $P < 0.05$). (*D*)

Dose–response curves of candidate ORs to sex pheromone components such as ethyl laurate

(EL), ethyl palmitate (EP), 2,3,5-trimethyl pyrazine (TMP), and 2,3,5,6-tetramethylpyrazine

(TTMP). (*E*) Dose–response curves of candidate ORs to food odorants such as ethyl benzoate

(EBE), ethyl butyrate (EBU), diethyl maleate (DM), and methyl eugenol (ME). In both cases, *n*

= 5. (*F*) Co-localization of Orco (red) and SkR1 (green) neurons in *B. dorsalis* antennae. Scale

bars = 50 μm.

Figure 5. Schematic representation of peripheral olfactory remodeling in OR neurons of the antenna that arbitrate between mating and foraging behavior in *B. dorsalis*. Starvation increases the abundance of SkR1 in the OR neurons (red arrows) and SkR1 signaling induces the expression of genes encoding ORs that sense food odors, resulting in successful foraging. Satiation suppresses *SkR1* expression and induces the expression of genes encoding ORs that sense opposite-sex pheromones, leading to successful mating (blue arrows). Dashed arrows represent the additional possible pathways that have not been tested in this study, but not excluded in the model. Please see the discussion for details of additional possible factors modulating odorant sensitivity relevant to satiety.