1

Secreted chemokines reveal diverse inflammatory and degenerative processes in the

intervertebral disc of the STZ-HFD mouse model of Type 2 diabetes

Christian E. Gonzalez[1], Rachana S. Vaidya[4], Sade W. Clayton[4], Simon Y. Tang[1-4]

[1]Department of Biomedical Engineering, Washington University in St. Louis, St. Louis, MO
[2]Institute of Material Science and Engineering, Washington University in St. Louis, St. Louis, MO
[3]Department of Mechanical Engineering and Materials Science, Washington University in St. Louis, St. Louis, MO
[4]Department of Orthopaedic Surgery, Washington University School of Medicine, St. Louis, MO

Corresponding Author:
Simon Y. Tang, Ph.D., MSCI
Associate Professor
Orthopedic Surgery, Biomedical Engineering,
Mechanical Engineering & Materials Science
Washington University in St. Louis
simon.tang@wustl.edu

Keywords: type 2 diabetes; intervertebral disc degeneration; streptozotocin-high-fat-diet; leptin receptor

deficiency; chronic inflammatory cytokines

Abstract

The chronic inflammation present in type 2 diabetes causes many chronic inflammatory comorbidities,

including cardiovascular, renal, and neuropathic complications. Type 2 diabetes is also associated with a

number of spinal pathologies, including intervertebral disc (IVD) degeneration and chronic neck and back

pain. Although confounding factors such as obesity are thought to increase the loads to the

musculoskeletal system and subsequent degeneration, studies have shown that even after adjusting age,

body mass index, and genetics (e.g. twins), patients with diabetes suffer from disproportionately more IVD

degeneration and back pain. Yet the tissue-specific responses of the IVD during diabetes remains

relatively unknown. We hypothesize that chronic diabetes fosters a proinflammatory microenvironment

within the IVD that accelerates degeneration and increases susceptibility to painful disorders. To test this

hypothesis, we evaluated two commonly used mouse models of diabetes – the leptin-receptor deficient

mouse (db/db) and the chronic high-fat diet in mice with impaired beta-cell function (STZ-HFD). The

db/db is a genetic model that spontaneous develop diabetes through hyperphagia, while the STZ-HFD

mouse first exhibits rapid obesity development under HFD and pronounced insulin resistance following

streptozotocin administration. Both animal models were allowed to develop sustained diabetes for at

least twelve weeks, as defined by elevated hemoglobin A1C, hyperglycemia, and glucose intolerance.

Following the twelve-week period, the IVDs were extracted in quantified in several measures including

tissue-specific secreted cytokines, viscoelastic mechanical behavior, structural composition, and

histopathologic degeneration. Although there were no differences in mechanical function or the overall

structure of the IVD, the STZ-HFD IVDs were more degenerated. More notably, the STZ-HFD model

shows a significantly higher fold increase for eight cytokines: CXCL2, CCL2, CCL3, CCL4, CCL12

(monocyte/macrophage associated), IL-2, CXCL9 (T-cell associated), and CCL5 (pleiotropic). Correlative

network analyses revealed that the expression of cytokines differentially regulated between the db/db and

the STZ-HFD models. Moreover, the STZ-HFD contained a fragmented and modular cytokine network,

indicating greater complexities in the regulatory network. Taken together, the STZ-HFD model of type 2

diabetes may better recapitulate the complexities of the chronic inflammatory processes in the IVD during

diabetes.

Introduction

{Figure 1}

Type 2 diabetes (T2D) is a prevalent metabolic disorder marked by insulin resistance and prolonged hyperglycemia, impacting millions around the globe and leading to significant healthcare expenses (Srinivasan and Ramarao, 2007; Boucher et al., 2014; Petersen and Shulman, 2018; United States, Center for Disease Control, 2022). This disease shares several characteristics with autoimmune disorders, including the chronic, systemic overexpression of immunomodulating cytokines, which can gradually lead to widespread accrual of tissue damage across multiple organ systems (Itariu and Stulnig, 2014; Chen et al., 2017; de Candia et al., 2019; Daryabor et al., 2020). Intervertebral disc (IVD) degeneration is a comorbidity of particular interest due to strong evidence for chronic-inflammatory etiology (Risbud and Shapiro, 2014; Molinos et al., 2015; Navone et al., 2017; Lyu et al., 2021; Pinto et al., 2023) and low back pain is a potently association with chronic T2D (Robinson et al., 1998; Jhawar et al., 2006; Sakellaridis, 2006; Liu et al., 2018; Alpantaki et al., 2019; Cannata et al., 2019; Broz et al., 2021). In T2D, chronic inflammation driven by a persistent milieu of chemokines may foster a pro-degenerative microenvironment within the IVD, potentially linking T2D-induced inflammation to accelerated IVD degeneration.

Various animal models are used to study T2D, primarily being rodent models due to their cost-effectiveness, ease of handling, and genetic similarities to humans. Models can be classified into genetic (spontaneously induced) and non-genetic (experimentally induced) types. Typical genetic models involve animal strains with inherent mutations predisposing the animal to developing T2D, while experimentally-induced models typically employ some combination of chemical induction, dietary manipulations, and surgical methods (Srinivasan and Ramarao, 2007; Islam and Wilson, 2012). Each model mimics different aspects of T2D pathogenesis, such as insulin resistance and beta-cell dysfunction, but none can entirely replicate the human condition. The Lepr^{db} (db/db) mouse, a particularly popular model for studying T2D, is characterized by a point mutation in the *db* gene, leading to the inactivation of the leptin receptor [**Fig. 1A**] (Chen et al., 1996; Lee et al., 1996). This model exhibits metabolic characteristics similar to human T2D, including severe obesity, hyperglycemia, insulin resistance, and hyperinsulinemia, making it a valuable model for studying T2D (Wang et al., 2014). The etiology of T2D in db/db mice differs from human T2D,

though, as it is driven by leptin receptor deficiency rather than a combination of genetic and lifestyle

factors commonly seen in humans. This distinction complicates the direct translation of findings from this

model to human T2D (Wang et al., 2014).

The db/db mouse model has been widely employed in spine research to further investigate the

consequences of T2D on spinal complications. Studies focusing on the IVDs of db/db mice have shown

that these mice exhibit signs of disc degeneration, including increased cell apoptosis and extracellular

matrix degradation (Li et al., 2020; Natelson et al., 2020; Lintz et al., 2022). These findings suggest that

the db/db model can be useful for studying T2D-induced IVD degeneration. However, the absence of

functional leptin signaling complicates interpretations of these changes as it may independently affect IVD

homeostasis. Leptin, beyond its well-known role in energy regulation and appetite control, influences

various systemic functions, including immune response modulation, bone formation, and reproductive

health (Francisco et al., 2018). In the musculoskeletal system, leptin affects chondrocyte proliferation,

osteoblast activity, the synthesis of extracellular matrix proteins, and many more tissue- and cell-level

functions crucial for maintaining homeostasis (Gruber et al., 2007; Li et al., 2013; Han et al., 2018;

Sharma, 2018; Segar et al., 2019; Curic, 2021). Leptin's responsibilities in the IVD specifically include

promoting anabolic processes and reducing catabolic activities, thereby supporting disc health (Li et al.,

2013; Han et al., 2018). Consequently, the lack of leptin signaling in db/db mice might mask or alter the

degenerative pathways activated by the diabetic condition, potentially confounding observations of the

diabetic milieu with those of leptin ablation. This complicates the interpretation of T2D-related changes in

the IVD observed in this model, as the effects could be due to the absence of leptin signaling rather than

diabetes alone.

In contrast, the Streptozotocin-High Fat Diet (STZ-HFD) model offers a non-genic approach to

replicating T2D. It induces the condition through pro-glycemic diet and low-dose streptozotocin-induced

pancreatic beta-cell dysfunction, avoiding genetic ablation of hormonal pathways like leptin (as seen in

the db/db model) [**Fig. 1A**]. This model is characterized by significant metabolic disturbances to glycemic

status, insulin resistance, and body weight, mirroring the human T2D phenotype more closely (Kusakabe

et al., 2009; Islam and Wilson, 2012). Additional metabolic characteristics reported in STZ-HFD mouse

studies include elevated serum insulin levels, dyslipidemia (increased triglycerides, LDL cholesterol, and

total cholesterol), and increased markers of inflammation and oxidative stress (Gilbert et al., 2011; Alquier

and Poitout, 2018; Yin et al., 2020). This sets this mouse model apart as valuable tool for studying the

complex interactions within diabetic complications without the confounding factor of complete leptin

signaling ablation (Kusakabe et al., 2009). Previously this model has been employed in studying diabetic

complications in bone (Eckhardt et al., 2020). Our study aims to uncover the mechanisms behind

inflammatory-pathway contribution to IVD degeneration and dysfunction, advancing the field's

understanding of T2D-related IVD complications.

Inflammatory cytokines are central to the pathophysiology of both IVD degeneration and T2D.

They not only mediate acute inflammatory responses but also perpetuate chronic inflammation, leading to

tissue degradation (Guest et al., 2008; Velikova et al., 2021). Key proteins such as ADAMTs and MMPs,

regulated by cytokines, are instrumental in the degradation of extracellular matrix in the IVD, furthering

degeneration (Bond et al., 1998; Malemud, 2019). The identification and characterization of these

cytokines in the context of T2D can elucidate potential therapeutic targets and markers for early

intervention (Al-Shukaili et al., 2013; Herder et al., 2013).

Studies have shown that inflammatory cytokines are involved in the degradation of IVD tissue,

contributing to conditions like pain and disc degeneration (Shamji et al., 2010; Risbud and Shapiro, 2014).

Research indicates that pro-inflammatory cytokines, such as TNF-α, IL-1β, and IL-6, are upregulated in

degenerated and herniated disc tissues, perpetuating the inflammatory response and leading to further

tissue damage (Wuertz and Haglund, 2013; Molinos et al., 2015; Navone et al., 2017; De Geer, 2018).

Moreover, these cytokines influence the expression of matrix-degrading enzymes like ADAMTS-4 and

MMP-9, which are crucial for the breakdown of extracellular matrix components, exacerbating disc

degeneration (Tian et al., 2013; Zhang et al., 2015).

The role of inflammatory cytokines in T2D is also well-documented. They contribute to insulin

resistance and beta-cell dysfunction, key features of T2D (Calle and Fernandez, 2012). Studies have

identified elevated levels of pro-inflammatory cytokines in T2D patients, suggesting their role in the

disease's progression and complications (Guest et al., 2008; Velikova et al., 2021). Targeting these

cytokines could provide novel therapeutic approaches for both T2D and IVD degeneration, highlighting

the interconnected nature of these conditions (Al-Shukaili et al., 2013; Herder et al., 2013).

This study seeks to shift the research focus towards using the STZ-HFD model by comparing the

overall IVD health between the db/db and STZ-HFD mouse models, with a particular focus on

inflammation and related complications. We will examine inflammation and broad genomic changes in

functional spine units to understand the molecular mechanisms at play. Additionally, degeneration and

morphological changes will be assessed to determine the extent of tissue damage and structural

alterations. Finally, the mechanical function of the intervertebral discs will be evaluated to understand the

impact of T2D. By investigating these comprehensive metrics [**Figure 1C**], we aim to dissect the specific

pathways through which T2D exacerbates IVD degeneration. Our findings will enhance the understanding

of the interplay between T2D and IVD homeostasis and validate the relevance of the STZ-HFD model.

Materials & Methods

Animals. In this study, we focused on skeletally-mature (12-week-old) male C57BL/6 mice (N = 20) for

their established susceptibility to Type 2 Diabetes (T2D) when exposed to a high-fat diet (HFD) and

treated with streptozotocin (STZ). Previous findings have indicated that this strain, particularly males,

exhibits rapid obesity development under HFD and pronounced insulin resistance following STZ

administration (Luo et al., 1998; Mu et al., 2006; Mu et al., 2009). As a result of estrogen-mediated

mechanisms of protection, female STZ-HFD mice are resistant to developing T2D and are thus excluded

from this study (Medrikova et al., 2012; Pettersson et al., 2012; Stubbins et al., 2012). To contrast the

STZ-HFD model's pathology with a well-characterized model of chronic T2D, we included a parallel

cohort of 3-month-old homozygous (db/db) male Lepr[db] mutant mice with heterozygous (db/+) littermate

controls (N = 18), as at this timepoint they have endured a similar duration of T2D symptoms; male mice

were used to avoid confounding effects of sex-based differences. All mice were group-housed (max. 5

mice per cage) under pathogen-free conditions in standard cages; the environment was controlled with a

stable temperature and a 12-hour light/dark cycle, with *ad libitum* access to food and water. All

procedures were approved by the Institutional Animal Care and Use Committee (IACUC) of Washington

University in St. Louis. Regular health and welfare assessments were conducted, including general

monitoring of weight, food supply, and behavior.

Study Design. The study was organized into two phases: initially, mice were maintained on a HFD for four

to six weeks, after which they received a one-time dose of STZ and continued on the HFD for an

additional 12 weeks. The STZ-HFD group (n = 13) was given a high-fat diet (Research Diets, Inc.,

D12492i, 60% kcal from fat) for the duration of the study, with the Control + Vehicle group (n = 7)

receiving standard mouse chow (5053 PicoLab® Rodent Diet 20, 13% kcal from fat). At the end of the

initial phase, baseline measurements of body weight and fasting blood glucose were collected. Following

the first phase, STZ-HFD mice were injected intraperitoneally with 100 mg/kg Streptozotocin

(MilliporeSigma) in 50 mM sodium citrate buffer (pH 4.5), with Con+Veh animals receiving the sodium

citrate buffer only. The two experimental groups received their respective diets for 12 weeks following the

injection (experimental phase), and animals were assessed for diabetic status via glucose tolerance.

Finally, animals were euthanized, and sterile coccygeal functional spine units (FSUs) including IVDs were

harvested from each animal for organ culture and terminal measurements. The overall study design is

outlined in the schematic in **Fig. 1B** and **Fig. 1C**.

Measures of diabetic status. As previously mentioned, several measures of diabetic status were collected

at various points throughout the experimental phase, including fasting blood glucose, % A1C, and glucose

tolerance. Blood glucose levels (mg/dL) were measured using a glucometer (GLUCOCARD Vital$^{®}$ Blood

Glucose Meter). Blood samples were drawn via superficial incision to the tail tip of fasted mice using a

scalpel; the tail was immediately treated with analgesic (Kwik Stop® Styptic Powder) after blood

collection. The percentage of glycated hemoglobin (% A1C) was measured using the A1CNow®+ system

(PTS Diagnostics) according to kit instructions. Blood samples were drawn in the same way as during the

blood glucose test. Finally, to assess glucose tolerance (oral glucose tolerance test, oGTT), mice were

fasted and had their blood glucose measured as described above to establish a baseline. Mice were then

injected intraperitoneally with 2 g/kg glucose in sterile water. Additional blood glucose measurements

were taken at 30 min., 60 min., and 90 min. post injection. The area-under-the-curve of blood glucose

(mg•h/dL) was calculated over the course of the test. For the purpose of evaluating diabetic status for

inclusion in the study, a cutoff of 435 mg•h/dL on the GTT, established in previous studies on human T2D

criteria (Sakaguchi et al., 2015), was adjusted for time and interspecies differences in blood glucose

levels.

Organ culture. Following extraction, FSUs were cultured in 2 mL Dulbecco's Modified Eagle

Medium/Nutrient Mixture F-12 Ham with L-glutamine and 15 mM HEPES (Sigma-Aldrich, D6421) .

Culture medium was supplemented with 20% fetal bovine serum (Gibco No. A5256801) and 1% penicillin-

streptomycin (Gibco No. 15140122). Cultures underwent a preconditioning period of 7 days with regular

media changes to account for the inflammatory response from extraction. Conditioned media was

collected 48 hours after the final media change at the end of the preconditioning period and immediately

frozen in -80° C.

Chemokine assay. A multiplex assay of remodeling factors and inflammatory chemokines (45-Plex Mouse

Cytokine Discovery Assay, Eve Technologies Assays; CCL-2,3,4,5,11,12,17,20,21,22; CSF-1,2,3; IL-

1α,1β,2,3,4,5,6,7,9,10,11,12A,12B,13,15,16,17; CXCL1,2,5,9,10; CX3CL1; IFN-γ,β1; TNFα; LIF; VEGF;

EPO; TIMP-1) was performed on conditioned media samples. Cytokine levels were analyzed using

Welch's t-Test. Cytokines with greater than 25% missingness across all experimental groups were

excluded from further analysis. Significantly upregulated cytokines in each model were selected for a

secondary fold change analysis, where protein expression levels for db/db and STZ-HFD mice were used

to calculate fold change for each cytokine over the corresponding average expression of the control (db/+

and Con+Veh respectively).

Cytokine Interaction Network Construction and Analysis. To further investigate the inflammatory profiles of

these T2D models, networks of cytokine interactions were constructed and analyzed using a custom

MATLAB (Version: 9.13.0.2080170 R2022b) script. Networks were generated by calculating a Pearson

correlation matrix for each experimental group based on cytokine expression data from the multiplex

panel of conditioned media. To isolate strong protein correlations, a threshold ($|r| > 0.7$) was applied to the

correlation matrices. The filtered matrices were used to create undirected graphs, with nodes

representing cytokines and edges representing significant interactions. Centrality measures were

calculated to determine the importance of each cytokine within the networks. Eigenvector centrality and

betweenness centrality were computed for each network using the centrality function. The resulting

centrality values were organized into tables and sorted to identify the top-ranking cytokines. For each

centrality metric, shared high-ranking cytokines between the diabetic models and unique cytokines for

each diabetic model were aggregated. Additionally, key network characteristics were extracted to

understand the structure and function of the cytokine networks. The average path length was determined

using the distances function to compute the shortest finite paths between all pairs of nodes. Modularity

and community structure were assessed using the Louvain community structure and modularity algorithm (Blondel et al., 2008). We also computed k-hop reachability to assess the extent to which cytokines can influence each other within one (k=1) or two (k=2) steps. The Jaccard index was used to compare the reachability matrices between different groups, providing a measure of similarity. Finally, the networks were visualized using force-directed layouts with nodes colored by eigenvector centrality and sized by betweenness centrality.

Histology. Following removal from culture, FSUs were fixed in 10% neutral-buffered formalin (Epredia™ 5735) overnight and decalcified in ImmunoCal (StatLab STL14141) for 72 hours. Samples were fixed in paraffin, sectioned in the sagittal plane at 10 μm thickness, and stained with Safranin-O/Fast Green prior to being imaged via Hamamatsu NanoZoomer with a 20x objective. Blinded histological images of the IVDs were evaluated for degeneration based on a standardized histopathological scoring system (Melgoza et al., 2021).

Contrast-enhanced Micro-computed Tomography. To prepare for imaging, functional spine units were incubated in a 175 mg/mL Ioversol solution (OptiRay 350; Guerbet, St. Louis) diluted in PBS at 37 °C. Following 4 hours of incubation, the samples underwent scanning with a Viva CT40 (Scanco Medical) at a 10-μm voxel size, using 45 kVp, 177 μA, high resolution, and a 300 ms integration time. CEμCT data was exported as a DICOM file for analysis in a custom MATLAB program (https://github.com/WashUMusculoskeletalCore/Washington-University-Musculoskeletal-Image-Analyses). After an initial Gaussian filter (kernel size = 5), functional spine units were segmented by drawing a contour around the perimeter of the IVD every 10 transverse slices and morphing using linear interpolation. This was defined as the whole disc mask. The NP was segmented from the whole disc by thresholding and performing morphological close and morphological open operations to fill interior holes and smooth NP boundaries. The volumes and intensities were calculated from the NP and whole disc regions. Disc height index (DHI) was measured by averaging the height-to-width ratio of the IVD over five slices in the mid-sagittal plane. Finally, the NP intensity/disc intensity (NI/DI) ratio and NP volume fraction (NPVF; NP volume / total volume) was computed using the intensity and volume metrics reported by the output analysis within the MATLAB program. All thresholding and analysis were performed in a blinded fashion. This approach is described in further detail in prior studies (Lin et al., 2016; Lin and Tang, 2017).

Mechanical Testing. Mechanical testing of functional spine units was performed using cyclic compression

on a microindentation system (BioDent; Active Life Scientific) with a 2.39 mm probe as previously

described (Liu et al., 2015). Samples were adhered to an aluminum plate and placed in a PBS bath prior

to aligning the sample beneath the probe with a 0.03 N preload. Each unit was then sinusoidally loaded in

compression at 1 Hz for 20 cycles with a 35 µm amplitude. A loading slope value was calculated from the

linear region of the force-displacement curve, and the tan delta (loss tangent) was calculated from the

phase delay between loading and displacement. This approach is described in further detail in prior

studies.

Matrix Protein Assays. Whole extracted discs were used to measure the biochemical content of various

matrix proteins. First, discs were digested overnight in a papain digestion buffer, after which the buffer

was collected for a 1,9-dimethylmethylene blue assay of sulfated glycosaminoglycan content with a

chondroitin sulfate standard (Liu et al., 2017). The disc was then subjected to high temperature bulk

hydrolyzation in 12 N HCl. Hydrolysates were desiccated and reconstituted with 0.1x phosphate-buffered

saline (PBS) and measured against a quinine standard for advanced glycation end product content (Liu et

al., 2017). Finally, a hydroxyproline assay was used to quantify collagen content as previously described

(Liu et al., 2017).

Results

{Figure 2}

The db/db and STZ-HFD Mouse Models Exhibit a Characteristically Diabetic Phenotype

Both db/db and STZ-HFD mice demonstrate hallmark features of diabetes, as evidenced by the

results in **Figure 2**. Both groups exhibit AUC GTT values above the defined threshold, indicating impaired

glucose tolerance [**Fig. 2A**]. The comparison between Con+Veh and STZ-HFD groups is highly significant

(p < 0.0001), indicating particularly severe glucose intolerance in the STZ-HFD model. The terminal body

weights show that db/db mice weigh significantly more than both db/+ and STZ-HFD mice, underscoring

the severe obesity associated with the db/db model [**Fig. 2B**]. This highlights the db/db mice as a

representative model of a more severely obese type 2 diabetic phenotype than the STZ-HFD model. The

terminal fasting blood glucose levels indicate a significant difference only between the STZ-HFD and

Con+Veh groups, demonstrating notable fasting hyperglycemia in the STZ-HFD model [**Fig. 2C**]. HbA1c

levels reveal no difference between db/db mice and STZ-HFD mice, but each group significantly differs

from their respective controls [**Fig. 2D**]. This indicates that both models exhibit chronic hyperglycemia,

confirming their relevance as models of T2D.

{Figure 3}

Histopathological Analysis Reveals IVD Degeneration in STZ-HFD Mice

The STZ-HFD models exhibited a more degenerative phenotype in a histopathological analysis of

both models. **Figure 3** presents a comparison of histological phenotypes across the four groups (db/+,

db/db, Con+Veh, and STZ-HFD), showing the range of histopathological scores within this study [**Fig.**

3A]. Histopathological scoring revealed that only the STZ-HFD mice exhibit significantly greater IVD

degeneration compared to the control group [**Fig. 3B**]. **Figure 3C** highlights common degenerative

characteristics in the IVD of STZ-HFD mice. In the Con+Veh samples, the annulus fibrosus has healthy,

convexed outer lamellae (➡) and well-organized, concentric inner lamellae (▼). In contrast, the STZ-HFD

samples show degenerate crimped, concave outer lamellae (⇨) and wavy, disorganized inner lamellae

(▽), indicating altered matrix structure in the AF.

{Figure 4}

CT Analysis, Matrix Assays, and Mechanical Testing All Show No Major Effects in IVD Physiology

The analysis of CEμCT data, mechanical testing, and matrix protein assays revealed only one significant

difference among the four groups across all nine measured outcomes. Specifically, parameters such as

NI/DI, NPVF, loading slope, hysteresis energy, tan delta, and biochemical content showed no variations

between groups. The only significant result was a difference in morphology between the db/db and STZ-

HFD groups, as indicated by the DHI. These findings suggest that the structural integrity, mechanical

behavior, and biochemical composition of the IVDs are mostly consistent across the db/db and STZ-HFD

models. The lack of significant differences broadly implies that, under the conditions tested, type 2

diabetes does not markedly affect IVD properties in these models, and cannot further establish the

degenerate features gleaned in the histopathological analysis. This highlights the need for more sensitive

measures or longer study durations to detect subtle changes related to T2D.

{Figure 5}

Cytokines Representing a Plethora of Immune Functions Are Chronically Upregulated STZ-HFD IVDs

The comparative analysis of cytokine expression levels between db/db and STZ-HFD models

provides novel insights into diabetic inflammation of the IVD. The initial comparative analysis of the two

models' protein expression levels revealed that two cytokines (CCL2,3) and sixteen cytokines

(CCL2,3,4,5,12; CXCL1,2,9,10; CX3CL1; IL-2,6,16; CSF-3; VEGF; LIF) were upregulated in the db/db

and STZ-HFD models respectively, and thus included in the fold change analysis. In **Fig. 4A**, the fold

change in cytokine levels is computed over their respective controls (db/+ for db/db, Con+Veh for STZ-

HFD). The STZ-HFD model exhibited a significantly higher fold increase compared to the db/db model for

8 cytokines: CXCL2, CCL2, CCL3, CCL4, CCL12 (monocyte/macrophage associated cytokines) (Sagar et

al., 2012; Arendt et al., 2013; Lança et al., 2013; Etna et al., 2014; Motwani and Gilroy, 2015; He et al.,

2016; Lim et al., 2016; DeLeon-Pennell et al., 2017; Ruytinx et al., 2018; Sindhu et al., 2019; Zhang et al.,

2019, 38; Huang et al., 2020; Pelisch et al., 2020; Yang et al., 2020; Xu et al., 2021a; Xu et al., 2021b;

Sheng et al., 2022), IL-2, CXCL9 (T-cell associated cytokines) (Chang and Radbruch, 2007; Venetz et al.,

2010; Shachar and Karin, 2013; Ochiai et al., 2015; Boff et al., 2018; Kuo et al., 2018; Mortara et al.,

2018; House et al., 2020; Marcovecchio et al., 2021; Markovics et al., 2022), and CCL5 (pleiotropic

cytokine) (Juhas et al., 2015; Atri et al., 2018; Kranjc et al., 2019; Chen et al., 2020; Zeng et al., 2022).

Figure 4B shows a Venn diagram illustrating the cytokine expression profiles. The outer circle represents

the STZ-HFD model, encompassing a large number of upregulated cytokines. The inner circle represents

the db/db model, containing only the two upregulated cytokines, both of which are also upregulated in the

STZ-HFD model. This indicates that the STZ-HFD model has a broader and more pronounced cytokine

response compared to the db/db model.

{Figure 6}

Differential Network Structures Reveal Unique Inflammatory Pathways of T2D in IVD

Our analysis of cytokine networks in these two models revealed several key insights into the

inflammatory pathways associated with T2D. CCL2 and CCL4 emerged as pivotal cytokines in both

models based on betweenness centrality, highlighting their crucial roles in maintaining network

connectivity (**Error! Reference source not found.**). In the db/db mouse model, unique cytokines were

identified that are likely influenced by the absence of leptin signaling. These included CSF-3

(betweenness centrality), CXCL-5, CXCL-9, CXCL-10, and IL-4 (eigenvector centrality), and IL-11 (both

centralities) (**Error! Reference source not found.**). In contrast, the STZ-HFD mouse model, which

maintains functional leptin signaling and mimics human T2D etiology, presented a different profile of

unique cytokines. CXCL2 and IL-6 (betweenness centrality), and IL-16, CCL11, and CSF3 (eigenvector

centrality) were central in this model (**Error! Reference source not found.**). This indicates that these

cytokines may be independently regulated through pathways that are either bypassed or inhibited in the

simultaneous presence of leptin signaling and T2D-associated chronic inflammation. The db/db model

demonstrated a shorter average path length compared to the STZ-HFD model (**Error! Reference source**

not found.), indicating more efficient communication within the cytokine network of the db/db model. The

longer path length in the STZ-HFD model suggests a more dispersed network structure, consistent with

the wide array of upregulated cytokines across multiple signaling cascades. Additionally, the STZ-HFD

network had the highest modularity (**Error! Reference source not found.**). Higher modularity in the STZ-

HFD model indicates well-defined communities within the cytokine network, reflecting distinct functional

signaling pathways in a broad inflammatory response. The lower modularity in the db/db model suggests

a less distinct community structure and a more generic inflammatory response. The Jaccard index values

for the STZ-HFD and WT comparisons were 0.144 (k=1) and 0.246 (k=2), indicating low similarity

between these networks. Comparisons between db/db and db/+ showed slightly higher values, indicating

closer similarity (**Error! Reference source not found.**). This suggests significant changes in the cytokine

networks of the STZ-HFD model, while the db/db model retains more similarity to its control, reflecting

less drastic alterations to the inflammatory signaling cascade.

Discussion

This study offers novel insights into the chronic inflammatory profiles and degeneration in the IVD of

murine T2D models, specifically comparing the db/db and STZ-HFD mouse models. Our findings reveal

critical differences between these models, underscoring the heightened relevance of the STZ-HFD model

for studying T2D-related complications and inflammation in the IVD. Both db/db and STZ-HFD mice

exhibit classic diabetic traits, specifically glucose intolerance and chronic hyperglycemia. These

phenotypic characteristics align with established literature on human T2D (Dalgaard and Pedersen, 2001;

Cefalu, 2006; Lin and Sun, 2010), validating the use of these models in diabetes research. However,

while both models mimic key aspects of T2D, the STZ-HFD model parallels the etiology of human T2D

(severe insulin resistance and pancreatic beta-cell dysfunction) more closely (Gilbert et al., 2011; Stott

and Marino, 2020).

In terms of IVD degeneration, our study demonstrates that STZ-HFD mice exhibit significantly

more IVD degeneration compared to controls, indicating a notable degenerative phenotype. As evidenced

by the structural integrity, mechanical behavior, and biochemical composition of the IVDs in these two

models, the degenerative features remain primarily encapsulated by the histopathological analysis and

are not further elucidated by these measures. In contrast, db/db mice, although characterized by severe

obesity, insulin resistance, and chronic hyperglycemia, show less severe IVD degeneration compared to

littermate controls. This fits within the broader picture of existing literature wherein db/db mice develop

moderate IVD degeneration, influenced by factors such as sex and specific metabolic disruptions. Studies

by Lintz et al. (2022), Natelson et al. (2020), and Li et al. (2020) support these observations, noting that

while db/db mice do exhibit IVD degeneration, its typically mild or nuanced in features.

Our novel findings on the inflammatory profiles of these models serve as the strongest evidence

to the superiority of the STZ-HFD model in mimicking clinically relevant T2D. The STZ-HFD model shows

a significantly more robust inflammatory response, upregulating 16 different cytokines, including those

associated with monocyte and macrophage recruitment, T-cell activation, and pleiotropic immune cell

effects. This extensive cytokine upregulation indicates a more complex inflammatory response, which

may contribute to the IVD degeneration observed in the STZ-HFD model. The relevance of these findings

is underscored by literature examining the inflammatory profile of human T2D. CXCL2 is elevated in T2D

patients (Rebuffat et al., 2018; Pan et al., 2021). Similarly, CCL2 and CCL3 are found at higher levels in

diabetic conditions (Neumeier et al., 2011; Arner et al., 2012; Sullivan et al., 2013; Sindhu et al., 2017;

Chang et al., 2021; Pan et al., 2021). The presence of CXCL9 and IL-2, aligns with findings in diabetic

patients, highlighting the potential role of T-cell-mediated inflammation in the T2D IVD (Higurashi et al.,

2009; Nawaz et al., 2013; Pan et al., 2021; Suri et al., 2022). CCL4 is also elevated in human studies

where this cytokine are implicated in chronic inflammation in T2D (Chang et al., 2021; Pan et al., 2021;

Mir et al., 2024). Moreover, the upregulation of the pleiotropic cytokine CCL5 in the STZ-HFD model, a

significant marker in human T2D inflammation, further underscores the physiological relevance of this

model (Keophiphath et al., 2010; Pettigrew et al., 2010; Inayat et al., 2019; Chang et al., 2021; Pan et al.,

2021; Alshammary et al., 2023; Mir et al., 2024). In summary, the STZ-HFD mouse model demonstrates a

more comprehensive inflammatory response, akin to the inflammatory profile observed in human T2D,

making it a more physiologically relevant model for studying diabetic inflammation in the IVD. The db/db

model, while still relevant, exhibits a more limited cytokine profile. This comprehensive inflammatory

characterization of the STZ-HFD model provides a valuable framework for future research into T2D-

associated IVD degeneration and related pathologies.

Our network analysis further identified the STZ-HFD IVD as having a distinct inflammatory profile,

revealing unique cytokine interactions and regulatory mechanisms. In the db/db model, the absence of

leptin signaling results in compensatory network configurations wherein cytokines like CSF-3, CXCL-5,

CXCL-9, CXCL-10, IL-4, and IL-11 are more responsible for network function. Leptin's regulatory effects

on the immune system and inflammation are crucial (Francisco et al., 2018); without them, the

inflammatory response to chronic stimuli from untreated T2D is stifled and incomplete. This is consistent

with the db/db network, which is indicative of a constrained inflammatory response, as revealed by the

short average path length and high similarity to control networks. Conversely, the STZ-HFD model, which

maintains leptin signaling and mirrors human T2D etiology, exhibits extensive cytokine upregulation. As

demonstrated by the cytokine network modeling, CXCL2, IL-6, IL-16, CCL11, and CSF3 are crucial

mediators of this response. CXCL2, with both high centrality and significantly upregulated expression,

may serve as a potential therapeutic target for inhibiting diabetic inflammation in the IVD. The presence of

leptin signaling supports a more comprehensive and complex inflammatory response, reflecting the

systemic impact of obesity and insulin resistance on the immune system (Francisco et al., 2018; Rebuffat

et al., 2018; Sharma, 2018; Segar et al., 2019; Pan et al., 2021). The STZ-HFD network has higher

modularity and longer average path length, suggesting a dispersed network with distinct functional

modules. This indicates a collective of specific pathways driving the broader inflammatory response,

aligning with literature above that suggests obesity and diabetes regulate a broad range of pathways.

Despite these significant findings, this study has several limitations. The selection of cytokines

examined was relatively small, potentially missing other important inflammatory mediators involved in IVD

degeneration. Additionally, the mechanistic link between inflammation and degeneration remains unclear

and warrants further investigation. While the STZ-HFD model provides a comprehensive inflammatory

profile, the specific pathways driving the observed IVD degeneration need to be elucidated through future

studies. Future directions for research based on our findings include expanding the panel of cytokines

and other inflammatory mediators examined in the STZ-HFD model to gain a more complete

understanding of the inflammatory landscape in T2D. Investigating the specific molecular and cellular

mechanisms linking inflammation to IVD degeneration will be crucial to furthering future therapeutic

approaches. Finally, exploring therapeutic interventions targeting the identified cytokine pathways could

provide insights into potential treatments for T2D-related IVD degeneration.

In conclusion, our findings establish the STZ-HFD model as a more physiologically relevant

model for studying T2D-related inflammation and IVD degeneration. The extensive cytokine upregulation

and significant degenerative phenotype observed in this model provide a valuable framework for future

research into T2D-associated pathologies. The db/db model, while still relevant, exhibits a more stunted

cytokine profile and limited IVD degeneration, making it less representative of the chronic inflammatory

environment seen in human T2D. These insights enhance our understanding of T2D-induced IVD

degeneration and identify key cytokine pathways for therapeutic development, emphasizing the need to

address these pathways holistically for effective intervention.

Conflicts of Interest

Authors declare no conflict of interest.

Acknowledgements

This work was supported by the NIH R01AR074441 and P30AR074992. Thank you to the Alafi Neuroimaging Lab for

Nanozoomer Access (supported by NIH S10RR027552). Multiplex cytokine panels were performed by Eve

Technologies Corp.

References

Alpantaki, K., Kampouroglou, A., Koutserimpas, C., Effraimidis, G. and Hadjipavlou, A. (2019).
Diabetes mellitus as a risk factor for intervertebral disc degeneration: a critical review. *Eur. Spine
J.* **28**, 2129–2144.

Alquier, T. and Poitout, V. (2018). Considerations and guidelines for mouse metabolic phenotyping in
diabetes research. *Diabetologia* **61**, 526–538.

Alshammary, A. F., Alshammari, A. M., Alsobaie, S. F., Alageel, A. A. and Ali Khan, I. (2023).
Evidence from genetic studies among rs2107538 variant in the *CCL5* gene and Saudi patients
diagnosed with type 2 diabetes mellitus. *Saudi J. Biol. Sci.* **30**, 103658.

Al-Shukaili, A., AL-Ghafri, S., Al-Marhoobi, S., Al-Abri, S., Al-Lawati, J. and Al-Maskari, M. (2013).
Analysis of Inflammatory Mediators in Type 2 Diabetes Patients. *Int. J. Endocrinol.* **2013**, 976810.

An In Vitro Organ Culture Model of the Murine Intervertebral Disc.

Arendt, L. M., McCready, J., Keller, P. J., Baker, D. D., Naber, S. P., Seewaldt, V. and Kuperwasser,
C. (2013). Obesity Promotes Breast Cancer by CCL2-Mediated Macrophage Recruitment and
Angiogenesis. *Cancer Res.* **73**, 6080–6093.

Arner, E., Mejhert, N., Kulyté, A., Balwierz, P. J., Pachkov, M., Cormont, M., Lorente-Cebrián, S.,
Ehrlund, A., Laurencikiene, J., Hedén, P., et al. (2012). Adipose Tissue MicroRNAs as
Regulators of CCL2 Production in Human Obesity. *Diabetes* **61**, 1986–1993.

Atri, C., Guerfali, F. Z. and Laouini, D. (2018). Role of Human Macrophage Polarization in Inflammation
during Infectious Diseases. *Int. J. Mol. Sci.* **19**, 1801.

Blondel, V. D., Guillaume, J.-L., Lambiotte, R. and Lefebvre, E. (2008). Fast unfolding of communities
in large networks. *J. Stat. Mech. Theory Exp.* **2008**, P10008.

Boff, D., Crijns, H., Janssens, R., Vanheule, V., Menezes, G. B., Macari, S., Silva, T. A., Amaral, F. A.
and Proost, P. (2018). The chemokine fragment CXCL9(74–103) diminishes neutrophil
recruitment and joint inflammation in antigen-induced arthritis. *J. Leukoc. Biol.* **104**, 413–422.

Bond, M., Fabunmi, R. P., Baker, A. H. and Newby, A. C. (1998). Synergistic upregulation of
metalloproteinase-9 by growth factors and inflammatory cytokines: an absolute requirement for
transcription factor NF-κB. *FEBS Lett.* **435**, 29–34.

Boucher, J., Kleinridders, A. and Kahn, C. R. (2014). Insulin Receptor Signaling in Normal and Insulin-
Resistant States. *Cold Spring Harb. Perspect. Biol.* **6**, a009191–a009191.

Broz, K., Walk, R. E. and Tang, S. Y. (2021). Complications in the spine associated with type 2 diabetes:
The role of advanced glycation end-products. *Med. Nov. Technol. Devices* **11**, 100065.

Calle, M. C. and Fernandez, M. L. (2012). Inflammation and type 2 diabetes. *Diabetes Metab.* **38**, 183–
191.

Cannata, F., Vadalà, G., Ambrosio, L., Fallucca, S., Napoli, N., Papalia, R., Pozzilli, P. and Denaro, V.
(2019). Intervertebral disc degeneration: A focus on obesity and type 2 diabetes. *Diabetes Metab.*
Res. Rev. **36**,.

Cefalu, W. T. (2006). Animal Models of Type 2 Diabetes: Clinical Presentation and Pathophysiological
Relevance to the Human Condition. *ILAR J.* **47**, 186–198.

Chang, H.-D. and Radbruch, A. (2007). The Pro- and Anti-Inflammatory Potential of Interleukin-12. *Ann.*
N. Y. Acad. Sci. **1109**, 40–46.

Chang, T.-T., Lin, L.-Y. and Chen, J.-W. (2021). A Novel Resolution of Diabetes: C-C Chemokine Motif
Ligand 4 Is a Common Target in Different Types of Diabetes by Protecting Pancreatic Islet Cell
and Modulating Inflammation. *Front. Immunol.* **12**,.

Chen, H., Charlat, O., Tartaglia, L. A., Woolf, E. A., Weng, X., Ellis, S. J., Lakey, N. D., Culpepper, J.,
More, K. J., Breitbart, R. E., et al. (1996). Evidence That the Diabetes Gene Encodes the Leptin
Receptor: Identification of a Mutation in the Leptin Receptor Gene in db/db Mice. *Cell* **84**, 491–
495.

Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X. and Zhao, L. (2017).
Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget* **9**, 7204–
7218.

Chen, L., Zhang, Q., Yu, C., Wang, F. and Kong, X. (2020). Functional roles of CCL5/RANTES in liver
disease. *Liver Res.* **4**, 28–34.

Curic, G. (2021). Intervertebral Disc and Adipokine Leptin—Loves Me, Loves Me Not. *Int. J. Mol. Sci.* **22**,
375.

Dalgaard, L. T. and Pedersen, O. (2001). Uncoupling proteins: functional characteristics and role in the
pathogenesis of obesity and Type II diabetes. *Diabetologia* **44**, 946–965.

Daryabor, G., Atashzar, M. R., Kabelitz, D., Meri, S. and Kalantar, K. (2020). The Effects of Type 2
Diabetes Mellitus on Organ Metabolism and the Immune System. *Front. Immunol.* **11**,.

de Candia, P., Prattichizzo, F., Garavelli, S., De Rosa, V., Galgani, M., Di Rella, F., Spagnuolo, M. I.,
Colamatteo, A., Fusco, C., Micillo, T., et al. (2019). Type 2 Diabetes: How Much of an
Autoimmune Disease? *Front. Endocrinol.* **10**,.

De Geer, C. M. (2018). Cytokine Involvement in Biological Inflammation Related to Degenerative
Disorders of the Intervertebral Disk: A Narrative Review. *J. Chiropr. Med.* **17**, 54–62.

DeLeon-Pennell, K. Y., Iyer, R. P., Ero, O. K., Cates, C. A., Flynn, E. R., Cannon, P. L., Jung, M.,
Shannon, D., Garrett, M. R., Buchanan, W., et al. Periodontal-induced chronic inflammation
triggers macrophage secretion of Ccl12 to inhibit fibroblast-mediated cardiac wound healing. *JCI*
Insight **2**, e94207.

Eckhardt, B. A., Rowsey, J. L., Thicke, B. S., Fraser, D. G., O'Grady, K. L., Bondar, O. P., Hines, J.
M., Singh, R. J., Thoreson, A. R., Rakshit, K., et al. Accelerated osteocyte senescence and
skeletal fragility in mice with type 2 diabetes. *JCI Insight* **5**, e135236.

Etna, M. P., Giacomini, E., Severa, M. and Coccia, E. M. (2014). Pro- and anti-inflammatory cytokines
in tuberculosis: A two-edged sword in TB pathogenesis. *Semin. Immunol.* **26**, 543–551.

Francisco, V., Pino, J., Campos-Cabaleiro, V., Ruiz-Fernández, C., Mera, A., Gonzalez-Gay, M. A.,
Gómez, R. and Gualillo, O. (2018). Obesity, Fat Mass and Immune System: Role for Leptin.
Front. Physiol. **9**, 640.

Gilbert, E. R., Fu, Z. and Liu, D. (2011). Development of a Nongenetic Mouse Model of Type 2 Diabetes.
J. Diabetes Res. **2011**, 416254.

Gruber, H. E., Ingram, J. A., Hoelscher, G. L. and Hanley, E. N. (2007). Leptin expression by annulus
cells in the human intervertebral disc. *Spine J.* **7**, 437–443.

Guest, C. B., Park, M. J., Johnson, D. R. and Freund, G. G. (2008). The implication of proinflammatory
cytokines in type 2 diabetes. *Front. Biosci.-Landmark* **13**, 5187–5194.

Han, Y.-C., Ma, B., Guo, S., Yang, M., Li, L.-J., Wang, S.-J. and Tan, J. (2018). Leptin regulates disc
cartilage endplate degeneration and ossification through activation of the MAPK-ERK signalling
pathway in vivo and in vitro. *J. Cell. Mol. Med.* **22**, 2098–2109.

He, M., Dong, H., Huang, Y., Lu, S., Zhang, S., Qian, Y. and Jin, W. (2016). Astrocyte-Derived CCL2 is
Associated with M1 Activation and Recruitment of Cultured Microglial Cells. *Cell. Physiol.*
Biochem. **38**, 859–870.

Herder, C., Carstensen, M. and Ouwens, D. M. (2013). Anti-inflammatory cytokines and risk of type 2
diabetes. *Diabetes Obes. Metab.* **15**, 39–50.

Higurashi, M., Ohya, Y., Joh, K., Muraguchi, M., Nishimura, M., Terawaki, H., Yagui, K., Hashimoto,
N., Saito, Y. and Yamada, K. (2009). Increased urinary levels of CXCL5, CXCL8 and CXCL9 in
patients with Type 2 diabetic nephropathy. *J. Diabetes Complications* **23**, 178–184.

House, I. G., Savas, P., Lai, J., Chen, A. X. Y., Oliver, A. J., Teo, Z. L., Todd, K. L., Henderson, M. A.,
Giuffrida, L., Petley, E. V., et al. (2020). Macrophage-Derived CXCL9 and CXCL10 Are Required
for Antitumor Immune Responses Following Immune Checkpoint Blockade. *Clin. Cancer Res.* **26**,
487–504.

Huang, J., Yang, G., Xiong, X., Wang, M., Yuan, J., Zhang, Q., Gong, C., Qiu, Z., Meng, Z., Xu, R., et
al. (2020). Age-related CCL12 Aggravates Intracerebral Hemorrhage-induced Brain Injury via
Recruitment of Macrophages and T Lymphocytes. *Aging Dis.* **11**, 1103–1115.

Inayat, H., Azim, M. K. and Baloch, A. A. (2019). Analysis of Inflammatory Gene Expression Profile of
Peripheral Blood Leukocytes in Type 2 Diabetes. *Immunol. Invest.* **48**, 618–631.

Islam, Md. S. and Wilson, R. D. (2012). Experimentally Induced Rodent Models of Type 2 Diabetes. In
Animal Models in Diabetes Research (ed. Joost, H.-G.), Al-Hasani, H.), and Schürmann, A.), pp.
161–174. Totowa, NJ: Humana Press.

Itariu, B. K. and Stulnig, T. M. (2014). Autoimmune Aspects of Type 2 Diabetes Mellitus - A Mini-Review.
Gerontology **60**, 189–196.

Jhawar, B. S., Fuchs, C. S., Colditz, G. A. and Stampfer, M. J. (2006). Cardiovascular risk factors for
physician-diagnosed lumbar disc herniation. *Spine J.* **6**, 684–691.

Juhas, U., Ryba-Stanisławowska, M., Szargiej, P. and Myśliwska, J. (2015). Different pathways of
macrophage activation and polarization. *Postępy Hig. Med. Dośw.* **69**, 496–502.

Keophiphath, M., Rouault, C., Divoux, A., Clément, K. and Lacasa, D. (2010). CCL5 Promotes
Macrophage Recruitment and Survival in Human Adipose Tissue. *Arterioscler. Thromb. Vasc.*
Biol. **30**, 39–45.

Kranjc, M. K., Novak, M., Pestell, R. G. and Lah, T. T. (2019). Cytokine CCL5 and receptor CCR5 axis
in glioblastoma multiforme. *Radiol. Oncol.* **53**, 397–406.

Kuo, P., Tuong, Z. K., Teoh, S. M., Frazer, I. H., Mattarollo, S. R. and Leggatt, G. R. (2018). HPV16E7-
Induced Hyperplasia Promotes CXCL9/10 Expression and Induces CXCR3+ T-Cell Migration to
Skin. *J. Invest. Dermatol.* **138**, 1348–1359.

Kusakabe, T., Tanioka, H., Ebihara, K., Hirata, M., Miyamoto, L., Miyanaga, F., Hige, H., Aotani, D.,
Fujisawa, T., Masuzaki, H., et al. (2009). Beneficial effects of leptin on glycaemic and lipid
control in a mouse model of type 2 diabetes with increased adiposity induced by streptozotocin
and a high-fat diet. *Diabetologia* **52**, 675–683.

Lança, T., Costa, M. F., Gonçalves-Sousa, N., Rei, M., Grosso, A. R., Penido, C. and Silva-Santos, B.
(2013). Protective Role of the Inflammatory CCR2/CCL2 Chemokine Pathway through
Recruitment of Type 1 Cytotoxic γδ T Lymphocytes to Tumor Beds. *J. Immunol.* **190**, 6673–6680.

Lee, G.-H., Proenca, R., Montez, J. M., Carroll, K. M., Darvishzadeh, J. G., Lee, J. I. and Friedman, J.
M. (1996). Abnormal splicing of the leptin receptor in diabetic mice. *Nature* **379**, 632–635.

Li, Z., Shen, J., Wu, W. K. K., Yu, X., Liang, J., Qiu, G. and Liu, J. (2013). The role of leptin on the
organization and expression of cytoskeleton elements in nucleus pulposus cells. *J. Orthop. Res.*
31, 847–857.

Li, X., Liu, X., Wang, Y., Cao, F., Chen, Z., Hu, Z., Yu, B., Feng, H., Ba, Z., Liu, T., et al. (2020).
Intervertebral disc degeneration in mice with type II diabetes induced by leptin receptor
deficiency. *BMC Musculoskelet. Disord.* **21**, 77.

Lim, S. Y., Yuzhalin, A. E., Gordon-Weeks, A. N. and Muschel, R. J. (2016). Targeting the CCL2-CCR2
signaling axis in cancer metastasis. *Oncotarget* **7**, 28697–28710.

Lin, Y. and Sun, Z. (2010). Current views on type 2 diabetes. *J. Endocrinol.* **204**, 1.

Lin, K. H. and Tang, S. Y. (2017). The Quantitative Structural and Compositional Analyses of
Degenerating Intervertebral Discs Using Magnetic Resonance Imaging and Contrast-Enhanced
Micro-Computed Tomography. *Ann. Biomed. Eng.* **45**, 2626–2634.

Lin, K. H., Wu, Q., Leib, D. J. and Tang, S. Y. (2016). A novel technique for the contrast-enhanced
microCT imaging of murine intervertebral discs. *J. Mech. Behav. Biomed. Mater.* **63**, 66–74.

Lintz, M., Walk, R. E., Tang, S. Y. and Bonassar, L. J. (2022). The degenerative impact of
hyperglycemia on the structure and mechanics of developing murine intervertebral discs. *JOR*
Spine **5**, e1191.

Liu, J. W., Abraham, A. C. and Y. Tang, S. (2015). The high-throughput phenotyping of the viscoelastic
behavior of whole mouse intervertebral discs using a novel method of dynamic mechanical
testing. *J. Biomech.* **48**, 2189–2194.

Liu, X., Pan, F., Ba, Z., Wang, S. and Wu, D. (2018). The potential effect of type 2 diabetes mellitus on
lumbar disc degeneration: a retrospective single-center study. *J. Orthop. Surg.* **13**, 52.

Luo, J., Quan, J., Tsai, J., Hobensack, C. K., Sullivan, C., Hector, R. and Reaven, G. M. (1998).
Nongenetic mouse models of non—insulin-dependent diabetes mellitus. *Metabolism* **47**, 663–
668.

Lyu, F.-J., Cui, H., Pan, H., MC Cheung, K., Cao, X., Iatridis, J. C. and Zheng, Z. (2021). Painful
intervertebral disc degeneration and inflammation: from laboratory evidence to clinical
interventions. *Bone Res.* **9**, 1–14.

Malemud, C. J. (2019). Inhibition of MMPs and ADAM/ADAMTS. *Biochem. Pharmacol.* **165**, 33–40.

Marcovecchio, P. M., Thomas, G. and Salek-Ardakani, S. (2021). CXCL9-expressing tumor-associated
macrophages: new players in the fight against cancer. *J. Immunother. Cancer* **9**, e002045.

Markovics, A., Rosenthal, K. S., Mikecz, K., Carambula, R. E., Ciemielewski, J. C. and Zimmerman,
D. H. (2022). Restoring the Balance between Pro-Inflammatory and Anti-Inflammatory Cytokines
in the Treatment of Rheumatoid Arthritis: New Insights from Animal Models. *Biomedicines* **10**, 44.

Medrikova, D., Jilkova, Z. M., Bardova, K., Janovska, P., Rossmeisl, M. and Kopecky, J. (2012). Sex
differences during the course of diet-induced obesity in mice: adipose tissue expandability and
glycemic control. *Int. J. Obes.* **36**, 262–272.

Melgoza, I. P., Chenna, S. S., Tessier, S., Zhang, Y., Tang, S. Y., Ohnishi, T., Novais, E. J., Kerr, G. J.,
Mohanty, S., Tam, V., et al. (2021). Development of a standardized histopathology scoring

system using machine learning algorithms for intervertebral disc degeneration in the mouse
model—An ORS spine section initiative. *JOR SPINE* **4**, e1164.

Mir, M. M., Alfaifi, J., Sohail, S. K., Rizvi, S. F., Akhtar, M. T., Alghamdi, M. A. A., Mir, R., Wani, J. I.,
Sabah, Z. U., Alhumaydhi, F. A., et al. (2024). The Role of Pro-Inflammatory Chemokines CCL-
1, 2, 4, and 5 in the Etiopathogenesis of Type 2 Diabetes Mellitus in Subjects from the Asir
Region of Saudi Arabia: Correlation with Different Degrees of Obesity. *J. Pers. Med.* **14**, 743.

Molinos, M., Almeida, C. R., Caldeira, J., Cunha, C., Gonçalves, R. M. and Barbosa, M. A. (2015).
Inflammation in intervertebral disc degeneration and regeneration. *J. R. Soc. Interface* **12**,
20141191.

Mortara, L., Balza, E., Bruno, A., Poggi, A., Orecchia, P. and Carnemolla, B. (2018). Anti-cancer
Therapies Employing IL-2 Cytokine Tumor Targeting: Contribution of Innate, Adaptive and
Immunosuppressive Cells in the Anti-tumor Efficacy. *Front. Immunol.* **9**,.

Motwani, M. and Gilroy, D. (2015). Macrophage development and polarization in chronic inflammation.
Semin. Immunol. **27**,.

Mu, J., Woods, J., Zhou, Y.-P., Roy, R. S., Li, Z., Zycband, E., Feng, Y., Zhu, L., Li, C., Howard, A. D.,
et al. (2006). Chronic Inhibition of Dipeptidyl Peptidase-4 With a Sitagliptin Analog Preserves
Pancreatic β-Cell Mass and Function in a Rodent Model of Type 2 Diabetes. *Diabetes* **55**, 1695–
1704.

Mu, J., Petrov, A., Eiermann, G. J., Woods, J., Zhou, Y.-P., Li, Z., Zycband, E., Feng, Y., Zhu, L., Roy,
R. S., et al. (2009). Inhibition of DPP-4 with sitagliptin improves glycemic control and restores
islet cell mass and function in a rodent model of type 2 diabetes. *Eur. J. Pharmacol.* **623**, 148–
154.

Natelson, D. M., Lai, A., Krishnamoorthy, D., Hoy, R. C., Iatridis, J. C. and Illien-Jünger, S. (2020).
Leptin signaling and the intervertebral disc: Sex dependent effects of leptin receptor deficiency
and Western diet on the spine in a type 2 diabetes mouse model. *PLOS ONE* **15**, e0227527.

National Diabetes Statistics Report | Diabetes | CDC (2022).

Navone, S. E., Marfia, G., Giannoni, A., Beretta, M., Guarnaccia, L., Gualtierotti, R., Nicoli, D.,
Rampini, P. and Campanella, R. (2017). Inflammatory mediators and signalling pathways
controlling intervertebral disc degeneration. *Histol. Histopathol.* 523–542.

Nawaz, M. I., Van Raemdonck, K., Mohammad, G., Kangave, D., Van Damme, J., Abu El-Asrar, A. M.
and Struyf, S. (2013). Autocrine CCL2, CXCL4, CXCL9 and CXCL10 signal in retinal endothelial
cells and are enhanced in diabetic retinopathy. *Exp. Eye Res.* **109**, 67–76.

Neumeier, M., Bauer, S., Brühl, H., Eisinger, K., Kopp, A., Abke, S., Walter, R., Schäffler, A. and
Buechler, C. (2011). Adiponectin stimulates release of CCL2, -3, -4 and -5 while the surface
abundance of CCR2 and -5 is simultaneously reduced in primary human monocytes. *Cytokine* **56**,
573–580.

Ochiai, E., Sa, Q., Brogli, M., Kudo, T., Wang, X., Dubey, J. P. and Suzuki, Y. (2015). CXCL9 Is
Important for Recruiting Immune T Cells into the Brain and Inducing an Accumulation of the T
Cells to the Areas of Tachyzoite Proliferation to Prevent Reactivation of Chronic Cerebral Infection
with *Toxoplasma gondii. Am. J. Pathol.* **185**, 314–324.

Pan, X., Kaminga, A. C., Wen, S. W. and Liu, A. (2021). Chemokines in Prediabetes and Type 2
Diabetes: A Meta-Analysis. *Front. Immunol.* **12**,.

Pelisch, N., Rosas Almanza, J., Stehlik, K. E., Aperi, B. V. and Kroner, A. (2020). CCL3 contributes to
secondary damage after spinal cord injury. *J. Neuroinflammation* **17**, 362.

Petersen, M. C. and Shulman, G. I. (2018). Mechanisms of Insulin Action and Insulin Resistance.
Physiol. Rev. **98**, 2133–2223.

Pettersson, U. S., Waldén, T. B., Carlsson, P.-O., Jansson, L. and Phillipson, M. (2012). Female Mice
are Protected against High-Fat Diet Induced Metabolic Syndrome and Increase the Regulatory T
Cell Population in Adipose Tissue. *PLOS ONE* **7**, e46057.

Pettigrew, K. A., McKnight, A. J., Patterson, C. C., Kilner, J., Sadlier, D. M. and Maxwell, A. P. (2010).
Resequencing of the CCL5 and CCR5 genes and investigation of variants for association with
diabetic nephropathy. *J. Hum. Genet.* **55**, 248–251.

Pinto, E. M., Neves, J. R., Laranjeira, M. and Reis, J. (2023). The importance of inflammatory
biomarkers in non-specific acute and chronic low back pain: a systematic review. *Eur. Spine J.*

Rebuffat, S. A., Sidot, E., Guzman, C., Azay-Milhau, J., Jover, B., Lajoix, A.-D. and Peraldi-Roux, S.
(2018). Adipose tissue derived-factors impaired pancreatic β-cell function in diabetes. *Biochim.*
Biophys. Acta BBA - Mol. Basis Dis. **1864**, 3378–3387.

Risbud, M. V. and Shapiro, Irving. M. (2014). Role of Cytokines in Intervertebral Disc Degeneration:
Pain and Disc-content. *Nat. Rev. Rheumatol.* **10**, 44–56.

Robinson, D., Mirovsky, Y., Halperin, N., Evron, Z. and Nevo, Z. (1998). Changes in Proteoglycans of
Intervertebral Disc in Diabetic Patients: A Possible Cause of Increased Back Pain. *Spine* **23**, 849.

Ruytinx, P., Proost, P., Van Damme, J. and Struyf, S. (2018). Chemokine-Induced Macrophage
Polarization in Inflammatory Conditions. *Front. Immunol.* **9**,.

Sagar, D., Lamontagne, A., Foss, C. A., Khan, Z. K., Pomper, M. G. and Jain, P. (2012). Dendritic cell
CNS recruitment correlates with disease severity in EAE via CCL2 chemotaxis at the blood–brain
barrier through paracellular transmigration and ERK activation. *J. Neuroinflammation* **9**, 245.

Sakaguchi, K., Takeda, K., Maeda, M., Ogawa, W., Sato, T., Okada, S., Ohnishi, Y., Nakajima, H. and
Kashiwagi, A. (2015). Glucose area under the curve during oral glucose tolerance test as an
index of glucose intolerance. *Diabetol. Int.* **7**, 53–58.

Sakellaridis, N. (2006). The influence of diabetes mellitus on lumbar intervertebral disk herniation. *Surg.*
Neurol. **66**, 152–154.

Segar, A. H., Fairbank, J. C. T. and Urban, J. (2019). Leptin and the intervertebral disc: a biochemical
link exists between obesity, intervertebral disc degeneration and low back pain—an in vitro study
in a bovine model. *Eur. Spine J.* **28**, 214–223.

Shachar, I. and Karin, N. (2013). The dual roles of inflammatory cytokines and chemokines in the
regulation of autoimmune diseases and their clinical implications. *J. Leukoc. Biol.* **93**, 51–61.

Shamji, M. F., Setton, L. A., Jarvis, W., So, S., Chen, J., Jing, L., Bullock, R., Isaacs, R. E., Brown, C.
and Richardson, W. J. (2010). Proinflammatory cytokine expression profile in degenerated and
herniated human intervertebral disc tissues. *Arthritis Rheum.* **62**, 1974–1982.

Sharma, A. (2018). The Role of Adipokines in Intervertebral Disc Degeneration. *Med. Sci.* **6**, 34.

Sheng, D., Ma, W., Zhang, R., Zhou, L., Deng, Q., Tu, J., Chen, W., Zhang, F., Gao, N., Dong, M., et
al. (2022). Ccl3 enhances docetaxel chemosensitivity in breast cancer by triggering
proinflammatory macrophage polarization. *J. Immunother. Cancer* **10**, e003793.

Sindhu, S., Akhter, N., Arefanian, H., Al-Roub, A. A., Ali, S., Wilson, A., Al-Hubail, A., Al-Beloushi,
S., Al-Zanki, S. and Ahmad, R. (2017). Increased circulatory levels of fractalkine (CX3CL1) are
associated with inflammatory chemokines and cytokines in individuals with type-2 diabetes. *J.*
Diabetes Metab. Disord. **16**, 15.

Sindhu, S., Kochumon, S., Shenouda, S., Wilson, A., Al-Mulla, F. and Ahmad, R. (2019). The
Cooperative Induction of CCL4 in Human Monocytic Cells by TNF-α and Palmitate Requires
MyD88 and Involves MAPK/NF-κB Signaling Pathways. *Int. J. Mol. Sci.* **20**, 4658.

Srinivasan, K. and Ramarao, P. (2007). Animal models in type 2 diabetes research: An overview. *Indian*
J. Med. Res. **125**, 451.

Stott, N. L. and Marino, J. S. (2020). High Fat Rodent Models of Type 2 Diabetes: From Rodent to
Human. *Nutrients* **12**, 3650.

Stubbins, R. E., Holcomb, V. B., Hong, J. and Núñez, N. P. (2012). Estrogen modulates abdominal
adiposity and protects female mice from obesity and impaired glucose tolerance. *Eur. J. Nutr.* **51**,
861–870.

Sullivan, T. J., Miao, Z., Zhao, B. N., Ertl, L. S., Wang, Y., Krasinski, A., Walters, M. J., Powers, J. P.,
Dairaghi, D. J., Baumgart, T., et al. (2013). Experimental evidence for the use of CCR2
antagonists in the treatment of type 2 diabetes. *Metabolism* **62**, 1623–1632.

Suri, S., Mitra, P., Abhilasha, A., Saxena, I., Garg, M. K., Bohra, G. K. and Sharma, P. (2022). Role of
interleukin-2 and interleukin-18 in newly diagnosed type 2 diabetes mellitus. *J. Basic Clin.*
Physiol. Pharmacol. **33**, 185–190.

Tian, Y., Yuan, W., Fujita, N., Wang, J., Wang, H., Shapiro, I. M. and Risbud, M. V. (2013).
Inflammatory Cytokines Associated with Degenerative Disc Disease Control Aggrecanase-1
(ADAMTS-4) Expression in Nucleus Pulposus Cells through MAPK and NF-κB. *Am. J. Pathol.*
182, 2310–2321.

Velikova, T. V., Kabakchieva, P. P., Assyov, Y. S. and Georgiev, T. A. (2021). Targeting Inflammatory
Cytokines to Improve Type 2 Diabetes Control. *BioMed Res. Int.* **2021**, 1–12.

Venetz, D., Ponzoni, M., Schiraldi, M., Ferreri, A. J. M., Bertoni, F., Doglioni, C. and Uguccioni, M.
(2010). Perivascular expression of CXCL9 and CXCL12 in primary central nervous system
lymphoma: T-cell infiltration and positioning of malignant B cells. *Int. J. Cancer* **127**, 2300–2312.

Wang, B., P., C. C. and Pippin, J. J. (2014). Leptin- and Leptin Receptor-Deficient Rodent Models:
Relevance for Human Type 2 Diabetes. *Curr. Diabetes Rev.* **10**, 131–145.

Wuertz, K. and Haglund, L. (2013). Inflammatory Mediators in Intervertebral Disk Degeneration and
Discogenic Pain. *Glob. Spine J.* **3**, 175–184.

Xu, P., Zhang, F., Chang, M., Zhong, C., Sun, C.-H., Zhu, H.-R., Yao, J.-C., Li, Z.-Z., Li, S.-T., Zhang,
W.-C., et al. (2021a). Recruitment of γδ T cells to the lesion via the CCL2/CCR2 signaling after
spinal cord injury. *J. Neuroinflammation* **18**, 64.

Figure 1. Experimental design, animal model, and workflow of the current study. (A) The db/db model arises due to a point mutation in the leptin receptor gene, while the STZ-HFD model develops diabetes through a pro-glycemic diet and beta cell impairment. Both models present symptoms of obesity, chronic hyperglycemia, and insulin resistance, though the magnitude can vary both between and within each model. **(B)** The experimental timeline outlines the progression of the study for both models. The db/db mice are acquired at skeletal maturity (12 weeks old) and sacrificed after metabolic measurements are collected. For the STZ-HFD model, mice undergo a lead-in phase of 4-6 weeks on a HFD followed by a single low dose of STZ. Subsequently, the experimental phase for the STZ-HFD mice continues with HFD for 12 weeks, with periodic assessments of fasting blood glucose, A1c levels, and glucose tolerance. **(C)** After sacrifice, FSUs, consisting of the intervertebral disc and the adjacent vertebral bodies, are extracted from mice. FSUs are then utilized for terminal measures pictured above.

Xu, L., Chen, Y., Nagashimada, M., Ni, Y., Zhuge, F., Chen, G., Li, H., Pan, T., Yamashita, T., Mukaida,
N., et al. (2021b). CC chemokine ligand 3 deficiency ameliorates diet-induced steatohepatitis by
regulating liver macrophage recruitment and M1/M2 status in mice. *Metabolism* **125**, 154914.

Yang, J., Agarwal, M., Ling, S., Teitz-Tennenbaum, S., Zemans, R. L., Osterholzer, J. J., Sisson, T.
H. and Kim, K. K. (2020). Diverse Injury Pathways Induce Alveolar Epithelial Cell CCL2/12,
Which Promotes Lung Fibrosis. *Am. J. Respir. Cell Mol. Biol.* **62**, 622–632.

Yin, R., Xue, Y., Hu, J., Hu, X. and Shen, Q. (2020). The effects of diet and streptozotocin on metabolism
and gut microbiota in a type 2 diabetes mellitus mouse model. *Food Agric. Immunol.* **31**, 723–
739.

Zeng, Z., Lan, T., Wei, Y. and Wei, X. (2022). CCL5/CCR5 axis in human diseases and related
treatments. *Genes Dis.* **9**, 12–27.

Zhang, Y., Lin, J. and Wei, F. (2015). The Function and Roles of ADAMTS-7 in Inflammatory Diseases.
Mediators Inflamm. **2015**, 801546.

Zhang, X., Fan, L., Wu, J., Xu, H., Leung, W. Y., Fu, K., Wu, J., Liu, K., Man, K., Yang, X., et al. (2019).
Macrophage p38α promotes nutritional steatohepatitis through M1 polarization. *J. Hepatol.* **71**,
163–174.

Figure 4. Comparative analysis of IVD structure, mechanics, and composition are similar between db/db and STZ-HFD
748 **mice. (A)-(C)** NPVF, NIDI, and DHI indicate few significant differences in structural integrity between the models. **(D)-(F)** Load slope, energy dissipated, and phase shift demonstrating no significant variations in viscoelastic mechanical behavior. **(G)-(I)** Biochemical content measurements, including collagen, s-GAG, and AGEs, show no significant differences.

Figure 2. Both db/db and STZ-HFD mice represent a characteristically T2D phenotype. (A) AUC-GLU shows elevated glucose
Figure 5. STZ-HFD IVD Produces a More Pro-Inflammatory Microenvironment than the db/db IVD in Comparative Analysis of Cytokine Expression. (A) The STZ-HFD model shows a significantly higher fold increase for eight cytokines: CXCL2, CCL2, CCL3, CCL4, CCL12 (monocyte/macrophage associated), IL-2, CXCL9 (T-cell associated), and CCL5 (pleiotropic). **(B)** The STZ-HFD model encompasses a broader and more pronounced cytokine response compared to the db/db model, highlighting the extensive upregulation of inflammatory cytokines in the STZ-HFD model.

Figure 3. STZ-HFD mice exhibit more severe histopathological IVD degeneration compared to db/db mice. (A) Comparison
Figure 6. STZ-HFD IVD Invokes Unique Inflammatory Signaling Pathways in Networks of Cytokine Expression. (A) The STZ-HFD model shows a distinct network structure, demonstrating the unique upregulation of various inflammatory pathways **(B)** The STZ-HFD and db/db networks each rely on a number of unique (red/blue) and shared (purple) cytokines, indicating both leptin-dependent and leptin-independent inflammatory signaling cascades **(C)** The STZ-HFD mouse model displays a fragmented and modular cytokine network, indicating the parallel signaling of multiple signaling pathways.

(A)

Best | Median | Worst

db/+
Score: 4 | Score: 7 | Score: 7

db/db
Score: 9 | Score: 10 | Score: 11

Con+Veh
Score: 4 | Score: 6 | Score: 6

STZHFD
Score: 7 | Score: 10 | Score: 12

(B) IVD Degeneration

Degeneration Score (0-35)

p = 0.0143

db/+   db/db   Con+Veh   STZHFD

NP
AF
IF
EP

(C)

(A)

Control+Vehicle

STZHFD

Eigenvector Centrality

High

Low

(B) Heatmap of Most Central Cytokines

Cytokines

CCL11
CSF3
CXCL2
IL16
IL6
CCL2
CCL4
CXCL10
CXCL5
CXCL9
IL11
IL4

STZHFD   db/db

Condition

Unique Centrality (Diabetic Δ Control)

Shared Centrality (STZHFD ∩ db/db)

No Centrality

(C)

Metric	Con+Veh	STZHFD	db/+	db/db
Path Length	1.3	2.3	1.3	1.3
Modularity	0.34	0.46	0.29	0.18

	Con+Veh vs STZHFD	db/db vs STZHFD	db/+ vs db/db	db/+ vs db/db
Jaccard Index (k=1)	0.14	0.13	0.21	0.15
Jaccard Index (k=2)	0.25	0.24	0.37	0.32