1
2
3
4   A distinct circuit for biasing visual perceptual decisions and modulating superior
5   colliculus activity through the mouse posterior striatum
6
7
8   Running title:
9   Posterior striatum biases visual decisions

15   Kara K. Cover*, Kerry Elliott, Sarah M. Preuss & Richard J. Krauzlis*
16
17
18
19
20   Laboratory of Sensorimotor Research, National Eye Institute
21   Bethesda, Maryland 20892 USA
22
23
24
25 *Corresponding authors:
26 Richard J. Krauzlis: richard.krauzlis@nih.gov
27 Kara K. Cover: kara.cover@nih.gov

28
29
30
31 48 pages
32 8 figures
33 3 tables
34
35 Abstract: 188 words
36 Introduction: 601 words
37 Discussion: 1290 words
38
39
40 Conflicts of interest: none.
41
42

43 Acknowledgements: This work is supported by the National Eye Institute Intramural
44 Research Program at the National Institutes of Health ZIA EY000511 (R.J.K.). The
45 authors thank Dr. Carlos Mejias-Aponte for assistance with histological analysis.

Abstract

47
The basal ganglia play a key role in visual perceptual decisions. Despite being

the primary target in the basal ganglia for inputs from the visual cortex, the posterior

striatum's (PS) involvement in visual perceptual behavior remains unknown in rodents.

We reveal that the PS direct pathway is largely segregated from the dorsomedial

striatum (DMS) direct pathway, the other major striatal target for visual cortex. We

investigated the role of the PS in visual perceptual decisions by optogenetically

stimulating striatal medium spiny neurons in the direct pathway (D1-MSNs) of mice

performing a visual change-detection task. PS D1-MSN activation robustly biased visual

decisions in a manner dependent on visual context, timing, and reward expectation. We

examined the effects of PS and DMS direct pathway activation on neuronal activity in

the superior colliculus (SC), a major output target of the basal ganglia. Activation of

either direct pathway rapidly modulated SC neurons, but mostly targeted different SC

neurons and had opposite effects. These results demonstrate that the PS in rodents

provides an important route for controlling visual decisions, in parallel with the better

known DMS, and with distinct anatomical and functional properties.

Significance Statement

The rodent posterior striatum (PS) is strongly innervated by visual cortex and

thalamus, but its functional role in visual behavior has not been explored. We show that

the PS initiates a direct pathway through the basal ganglia that is anatomically distinct

from the more commonly studied dorsomedial striatum (DMS). Activating the PS direct

pathway selectively biases decisions for expected, valued visual events. We also show

that both DMS and PS direct pathways modulate neuronal activity in the superior

colliculus, a structure critical for visual processing and sensorimotor function, and

preferentially modulate collicular units with properties relevant in the visual detection

task. These findings identify a distinct and novel circuit through the basal ganglia for

controlling visually guided perceptual decisions.

Introduction

The basal ganglia comprise a set of interconnected subcortical nuclei that govern a range of complex functions. The best-known functions are related to motor control. The striatum, the input nucleus of the basal ganglia, facilitates elements of action execution including movement (Freeze et al., 2013), goal-directed action selection (Tai et al., 2012) and orienting to valued sensory stimuli (Hikosaka et al., 2019). However, a growing body of evidence implicates the basal ganglia in aspects of perceptual decision-making that are distinct from motor control (Ding, 2023). Caudate neurons in primates exhibit spatially selective modulation in the absence of goal-directed actions (Arcizet and Krauzlis, 2018). Neurons in the caudate of primates (Ding and Gold, 2010) and dorsal striatum of rats (Yartsev et al., 2018) show activity consistent with sensory evidence accumulation. Moreover, deficits in sensory perception are noted in disorders of basal ganglia function, such as Parkinson's (Permezel et al., 2023) and Huntington's (Profant et al., 2017) diseases.

Sensory information is routed to the striatum through cortical and thalamic nuclei associated with somatosensory, visual, and auditory modalities (Hintiryan et al., 2016; Hunnicutt et al., 2016). The majority of studies on striatal involvement in visual perceptual behavior have focused on the dorsomedial striatum (DMS) in rodents and the head of the caudate nucleus in primates which receive input from visual cortex (Yeterian and Pandya, 1995; Khibnik et al., 2014) as well as visual thalamus (Smith and Parent, 1986; Kamishina et al., 2008). Dopamine D1-receptor expressing medium spiny neurons (D1-MSNs) in the striatum initiate the direct pathway, a disynaptic circuit that is believed to gate behaviors by disinhibiting basal ganglia targets in the thalamus and

midbrain. Activating DMS D1-MSNs in mice biases visual decisions in a manner linked

to the allocation of spatial attention (Wang et al., 2018; Wang and Krauzlis, 2020). Thus,

several lines of evidence have established the direct pathway arising from the dorsal

striatum as an important circuit for visual perceptual decisions.

The posterior striatum (PS) is also innervated by visual cortex and thalamus

(Khibnik et al., 2014; Hunnicutt et al., 2016; Griggs et al., 2017; Lee et al., 2023). In the

primate homologue, the tail of the caudate nucleus, neurons in the direct pathway

encode visual cue value and bias the generation of saccades (Kim et al., 2017; Amita et

al., 2019, 2020). In the rodent, a role for PS in visually guided behavior remains

untested. PS D1-MSN signaling is causally involved in auditory decisions in mice (Guo

et al., 2018; Chen et al., 2022). Also, dopamine axons in the PS respond to light

(Menegas et al., 2018). It thus seems plausible that visual circuits through the PS could

be a major contributor to visual perceptual function, but this has yet to be explored in

rodents.

Here, we investigated a functional role for the PS direct pathway in visual

perceptual decisions. We first characterized this circuit using anatomical methods and

then tested its functional role by combining optogenetic PS or DMS D1-MSN

manipulations with SC neuronal recordings in mice making visual decisions. We found

that the direct pathway from the PS is largely segregated from that of the DMS and is

more broadly and inclusively innervated by visual cortex. Activation of this PS direct

pathway circuit biased performance in the visual change-detection task and modulated

the activity of SC neurons. Whereas the predominate neuronal effect of PS or DMS D1-

MSN activation was excitatory modulation of SC units, PS D1-MSN stimulation induced

inhibitory effects more frequently than that of DMS. Together, these findings

demonstrate that the PS provides a functionally significant and distinct route through the

basal ganglia for visual perceptual decisions.

Materials and Methods

127
Subjects

Male and female C57BL/6J (wild-type; Jackson Laboratory, #000664) or Drd1a-

Cre heterozygous (B6.FVB(Cg)-Tg(Drd1a-cre)EY217Gsat/Mmucd; MMRRC #034258-

UCD) mice were housed a in temperature and humidity-controlled vivarium under a 12-

hour reversed light/dark cycle (lights off at 0800 hours). All experimental procedures

occurred during the dark cycle. Mice were housed with littermates (two to five per cage),

except for those singly housed following head post and optic fiber implantation. Mice

performing behavioral tasks were weighed and fed daily to maintain (at least) 85% _ad_

libitum weight and the health status of each mouse was monitored daily throughout the

study. These mice had free access to water, but their intake of dry food was controlled

and were additionally given access to a nutritionally complete 8% soy-based infant

formula (Similac, Abbott, IL). All experimental and animal husbandry procedures were

approved by the NIH Institutional Animal Care and Use Committee (IACUC) and

complied with the Public Health Service policy on the humane care and use of

laboratory animals.

Surgical procedures

Mice of at least 6 weeks of age were anesthetized with isoflurane (4% induction,

0.8-1.5% maintenance) and secured by a stereotaxic frame with ear bars. A feedback-

controlled heating pad was used to maintain the body temperature at 37°C and

ophthalmic ointment was applied to the eyes. Dexamethasone (1.6 mg/kg, s.c.) and

meloxicam (2.0 mg/kg, s.c.) were administered to reduce inflammation and discomfort,

respectively. Viruses were infused at a rate of 2 nl/s (50 nl/min maximum) using a

microinjector (Nanoject III, Drummond Scientific, Broomall, PA).

For basal ganglia direct pathway tracing in Drd1-cre heterozygous mice, either

AAV1-Ef1A-DIO-iChloc-2a-dsRed (NIDA Genetic Engineering and Viral Vector Core) or

AAV1-Ef1A-DIO-eYFP (Addgene #27056-AAV1) was unilaterally injected in the DMS

(200 nl; distance from bregma in mm, [anterior-posterior, AP] 0.0, [medial-lateral, ML]

2.0, [dorsal-ventral, DV] -2.1) and the other virus injected in the PS (100 nl at each

depth; [AP] -1.5, [ML] 3.5, [DV] -2.5 and -3.1). The viruses injected in DMS and PS were

counter-balanced across 3 mice.

For di-synaptic tracing of DMS and PS direct pathways through the SN in wild-

type mice, either AAV1-hSyn-cre-WPRE-hGHpA (Addgene #105553-AAV1) or AAV1-

Ef1a-flp (Addgene #55637-AAV1) was unilaterally injected in the DMS and the other

virus in the PS (same volumes and coordinates as listed above), as well as Fluorogold

(30 nl, 1.5%, Fluorochrome LLC, Denver, CO) to label the injection site. In the same

mice, both AAV1-Ef1a-fDIO-mCherry (Addgene #114471-AAV1) and AAV1-Ef1a-DIO-

eYFP (Addgene #27056-AAV1) were injected in the SNR (150nl each; [AP] -3.2, [ML]

1.6, [DV] -4.4). The viruses injected in DMS and PS were counter-balanced across 4

mice. For di-synaptic tracing of the DMS direct pathway through the EP (200 nl; [AP] -

1.25, [ML] 1.9, [DV] -4.3), a single set of viruses were used (either AAV1-cre and AAV1-

DIO-eYFP, or AAV1-flp and AAV1-fDIO-mCherry), counter-balanced across 4 mice.

For retrograde tracing of projections to the striatum, Fluorogold (3%) was

unilaterally injected in the DMS (80 nl; [AP] 0.0, [ML] 2.15, [DV] -2.1) of 2 mice or in the

PS (40 nl at each depth; [AP] -1.5, [ML] 3.5, [DV] -2.5 and -2.95) of 2 mice.

For optogenetic manipulation of the direct pathway in mice performing behavioral

tasks, AAV2-Ef1a-DIO-hChR2(H134R)-mCherry (University of North Carolina viral core)

or a control virus AAV1-hSyn-DIO-mCherry (Addgene #50459-AAV1) was injected

unilaterally in the DMS (250 nl; [AP] 0.0, [ML] 1.7, [DV] -2.1) and the PS (80 nl at each

depth; [AP] -1.5, [ML] 3.25, [DV] -2.25 and -2.75). Additionally, optic fibers (200 µm core;

Plexon, Dallas, TX) were inserted in the DMS ([AP] 0.0, [ML] 1.7, [DV] -1.7) and PS

([AP] -1.5, [ML] 3.25, [DV] -2.15) and a custom-designed titanium head post was

secured to the skull using Metabond (Parkell Inc., Edgewood, NY). The skin wound

edge was then closed with surgical adhesive and treated with antibiotic ointment. Mice

assigned to behavioral testing began food restriction 18-20 days following surgery.

After mice were trained and tested on the visual detection task with optogenetic

manipulation, a second surgery was performed to implant electrophysiology microwire

bundles. Following the same anesthetic and analgesic procedure as detailed above, a

small craniotomy was made to implant a drivable 16-wire microwire bundle (Innovative

Neurophysiology, Durham, NC). The tip of the stainless-steel cannula of the bundle was

implanted at [AP] -3.4 – -3.9, [ML] 0.8 – 1.2 and [DV] -0.2. The cranial opening was

sealed with Vaseline petroleum jelly and the microwire bundle assembly was secured

on the skull with Metabond.

Immunohistochemistry was performed to confirm virus expression and fiber

placement. Animals were excluded from analysis for poor virus expression at target

regions or viral expression in non-target regions (i.e. cortex).

Immunohistochemistry

Mice were deeply anesthetized with isoflurane prior to transcardial perfusion with

0.1 M PBS, pH 7.3, followed by ice cold 4% paraformaldehyde in PBS. Brains were

extracted and stored in 4% paraformaldehyde at 4°C. 50 μm coronal sections were cut

using either a Campden Instruments 7000smz-2 vibrating microtome (Lafayette, IN) at

room temperature or, following transfer to 20% sucrose solution, a Leica SM2010 R

freezing microtome (Deer Park, IL). Rabbit anti-GFP (1:1000; G10362, Invitrogen,

Carlsbad, CA) and donkey anti-rabbit conjugated to Alexa Fluor 488 (1:1000; A21206,

Invitrogen) were used to amplify eYFP expression. Chicken anti-mCherry (1:2000;

NBP2-25158, Novus Biologicals, Littleton, CO) and donkey anti-chicken conjugated to

Alexa Fluor 594 (1:1000; 703-585-155, Jackson ImmunoResearch, West Grove, PA)

were used to amplify dsRed and mCherry.

Behavioral testing

Behavioral apparatus

Behavioral testing was conducted in a custom-built booth that displayed visual

stimuli on the left- and right-hand sides of the head-fixed mouse that was coupled to

their locomotion on a custom-built Styrofoam linear wheel(Krauzlis et al., 2020). A pair

of LCD displays (VS17287; ViewSonic, Brea, CA) were positioned at a 45° angle from

the midline of the animal so that the displays were centered on the right or left eye and

subtended by ~90° horizontal and ~55° vertical of the visual hemifield. The viewing

distance from the mouse to the screen was ~27.5 cm. Sound-absorbent material lined

the inside of the booth to reduce noise.

Experiments were controlled by a computer using a modified version the

PLDAPS system (Eastman and Huk, 2012), consisting of a Datapixx peripheral (Vpixx

Technologies, Saint-Bruno, QC, Canada) and the Psychophysics Toolbox (Brainard,

1997; Pelli, 1997) for MATLAB (MathWorks, Natick, MA), controlled by MATLAB-based

routines run on a PC running Ubuntu 18.04 or on a Mac Pro (Apple, Cupertino, CA).

The Datapixx device provided autonomously timed control of analog and digital inputs

and outputs synchronized to the visual display stimuli. A spout delivering liquid rewards

was placed near the mouth of the mouse and lick contacts on the spout were detected

by a piezo sensor using custom electronics. ~5 - 10 μL of 8% solution of infant formula

were used as rewards for correct task performance and were delivered via a peristaltic

pump.

Visual detection tasks

The detection task was modified from our previously published study (Wang et

al., 2018). Animals were run in daily sessions producing 153 to 421 trials, with data

pooled across sessions. Experiments were organized in blocks of randomly shuffled

trials. Each trial consisted of a sequence of epochs that were defined by the stimuli

presented on the visual displays. Epoch durations were determined by either the

randomized timing of the epoch or the time that it took for the mouse to travel a

randomized distance on the wheel. Mice were trained to detect a change in the

orientation of a Gabor patch visual stimulus, which the mouse reported by contacting

the lick spout to receive a reward.

The trial sequence of the primary detection task consisted of four epochs (Figure

5A). The average luminance across each visual display in all epochs was 4 – 8 cd/m^2.

In the first epoch ("Pink noise"), the uniform gray of the intertrial interval was changed to

pink visual noise with a root mean square contrast of 3.3%. In the second epoch

("Gabor patch cue") a vertically oriented Gabor patch was added to the pink noise,

centered on either the left or right visual display. The sinusoidal grating of the Gabor

patch (93.7% Michelson contrast) had a spatial frequency of 0.1 cycles per degree and

was modulated by a Gaussian envelope with a full width at half maximum of 18°

(standard deviation = 7.5°). The phase of the grating was incremented in proportion to

the wheel rotation and updated on every monitor refresh so that the sinusoidal pattern

displaced on the screen matched the distance that the mouse traveled on the wheel.

The Gabor patch on the left (right) drifted leftward (rightward), consistent with optic flow

during locomotion. The third epoch ("Delay epoch") consisted of vertical Gabor patches

appearing in both left and right visual displays. The stimuli in the fourth epoch depended

on whether the trial was a "Change" or "Catch" condition; the two trial types were

equally likely and randomly interleaved within a block. On Change trials, the Gabor

patch on the same side as the patch present in second epoch changed its orientation

(10-13°, depending on the mouse). The direction of the orientation change was

clockwise for the left patch and counterclockwise for the right patch. On catch no-

change trials, neither Gabor patch changed orientation, such that the epoch proceeded

as a seamless extension of the previous delay epoch.

Mice were to lick the spout upon detection of a change in the orientation of the

Gabor patch and otherwise withhold licking. Mice were required to lick within a 550 ms

response window starting 220 ms after the onset of the orientation change to score a

"hit" and receive a reward. If the mouse failed to lick within the window, the trial was

scored as a "miss" and no reward was given. On catch trials, if the mouse licked with

the same response window aligned on the no-change epoch onset, the trial was scored

as a "false alarm" and was used to calculate lick probability rates across trial types. If

the mouse correctly withheld licks during this window, the trial extended to include an

additional "Safety" epoch in which the initially appearing Gabor underwent a 30°

suprathreshold orientation change and the mouse could receive a reward by licking

within a comparable response window. This safety epoch was intended to maintain

motivation by rewarding mice for correct behavior without violating the task rule that

they should lick only for orientation changes. False alarms or two or more premature

licks before the response window resulted in termination of the trial and a three second

timeout.

In half of the trials, 465 nm blue light was delivered unilaterally to either the PS or

DMS using an LED system (Plexon). For half of these trials, light was delivered during

the delay epoch ("Early delay") for 150 ms, starting 200 ms after the onset of the epoch.

For the remaining 50% of stimulation trials, light was delivered during the fourth epoch,

either during catch ("Late delay") or change trials with equal frequency, for 150 ms,

starting 50 ms after the onset of the epoch. Light power was set at a fixed intensity per

striatal target per mouse to induce a non-saturating increase in lick probability (i.e., less

that 100%) and ranged from 0.05 to 1.15 mW across the ChR2-expressing cohort.

Similar light intensities were used for mice expressing the control mCherry fluorophore.

Trials were organized into alternating 40-trial blocks in which the orientation

change occurred for either the left or right Gabor patch. Each block had an equal

number of trials with and without orientation changes, and with and without optogenetic

light delivery, all randomly shuffled. Each session started with a mini-block of 20 trials

without light delivery trials which were excluded from behavioral analysis.

A "sequence arrested" control experiment consisted of blocks of 54 trials, 40 of

which followed the standard four-epoch trial sequence: 20 with an orientation change

(half of which were paired with optogenetic stimulation during the change) and 20 no-

change catch trials (half of which were paired with optogenetic stimulation during the

late delay catch epoch). Two additional trial types were introduced in which the visual

sequence of epochs stopped once the trial reached a particular epoch: seven stopped

at the single Gabor patch and seven stopped at the pink noise. These sequence-

arrested trials had the same durations and matched timing for optogenetic light delivery

as the standard four-epoch trial sequence. We recorded 2-3 sessions per mouse with

PS D1-MSN stimulation and pooled across sessions.

To test the interaction of PS D1-MSN stimulation on unreinforced change

detection responses, we conducted a version of the detection task with four trial types:

catch and change trials (50% of each), with stimulation during the change or late delay

on half of each. Instead of blocks of trials switching between left- and right-side

orientation changes, we only presented trials with the Gabor patch cue and orientation

change occurring in the visual field contralateral to the hemisphere receiving

optogenetic stimulation. After a 60-trial block in which correct change detection was

rewarded with soymilk rewards, the remaining trials were not reinforced, and the

session was terminated after the mouse extinguished lick responses. We recorded 2-3

sessions per mouse and pooled across sessions and mice for subsequent analysis.

In vivo electrophysiology

In the cohort of Drd1a-cre mice expressing DIO-ChR2-mCherry and optic fibers

in PS and DMS, we obtained electrophysiological recordings from isolated neurons in

the SC while mice performed the orientation change detection task with optogenetic

activation of PS and DMS D1-MSNs in the same hemisphere. The extracellular activity

of SC neurons was recorded with microwire bundles. Electrophysiological signals were

acquired through an RZ5D processor (Tucker-Davis Technologies, Alachua, FL) with

spikes filtered between 0.3 and 5 kHz and sampled at 25 kHz. Single units were sorted

offline using KiloSort (Pachitariu et al., 2016). The SC surface was determined based on

the first visual responses detected while advancing the microwire bundle. Sessions

alternated between optogenetic stimulation of either PS or DMS D1-MSNs.

Visual receptive field maps were obtained at the end of every recording session,

during which white circular disks (118.5 cd/m^2) of 10° in diameter were flashed against a

gray background (7.2 cd/m^2) in the visual display contralateral to the recording side. We

sampled visual locations pseudo-randomly drawn from a 3 x 7 isotropic grid that

extended from -25° to 25° in elevation and 0° to 90° in azimuth of the contralateral

visual field. An individual trial consisted of 8 consecutive 250 ms flashes, which flash

followed by a 250 ms blank interval. At least 9 flash repetitions for each grid location

were presented in each mapping session. Receptive fields of individual neurons were

estimated from the mean spike counts 50 – 150 ms after the flash onset in each grid

location, after subtracting baseline activity. The baseline in each trial was defined as the

mean spike count within the 100 ms period before the presentation of the first flash.

Baseline-subtracted mean spike counts were linearly interpolated between grid

locations with 1° resolution using the *scatteredInterpolant* function in Matlab.

Following the last recording session, we performed electrolytic lesions to aid in

histological reconstruction of bundle location within the SC. Under isoflurane

anesthesia, 80 µA current was delivered for 7-10 s to two wires per bundle. Mice were

administered meloxicam and perfused one week later.

Eye movement monitoring

To verify that changes in SC neuronal activity were not simply due to eye

movements, a 120Hz video eye tracking system (ETL-200, ISCAN, Woburn, MA) was

used to monitor eye position and pupil size of head-fixed mice during electrophysiology

experiments with optogenetic manipulation. The frequency of eye movements occurring

in trials with and without optogenetic D1-MSN stimulation per stimulation site, per

mouse, was statistically tested with chi-square tests.

Quantification

Behavioral analyses

Lick probability was quantified separately for contralaterally and ipsilaterally cued

trials and light delivery and no light delivery trials. Probability was calculated from the

number of trials in which there were any licks within a 750 ms window starting at onset

of light delivery for light delivery trials and a matched time window for non-light delivery

trials (i.e. 200 ms to 950 ms from onset of the early delay epoch and 50 ms to 800 ms

from the onset of the late delay catch epoch) divided by the total number of trials within

the assessed trial condition, per mouse. Reaction time was calculated from the first lick

following the orientation change per trial condition per mouse.

To compare behavioral extinction rates we separately fit binary responses (1 =

hit, 0 = miss) to the orientation change for non-stimulation and PS D1-MSN stimulation

trials using a four-parameter logistic regression model (Subramanian et al., 2023):

$$y = \max + \frac{min - max}{1 + (\frac{x}{thresh})^{slope}}$$

In which *x* is the trial number relative to reward non-reinforcement (1 = the first

unrewarded trial), *y* is the value of the function, *min* is the minimal value of the curve,

max is the value of y at the lowest value of x, *thresh* is the inflection point, and *slope* is

the slope of the function. The four parameter terms were fit to binary data by minimizing

mean squared error. For each data set, 95% CIs for the four parameters were estimated

using a standard bootstrap procedure with 1000 random draws. Significant differences

between the fitted curves for non-stimulation and PS stimulation trials were assessed

with z-tests for specific values of y (probability of a hit).

Neuronal analyses

Firing rates for each neuron were summarized as peri-stimulus time histograms

(PSTHs) by counting spikes within 20 ms non-overlapping bins aligned to key task

events. When pooling firing rates across neurons, these spike counts were converted to

normalized z-score values by subtracting the mean spike count from the PSTH and

dividing by the standard deviation, using the mean and standard deviation of the spike

counts calculated from the 20 ms bins within the trials across the entire recording

session. Population PSTHs presented in figures were generated from spike counts in 20

ms bins sliding in 1 ms intervals and normalized to z-scores, with error bars

representing standard error.

Phasic responses to the Gabor patch cue were tested by either Wilcoxon signed

rank test on spike counts in the interval 50 – 150 ms after Gabor patch cue epoch onset

to spike counts -100 – 0 ms before Gabor patch cue epoch onset on contralaterally

cued trials (relative to the recorded hemisphere) or Wilcoxon rank sum test on spike

counts in the interval 50 – 150 ms after Gabor patch cue epoch onset on contralaterally

cued trials and spike counts in the same interval for ipsilaterally cued trials. Delay-

related activity was tested by Wilcoxon signed rank test on spike counts in the interval

200 – 300 ms after delay epoch onset to spike counts in the final 100 ms of the same

epoch. Spatial cue -related modulation was tested by unpaired t-test of spike counts in

the last 200 ms of the delay epoch on contralaterally cued trials to that of ipsilaterally

cued trials. Phasic response to the orientation change was tested by Wilcoxon signed

rank test of spike counts in the interval 50 – 150 ms after change epoch onset to -100 –

0 ms before change epoch onset on contralaterally cued trials. Lick related activity was

tested by Wilcoxon signed rank test of spike counts in the interval -50 – 50 ms relative

to first detection lick during the change epoch to spike counts in the interval -100 to 0

ms prior to change epoch onset. Significant differences in the proportions of units

exhibiting task-related properties were assessed by chi-square tests.

SC neuronal responses to optogenetic direct pathway stimulation were tested by

Wilcoxon signed rank test on spike counts in the interval 0 – 150 ms relative to light

delivery to spike counts in a time-matched interval on non-stimulation trials and

manually validated. Responses in each stimulation epoch (early and late delay,

contralateral and ipsilaterally cued trials) were separately assessed for optogenetic

modulation.

Histological analysis

Fluorescence density maps were generated as followed. Micrographs were

converted to 8-bit and underwent contrast stretching to expand fluorescence intensity

range from 0 to 255 using ImageJ Fiji (v. 1.54k). Images were loaded into Python (v.

3.11) and pixel intensity was converted to a z-score value calculated from all pixel

values within the entire image of the micrographs of the SN or within the boundaries of

the SC. A 50 um grid was applied to each image. A threshold of 1 standard deviation

was then applied and for each 50 by 50 μm square, the pixels above this threshold were

summed and divided by the total number of pixels, then multiplied by 100 to produce a

percentage of pixels above threshold. This percentage is represented in grayscale,

ranging from 0 (white) to 100% (black).

<u>*Experimental design and statistical analyses*</u>

Sample sizes were determined based on publication standards. Studies were

completed across multiple cohorts that produced homogenous experimental findings.

For behavioral experiments, animals were randomly assigned to control and

experimental groups. Group assignment was also balanced to ensure equal numbers of

males and females.

Statistical tests were performed with Matlab (R2022a and R2023a) or GraphPad

Prism 10 (Boston, MA) software. Error bars in figures represent 95% CI or standard

error of the mean (s.e.m.) as indicated in figure legends. 95% CIs in figure plots were

generated with the *binofit* Matlab function. The value of N represents the number of

animals and n represents the number of units. All statistical tests were conducted two-

sided, when applicable. Post-hoc Holm-Ŝidàk tests were performed to correct for

multiple comparisons for ANOVAs; significant test results are indicated in the figures. All

asterisks in figures indicate: * $p<0.05$, ** $p<0.01$, *** $p<0.001$, and **** $p<0.0001$.

420   Effect sizes were calculated for significant statistical comparisons. The eta

squared ($\eta2=SS_{Effect}/SS_{Total}$) was calculated for 1-way ANOVA and the partial eta

squared (partial $\eta2=SS_{Effect}/(SS_{Error}+ SS_{Effect})$) was calculated for N-way ANOVA main

effects and interactions, where SS = sum of squares. Cohen's d was calculated for post-

hoc pairwise comparisons (d = mean difference/pooled standard deviation). For

comparisons assessed with Chi-square tests, Cohen's d was calculated from the chi

square test statistic and sample sizes (n), as follows: $d = 2\sqrt{\frac{\chi^2}{(n1+n2)-\chi^2}}$.

427
428   **Results**
429
430   _The posterior striatum provides a unique basal ganglia direct pathway for visual signals_

431   The rodent striatum is canonically divided into dorsomedial, dorsolateral, ventral,

and posterior (or tail) subregions. D1-MSNs populate all regions of the striatum and

provide the origin of the direct pathway of the basal ganglia. We sought to examine

whether the direct pathway originating from the PS is anatomically distinct or shared

with that of the DMS. To do this, we expressed cre-dependent anterograde fluorophore -

expressing viruses in the PS and DMS of Drd1a-cre heterozygous mice (Figure 1). We

found that DMS labeled D1-MSNs terminated in the ventral medial segment of the

substantia nigra (SN) and in the entopeduncular nucleus (EP; homolog to the primate

internal segment of the globus pallidus). In contrast, PS labeled D1-MSNs innervated

the dorsal lateral region of the SN, with no labeling in the EP.

We then used a viral strategy to selectively label the neurons in the SN that are

postsynaptic to PS and DMS D1-MSNs. We injected anterograde trans-synaptic AAV1

viruses in the PS and DMS of wild-type mice that delivered cre or flp recombinases to

all efferent cells. We then co-injected cre- and flp-dependent non-trans-synaptic

fluorophore-expressing viruses in the SN to locally label all recombinase-expressing

cells putatively postsynaptic to D1-MSNs (Figure 2). This strategy selectively targets

D1-MSNs because D2-MSNs do not project to SN (Lein et al., 2007; Kreitzer and

Malenka, 2008; Allen Institute for Brain Science, 2025a, 2025b). We found neurons

targeted by PS D1-MSNs were largely confined to the substantia nigra pars lateralis

(SNl) and adjacent peripeduncular nucleus (PP) (Figure 2B). In contrast to the PS,

neurons targeted by DMS D1-MSNs resided nearly exclusively in the substantia nigra

pars reticulata (SNr) and in the overlying substantia nigra pars compacta (SNc) (Figure

2C). The presence of labeled cells in the SNc, a region that appeared to be sparsely

innervated along the SNr border from DMS D1-MSN axons (Figure 1F), may be a result

of retrograde labeling of striatal afferents which is noted to occur in recombinase-

expressing AAV1 viruses (Zingg et al., 2020). Density mapping of fluorescent

expression further demonstrated a general segregation of neuronal populations

innervated by either PS or DMS (Figure 2D-E). By following the axonal expression of

the neurons targeted by the PS and DMS D1-MSNs, we found a mixture of both

exclusive and shared output targets in the diencephalon and midbrain (Figure 2F-L, O-

P). To characterize the second direct pathway originating from the DMS and determine

its efferent targets, in a separate set of cases we applied this di-synaptic labeling

approach to confirm the projection from DMS D1-MSNs to neurons in the EP, with axons

that innervate several midbrain nuclei and the habenula (Figure 3).

Although direct pathways originating from both DMS and PS innervate the SC,

we observed striking differences. DMS-originating terminals showed concentrated

innervation of SC intermediate layers (Figure 2N,R; Figure 3D-F), whereas PS-

originating terminals exhibited diffuse innervation across all layers with relatively greater

expression laterally and in more caudal sections of the SC (Figure 2M,Q).

Having established that D1-MSNs in the PS and DMS give rise to anatomically

distinct pathways through the basal ganglia, we next compared the extent of visual

cortical innervation to these two striatal subregions. Prior work indicates that the PS and

DMS are the exclusive striatal targets of visual cortex (Hunnicutt et al., 2016). To identify

the regions of visual cortex that project to these striatal regions, we injected the

retrograde tracer Fluorogold unilaterally in either the PS or DMS of wild-type mice. We

observed strong labeling of PS-projecting cells in what we identified as layers 5 and 6

across the entire rostral-caudal axis of the visual cortex, both in primary and secondary

visual areas (Figure 4A-D). In contrast, we found strong labeling of DMS-projecting cells

in layer 5 in rostral visual cortex but the labeling was weaker in more caudal sections,

especially for primary visual cortex (Figure 4E-H). We also observed weak labeling of

putative layer 3 cells for both striatal regions (PS, Figure 4B; DMS, Figure 4F). These

results agree with similar findings in rats demonstrating stronger innervation from the

visual cortex and thalamus to the PS than to the DMS (Jiang and Kim, 2018).

Collectively these experiments establish the PS as a distinct region within the

striatum for influencing visual behavior. Unlike the DMS, the PS is strongly innervated

by multiple layers through the entirety of visual cortex. By routing through the SNI and

PP, the PS initiates a direct pathway largely segregated from that of the DMS. Lastly,

we identified three distinct circuits by which the DMS and PS may modulate

downstream targets, including the superior colliculus.

Stimulation of the posterior striatum biases visual decisions

Our neuroanatomical results establish that the PS direct pathway provides a

major route for visual signals through the basal ganglia that is distinct from the circuits

emanating from the DMS. We next probed for functional roles of this striatal region by

optogenetically activating PS D1-MSNs in mice performing a spatially cued orientation

change-detection task (Wang and Krauzlis, 2018; Wang et al., 2018). Briefly, head-fixed

mice were presented a vertical Gabor patch cue in one visual field that drifted at a rate

coupled to running speed (Figure 5A). As the trial proceeded, a second Gabor patch

appeared in the opposite visual field. After a variable time, the patch that had appeared

first either changed orientation or remained vertical ('catch'; 50% occurrence for each).

Mice were trained to report their detection of the orientation change by licking a centrally

placed spout to receive a soymilk reward and otherwise suppressed licks during other

trial epochs (Figure 5B). Sessions were run in 40-trial blocks in which the orientation

changed occurred exclusively in one visual field (left or right) before switching to the

other location.

We unilaterally stimulated channelrhodopsin (ChR2)-mCherry -expressing PS or

DMS D1-MSNs in Drd1a-cre mice during one of three epochs of the task (Figure 5C-D).

On 50% of the trials, 150 ms blue light was delivered either shortly after the appearance

of the second Gabor patch ('early delay'), during the change event ('change'), or at a

time-matched period on catch trials ('late delay'). The visual conditions were identical

between the early delay and late delay conditions (the presence of two vertical Gabor

patches). We assessed lick responses within 750 ms of light delivery onset (or an

equivalent time window on no-stimulation trials).

Light delivery to the PS during the early delay produced a small but significant

increase in lick probability ('yes' response) in ChR2-mCherry-expressing mice (N = 6)

compared to non-stimulation trials (Figure 5E-F; Table 1). Delivering light to the PS of

mice expressing a control mCherry fluorophore (mCherry; N = 4) did not result in

significant changes in lick probability (Figure 5F; Table 1). Delivering light during the late

delay (in the absence of an orientation change) induced a larger significant increase in

lick probability for ChR2-mCherry mice (Figure 5E,G, Table 1). Again, the control

mCherry mice showed no effect (Figure 5G; Table 1). Comparing the effects across the

factors, repeated measures 4-way ANOVA revealed significant main effects for epoch,

light delivery, and group, but not cued side, and included several significant interactions

across factors (Table 1). In particular, we found significant interactions between group

and all other factors (light delivery, cued side, and delay epoch). Thus, our main finding

with optogenetic PS D1-MSN stimulation during the delay period of the task was that

light-evoked increases in lick probability were significantly greater during the late delay

than the early delay.

For light delivered to the PS during the orientation-change event, because

performance was already at ceiling, it was not possible to observer increases in lick

rate. However, we did find effects on lick reaction time. Delivering light to the PS during

the change event significantly reduced the median response latency in ChR2-mCherry

mice but not in mCherry control mice (Figure 5H; Table 2). A repeated measures 3-way

ANOVA showed a significant main effect of light delivery, but not of group nor cued side

(Table 2).

Comparison with stimulation of the dorsomedial striatum

In the same mice, we also tested the effects of DMS direct pathway stimulation

during the task. Our group previously demonstrated that DMS D1-MSN activation

biases lick responses in a manner dependent on spatial cues and reward

conditions(Wang et al., 2018; Wang and Krauzlis, 2020). Here, to compare with our PS

activation, we tested whether temporal expectation also influenced the behavioral

effects of DMS direct pathway activation. We found that light delivery to the DMS during

the early delay did not significantly change lick probability for either ChR2-mCherry mice

or mCherry control mice (Table 1). Light delivery to the DMS during the late delay did

significantly increase lick probability in ChR2-mCherry mice but not mCherry control

mice (Table 1). As with PS, a repeated measures 4-way ANOVA revealed significant

main effects for epoch, light delivery, and group, but not cued side, and included several

significant interactions (Table 1). Our group has previously observed an effect of spatial

cueing on DMS-evoked licking (Wang et al., 2018; Wang and Krauzlis, 2020). The lack

of effect in the present cohort may be due to a smaller group size, as not all mice

demonstrate cueing effects, even in the original studies.

We also examined the effects of DMS direct pathway stimulation on the response

to the change event. Unlike our results with PS stimulation, we did not find a significant

effect of light delivery on median response latency for ChR2-mCherry mice or mCherry

mice (Table 2). A repeated measures 3-way ANOVA revealed a trending but insignificant

main effect of light delivery (Table 2). Our group has previously demonstrated that

activating DMS D1-MSNs significantly reduces event detection latency (Wang et al.,

2018); the negative result here may be due to an underpowered cohort size.

Thus, stimulation of either the PS or DMS biased decisions in our visual change-

detection task, and both showed larger effects when the stimulation was applied late in

the delay period, when the change event was expected to occur. These results show

that stimulation of either PS or DMS does not automatically evoke a lick response but

instead depends on the context and temporal structure that the mice have learned to

associate with rewards in our visual task.

Effects of stimulating PS depend on the visual context and reward reinforcement

To further elucidate the conditions in which PS direct pathway activation biases

visual decisions we performed two additional behavioral experiments in the same

cohort. First, we tested whether the induced lick response depended on the presence of

choice-relevant visual stimuli (i.e. the Gabor patch). Using our standard visual change-

detection task, we introduced a minority of trials in which the stimulation was paired with

either the one-patch or pink noise stimuli, but at a time temporally matched to

stimulation during the late delay or change epoch (Figure 6A *left*). This experiment

replicated our earlier finding that stimulation during contralaterally cued two-patch visual

presentation increased the mean lick probability (no light: 0.067 ± 0.034 s.e.m. vs. light:

0.491 ± 0.044). In contrast, pairing PS D1-MSN activation with the contralateral one-

patch stimulus resulted in a lick probability of 0.249 ± 0.054, approximately half as much

as the two-patch, and light delivery paired with pink noise produced a lick probability of

0.018 ± 0.012, essentially zero (Figure 6A *right*). A repeated measures 1-way ANOVA

showed a significant main effect of visual stimulus ($F(3,15) = 40.17$, $P < 0.0001$, $d =$

4.23) on lick probability. Thus, the effects of PS stimulation were not simply due to

timing in the task but depended on visual and behavioral context. Evoked-lick rates

were highest for the two-patch presentation, which is the context associated with the

possibility of a reward-related visual event.

We next asked whether the decision bias mediated by PS stimulation also

depended on reward reinforcement. Specifically, if rewards were withheld for correct

responses in our visual task, would PS D1-MSN activation still evoke licks, or otherwise

delay the emergence of operant extinction? To test this, we ran mice on a modified

version of the task in which only contralaterally cued trials were presented (50%

resulted in the change event; 50% of both trial types were paired with PS D1-MSN

stimulation). After 60 trials in which correct detection responses were rewarded, trials

continued as before but with no rewards administered for the remainder of the session

(Figure 6B *left*). Pooling across 10 sessions from 4 mice, we quantified the extinction

rates separately for non-stimulation and PS D1-MSN stimulation trials by fitting the hits

and misses relative to the first non-rewarded trial in a four-parameter logistic regression

model with bootstrapped computed 95% confidence intervals (CIs) (Figure 6B *right*). We

found that the fitted model curve for stimulation trials did not significantly differ from that

of non-stimulation trials. A predicted hit rate of 0.5 occurred at 8.9 trials (95% CIs: 5,

11.8) post- reward non-reinforced on the curve fitted to the non-stimulation trials and 5.8

trials (CIs: 1.4, 9.1) on the curve fitted to stimulation trials (z-test z = -1.673, P = 0.094).

Thus, PS stimulation did not change the time course of operant extinction, and the

effects of PS stimulation disappeared as extinction took hold.

The basal ganglia direct pathway modulates superior colliculus neuronal activity during

visual decision-making

Having demonstrated that both PS and DMS D1-MSN activation robustly affects

visual decisions in our task, we next investigated one of the major output targets of

these basal ganglia circuits and how neuronal activity was modulated during the visual

detection task. The SC is a shared target of both PS- and DMS-originating direct

pathways (Figure 2K-R, Figure 3C-F) and plays a crucial role in visual processing and

visually guided decisions in rodents (Cang et al., 2018; Wang et al., 2020). We collected

extracellular electrophysiological recordings in the SC of our cohort of PS and DMS D1-

MSN ChR2-mCherry expressing mice (N = 6) performing the orientation change-

detection task with and without optogenetic stimulation. PS or DMS D1-MSN activation

was tested in separate recording sessions.

We recorded a total of 530 single units across the layers of the SC (Figure 7A-B).

Grouping these units by their estimated location in either superficial (sSC) or deeper

(dSC) layers, we observed neuronal responses to key features of the spatially cued

change-detection task (Figure 7C-D; Table 3). Phasic responses to the onset of the cue

patch, when presented in the contralateral visual field, were present in the majority of

SC neurons: 78.4% of sSC units and 55.7% of dSC units. Delay activity (defined as a

significant difference in firing rates in the late vs early delay epochs) was also observed

in many neurons: 46.6% of sSC units and 37.3% of dSC units. We observed spatial

cue-related modulation (defined as the difference in firing rate between contra- and ipsi-

laterally cued trials late in the delay epoch) but this was present in a minority of

neurons: 18.2% of sSC units and 20.6% of dSC units. Similarly, phasic responses to the

orientation change were observed in 18.2% of sSC units and 22.4% of dSC units.

Lastly, we found lick-related activity (defined as the difference in firing rate in a 100 ms

window centered on the first detection lick compared to pre-change event firing activity)

in a small minority of SC neurons: 1.1% of sSC units and 6.6% of dSC units. Comparing

the modulation between the superficial and deep layers, we found that sSC units were

significantly more likely to have a cue response than dSC units ($\chi^2 = 15.76$, P < 0.0001,

d = 0.35) whereas dSC units were significantly more likely to have lick-related activity

($\chi^2 = 4.04$, P = 0.044, d = 0.18).

We next examined how optogenetic direct pathway activation modulated

ipsilateral SC neuronal activity. We classified units as modulated by D1-MSN activation

if there was a significant difference in firing during the 150-ms light-delivery period

compared to an equivalent window during non-stimulation trials. Because the basal

ganglia direct pathway reaches the SC by a series of two inhibitory projections, we

expected activation of D1-MSNs to have a predominately excitatory effect on SC

neuronal activity (Hikosaka and Wurtz, 1983; Hikosaka et al., 2000; Kaneda et al., 2008;

Freeze et al., 2013). Indeed, the majority of units modulated by PS or DMS direct

pathway activation showed excitatory changes in firing rate (Figure 8A-B *left*). However,

we also observed populations of units that were exclusively inhibited by direct pathway

activation (Figure 8A-B *middle*). This effect was more prevalent for PS. Stimulation

induced an inhibitory effect in a larger percentage of dSC units modulated by PS

(27.5%; 15.9, 41.7 95% CI) than modulated by DMS (13.3%; 6.8, 22.5; χ^2 = 4.20, P =

0.04, d = 0.36). Finally, we observed a small number of units that demonstrated both

excitation and inhibition, either in different test epochs or within the same epoch (Figure

8A-B *right*).

We examined how SC units that exhibited different task-related properties were

modulated by direct pathway stimulation. For the sSC, we found very few units to be

modulated by either PS (4 out of 35 units) or DMS (3 out of 45 units) D1-MSN

activation. Therefore, we restricted our analyses to dSC units, among which 51 out of

186 units were modulated by PS direct pathway activation and 83 out of 228 units were

modulated by DMS direct pathway activation. We observed that activation of the striatal

direct pathways did not affect all dSC units equally. For the PS, we found that D1-MSN

activation was more likely to modulate units that exhibited delay activity (χ^2 = 4.46, P =

0.035, d = 0.31), showed spatial cue-related modulation (χ^2 = 5.92, P = 0.015, d = 0.36),

or responded to the orientation change (χ^2 = 16.74, P < 0.0001, d = 0.63) (Figure 8C

left; Table 3). Similarly, for the DMS, D1-MSN activation preferentially modulated units

with delay activity (χ^2 = 9.52, P = 0.002, d = 0.42), spatial cue-related modulation (χ^2 =

12.66, P = 0.0004, d = 0.48), or that responded to the orientation change (χ^2 = 17.89, P

< 0.0001, d = 0.58). DMS D1-MSN activation was also more likely to modulate units that

did not show lick related activity (χ^2 = 6.60, P = 0.0102, d = 0.35) (Figure 8C *right*; Table

3). However, we note that we had relatively few units with lick-related activity in our

recordings, likely because most of our recording sites were in the central SC, rather

than the lateral SC, where lick-related units tend to be concentrated (Thomas et al.,

2023).

We also examined whether dSC modulation by direct pathway activation

depended on the stimulated epoch. For PS stimulation, we found that approximately

half of D1-MSN modulated dSC units were modulated during both the early and late

delay epochs (54.9%), whereas 39.2% were exclusively modulated during the early

delay and only a small percentage were modulated exclusively during the late delay

(5.9%) (Figure 8D). dSC units modulated by DMS D1-MSN stimulation showed similar

distributions: 48.2% for both epochs, 45.8% exclusively during the early delay, and 6.0%

exclusively during the late delay (Figure 8D).

Direct comparison of PS and DMS activation on SC unit activity

We also were able to directly compare the effects of PS and DMS direct pathway

activation on SC units. This comparison at the neuronal level is of particular interest

because our histological tracing of PS and DMS direct pathways targeting the SC

showed distinct but overlapping patterns of axonal innervation in the SC. We

investigated whether these pathways target the same neuronal populations within the

SC. In two mice, we recorded from SC units while stimulating PS or DMS D1-MSNs

during the early delay in the same session. We recorded a total of 30 optogenetically

modulated single units: 40% were exclusively modulated by DMS direct pathway

activation, 20% were exclusively modulated by PS direct pathway activation, and 40%

were modulated by both pathways (Figure 8E). Of the units that were modulated by

both direct pathways, 58.3% exhibited different modulatory effects, that is, these units

were excited by one pathway and inhibited by the other. This finding suggests that the

PS and DMS direct pathways may innervate different SC populations.

Discussion

The present study investigated the contributions of the mouse PS to visual

behavior. Our results demonstrate that the PS is uniquely anatomically positioned to

relay visual signals through the basal ganglia. Compared to the DMS, which has been

studied more frequently in visual tasks, the PS is more strongly innervated by visual

cortex and gives rise to an alternative direct pathway for visual processing. Our results

show that activating this pathway robustly and selectively modulates visual decisions.

We also identified that both PS and DMS direct pathway activation targeted neurons in

the SC that were sensitive to context and reward-associated stimuli. Together, these

findings further demonstrate distinct circuits through the basal ganglia for visual

perceptual decision-making.

The effects of PS direct pathway manipulation on visual decisions were similar to

that of the DMS (Wang et al., 2018), suggesting a comparable influence on perceptual

behavior, at least as revealed in our visual task. Similar to the DMS, we found that

stimulating the PS direct pathway does not drive an obligatory and fixed lick response.

Optogenetic activation induced 'yes' perceptual responses that depended on visual and

temporal context. Moreover, PS D1-MSN stimulation did not delay extinction of the

response to the devalued change event when rewards were withheld. These findings

are consistent with the view that the direct pathway provides a permissive gating

function for actions driven by circuits extrinsic to the basal ganglia (Hormigo et al.,

2021). In the context of perceptual decision-making, the convergence of cortical and

subcortical inputs to the striatum is proposed to similarly enable the selection and

implementation of visual-based decision policies (Ding, 2023).

The PS direct pathway has also been shown to bias auditory-based decisions

(Guo et al., 2018; Chen et al., 2022). Dopaminergic signaling in this region reinforces

the avoidance of novel or threatening stimuli of multiple sensory modalities (Menegas et

al., 2018) and is also shown to scale with auditory uncertainty (Chen et al., 2022).

Moreover, activating PS D1-MSNs in freely moving mice failed to induce orienting

responses which are reliably observed with dorsal striatal direct pathway activation

(Guo et al., 2018). The combination of sensitivity to multimodal stimuli and distinct

dopaminergic activity positions the PS to uniquely contribute to decision-making and

behavior.

Although PS and DMS stimulation yielded comparable behavioral effects in our

paradigm, differences in the connectivity of these striatal subregions hint at potential

functional specializations, starting with the sources of their cortical afferents. The DMS

was preferentially innervated by anterior primary and secondary visual areas. Neurons

in mouse anterior secondary areas are noted to preferentially respond to stimuli of lower

spatial and higher temporal frequencies, exhibit stronger orientation selectivity, and

prefer fast-moving stimuli (Han et al., 2022). In contrast, we found that the PS was

innervated by all of secondary visual cortex, including posterior areas that are

characterized by a preference for higher spatial, lower temporal, and slow-moving

stimuli (Han et al., 2022). Whether this broader anatomical connection confers the PS a

relative sensitivity or contextual selectively for visual input remains to be determined.

Differences in striatal afferents extend beyond the visual cortex. Whereas the DMS is

preferentially innervated by frontal cortices including anterior cingulate, prelimbic and

orbital cortices, the PS notably receives input from all sensory cortical areas (Hintiryan

et al., 2016; Hunnicutt et al., 2016; Lee et al., 2023).

There are also differences in the efferent pathways from the PS and DMS, giving

rise to distinct circuits through the basal ganglia that target different topographical

subregions of the output nuclei (Lee et al., 2020; Miyamoto and Fukuda, 2022). The

DMS direct pathway is routed through the SNr and EP, whereas the PS direct pathway

innervates the SNl and PP. Although the specific functions of the SNl and PP are not yet

well defined, our findings of robust behavioral effects induced by PS D1-MSN activation

indicate that these nuclei should be considered among the basal ganglia nuclei involved

in perceptual decision-making. Previous studies of the SNl and PP that have examined

the anatomical properties of these nuclei suggest a range of possible circuit

mechanisms. The SNl consists of dopaminergic and GABAergic neurons that

differentially project to sensory nuclei, motor-related regions, the amygdala, as well as

forming loops back to the basal ganglia (Kaelber and Afifi, 1979; Moriizumi et al., 1992;

Takada, 1992; Todd et al., 2022). The adjacent PP innervates a range of midbrain,

cortical, and diencephalon nuclei (Arnault and Roger, 1987; Cai et al., 2024).

Given that the PS and DMS route through different output nuclei, it is not

surprising that these two striatal regions yield divergent efferent connectivity in the

thalamus and midbrain (Benavidez et al., 2021; McElvain et al., 2021). We found

marked differences in the innervation of one prominent target, the SC. Compared to the

DMS-originating direct pathways, the PS to SNl/PP circuit exhibited preferential

innervation of the lateral SC, which is notably connected with cortical, thalamic, and

hindbrain nuclei associated with visuomotor and somatic sensorimotor functions

(Benavidez et al., 2021). In contrast, the direct pathways initiated from the DMS did not

show medial-lateral differences in SC innervation, and this more uniform innervation of

SC deeper layers suggests engagement with additional anatomical networks including

those involving the dorsal lateral geniculate nucleus as well as secondary visual and

association cortical areas (Benavidez et al., 2021).

Related to these different efferent circuit projections, we also identified

differences in how activation of the direct pathways alters neuronal activity in the SC.

We observed that PS direct pathway activation induced inhibition in a relatively larger

population of dSC units. Given the established disinhibitory effect of the direct pathway,

excitation was the expected, and indeed, predominate modulatory effect on SC unit

firing. However, suppression may occur if the recorded SC unit is efferent to a local

interneuron that is itself disinhibited by the direct pathway resulting in a tri-synaptic

inhibitory configuration (e.g., D1-MSN→SN/EP→SC interneuron→recorded SC unit). A

larger proportion of SC units inhibited by PS D1-MSN activation suggests that the

innervation of the SC by this direct pathway may be more biased toward interneurons.

We found that both PS and DMS direct pathways preferentially modulated dSC

units with specific types of task-related activity. Units that exhibited delay-related activity,

spatial cue-related modulation, or phasically responded to the reward-associated

orientation change were more likely to be modulated by direct pathway activation.

Interestingly, we did not find a similar preference for units that responded to the

appearance of the Gabor patch cue. This suggests that the direct pathway bias is not

merely for visually responsive SC units, but for those processing aspects of visual

signals related to a goal or outcome. Our group has previously shown that dSC cue-

related modulation contributes to spatial attention by biasing signal processing for

reward-associated event locations (Wang et al., 2022). Preferential innervation of

neurons showing attention-like activity may be one mechanism by which the direct

pathway biases decisions in manner linked to the allocation of spatial attention (Wang

and Krauzlis, 2020).

In summary, the results of the present study provide the first evidence for

involvement of the mouse PS in visual behavior. Here, we monitored the neuronal

modulation of one target of the basal ganglia direct pathway, the SC, during a visual

decision-making task. In addition to the several circuit routes comprising the direct

pathway that we have already noted, there are also many output targets that may

contribute to perceptual decisions. The basal ganglia direct pathways also target the

motor thalamus, reticular formation, and pedunculopontine nucleus, which may each

also contribute to the behavioral response (van der Kooy and Carter, 1981; McElvain et

al., 2021). Although it remains an open challenge to understand how these complex

circuits operate together for visual perception, our results show that any description

should include circuits that originate in the posterior striatum.

References

Allen Institute for Brain Science (2025a) Allen Mouse Brain Connectivity Atlas
[#155737254]. Available at: https://connectivity.brain-
map.org/projection/experiment/155737254.

Allen Institute for Brain Science (2025b) Allen Mouse Brain Connectivity Atlas
[#175732996]. Available at: https://connectivity.brain-
map.org/projection/experiment/175732996.

Amita H, Kim HF, Inoue K-I, Takada M, Hikosaka O (2020) Optogenetic manipulation of
a value-coding pathway from the primate caudate tail facilitates saccadic gaze
shift. Nat Commun 11:1876.

Amita H, Kim HF, Smith MK, Gopal A, Hikosaka O (2019) Neuronal connections of
direct and indirect pathways for stable value memory in caudal basal ganglia. Eur
J Neurosci 49:712–725.

Arcizet F, Krauzlis RJ (2018) Covert spatial selection in primate basal ganglia. PLoS
Biol 16:e2005930.

Arnault P, Roger M (1987) The connections of the peripeduncular area studied by
retrograde and anterograde transport in the rat. J Comp Neurol 258:463–476.

Benavidez NL et al. (2021) Organization of the inputs and outputs of the mouse superior
colliculus. Nat Commun 12:4004.

Brainard DH (1997) The Psychophysics Toolbox. Spat Vis 10:433–436.

Cai H-R, Chen S-Q, Xiang X-J, Zhang X-Q, Ma R-Z, Zhu G, Ding S-L (2024)
Comparison of the connectivity of the posterior intralaminar thalamic nucleus and
peripeduncular nucleus in rats and mice. Front Neural Circuits 18:1384621.

Cang J, Savier E, Barchini J, Liu X (2018) Visual Function, Organization, and
Development of the Mouse Superior Colliculus. Annu Rev Vis Sci 4:239–262.

Chen APF, Malgady JM, Chen L, Shi KW, Cheng E, Plotkin JL, Ge S, Xiong Q (2022)
Nigrostriatal dopamine pathway regulates auditory discrimination behavior. Nat
Commun 13:5942.

Ding L (2023) Contributions of the Basal Ganglia to Visual Perceptual Decisions. Annu
Rev Vis Sci 9:385–407.

Ding L, Gold JI (2010) Caudate encodes multiple computations for perceptual decisions.
J Neurosci Off J Soc Neurosci 30:15747–15759.

Eastman KM, Huk AC (2012) PLDAPS: A Hardware Architecture and Software Toolbox
for Neurophysiology Requiring Complex Visual Stimuli and Online Behavioral

Control. Front Neuroinformatics 6 Available at:
http://journal.frontiersin.org/article/10.3389/fninf.2012.00001/abstract [Accessed
July 11, 2024].

Franklin KBJ, Paxinos G (2008) The mouse brain in stereotaxic coordinates, 3. ed.
Amsterdam: Elsevier, AP.

Freeze BS, Kravitz AV, Hammack N, Berke JD, Kreitzer AC (2013) Control of basal
ganglia output by direct and indirect pathway projection neurons. J Neurosci Off J
Soc Neurosci 33:18531–18539.

Griggs WS, Kim HF, Ghazizadeh A, Costello MG, Wall KM, Hikosaka O (2017) Flexible
and Stable Value Coding Areas in Caudate Head and Tail Receive Anatomically
Distinct Cortical and Subcortical Inputs. Front Neuroanat 11:106.

Guo L, Walker WI, Ponvert ND, Penix PL, Jaramillo S (2018) Stable representation of
sounds in the posterior striatum during flexible auditory decisions. Nat Commun
9:1534.

Han X, Vermaercke B, Bonin V (2022) Diversity of spatiotemporal coding reveals
specialized visual processing streams in the mouse cortex. Nat Commun
13:3249.

Hikosaka O, Kim HF, Amita H, Yasuda M, Isoda M, Tachibana Y, Yoshida A (2019) Direct
and indirect pathways for choosing objects and actions. Eur J Neurosci 49:637–
645.

Hikosaka O, Takikawa Y, Kawagoe R (2000) Role of the basal ganglia in the control of
purposive saccadic eye movements. Physiol Rev 80:953–978.

Hikosaka O, Wurtz RH (1983) Visual and oculomotor functions of monkey substantia
nigra pars reticulata. IV. Relation of substantia nigra to superior colliculus. J
Neurophysiol 49:1285–1301.

Hintiryan H, Foster NN, Bowman I, Bay M, Song MY, Gou L, Yamashita S, Bienkowski
MS, Zingg B, Zhu M, Yang XW, Shih JC, Toga AW, Dong H-W (2016) The mouse
cortico-striatal projectome. Nat Neurosci 19:1100–1114.

Hormigo S, Zhou J, Chabbert D, Shanmugasundaram B, Castro-Alamancos MA (2021)
Basal Ganglia Output Has a Permissive Non-Driving Role in a Signaled
Locomotor Action Mediated by the Midbrain. J Neurosci Off J Soc Neurosci
41:1529–1552.

Hunnicutt BJ, Jongbloets BC, Birdsong WT, Gertz KJ, Zhong H, Mao T (2016) A
comprehensive excitatory input map of the striatum reveals novel functional
organization. eLife 5:e19103.

Jiang H, Kim HF (2018) Anatomical Inputs from the Sensory and Value Structures to the
Tail of the Rat Striatum. Front Neuroanat 12:30.

Kaelber WW, Afifi AK (1979) Efferent connexions of the pars lateralis of the substantia
nigra (SNL). J Anat 129:405–412.

Kamishina H, Yurcisin GH, Corwin JV, Reep RL (2008) Striatal projections from the rat
lateral posterior thalamic nucleus. Brain Res 1204:24–39.

Kaneda K, Isa K, Yanagawa Y, Isa T (2008) Nigral inhibition of GABAergic neurons in
mouse superior colliculus. J Neurosci Off J Soc Neurosci 28:11071–11078.

Khibnik LA, Tritsch NX, Sabatini BL (2014) A direct projection from mouse primary visual
cortex to dorsomedial striatum. PloS One 9:e104501.

Kim HF, Amita H, Hikosaka O (2017) Indirect Pathway of Caudal Basal Ganglia for
Rejection of Valueless Visual Objects. Neuron 94:920-930.e3.

Krauzlis RJ, Nichols N, Rangarajan KV, McAlonan K, Goldstein S, Yochelson D, Wang L
(2020) Visual Psychophysics in Head-Fixed Mice. Curr Protoc Neurosci 92:e95.

Kreitzer AC, Malenka RC (2008) Striatal plasticity and basal ganglia circuit function.
Neuron 60:543–554.

Lee J, Wang W, Sabatini BL (2020) Anatomically segregated basal ganglia pathways
allow parallel behavioral modulation. Nat Neurosci 23:1388–1398.

Lee K, An S-Y, Park J, Lee S, Kim HF (2023) Anatomical and Functional Comparison of
the Caudate Tail in Primates and the Tail of the Striatum in Rodents: Implications
for Sensory Information Processing and Habitual Behavior. Mol Cells 46:461–
469.

Lein ES et al. (2007) Genome-wide atlas of gene expression in the adult mouse brain.
Nature 445:168–176.

McElvain LE, Chen Y, Moore JD, Brigidi GS, Bloodgood BL, Lim BK, Costa RM,
Kleinfeld D (2021) Specific populations of basal ganglia output neurons target
distinct brain stem areas while collateralizing throughout the diencephalon.
Neuron 109:1721-1738.e4.

Menegas W, Akiti K, Amo R, Uchida N, Watabe-Uchida M (2018) Dopamine neurons
projecting to the posterior striatum reinforce avoidance of threatening stimuli. Nat
Neurosci 21:1421–1430.

Miyamoto Y, Fukuda T (2022) New Subregions of the Mouse Entopeduncular Nucleus
Defined by the Complementary Immunoreactivities for Substance P and
Cannabinoid Type-1 Receptor Combined with Distributions of Different Neuronal
Types. eNeuro 9:ENEURO.0208-22.2022.

Moriizumi T, Leduc-Cross B, Wu JY, Hattori T (1992) Separate neuronal populations of
the rat substantia nigra pars lateralis with distinct projection sites and transmitter
phenotypes. Neuroscience 46:711–720.

Pachitariu M, Steinmetz N, Kadir S, Carandini M, Harris K (2016) Fast and accurate
spike sorting of high-channel count probes with KiloSort. Adv Neural Inf Process
Syst 29.

Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming
numbers into movies. Spat Vis 10:437–442.

Permezel F, Alty J, Harding IH, Thyagarajan D (2023) Brain Networks Involved in
Sensory Perception in Parkinson's Disease: A Scoping Review. Brain Sci
13:1552.

Profant O, Roth J, Bureš Z, Balogová Z, Lišková I, Betka J, Syka J (2017) Auditory
dysfunction in patients with Huntington's disease. Clin Neurophysiol Off J Int Fed
Clin Neurophysiol 128:1946–1953.

Smith Y, Parent A (1986) Differential connections of caudate nucleus and putamen in
the squirrel monkey (Saimiri sciureus). Neuroscience 18:347–371.

Subramanian D, Pearson JM, Sommer MA (2023) Bayesian and Discriminative Models
for Active Visual Perception across Saccades. eNeuro 10:ENEURO.0403-
22.2023.

Tai L-H, Lee AM, Benavidez N, Bonci A, Wilbrecht L (2012) Transient stimulation of
distinct subpopulations of striatal neurons mimics changes in action value. Nat
Neurosci 15:1281–1289.

Takada M (1992) The lateroposterior thalamic nucleus and substantia nigra pars
lateralis: Origin of dual innervation over the visual system and basal ganglia.
Neurosci Lett 139:153–156.

Thomas A, Yang W, Wang C, Tipparaju SL, Chen G, Sullivan B, Swiekatowski K, Tatam
M, Gerfen C, Li N (2023) Superior colliculus bidirectionally modulates choice
activity in frontal cortex. Nat Commun 14:7358.

Todd KL, Lipski J, Freestone PS (2022) Subthalamic nucleus exclusively evokes
dopamine release in the tail of the striatum. J Neurochem 162:417–429.

van der Kooy D, Carter DA (1981) The organization of the efferent projections and
striatal afferents of the entopeduncular nucleus and adjacent areas in the rat.
Brain Res 211:15–36.

Wang L, Herman JP, Krauzlis RJ (2022) Neuronal modulation in the mouse superior
colliculus during covert visual selective attention. Sci Rep 12:2482.

Wang L, Krauzlis RJ (2018) Visual Selective Attention in Mice. Curr Biol 28:676-685.e4.

Wang L, Krauzlis RJ (2020) Involvement of Striatal Direct Pathway in Visual Spatial
Attention in Mice. Curr Biol 30:4739-4744.e5.

Wang L, McAlonan K, Goldstein S, Gerfen CR, Krauzlis RJ (2020) A Causal Role for
Mouse Superior Colliculus in Visual Perceptual Decision-Making. J Neurosci
40:3768–3782.

Wang L, Rangarajan KV, Gerfen CR, Krauzlis RJ (2018) Activation of Striatal Neurons
Causes a Perceptual Decision Bias during Visual Change Detection in Mice.
Neuron 97:1369-1381.e5.

Yartsev MM, Hanks TD, Yoon AM, Brody CD (2018) Causal contribution and dynamical
encoding in the striatum during evidence accumulation. eLife 7:e34929.

Yeterian EH, Pandya DN (1995) Corticostriatal connections of extrastriate visual areas
in rhesus monkeys. J Comp Neurol 352:436–457.

Zingg B, Peng B, Huang J, Tao HW, Zhang LI (2020) Synaptic Specificity and
Application of Anterograde Transsynaptic AAV for Probing Neural Circuitry. J
Neurosci Off J Soc Neurosci 40:3250–3267.

Figure legends

Figure 1. PS and DMS D1-MSNs innervate different basal ganglia output nuclei.

A. Viral strategy to dual-label posterior (PS) and dorsomedial (DMS) D1-MSNs using cre-dependent anterograde viruses in D1-cre transgenic mice. B. Injection site of eYFP-expressing virus in DMS (blue). C. Injection site of dsRed-expressing virus in PS (yellow); axonal labeling of DMS-originating D1-MSNs in the entopeduncular nucleus (EP; blue). D-F. Axonal labeling of PS and DMS -originating D1-MSNs in the substantia nigra (SN). Anatomical boundaries approximated from Franklin & Paxinos Atlas (2008). Scale bars: 500 μm. Abbreviations: AP (anterior-posterior), PP (peripeduncular nucleus), SNl (substantia nigra pars lateralis), SNr (substantia nigra pars reticulata).

Figure 2. PS and DMS direct pathway circuits through the nigra are anatomically distinct but have shared output targets.

A. Viral strategy to dual label PS and DMS -originating SN disynaptic pathways. B. Expression of DMS-innervated eYFP-labeled cells in the SNc and SNr. C. Expression of PS-innervated tdTom-labeled cells in the SNl and PP. D-E. Fluorescence density maps for cells and associated axons postsynaptic to the PS (D) and DMS (E). Grayscale represents the % of pixels within a 50 by 50 um area that exceed one standard deviation of z-scored fluorescent intensity (black = 100%). F-L, O-P. Terminal expression of PS→SN (tdTom) and DMS→SN (eYFP) cells in thalamus and midbrain. M-N, Q-R. Density maps of PS or DMS -initiating direct pathway terminal innervation of the superior colliculus (SC). Scale bars: 500 μm. Abbreviations: FG (Fluorogold).

Figure 3. The DMS direct pathway through the EP innervates the SC.

A. Viral strategy to label the DMS -originating EP disynaptic pathway. B. Labeled EP

cells innervated by the DMS. C,E. Terminal expression of DMS→EP cells in the

midbrain. D,F. Fluorescence density maps of terminal expression in the SC. Scale bars:

500 μm.

Figure 4. Visual cortex innervates PS more broadly and inclusively than DMS.

A. Injection site of Fluorogold (FG) retrograde tracer in the PS. B-D. PS projecting FG-

labeled neurons in visual cortex. E. Injection site of FG in the DMS. F-H. DMS projecting

FG-labeled neurons in visual cortex. Cortical divisions approximated from Franklin &

Paxinos and Allen Brain Atlases. Scale bars: 500 μm. Abbreviations: Au1 (primary

auditory cortex), AuD (dorsal secondary auditory cortex), AuV (ventral secondary

auditory cortex), CTX (cortex), Ect (ectorhinal cortex), PrS (presubiculum), RSD

(retrosplenial dysgranular cortex), RSG (retrosplenial granular cortex), TeA (temporal

association cortex), V1 (primary visual cortex), V2al (anterolateral secondary visual

cortex), V2am (anteromedial secondary visual cortex), V2l (lateral secondary visual

cortex), V2pm (posteromedial secondary visual cortex).

Figure 5. PS direct pathway activation biases visual decisions.

A. Schematic of spatially cued orientation change-detection task trial structure and left

and right visual displays. B. Peri-stimulus event histograms (PSTHs) of mean lick rates

in 40-ms bins aligned to trial epoch onsets for change (top) and catch (bottom) trials

(N=10). C. Representative expression of cre-dependent Channelrhodopsin (ChR2-

mCherry) and optic fiber placement in the PS (top) and DMS (bottom) of a Drd1a-cre

mouse. D. Schematic of light delivery to PS or DMS during the early delay (left) and late

delay or change epochs (right). E. Cumulative lick probabilities from an example ChR2-

expressing Drd1a-cre mouse. The first lick following PS light delivery onset (blue) or at a

matched time on non-stimulation trials (gray) was quantified for the early delay epoch

(left column) and late delay epoch (right column), separately for contralaterally cued

trials (top row) and ipsilaterally cued trials (bottom row). F. Light delivery to the PS

during the early delay epoch produced a small but significant increase in lick probability

for ChR2-mCherry (purple and gold; N = 6) but not mCherry control (black and gray; N =

4) mice. G. PS light delivery during the late delay epoch significantly increased lick

probability for ChR2-mCherry but not mCherry control mice. H. Light delivery during the

change event reduced the median response latency for ChR2-mCherry but not mCherry

control mice. Scale bars: 500 µm. Four-way repeated measures ANOVA (F, G); three-

way repeated measures ANOVA (H). Error bars represent standard error of the mean

(s.e.m.). d = Cohen's d effect size. Abbreviations: ns (not significant).

1019

Figure 6. PS direct pathway bias of visual decisions depends on visual context

and reward reinforcement.

A. Left: Schematic of visual displays and timing of light delivery for two-patch (late

delay) and sequence-arrested one-patch and pink noise control trials. Right: Lick

probability observed following PS D1-MSN stimulation paired with the different visual

conditions on contralaterally cued trials (N = 6 mice). B. Left: Schematic of operant

extinction session structure, with change event paired or unpaired with PS D1-MSN

stimulation Right: Omitting rewards for correct detection responses resulted in similar

extinction rates for PS D1-MSN stimulation (blue) and non-stimulation (gray) trials.

Extinction rates derived from trial hits and misses (open circles; combined sessions of 4

mice). One-way repeated measures ANOVA (A). Error bars represent s.e.m. (A);

dashed lines represent 95% confidence intervals (CIs) (B).

1032

Figure 7. PS and DMS direct pathways preferentially modulate SC neurons

responsive to visual task properties.

A. Approximated regions of the SC recorded from 6 mice on a standard atlas (Franklin

and Paxinos, 2008). Transparent bar represents the anterior-posterior and medial-

lateral location of the recording bundle center, with approximate DV range of recorded

units relative to the division between superficial (sSC) and deeper (dSC) layers (solid

bar). Filled circles indicate location of electrolytic lesioned wire tips, color-coded per

mouse. B. Distribution of recorded single units across the SC relative to the estimated

division of sSC and dSC layers (n = 530 units; N = 6 mice). C. Left: PSTHs of sSC

single unit responses to the spatially cued orientation change task stimuli (n = 88).

Right: PSTHs of dSC single unit responses with time windows for quantifying task-

related activity (n = 442). D. Percentages of sSC (light gray) and dSC (dark gray) units

that exhibited task-related activity. Chi-square tests (D). Error bars represent s.e.m. (C)

or 95% CI (D). Scale bar = 500 um. Abbreviations: D (dorsal), L (lateral).

1047

Figure 8. Superior colliculus neuronal responses to PS and DMS direct pathway

activation.

A. Mean firing rates for example dSC units that were excited (left), inhibited (center), or

both excited and inhibited (right) by PS D1-MSN optogenetic stimulation (blue)

compared to activity on non-stimulation trials (gray). Optogenetic light delivery indicated

by light blue shading. B. Same as A but for dSC units modulated by DMS D1-MSN

stimulation. C. Percentages of dSC units exhibiting task properties (colored bars) or not

(white bars) that were modulated by optogenetic activation of PS (left) or DMS (right)

D1-MSNs. D. Proportions of PS (top) and DMS (bottom) D1-MSN modulated dSC units

that were modulated during early delay (white), late delay (dark gray), or both epochs

(light gray). E. Left: from recording sessions with both DMS and PS D1-MSN

stimulation, proportions of dSC single units that were modulated by PS, DMS or both

D1-MSN pathways. Center and right: the type of modulatory effect on dSC firing by PS

and DMS D1-MSN stimulation.

1062

1063 **Tables**

 Table 1. Effects of D1-MSN activation on lick probability (related to Figure 5F-G).

	Contra-cued trials		Ipsi-cued trials	
	No light	**Light**	**No Light**	**Light**
Mean (s.e.m.)	PS ChR2-mCherry (N = 6)			
Early delay	0.026 (0.004)	0.129 (0.029)	0.043 (0.014)	0.105 (0.031)
Late delay	0.086 (0.026)	0.533 (0.056)	0.083 (0.022)	0.352 (0.053)
	PS mCherry (N = 4)			
Early delay	0.037 (0.032)	0.019 (0.018)	0.061 (0.017)	0.044 (0.032)
Late delay	0.064 (0.047)	0.052 (0.033)	0.031 (0.017)	0.084 (0.081)
	DMS ChR2-mCherry (N = 5)			
Early delay	0.030 (0.011)	0.041 (0.007)	0.064 (0.013)	0.079 (0.021)
Late delay	0.089 (0.018)	0.603 (0.080)	0.100 (0.032)	0.513 (0.103)
	DMS mCherry (N = 4)			
Early delay	0.083 (0.033)	0.058 (0.013)	0.044 (0.005)	0.043 (0.008)
Late delay	0.038 (0.014)	0.075 (0.026)	0.066 (0.017)	0.084 (0.029

	df	F	p	Partial η^2
PS 4-way ANOVA				
Epoch	1,7	48.32	0.0001	0.93
Light delivery	1,7	59.93	0.0001	0.94
Group	1,7	24.59	0.0011	0.95
Cued side	1,7	0.72	0.42	*N/A*
Group x Light	1,7	58.19	0.0001	0.94
Group x Epoch	1,7	33.45	0.0004	0.91
Epoch x Light	1,7	45.13	0.0001	0.89
Group x Light x Cued side	1,7	6.04	0.0394	0.62
Group x Light x Epoch	1,7	26.10	0.0009	0.82
Group x Light x Cued side x Epoch	1,7	6.67	0.0325	0.45
DMS 4-way ANOVA				
Epoch	1,6	20.57	0.0027	0.97
Light delivery	1,6	33.23	0.0007	0.96
Group	1,6	18.32	0.0037	0.96
Cued side	1,6	0.03	0.87	*N/A*
Group x Light	1,6	29.48	0.001	0.96
Group x Epoch	1,6	18.15	0.0037	0.97
Epoch x Light	1,6	37.53	0.0005	0.96
Group x Light x Cued side	1,6	1.30	0.29	*N/A*
Group x Light x Epoch	1,6	26.38	0.0013	0.94
Group x Light x Cued side x Epoch	1,6	0.66	0.44	*N/A*

1065 Table 2. Effects of D1-MSN activation on change event reaction time (ms) (related to
1066 Figure 5H)

	Contra-cued trials		Ipsi-cued trials	
	No light	**Light**	**No Light**	**Light**
Mean (s.e.m.)				
PS ChR2-mCherry (N = 6)	333 (19)	311 (16)	342 (18)	323 (20)
PS mCherry (N = 4)	348 (14)	341 (13)	361 (8)	343 (12)
DMS ChR2-mCherry (N = 5)	334 (21)	320 (18)	352 (18)	328 (15)
DMS mCherry (N = 4)	358 (29)	358 (31)	337 (24)	333 (21)

	df	F	p	Partial η2
PS 3-way ANOVA				
Light delivery	1,8	50.27	0.0001	0.75
Group	1,8	0.78	0.40	*N/A*
Cued side	1,8	2.98	0.12	*N/A*
DMS 3-way ANOVA				
Light delivery	1,7	4.90	0.06	*N/A*
Group	1,7	0.21	0.66	*N/A*
Cued side	1,7	0.27	0.62	*N/A*

1067

1068   Table 3. Percentages (95% CIs) of SC units modulated by task properties and D1-MSN
1069   activation (related to Figure 7D and Figure 8C).

	Cue Event	Delay Activity	Spatial Cue Mod.	Change Event	Lick Activity
All sSC units	78.4 (68.4, 86.5)	46.6 (35.9, 57.5)	18.2 (10.8, 27.8)	18.2 (10.8, 27.8)	1.1 (0.03, 6.2)
All dSC units	55.7 (50.9, 60.4)	37.3 (32.8, 42.0)	20.6 (16.9, 24.7)	22.4 (18.6, 26.6)	6.6 (4.4, 9.3)
PS mod. & Task non-mod. dSC units	21.3 (12.7, 32.3)	21.9 (14.7, 30.7)	22.9 (16.2, 30.7)	19.9 (13.6, 27.4)	27.6 (21.2, 34.6)
PS mod. & Task mod. dSC units	31.5 (23.0, 41.0)	36.1 (25.1, 48.3)	41.3 (27.0, 56.8)	51.1 (35.8, 66.3)	0 (0, 97.5)
DMS mod. & Task non-mod. dSC units	36.1 (26.6, 46.5)	29.0 (21.7, 37.1)	31.4 (24.9, 38.5)	29.4 (22.9, 36.7)	38.9 (32.3, 45.9)
DMS mod. & Task mod. dSC units	36.6 (28.4, 45.5)	49.4 (38.2, 60.6)	62.2 (47.4, 76.1)	62.5 (47.4, 76.1)	10.0 (1.2, 31.7)

1070

Figure 1

Figure 2

A — AAV1-cre + FG (DMS), AAV1-flp + FG (PS), AAV1-fDIO-tdTom, AAV1-DIO-eYFP, cre, flp, SN, (Wild-type)

B — tdTom (PS), PP, SNl, AP: -3.1 mm

C — eYFP (DMS), SNc, SNr

D — PS→SN, 100%, 0%

E — DMS→SN

F — tdTom (PS), AP: -1.1 mm

G — eYFP (DMS)

H — tdTom (PS), AP: -1.6 mm

I — eYFP (DMS)

K — tdTom (PS), AP: -3.6 mm

L — eYFP (DMS)

M — PS→SN→ anterior SC

N — DMS→SN→ anterior SC

O — tdTom (PS), AP: -4.4 mm

P — eYFP (DMS)

Q — PS→SN→ posterior SC

R — DMS→SN→ posterior SC

Figure 3

A

AAV1-cre
+ FG

DMS

cre

AAV1-DIO
-eYFP

EP

(Wild-type)

B

eYFP

EP

AP: -1.4 mm

C

eYFP

AP: -3.6 mm

D

DMS→EP→anterior SC

100%

0%

E

eYFP

AP: -4.4 mm

F

DMS→EP→posterior SC

Figure 4

Figure 5

A

Pink noise
0.2 - 0.3 s

Gabor patch cue
0.4 - 1.0 s

"Early delay"
Delay epoch
0.8 - 2.2 s

50%

10-13° orientation change

"Late delay"
Catch epoch
0.78 - 1.56 m

B

Change trials:

Left side cued
Right side cued

Catch trials:

C

PS

mCherry

DMS

mCherry

D

D1-MSN stimulation
during early delay

Trial
Epochs

Pink
Noise   Cue   Delay   Change or
catch

Delay

Light
Delivery
(25% of
trials)

0.05-1.0 mW
465 nm blue light

0   200 350   800+

Time from delay epoch onset (ms)

D1-MSN stimulation during
late delay or change event

Trial
Epochs

Noise Cue Delay   Change or
catch

Change or
catch

Light
Delivery
(25% of
trials)

0 50 200

Time from change (or catch)
epoch onset (ms)

E Example ChR2-expressing mouse:

Early delay   Late delay

Contra-cued
trials

No light
Light

Ipsi-cued
trials

F Early delay

*
d=1.55   ns

Contra-cued trials   Contra-cued trials
Ipsi-cued trials   Ipsi-cued trials

G Late delay

***
d=3.01   ns

H Change event

* d=0.50   ns
* d=0.42   ns

Figure 6

Figure 7

Figure 8

A — Example units modulated by PS D1-MSN stimulation

B — Example units modulated by DMS D1-MSN stimulation

C — PS D1-MSN activation; DMS D1-MSN activation

Legend:
- Units without task property
- Units with task property:
  - Inhibited by D1-MSN activation
  - Excited and inhibited by D1-MSN activation
  - Excited by D1-MSN activation

D — PS: 39.2%, 54.9%, 5.9%; DMS: 45.8%, 48.2%, 6.0%
- Early delay
- Early & late delay
- Late delay

E — All direct pathway modulated dSC units (n = 30 units): DMS, PS, Both

Units modulated by PS only: Excit. by PS 100%

Units modulated by DMS only: Excit. by DMS 92%, Inhib. by DMS 8%

Units modulated by both PS and DMS: Inhib. by both 8%, Inhib. by DMS 8%, Excit. by PS 8%, Excit. by DMS Inhib. by PS 50%, Excit. by both 33%