Honey bulk DNA metagenomic analysis to identify honey biological composition and monitor honey bee pathogens

Priit Paluoja[1,2,*], Mihkel Vaher[1], Hindrek Teder[1], Kaarel Krjutškov[1,2], Andres Salumets[1,2,3], Kairi Raime[1,4]

[1] Celvia CC AS, Tartu, Estonia
[2] Department of Obstetrics and Gynaecology, Institute of Clinical Medicine, University of Tartu, Tartu, Estonia
[3] Division of Obstetrics and Gynecology, Department of Clinical Science, Intervention and Technology, Karolinska Institutet and Karolinska University Hospital, Stockholm, Sweden
[4] Institute of Molecular and Cell Biology, University of Tartu, Estonia

* Corresponding author

E-mail: priit.paluoja@ccht.ee

Abstract

Honey's DNA mixture originates from various organismal groups like plants, arthropods, fungi, bacteria, and viruses. Conventional methods like melissopalynological analysis and targeted honey DNA metabarcoding offer a limited view of honey's biological composition. We conducted a honey bulk DNA metagenomic analysis to characterize the honey's taxonomic composition and identify honey bee-related pathogens and parasites based on 266 Estonian and 103 foreign honey samples. 70.4% of the DNA in Estonian honey was derived from green plant families like *Brassicaceae*, *Rosaceae*, *Fabaceae*, and *Pinaceae*. Geographical distribution analysis revealed distinct botanical compositions between Estonian mainland and island samples. The bacterial family *Lactobacillaceae* was prevalent overall, reflecting the honey bee microbiota in honey. We detected 12 honey bee pathogens and parasites, including *Paenibacillus larvae* and *Nosema ceranae*. In conclusion, the study underscores the potential of bulk DNA-based and non-targeted metagenomic approaches for monitoring honey bee health, environmental quality, and honey composition, origin, and authenticity.

Introduction

Honey bees are considered effective large-scale environmental monitors due to their large-scale foraging activities. Their hive products, especially honey, provide a snapshot of the honey bee and honey production environment, containing nectar and pollen DNA from various plant species and DNA sequences from arthropods, fungi, bacteria, and viruses[1,2]. Previous studies focusing on North European honey biological composition based on DNA analysis have identified predominant floral sources such as *Brassica, Trifolium, Malus, Prunus, Fragaria, Medicago, Populus*, and *Solanum*[3,4], that are widely spread plant genera also in Estonian nature. *Apis mellifera*, as anticipated, is the most commonly detected arthropod species in honey DNA analyses [1,3]. Additionally, DNA from several other arthropods from honey bee foraging environments, like plant-sucking and honeydew-producing insects, aphids from the order *Hemiptera* have been detected not only from honeydew honey but also from blossom honey[5]. From viruses, mainly *Apis mellifera* filamentous virus (AmFV) has also been identified within honey DNA, which is known to be a ubiquitous dsDNA virus that affects many apiaries throughout Europe and can have mild pathogenetic effects on honey bees[6]. Also, fungi, mostly yeasts, that are known to tolerate high sugar concentration and recognised for their roles in food and beverage production as fermentative agents, such as species from *Zygosaccharomyces*, and fungal pathogens affecting insects or plants, such as *Metarhizium* spp., *Aspergillus* spp., *Nosema (Vairimorpha) ceranae*, *Bettsia alvei*, or *Alternaria alternata*, have also been observed in honey samples[1,3]. Honey DNA has been found to contain common microorganisms from the honey bee gut microbiota, such as *Lactobacillus kunkeei*, as well as pathogens affecting honey bees or plants, and ubiquitous species like *Escherichia coli*[1]. Honey DNA analysis has been used to detect several potential honey bee pathogens, such as *Paenibacillus larvae* – the causative agent of American Foulbrood, *Melissococcus plutonius* – the aetiological agent of European Foulbrood, and *Spiroplasma* species – the agent of the spiroplasmosis[1,3,7]. Screening for pathogens is essential for several reasons. This detection aids in identifying and managing diseases affecting honey bee colonies that are already at an early stage. Colony losses have been linked to pathogens such as *Varroa destructor* or *Nosema ceranae*[8]. Sensitive bulk DNA-based screening allows the detection of infections before visual

symptoms appear. For hive health, early detection of pathogens can facilitate timely intervention,

potentially saving colonies from devastating diseases. Additionally, understanding the prevalence and

spread of pathogens locally and on larger scales can help monitor and manage diseases and invasive

honey bee parasites.

Considering the above, the honey composition reflects the surrounding ecological landscape. It helps

detect pathogens, map hive health, describe the honey bee foraging and honey production

environment, and describe geographical peculiarities, creating a fingerprint of common regional honey

and combating food fraud. Traditional methods, such as melissopalynological analysis or DNA

metabarcoding, offer a limited view of honey composition. The melissopalynological analysis is

restricted to detecting pollen plants, ignoring nectar and honeydew plants and other organisms,

including pathogens, that leave DNA traces in honey[9]. DNA metabarcoding expands this scope by

targeting a broader range of organisms, but it remains a targeted approach, limited to detecting only

targeted taxa based on a few successfully preamplified genomic regions[10]. To use an unbiased

approach, we used shotgun metagenomics sequencing of all DNA extracted from honey sample, which

describes the complexity of samples containing thousands of distinct species belonging to different

kingdoms or phyla[10]. We conducted a thorough all-DNA-sequencing-based metagenomic analysis on

266 Estonian centrifugally-extracted honey samples to map the botanical composition of Estonian

honey with geographical distribution analysis. Additionally, we included 103 foreign honey samples

for comprehensive honey bee-related pathogens and parasites analysis, as not all honey bee-related

pathogens and parasites of interest are present in Estonia.

Methods

Honey samples

A total of 264 honey samples were collected from various regions across Estonia to describe the DNA taxonomic composition of Estonian honey (**S1 Fig**). Additionally, two positive control samples from the hives with diagnosed American Foulbrood infection caused by *Paenibacillus larvae* were included, although their specific locations were not disclosed and are therefore included in honey bee pathogen analysis but not in the analysis of Estonian honey DNA botanical composition and geographical distribution. For honey bee pathogen analysis, in addition to the Estonian honey samples, 103 foreign samples were obtained directly from beekeepers, shops, or honey markets (Error! Reference source not found.). All samples were produced during the summers of 2020 to 2022 and collected for analysis between 2020 and 2023. All honey samples were collected from centrifugal extracted honey and not directly from the honeycomb. Centrifugally extracted honey samples contain DNA traces from several honeycombs and several hives in the apiary and provide a more comprehensive DNA taxonomical composition picture of the honey that is sold on a market as well as the honey bees' foraging, hives', and honey production environment in an apiary.

DNA extraction and sequencing

Each honey sample was preheated at 40°C and homogenized by mixing with a clean spoon. 30 g honey was weighed into a 50 ml centrifuge tube and diluted in 25 ml of preheated MilliQ water. After centrifugation at 4000 rpm, the supernatant was removed, and bulk DNA from the pellet was extracted by NucleoSpin Food Mini kit (MACHEREY□NAGEL). The DNA was fragmented down to 150-200 bp fragments by Covaris M220 focused-ultrasonicator (Covaris) and concentrated by NucleoSpin Gel and PCR Clean-up kit (MACHEREY□NAGEL). The quality and quantity of the DNA fragments were assessed on Agilent 2200 TapeStation (Agilent Technologies). Illumina-compatible DNA libraries were prepared using the Celvia CC AS in-house developed FOCUS protocol. Briefly, fragmented 25 µl honey bulk DNA (1 ng/µl) was end-repaired and A-tailed by a specific enzymatic mixture. Short double-stranded and index-labelled DNA adapters were ligated to both ends of pre-treated DNA

108 fragments. The full adapter sequence and sufficient ready-made Illumina-compatible library were

109 ensured by 12-cycle PCR. 36 samples were pooled equimolarly, and the quality and quantity of the

110 pool were assessed on Agilent 2200 TapeStation (Agilent Technologies). The honey bulk DNA pooled

111 library was sequenced using the Illumina NextSeq 500 instrument (Illumina Inc.) and 85 bp single-

112 read protocol.

Metagenomic analysis

Sequencing read counts ranged from 1 to 27 million, with a median of 13.7 million reads per sample.

Sequencing data analysis was managed with Nextflow (23.09.3-edge) and conducted using the

computational resources of the High Performance Computing Center of the University of Tartu[11,12].

Sample sequencing lanes were concatenated, assessed for quality using SeqKit stats (2.4.0), and

subsequently analysed for taxonomic composition[13]. No additional pre-processing was applied to the

FASTQ files.

To classify the taxonomic composition of the Estonian honey by assigning taxonomic labels to

sequence reads, we utilized the taxonomic sequence classifier Kraken 2 (2.1.3) with a custom

reference database[14]. The minimum hit groups required for classification were set to 3, and the

confidence threshold representing a non-probabilistic scoring scheme for taxonomic assignment was

set to 0.5. Across 264 Estonian honey samples, the median percentage of classified sequencing reads

was 36.5%, ranging from 9.12% to 71.5%, with a standard deviation of 10.1%.

We constructed a comprehensive reference database by combining data from three sources: the NCBI

nucleotide (nt) collection, the One Thousand Plants Project (1KP), and the NCBI Sequence Read

Archive (SRA)[15–17]. To ensure the database is highly relevant, we focused on species commonly

associated with honey. This process involved two steps: identifying gaps in the coverage of known

honey plants and analysing unclassified DNA reads from honey samples. For the latter, we used

computational tools to match these unclassified sequences with entries in the SRA database[17]. By

selecting the most relevant sequences, we enriched the database to better capture the diversity of

organisms found in honey. Sequences from the SRA were further processed to ensure high quality

(fastp with the parameters cut_front, cut_right, and correction)[18]. Raw sequencing reads were cleaned

and assembled into longer fragments (SPAdes 3.14.1), which were then screened to remove

contaminants such as human or bacterial DNA[19]. This ensured that the final database included only

meaningful sequences for honey DNA analysis. To improve efficiency, we reduced redundancy in the

dataset by removing duplicate or overlapping sequences within each species.

To describe and estimate the abundances of honey bee pathogens and parasites in honey DNA on the

species level, we used Bracken (2.8) with the read length set to 80, taxonomic level to species, and

threshold for the abundance estimation set to 10^{20}. We analysed the presence of the following 20 honey

bee-related parasites and pathogens: *Acarapis woodi, Acarus siro, Achroia grisella, Aethina tumida,*

Ascosphaera apis, Bettsia alvei, Braula coeca, Forficula auricularia, Galleria mellonella,

Melissococcus plutonius, Nosema apis, Nosema ceranae, Oplostomus fuligineus, Paenibacillus larvae,

Senotainia tricuspis, Spiroplasma apis, Spiroplasma melliferum, Tropilaelaps clareae, Tropilaelaps

mercedesae, and *Varroa destructor.*

Finally, read counts were normalised by total read count to ensure comparability across samples.

For the geographical distribution analysis for plant genera, read counts were further normalised by the

number of samples analysed within each county. Statistical analysis of the Bracken and Kraken 2

outputs, as well as data visualisation, was conducted in R (4.4.1)[21].

Results

152 Our study presents a metagenomic analysis of honey bulk DNA to identify its taxonomic composition

153 and monitor honey bee pathogens.

Estonian honey DNA taxonomic composition

155 In our analysis of the Estonian honey DNA taxonomic composition, we characterised the proportions

156 of bacteria, fungi, animals (Animalia, Metazoa), green plants (Viridiplantae), and viruses (**Fig 1**). As

157 anticipated, most of the DNA was derived from green plants (70.4% ± 0.12), with bacteria constituting

158 a secondary component (22.7% ± 0.07).

2.9% ± 0.02

22.7% ± 0.07

1.6% ± 0.02

2.4% ± 0.02

70.4% ± 0.12

Bacteria
Fungi
Metazoa
Viridiplantae
Viruses

Fig 1. Bulk DNA taxonomic composition of Estonian honey. The symbol "±" represents the standard deviation (s.d.).

Although Viridiplantae dominated the honey composition, the dominant family identified was the bacterial *Lactobacillaceae* (average relative read abundance of 19.5%) (**Fig 2A**). Within the family *Lactobacillaceae*, the prevalent genus was *Apilactobacillus* (**S2 Fig**). Other common bacterial families were *Pseudomonadaceae* (1.7%) and *Erwiniaceae* (1.5%). The top five prevalent families of Viridiplantae in Estonian honey were *Brassicaceae* (19.1%), *Rosaceae* (13.1%), *Fabaceae* (12.0%), *Pinaceae* (9.1%), and *Salicaceae* (7.4%) (**Fig 2A**). As expected, the genera with the highest abundance were *Brassica*, *Picea*, *Trifolium*, *Rubus*, and *Salix* (**S2 Fig**).

The prominent Animalia families detected in honey DNA were *Hominidae* and *Apidae*, containing among others human (genus *Homo*), honey bee (genus *Apis*), and bumblebee (genus *Bombus*) DNA (**Fig 2A,B, S2 Fig**). Interestingly, the analysis revealed DNA traces belonging to the mammal families *Canidae* and *Bovidae*, albeit in proportions under 0.2% (**Fig 2A,B, S2 Fig**). Also, DNA from arthropod families containing honey bee or hive parasites or pests can be detected, e.g., *Varroidae*, *Pyralidae, and Vespidae* (**Fig 2B**). The prepromorant fungal families detected in honey DNA were *Saccharomycetaceae* and *Metschnikowiaceae*, mainly from yeasts' genera *Zygosaccharomyces*,

Saccharomyces, and *Metschnikowia* (**S2 Fig**). Viral DNA was predominantly from the *Apis mellifera*

*filamentous virus (***S2 Fig**).

Fig 2. Average relative read abundances of bacteria, fungi, animals (Animalia), and green plants
(Plantae) detected in Estonian honey. Panel A shows the relative read abundances for families with
an average abundance greater than 0.2% across all Estonian honey samples. Panel B shows families
with an average abundance less than or equal to 0.2%.

Estonian honey bulk DNA botanical composition and geographical distribution of plant genera

We investigated the geographical distribution of different plant genera of the family Viridiplantae in Estonian honey samples, based on their average relative sequencing read abundances (**Fig 3**). The most widely distributed genus in Estonian honey DNA was *Brassica*, as confirmed by **Fig 2**, where the family *Brassicaceae* was the most common Viridiplantae. While *Brassica* was common and contributed in most areas, there were exceptions. For example, *Brassica* was not dominant on Estonian islands like Vilsandi, Ruhnu, Muhu, Kihnu, and Vormsi (**Fig 3**). Additionally, the islands had different prevalent plant genera compared to the Estonian mainland, such as *Frangula*, *Geum, Rhamnus*, and a considerable proportion of other plant genera (categorised as "Other") (**Fig 3**). From north to south, the mainland featured common plant genera such as *Brassica, Picea, Trifolium, Salix* and *Rubus*. From east to west, there was an increase in *Rhamnus* and *Frangula* prevalence. Other plant genera, such as *Aegopodium*, *Vicia*, and *Melilotus*, were also prevalent in Estonian honey.

Fig 3. Honey bulk DNA botanical composition and geographical distribution of 16 most prevalent plant genera across Estonia. Each pie chart represents a county, showing the proportional composition of plant genera identified in honey samples. The color-coded legend indicates the corresponding plant genera.

Honey bee pathogens and parasites in honey bulk DNA

In addition to plants, our methodology also detects DNA traces from animals, fungi and bacteria, including honey bee-related pathogens and parasites. We pre-selected and monitored 20 honey bee pathogens and parasites in Estonian and foreign honey samples (see **Methods**). Specific DNA sequences from 12 pathogens or parasites (out of 20 monitored) were detected in numerous samples with either laboratory-confirmed pathogens, visually confirmed parasites, beekeeper-suspected issues, or samples without confirmation (**Fig 4, Fig 5**). For instance, DNA proof from the bacterium *Paenibacillus larvae*, which causes honey bee disease American Foulbrood, was detected in both two

laboratory-confirmed control honey samples, each with a fraction of sequencing reads exceeding 2%.

In all the samples where the microsporidian parasite *Nosema* sp. was detected, including two samples

from the hives suspected of nosematosis, only *Nosema ceranae* was detected but not *Nosema apis*. As

expected, DNA traces of *Aethina tumida* (small hive beetle) were only observed in some foreign

samples, from USA, Spain and Ghana, as this beetle is not present in Estonia or other European

countries. DNA traces from flour mite *Acarus siro* were detected in one Estonian honey sample. The

widespread parasitic honey bee mite (*Varroa destructor*) and the greater wax moth (*Galleria*

mellonella) were found in many Estonian and foreign apiaries (**Fig 5**). Also, DNA sequences from

honey bee pathogens or pests like *Ascosphaera apis* (fungus causing Chalkbrood), *Melissococcus*

plutonius (causing European Foulbrood), *Spiroplasma* species (related to spiroplasmosis, May

disease), *Bettsia alvei* (causing pollen mold), and even from *Forticula auricularia* (insect, European

earwig) were detected in numerous Estonian and foreign honey samples (**Fig 4, Fig 5**).

Fig 4. Detection of pathogens and parasites in Estonian (A) and foreign (B) honey samples. Red triangles indicate laboratory-confirmed pathogens or visually confirmed parasites, while orange triangles represent beekeeper-suspected issues. Grey points ("No information") depict samples with no information. Honey samples that did not yield any sequencing reads assigned to the pathogens listed in the Methods section are excluded from this figure. A fraction close to 0% signifies a very low proportion of sequencing reads assigned to a particular pathogen but indicates presence. Notably, certain pathogens were detected exclusively in either Estonian or foreign honey samples. For example, *Aethina tumida* presence was found only in foreign samples (panel B), whereas *Acarus siro* was detected in only one Estonian sample (panel A). Sequencing reads originating from *Acarapis woodi* were not detected in any of the samples analysed.

Fig 5. Comparison of the percentage of honey samples from Estonia and foreign origins affected by screened pathogens and parasites.

Discussion

The honey bulk DNA metagenomic analysis provides a more unbiased and less restricted overview of honey's biological composition compared to the targeted DNA-based approaches. Unlike the DNA metabarcoding method, which targets limited selected gene(s) of the specific organism(s), the honey bulk DNA approach provides a comprehensive overview of honey botanical, microbial, fungal, viral, and animal (including entomological) diversity, including honey bee pathogens and parasites[22]. We conducted thorough analyses on 266 Estonian and 103 foreign honey samples. Unlike honeycomb-scrapped samples, these samples were collected from centrifugally extracted honey, which contains honey DNA from various honeycombs and hives of the apiaries from different locations. The amount of at least one million metagenomic DNA sequencing reads per honey sample enables us to describe the biological environment of honey bee foraging and honey production. In addition to the DNA only from plant pollen, this method analyses all DNA traces in the sample, including cell-free DNA, which allows us to detect pollen, nectar and honeydew plants as well as DNA from other organisms.

We demonstrate that green plants (Viridiplantae) constitute the majority of the DNA content in honey, accounting for 70.4% of the total honey DNA composition, with *Brassicaceae*, *Rosaceae*, *Fabaceae*, *Pinaceae*, and *Salicaceae* being the most common plant families identified in Estonian honey (**Fig 1, Fig 2**). The most common plant genera were expectedly *Brassica*, *Picea*, *Trifolium*, *Rubus*, and *Salix* (**S2 Fig**). These results concord with the observations made for the composition of honey pollen plants in Estonia[23], indicating that a significant part of plant DNA in honey may originate from plant pollen.

Interestingly, the most predominant genus detected in honey DNA based on the amount of sequencing reads was not from the plants but the bacterial genus *Apilactobacillus*, aligning with its known association with honey bee microbiota (**S2 Fig**), as also shown by the past study[3]. Although in much lower proportions, also other notable bacterial families, like *Pseudomonadaceae* and *Erwiniaceae* (1.7% and 1.5%, respectively), were detected, both of which include species known for their roles in various ecological functions and interactions with plants and insects (**Fig 2**)[24]. These findings demonstrate that the taxonomic diversity of plant genera in honey DNA surpasses that of bacterial

genera. Moreover, bacterial taxa are often represented by a few dominant families, while plant DNA is

more evenly distributed across numerous genera.

As expected, the most common Animalia families detected in honey DNA were the mammal's family

Hominidae and the arthropods' family *Apidae*, containing mostly human (genus *Homo*), honey bee

(genus *Apis*), and bumblebee (genus *Bombus*) DNA from honey bee foraging and honey production

environment. Interestingly, DNA from arthropod families containing common honey bee or hive

parasites or pests from the honey bee or honey production environment can be detected in honey

DNA, e.g., *Varroidae*, *Pyralidae, and Vespidae* (**Fig 2B**). The family *Vespidae* includes species

detrimental to honey bees, such as hornets. Although hornet DNA detected in our samples was mainly

from the European hornet *Vespa crabro*, this finding could be valuable when searching methods for

monitoring and early detection of the Asian hornet (*Vespa velutina*), a species known to be devastating

for honey bee populations in warmer areas of Europe, but not yet detected in Estonia [25]. The

widespread parasitic honey bee mite (*Varroa destructor*) from the arthropod family *Varroidae* and the

greater wax moth (*Galleria mellonella*) from the family *Pyralidae* were detected both in many

Estonian and foreign honey samples (**Fig 4, Fig 5**)[26,27].

In contrast, the small hive beetle (*Aethina tumida*), known to cause colony collapses in weak colonies,

was only found in three samples, according to the label originating from the US, Spain, Ghana, and

two honey blends of undetermined geographical origins (**Fig 4**)[28]. Importantly, *Aethina tumida*, known

to be absent in Estonia, was not detected in any Estonian honey samples (**Fig 4**). This approach

demonstrates that the honey bulk DNA metagenomic analysis could be a valuable screening tool to

monitor agriculturally significant honey bee parasites' prevalence and geographical distribution.

Our analysis revealed the presence of several other honey bee-related pathogens and parasites (**Fig 4**).

Notably, the bacteria species *Paenibacillus larvae*, which is known to cause American Foulbrood

disease in honey bees, was detected in several samples, including two positive control honey samples

from the hives that were confirmed to have American Foulbrood disease[29]. In both control samples, a

substantial proportion of sequencing reads were attributed to *Paenibacillus larvae* (**Fig 4,** 8.5% and

2.4%). Also, DNA traces from honey bee pathogens or parasites like *Ascosphaera apis* (fungus

causing Chalkbrood), *Melissococcus plutonius* (causing European Foulbrood), *Nosema ceranae*

(microsporidian parasite, causing Nosematosis), *Spiroplasma* species (related to spiroplasmosis, May

disease), *Bettsia alvei* (causing pollen mold), and even from *Forticula auricularia* (insect, European

earwig) were detected in several Estonian and foreign honey samples. We did not detect DNA of the

following honey bee pathogens or parasites in any analysed Estonian or foreign honey sample:

Acarapis woodi (parasitic honey bee mite, causes acarapiosis), *Achroia grisella* (lesser wax moth),

Braula coeca (Braula fly, bee louse), *Oplostomus fuligineus* (large African hive beetle), *Senotainia*

tricuspis (fly, causes senotainiosis), *Tropilaelaps clareae* (parasitic honey bee mite, causes

tropilaelapsosis) and *Tropilaelaps mercedesae* (parasitic honey bee mite, causes tropilaelapsosis). This

might be because these important honey bee pathogens and parasites species are not widespread

worldwide, and none of these have been seen in Estonian apiaries yet. We also did not detect the

microsporidian parasite *Nosema apis* in our samples, even though it has been identified as the primary

Nosema species responsible for Nosematosis in Estonia[30]. Research has shown that *N. ceranae* has

replaced *N. apis* in many countries including Italy, Argentina or even northern countries such as

Lithuania[30–34]. Essentially, *N. ceranae* has spread rapidly worldwide[30]. Therefore, it is possible that *N.*

ceranae has also replaced *N. apis* in Estonia by now.

Interestingly, we even detected trace amounts of DNA sequences from mammals, probably originating

from domestic or pest animals, possibly due to the contamination from the honey bee foraging,

honeycombs´ storage, hove or honey production environment. For example, honey bees often collect

brackish water enriched with mineral salts, which could be contaminated by mammal excreta and

DNA (*Canidae* and *Bovidae,* **Fig 2**) [35]. This result shows the sensitivity of DNA analysis and indicates

the possible DNA transfer through honey bees' diet. This is in accordance with the study that has

demonstrated the presence of DNA from plant-sucking insects in honey DNA that produce the sticky

excretion collected by honey bees[5]. DNA contamination from pest animals, such as mice representing

<0.2% of sequencing reads, may result from their contact with the honeycombs or the hive

environment.

The fungal community was primarily represented by *Saccharomycetaceae* and *Metschnikowiaceae*,

families of yeasts, mainly genera *Zygosaccharomyces*, *Saccharomyces*, and *Metschnikowia*, commonly

involved in fermentation processes (**S2 Fig**). The presence of *Saccharomycetaceae* has also been

detected in previous honey-related studies[1,3,36]. We also detected viral DNA, predominantly from the

Apis mellifera filamentous virus (**S2 Fig**), which is known to infect honey bees but is little to no

pathogenic and has been detected in past studies[6,37]. The difference between our finding of 2.9%

sequencing reads assigned to DNA viruses and the 40.2% ($\pm$ 30.0%), as reported by Wirta *et al.* [3], can

be explained by differences in the reference database and the number of samples analysed (**Fig 1**).

However, most honey bee disease-causing, such as Deformed Wing Virus and many other, are RNA

viruses that need RNA-based screening methods[38].

We also investigated Estonian honey DNA botanical composition and geographical distribution of

most prevalent plant genera (**Fig 3**). Consistent with previous findings, we also observed frequent

occurrences of *Brassica*, *Malus,* and *Trifolium*, aligning with previous records from North European

honey (**Fig 3**)[3,23,39]. Interestingly, we observed distinct differences in the plant genera compositions

between the Estonian mainland and the islands, with the islands showing a higher proportion of

Frangula and species categorised as "Other" compared to the mainland (**Fig 3**). In the honey DNA

samples from small islands in Estonia, the proportion of *Brassica* was substantially lower compared to

the other regions. This could be explained by the lack of large agricultural fields on small islands in

Estonia. Furthermore, the diverse DNA taxonomical composition of honey creates a unique fingerprint

for every honey sample containing hundreds of different species of plants, bacteria, fungi,

animals/arthropods/ and other organisms. Therefore, we hypothesise that metagenomic analysis of all

extracted DNA could be utilised to analyse the authenticity and geographical origin of honey (**Fig 2**,

Fig 3).

Metagenomic analysis of honey DNA presents inherent challenges, primarily because the accuracy of

the results heavily relies on the public reference database used for analysis, as also pointed out by

other researchers [40]. If a genus is absent from the database, it can introduce biases and potentially

reduce the accuracy of the analysis [40]. As comprehensive databases for plants are still under

development and there is a predominance of complete genome sequences for bacteria and viruses in existing databases, we created a custom Kraken 2 reference database in our study (including partial genome sequences), with the extended numbers of honey-related plants. Our Kraken 2 reference database was sourced from three main collections: NCBI nt collection, The One Thousand Plants Project, and NCBI's Sequence Read Archive [15–17]. This approach enables the detection of an increased number of plants in honey DNA. In addition, the majority of foreign honey samples used in the pathogen analysis in this study were acquired from shops, the contents of the honey jars were not validated, and we had to rely on the label information. However, as we were using foreign honey samples only for pathogen analysis in this study, the accuracy of the label did not affect the proof-of-concept of detecting known pathogens in honey samples.

In conclusion, our metagenomic analysis of honey DNA provided a detailed and comprehensive overview of its biological composition, highlighting its significant diversity of botanical composition, and honey bee-related pathogens and parasites. This study mapped the botanical composition of Estonian honey with geographical distribution of different plant genera in Estonia, and conducted honey bee-related pathogen analysis, underscoring the potential of all DNA sequencing-based metagenomic approaches not only for describing the botanical composition of honey, monitoring honey bee health and apiary environment but also for identifying authenticity and origin of honey by using untargeted analysis of all DNA sequences extracted from honey.

Data availability

The data generated during this study is available in the Sequence Read Archive (SRA) repository under BioProject PRJNA1135913 (https://www.ncbi.nlm.nih.gov/sra/PRJNA1135913).

Funding

This work was supported by the European Agricultural Fund for Rural Development (Estonian Rural Development Plan 2014-2020, 616219790085).

References

1. Bovo S, Ribani A, Utzeri VJ, Schiavo G, Bertolini F, Fontanesi L. Shotgun metagenomics of honey DNA: Evaluation of a methodological approach to describe a multi-kingdom honey bee derived environmental DNA signature. *PLoS One*. 2018;13(10):e0205575. doi:10.1371/JOURNAL.PONE.0205575

2. Wirta H, Abrego N, Miller K, Roslin T, Vesterinen E. DNA traces the origin of honey by identifying plants, bacteria and fungi. *Sci Reports 2021 111*. 2021;11(1):1-14. doi:10.1038/s41598-021-84174-0

3. Wirta HK, Bahram M, Miller K, Roslin T, Vesterinen E. Reconstructing the ecosystem context of a species: Honey-borne DNA reveals the roles of the honeybee. *PLoS One*. 2022;17(7):e0268250. doi:10.1371/JOURNAL.PONE.0268250

4. Varis A-L, Helenius J, Koivulehto K. Pollen spectrum of Finnish honey. *Agric Food Sci*. 1982;54(5):403-420. doi:10.23986/AFSCI.72111

5. Utzeri VJ, Schiavo G, Ribani A, et al. Entomological signatures in honey: an environmental DNA metabarcoding approach can disclose information on plant-sucking insects in agricultural and forest landscapes. *Sci Rep*. 2018;8(1):9996. doi:10.1038/S41598-018-27933-W

6. Papp M, Tóth AG, Békési L, et al. Apis mellifera filamentous virus from a honey bee gut microbiome survey in Hungary. *Sci Reports 2024 141*. 2024;14(1):1-8. doi:10.1038/s41598-024-56320-x

7. Ebeling J, Knispel H, Hertlein G, Fünfhaus A, Genersch E. Biology of Paenibacillus larvae, a deadly pathogen of honey bee larvae. *Appl Microbiol Biotechnol*. 2016;100(17):7387-7395. doi:10.1007/S00253-016-7716-0/FIGURES/1

8. Ravoet J, Maharramov J, Meeus I, et al. Comprehensive Bee Pathogen Screening in Belgium Reveals Crithidia mellificae as a New Contributory Factor to Winter Mortality. *PLoS One*. 2013;8(8):e72443. doi:10.1371/JOURNAL.PONE.0072443

9.   Louveaux J, Maurizio A, Vorwohl G. Methods of Melissopalynology. *Bee World*.
1978;59(4):139-157. doi:10.1080/0005772X.1978.11097714

10.   Porter TM, Hajibabaei M. Scaling up: A guide to high-throughput genomic approaches for
biodiversity analysis. *Mol Ecol*. 2018;27(2):313-338. doi:10.1111/MEC.14478

11.   Tartu U of. UT Rocket. Published online 2018. doi:10.23673/PH6N-0144

12.   DI Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow
enables reproducible computational workflows. *Nat Biotechnol 2017 354*. 2017;35(4):316-319.
doi:10.1038/nbt.3820

13.   Shen W, Sipos B, Zhao L. SeqKit2: A Swiss army knife for sequence and alignment
processing. *iMeta*. 2024;3(3):e191. doi:10.1002/IMT2.191

14.   Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. *Genome Biol*.
2019;20(1):1-13. doi:10.1186/S13059-019-1891-0/FIGURES/2

15.   Benson DA, Karsch-Mizrachi I, Lipman DJ, Ostell J, Wheeler DL. GenBank. *Nucleic Acids
Res*. 2007;35(suppl_1):D21-D25. doi:10.1093/NAR/GKL986

16.   Leebens-Mack JH, Barker MS, Carpenter EJ, et al. One thousand plant transcriptomes and
the phylogenomics of green plants. *Nat 2019 5747780*. 2019;574(7780):679-685.
doi:10.1038/s41586-019-1693-2

17.   Kodama Y, Collaboration on behalf of the INSD, Shumway M, Collaboration on behalf of the
INSD, Leinonen R, Collaboration   on behalf of the INSD. The sequence read archive:
explosive   growth   of   sequencing   data.   *Nucleic   Acids   Res*.   2012;40(D1):D54-D56.
doi:10.1093/NAR/GKR854

18.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics. 2018;34(17):i884-i890. doi:10.1093/BIOINFORMATICS/BTY560

19.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo
Assembler. *Curr Protoc Bioinforma*. 2020;70(1):e102. doi:10.1002/CPBI.102

20. Lu J, Breitwieser FP, Thielen P, Salzberg SL. Bracken: Estimating species abundance in

metagenomics data. *PeerJ Comput Sci*. 2017;2017(1):e104. doi:10.7717/PEERJ-CS.104/SUPP-

5

21. R Core Team. R: A Language and Environment for Statistical Computing. Published online

2024. https://www.r-project.org/

22. Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E. Towards next-generation

biodiversity assessment using DNA metabarcoding. *Mol Ecol*. 2012;21(8):2045-2050.

doi:10.1111/J.1365-294X.2012.05470.X

23. Puusepp L, Koff T. Pollen analysis of honey from the Baltic region, Estonia. *Grana*.

2014;53(1):54-61. doi:10.1080/00173134.2013.850532

24. Cambronero-Heinrichs JC, Battisti A, Biedermann PHW, et al. Erwiniaceae bacteria play

defensive and nutritional roles in two widespread ambrosia beetles. *FEMS Microbiol Ecol*.

2023;99(12):1-11. doi:10.1093/FEMSEC/FIAD144

25. Requier F, Rome Q, Chiron G, et al. Predation of the invasive Asian hornet affects foraging

activity and survival probability of honey bees in Western Europe. *J Pest Sci (2004)*.

2019;92(2):567-578. doi:10.1007/S10340-018-1063-0/FIGURES/7

26. Rosenkranz P, Aumeier P, Ziegelmann B. Biology and control of Varroa destructor. *J Invertebr*

Pathol. 2010;103(SUPPL. 1):S96-S119. doi:10.1016/J.JIP.2009.07.016

27. Kwadha CA, Ong'amo GO, Ndegwa PN, et al. The Biology and Control of the Greater Wax

Moth, Galleria mellonella. *Insects 2017, Vol 8, Page 61*. 2017;8(2):61.

doi:10.3390/INSECTS8020061

28. Claing G, Dubreuil P, Bernier M, et al. Prevalence of pathogens in honey bee colonies and

association with clinical signs in southwestern Quebec, Canada. *Can J Vet Res*. 2024;88(2):45.

Accessed July 8, 2024. /pmc/articles/PMC11000428/

29. Hansen H, Brødsgaard CJ. American foulbrood: a review of its biology, diagnosis and control.

444   *Bee World*. 1999;80(1):5-23. doi:10.1080/0005772X.1999.11099415

445 30. Naudi S, Šteiselis J, Jürison M, et al. Variation in the Distribution of Nosema Species in

446   Honeybees (Apis mellifera Linnaeus) between the Neighboring Countries Estonia and Latvia.

447   *Vet Sci 2021, Vol 8, Page 58*. 2021;8(4):58. doi:10.3390/VETSCI8040058

448 31. Pacini A, Mira A, Molineri A, et al. Distribution and prevalence of Nosema apis and N.

449   ceranae in temperate and subtropical eco-regions of Argentina. *J Invertebr Pathol*.

450   2016;141:34-37. doi:10.1016/J.JIP.2016.11.002

451 32. Papini R, Mancianti F, Canovai R, et al. Prevalence of the microsporidian Nosema ceranae in

452   honeybee (Apis mellifera) apiaries in Central Italy. *Saudi J Biol Sci*. 2017;24(5):979-982.

453   doi:10.1016/J.SJBS.2017.01.010

454 33. Klee J, Besana AM, Genersch E, et al. Widespread dispersal of the microsporidian Nosema

455   ceranae, an emergent pathogen of the western honey bee, Apis mellifera. *J Invertebr Pathol*.

456   2007;96(1):1-10. doi:10.1016/J.JIP.2007.02.014

457 34. Sinpoo C, Paxton RJ, Disayathanoowat T, Krongdang S, Chantawannakul P. Impact of

458   Nosema ceranae and Nosema apis on individual worker bees of the two host species (Apis

459   cerana and Apis mellifera) and regulation of host immune response. *J Insect Physiol*.

460   2018;105:1-8. doi:10.1016/J.JINSPHYS.2017.12.010

461 35. Mahefarisoa KL, Simon Delso N, Zaninotto V, Colin ME, Bonmatin JM. The threat of

462   veterinary medicinal products and biocides on pollinators: A One Health perspective. *One*

463   *Heal*. 2021;12:100237. doi:10.1016/J.ONEHLT.2021.100237

464 36. Marvin GE. The Occurrence and Characteristics of Certain Yeasts Found in Fermented Honey.

465   *J Econ Entomol*. 1928;21(2):363-370. doi:10.1093/JEE/21.2.363

466 37. Hartmann U, Forsgren E, Charrière JD, Neumann P, Gauthier L. Dynamics of Apis mellifera

467   Filamentous Virus (AmFV) Infections in Honey Bees and Relationships with Other Parasites.

468   *Viruses 2015, Vol 7, Pages 2654-2667*. 2015;7(5):2654-2667. doi:10.3390/V7052654

38. Highfield AC, El Nagar A, Mackinder LCM, et al. Deformed Wing Virus Implicated in
Overwintering Honeybee Colony Losses. *Appl Environ Microbiol*. 2009;75(22):7212.
doi:10.1128/AEM.02227-09

39. Salonen A, Ollikka T, Grönlund E, Ruottinen L, Julkunen-Tiitto R. Pollen analyses of honey
from Finland. *Grana*. 2009;48(4):281-289. doi:10.1080/00173130903363550

40. Lu J, Rincon N, Wood DE, et al. Metagenome analysis using the Kraken software suite. *Nat
Protoc 2022 1712*. 2022;17(12):2815-2839. doi:10.1038/s41596-022-00738-y

Supporting information

S1 Fig. Geographical distribution of 264 Estonian honey samples used in the study.

481
482 **S2 Fig. Common genera of Bacteria, Fungi, Animalia, Plantae, and Viruses from the DNA of**

483 **Estonian honey.**

485   **Table S1. Origins of 103 foreign honey samples utilised in the pathogen analysis.**

Origin	N
Austria	1
Bulgaria	9
China	10
Denmark	2
England	2
Finland	3
Germany	1
Ghana	1
Greece (including Rhodes Island)	4
India	4
Italy	1
Kazakhstan	1
Latvia	4
Mix of EU and non-EU honey	18
Montenegro	1
Non-EU honey (including Ukraine, Central and South American honey)	7
Poland	3
Portugal	1
Saudi Arabia	1
Scotland	3
Spain	2
Switzerland	2
Ukraine	17
United Arab Emirates	1
Unspecified EU honey	2
USA	1
Yemen	1