Nanopore-based analysis unravels the genetic landscape and phylogenetic placement

of human-infecting *Trichuris* species in Côte d'Ivoire, Tanzania, Uganda, and Laos

Nurudeen Rahman[1,2], **Max A. Bär**[1,2], **Julian Dommann**[1,2], **Eveline Hürlimann**[1,2], **Jean T.**

Coulibaly[1,2,3,4], **Said Ali**[5], **Somphou Sayasone**[1,2,6], **Prudence Beinamaryo**[7], **Jennifer**

Keiser[*1,2], **Pierre H.H. Schneeberger**[*1,2]

*These authors contributed equally as senior authors

[1] Department of Medical Parasitology and Infection Biology, Swiss Tropical and Public

Health Institute, Allschwil, Switzerland

[2] University of Basel, Basel, Switzerland

[3] Unité de Formation et de Recherche Biosciences, Université Felix Houphouët-Boigny,

Abidjan, Côte d'Ivoire

[4] Centre Suisse de Recherches Scientifiques en Côte d'Ivoire, Abidjan, Côte d'Ivoire

[5] Public Health Laboratory Ivo de Carneri, Chake Chake, Pemba, Zanzibar, Tanzania

[6] Lao Tropical and Public Health Institute, Vientiane, Laos

[7] Vector Borne and Neglected Tropical Diseases Division, Ministry of Health, Kampala,

Uganda

Abstract

Soil-transmitted helminthiases (STH), including trichuriasis, pose a significant global burden,

affecting numerous mammalian hosts. Traditional diagnostic methods have struggled to

differentiate species within the *Trichuris* genus, prompting the use of molecular techniques to

understand the diversity and zoonotic transmission of *Trichuris* spp. among humans and

non-human primates (NHP), dogs, and pigs. In this study, we used nanopore-based full-

length ITS2 rDNA sequencing to genetically characterize and confirm the phylogenetic

placement of the newly identified human-infecting *Trichuris incognita* and the well-known

Trichuris trichiura. Using 670 samples from clinical studies in Côte d'Ivoire, Laos, Tanzania,

and Uganda, along with publicly available ITS2 sequences, we confirmed two distinct

clusters of human-infecting *Trichuris* species. We found a varying prevalence of *T. incognita*

and *T. trichiura* across the four countries and that these were also present in NHP. Our

findings demonstrated that ITS2 fragment length accurately differentiates both human-

infecting *Trichuris* species and thus makes it a good diagnostic marker for future point-of-

care applications in resource-constrained settings.

Introduction

Trichuriasis, an infection caused by the whipworm *Trichuris spp.* is of considerable public

health significance and is considered a Neglected Tropical Disease (NTD) (1). This particular

NTD has been estimated to disproportionately infect over 465 million people worldwide,

mainly prevalent among children living in impoverished communities with limited hygiene or

sanitation facilities, leading to a significant burden of 0.64 million disability-adjusted life

years (DALYs) lost annually (2). However, several other mammals such as non-human

primates (NHP), rodents, ruminants, and marsupials to name a few encompass the more

than 20 specific taxonomic groups of hosts associated specifically with the genus *Trichuris,*

meaning they are multi-host parasites (3,4).

Previous research approaches using traditional diagnostic methods like Kato-Katz have

always regarded *Trichuris sp.* found in humans and NHP as *Trichiura trichiura* except in

host-specific cases like the NHP Colobus monkeys (*Trichuris colobae*) (5) and Chacma

baboons (*Trichuris ursinus*) (6) where there is a partial understanding of the *Trichuris* multi-

host complexity. Morphologically, both eggs and adult worms of *T. trichiura* from humans

and NHP are indistinguishable making molecular methods a promising approach (7–10).

Recent efforts employed mostly sequences from the internal transcribed spacer (ITS)

regions 1 and 2 (ribosomal DNA), mitochondrial markers and β tubulin gene to delineate this

cryptic difference, hence resolving the *Trichuris* complex or multi-host specificity (5,9–15).

These sequences were used to prove the existence of two separate clusters of *Trichuris* in

humans with the consensus being that they both were *T. trichiura* (6,9–15). However, a

recent study in Côte d'Ivoire revealed a novel *Trichuris* species, called *T. incognita,* that was

observed in a monophyletic cluster with *T.suis* and *T. colobae* (16).   ITS sequences

corresponding to this species have been observed in single human patients in Uganda, and

Cameroon, as well as several NHP (14). This sequence has also been shown to be a

reliable marker to distinguish between *T. suis* isolated from pigs, *T. incognita* isolated from

humans, and *T. trichiura* isolated from humans and non-human primates (16,17).

In this study, we sought to investigate the genetic diversity and phylogenetic placement of

human-infecting *Trichuris* species in four Low- and middle-income countries (LMIC) across

the globe. To do this, we analyzed the full ITS2 region (~500-600bp) from whipworms in

stool samples collected from a total of 670 patients from clinical studies in Côte d'Ivoire,

Tanzania, Uganda, and Laos. These samples were collected as part of randomized

controlled trials (18,19) aiming at testing the efficacy of anthelmintic treatments. We next

integrated the dataset generated from these studies with all publicly available and complete

ITS2 sequences for the *Trichuris* genus, including sequences derived from humans, non-

human primates, pigs, livestock, and rodents from multiple locations. Subsequently, we

investigated the genetic makeup and phylogenetic relationships within our study populations

and in a broader context. Finally, we identified a potential PCR-based diagnostic marker

using the ITS2 fragment length that effectively differentiates between the two known human-

infecting species, *Trichuris trichiura* and *Trichuris incognita*.

Results

Description of the study population

The population studied consisted of patients from the Pujini and Dodo shehias on Pemba

Island, Tanzania; the Kisoro and Kabale districts in Uganda; the Dabou and Jacqueville

regions around the Lagunes district in Côte d'Ivoire; and the Nambak district and Luang

Prabang province in Laos (**Fig. 1,** (adapted from Keller et al. (20)). This study included

frozen faecal samples from participants aged 6 to 60 years in Côte d'Ivoire, Tanzania and

Laos, as well as ethanol-preserved faecal samples from school-aged children (6-12 years) in

Uganda.

Fig. 1: Sampling locations of the study population. The map of the world highlights the four different
study locations: the Lagunes region in Côte d'Ivoire, Pujini and Dodo Shehias on Pemba Island in
Tanzania, Nambak districts and Luang Prabang province in Laos PDR (adapted from Keller et al. (20)
), and the Kabale and Kisoro districts in Uganda each depicted in varying colours on the map.

Stool samples used in this study were collected in the framework of a double-blind, placebo-

controlled randomized trial conducted between September 2018 and June 2020 in Côte

d'Ivoire, Tanzania, and Laos (18,21,22) as well as a parallel open-label randomized

controlled superiority trial conducted between October to November 2023 in Uganda

(registered as NCT06037876; clinicaltrials.gov/). The metadata for the demographics and

helminth infection characteristics of the 670 study participants are shown in **Table 1**. The

infection intensities as diagnosed with Kato-Katz (KK) and sex distribution of the study

participants depict a good balance between treatment groups. Of the 670 participants,

56.0% (375/670) were female and 44.0% (295/670) were male across the four countries.

Overall, the Lao participants were older (mean age 23·1 years [SD 15·2]) than the other

three cohorts (mean age 16·5 years [13·0] in Côte d'Ivoire, 14·1 years [9.8] in Tanzania, and

15.8 years [12.6] in Uganda). Samples of all 670 participants were included in PCR-based

investigations of the internal transcribed spacer 2 (ITS2) region.

Table 1 Description of the study population

	Côte d'Ivoire (n=189)	Laos (n=156)	Tanzania (n=190)	Uganda (n=135)
Age in Years				
6 -12	116 (61.4%)	71 (45.5%)	119 (62.6%)	135 (100%)
13 – 24	32 (16.9%)	14 (9.0%)	49 (25.8%)	0
≥25	41 (21.7%)	71 (45.5%)	22 (11.6%)	0
Sex				
Female	96 (50.8%)	84 (53.8%)	107 (56.4%)	88 (65.2%)
Male	93 (49.2%)	72 (46.2%)	83 (43.6%)	47 (34.8%)
Infection intensity[+]				
Light	144 (76.2%)	136 (87.2%)	144 (75.8%)	118 (87.4%)
Moderate	42 (22.2%)	20 (12.8)	43 (22.6%)	17 (12.6%)
Heavy	3 (1.6%)	0	3 (1.6%)	0

[+] Infection intensity was classified according to mean egg per gram EPG into light (1–999 EPG),

moderate (1000–9999 EPG), and heavy (≥10 000 EPG).

Molecular Analysis of ITS2 Nuclear Marker

Following DNA extraction of the faecal samples from the patients partaking in the clinical

trials across the four countries, we successfully produced amplicons from 188, 154, 174 and

113 participants in Tanzania, Laos, Côte d'Ivoire, and Uganda, respectively. The resulting

ITS2 amplicons were sequenced using a MinION platform to achieve 5 to 9896 unprocessed

reads per sample. Using a stringent Divisive Amplicon Denoising Algorithm Version 2,

DADA2-based pipeline, we retained a total of 198,792 reads (ranging between 5 to 9420

high-quality reads per sample) that were processed into 474 unique amplicon sequence

variants (ASVs) (**Additional file 2: Fig. S1**).

Population Differentiation and Network Analysis

Analyses of haplotypes showed the presence of 233 haplotypes of the *Trichuris* population

across the four different countries (**Additional file 2: Table S1**). Ten haplotypes were

shared among the ASVs from the different countries, with four haplotypes shared within the

four countries, while 223 haplotypes were revealed across all settings (51, 63, 74 and 35

from Côte d'Ivoire, Laos, Tanzania and Uganda, respectively). A few of the haplotypes

analyzed showed high frequencies, such as Hap 1 in Laos (19%), Hap 2 in Côte d'Ivoire

(31%), Hap 3 in Tanzania (14%) and Hap4 in Laos and Tanzania (18% and 24%

respectively), although these haplotypes were also present in varying proportions across the

other countries (**Additional file 2: Table S1**). Comparison of these observed haplotypes

across the four countries through NCBI BLAST showed that five haplotypes showing 100%

identity with already described haplotypes: Hap1, Hap3 and Hap 4 were identical to *T.*

trichiura isolated from samples analyzed from Ugandan patients (Accession number

KJ588133 and KJ588134 respectively, (9)); Hap2 and Hap9 identified as *T. trichiura* isolated

in worms expelled from patients in Cameroon (Accession number GQ301555, (14)). Also,

the samples analyzed in Hap2 and Hap9 had a 96% to 100% identity match to ITS2

sequences from the newly described *T. incognita* species (17) following NCBI BLAST to its

sequence. The comprehensive list encompassing the BLAST results is available as

supplementary material (**Additional file 2: Table S2**).

Table 2: Genetic diversity estimations of *Trichuris* population per region

Country	N	N_H	Gene diversity	π	H_e
Côte d'Ivoire	189	58	0.9889±0.005	0.236±0.113	0.20
Laos	156	70	0.9951±0.003	0.153±0.073	0.13

Tanzania	190	83	0.9951±0.016	0.143±0.069	0.13
Uganda	135	42	0.9929±0.050	0.173±0.083	0.16

N = Number of samples sequenced; N_H = Number of haplotypes; π = Nucleotide diversity ; H_e = Expected heterozygosity

The analysis of genetic diversity indexes revealed that gene diversity within the populations

ranged from 0.9889 to 0.9951, with the highest levels in Laos and Tanzania (both 0.9951),

and the lowest in Côte d'Ivoire (0.9889). Conversely, nucleotide diversity (π) and expected

heterozygosity (H_e) showed a different pattern, with the sample population from Côte d'Ivoire

and Uganda exhibiting higher diversity. 58 haplotypes from Côte d'Ivoire had the highest

nucleotide diversity (π = 0.236±0.113) and expected heterozygosity (H_e = 0.20), followed by

42 haplotypes from Uganda (π = 0.173±0.083, H_e = 0.16). In contrast, 70 haplotypes from

Laos and 83 from Tanzania had lower nucleotide diversity (π = 0.153±0.0730 and

0.143±0.069 respectively) and expected heterozygosity (H_e = 0.13 for both), indicating fewer

Fig. 2: Underlying population genetic structure and network analysis among the four countries A. Haplotype network inferred from ASVs of *Trichuris* species haplotype sequences

from Côte d'Ivoire, Laos, Tanzania and Uganda. Each circle represents a unique haplotype and the size of the circle is proportional to the corresponding haplotype frequency. The colours of the

pie charts represent the country of origin (red, Côte d'Ivoire; yellow, Laos; purple, Tanzania; blue, Uganda). Hatch marks along the network branches indicate the number of nucleotide differences

differences between sequences in these populations. This suggests that while 153

haplotypes from Laos and Tanzania have extensive allelic variation, the alleles are more

similar, whereas 100 haplotypes from Côte d'Ivoire and Uganda have more distinct

variations at the nucleotide level (**Table 2**).

Analysis of the haplotype network analyses revealed a reticulate relationship among the

ASVs in the four countries scattered in separate clusters (**Fig. 2A**). Specifically, the ASVs

present in *Trichuris* populations of Côte d'Ivoire, Laos, Tanzania and Uganda were observed

in the two clusters separated by 40 point mutations or nucleotide differences. By

comparison, a further internal subdivision among haplotypes with a higher proportion was

recorded in Tanzania than in the other three countries while the other sub-cluster revealed a

higher proportion of haplotypes from Laos study participants than in the other three

countries. The remaining haplotypes were observed in a different cluster, which represented

a greater proportion of the *Trichuris* population in Côte d'Ivoire than those found in the

Tanzania, Laos and Uganda populations.

Comparison of the pairwise fixation index (FST) values between the sampled populations

revealed a range from -0.0007 to 0.1774 (**Fig. 2B**). ASVs of the sampled population from

Côte d'Ivoire were highly differentiated from those in Tanzania, Laos and Uganda with FSTs

of 0.1774, 0.1692 and 0.14112 respectively (P < 0.001 for all FST values). Consequently,

ASVs between Laos and Uganda, Laos and Tanzania, as well as Uganda and Tanzania

represented an undifferentiated population (FST < 0.010, P > 0.001).

Phylogenetic and comparative sequence analysis

Phylogenetic analysis of the ITS2 ASVs from the four countries together with publicly

available ITS2 sequences of other related species was inferred using neighbour-joining (NJ),

and Maximum Likelihood (ML) algorithms and their corresponding tree topologies provided a

consistent, robust phylogenetic discrimination (**Fig. 3A**). We identified five clades, each

corresponding to different but taxonomically related host species and broadly matching

published whole genome sequences (WGS) of *Trichuris* species. All the pairwise nucleotide

identities between sequences generated in this study and publicly available sequences are

summarized in **Fig. 3B**. Clade A consisted of published ITS2 sequences of *Trichuris* species

of Indian bison, camel, deer and sheep but did not contain any well-characterized WGS of a

specific *Trichuris* species.   Clade B consisted of published sequences of *Trichuris* species

of arvicolids and murid rodents – and included the sequence of an ITS2 region from a

previously published WGS of *T. muris* (23,24). Clade C consisted of published sequences of

Trichuris species of pigs which correctly matched the WGS of *T. suis* (25). Clade D was

distinctly differentiated from *T. suis* sequences with nucleotide similarities ranging from 82 to

88% and contains ITS2 sequences from 20.5% (N = 97 samples) included in this study, as

well as 14 publicly available sequences of human and NHP species. Sequences from clade

D shared approximately 98% - 100% nucleotide sequence identity with the WGS sequence

of novel species of human-infecting *Trichuris*, named *T. incognita*, observed in Côte d'Ivoire

(17), as well as other published partial sequences from humans in Cameroon (accession

number GQ301555) (14), non-human primates (NHP) from Italy and Uganda, and more

distantly with Chacma baboons in South Africa (accession number GQ301554) (14) with

90% to 94% nucleotide sequence identity. It is noteworthy that over 52.6% of samples

contained in this clade were from Côte d'Ivoire (N = 51) but it also contained samples from

other countries, Laos (N = 15 samples), Tanzania (N = 21 samples), and Uganda (N = 10

samples) (**Fig. 3C**). Clade E contained 377 *Trichuris* sequences derived from the human

samples in our study as well as 36 ITS2 reference sequences of human and NHP species

and 1 WGS sequence corresponding to the most common human-infecting species, *T.*

trichiura. Clade E shared 96% - 100% nucleotide sequence identity to published sequences

from humans in Ecuador and Uganda (9,23), as well as sequences from captive and non-

captive NHP from Uganda, China, South Africa, Italy, Spain, and St. Kitts & Nevis

(10,11,14,26,27). It showed a more distant nucleotide sequence identity (79% to 86%) with

sequences obtained from Francois Leaf monkeys in China (26). The majority of sequences

clustering in this clade were from Tanzania (N = 148), Laos (N =108), and Uganda (N = 59),

but also contained approximately 16.4% (N

Fig. 3: Phylogenetic and sequence characterization of *Trichuris* population A. Maximum likelihood tree based on the ITS2 rDNA using Tamura-Nei with gamma distribution as the substitution model and *Trichinella spiralis* (GenBank accession: KC006432) as an outgroup. Coloured branches correspond to major clades A to E. External rings (from outer to inner one) represent metadata from the samples (as well as other *Trichuris* reference sequences available from rodents, livestock, pigs, humans, and non-human primates), namely continent, country of origin, host species and host-type respectively. **B.** Pairwise comparison of sequence similarity between human samples (x-axis) from the four countries and reference sequences (y-axis) incorporated in the phylogenetic tree using ward clustering method and Euclidean correlation on OriginPro 2021 graphing software. **C.** Proportion of human-infecting *Trichuris* species observed within Côte d'Ivoire, Uganda Laos, and Tanzania.

204   = 62) of sequences generated from samples collected in Côte d'Ivoire. Furthermore, clade

205   E was also more distant from clades C and D with nucleotide identity values ranging from

206   64% to 69%.

Fragment length as a robust diagnostic marker to differentiate human-infecting
Trichuris species

209   The fragment lengths - in basepair (bp) - of the ITS2 regions corresponding to the clade

210   structure (clades A to E) are summarized in **Fig. 4**. The length of the ITS2 regions ranged

211   from 424.9±47.6bp for _T. muris_-related sequences, 443.7±7.9bp for _Trichuris_ species found

212   in livestock, 579.9±2.4bp for _T. suis_-related sequences, 603.7±13.5bp for _T. incognita_-

213   related sequences, and 539.1±30.4bp for _T. trichiura_ related sequences. We observed

**Figure 4: Fragment length could distinguish between different human-infecting _Trichuris_
species**. **A** Comparison of the fragment length between the five clades. Box plots include the
median line, p-value estimated by the one-tailed Mann–Whitney U test. **B** ROC curve showing the
performance of random-forest models built by combining the 2 signatures features of fragment
length using the lose-1-keep1- m. bp: base pair.

significant differences in fragment length between all clades, except between _T. muris_ and

the livestock-related clade A.

We then evaluated the predictive power of the ITS2 region length to discriminate between

the two human-infecting species, _T. incognita_ and _T. trichiura_. For this purpose, we used a

random forest model on our dataset as shown in **Fig. 4B**. The model exhibited comparable

classification performance with an area under the receiver operating curve of this model of

99.7% (95% CI: 99.4-100). The fragment length-based random forest model accurately

classified 95.4% (Kappa = 85.9%) of samples into their respective *Trichuris* species. Taken

together, our results showed that the fragment length is an accurate diagnostic marker to

differentiate between the two human-infecting *Trichuris* species.

Human-infecting *Trichuris* species and metadata associations

By focusing on a sub-group of study participants between the ages of 6-18 years, we sought

to understand whether there were differences in the metadata associated with both, *T.*

incognita and *T. trichiura*. We did not find any association between age and *Trichuris*

species in any of the settings, revealing a homogeneous distribution of both species in the

population traditionally most at risk, school-aged children (**Fig. 5A**). Similarly, we did not find

any differences in infection intensity (as a categorical variable defined by WHO) between

both species (**Fig. 5B/C**). Finally, we found that in 3 out of 4 settings (Laos, Tanzania, and

Uganda), females were more likely to be infected with *T. incognita* than males (**Fig. 5D**).

Discussion

The etiological agent of human trichuriasis has traditionally been identified as *T. trichiura*

based on morphological characteristics. However, morphological traits alone are often

insufficient to distinguish closely related species, especially when adaptive changes or when

species divergence has occurred relatively recently. Molecular epidemiological investigations

using coding and non-coding regions of nuclear ribosomal DNA have revealed a second

Trichuris species infecting humans, closely resembling those found in NHP and pigs

(14,16,17). Here, we utilized the nuclear ITS2 region to molecularly identify *Trichuris* species

infecting humans from samples from randomized controlled trials across diverse

geographical regions.

Our phylogenetic analysis of ASVs revealed that *Trichuris* populations from humans were

grouped into two distinct *Trichuris* genotypes across these study sites, which we distinguish

as Clade D and Clade E (**Fig. 3A**). Haplotype networks, pairwise FST, and pairwise distance

analysis (**Fig. 2A-B**; **Fig. 3B**) corroborated these results, indicating well-differentiated and

genetically divergent clades across the study sites. These findings align with previous

studies using ITS2 rDNA, demonstrating the multi-host affiliation of *Trichuris* species among

humans and NHP (9–11,14–16,28,29).

Clade D contained majorly of the ITS2 ASVs from Côte d'Ivoire indicating the establishment

of this clade in this setting, which is also where the first human samples came from that led

to the specification of this new species (16,17). However, our study not only incorporates

the enhanced resolution of full-length ITS2 sequencing but also significantly broadens the

geographical scope including samples from another district Côte d'Ivoire and an entirely new

country, Uganda, thereby enriching the diversity and depth of our data. To the best of our

knowledge, this is the first time that *T. incognita* has been described in Tanzania, Laos, and

Uganda albeit in a smaller proportion suggesting that this species has been infecting

humans and bypassing standard microscopy-based diagnostic techniques across at least

two continents with endemic trichuriasis for a significant amount of time.

Furthermore, Clade D shares a monophyletic lineage with the *T. suis* clade, with an

estimated 82 to 88% nucleotide sequence similarity. Although both share similar

physiological features, including overall morphology, egg production, and range of hosts, the

evidence derived from the nucleotide sequence similarity between the clades suggests that

T. incognita and *T. trichiura* evolved in parallel from a distant common ancestor.

In addition, we showed that Clade D contains sequences from samples collected previously

from 13 primate species spanning multiple studies and countries, as well as one human

patient in Cameroon. With approximately 98-100% nucleotide identity of the ITS2 region

among these sequences, it shows that *T. incognita* consists of a multi-host lineage

suggesting that it can infect multiple hosts just like *T. trichiura* thus justifying its potential

zoonotic origin.

In contrast to Clade D, Clade E contained over 70% of the ASVs generated from the human

samples in our study. These sequences clustered closely with those derived from 36 human

and NHP species, as well as one whole-genome sequence (WGS). Clade E sequences

conformed to the most common human-infecting species, sharing 96% - 100% nucleotide

similarity with sequences from humans in Ecuador and Uganda (9,23). Additionally, they

matched sequences from both captive and non-captive NHP in Uganda, China, South Africa,

Italy, Spain, and St. Kitts & Nevis (10,11,14,26,27). This result is in agreement with a

recently published study using ITS2 rDNA (16) and further supports that whipworms in

primates comprise a *Trichuris* species complex. Additionally, within this clade, results based

on ITS2 revealed an interpopulation similarity of approximately 79% to 86% between *T.*

trichiura populations and *Trichuris sp.* from Francois Leaf monkeys, hence, agreeing with Liu

et al. (26) that *Trichuris* populations from Francois Leaf monkeys potentially represent a new

Trichuris species (**Fig. 3A-B**).

Despite earlier theories suggesting zoonotic transmission between pigs, humans, and NHP

from an African origin to Asia and subsequently to South America (30,31), recent evidence

suggests limited transmission influenced by natural host dispersion and anthropogenic

factors in recent evolutionary history (30,32). Furthermore, comparative analyses of gene

duplication, gain, and loss patterns in *T. incognita* compared to *T. suis* and *T. trichiura*

indicate a more ancient and divergent lineage (17). All these observations collectively

indicate a complex evolutionary history, resulting in a rich diversity of co-evolving species

within the *Trichuris* genus. With reports of drug resistance emerging in the veterinary world

(33–37), coupled with the recent discovery of *T. incognita* as a result of ivermectin-

albendazole unsatisfactory drug efficacy in Côte d'Ivoire compared to other settings (16) as

well as expected resistance in human trichuriasis within the next decade (38), it is critical to

understand the distribution, transmission of *Trichuris* etiologies and potential factors for

reduced efficacy or drug resistance including parasite-related factors (strain analysis). This

understanding is essential to sustain the development of effective control strategies and

achieve the WHO's 2030 goal of disease elimination as a public health problem (39).

To that end, while next-generation sequencing poses challenges for adopting species-level

diagnostic tools in resource-constrained environments, we have identified a potential PCR-

based diagnostic marker that can accurately differentiate between the two human-infecting

Trichuris species, *T. incognita* and *T. trichiura*. This marker simplifies the detection process,

eliminating the need for expensive sequencing platforms, and establishes a cost-effective,

robust method for distinguishing these morphologically similar species. By using the length

of ITS2 fragments, a random forest model validated its diagnostic utility, achieving a 95.4%

classification accuracy (Kappa = 85.9%). This demonstrates the reliability of fragment length

as a marker in epidemiological studies and control programs, complementing previous

findings on ITS2 rDNA's effectiveness in distinguishing closely related nematode species

(11,13,16,28,40–42). The potential use of ITS2 length as a diagnostic marker is particularly

suitable for resource-constrained environments and could be combined with PCR or further

developed into LAMP assays. While MinION sequencing is relatively affordable and allows

high multiplexing with our approach, a length-based diagnostic tool is simpler and more

accessible for widespread application (43–46).

Our study has a few drawbacks. First, our observations are based on samples collected from

clinical studies in Côte d'Ivoire, Laos, Tanzania, and Uganda, which may not fully capture

the global diversity of the newly identified human-infecting *Trichuris* species due to the

limited geographical scope. Second, our dependence on nuclear ribosomal DNA for species

identification, without comprehensive morphological analysis or whole-genome sequencing,

might miss other genetic variations and phenotypic traits. Furthermore, we did not evaluate

temporal aspects of infections, such as seasonal variations or long-term changes, and our

study did not include an extensive analysis of environmental factors that could influence the

distribution and genetic diversity of *Trichuris* species. Importantly, the samples were

collected within the context of clinical trials but these represented the status before drug use

hence we did not explore how genetic diversity might relate to decreased drug efficacy.

While we believe our findings are robust, further studies with broader geographical

coverage, and integrated genomic, morphological, and environmental approaches are

needed to expand upon these findings.

In conclusion, our study utilized an enhanced resolution of full-length ITS2 rDNA sequencing

to identify two distinct human-infecting *Trichuris* species, *T. incognita* and *T. trichiura*, across

diverse regions. Our research expands the geographical scope of previous studies, including

new regions and countries such as Uganda. We also identified *T. incognita* in Tanzania, and

Laos, for the first time, reinvigorating the delimitations of traditional diagnostics in

distinguishing between different *Trichuris* species. Additionally, we identified a PCR-based

marker using ITS2 fragment length as a powerful diagnostic marker for distinguishing

human-infecting *Trichuris* species. This method demonstrates a simplified, cost-effective,

and robust alternative to expensive sequencing approaches and is especially suitable for

resource-limited settings. Furthermore, it promises improved epidemiological studies and

control programs, hence, aiding the WHO 2030 trichuriasis elimination goal. Future research

should refine this diagnostic method and explore other genetic markers in conjunction with

ITS2 fragment length to enhance diagnostic precision and provide deeper insights into the

evolutionary relationships within the *Trichuris* genus.

Methods

Study design and sample collection

The data and samples used in this study were derived from two studies. The first study was

a double-blind, placebo-controlled randomized trial conducted between September 2018 and

June 2020 in Côte d'Ivoire, Laos, and Tanzania. The study examined the efficacy and safety

of co-administered albendazole and ivermectin versus albendazole monotherapy against

whipworm infections in children and adults with the rationale, procedures, and results of the

study published previously (18,21,22).

Briefly, a fresh faecal sample was collected from community members aged between 6 and

60 years identified as infected with *T. trichiura* based on a prior stool examination and

considered eligible for the trial with an infection intensity of at least 100 eggs per gram of

stool. From every sample, about 1g of faeces was transferred into a 2ml screw cap cryotube

using a UV sterilized plastic spatula and immediately frozen at -20°C, which were then

shipped to Swiss TPH on dry ice at the end of the trial. Samples considered for this analysis

represent baseline stool samples provided before any drug administration.

Additionally, data was also derived from a parallel open-label randomized controlled

superiority trial conducted between October to November 2023 in two primary schools in

Kabale and Kisoro districts, southwestern Uganda among individuals aged 6-12 years to

investigate the superiority of co-administered ivermectin plus albendazole to albendazole

monotherapy in terms of cure rates against *T. trichiura* infections (registered as

NCT06037876; clinicaltrials.gov/). At baseline screening of the trial, two stool samples were

collected at baseline and children found to be infected with *T. trichiura* based on

quadruplicate Kato-Katz thick smear readings were enrolled into the study. A portion of the

second baseline stool sample (1.5-2 g) from 135 eligible participants was preserved in 95%

ethanol, shipped to Swiss TPH in Basel and subjected to amplicon deep sequencing for

characterization of *Trichuris* species. Faecal egg count and epidemiological data are

available in **(Additional file 2: Table S3)**.

DNA isolation and sample processing

Two separate extraction approaches were used in the isolation of DNA from stool received

from the multi-country study and the study from Uganda. For the multi-country study, a total

of 658 (226, 191, and 221 respectively from Côte d'Ivoire, Laos, and Tanzania) frozen 100

mg of stool samples from the baseline were extracted using a QiAmp PowerFecal DNA

extraction kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol with slight

modification. For the faecal samples collected in Southwestern Uganda, 150 - 250mg of

samples preserved in Ethanol was used in the extraction of the DNA. Before extraction with

the QiAmp PowerFecal DNA extraction kit, a few modification steps were taken to

mechanically rupture the *Trichuris* egg shells which include the removal of ethanol from the

stool, washing with phosphate buffer saline (PBS), as well as a freeze-thaw-boiling step to

enhance egg shell rupture and minimize inhibition (16,47) (see **Additional file 3: Appendix**

S1 for more details of the modified ethanol-preserved sample extraction protocol).

PCR Amplification, Quantification and Purification

Primers   used   in   the   ITS2   rDNA   reactions   were   Trichuris_ITS2_F   5'-

ATGTCGACGCTACGCCTGTC-3' and Trichuris_ITS2_R 5'-TAGCCTCGTCTGATCTGAGG-

3' (16) with a reverse primer containing an added custom barcodes that were adapted from

the Illumina 12bp barcode sequences (48,49) (See **Additional file 2: Table S4** for full

primer sequences). For each run, 96-well plates were multiplexed with the Trichuris_ITS2_F

primers and eight custom barcoded reverse primers to amplify the ITS2- regions yielding

unique barcodes of 592 bp ITS2 regions for each sample per plate column. LongAmp® Hot

Start Taq 2X Master Mix (cat no. M0533S, New England BioLabs, USA) and 2µL of source

DNA were used in each amplification. Thermocycling conditions included the following: 94°C

for 30 sec initially, 40 cycles of 94°C for 30 sec, 65°C for 15 sec, 65°C for 50 sec, followed

by a final extension at 65°C for 10 min carried out on an Eppendorf Mastercycler Nexus

Gradient Thermal Cycler (Eppendorf AG, Hamburg, Germany). Barcoded amplicons were

quantified directly using a dsDNA HS Assay kit on a Qubit 4 Fluorometer (Invitrogen, USA)

according to the manufacturer's protocol to ensure even throughput in the DNA pool using 2

µL of PCR reaction as input. 1% agarose and capillary gel electrophoresis (100V, 20min)

were used to verify the amplification of the fragment at the expected band size. Afterwards,

sample plates were stored overnight for a short period at 4°C overnight or at -20°C for long-

term storage.

Library preparation and sequencing

Nanopore sequencing libraries were prepared using the Native Barcoding Kit 24 v14 (SQK-

NBD114.24, Oxford Nanopore Technologies, UK) according to the Native Barcoding

Genomic DNA protocol with slight modifications aiming to normalize sample input, allowing

for multiplexing as described in more details elsewhere (49). Briefly, samples were pooled

column-wise from a 96-well plate before the end-repair step, requiring 12 different outer

barcodes for every 8 samples. Each column, representing 8 samples, was processed

together with approximately 80 ng input corresponding to 200 fmol of the expected 0.59 kb

fragments for the end-repair steps. Following end repair, samples were assessed using the

1X dsDNA HS Assay kit on a Qubit 4 fluorometer. Equimolar masses of the pooled reactions

were used for native barcode ligation to ensure equal read numbers per barcode. Adapter

ligation and clean-up followed the protocol, utilizing a short-fragment buffer to preserve the

ITS2 rDNA fragments. Afterwards, 1 µl of the prepared library was quantified on the Qubit

4.0 (Invitrogen, USA) using the Qubit dsDNA HS Assay Kit (Invitrogen, USA). The protocol

from ONT recommends loading 10-20 fmol of amplicon product onto the Flongle Flow Cell

(Flongle Flow Cell, R10 Version, FLO-FLG114, Oxford Nanopore Technologies, UK), so we

diluted to a concentration of 2-3 ng/µL in Elution Buffer to load approximately 15 fmol of the

library onto the Flongle Flow Cell. The Flongle Flow Cell was primed for loading by flushing

the flow cell with 120 µL priming mix (3 µl of Flush Tether in 117 µL of Flow Cell Flush),

taking care to avoid the introduction of air bubbles. The library was prepared for loading by

mixing 15 µl Sequencing Buffer, 10 µl Loading Beads and 5 µl diluted DNA library, after

which the sample was added to a Flongle Flow Cell on a MinION Mk1C (Oxford Nanopore

Technologies, UK) and sequenced at depth for 24 h

Read processing and sequencing analysis

All POD5 data resulting from the sequencing of the amplicons were already basecalled and

demultiplexed based on the native ONT barcode in real-time on the MinKNow v3.6.5 and the

built-in Guppy (v6.5.7) using the high accuracy basecalling model. The base-called POD5

files   were   then   re-run   through   Dorado   (v. 0.5.1;   base-caller   model

dna_r10.4.1_e8.2_400bps_sup@v4.3.0) generating high-quality simplex reads) which was

fed through our custom demultiplexing scripts that combine seqkit (v.2.6.1) (50) and custom

Python   scripts   available   at

https://github.com/STPHxBioinformatics/ITS2_Trichuriasis_nanopore_processing. The script

demultiplexes, quality filters, and length filters, as well as utilizes a minimum and maximum,

trimmed-read cutoff of 450 bp and 750 bp, respectively with zero mismatches.

Consequently, it trims the reads based on the inner PCR barcode (the reverse barcode in

this case) ensuring the barcodes are found in the correct orientation. NanoPlot v1.30.1 (51)

was then used for descriptive statistics on the runs. Demultiplexed ITS2 reads were then

subjected to the R package 'DADA2' version 1.3.2 (52) pipeline to include further filtering to

remove reads containing unresolved nucleotides (maxN = 0) as well as reads exceeding

the expected error number (maxEE = 15) and size range (425–700 bp). The dataset

generated was then used as input to define the error rates and perform the removal of

identical reads (derepFastq) while inferring the composition of the sequenced pool using the

dereplicated sequence dataset as input (dadaFs), and removal of chimeric sequences.

Following this, Amplicon sequence Variants (ASVs) generated through the DADA2 pipeline

with ASVs below 5 reads across all samples were removed.

Genetic Variation and Phylogenetic Analysis

A curated database of reference ITS2 sequences was created by downloading all publicly

available full ITS1-5.8S-ITS2 reference sequences of closely related species on NCBI using

the search term "Trichuris"[porgn:__txid36086] as well as from the WGS data of *Trichuris*

muris (23,24), *Trichuris trichiura* (23), *Trichuris suis* (25) and an adult worm from Côte

d'Ivoire (17). Afterwards, seqkit (50) was used to extract only ITS2 sequences within the

forward and reverse primers used in this study with the command seqkit -j 4 amplicon –m 2 -

p   primers.tab   -r   1:-1   20240201_Trichuris.txid_36086.fasta   --bed   >

20240201_Trichuris_txid_36086.bed. The ASVs were then subjected to multiple sequence

alignments with the newly curated reference sequences **(Additional file 2: Table S5)**, as

well as an outgroup species *Trichinella spiralis* (Accession No: KC006432), using the

MAFFT tool (Katoh et al., 2019). Phylogenetic relatedness was inferred using NJ in MEGA

v11.0 (53) and the Maximum Likelihood method using RaxML v8.0 (54) employing the

jModelTest v0.1.1 (55) to choose the best-to-fit model.

Population Genetic Structure

Alignment from good quality ITS2 ASVs was used for population genetic data and haplotype

polymorphic analysis such as numbers of variable sites, number of haplotypes, nucleotide

diversity (π), expected heterozygosity, and pairwise FST calculations between the *Trichuris*

populations in the four countries were calculated using DnaSP v6.0 (56) and Arlequin v3.5.2

(57) using 1000 permutations. Statistical Parsimonious networks were inferred and

visualized using Population Analysis with Reticulate Trees, POPART v1.7 (58) with the

connection limit set to 95% and gaps being treated as missing. Furthermore, the number of

nucleotide differences per sequence in each country and other *Trichuris* species

incorporated in the phylogenetic analysis was analyzed using the Compute Pairwise

Distance prompt based on the number of differences method of MEGA v11 (53) to assess

the sequence similarity.

Availability of data and materials

The custom demultiplexing script and DADA2 analysis code used for this study are available

at: https://github.com/STPHxBioinformatics/HITS2. The raw ONT sequence data (ITS2 rDNA

amplicon sequencing) generated in this study is available in the NCBI Short Read Archive

under accession (PRJNA1141773: https://www.ncbi.nlm.nih.gov/bioproject/PRJNA547591).

Acknowledgements

We want to acknowledge the participants and investigators of the IVM-ALB multi-country and

FACE-IT efficacy trial study. The authors also thank all the Keiser's lab members for their

valuable discussions. Computational analyses were performed using the sciCORE

(https://scicore.unibas.ch) scientific computing center at the University of Basel. We are

grateful to the European Research Council (No. 101019223) for financial support.

References

1. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-
transmitted helminth infections: ascariasis, trichuriasis, and hookworm. The Lancet.
2006 May 6;367(9521):1521–32.

2. Pullan RL, Smith JL, Jasrasaria R, Brooker SJ. Global numbers of infection and disease
burden of soil transmitted helminth infections in 2010. Parasites Vectors. 2014 Jan
21;7(1):37.

3. Bundy DAP, Cooper ES. *Trichuris* and Trichuriasis in Humans. In: Baker JR, Muller R,
editors. Advances in Parasitology [Internet]. Academic Press; 1989 [cited 2024 Jul 18].
p. 107–73. Available from:
https://www.sciencedirect.com/science/article/pii/S0065308X08603322

4. Callejón R, Robles MDR, Panei CJ, Cutillas C. Molecular diversification of Trichuris spp.
from Sigmodontinae (Cricetidae) rodents from Argentina based on mitochondrial DNA
sequences. Parasitol Res. 2016 Aug 1;115(8):2933–45.

5. Cutillas C, de Rojas M, Zurita A, Oliveros R, Callejón R. Trichuris colobae n. sp.
(Nematoda: Trichuridae), a new species of Trichuris from Colobus guereza kikuyensis.
Parasitol Res. 2014 Jul 1;113(7):2725–32.

6. Callejón R, Halajian A, Cutillas C. Description of a new species, *Trichuris ursinus* n. sp.
(Nematoda: Trichuridae) from *Papio ursinus* Keer, 1792 from South Africa. Infection,
Genetics and Evolution. 2017 Jul 1;51:182–93.

7. Oliveros R, Cutillas C, De Rojas M, Arias P. Characterization of four species of Trichuris
(Nematoda: Enoplida) by their second internal transcribed spacer ribosomal DNA
sequence. Parasitol Res. 2000 Nov 1;86(12):1008–13.

8. Ooi HK, Tenora F, Itoh K, Kamiya M. Comparative Study of Trichuris trichiura from Non-
Human Primates and from Man, and Their Difference with T. suis. Journal of Veterinary
Medical Science. 1993;55(3):363–6.

9. Ghai RR, Simons ND, Chapman CA, Omeja PA, Davies TJ, Ting N, et al. Hidden
Population Structure and Cross-species Transmission of Whipworms (Trichuris sp.) in
Humans and Non-human Primates in Uganda. Gasser RB, editor. PLoS Negl Trop Dis.
2014 Oct 23;8(10):e3256.

10. Rivero J, Cutillas C, Callejón R. Trichuris trichiura (Linnaeus, 1771) From Human and
Non-human Primates: Morphology, Biometry, Host Specificity, Molecular
Characterization, and Phylogeny. Front Vet Sci. 2021 Feb 9;7:626120.

11. Cavallero S, De Liberato C, Friedrich KG, Di Cave D, Masella V, D'Amelio S, et al.
Genetic heterogeneity and phylogeny of Trichuris spp. from captive non-human primates
based on ribosomal DNA sequence data. Infection, Genetics and Evolution. 2015
Aug;34:450–6.

12. Cavallero S, Montalbano Di Filippo M, Rondón S, Liberato CD, D'Amelio S, Friedrich
KG, et al. Nuclear and Mitochondrial Data on Trichuris from Macaca fuscata Support
Evidence of Host Specificity. Life. 2020 Dec 31;11(1):18.

13. Nissen S, Al-Jubury A, Hansen TVA, Olsen A, Christensen H, Thamsborg SM, et al.
Genetic analysis of Trichuris suis and Trichuris trichiura recovered from humans and
pigs in a sympatric setting in Uganda. Veterinary Parasitology. 2012 Aug;188(1–2):68–
77.

14. Ravasi DF, O'Riain MJ, Davids F, Illing N. Phylogenetic Evidence That Two Distinct
Trichuris Genotypes Infect both Humans and Non-Human Primates. Parkinson J, editor.
PLoS ONE. 2012 Aug 28;7(8):e44187.

15. Xie Y, Zhao B, Hoberg EP, Li M, Zhou X, Gu X, et al. Genetic characterisation and
phylogenetic status of whipworms (Trichuris spp.) from captive non-human primates in
China, determined by nuclear and mitochondrial sequencing. Parasites Vectors. 2018
Dec;11(1):516.

16. Venkatesan A, Chen R, Baer M, Schneeberger PHH, Reimer B, Hürlimann E, et al.
Discovery of a novel albendazole-ivermectin refractory Trichuris species, T. incognita, in
Côte d'Ivoire patients through fecal DNA metabarcoding [Internet]. bioRxiv; 2024 [cited
2024 Jul 16]. p. 2024.07.04.602123. Available from:
https://www.biorxiv.org/content/10.1101/2024.07.04.602123v2

17. Bär MA, Kouamé NA, Touré S, Coulibaly JT, Schneeberger PHH, Keiser J. Genetic
makeup of *Trichuris incognita* , a novel *Trichuris* species naturally infecting humans and
displaying signs of resistance to drug treatment [Internet]. 2024 [cited 2024 Jul 30].
Available from: http://biorxiv.org/lookup/doi/10.1101/2024.06.11.598441

18. Hürlimann E, Keller L, Patel C, Welsche S, Hattendorf J, Ali SM, et al. Efficacy and
safety of co-administered ivermectin and albendazole in school-aged children and adults
infected with Trichuris trichiura in Côte d'Ivoire, Laos, and Pemba Island, Tanzania: a
double-blind, parallel-group, phase 3, randomised controlled trial. The Lancet Infectious
Diseases. 2022 Jan;22(1):123–35.

19. Hürlimann E, Palmeirim, Marta. Efficacy of Ivermectin-albendazole vs Albendazole
Alone in School-aged Children Infected With Trichuris Trichiura in Uganda. 2024.

20. Keller L, Hürlimann E, Patel C, Welsche S, Sayasone S, Ali SM, et al. Insights gained
from conducting a randomised controlled trial on Ivermectin-Albendazole against
Trichuris trichiura in Côte d'Ivoire, Lao PDR and Pemba Island. In: Rollinson D, Stothard
JR, editors. Advances in Parasitology [Internet]. Academic Press; 2021 [cited 2024 Jul
25]. p. 253–76. Available from:
https://www.sciencedirect.com/science/article/pii/S0065308X2030110X

21. Keller L, Welsche S, Patel C, Sayasone S, Ali SM, Ame SM, et al. Long-term outcomes
of ivermectin-albendazole versus albendazole alone against soil-transmitted helminths:
Results from randomized controlled trials in Lao PDR and Pemba Island, Tanzania.
Fairfax KC, editor. PLoS Negl Trop Dis. 2021 Jun 30;15(6):e0009561.

22. Patel C, Hürlimann E, Keller L, Hattendorf J, Sayasone S, Ali SM, et al. Efficacy and
safety of ivermectin and albendazole co-administration in school-aged children and
adults infected with Trichuris trichiura: study protocol for a multi-country randomized
controlled double-blind trial. BMC Infect Dis. 2019 Dec;19(1):262.

23. Foth BJ, Tsai IJ, Reid AJ, Bancroft AJ, Nichol S, Tracey A, et al. Whipworm genome and
dual-species transcriptome analyses provide molecular insights into an intimate host-
parasite interaction. Nat Genet. 2014 Jul;46(7):693–700.

24. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite – a
comprehensive resource for helminth genomics. Molecular and Biochemical
Parasitology. 2017 Jul 1;215:2–10.

25. Jex AR, Nejsum P, Schwarz EM, Hu L, Young ND, Hall RS, et al. Genome and
transcriptome of the porcine whipworm Trichuris suis. Nat Genet. 2014 Jul;46(7):701–6.

26. Liu GH, Gasser RB, Nejsum P, Wang Y, Chen Q, Song HQ, et al. Mitochondrial and
Nuclear Ribosomal DNA Evidence Supports the Existence of a New Trichuris Species in
the Endangered François' Leaf-Monkey. Janke A, editor. PLoS ONE. 2013 Jun
20;8(6):e66249.

27. Yao C, Walkush J, Shim D, Cruz K, Ketzis J. Molecular species identification of Trichuris
trichiura in African green monkey on St. Kitts, West Indies. Vet Parasitol Reg Stud
Reports. 2018 Jan;11:22–6.

28. Liu GH, Zhou W, Nisbet AJ, Xu MJ, Zhou DH, Zhao GH, et al. Characterization of
Trichuris trichiura from humans and *T. suis* from pigs in China using internal transcribed
spacers of nuclear ribosomal DNA. J Helminthol. 2014 Mar;88(1):64–8.

29. Sirima C, Bizet C, Hamou H, Červená B, Lemarcis T, Esteban A, et al. Soil-transmitted
helminth infections in free-ranging non-human primates from Cameroon and Gabon.
Parasites Vectors. 2021 Dec;14(1):354.

30. Hawash MBF, Betson M, Al-Jubury A, Ketzis J, LeeWillingham A, Bertelsen MF, et al.
Whipworms in humans and pigs: origins and demography. Parasites Vectors. 2016
Dec;9(1):37.

31. Nejsum P, Betson M, Bendall RP, Thamsborg SM, Stothard JR. Assessing the zoonotic
potential of Ascaris suum and Trichuris suis: looking to the future from an analysis of the
past. Journal of Helminthology. 2012 Jun;86(2):148–55.

32. Hawash MBF, Andersen LO, Gasser RB, Stensvold CR, Nejsum P. Mitochondrial
Genome Analyses Suggest Multiple Trichuris Species in Humans, Baboons, and Pigs
from Different Geographical Regions. Lok JB, editor. PLoS Negl Trop Dis. 2015 Sep
14;9(9):e0004059.

33. Beleckė A, Kupčinskas T, Stadalienė I, Höglund J, Thamsborg SM, Stuen S, et al.
Anthelmintic resistance in small ruminants in the Nordic-Baltic region. Acta Veterinaria
Scandinavica. 2021 Apr 27;63(1):18.

34. Fairweather I, Brennan GP, Hanna REB, Robinson MW, Skuce PJ. Drug resistance in
liver flukes. International Journal for Parasitology: Drugs and Drug Resistance. 2020 Apr
1;12:39–59.

35. Kelleher AC, Good B, de Waal T, Keane OM. Anthelmintic resistance among
gastrointestinal nematodes of cattle on dairy calf to beef farms in Ireland. Irish Veterinary
Journal. 2020 Jul 1;73(1):12.

36. Rose Vineer H. What Modeling Parasites, Transmission, and Resistance Can Teach Us.
Veterinary Clinics of North America: Food Animal Practice. 2020 Mar 1;36(1):145–58.

37. Kotze AC, Gilleard JS, Doyle SR, Prichard RK. Challenges and opportunities for the
adoption of molecular diagnostics for anthelmintic resistance. International Journal for
Parasitology: Drugs and Drug Resistance. 2020 Dec 1;14:264–73.

38. Else KJ, Keiser J, Holland CV, Grencis RK, Sattelle DB, Fujiwara RT, et al. Whipworm
and roundworm infections. Nat Rev Dis Primers. 2020 May 28;6(1):44.

39. World Health Organization. Ending the neglect to attain the Sustainable Development
Goals: A road map for neglected tropical diseases 2021–2030 [Internet]. Geneva: World
Health Organization; 2021 Jan [cited 2024 Jul 22]. Available from:
https://www.who.int/publications/i/item/9789240010352

40. Ceballos-Mendiola G, Valero A, Polo-Vico R, Tejada M, Abattouy N, Karl H, et al.
Genetic variability of Anisakis simplex s.s. parasitizing European hake (Merluccius
merluccius) in the Little Sole Bank area in the Northeast Atlantic. Parasitol Res. 2010
Nov 1;107(6):1399–404.

41. Cutillas C, Callejón R, de Rojas M, Tewes B, Ubeda JM, Ariza C, et al. *Trichuris suis*
and *Trichuris trichiura* are different nematode species. Acta Tropica. 2009 Sep
1;111(3):299–307.

42. Wickramasinghe S, Yatawara L, Rajapakse RPVJ, Agatsuma T. Toxocara canis and
Toxocara vitulorum: molecular characterization, discrimination, and phylogenetic
analysis based on mitochondrial (ATP synthase subunit 6 and 12S) and nuclear
ribosomal (ITS-2 and 28S) genes. Parasitol Res. 2009 May 1;104(6):1425–30.

43. George S, Geldhof P, Albonico M, Ame SM, Bethony JM, Engels D, et al. Molecular
speciation of soil-transmitted helminths egg isolates collected during six drug efficacy
trials in endemic countries. Trans R Soc Trop Med Hyg. 2016 Nov 1;110(11):657–63.

44. Hassan NA, Noor Badi FA, Mohd-Shaharuddin N, Wan Yusoff WS, Lim Y a. L, Chua KH,
et al. A conventional multiplex PCR for the detection of four common soil-transmitted
nematodes in human feces: development and validation. Trop Biomed. 2022 Mar
1;39(1):135–43.

45. Alomashi GBA, Khudhur HR, Jebur LS. Conventional Multiplex PCR for Identifying Soil
Transmitted Helminthes (*Ascaris Lumbricoides*, *Trichuris Trichura* and *Ancylostoma*
Duodenale) in Fecal Specimens. Nano Biomedicine and Engineering. 2022 Dec
31;14(4):289–94.

46. Kaisar MMM, Brienen E a. T, Djuardi Y, Sartono E, Yazdanbakhsh M, Verweij JJ, et al.
Improved diagnosis of Trichuris trichiura by using a bead-beating procedure on ethanol
preserved stool samples prior to DNA isolation and the performance of multiplex real-
time PCR for intestinal parasites. Parasitology. 2017 Jun;144(7):965–74.

47. Ayana M, Cools P, Mekonnen Z, Biruksew A, Dana D, Rashwan N, et al. Comparison of
four DNA extraction and three preservation protocols for the molecular detection and
quantification of soil-transmitted helminths in stool. Cantacessi C, editor. PLoS Negl
Trop Dis. 2019 Oct 28;13(10):e0007778.

48. Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et al. Ultra-
high-throughput microbial community analysis on the Illumina HiSeq and MiSeq
platforms. The ISME Journal. 2012 Aug 1;6(8):1621–4.

49. Dommann J, Kerbl-Knapp J, Albertos Torres D, Egli A, Keiser J, Schneeberger PHH. A
novel barcoded nanopore sequencing workflow of high-quality, full-length bacterial 16S
amplicons for taxonomic annotation of bacterial isolates and complex microbial
communities. mSystems. 2024 Sep 10;0(0):e00859-24.

50. Shen W, Le S, Li Y, Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q
File Manipulation. Zou Q, editor. PLoS ONE. 2016 Oct 5;11(10):e0163962.

51. De Coster W, Rademakers R. NanoPack2: population-scale evaluation of long-read
sequencing data. Alkan C, editor. Bioinformatics. 2023 May 4;39(5):btad311.

52. Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, et al. High-throughput
amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution.
Nucleic Acids Research. 2019 Oct 10;47(18):e103–e103.

53. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis
Version 11. Battistuzzi FU, editor. Molecular Biology and Evolution. 2021 Jun
25;38(7):3022–7.

54. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics. 2014 May 1;30(9):1312–3.

55. Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. ModelTest-NG: A
New and Scalable Tool for the Selection of DNA and Protein Evolutionary Models.
Crandall K, editor. Molecular Biology and Evolution. 2020 Jan 1;37(1):291–4.

56. Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-
Onsins SE, et al. DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data Sets.
Molecular Biology and Evolution. 2017 Dec 1;34(12):3299–302.

57. Excoffier L, Lischer HEL. Arlequin suite ver 3.5: a new series of programs to perform
population genetics analyses under Linux and Windows. Molecular Ecology Resources.
2010 May;10(3):564–7.

58. Leigh JW, Bryant D. popart: full-feature software for haplotype network construction.
Methods in Ecology and Evolution. 2015;6(9):1110–6.