Comparative lifespan and healthspan of nonhuman primate species common to biomedical research

Hillary F Huber[1]*, Hannah C Ainsworth[2]*, Ellen Quillen[2], Adam Salmon[3], Corinna Ross[1], Adinda D Azhar[4],

Karen Bales[5,6], Michele A Basso[7], Kris Coleman[8,9], Ricki Colman[10], Huda S Darusman[4,11], William Hopkins[12,13],

Charlotte E Hotchkiss[7], Matt Jorgensen[2], Kylie Kavanagh[2,15], Cun Li[14], Julie A Mattison[16], Peter W

Nathanielsz[1,14], Suryo Saputro[4], Diana Scorpio[1,17], Paul-Michael Sosa[5], Eric Vallender[18,19], Yaomin Wang[2],

Caroline Zeiss[20], Carol Shively[2]*, Laura A Cox[2]*

*Equal contributions

[1]Texas Biomedical Research Institute, San Antonio, TX, USA

[2]Wake Forest Univ School of Medicine, Winston-Salem, NC, USA

[3]University of Texas Health Science Center, San Antonio, TX, USA

[4]Primate Research Center IPB University, Bogor, Indonesia

[5]California National Primate Research Center, Davis, CA, USA

[6]University of California, Davis, CA, USA

[7]Washington National Primate Research Center, Seattle, WA, USA

[8]Oregon National Primate Research Center, Hillsboro, OR, USA

[9]Oregon Health & Science University, Portland, OR, USA

[10]Wisconsin National Primate Research Center, Madison, WI, USA

[11]School of Veterinary Medicine and Biomedical Sciences IPB University, Bogor, Indonesia

[12]The University of Texas MD Anderson Cancer Center, Bastrop, TX, USA

[13]Emory National Primate Research Center, Atlanta, GA, USA

[14]University of Wyoming, Laramie, WY, USA

[15]University of Tasmania, Hobart, Tasmania, Australia

[16]National Institute on Aging, National Institutes of Health, Gaithersburg, MD, USA

[17]Envol Biomedical, Immokalee, FL, USA

18Tulane National Primate Research Center, Covington, LA, USA

19New England Primate Research Center, Southborough, MA, USA

20Yale University School of Medicine, New Haven, CT, USA

Abstract

There is a critical need to generate age- and sex-specific survival curves to characterize chronological aging consistently across nonhuman primates (NHP) used in biomedical research. Accurate measures of chronological aging are essential for inferences into genetic, demographic, and physiological variables driving differences in NHP lifespan within and between species. Understanding NHP lifespans is relevant to public health because unraveling the demographic, molecular, and clinical bases of health across the life course in translationally relevant NHP species is fundamentally important to the study of human aging. Data from more than 110,000 captive individual NHP were contributed by 15 major research institutions to generate sex-specific Kaplan-Meier survival curves using uniform methods in 12 translational aging models: *Callithrix jacchus* (common marmoset), *Chlorocebus aethiops sabaeus* (vervet/African green), *Macaca fascicularis* (cynomolgus macaque), *M. fuscata* (Japanese macaque), *M. mulatta* (rhesus macaque), *M. nemestrina* (pigtail macaque), *M. radiata* (bonnet macaque), *Pan troglodytes* spp. (chimpanzee), *Papio hamadryas* spp. (baboon), *Plecturocebus cupreus* (coppery titi monkey), *Saguinus oedipus* (cotton-top tamarin), and *Saimiri* spp. (squirrel monkey). After employing strict inclusion criteria, primary analysis results are based on 12,269 NHP that survived to adulthood and died of natural/health-related causes. A secondary analysis was completed for 32,616 NHP that died of any cause. For the primary analyses, we report ages of 25th, 50th, 75th, and 85th percentiles of survival, maximum observed ages, rates of survivorship, and sex-based differences captured by quantile regression models and Kolmogorov-Smirnov tests. Our findings show a pattern of reduced male survival among catarrhines (African and Asian primates), especially macaques, but not platyrrhines (Central and South American primates). For many species, median lifespans were lower than previously reported. An important consideration is that these analyses may offer a better reflection of healthspan than lifespan. Captive NHP used in research are typically euthanized for humane welfare reasons before their natural end of life, often after diagnosis of their first major disease requiring long-term treatment with reduced quality of life (e.g., endometriosis, cancer, osteoarthritis).

Supporting the idea that these data are capturing healthspan, for several species typical age at onset of chronic disease is similar to the median lifespan estimates. This data resource represents the most comprehensive characterization of sex-specific lifespan and age-at-death distributions for 12 biomedically relevant species, to date. The results clarify the relationships among NHP ages and will provide a valuable resource for the aging research community, improving human-NHP age equivalencies, informing investigators of the expected survival rates of NHP assigned to studies, providing a metric for comparisons in future studies, and contributing to our understanding of the factors that drive lifespan differences within and among species.

Introduction

Nonhuman primates (NHPs) are genetically, physiologically, and behaviorally the best translational models for human aging as their genomes, developmental trajectory, reproductive strategies, and aging-related changes in physical function, cognitive function, and disease development are more similar to humans than those of other mammals.[1–4] Yet we know little about longevity in the NHPs most commonly used as translational models. Few studies have attempted cross-species comparisons and reports are often contradictory, likely due to the use of different methodological approaches (e.g., inclusion criteria). To determine how NHP ages correspond with human age, it is essential to fully characterize the demography of NHP longevity within each species, rather than focusing on individual reports of maximum longevity. Numerous publications list NHP maximum lifespans in tables that include a variety of other life history features, but few cite primary sources. This leads to overreporting of the same statistics without verifying the validity of the measure or the relevance to animals under study. For example, 37.5 years is often cited as the lifespan of baboons (*Papio hamadryas* spp.).[5–8] However, tracing citations to the primary source reveals that this statistic comes from a single baboon that died at the Brookfield Zoo in 1972; the birth date is given as June 1, 1935 (one year after the zoo opened), but it is not documented whether this date is known or estimated.[9] This estimate of maximum longevity in baboons is not particularly useful without additional context indicating how many baboons survive to the maximum or what the median baboon lifespan is. Median captive baboon lifespan has been reported as 21[10] or 11[11] years but the report of maximum longevity is more frequently cited. It is likely that the discrepancy in median baboon lifespan reflects differences in methodological approaches to data analysis. This example in

baboons highlights how differences in analytic approaches across studies make it difficult to compare reports

within or across species. The unclear data on NHP lifespan, such as the reporting of maximum longevity to

indicate "lifespan," creates confusion in scientific analysis and in the peer review process.

Cross-species comparisons are a major goal of aging research since they can reveal factors contributing to

variation in lifespans. Inconsistent lifespan estimates are problematic when looking at a single species, and the

problem is compounded by cross-species comparisons. We address this knowledge gap by creating rigorous

and reproducible survivorship data, identifying mortality risk and its relationship to biological age at different

chronological ages, and examining the shape of mortality and healthspan curves across 12 captive NHP

species. The initial dataset, prior to quality control and filtering, included lifespan data from 114,255 animals

from 58 species at 15 institutions. We highlight that while maximum age is an easily reported statistic as it is

purely observational, calculating median lifespan is more challenging, as methodological decisions about

inclusion and exclusion criteria vary among studies, producing substantial discrepancies across cohorts and

species. With the data herein, we have the unique ability to calculate survival probabilities using the same

criteria for all 12 species, producing the most methodologically consistent cross-species comparison to date.

The value of such a large dataset is the ability to filter the data to the most representative sample and retain

adequate sample sizes for statistical analyses. In this study, survival curves were generated on animals that

survived to at least adulthood (defined in Methods) because, as in most mammals including humans, risk of

death in infancy is substantial and strongly biases the median lifespan. Primary results and comparisons by

sex are built using data from animals that died of natural causes or were euthanized for clinical/health reasons.

This report provides comprehensive data summaries and tools to improve biomedical research involving NHPs

within and beyond the field of aging.

Results

Primary analyses. Sample counts of primary analysis datasets, featuring natural or health-related deaths only,

are shown in **Table 1**. Maximum observed age including all types of deaths (e.g., research-related sacrifice,

clinical/health-related euthanasia, and natural), as well as median age at death calculated from only natural

and clinical deaths, are summarized by sex and species in **Table 2**. **Figure 1** shows a scatter plot of datapoints

for natural and clinical deaths, with medians, interquartile ranges, and proportions of data by sex and species.

Combined survival curves for all 12 species in males and females are shown in **Figure 2**. To evaluate the rate

of decline for the survivorship curves, data from the first and last quartiles of the Kaplan-Meier survivorship

function were fit to an exponential model that captures rate of decay (i.e., change in probability of death), and

species were then compared within and between sexes. Comparing first and last quartiles illustrated that

species predominantly experienced faster rates of death within the first quartile of adulthood. Comparing male

and female rates of decline within both quartiles highlighted the faster rates of decline for males within the first

quartile. However, in the last quartile, this pattern was nearly reversed; the majority of species (except cotton-

top tamarin, vervet/African green monkey, and common marmoset) exhibited slower rates of decline in males

compared to females (**Figure 3**).

For each species, individual survival curves are shown in **Figure 4** and sex-based comparisons in **Table 3**. In

most species, males showed reduced survival compared to females. Among vervets, Japanese macaques,

and chimpanzees, males showed reduced survival at every age with a different overall distribution of age at

death. *C*ynomolgus macaque and baboon males showed reduced survival compared to females at younger

ages (25th and 50th percentiles), but there was no difference in survival at later stages of life. Rhesus macaque

males showed reduced survival compared to females at the 25th, 50th, and 75th percentiles, but females had

lower age of survival at the 85th percentile. There was a strong difference in the distribution of age at death

between males and females (P-value=2.20×10^{-16}). Pig-tailed macaque males showed reduced survival

compared to females early in life (25%) but the sexes were similar at other ages. In contrast, females showed

reduced survival compared to males at every age in common marmosets. Male and female survival was similar

at every age with no difference in the distribution of age at death between sexes for cotton-top tamarins and

squirrel monkeys. There was also no difference in distributions for coppery titi monkeys and bonnet macaques;

however, the modest sample size for the species limits power to detect small differences.

Secondary analyses. *C*ensored data (deaths due to research sacrifice and colony management) were biased

by sex (**Extended Data Figure 1**) and prevented statistical comparisons between males and females when

including censored data.[12] However, as a secondary analysis, survival curves that include censored events are

presented as extended data (**Extended Data Figure 2**) for reference. Across species, inclusion of additional

datapoints from censored events increased median lifespan estimates. We note that the high proportion of

censored events (**Extended Data Figure 1**), especially in some species (i.e., greater than 50% of deaths in

baboons, cynomolgus, pigtails, rhesus, squirrel monkeys, and vervets), yielded survivorship functions that

never reach zero, limiting utility and inference for the full lifespan.

144   **Discussion**

Lifespan vs healthspan. A major consideration of note for this study is that few research NHPs live until

natural death. Most are humanely euthanized due to study protocols or clinical determinations based on quality

of life. The issues considered by veterinarians in making euthanasia decisions vary by facility and study

protocol, but a common approach is to euthanize at the first diagnosis of major disease or injury requiring long-

term treatment with reduced quality of life. Reasons for humane euthanasia may include such diverse

conditions as advanced spinal or knee osteoarthritis, endometriosis, broken limbs, tumors, and meningitis – not

all of which are the result of aging-related diseases. Therefore, we posit that these findings may be measuring

healthspan rather than lifespan in NHP cohorts housed at research facilities. For our survival analyses, this

potential limitation is partially mediated by our very large database, which enabled analyses even after

removing experimental and other non-clinical deaths.

Supporting the idea that we are measuring healthspan rather than lifespan, for several species, typical age at

onset of chronic disease is similar to the median lifespan estimates. Among baboons, age-related diseases are

apparent around 9 years old (e.g., edema, kyphosis, prolapse, myocarditis), and by 12 years many more are

evident (e.g., pancreatitis, stricture, lymphosarcoma).[13] Median baboon lifespan in this report is 10.1 years for

males and 11.1 years for females. Marmoset age-related diseases tend to emerge in animals >6 years old,

including cardiovascular disease, diabetes, and neoplasias.[14] Median marmoset lifespan in our study is 5.5

years in males and 5.0 years in females. Rhesus macaques are on average diagnosed with the first chronic condition at age 9.0 years and the second at age 10.7 years.[15] Median rhesus lifespan in our study is 9.1 years in males and 10.6 years in females. Differences in veterinary care for these conditions mean that some pathologies in some species may be treated medically, whereas others proceed to veterinarian-suggested euthanasia. We speculate that zoo NHPs may be treated for more chronic conditions than research NHPs and would make a useful lifespan and healthspan comparison to humans.

The ability to make more accurate comparisons between NHP age and the human equivalent was a primary goal of the current analyses. Since the NHP estimates herein may be closer to healthspan than lifespan, it is useful to consider them in relation to human healthspan. The most frequently studied measures of human healthspan are deficit accumulation indices, which measure accumulation of health deficits and decline in physical function or frailty.[16–20] In one study of 66,589 Canadians in the National Population Health Survey, accumulation of health deficits was gradual before age 46 years, with 40% of 45-50 year-olds having a frailty index score of 0 (no health deficits); starting at age 46, deficit accumulation was much more rapid, and at age 80, only 5% still had a score of 0.[20,21] Among 73,396 people from the Longitudinal Ageing Study in India, average age of onset of any chronic disease was 53 years.[22] We speculate that our NHP median lifespan estimates may align better with human onset and accumulation of health deficits, rather than human lifespan. However, our analysis does not address onset of health deficits, and we are unable to distinguish between which NHPs died at the end of their lifespan versus those which died at the end of their healthspan. Therefore, we are unable to make specific comparisons between human and NHP healthspans.

Sources of variation within and between species. Our findings show great variation in adult life expectancy among all 12 species, in contrast to a prior cross-species analysis of six primate species that found little variation in adult survival.[23] Many factors contribute to variation in adult survival. Some may assume that in captive research populations, quality of veterinary care is a major driving force. While this may have been important in the early years of NHP research, most species have been in captivity for decades and quality care is well defined. Institutional management practices are important factors, such as how decisions are made

about euthanizing animals due to illness or reproductive capacity. The goals of the research are also important

to consider. For example, rhesus monkeys have been the subjects in two longevity studies in which survival

time was an outcome variable. Here, additional measures were taken to maintain older animals, which explains

the extreme maximum age of rhesus macaques – 44.2 years – relative to other the other four macaque

species, which show maximum ages in the 20s and 30s.[24,25] Another potential source of bias is the way

animals are selected for studies. NHPs go through health checks beforehand, and healthy animals may be

preferentially selected. In our study, many of the longest-lived animals were excluded from lifespan

calculations because their endpoints were research-related (**Extended Data Figure 1**). Thus, limiting the

analyses to natural deaths seems to influence lifespan calculations towards younger ages.

Within species, life history features can influence lifespan. It has been proposed that reproductive strategies

play an evolutionary role in regulating lifespan, since there may be tradeoffs between female fertility,

investment in offspring, and longevity,[26] although this long-held view has been challenged since the

relationships between reproduction and longevity are not consistent across species.[27,28] Adult body size also

factors into survival because a longer period of growth will likely result in later reproductive maturity and a

greater need for investment in offspring. In our data, common marmosets have the shortest maximum and

median lifespan of all 12 species. Marmosets are also the smallest species (average weight 350-400 g), reach

adulthood at the youngest age (1.5 years), and usually give birth to twins.[14,29] However, cotton-top tamarins,

the other small (average weight in captivity 565.7 g), quickly maturing (2.5 years at adulthood), twinning

callitrichine[30] in this study, has maximum and median lifespan resembling that of several larger bodied, slower

maturing species that give birth to singletons, including squirrel monkeys, baboons, vervets, and macaques. It

is unclear to what extent these patterns are driven by inherent species characteristics versus institutional

practices, but it would be advantageous to explore this question in future studies.

Sex-based differences. Among primates, males have been shown to have higher age-specific mortality than

females throughout adulthood.[31] We see this in some species included in the current study. One pattern is

shorter lifespan among macaque males. Five macaque species (*Macaca* spp.) are reported here. In three

species males have shorter median lifespan than females (cynomolgus, Japanese, and rhesus macaques). In pigtails, males have lower survival probability in early adulthood (25%) but similar survival probability at older ages, and in bonnet macaques male lifespan appears shorter in the curves and estimates, but sample size may be too small to detect a difference (female n=43, male n=19). This pattern seems to extend to all of the parvorder Catarrhini (Old World monkeys- Cercopithecoidea and apes- Hominoidea). Vervets have the largest sex-based differential with median age of 8.3 years for males and 17.9 years for females. For baboons, males show borderline lower survival probability at the 25[th] and 75[th] percentiles. Male chimpanzees also have lower survival probability relative to females at every life stage.

In contrast, in the parvorder Platyrrhini (Central and South American monkeys), there is generally no difference between males and females in survival estimates. For context, a phylogenetic tree for the 12 species in this study is shown in **Extended Data Figure 3**.[32] The exception is the common marmoset, with lower female survival at every age, replicating the findings of another marmoset report.[14] The relatively short female marmoset lifespan is related to their high fertility rates.[26,29] There are no differences in survival between males and females in coppery titi monkeys, squirrel monkeys, or cotton-top tamarins. A prior primate lifespan comparison that suggested female primates have longer lifespan than males included several catarrhine species but few data from platyrrhine species.[31] A recent study of coppery titi monkey lifespan showed a trend toward longer lifespan in males relative to females using the same population of monkeys in the current study but with different inclusion criteria.[33]

It is difficult to know if the observed sex-based differences between catarrhine versus platyrrhine species are due to inherent species characteristics, institutional practices, or their interactions. For example, in catarrhine monkeys, it is common to house a single breeding or vasectomized male with multiple females. Fewer males than females are needed for breeding programs because males will mate with multiple females. In some species, especially baboons, males are much larger than females, requiring more space and resources. These factors and more mean males and females are not equally distributed and are subject to different animal selection practices in research institutions. The difference is also evident in the sample size. Before data

filtering, the sample size included 44,704 females and 43,413 males. After data filtering, there were 8,296

females and 3,973 males. A larger proportion of the males were filtered out of the analyses because of

research-related endpoints or humane euthanasia for management reasons, reflecting bias in how sexes are

deployed in research.

Comparison with prior reports of captive NHP lifespan. As mentioned in the introduction, captive baboon

maximum lifespan has been reported as 37.5 years,[5–8] and median lifespan as 21[10] or 11[11] years. Our median

lifespan findings align with the lowest of those estimates, and close inspection of the methods used to arrive at

that estimate reveals that the study employed similar inclusion and exclusion criteria as the current study.[11]

The 37.5 year estimate is based on a single zoo baboon[9] and is a rare case of extreme maximum longevity.

The 21-year baboon lifespan estimate uses different methods from the current study, such as inclusion of live

animals as right censored datapoints.[10] In another report that includes 4,480 zoo baboons, male *P. hamadryas*

were estimated to live 13.2 years and females 17.1 years from birth.[23] We expect that this difference is due to

both methodological differences in calculating median lifespan and differences in the veterinary care for the

small numbers of baboons in zoo settings, e.g., they frequently receive long-term treatment for chronic

diseases. It may also be due to differences between hamadryas and the mixed baboons in our study. Prior

reports of lifespan of rhesus macaques have hovered around a median lifespan of 25 years and maximum 40

years, but again, these studies employed right censored data approaches.[24,34–36] In contrast, our median

lifespan estimate for rhesus is 7.9 years in males and 10.3 years in females using data only from animals with

known ages at death, rather than including ages from still living animals with a right censored approach. To

highlight this methodological difference, we provide survivorship probabilities with censored data for reference

(**Extended Data Figure 2**). A prior study of common marmosets at a single institution estimated median

lifespan of 6.5 years in animals that survived to at least two years (compared with our starting age of 1.5

years).[14] Another marmoset study from a different institution estimated median lifespan at four years in

marmosets that survived for 60 days; the same study reported cotton-top tamarin median life expectancy of 7.2

years.[37] Our estimates from marmosets at 4 different institutions are 5.3 years in females and 6.0 years in

males. For cotton-top tamarins, our estimates of median lifespan (from animals living at one institution) are 9.6

years for males and 8.9 years for females. Chimpanzee median survival in a biomedical research population has been reported as 31.0 years in males and 38.8 years in females among individuals who reached 1 year of age.[38] In a zoo population, male chimpanzees lived a median of 26.0 years and females 30.5 years from birth.[23] Our estimates are 33.0 years in males and 44.0 years in females among individuals who reached ten years of age and are therefore fairly consistent with previous reports. For coppery titi monkeys, median lifespan has been reported as 14.9 years in males and 11.4 years in females among individuals surviving to 31 days,[33] compared with our estimates of 8.6 years for males and 9.2 years for females. Once again, the differences between estimates in our studies and prior reports likely arise methodologically, such as choices made about age of inclusion and use of a right censored approach to include individuals still alive and/or those euthanized for research-related endpoints. A major strength of the current study is the use of uniform methods across 12 different NHP species.

Importance of data filtering. This study highlights the necessity of thorough methodological documentation in NHP lifespan studies. As illustrated with our primary and secondary analyses, filtering and methodological decisions impact the results and interpretation. The simplest example is the minimum age threshold for computing the survivorship functions. Including juveniles dramatically lowers median lifespan due to high rates of juvenile mortality among primates. Additionally, by including only animals that were born and died at the same institute, it sometimes eliminated the oldest known individuals from the dataset, such as two 19-year-old SNPRC marmosets; however, these instances were rare in our very large sample. Decisions that greatly reduced our analysis sample size, such as date-of-birth (DOB) cutoffs, are a privilege of a large initial (pre-filtered) dataset. So, while the DOB cutoffs greatly reduced our final sample size, it removed bias associated with very early deaths (since our dataset did not include currently alive animals). Overall, given the impact of filtering decisions, we emphasize the need for robust reporting of the decision criteria in NHP survival studies. We encourage authors to follow the ARRIVE guidelines (Animal Research: Reporting of In Vivo Experiments; https://arriveguidelines.org/), a checklist for full and transparent reporting aimed at improving rigor, transparency, and reproducibility in animal research.[39] In longevity research, it is particularly crucial to report inclusion and exclusion criteria in addition to the details of statistical approaches.

Limitations. One limitation of the study is that the stringent inclusion criteria reduced our starting sample size

by 86%. This was necessary to ensure appropriate comparisons across institutions and species. For example,

some species (cynomolgus, pigtails, baboons) have a very high percentage of deaths by research sacrifice,

rather than by natural or health-related causes. Including research-related deaths as right censored data

results in highly skewed models with limited utility for these species (e.g., survival curves for female baboons

do not converge past the median survivorship when including censored data). Further, censoring was biased

by sex because of the differences in research utilization and breeding needs, statistically hindering the

possibility of comparisons between males and females. Therefore, primary analyses were limited to data from

natural or clinical deaths, eliminating the need for right censoring. Another constraint of the study is our limited

knowledge of specific cause of death. Differences in institutional death coding systems make it difficult to easily

determine cause of death, since some record systems group many types of deaths, while others have more

granular codes to distinguish among death types.

Conclusions. The need for comparative analyses of lifespans across species has been widely

acknowledged.[40] Investigators need access to reliable lifespan tables, survivorship graphs, and maximum

lifespan measurements to conduct relevant translational aging studies. Here we provide the largest dataset yet

assembled from captive research NHPs. These data provide a valuable comparative resource for translational

NHP research, primary data on multispecies NHP lifespan in captivity, and context for consideration of

morbidity and mortality in the study of diverse diseases.

Methods

Species

Twelve NHP species for analyses are shown in **Table 1**. We are considering all members of the genus *Papio* a

single species and considering Indian- and Chinese-origin rhesus macaques together, as captive research

baboons have a high degree of morphotype mixing[41,42] and captive rhesus are similarly highly admixed from

these geographic source populations.[43] We included chimpanzees (*Pan troglodytes* spp.), but it must be noted

that biomedical research with great apes is heavily restricted across the world. Still, many retired chimpanzees

reside at research facilities and they provide a valuable comparison since their estimated lifespan is between

that of humans and the monkey species commonly found at biomedical research facilities. Similarly, while

cotton-top tamarins (*Saguinus oedipus*) were at one time biomedical research models, they have not been

used for that purpose since 2008 when deforestation resulted in animals being listed as critically endangered.

Participating institutions

Data from eight United States National Primate Research Centers (NPRCs) are included: California (CNPRC),

Emory (ENPRC), New England (NEPRC; this center is no longer open but we obtained archival data), Oregon

(ONPRC), Southwest (SNPRC), Tulane (TNPRC), Washington (WaNPRC), and Wisconsin (WNPRC). Data

also originated from Primate Research Center IPB University in Indonesia, Keeling Center for Comparative

Medicine and Research at The University of Texas MD Anderson Cancer Center, National Institute on Aging

Intramural Research Program, Sam and Ann Barshop Institute for Longevity and Aging Studies at UT Health

San Antonio, Vervet Research Colony at Wake Forest University, and Yale University. **Table S1** shows

species sample sizes contributed by each institute. A data extraction standard operating protocol (SOP) was

developed to ensure consistency among institutions. The SOP requested data from all NHPs that were born

and died at the same institute going back through all historical records, along with sex, species, date of birth,

date of death, and disposition (i.e., death) code and description. We received data from 27 species categories

at the Duke Lemur Center, but ultimately did not include these data herein because they did not meet stage 1

filtering requirements of this study. We also note that life history profiles for these animals are published[44] and

the data are available for public download (https://lemur.duke.edu/duke-lemur-center-database/).

Data Filtering and Quality Control

Received data were first processed via a series of quality control checks for non-NHP species labels,

inconsistent or undefined codes, and duplicated records. We attempted to resolve inconsistencies or undefined

codes via follow-up with the original data source. Records that were unable to be resolved were removed from

subsequent analyses. The resulting data were then parsed through a two-stage filtering process. Stage One

filtering retained records with: 1) sex classified as male or female, 2) known date of birth (not estimated), and 3) survived at least 30 days (removing neonatal deaths). Species were then filtered to only include those which retained at least 150 animals. These Stage One filtered data yielded over 77,000 animals across 12 species. Stage Two filtering retained 1) animals that survived to adulthood using the National Institutes of Health Nonhuman Primate Evaluation and Analysis table of NHP life stages (**Table 1**).[45] The earliest age listed as adult for each species was used, supplemented by additional references for two species not present in the table, chimpanzees[46] and coppery titi monkeys.[33] Stage Two filtering also implemented a date of birth (DOB) cutoff. This step was critical for survival analyses and lifespan inference as received data did not include records on alive animals. Removing later (more recent) births avoided skewing results towards earlier deaths, and inference was thus based on the dataset of animals that had greatest opportunity to live to their maximum ages (**Figure S1**). The DOB threshold was implemented by retaining animals born before 2023 minus the number of years corresponding to the initial assessment of the 85th percentile of lifespan for that species (combined sexes; non-natural deaths as censored events). In total, this filtering stage yielded a dataset of 32,616 animals, across 12 species.

Defining censored events by death types. Given that these data did not include alive animals, for survival analyses, censored events were based on death type, as follows: 1) death types pertaining to research sacrifice and colony management were categorized as right censored events; 2) death types pertaining to natural causes or humane euthanasia for health reasons were coded as un-censored events. Right censoring is a statistical approach in survival analysis that enables inclusion of the knowledge that the subject survived at least to that point.[12] Treating deaths related to research sacrifice and colony management as right-censored events enabled animals to contribute to the survivorship model up until age of censoring. That is, this accounts for the lack of knowledge of how long the animal would have lived until a natural or health-related death. The final Stage Two filtered dataset was comprised of 12,269 events and 20,347 censored events.

Statistical analyses

We computed the Kaplan-Meier estimator[47] of the survivorship function for each species and sex, using the

ggsurvfit package[48] in R version 4.1.2. Survival curves and median lifespan estimates were calculated for both

including and excluding censored (research sacrifice; colony management death types) data. A critical analytic

consideration was that censoring was greatly biased by sex. Thus, the primary analyses presented with

comparisons by sex were limited to natural/health-related deaths only (no censored data). For many species,

proportional hazards assumptions were violated (preventing usage of the cox-proportional hazards model), but

since the primary analysis datasets were absent of censored events, analyses were not restricted to methods

for censored data. The analysis plan followed one that was applicable across all twelve species of various

sample sizes. For each species, maximum ages were compared between males and females using two

analytic approaches. First, quantile regression models were analyzed in SAS version 9.2 at the 25^{th}, 50^{th}, 75^{th},

and 85^{th} maximum age percentiles with sex as the predictor and primate center was included as a covariate.

For each species, we also tested for differences in the maximum age distributions by sex using the

nonparametric two-sample Kolmogorov-Smirnov test. [47] Finally, to evaluate the uniformity of the rate of decline

across survivorship curves, we fit an exponential model (e^{β}), separately, to the first and last quartiles of the

Kaplan-Meier survival curves using the nonlinear least squares function in R (version 4.1.2), shown in **Figure**

S3. As β captures the function's rate of decay, we illustrated trends across species, by sex, by plotting the

magnitude of β for these two quartiles. Computations were performed using the Wake Forest University (WFU)

High Performance Computing Facility.[49]

Data Availability

Raw, de-identified data are available via the password-protected database MIDAS (Monkey Inventory and

DAta management of Samples), request for access available from https://midas.wakehealth.edu/MIDAS. The

MIDAS database will include the same information provided in the manuscript and extended tables, and

provide tools for species comparisons, which will make this a user-friendly resource accessible to researchers.

Data sharing will be limited to scientific uses.

Acknowledgements

This work was supported by the National Institutes of Health: P40-OD010965 (MJJ), P51-OD011133 (CR; SNPRC), P51-OD011106 (RC; WNPRC), P51-OD011103 (EJV;NEPRC), P51-OD011104 (EJV,TNPRC), U42OD011123 (CEH; WaNPRC), P51OD010425 (CEH; WaNPRC), P51OD011092 (KC), OD011107 (KB), and the National Institute on Aging: U19AG057758 (LAC), U34AGAG068482 (AS), P30AG013319 (AS), P30AG044271 (AS), R01AG050797 (AS), AG-067419 (BH), NIH-NIA Intramural Research Program (JAM). Computations were performed using the Wake Forest University (WFU) High Performance Computing Facility, a centrally managed computational resource available to WFU researchers including faculty, staff, students, and collaborators.

References

1. Frye BM, Craft S, Latimer CS, et al. Aging-related Alzheimer's disease-like neuropathology and functional decline in captive vervet monkeys (Chlorocebus aethiops sabaeus). *Am J Primatol*. 2021;83(11):e23260.

2. Cox LA, Comuzzie AG, Havill LM, et al. Baboons as a model to study genetics and epigenetics of human disease. *ILAR J*. 2013;54(2):106-121.

3. Ross CN, Salmon AB. Aging research using the common marmoset: Focus on aging interventions. *Nutr Healthy Aging*. 2019;5(2):97-109.

4. Mattison JA, Vaughan KL. An overview of nonhuman primates in aging research. *Exp Gerontol*. 2017;94:41-45.

5. Saltzman W, Tardif SD, Rutherford JN. Chapter 13 - Hormones and Reproductive Cycles in Primates. In: Norris DO, Lopez KH, eds. *Hormones and Reproduction of Vertebrates*. Academic Press; 2011:291-327.

6. Zimmermann E, Radespiel U. Primate Life Histories. In: Henke W, Tattersall I, eds. *Handbook of Paleoanthropology*. Springer; 2015:1527-1592.

7. Tacutu R, Thornton D, Johnson E, et al. Human Ageing Genomic Resources: new and updated databases. *Nucleic Acids Res*. 2018;46(D1):D1083-D1090.

8. Hakeem AY, Sandoval GR, Jones M, Allman JM. Brain and life span in primates. In: *Handbook of the*
Psychology of Aging. Academic Press, Inc; 1996:78-104.

9. Weigl R. *Longevity of Mammals in Captivity; from the Living Collections of the World*. Kleine Senckenberg-
Reihe; 2005.

10. Bronikowski AM, Alberts SC, Altmann J, Packer C, Carey KD, Tatar M. The aging baboon: Comparative
demography in a non-human primate. *PNAS*. 2002;99(14):9591-9595.

11. Martin LJ, Mahaney MC, Bronikowski AM, Carey KD, Dyke B, Comuzzie AG. Lifespan in captive baboons
is heritable. *Mech Ageing Dev*. 2002;123(11):1461-1467.

12. Lagakos SW. General right censoring and its impact on the analysis of survival data. *Biometrics*.
1979;35(1):139-156.

13. Dick EJ, Owston MA, David JM, Sharp RM, Rouse S, Hubbard GB. Mortality in captive baboons (Papio
spp.): a-23-year study. *J Med Primatol*. 2014;43(3):169-196.

14. Tardif SD, Mansfield KG, Ratnam R, Ross CN, Ziegler TE. The marmoset as a model of aging and age-
related diseases. *ILAR J*. 2011;52(1):54-65.

15. Schaaf GW, Justice JN, Quillen EE, Cline JM. Resilience, aging, and response to radiation exposure
(RARRE) in nonhuman primates: a resource review. *GeroScience*. 2023;45(6):3371-3379.

16. Zedda N, Bramanti B, Gualdi-Russo E, Ceraico E, Rinaldo N. The biological index of frailty: A new index
for the assessment of frailty in human skeletal remains. *American Journal of Physical Anthropology*.
2021;176(3):459-473.

17. Searle SD, Mitnitski A, Gahbauer EA, Gill TM, Rockwood K. A standard procedure for creating a frailty
index. *BMC Geriatr*. 2008;8:24.

18. Zeng A, Song X, Dong J, et al. Mortality in relation to frailty in patients admitted to a specialized geriatric
intensive care unit. *J Gerontol A Biol Sci Med Sci*. 2015;70(12):1586-1594.

19. Shi SM, Olivieri-Mui B, McCarthy EP, Kim DH. Changes in a Frailty Index and Association with Mortality. *J*

Am Geriatr Soc. 2021;69(4):1057-1062.

20. Rockwood K, Howlett SE. Age-related deficit accumulation and the diseases of ageing. *Mech Ageing Dev*.

2019;180:107-116.

21. Rockwood K, Mogilner A, Mitnitski A. Changes with age in the distribution of a frailty index. *Mechanisms of*

Ageing and Development. 2004;125(7):517-519.

22. Rashmi R, Mohanty SK. Examining chronic disease onset across varying age groups of Indian adults using

competing risk analysis. *Sci Rep*. 2023;13:5848.

23. Colchero F, Aburto JM, Archie EA, et al. The long lives of primates and the "invariant rate of ageing"

hypothesis. *Nat Commun*. 2021;12(1):3666.

24. Mattison JA, Roth GS, Beasley TM, et al. Impact of caloric restriction on health and survival in rhesus

monkeys from the NIA study. *Nature*. 2012;489(7415):318-321.

25. Mattison JA, Colman RJ, Beasley TM, et al. Caloric restriction improves health and survival of rhesus

monkeys. *Nature Communications*. 2017;8:14063.

26. Kirkwood TB, Holliday R. The evolution of ageing and longevity. *Proc R Soc Lond B Biol Sci*.

1979;205(1161):531-546.

27. Maklakov AA, Chapman T. Evolution of ageing as a tangle of trade-offs: energy versus function. *Proc Biol*

Sci. 2019;286(1911):20191604.

28. Cohen AA, Coste CFD, Li XY, Bourg S, Pavard S. Are trade-offs really the key drivers of ageing and life

span? *Functional Ecology*. 2020;34(1):153-166.

29. Cawthon Lang K. Primate Factsheets: Common marmoset (Callithrix jacchus). Primate Info Net, Wisconsin

National Primate Research Center.

30. Cawthon Lang K. Primate Factsheets: Cotton-top tamarin (Saguinus oedipus). Primate Info Net, Wisconsin
National Primate Research Center.

31. Bronikowski AM, Altmann J, Brockman DK, et al. Aging in the natural world: comparative data reveal
similar mortality patterns across primates. *Science*. 2011;331(6022):1325-1328.

32. Arnold C, Matthews LJ, Nunn CL. The 10kTrees website: A new online resource for primate phylogeny.
Evol Anthropol. 2010;19(3):114-118.

33. Zablocki-Thomas P, Rebout N, Karaskiewicz CL, Bales KL. Survival rates and mortality risks of
Plecturocebus cupreus at the California National Primate Research Center. *Am J Primatol*. Published
online July 9, 2023:e23531.

34. Chiou KL, Montague MJ, Goldman EA, et al. Rhesus macaques as a tractable physiological model of
human ageing. *Philos Trans R Soc Lond B Biol Sci*. 2020;375(1811):20190612.

35. Roth GS, Mattison JA, Ottinger MA, Chachich ME, Lane MA, Ingram DK. Aging in rhesus monkeys:
relevance to human health interventions. *Science*. 2004;305(5689):1423-1426.

36. Colman RJ, Beasley TM, Kemnitz JW, Johnson SC, Weindruch R, Anderson RM. Caloric restriction
reduces age-related and all-cause mortality in rhesus monkeys. *Nat Commun*. 2014;5:3557.

37. Ward JM, Buslov AM, Vallender EJ. Twinning and survivorship of captive common marmosets (Callithrix
jacchus) and cotton-top tamarins (Saguinus oedipus). *J Am Assoc Lab Anim Sci*. 2014;53(1):7-11.

38. Arbogast DM, Crews DE, McGraw WS, Ely JJ. Demography and epidemiology of captive former
biomedical research chimpanzees (Pan troglodytes) 1: Survival and mortality. *American Journal of
Primatology*. 2023;85(4):e23466.

39. Sert NP du, Hurst V, Ahluwalia A, et al. The ARRIVE guidelines 2.0: Updated guidelines for reporting
animal research. *PLOS Biology*. 2020;18(7):e3000410.

40. Riddle NC, Biga PR, Bronikowski AM, et al. Comparative analysis of animal lifespan. *GeroScience.*
2024;46(1):171-181.

41. Keller C, Roos C, Groeneveld LF, Fischer J, Zinner D. Introgressive hybridization in southern African
baboons shapes patterns of mtDNA variation. *Am J Phys Anthropol.* 2010;142(1):125-136.

42. Robinson JA, Belsare S, Birnbaum S, et al. Analysis of 100 high-coverage genomes from a pedigreed
captive baboon colony. *Genome Res.* 2019;29(5):848-856.

43. Kanthaswamy S, Trask JS, Ross CT, et al. A large-scale SNP-based genomic admixture analysis of the
captive rhesus macaque colony at the California National Primate Research Center. *Am J Primatol.*
2012;74(8):747-757.

44. Zehr SM, Roach RG, Haring D, Taylor J, Cameron FH, Yoder AD. Life history profiles for 27 strepsirrhine
primate taxa generated using captive data from the Duke Lemur Center. *Sci Data.* 2014;1(1):140019.

45. Feister AJ, DiPietrantonio A, Yuenger J, Ireland K, Rao A. *Nonhuman Primate Evaluation and Analysis*
Part 1: Analysis of Future Demand and Supply | Office of Research Infrastructure Programs (ORIP) –
DPCPSI – NIH.; 2018.

46. Stumpf R. Chimpanzees and Bonobos: Diversity within and between species. In: Campbell CJ, Fuentes A,
MacKinnon KC, Panger M, Bearder SK, eds. *Primates in Perspective.* Oxford University Press; 2007:321-
344.

47. Hosmer DW, Lemeshow S, May S. *Applied Survival Analysis: Regression Modeling of Time-to-Event Data.*
John Wiley & Sons, Ltd; 2008.

48. Sjoberg D, Baillie M, Fruechtenicht C, Haesendonckx S, Treis T. pharmaverse/ggsurvfit. Published online
February 29, 2024.

49. WFU High Performance Computing Facility. Information Systems and Wake Forest University.

Comparative lifespan and healthspan of nonhuman primate species common to biomedical research.

Primary Display Items and Extended Data Figures

Table of Contents

Primary Figures

Figure 1. Scatter plot of data points for natural and health-related euthanasia deaths by species.

Boxplot overlay depicts median and interquartile range by species and sex. Proportion of data by sex and species also shown. The vertical dashed line denotes equal counts of males and females by species.

Figure 2. Combined survival curves for females (A) and males (B) of all 12 species.

Data shown are for animals with deaths resulting from natural causes or humane euthanasia for health-related reasons.

Figure 3. Comparison of rate of survivorship decline by quartile and sex.

Rates of decline were calculated from fitting an exponential model to the first and last quartiles of the sex-specific Kaplan-Meier survival curves. Males and females are compared by quartile. Rate of decline was generally faster in males within the first quartile with the pattern nearly reversed by sex in the last quartile.

Figure 4. Kaplan-Meier survival curves by sex and species for natural deaths or humane euthanasia for health-related reasons.

For each plot, the X-axis scaling (maximum age) is species-specific.

Primary Tables

Table 1. Sample sizes of primary analysis datasets and species-specific age categories.
For each species, age categories and estimated age ranges are shown.[33,44,45]

Common Name	Species name	Post-filtering sample size*		Age categories			
		Male	Female	Infant	Juvenile	Adult	Geriatric
Baboon	*Papio hamadryas* spp.	334	669	<12 months	1-4 years	4-15 years	>15 years
Bonnet macaque	*Macaca radiata*	19	43	<12 months	1-4 years	4-15 years	>15 years
Chimpanzee	*Pan troglodytes* spp.	48	50	<12 months	1-10 years	10-35 years	>35 years
Common marmoset	*Callithrix jacchus*	378	453	<6 months	6-18 months	1.5-8 years	>8 years
Coppery titi monkey	*Plecturocebus cupreus*	32	33	<12 months	1-4 years	4-10 years	>10 years
Cotton-top tamarin	*Saguinus oedipus*	155	191	<7 months	7-30 months	2.5-10 years	>10 years
Cynomolgus macaque	*Macaca fascicularis*	82	132	<12 months	1-4 years	4-17 years	>17 years
Japanese macaque	*Macaca fuscata*	174	196	<12 months	1-4 years	4-15 years	>15 years
Pig-tailed macaque	*Macaca nemestrina*	173	596	<12 months	1-4 years	4-15 years	>15 years
Rhesus macaque	*Macaca mulatta*	2465	5742	<12 months	1-4 years	4-17 years	>17 years
Squirrel monkey	*Saimiri* spp.	53	47	<12 months	1-4 years	4-15 years	>15 years
Vervet/African green	*Chlorocebus aethiops sabaeus*	60	144	<12 months	1-4 years	4-15 years	>15 years

Natural or Health-related deaths only

Table 2. Maximum and median age at death by sex and species

Common Name	Species name	Maximum observed age in years*		Median age at death in years (range)*	
		Male	Female	Male	Female
Baboon	*P. hamadryas* spp.	30.3	30.6	11.29(10.41-12.47)	11.65(11.08-12.44)
Bonnet macaque	*M. radiata*	32.8	21.4	7.93(5.70-14.54)	9.22(7.81-13.49)
Chimpanzee	*P. troglodytes* spp.	53.3	58.8	33.00(28.41-38.33)	43.96(41.66-45.82)
Common marmoset	*C. jacchus*	17.3	17.1	5.97(5.41-6.74)	5.31(4.92-5.66)
Coppery titi monkey	*P. cupreus*	24.4	23.2	8.59(6.92-12.13)	9.16(7.35-14.13)
Cotton-top tamarin	*S. oedipus*	24.7	23.1	9.60(7.87-11.27)	8.87(7.67-10.57)
Cynomolgus macaque	*M. fascicularis*	28.4	23.5	6.93(6.21-8.18)	8.62(7.72-9.84)
Japanese macaque	*M. fuscata*	38.4	30.1	8.19(7.48-9.36)	11.41(10.27-12.70)
Pig-tailed macaque	*M. nemestrina*	27.9	29.2	8.43(7.49-9.12)	8.96(8.43-9.59)
Rhesus macaque	*M. mulatta*	44.2	42	7.89(7.65-8.24)	10.26(10.03-10.49)
Squirrel monkey	*Saimiri* spp.	22.7	21.8	8.78(6.97-10.09)	9.22(6.55-11.19
Vervet/African green	*C. aethiops sabaeus*	24.1	30.6	8.34(7.57-10.71)	17.87(15.24-20.23)

Median age at death is calculated from natural and clinical deaths only; maximum observed age includes animals with any type of death. Maximum ages are from the current dataset only; there are known older animals of some of these species at research institutes, such as a 29-year-old titi monkey male at CNPRC and two 19-year-old male marmosets at SNPRC.

Table 3. Sex-based comparisons of age by species.

Quantile regression for 25th, 50th, 75th, and 85th percentiles. Regression models adjusted for primate location (data source). Distribution of ages by sex were assessed using the Kolmogorov Smirnov test. Complete data used for analyses (natural or clinical deaths) with no censoring.

Species	Max Age Percentile	Years of Age (Male)	Years of Age (Female)	Quantile Regression Estimate	Standard Error	Quantile Regression P-value	Kolmogorov-Smirnov P-Value
Baboon	25th	7.42 (6.68-7.92)	7.57 (7.09-8.10)	0.68	0.38	0.073	
(N=334 M, 669 F)	50th	11.29 (10.41-12.47)	11.65 (11.08-12.44)	0.96	0.65	0.141	0.352
	75th	15.68 (14.80-16.53)	16.40 (15.86-17.12)	0.94	0.57	0.097	
	85th	17.47 (16.97-18.79)	18.28 (17.70-19.14)	0.62	0.47	0.185	
Bonnet macaque	25th	5.42 (4.23-7.93)	5.72 (5.03-8.40)	0.30	1.16	0.798	
(N=19 M, 43 F)	50th	7.93 (5.70-14.54)	9.22 (7.81-13.49)	1.29	2.38	0.591	0.794
	75th	14.54 (7.93-19.23)	15.32 (11.57-17.76)	0.78	2.76	0.778	
	85th	16.61 (12.34-21.32)	17.10 (15.32-21.40)	0.50	3.05	0.871	
Chimpanzee	25th	24.30 (17.71-28.41)	37.47 (29.23-41.66)	14.32	4.42	1.65×10^{-3}	
(N=48 M, 50 F)	50th	33.00 (28.41-38.33)	43.96 (41.66-45.82)	10.59	2.61	1.03×10^{-4}	1.78×10^{-5}
	75th	39.84 (37.52-47.14)	48.84 (45.67-51.77)	7.77	3.07	0.013	
	85th	44.96 (39.47-48.67)	51.57 (48.84-54.32)	5.15	2.58	0.049	
Common marmoset	25th	3.56 (3.08-4.00)	3.42 (3.08-3.67)	-0.29	0.26	0.274	
(N=378 M, 453 F)	50th	5.97 (5.41-6.74)	5.31 (4.92-5.66)	-0.59	0.29	0.040	0.002
	75th	8.71 (8.35-9.19)	7.98 (7.19-8.71)	-0.68	0.35	0.048	
	85th	10.00 (9.24-10.47)	9.56 (9.11-10.41)	-0.38	0.37	0.303	
Coppery titi monkey	25th	5.90 (5.18-7.27)	7.04 (4.28-7.81)	1.05	1.43	0.469	
(N=32 N, 33 F)	50th	8.59 (6.92-12.13)	9.16 (7.35-14.13)	1.04	2.42	0.669	0.322
	75th	12.32 (9.87-17.77)	16.19 (13.11-18.80)	3.88	2.38	0.108	
	85th	16.72 (12.31-24.43)	18.43 (15.74-23.23)	1.72	2.47	0.490	
Cotton-top tamarin	25th	5.69 (4.80-6.52)	5.30 (4.63-6.19)	-0.39	0.55	0.477	
(N=155 M, 191 F)	50th	9.60 (7.87-11.27)	8.87 (7.67-10.57)	-0.73	0.96	0.446	0.874
	75th	14.70 (13.35-16.13)	14.17 (12.64-15.28)	-0.53	0.94	0.574	
	85th	16.74 (15.85-17.68)	16.21 (14.71-17.14)	-0.53	0.74	0.480	
Cynomolgus macaque	25th	4.89 (4.47-5.61)	5.91 (5.23-6.60)	0.93	0.53	0.082	
	50th	6.93 (6.21-8.18)	8.62 (7.72-9.84)	1.58	0.72	0.028	0.034
(N=82 M, 132 F)	75th	10.43 (8.45-15.73)	12.24 (11.01-13.61)	1.97	1.41	0.165	
	85th	15.73 (12.99-24.63)	13.94 (12.75-15.37)	-2.03	1.86	0.278	
Japanese macaque	25th	6.23 (5.62-6.58)	7.33 (6.75-8.56)	1.08	0.46	0.021	
(N=174 M, 196 F)	50th	8.19 (7.48-9.36)	11.41 (10.27-12.70)	3.26	0.81	6.88×10^{-5}	4.66×10^{-5}
	75th	13.41 (12.00-15.13)	16.81 (15.40-18.86)	3.23	1.23	0.009	
	85th	16.33 (14.68-18.34)	19.44 (18.48-21.93)	3.10	0.97	0.002	
Pigtail macaque	25th	5.39 (5.14-5.94)	6.27 (5.75-6.73)	0.90	0.36	0.013	
(N=173 M, 596 F)	50th	8.43 (7.49-9.12)	8.96 (8.43-9.59)	0.63	0.55	0.254	0.134
	75th	12.80 (11.08-14.63)	12.30 (11.70-12.90)	-0.46	0.69	0.510	
	85th	15.56 (13.77-17.48)	14.17 (13.65-14.90)	-1.19	0.70	0.091	
Rhesus macaque	25th	5.55 (5.45-5.66)	6.85 (6.66-7.01)	1.22	0.10	4.21×10^{-37}	
(N=2465 M, 5742 F)	50th	7.89 (7.65-8.24)	10.26 (10.03-10.49)	1.89	0.14	1.27×10^{-40}	2.20×10^{-16}
	75th	13.98 (13.33-14.74)	14.70 (14.41-14.88)	0.90	0.20	7.02×10^{-6}	
	85th	17.73 (17.14-18.41)	16.97 (16.72-17.29)	0.24	0.26	0.355	
Squirrel monkey	25th	5.72 (4.78-6.97)	5.40 (4.95-6.67)	-0.05	0.58	0.934	
(N=53 M, 47 F)	50th	8.78 (6.97-10.09)	9.22 (6.55-11.19	0.84	1.00	0.401	0.585
	75th	12.76 (10.07-15.46	13.39 (10.65-14.91	0.79	1.61	0.625	
	85th	15.25 (12.76-21.18)	13.84 (13.39-18.43)	-0.34	1.91	0.859	
Vervet/African green monkey	25th	6.86 (5.80-7.44)	10.57 (9.54-12.21)	3.4	0.88	1.49×10^{-4}	
	50th	8.34 (7.57-10.71)	17.87 (15.24-20.23)	8.93	1.49	8.98×10^{-9}	7.92×10^{-10}
(N=60 M, 144 F)	75th	12.93 (10.71-14.51)	23.12 (21.99-24.60)	10.26	1.16	4.51×10^{-16}	
	85th	13.88 (13.00-16.70)	24.81 (24.25-26.34)	10.98	1.06	2.45×10^{-20}	

Extended Data Figures (Active Links within Article)

Extended Data Figure 1: Distribution and proportions of sex and censored data points for filtered data (n=32,616).

Deaths due to natural causes or health-related reasons are labeled as 'natural.' Deaths due to research sacrifice or colony management are labeled as 'censored.' Multiple species had very high proportions of censored events.

Extended Data Figure 2: Kaplan-Meier survival curves for 12 species, by sex, including censored events.

Graphs of Kaplan-Meier estimates of the survivorship function for all data passing quality control filtering are provided. Presented are survival curves where deaths from research sacrifice and colony management are included as censored events; censoring was biased by sex, so sex-specific comparisons were not computed for these data and were instead limited to data containing natural deaths, only. For many species (e.g., baboons), the high proportion of censored events yielded unstable survival probability estimates for latter ages. For each plot, the X-axis scaling (maximum age) is species-specific.

Extended Data Figure 3: Phylogenetic tree of 12 species analyzed in study.

This tree was generated with the 10kTrees Project and modified to match taxonomic names with those used in our study and to simplify the presentation.[32] Only the 12 species studied herein are represented in the tree; there are many other species of primates in these clades not pictured.