Title: Motor sequence learning-induced corticospinal plasticity is biased towards

sensorimotor mu rhythm peak phases

Abbreviated title: Learning-induced mu phase-dependent corticospinal plasticity

Authors: Tharan Suresh[1], Fumiaki Iwane[2,3], Minsu Zhang[3], Margaret McElmurry[1], Muskan

Manesiya[1], Michael V Freedberg[1], Sara J Hussain[1]*

Affiliations:

[1]Department of Kinesiology & Health Education, The University of Texas at Austin. Austin,

TX, USA. [2]Human Cortical Physiology and Neurorehabilitation Section, National Institute of

Neurological Disorders and Stroke, National Institutes of Health. Bethesda, MD, USA.

[3]Department of Electrical & Computer Engineering, The University of Texas at Austin.

Austin, TX, USA.

*Corresponding author: sara.hussain@austin.utexas.edu

Number of pages: 40

Number of figures: 6

Number of words for Abstract: 243

Number of words for Introduction: 757

Number of words for Discussion: 2131

Conflict of interest: The authors declare no competing financial interests.

Acknowledgements: SJH was supported by K12HD093427 and R21NS133605 and TS

was supported by R25HD105583. The authors thank Kristen Pulliam for assistance with data

collection.

Abstract

Motor cortical (M1) transcranial magnetic stimulation (TMS) increases corticospinal output
and improves motor learning when delivered during sensorimotor mu rhythm trough but not
peak phases, suggesting that mechanisms supporting motor learning may be most active
during mu trough phases. If so, learning-related corticospinal plasticity should be most
evident during mu trough phases. Healthy adults were assigned to either a sequence or
control group. Participants in the sequence group practiced the implicit serial reaction time
task (SRTT), which contained an embedded, repeating 12-item sequence. Participants in the
control group practiced a version of the SRTT that contained no sequence. We measured
mu phase-independent and phase-dependent MEP amplitudes using EEG-informed single-
pulse TMS before, immediately, and 30 minutes after the SRTT in both groups. All
participants performed a retention test one hour after SRTT acquisition. In both groups, mu
phase-independent MEP amplitudes increased following SRTT acquisition, but the pattern of
mu phase-dependent MEP amplitude increases after SRTT acquisition differed between
groups. MEP amplitude changes from baseline to 30 minutes after SRTT acquisition more
strongly differed across phases in the control relative to the sequence group, with the control
group showing smaller increases in peak- than trough-specific MEPs. Contrary to our
original hypothesis, results revealed that sequence learning recruits peak- rather than
trough-specific neurophysiological mechanisms. Overall, these findings suggest that mu
peak phases may provide protected time windows for motor memory consolidation and
demonstrate the presence of a mu phase-dependent motor learning mechanism in the
human brain.

Keywords: sensorimotor rhythms, motor learning, brain stimulation, phase, motor cortex

Significance statement

Recent work suggests that the neurophysiological mechanisms supporting motor learning may be most active during sensorimotor mu rhythm trough phases. Here, we evaluated this possibility by measuring mu phase-dependent corticospinal plasticity induced by motor sequence learning. Results provide first evidence that motor sequence learning produced corticospinal plasticity that was more pronounced during mu peak than trough phases, demonstrating the presence of a phase-dependent learning mechanism within the human motor system.

Introduction

The sensorimotor cortex is a hub within brain networks controlling human motor skill acquisition. Human sensorimotor cortical activity dynamically oscillates in the mu (8-13 Hz) frequency range, with distinct phases of this rhythm reflecting brief windows of relative excitation and inhibition (Bergmann et al., 2019; Hussain et al., 2019; Suresh & Hussain, 2023; Zrenner et al., 2018). In non-human primates, motor cortical (M1) neuronal spiking rates are highest during mu rhythm trough phases and lowest during opposing peak phases (Haegens et al., 2011). In humans, corticospinal output (Bergmann et al., 2019; Hussain et al., 2019; Suresh & Hussain, 2023; Wischnewski et al., 2022; Zrenner et al., 2018) and interhemispheric communication between homologous M1 subregions (Stefanou et al., 2018) are also increased during sensorimotor mu rhythm trough relative to peak phases.

In addition to shaping single-neuron spiking rates, corticospinal output, and inter-hemispheric communication, sensorimotor mu rhythm phase also determines the likelihood of inducing long-term potentiation-like (LTP-like) corticospinal plasticity in humans. For example, transcranial magnetic stimulation (TMS) interventions applied to M1 during mu rhythm trough phases preferentially induce LTP-like corticospinal plasticity, measured as increases in corticospinal output (Baur et al., 2020; Zrenner et al., 2018). In contrast, identical TMS interventions applied during mu rhythm peak phases induce weak long-term depression-like (LTD-like) corticospinal plasticity, measured as decreases in corticospinal output (Baur et al., 2020; Zrenner et al., 2018). These findings suggest that corticospinal plasticity induction is causally related to sensorimotor mu rhythm phase, with LTP-like and weak LTD-like corticospinal plasticity occurring preferentially during mu rhythm trough and peak phases, respectively. Given that LTP-like plasticity processes have been proposed to support human motor learning (Cantarero et al., 2013a; Cantarero et al., 2013b), and TMS interventions preferentially enhance motor learning when applied during mu rhythm trough phases (Hussain et al., 2021), the neurophysiological mechanisms underlying human motor learning may thus be most active during sensorimotor mu rhythm trough phases.

Outside of the motor system, brain oscillatory activity is thought to rhythmically control

various attentional (Busch & VanRullen, 2010; VanRullen et al., 2011), perceptual

(Baumgarten et al., 2015; Busch et al., 2009; Dugué et al., 2011), and memory processes

(Kerrén et al., 2018; Siegel et al., 2009). For example, the formation and retrieval of

declarative memories has been proposed to occur during opposing hippocampal oscillatory

phases (Hasselmo et al., 2002; Kerrén et al., 2018), object memory formation preferentially

occurs during distinct prefrontal oscillatory phases (Siegel et al., 2009), and replay of whole-

brain neuronal activity patterns associated with working memory are phase-locked to the

theta rhythm (Fuentemilla et al., 2010). These findings suggest that oscillatory phase-

dependent memory processing may be a widespread learning mechanism present across

neural domains (Watrous et al., 2015). If so, phase-dependent neurophysiological signatures

of motor learning should also be detectable within the human motor system.

Here, we reasoned that, if motor learning relies on a sensorimotor mu phase-dependent

mechanism, then learning-related corticospinal plasticity should be more evident during

some mu rhythm phases than others. That is, expression of learning-related corticospinal

plasticity should vary significantly across mu phases. Based on previous studies showing

that corticospinal output (Hussain et al., 2019; Suresh & Hussain, 2023; Zrenner et al.,

2018), interhemispheric communication (Stefanou et al., 2018), corticospinal plasticity (Baur

et al., 2020; Zrenner et al., 2018), and the effects of TMS interventions on motor skill

learning (Hussain et al., 2021) are all enhanced during mu rhythm trough relative to peak

phases, we predicted that learning-related changes in corticospinal output would also be

more evident during mu rhythm trough than peak phases.

To test this possibility, we measured sensorimotor mu phase-independent and phase-

dependent corticospinal output using single-pulse, real-time EEG-informed TMS immediately

before and after healthy adults practiced a serial reaction time task (SRTT) that included

either an embedded, repeating 12-item motor sequence (sequence group) or a control

version of the same task that included no sequence (control group). As expected, our results

show that participants in the sequence group exhibited greater sequence-specific learning

than those the control group. In both groups, mu phase-independent corticospinal output

increased after SRTT acquisition, but these increases did not differ between groups. In

contrast, corticospinal plasticity induced by SRTT acquisition more strongly varied across mu

rhythm phases for the control group than sequence group, such that the control group

showed significantly smaller increases in peak- than trough-specific corticospinal output.

Surprisingly, these findings indicate that corticospinal plasticity induced by motor sequence

learning is more prominent during mu rhythm peak than trough phases, highlighting the

mechanistic role of mu rhythm phase in shaping behaviorally-relevant corticospinal plasticity

within the human motor system.

Methods

Participants

Thirty-six naïve right-handed healthy adult participants (age = 20.9 ± 0.52, 25 F and 11 M)

participated in a single-session study that involved single-pulse transcranial magnetic

stimulation (TMS), electroencephalography (EEG), electromyography (EMG), and behavioral

testing. Right-handedness was confirmed using the Edinburgh Handedness Inventory

(Oldfield, 1971). This study was approved by the Institutional Review Board at the University

of Texas at Austin and all participants provided their written informed consent. Participants

were randomly assigned to either the sequence group (N = 18, age = 20 ± 0.37 years, 13 F

and 5 M) or control group (N=18, age = 21.8 ± 0.94 years, 12 F and 6 M) prior to beginning

experimental procedures. This sample size was chosen based on previous studies that

identified significant mu phase-dependent variation in corticospinal output and corticospinal

plasticity, as well as phase-dependent enhancement of motor learning (N = 12-23, Zrenner

et al., 2018; Bergmann et al., 2019; Hussain et al., 2019; Suresh & Hussain, 2023; Hussain

et al., 2021).

Experimental design and timeline

After EEG and EMG preparation, the scalp hotspot for the right first dorsal interosseous

muscle (FDI) and the stimulation intensity needed to elicit motor-evoked potentials (MEPs)

within this muscle were empirically determined for each subject. Participants then performed

a choice reaction time task (cRT), after which parameters for real-time targeting of

sensorimotor mu rhythm phases were individually optimized. Phase targeting accuracy was

assessed during EEG recordings while participants rested quietly with their eyes open.

Then, single-pulse TMS was delivered over the scalp hotspot during sensorimotor mu

rhythm peak, trough, and random phases. Participants subsequently practiced one of two

versions of the serial reaction time task (SRTT; see below for details). Immediately and 30

minutes after task completion, single-pulse TMS was again delivered during mu peak, trough

and random phases. One hour after completing the SRTT, both groups performed a brief

retention test that involved the same SRTT version performed during practice. After retention

testing, all participants completed the Process Dissociation Procedure (PDP; Wilkinson &

Shanks, 2004). The PDP was used to evaluate the development of explicit sequence

awareness, given that corticospinal plasticity following motor sequence learning under

implicit and explicit task conditions may differ (Tunovic et al., 2014). Data from the cRT are

not reported in this manuscript. See Figure 1 for experimental design and timeline.

Figure 1. Experimental timeline and serial reaction time task. A) Experimental design
and timeline. Real-time EEG analysis algorithm parameters were individually optimized
during resting EEG recordings. After acquisition of phase-independent (random) and phase-
dependent (peak and trough) MEPs at baseline, subjects in both groups performed the
SRTT. Post-SRTT phase-independent and phase-dependent MEPs were recorded
immediately and 30 minutes after SRTT acquisition. SRTT retention was evaluated 1 hour
after SRTT acquisition. **B)** Depiction of implicit serial reaction time task (SRTT). During
acquisition, participants practiced 14 blocks (120 trials per block) of the implicit SRTT.
During retention, participants practiced 7 blocks (120 trials per block). The sequence group
practiced the SRTT with an embedded 12-item sequence, while the control group practiced
the task with no sequence (all random blocks).

Data acquisition

EEG and EMG recording. 64-channel EEG and bipolar EMG signals were recorded using

TMS-compatible amplifiers (NeurOne Tesla with real-time digital out option, Bittium

Biosignals, Finland). EEG impedances were maintained below 10 kΩ. EMG signals were

recorded from the right FDI using disposable adhesive electrodes placed in a belly-tendon

montage. EEG and EMG were recorded at a sampling rate of 5 kHz, with low-pass hardware

filtering at 1250 Hz and a resolution of 0.001 µV.

Transcranial magnetic stimulation. We empirically determined the motor hotspot as the scalp

site at which suprathreshold, single-pulse TMS elicited a reliable, focal twitch within the right

FDI as well as a peak-to-peak motor evoked potential (MEP) amplitude of at least 50 µV.

After hotspot identification, resting motor threshold (RMT) was determined using an

automatic threshold-tracking algorithm (Borckardt et al., 2006). On average, RMT was 64.3 ±

3.8% of maximum stimulator output for the sequence group and 61.2 ± 3.1% for the control

group. All TMS procedures were performed using a figure-of-eight coil held at ~45° relative

to the midsagittal line (DuoMAG XT-100, Deymed Diagnostic, biphasic pulse shape). Coil

position accuracy was monitored online using frameless neuronavigation (BrainSight, Rogue

Research) during hotspot identification, RMT determination, and mu phase-independent and

mu phase-dependent single-pulse TMS.

Real-time EEG analysis. Real-time EEG-triggered TMS was performed using an established

method (Zrenner et al., 2018; Shirinpour et al., 2020; Hussain et al., 2021; Suresh &

Hussain, 2023). The EEG system was configured to stream data from 5 pre-selected EEG

channels overlying the left sensorimotor cortex (C3, FC3, FC5, CP3 and CP5) using a local

wired network connection (packet rate = 1 kHz, 5 packets per sample) between the EEG

system and a dedicated PC for signal processing (Microsoft Windows 10, 10 cores, Intel i9

processor, 16 GB RAM) using Lab Streaming Layer software (Kothe, 2014) in real-time.

Custom MATLAB scripts were used to receive EEG data, process it, and send TTL triggers

to the TMS device upon detection of the target mu rhythm phase. During data streaming,

EEG data was first downsampled to 1 kHz. To attenuate the effects of volume conduction,

the Hjorth transformation was applied by subtracting the mean of the surrounding electrodes

(FC3, FC5, CP3 and CP5) from the center electrode (C3; Hjorth 1975). The resulting signal

was then buffered to create overlapping 500 ms windows for real-time analysis. At each time

step, the current data window was zero-phase bandpass filtered using a Finite Impulse

Response (FIR) filter (8-13 Hz, order = 128) and 64 ms data segments at the beginning and

end of the 500 ms window were removed to eliminate edge effects. The remaining 372 ms

data window was then used to fit an autoregressive model (Yule-Walker method, order: 30)

that forward predicted the bandpass filtered signal 128 ms into the future. After forward

prediction, the Hilbert transform was used to generate an estimated phase angle timeseries

signal. The phase estimation algorithm was also configured to trigger TMS only when each

participant's sensorimotor mu power exceeded a predefined threshold. To calculate this

threshold, resting state EEG data was divided into non-overlapping 500 ms segments,

bandpass filtered using a Finite Impulse Response (FIR) filter (8-13 Hz, order = 128), Hjorth

transformed, downsampled to 1 kHz and then spectrally decomposed (Welch's method, 4-

100 Hz with 0.25 Hz resolution excluding frequencies between 57.75 – 62 Hz to reduce the

influence of line noise). For each participant, the median of the resulting frequency-domain

signal between 8 and13 Hz was defined as the power threshold. When the estimated

instantaneous phase angle at the time point of interest matched the pre-determined mu

phase condition and mu power exceeded each participant's threshold, a single TTL pulse

was sent to the TMS system and the EEG system to trigger single-pulse TMS and mark the

time of TMS delivery, respectively.

Phase targeting accuracy and real-time EEG-informed single-pulse TMS. Prior to delivering

real-time EEG-informed TMS, we first evaluated the accuracy of phase targeting in the

absence of TMS delivery in each participant. 25 TTL pulses were sent to the EEG system at

mu peak, trough and random phases, but no TMS was delivered (minimum inter-stimulus

interval = 5 s + random jitter). During mu phase-independent single-pulse TMS, blocks of 25

TMS pulses were delivered to the scalp hotspot for the right FDI during random mu phases

(120% RMT, inter-stimulus interval = 5 s + random jitter). During mu phase-dependent

single-pulse TMS, blocks of 25 TMS pulses were delivered during mu peak and trough

phases (120% RMT, minimum inter-stimulus interval = 5 s + random jitter). Real-time EEG-

informed single-pulse TMS was delivered at baseline, immediately after SRTT acquisition,

and 30 minutes after SRTT acquisition. The order of phases targeted per block (i.e., random,

peak, and trough) was counterbalanced across participants in both groups.

<u>Serial Reaction Time Task (SRTT)</u>. Participants in both the sequence and control groups

were seated in front of a computer monitor with their right arm resting on a tabletop and their

right hand positioned atop a serial response box (Chronos Response Device, Psychology

Software Tools, see Figure 1). Participants performed one of two modified versions

(sequence or control) of the SRTT task (Nissen & Bullemer, 1987; Perez et al., 2007)

programmed using E-prime 3 (Psychology Software Tools). The control group was included

in this study to experimentally account for non-specific effects of task engagement on

corticospinal output. During the SRTT, the monitor displayed four equally-spaced green

dashes against a black background. On each trial, one of these dashes was replaced by a

circle. Prior to beginning the task, participants were instructed to respond to the presentation

of the circle as fast and accurately as possible by pressing one of the four keys with the

corresponding finger of their right hand, as shown in Figure 1. When the participant pressed

the correct key, the circle disappeared, and the trial was terminated. If an incorrect key was

pressed, the circle remained on the monitor and the trial continued until the correct key was

pressed. 500 ms elapsed between trials. No feedback on keypress accuracy or keypress

latency was provided.

Participants became familiarized with the task during a practice block that included 10 trials.

During SRTT acquisition, participants in the sequence group performed 14 blocks of

keypresses, with one minute of rest between blocks. Each block was composed of 10

repetitions of a 12-item sequence, amounting to a total of 120 keypresses per block. Blocks

3-7 and 9-13 included a pre-determined, embedded 12-item sequence (2-3-1-4-3-2-4-1-3-4-

2-1). Blocks 1, 2, 8 and 14 were random blocks that preceded and followed the sequence

blocks and contained no embedded sequence. During SRTT retention, participants in the

sequence group performed 7 blocks of 120 keypresses each, again resting for one minute

between blocks. Blocks 1 and 7 were random blocks while blocks 2-6 were sequence

blocks.

Random blocks contained no consecutive item repeats and each item had approximately the

same frequency of appearance. Participants in the control group experienced keypresses

that were randomized. Participants were not informed about the presence or absence of the

sequence or its length.

Process Dissociation Procedure (PDP). After SRTT retention, participants completed the

PDP. The PDP assesses conscious control over sequence knowledge that may develop

during implicit SRTT paradigms (Destrebecqz et al., 2005; Wilkinson et al., 2009, 2015;

Wilkinson & Jahanshahi, 2007; Wilkinson & Shanks, 2004) and is based on the rationale that

conscious knowledge is both reportable upon request and under intentional control. The

PDP consists of an inclusion and exclusion test, each of which involve 12 unique three-item

fragments of the 12-item sequence that were embedded within the sequence version of the

SRTT (e.g., 2-3-1, 3-1-4, 1-4-3, etc.).

The inclusion test evaluates reportable, consciously accessible knowledge. During the

inclusion test, participants were presented with two elements of each of 12 three-item

fragments and were asked to produce the next element in the fragment by pressing the

corresponding button. The exclusion test evaluates intentional control. During the exclusion

test, participants were again presented with two elements of each of 12 three-item fragments

and were asked to *avoid* producing the next element in the fragment by pressing any button

other than the one that corresponded to the next element in the fragment. At the beginning

of the inclusion and exclusion tests, participants were informed that repeated element

sequences (i.e., a 2 immediately after a 2) were not allowed. For each trial of the inclusion

test, there was a 1/3 probability of pressing the correct key by chance. For each trial of the

exclusion test, there was a 2/3 probability of pressing the correct key by chance. The order

of inclusion and exclusion tests was counterbalanced across participants in both groups.

Data analysis

<u>Serial Reaction Time Task (SRTT).</u>

Reaction times (RTs) were averaged within each block. We defined acquisition and retention

skill as the difference between the mean RT of the last random block and last sequence

block during SRTT acquisition and the difference between the mean RT of the first random

block and first sequence block during SRTT retention, respectively. We also defined offline

changes in skill as the difference between acquisition skill and retention skill. Here, positive

values reflect offline losses while negative values reflect offline gains.

$$Acquisition\ skill = RT_{Acquisition\ block\ 14} - RT_{Acquisition\ block\ 13}$$

$$Retention\ skill = RT_{Retention\ block\ 1} - RT_{Retention\ block\ 2}$$

$$Offline\ skill\ changes = Acquisition\ skill - Retention\ skill$$

<u>Process Dissociation Procedure.</u> Keypresses during inclusion trials were considered

accurate when the participant correctly completed the 3-element fragment by pressing the

correct key in the sequence. Keypresses during exclusion trials were considered accurate

when the participant correctly avoided the next element in the fragment by pressing one of

the two other key choices. The total number of accurate trials in the inclusion and exclusion

conditions were compared to the corresponding chance levels (i.e., 4/12 for the inclusion test

and 8/12 for the exclusion test) to test for explicit awareness of the embedded sequence.

<u>EEG analysis.</u> EEG signals were processed using a combination of Fieldtrip (Oostenveld et

al., 2010) and custom-written MATLAB scripts. To evaluate phase targeting accuracy, we

segmented EEG data into 6 s segments (± 3 s relative to the targeted phase), and then

demeaned and detrended these segments. To ensure that all offline data analysis

procedures mimicked those applied during real-time phase targeting, we applied the Hjorth

transformation to the EEG data by subtracting the mean of four surrounding sensorimotor

electrodes (FC3, FC5, CP3 and CP5) from the center sensorimotor electrode (C3). Resulting

data segments were zero-phase bandpass filtered using a FIR filter (8-13 Hz). The

instantaneous phase of the filtered signal at the time of phase targeting was identified using

the Hilbert transformation. We then combined phase angles across all subjects per group

(see Figure 2).

To calculate pre-stimulus sensorimotor mu rhythm power, continuous EEG data obtained

during TMS delivery were segmented into 500 ms segments (0.51 to 0.01 seconds prior to

each TMS pulse), demeaned, linearly detrended and Hjorth transformed (center = C3,

surround = FC1, FC5, CP1, CP5; Hjorth, 1975; Zrenner et al., 2018; Hussain et al., 2021;

Suresh & Hussain, 2023). The resulting Hjorth-transformed signals were then spectrally

decomposed (Welch's method, 4-50 Hz with 0.25 Hz resolution, Suresh & Hussain, 2023).

Each trial's power values were averaged in the mu (8-13 Hz) range to generate a mixed mu

power estimate (i.e., the combination of aperiodic and periodic components of mu power;

Donoghue et al., 2020). Mixed mu power was calculated instead of periodic mu power as our

previous work demonstrated that statistical models including mu rhythm phase and mixed

mu power explain MEP amplitude variation better than those including mu phase and

periodic mu power (Suresh & Hussain, 2023).

MEP analysis. Pre-stimulus EMG activity was filtered with a discrete Fourier transform filter

to reduce line noise. All trials were visually inspected, and those that precluded reliable

calculation of peak-to-peak MEP amplitudes were rejected (3.3% of all trials). We calculated

background EMG as the root-mean-square of pre-stimulus EMG activity between 100 and

25 ms before each TMS pulse. Peak-to-peak MEP amplitudes were calculated as the

maximum voltage deflection between 20 and 40 ms after each TMS pulse. MEP amplitudes

for rejected trials were treated as missing data during statistical analysis. MEP amplitudes

were log-transformed to reduce positive skew and improve statistical model fits (Hussain et

al., 2019; Suresh & Hussain, 2023).

Statistical analysis

Phase targeting accuracy. Phase angle distributions at the time of phase targeting were

evaluated for deviations from a circular uniform distribution in the 90° (peak phases) and

270° (trough phases) directions using separate V-tests for each group. For random phases,

general deviations from circular uniformity were tested using Rayleigh's test. For each group,

mu peak and trough phase angle distributions were compared using the Watson-Williams

test. Phase angle distributions for each condition (peak, trough, and random) were also

compared between groups using the Watson-Williams test.

SRTT acquisition and retention. To evaluate changes in mean RT during SRTT acquisition

and retention, we fit separate linear mixed-effects models (LMMs) for acquisition and

retention. These included mean RT as the response variable, BLOCK (categorical variable:

acquisition blocks 1-14, retention blocks 1-7), GROUP (categorical variable: sequence,

control), and BLOCK x GROUP interactions as fixed effects. PARTICIPANT was included as

random intercept. To evaluate changes in skill from SRTT acquisition to retention, we fit an

LMM that included the change in mean RT as the response variable, TASK (categorical

variable: acquisition, retention, offline change), GROUP (categorical variable: sequence,

control), and TASK x GROUP interactions as fixed effects. Post-hoc pairwise comparisons

were used to compare SRTT acquisition and SRTT retention between groups (sequence

versus control). To evaluate offline skill, we compared offline changes to zero using separate

single-sample t-tests for the sequence and control group.

Process dissociation procedure (PDP). To assess explicit sequence awareness, we

compared fragment completion accuracy to chance using separate single-sample right-tailed

t-tests for the inclusion and exclusion tests. One participant in the control group did not

complete the PDP.

MEP analysis. We first examined the influence of SRTT acquisition on phase-independent

MEP amplitudes using an LMM with GROUP (categorical variable: sequence, control), TIME

(categorical variable: baseline, immediately after SRTT acquisition, 30 minutes after SRTT

acquisition), and the GROUP x TIME interaction as fixed factors, natural log-transformed

POWER (continuous variable) and background EMG (continuous variable) as covariates,

and natural log-transformed MEP amplitudes elicited during random mu phases as the

response variable. The random intercept of PARTICIPANT was also included (Hussain et

al., 2019; Hussain et al., 2021; Suresh & Hussain, 2023).

Next, we evaluated the influence of SRTT task performance on phase-dependent MEP

amplitudes by fitting an LMM that included fixed effects of GROUP (categorical variable:

sequence, control), PHASE (categorical variable: peak, trough), TIME (categorical variable:

baseline, immediately after SRTT acquisition, 30 minutes after SRTT acquisition) and the

GROUP x PHASE x TIME interaction as fixed factors, the PHASE x natural log-transformed

POWER interaction and background EMG (continuous variable) as covariates, and natural

log-transformed MEP amplitudes as the response variable. PARTICIPANT was included as

the random intercept.

Given the PHASE x POWER interaction identified in our previous work and that of others

(Bigoni et al., 2024; Hussain et al., 2019; Ozdemir et al., 2022; Suresh & Hussain, 2023) and

our primary interest in evaluating phase-dependent corticospinal plasticity, we included the

PHASE x natural log-transformed POWER interaction as a covariate in all our LMMs to

control for this known interaction effect (Hussain et al., 2019; Suresh & Hussain, 2023).

Furthermore, it is well established that pre-stimulus voluntary muscle activation can increase

MEP amplitudes (Hess et al., 1996). To account for any variation in MEP amplitudes caused

by pre-stimulus muscle activation, we also included pre-stimulus background EMG activity

as a covariate in our LMMs. The significance of fixed effects was determined using likelihood

ratio tests. All reported β-coefficients represent the difference in estimates between groups

and phases.

All statistical analyses were performed in R (version 4.3.1). Model fits were visually

inspected using histograms of residuals and quantile-quantile plots. Alpha was equal to 0.05

for all comparisons and p-values were corrected using the Tukey's Honestly Significant

Difference test for multiple comparisons where appropriate. Data are expressed as mean ±

standard error of the mean.

Results

Phase targeting accuracy

Real-time EEG analysis accurately identified and targeted mu peak (Figure 2, target phase =

90°, observed mean phase for sequence group = 85.7° [95% confidence interval {CI} =

80.27°– 91.12°], V = 237.9, p < 0.001; observed mean phase for control group = 86.93°

[95% CI = 80.77°– 93.10°], V = 217.52, p < 0.001) and mu trough phases (target phase =

270°, observed mean phase for sequence group = 271.72° [95% CI = 266.17° – 277.25°], V

= 56.01, p < 0.001; observed mean phase for control group = 269.66° [95% CI = 263.99° –

275.31°], V = 45.71, p = 0.001). As expected, targeted phases differed significantly between

peak and trough conditions in both the sequence group (F = 1687.0, p < 0.001) and control

group (F = 1495.8, p < 0.001). For random phases, we observed a deviation from circular

uniformity in the 20° direction in the sequence group (R = 5.56, p = 0.003, observed mean

phase = 19.05° [95% CI = 343.15° – 54.95°]) and no significant deviation from uniformity in

the control group (R = 2.47, p = 0.08, observed mean phase = 12.93° [95% CI = 311.2° –

74.66°]). However, we observed no significant differences in phase targeting accuracy

between the sequence and control group for any phase (peak: F = 0.11, p = 0.741; trough: F

= 0.341, p = 0.559; random: F = 0.188, p = 0.665).

Figure 2. Phase targeting accuracy. A, B) Group-averaged raw and filtered (8-13 Hz) EEG
signals recorded over left sensorimotor cortex during evaluation of mu phase targeting
accuracy in the sequence (A) and control (B) groups. Black vertical lines reflect the time
point at which the real-time EEG analysis algorithm identified mu peak, trough, and random
phases. Shading represents mean ± 1 SEM. **C, D)** Circular histograms indicate the phase
angle of the sensorimotor mu (8-13 Hz) rhythm at the time of phase targeting for peak,
trough, and random phases in the sequence (C) and control (D) groups. **E, F)** Scalp
distribution of C3-Hjorth transformed and band-pass filtered EEG activity during mu peak,
trough and random phases at the time of phase targeting for peak, trough, and random
phases in the sequence (E) and control (F) groups. Topographical plots reflect average
scalp-recorded EEG signals within a 5 ms window centered on the time of phase targeting.
Bolded channels indicate those used to calculate C3-Hjorth transformed EEG data. Asterisks
reflect significance at $p < 0.05$.

SRTT

During SRTT acquisition, participants in both the sequence and control groups decreased

their mean RTs across blocks (Figure 3, main effect of BLOCK: $F = 12.96$ and $p < 0.001$).

RTs did not differ between groups for random blocks, but participants in the sequence group

tended to show faster mean RTs during the sequence blocks than the control group during

the corresponding blocks (BLOCK x GROUP interaction: $F = 9.37$ and $p < 0.001$; no main

effect of GROUP: $F = 0.63$ and $p = 0.42$). However, post hoc pairwise comparisons did not

identify any significant differences in mean RTs between groups during these blocks (i.e.,

blocks 3-7 or 9-13, see Figure 3). For both groups, mean RTs were not statistically

distinguishable between the first and last random blocks (blocks 1 vs. 14, t < 0.165, p =

0.99). For the sequence group, mean RTs were faster during the last sequence block than

during the last random block (block 13 vs. 14, t ratio = -8.51, p < 0.001), but this was not the

case for the control group (t ratio = 0.16, p = 0.99). Finally, acquisition skill was significantly

greater in the sequence than control group (Figure 3, t ratio = 5.94, p < 0.001). Thus, as

expected, participants in the sequence group experienced sequence-specific learning.

During SRTT retention, participants in both groups decreased their mean RTs across blocks

(Figure 3, main effect of BLOCK: F = 18.91 and p < 0.001) and mean RTs were overall

faster for the sequence than the control group (main effect of GROUP: F = 6.04, p = 0.01).

Mean RTs were not statistically distinguishable between groups during random blocks but

were significantly faster for the sequence group during sequence blocks (GROUP x BLOCK

interaction: F = 21.14, p < 0.001). Post hoc pairwise comparisons revealed that mean RTs

were faster for the sequence group than for the control group during blocks 4-6 (t > 3.59, p

values < 0.03). In addition, retention skill was significantly greater in the sequence than the

control group (Figure 3, t ratio = 5.17, p < 0.001). Thus, participants in the sequence group

retained sequence-specific skill obtained during SRTT acquisition. Finally, participants in the

sequence group showed significant offline losses (t = 5.23 p < 0.0001, mean = 98.17 ms

[95% CI = 58.61 ms - 137.72 ms] while those in the control group did not (t = -0.51, p =

0.616, mean = -1.95 ms [95% CI = -10.01 ms - 6.11ms]). However, direct comparison of the

magnitude of these offline changes revealed no significant difference between groups

(Figure 3, t = 0.76, p = 0.97).

Figure 3. **SRTT performance. A)** Participants in the sequence group showed sequence-
specific decreases in mean RTs during SRTT acquisition and retention. Vertical dotted lines
indicate random blocks during SRTT (practice block not shown). Colored asterisks reflect
significance at p < 0.05 for post hoc pairwise comparisons between blocks 13 and 14 during
SRTT acquisition and blocks 1 and 2 during retention (green for sequence group). Shading
represents the mean ± 1 SEM. **B)** Sequence-specific skill acquisition and retention were
greater in the sequence than in the control group. Asterisks reflect significance at p < 0.05
for post hoc comparisons of acquisition, retention and offline skill changes between groups.
Note that for offline changes, positive values reflect offline losses while negative values
reflect offline gains. For boxplots, black lines represent median, white dots represent mean,
black dots represent individual participants, and error bars represent upper and lower inter-
quartile ranges.
492

Explicit sequence awareness

In the sequence group, mean fragment completion accuracies exceeded chance for the

inclusion test (6 ± 0.66 [SEM], t = 3.02, p = 0.007, chance = 4) but not the exclusion test

(7.72 ± 0.54, t = -0.51, p = 0.61, chance = 8). In the control group, fragment completion

accuracies did not exceed chance for either the inclusion or exclusion tests (Figure 4,

inclusion = 3.88 ± 0.45, exclusion = 7.7 ± 0.54, t < -0.52 for both, p > 0.59 for both). Given

that explicit awareness requires both conscious knowledge of the sequence and the ability to

intentionally control sequence production, there was weak evidence of explicit sequence

awareness in both groups.

Figure 4. Process Dissociation Procedure. Mean number of fragments completed or
avoided during the inclusion and exclusion portions of the PDP. The dashed lines reflect
chance performance (4/12 for inclusion and 8/12 for exclusion. For boxplots, black lines
represent group median, white dots represent group mean, black dots represent individual
participants, and error bars represent upper and lower inter-quartile ranges.

Phase-independent changes in MEP amplitudes

We first characterized the effects of SRTT acquisition on mu phase-independent MEP

amplitudes elicited during random mu phases. In both groups, MEP amplitudes increased

following SRTT acquisition (significant main effect of TIME; F = 27.05, p < 0.001, no main

effect of GROUP; F = 0.67, p = 0.41, no GROUP x TIME interaction, F = 1.25, p = 0.28).

Post hoc pairwise comparisons revealed that random, phase-independent MEP amplitudes

increased from baseline to immediately after SRTT acquisition (β = -0.08, t ratio = -2.47, p =

0.03), from baseline to 30 minutes after SRTT acquisition (β = -0.24, t ratio = -7.24, p <

0.0001), and from immediately to 30 minutes after SRTT acquisition (β = -0.16, t ratio = -

4.76, p < 0.001). Thus, phase-independent MEP amplitudes increased following SRTT

acquisition in both groups.

Phase-dependent changes in MEP amplitudes

We next evaluated the effects of SRTT acquisition on mu phase-dependent MEP amplitudes

elicited during mu peak and trough phases. We found a significant three-way GROUP x

TIME x PHASE interaction (Figure 5, likelihood ratio test: F = 8.53, p = 0.0001). However,

this three-way interaction was only present when comparing changes in MEP amplitudes

from baseline to 30 minutes after SRTT acquisition (no GROUP x TIME x PHASE interaction

for immediately after SRTT: $t = -1.691$, $p = 0.09$, significant GROUP x TIME x PHASE

interaction for 30 minutes after SRTT: $t = 2.42$, $p = 0.01$), indicating that the magnitude of

MEP amplitude changes from baseline to 30 minutes after SRTT acquisition more strongly

differed across phases in one group than the other. As a result, post hoc pairwise

comparisons were only performed for changes in MEP amplitudes from baseline to 30

minutes after SRTT acquisition. For the control group, there was a significant difference

between mu peak and trough MEP amplitudes 30 minutes after SRTT acquisition ($\beta = -0.22$,

z ratio = -4.42, $p = 0.0006$). However, for the sequence group, pairwise comparisons showed

no difference between mu peak and trough MEP amplitudes 30 minutes after SRTT

acquisition ($\beta = -0.03$, z ratio = -0.76, $p = 0.99$). Thus, the sequence group showed

significantly less phase-dependent variation in MEP amplitude 30 minutes after SRTT

acquisition than the control group.

In the control group, mu peak MEP amplitudes increased from baseline to immediately after

SRTT acquisition ($\beta = -0.21$, $z = -4.24$, $p = 0.001$) and from baseline to 30 minutes after

SRTT acquisition ($\beta = -0.27$, $z = -5.47$, $p < 0.0001$). Mu peak MEP amplitudes did not

significantly change between immediately and 30 minutes after SRTT acquisition ($\beta = -0.05$,

$z = -1.1$, $p = 0.99$). In contrast, mu trough MEP amplitudes increased from baseline to

immediately after SRTT acquisition ($\beta = -0.18$, $z = -3.7$, $p = 0.0013$), baseline to 30 minutes

after SRTT acquisition ($\beta = -0.42$, $z = -8.48$, $p < 0.0001$), and immediately after to 30 minutes

after SRTT acquisition ($\beta = -0.23$, $z = -4.79$, $p = 0.0001$). Overall, participants in the control

group exhibited significantly smaller increases in MEP amplitudes during mu peak than

trough phases 30 minutes after SRTT acquisition.

In the sequence group, mu peak MEP amplitudes increased from baseline to 30 minutes

after SRTT acquisition ($\beta = -0.35$, z ratio = -7.12, $p < 0.0001$) and immediately to 30 minutes

after SRTT acquisition (β = -0.25, z = -5.07, p < 0.0001). Mu peak MEP amplitudes did not

significantly change between baseline and immediately after SRTT acquisition (β = -0.1, z =

-2.01, p = 0.68). In addition, mu trough MEP amplitudes increased from baseline to

immediately after SRTT acquisition (β = -0.23, z = -4.78, p = 0.001) and baseline to 30

minutes after SRTT acquisition (β = -0.26, z = -5.28, p < 0.0001) with no significant change

between immediately and 30 minutes after SRTT acquisition (β = -0.02, z = -0.50, p = 0.99).

Overall, participants in the sequence group did not show significantly different increases in

MEP amplitudes between mu peak and trough phases at any time point.

To visualize differences in mu phase-dependent MEP amplitude changes between groups,

we subtracted control group model-fitted MEP amplitudes from sequence group model-fitted

MEP amplitudes (Figure 6). This comparison revealed that the observed GROUP x PHASE

x TIME interaction was driven by sequence learning-related increases mu peak-specific

corticospinal plasticity.

Figure 5. Phase-independent and phase-dependent changes in corticospinal output.
A) Group-level linear mixed-effects model coefficients depicting relationships between
phase-independent MEP amplitudes and group at baseline, immediately and 30 minutes
after SRTT acquisition. **B)** Group-level linear mixed-effects model coefficients depicting
relationships between phase-dependent MEP amplitudes and group at baseline,
immediately, and 30 minutes after SRTT acquisition. For A and B, dots represent model
coefficients and error bars indicate SEM. For A, black asterisks reflect significant post hoc
pairwise comparisons across time points. For B, colored asterisks reflect significant post hoc
pairwise comparisons across time points for each phase (orange for peak, blue for trough).

Discussion

We hypothesized that learning-related corticospinal plasticity would be more evident when

measured during sensorimotor mu rhythm trough than peak phases. To test this hypothesis,

we delivered single-pulse, real-time EEG-informed TMS during mu rhythm peak, trough, and

random phases immediately before and after healthy adults performed a version of the

implicit SRTT that included either an embedded, repeating motor sequence (sequence

group) or no sequence (control group). Behaviorally, participants in the sequence group

acquired greater sequence-specific knowledge than participants in the control group and

retained this sequence-specific knowledge over time. Physiologically, participants in both the

sequence and control groups showed phase-independent increases in corticospinal output

following SRTT acquisition, reflecting the occurrence of significant phase-independent

corticospinal plasticity that did not differ between groups. However, the two groups showed

different patterns of phase-dependent corticospinal plasticity following SRTT acquisition.

Although we did observe that motor sequence learning elicited phase-dependent

corticospinal plasticity, our results show that corticospinal plasticity induced by motor

sequence learning was more prominent during mu rhythm *peak* rather than trough phases.

Motor learning potentiates corticospinal output (Pascual-Leone et al., 1994; Perez et al.,

2007) and occludes subsequent induction of LTP-like corticospinal plasticity (Cantarero et

al., 2013a; Cantarero et al., 2013b). This occlusion is thought to support retention of newly-

acquired motor skills, such that its experimental reversal significantly diminishes retention

(Cantarero et al., 2013a). Thus, LTP-like corticospinal plasticity and its occlusion both play

key mechanistic roles in motor learning. Like motor learning, repetitive TMS interventions

can also induce corticospinal plasticity. Specifically, recent studies using real-time EEG-TMS

have shown that both low- and high-frequency repetitive TMS interventions preferentially

induce LTP-like corticospinal plasticity when applied during sensorimotor mu trough but not

peak phases (Baur et al., 2020; Zrenner et al., 2018). Such mu phase-dependent

corticospinal plasticity could therefore reflect a common neurophysiological mechanism

underlying the effects of repetitive TMS interventions and motor learning. If mu phase-

dependent corticospinal plasticity does support motor learning, then TMS interventions

should more strongly affect motor learning when delivered during specific mu phases. We

previously evaluated this possibility by measuring the effects of a low-frequency mu phase-

dependent repetitive TMS intervention on offline changes in explicit motor sequence task

performance (Hussain et al., 2021). In that study, repetitive TMS delivered during mu trough

but not peak phases enhanced offline gains in motor sequence performance. These findings

raised the intriguing possibility that the neurophysiological mechanisms underlying motor

learning may be most active and thus most susceptible to modification during sensorimotor

mu trough phases. However, in the current study, mu phase-dependent corticospinal

plasticity was less evident during mu peak than trough phases in the control group, but

similarly evident across phases in the sequence group.

What could explain these surprising findings? Our experimental design directly compared

changes in MEP amplitudes following performance of one of two related motor tasks. In the

sequence group, participants performed repeated keypresses while also acquiring

knowledge of an embedded sequence, while in the control group, participants performed

repeated keypresses in the absence of any embedded sequence. To isolate the effects of

sequence learning on mu phase-dependent corticospinal plasticity, we therefore subtracted

control group model-fitted MEP amplitudes from sequence group model-fitted MEP

amplitudes. This contrast revealed that sequence learning, after accounting for the effects of

repeated keypresses, induced mu phase-dependent corticospinal plasticity that was more

evident during mu *peak* phases. We previously proposed that mu rhythm peak phases

reflect a neutral excitability state that is not strongly affected by synchronization of

corticothalamic inputs to layer V pyramidal cells (Hussain et al., 2019). Specifically, M1

single neuron spiking rates are lower during mu rhythm peak than trough phases in non-

human primates (Haegens et al., 2011), which may correspond to the decreased

corticospinal output observed during mu peak relative to trough phases in humans

(Bergmann et al., 2019; Hussain et al., 2019; Suresh & Hussain, 2023). Indeed, Bergmann

et al. (2019) showed that mu peak phases reflect a lack of pulsed excitation rather than

pulsed inhibition of corticospinal output. Together, these previous findings suggest that mu

peaks reflect brief quiescent windows during which corticospinal and sensorimotor network

communication is diminished relative to mu troughs. As a result, neural activity should only

weakly propagate through an extended sensorimotor network during mu peak phases. We

and others previously provided empirical support for this possibility by showing that (1) TMS-

related whole-brain broadband power, an indirect marker of population-level neuronal

spiking (Gao et al., 2017; Manning et al., 2009), is weaker when M1 TMS delivery coincides

with mu rhythm peak than trough phases (Hussain et al., 2021), and (2) interhemispheric

communication between bilateral motor cortices is lower during mu peak than trough phases

(Stefanou et al., 2018). The decreased network communication that occurs during mu peak

phases may prevent motor sequence memories from being disrupted by task-irrelevant

sensory information carried by thalamocortical projections, such that sequence memories

can be replayed and consolidated by sensorimotor and hippocampal networks without

interference (Buch et al., 2021; Jacobacci et al., 2020; Robertson & Genzel, 2020). This idea

aligns with previous work showing that wakeful rest after motor sequence learning supports

consolidation and retention of the learned sequence (Bönstrup et al., 2019, 2020; Buch et

al., 2021; Robertson, 2019; Robertson et al., 2004; Walker et al., 2003). Mu peak phases

could therefore provide a protected time window during which replay and consolidation of

motor sequence memories may occur. Such a scenario could explain why sequence

learning-related corticospinal plasticity was more pronounced during mu peak phases in the

current study. Future studies could directly evaluate this possibility by examining

sensorimotor mu phase-dependency of motor sequence replay or altering mu peak phase

properties immediately following motor sequence learning using closed-loop neurofeedback

paradigms.

Figure 6. Effects of motor sequence learning on mu phase-dependent corticospinal plasticity. At each time point, group-level model fitted MEP amplitudes obtained during mu peak and trough phases in the control group were subtracted from corresponding group-level model fitted MEP amplitudes in the sequence group. Asterisk reflects significant 3-way interaction of GROUP X PHASE X TIME.

Although we previously suggested that the neurophysiological mechanisms underlying motor

sequence learning may be most active during sensorimotor mu trough phases (Hussain et

al., 2021), here we report that corticospinal plasticity following motor sequence learning is

more evident during mu peak than trough phases. Two key differences between our

previous work and the current study could explain these diverging results. First, the current

study used the implicit SRTT, while our previous work used the explicit motor sequence

learning task. Although both tasks involve acquisition of sequential knowledge, they rely on

dissimilar sensorimotor networks. Specifically, the implicit SRTT likely more heavily recruits

corticostriatal circuits (Grafton et al., 1995; Hazeltine et al., 1997), while the explicit motor

sequence learning task may more strongly engage prefrontal and premotor areas (Pinsard et

al., 2019; Yokoi & Diedrichsen, 2019). Given the fundamental role of oscillatory phase in

inter-regional neural communication (Fries, 2005, 2015; Womelsdorf et al., 2007), these two

brain networks may optimally interact with sensorimotor cortical neuronal populations during

different mu rhythm phases. Second, the current study directly assessed within-session

phase-dependent neurophysiological changes induced by motor sequence learning,

whereas our previous work examined the influence of a phase-dependent TMS intervention

on between-session behavioral changes in motor sequence performance. Thus,

corticospinal plasticity supporting motor sequence learning may be most evident during mu

peak phases, while TMS interventions may best improve motor sequence learning during mu

trough phases. Along these lines, it is possible that the neurophysiological mechanisms

supporting motor sequence learning are maximally active during mu peak phases, such that

peak-specific TMS interventions cannot further upregulate them. If so, the same

neurophysiological mechanisms may be only partially active during mu trough phases,

allowing them to be further enhanced by trough-specific TMS interventions. Regardless, our

current findings and previous work both support the notion that the neurophysiological

mechanisms underlying motor sequence learning are sensorimotor mu phase-dependent,

although the precise contributions of mu peak and trough phases to different aspects of such

learning require further investigation.

Unexpectedly, we observed that mu phase-dependent variation in corticospinal output

emerged 30 minutes after SRTT acquisition in the control group. One possible explanation

for this observation is that the simple repetitive keypress movements performed during the

control version of the SRTT version induced use-dependent plasticity that is biased towards

mu rhythm trough phases. Given that acquiring real-world motor skills involves multiple

learning mechanisms, sequence-dependent learning mechanisms that are most evident

during mu peak phases could co-occur with use-dependent plasticity mechanisms that may

be biased towards mu trough phases. However, our study was not specifically designed to

address this possibility. Future work aimed at establishing relationships between use-

dependent plasticity and sensorimotor rhythm phase should specifically examine phase-

dependent directional biases of TMS-evoked movements, as in previous studies (Bütefisch

et al., 2000; Mawase et al., 2017).

Our findings may also have implications for motor learning after stroke. Stroke-related brain

lesions can either directly or indirectly alter sensorimotor rhythms by disrupting their cortical

sources (Salmelin & Hari, 1994), their subcortical pacemaker cells (Hughes & Crunelli,

2005), or anatomical connections between them. For example, sensorimotor mu rhythms

weaken and slow after stroke (Hussain et al., 2020; Johnston et al., 2023). Given the role of

sensorimotor mu rhythm phase in learning-related corticospinal plasticity identified here,

stroke-related changes in mu rhythm properties could interfere with the neurophysiological

mechanisms underlying motor learning, limiting motor skill (re)acquisition and retention

during neurorehabilitation. Accordingly, the presence or absence of mu phase-dependent

corticospinal plasticity in stroke survivors could predict each individual's capacity for

rehabilitation-related motor recovery, and strategically delivering mu phase-specific TMS

interventions prior to rehabilitative training could enhance rehabilitation's effects in

individuals with intact sensorimotor mu rhythm-generating circuits. Indeed, the combined

effects of mu phase-dependent TMS interventions and motor rehabilitation on poststroke

upper extremity motor recovery are currently being evaluated via a multi-center randomized

controlled trial (Lieb et al., 2023). However, nearly all studies evaluating sensorimotor mu

phase-dependency within the human motor system have focused on neurotypical adults

(Baur et al., 2020; Bergmann et al., 2019; Stefanou et al., 2018; Suresh & Hussain, 2023;

Zrenner et al., 2018), and whether similar mu phase-dependent mechanisms support TMS-

induced and learning-related corticospinal plasticity after stroke is still unknown.

Several strengths of this study exist. First, our experimental design included two groups: one

that experienced sequence learning and one that did not. This design allowed us to

dissociate the effects of sequence learning and repeated keypresses on phase-dependent

corticospinal plasticity. Of note, non-specific effects of task engagement are often not

considered in motor sequence learning studies, including our own previous work (Hussain et

al., 2021; Bönstrup et al., 2019, 2020; Perez et al., 2007; Walker et al., 2003). Without the

control group, the current study would have failed to identify any role of phase-dependent

corticospinal plasticity in motor sequence learning. Second, participants did not have to meet

any pre-defined neurophysiological criteria to participate in this study. Thus, our results are

likely to be more generalizable across diverse groups of participants than previous studies

using similar techniques (Baur et al., 2020; Zrenner et al., 2018). Finally, rather than inferring

the existence of a phase-dependent learning mechanism from interventional studies

(Hussain et al., 2021; Baur et al., 2020; Zrenner et al., 2018), here we specifically quantified

phase-dependent neurophysiological changes induced by sequence learning itself. Thus,

the results of the current study can be directly attributed to sequence learning *per se* rather

than the combined effects of both sequence learning and a phase-dependent TMS

intervention.

While our study provides novel insights into phase-dependent mechanisms of motor

learning, it is not without limitations. First, previous studies have shown that mu rhythm

phase and power interdependently influence corticospinal output at rest, but here we

evaluated only mu phase-dependent changes in corticospinal output. Although this was

achieved by introducing a covariate into our statistical models that controlled for mu phase-

power interactions, this approach does not capture more complex learning-related changes

in mu phase-power interactions. Second, we did not measure changes in corticospinal

output during other mu phases (Wischnewski et al., 2022; Zrenner et al., 2023), other

sensorimotor rhythms (Hussain et al., 2019; Wischnewski et al., 2022) or broadband activity

(Davis et al., 2022). Experimental designs using phase-independent measures of

corticospinal excitability and post hoc trial sorting approaches may be better suited to

explore the contribution of other brain rhythms and phases to learning-related corticospinal

plasticity. Finally, the SRTT requires participants to activate several intrinsic and extrinsic

hand muscles, and we only recorded MEP amplitudes from the FDI. Future studies that

record MEPs from multiple task-relevant muscles could provide richer information on

learning-related corticospinal plasticity.

In conclusion, we report that corticospinal plasticity induced by motor sequence learning is

increased during sensorimotor mu rhythm peak relative to trough phases, providing first

direct evidence that motor learning induces phase-dependent corticospinal plasticity in

humans. Overall, our findings suggest that mu peak phases may provide brief protected time

windows during which motor sequence memory consolidation occurs and establish the

mechanistic role of mu rhythm phase in behaviorally-relevant corticospinal plasticity.

References

Baumgarten, T. J., Schnitzler, A., & Lange, J. (2015). Beta oscillations define discrete
  perceptual cycles in the somatosensory domain. *Proceedings of the National
  Academy of Sciences*, *112*(39), 12187–12192.
  https://doi.org/10.1073/pnas.1501438112

Baur, D., Galevska, D., Hussain, S., Cohen, L. G., Ziemann, U., & Zrenner, C. (2020).
  Induction of LTD-like corticospinal plasticity by low-frequency rTMS depends on pre-
  stimulus phase of sensorimotor µ-rhythm. *Brain Stimulation*, *13*(6), 1580–1587.
  https://doi.org/10.1016/j.brs.2020.09.005

Bergmann, T. O., Lieb, A., Zrenner, C., & Ziemann, U. (2019). Pulsed Facilitation of
  Corticospinal Excitability by the Sensorimotor µ-Alpha Rhythm. *Journal of
  Neuroscience*, *39*(50), 10034–10043. https://doi.org/10.1523/JNEUROSCI.1730-
  19.2019

Bigoni, C., Pagnamenta, S., Cadic-Melchior, A., Bevilacqua, M., Harquel, S., Raffin, E., &
  Hummel, F. C. (2024). MEP and TEP features variability: Is it just the brain-state?
  Journal of Neural Engineering. https://doi.org/10.1088/1741-2552/ad1dc2

Bönstrup, M., Iturrate, I., Hebart, M. N., Censor, N., & Cohen, L. G. (2020). Mechanisms of
  offline motor learning at a microscale of seconds in large-scale crowdsourced data.
  Npj Science of Learning, *5*(1), Article 1. https://doi.org/10.1038/s41539-020-0066-9

Bönstrup, M., Iturrate, I., Thompson, R., Cruciani, G., Censor, N., & Cohen, L. G. (2019). A
  Rapid Form of Offline Consolidation in Skill Learning. *Current Biology*, *29*(8), 1346-
  1351.e4. https://doi.org/10.1016/j.cub.2019.02.049

Borckardt, J. J., Nahas, Z., Koola, J., & George, M. S. (2006). Estimating Resting Motor
  Thresholds in Transcranial Magnetic Stimulation Research and Practice: A computer
  Simulation Evaluation of Best Methods. *The Journal of ECT*, *22*(3), 169.
  https://doi.org/10.1097/01.yct.0000235923.52741.72

Buch, E. R., Claudino, L., Quentin, R., Bönstrup, M., & Cohen, L. G. (2021). Consolidation of human skill linked to waking hippocampo-neocortical replay. *Cell Reports*, *35*(10). https://doi.org/10.1016/j.celrep.2021.109193

Busch, N. A., Dubois, J., & VanRullen, R. (2009). The Phase of Ongoing EEG Oscillations Predicts Visual Perception. *Journal of Neuroscience*, *29*(24), 7869–7876. https://doi.org/10.1523/JNEUROSCI.0113-09.2009

Busch, N. A., & VanRullen, R. (2010). Spontaneous EEG oscillations reveal periodic sampling of visual attention. *Proceedings of the National Academy of Sciences*, *107*(37), 16048–16053. https://doi.org/10.1073/pnas.1004801107

Bütefisch, C. M., Davis, B. C., Wise, S. P., Sawaki, L., Kopylev, L., Classen, J., & Cohen, L. G. (2000). Mechanisms of use-dependent plasticity in the human motor cortex. *Proceedings of the National Academy of Sciences*, *97*(7), 3661–3665. https://doi.org/10.1073/pnas.97.7.3661

Cantarero, G., Lloyd, A., & Celnik, P. (2013b). Reversal of Long-Term Potentiation-Like Plasticity Processes after Motor Learning Disrupts Skill Retention. *Journal of Neuroscience*, *33*(31), 12862–12869. https://doi.org/10.1523/JNEUROSCI.1399-13.2013

Cantarero, G., Tang, B., O'Malley, R., Salas, R., & Celnik, P. (2013a). Motor Learning Interference Is Proportional to Occlusion of LTP-Like Plasticity. *Journal of Neuroscience*, *33*(11), 4634–4641. https://doi.org/10.1523/JNEUROSCI.4706-12.2013

Davis, Z. W., Muller, L., & Reynolds, J. H. (2022). Spontaneous Spiking Is Governed by Broadband Fluctuations. *The Journal of Neuroscience*, *42*(26), 5159–5172. https://doi.org/10.1523/JNEUROSCI.1899-21.2022

Destrebecqz, A., Peigneux, P., Laureys, S., Degueldre, C., Fiore, G. D., Aerts, J., Luxen, A., Linden, M. V. D., Cleeremans, A., & Maquet, P. (2005). The neural correlates of implicit and explicit sequence learning: Interacting networks revealed by the process

dissociation procedure. *Learning & Memory*, *12*(5), 480–490.

https://doi.org/10.1101/lm.95605

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara,

A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing

neural power spectra into periodic and aperiodic components. *Nature Neuroscience*,

23(12), Article 12. https://doi.org/10.1038/s41593-020-00744-x

Dugué, L., Marque, P., & VanRullen, R. (2011). The Phase of Ongoing Oscillations Mediates

the Causal Relation between Brain Excitation and Visual Perception. *Journal of*

Neuroscience, *31*(33), 11889–11893. https://doi.org/10.1523/JNEUROSCI.1161-

11.2011

Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through

neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480.

https://doi.org/10.1016/j.tics.2005.08.011

Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. *Neuron*, *88*(1),

220–235. https://doi.org/10.1016/j.neuron.2015.09.034

Fuentemilla, L., Penny, W. D., Cashdollar, N., Bunzeck, N., & Düzel, E. (2010). Theta-

Coupled Periodic Replay in Working Memory. *Current Biology*, *20*(7), 606–612.

https://doi.org/10.1016/j.cub.2010.01.057

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance

from field potentials. *NeuroImage*, *158*, 70–78.

https://doi.org/10.1016/j.neuroimage.2017.06.078

Grafton, S. T., Hazeltine, E., & Ivry, R. (1995). Functional Mapping of Sequence Learning in

Normal Humans. *Journal of Cognitive Neuroscience*, *7*(4), 497–510.

https://doi.org/10.1162/jocn.1995.7.4.497

Haegens, S., Nácher, V., Luna, R., Romo, R., & Jensen, O. (2011). α-Oscillations in the

monkey sensorimotor network influence discrimination performance by rhythmical

inhibition of neuronal spiking. *Proceedings of the National Academy of Sciences*,

108(48), 19377–19382. https://doi.org/10.1073/pnas.1117190108

Hasselmo, M. E., Bodelón, C., & Wyble, B. P. (2002). A Proposed Function for Hippocampal

Theta Rhythm: Separate Phases of Encoding and Retrieval Enhance Reversal of

Prior Learning. *Neural Computation*, *14*(4), 793–817.

https://doi.org/10.1162/089976602317318965

Hazeltine, E., Grafton, S. T., & Ivry, R. (1997). Attention and stimulus characteristics

determine the locus of motor-sequence encoding. A PET study. *Brain*, *120*(1), 123–

140. https://doi.org/10.1093/brain/120.1.123

Hess, G., Aizenman, C. D., & Donoghue, J. P. (1996). Conditions for the induction of long-

term potentiation in layer II/III horizontal connections of the rat motor cortex. *Journal*

of Neurophysiology, *75*(5), 1765–1778. https://doi.org/10.1152/jn.1996.75.5.1765

Hjorth, B. (1975). An on-line transformation of EEG scalp potentials into orthogonal source

derivations. *Electroencephalography and Clinical Neurophysiology*, *39*(5), 526–530.

https://doi.org/10.1016/0013-4694(75)90056-5

Hughes, S. W., & Crunelli, V. (2005). Thalamic Mechanisms of EEG Alpha Rhythms and

Their Pathological Implications. *The Neuroscientist*, *11*(4), 357–372.

https://doi.org/10.1177/1073858405277450

Hussain, S. J., Claudino, L., Bönstrup, M., Norato, G., Cruciani, G., Thompson, R., Zrenner,

C., Ziemann, U., Buch, E., & Cohen, L. G. (2019). Sensorimotor Oscillatory Phase–

Power Interaction Gates Resting Human Corticospinal Output. *Cerebral Cortex*,

29(9), 3766–3777. https://doi.org/10.1093/cercor/bhy255

Hussain, S. J., Cohen, L. G., & Bönstrup, M. (2019). Beta rhythm events predict corticospinal

motor output. *Scientific Reports*, *9*(1), 18305. https://doi.org/10.1038/s41598-019-

54706-w

Hussain, S. J., Hayward, W., Fourcand, F., Zrenner, C., Ziemann, U., Buch, E. R., Hayward,

M. K., & Cohen, L. G. (2020). Phase-dependent transcranial magnetic stimulation of

the lesioned hemisphere is accurate after stroke. *Brain Stimulation: Basic,*

Translational, and Clinical Research in Neuromodulation, *13*(5), 1354–1357.

https://doi.org/10.1016/j.brs.2020.07.005

Hussain, S. J., Vollmer, M. K., Stimely, J., Norato, G., Zrenner, C., Ziemann, U., Buch, E. R.,

& Cohen, L. G. (2021). Phase-dependent offline enhancement of human motor

memory. *Brain Stimulation*, *14*(4), 873–883. https://doi.org/10.1016/j.brs.2021.05.009

Jacobacci, F., Armony, J. L., Yeffal, A., Lerner, G., Amaro, E., Jovicich, J., Doyon, J., &

Della-Maggiore, V. (2020). Rapid hippocampal plasticity supports motor sequence

learning. *Proceedings of the National Academy of Sciences*, *117*(38), 23898–23903.

https://doi.org/10.1073/pnas.2009576117

Johnston, P. R., McIntosh, A. R., & Meltzer, J. A. (2023). Spectral slowing in chronic stroke

reflects abnormalities in both periodic and aperiodic neural dynamics. *NeuroImage:*

Clinical, *37*, 103277. https://doi.org/10.1016/j.nicl.2022.103277

Kerrén, C., Linde-Domingo, J., Hanslmayr, S., & Wimber, M. (2018). An Optimal Oscillatory

Phase for Pattern Reactivation during Memory Retrieval. *Current Biology*, *28*(21),

3383-3392.e6. https://doi.org/10.1016/j.cub.2018.08.065

Kothe, C. (2014). *Lab streaming layer (LSL)*.

Lieb, A., Zrenner, B., Zrenner, C., Kozák, G., Martus, P., Grefkes, C., & Ziemann, U. (2023).

Brain-oscillation-synchronized stimulation to enhance motor recovery in early

subacute stroke: A randomized controlled double-blind three- arm parallel-group

exploratory trial comparing personalized, non- personalized and sham repetitive

transcranial magnetic stimulation (Acronym: BOSS-STROKE). *BMC Neurology*,

23(1), 204. https://doi.org/10.1186/s12883-023-03235-1

Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband Shifts in Local Field

Potential Power Spectra Are Correlated with Single-Neuron Spiking in Humans.

Journal of Neuroscience, *29*(43), 13613–13620.

https://doi.org/10.1523/JNEUROSCI.2041-09.2009

Mawase, F., Uehara, S., Bastian, A. J., & Celnik, P. (2017). Motor Learning Enhances Use-

Dependent Plasticity. *Journal of Neuroscience*, *37*(10), 2673–2685.

https://doi.org/10.1523/JNEUROSCI.3303-16.2017

Nissen, M. J., & Bullemer, P. (1987). Attentional requirements of learning: Evidence from

performance measures. *Cognitive Psychology*, *19*(1), 1–32.

https://doi.org/10.1016/0010-0285(87)90002-8

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh

inventory. *Neuropsychologia*, *9*(1), 97–113. https://doi.org/10.1016/0028-

3932(71)90067-4

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological

Data. *Computational Intelligence and Neuroscience*, *2011*, e156869.

https://doi.org/10.1155/2011/156869

Ozdemir, R. A., Kirkman, S., Magnuson, J. R., Fried, P. J., Pascual-Leone, A., & Shafi, M.

M. (2022). Phase matters when there is power: Phasic modulation of corticospinal

excitability occurs at high amplitude sensorimotor mu-oscillations. *Neuroimage:*

Reports, *2*(4), 100132. https://doi.org/10.1016/j.ynirp.2022.100132

Pascual-Leone, A., Grafman, J., & Hallett, M. (1994). Modulation of cortical motor output

maps during development of implicit and explicit knowledge. *Science*, *263*(5151),

1287–1289. https://doi.org/10.1126/science.8122113

Perez, M. A., Wise, S. P., Willingham, D. T., & Cohen, L. G. (2007). Neurophysiological

Mechanisms Involved in Transfer of Procedural Knowledge. *Journal of Neuroscience*,

27(5), 1045–1053. https://doi.org/10.1523/JNEUROSCI.4128-06.2007

Pinsard, B., Boutin, A., Gabitov, E., Lungu, O., Benali, H., & Doyon, J. (2019). Consolidation

alters motor sequence-specific distributed representations. *eLife*, *8*, e39324.

https://doi.org/10.7554/eLife.39324

Robertson, E. M. (2019). Skill Memory: Mind the Ever-Decreasing Gap for Offline

Processing. *Current Biology*, *29*(8), R287–R289.

https://doi.org/10.1016/j.cub.2019.03.007

Robertson, E. M., & Genzel, L. (2020). Memories replayed: Reactivating past successes and

new dilemmas. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *375*(1799), 20190226. https://doi.org/10.1098/rstb.2019.0226

Robertson, E. M., Pascual-Leone, A., & Miall, R. C. (2004). Current concepts in procedural

consolidation. *Nature Reviews Neuroscience*, *5*(7), 576–582.

https://doi.org/10.1038/nrn1426

Salmelin, R., & Hari, R. (1994). Spatiotemporal characteristics of sensorimotor

neuromagnetic rhythms related to thumb movement. *Neuroscience*, *60*(2), 537–550.

https://doi.org/10.1016/0306-4522(94)90263-1

Shirinpour, S., Alekseichuk, I., Mantell, K., & Opitz, A. (2020). Experimental evaluation of

methods for real-time EEG phase-specific transcranial magnetic stimulation. *Journal*

of Neural Engineering, *17*(4), 046002. https://doi.org/10.1088/1741-2552/ab9dba

Siegel, M., Warden, M. R., & Miller, E. K. (2009). Phase-dependent neuronal coding of

objects in short-term memory. *Proceedings of the National Academy of Sciences*,

106(50), 21341–21346. https://doi.org/10.1073/pnas.0908193106

Stefanou, M.-I., Desideri, D., Belardinelli, P., Zrenner, C., & Ziemann, U. (2018). Phase

Synchronicity of μ-Rhythm Determines Efficacy of Interhemispheric Communication

Between Human Motor Cortices. *The Journal of Neuroscience*, *38*(49), 10525–

10534. https://doi.org/10.1523/JNEUROSCI.1470-18.2018

Suresh, T., & Hussain, S. J. (2023). Re-evaluating the contribution of sensorimotor mu

rhythm phase and power to human corticospinal output: A replication study. *Brain*

Stimulation. https://doi.org/10.1016/j.brs.2023.05.022

Tunovic, S., Press, D. Z., & Robertson, E. M. (2014). A Physiological Signal That Prevents

Motor Skill Improvements during Consolidation. *Journal of Neuroscience*, *34*(15),

5302–5310. https://doi.org/10.1523/JNEUROSCI.3497-13.2014

VanRullen, R., Busch, N. A., Drewes, J., & Dubois, J. (2011). Ongoing EEG Phase as a

Trial-by-Trial Predictor of Perceptual and Attentional Variability. *Frontiers in*

Psychology, *2*, 60. https://doi.org/10.3389/fpsyg.2011.00060

Walker, M. P., Brakefield, T., Allan Hobson, J., & Stickgold, R. (2003). Dissociable stages of

human memory consolidation and reconsolidation. *Nature*, *425*(6958), 616–620.

https://doi.org/10.1038/nature01930

Watrous, A. J., Fell, J., Ekstrom, A. D., & Axmacher, N. (2015). More than spikes: Common

oscillatory mechanisms for content specific neural representations during perception

and memory. *Current Opinion in Neurobiology*, *31*, 33–39.

https://doi.org/10.1016/j.conb.2014.07.024

Wilkinson, L., & Jahanshahi, M. (2007). The striatum and probabilistic implicit sequence

learning. *Brain Research*, *1137*, 117–130.

https://doi.org/10.1016/j.brainres.2006.12.051

Wilkinson, L., Khan, Z., & Jahanshahi, M. (2009). The role of the basal ganglia and its

cortical connections in sequence learning: Evidence from implicit and explicit

sequence learning in Parkinson's disease. *Neuropsychologia*, *47*(12), 2564–2573.

https://doi.org/10.1016/j.neuropsychologia.2009.05.003

Wilkinson, L., & Shanks, D. R. (2004). Intentional Control and Implicit Sequence Learning.

Journal of Experimental Psychology: Learning, Memory, and Cognition, *30*(2), 354–

369. https://doi.org/10.1037/0278-7393.30.2.354

Wilkinson, L., Steel, A., Mooshagian, E., Zimmermann, T., Keisler, A., Lewis, J. D., &

Wassermann, E. M. (2015). Online feedback enhances early consolidation of motor

sequence learning and reverses recall deficit from transcranial stimulation of motor

cortex. *Cortex*, *71*, 134–147. https://doi.org/10.1016/j.cortex.2015.06.012

Wischnewski, M., Haigh, Z. J., Shirinpour, S., Alekseichuk, I., & Opitz, A. (2022). The phase

of sensorimotor mu and beta oscillations has the opposite effect on corticospinal

excitability. *Brain Stimulation: Basic, Translational, and Clinical Research in

Neuromodulation*, *15*(5), 1093–1100. https://doi.org/10.1016/j.brs.2022.08.005

Womelsdorf, T., Schoffelen, J.-M., Oostenveld, R., Singer, W., Desimone, R., Engel, A. K., &

Fries, P. (2007). Modulation of Neuronal Interactions Through Neuronal

Synchronization. *Science*, *316*(5831), 1609–1612.

https://doi.org/10.1126/science.1139597

Yokoi, A., & Diedrichsen, J. (2019). Neural Organization of Hierarchical Motor Sequence

Representations in the Human Neocortex. *Neuron*, *103*(6), 1178-1190.e7.

https://doi.org/10.1016/j.neuron.2019.06.017

Zrenner, C., Desideri, D., Belardinelli, P., & Ziemann, U. (2018). Real-time EEG-defined

excitability states determine efficacy of TMS-induced plasticity in human motor

cortex. *Brain Stimulation*, *11*(2), 374–389. https://doi.org/10.1016/j.brs.2017.11.016

Zrenner, C., Kozák, G., Schaworonkow, N., Metsomaa, J., Baur, D., Vetter, D., Blumberger,

D. M., Ziemann, U., & Belardinelli, P. (2023). Corticospinal excitability is highest at

the early rising phase of sensorimotor μ-rhythm. *NeuroImage*, *266*, 119805.

https://doi.org/10.1016/j.neuroimage.2022.119805

1000