**Title:** Motor sequence learning elicits mu peak-specific corticospinal plasticity

**Abbreviated title:** Sequence learning and mu phase-dependent corticospinal plasticity

**Authors:** Tharan Suresh[1], Fumiaki Iwane[2,3], Minsu Zhang[3], Margaret McElmurry[1], Muskan

Manesiya[1], Michael V Freedberg[1], Sara J Hussain[1]*

**Affiliations:** [1]Department of Kinesiology & Health Education, The University of Texas at

Austin. Austin, TX, USA. [2]Human Cortical Physiology and Neurorehabilitation Section,

National Institute of Neurological Disorders and Stroke, National Institutes of Health.

Bethesda, MD, USA. [3]Department of Electrical & Computer Engineering, The University of

Texas at Austin. Austin, TX, USA.

**\*Corresponding author:** sara.hussain@austin.utexas.edu

**Abstract**

Motor cortical (M1) transcranial magnetic stimulation (TMS) interventions increase

corticospinal output and improve motor learning when delivered during sensorimotor mu

rhythm trough but not peak phases, suggesting that the mechanisms supporting motor

learning may be most active during mu trough phases. Based on these findings, we

predicted that motor sequence learning-related corticospinal plasticity would be most evident

when measured during mu trough phases. Healthy adults were assigned to either a

sequence or no-sequence group. Participants in the sequence group practiced the implicit

serial reaction time task (SRTT), which contained an embedded, repeating 12-item

sequence. Participants in the no-sequence group practiced a version of the SRTT that

contained no sequence. We measured mu phase-independent and mu phase-dependent

MEP amplitudes using EEG-informed single-pulse TMS before, immediately after, and 30

minutes after the SRTT in both groups. All participants performed a retention test one hour

after SRTT acquisition. In both groups, mu phase-independent MEP amplitudes increased

following SRTT acquisition, but the pattern of mu phase-dependent MEP amplitude changes

after SRTT acquisition differed between groups. Relative to the no-sequence group, the

sequence group showed greater peak-specific MEP amplitude increases 30 minutes after

SRTT acquisition. Further, the magnitude of these peak-specific MEP amplitude increases

was negatively associated with the magnitude of sequence-specific learning. Contrary to our

original hypothesis, results revealed that motor sequence-specific learning elicits peak-

specific corticospinal plasticity. Findings provide first direct evidence for the presence of a

mu phase-dependent motor learning mechanism in the human brain.

*Keywords:* sensorimotor rhythms, motor learning, brain stimulation, phase, motor cortex

**New and Noteworthy**

Recent work suggests that motor learning's neural mechanisms may be most active during specific sensorimotor mu rhythm phases. If so, motor sequence learning-induced corticospinal plasticity should be more evident during some mu phases than others. Our results show that motor sequence-specific learning elicits corticospinal plasticity that is most prominent during mu peak phases. Further, this peak-specific plasticity correlates with learning. Findings establish the presence of a mu phase-dependent motor learning mechanism in the human brain.

**Introduction**

The sensorimotor cortex is a hub within the human motor skill acquisition network. Human sensorimotor cortical activity dynamically oscillates in the mu (8-13 Hz) frequency range, with distinct phases of this rhythm reflecting brief windows of relative excitation and inhibition (Bergmann et al., 2019; Hussain et al., 2019; Suresh & Hussain, 2023; Zrenner et al., 2018). In non-human primates, motor cortical (M1) neuronal spiking rates are highest during mu rhythm trough phases and lowest during mu rhythm peak phases (Haegens et al., 2011). In humans, corticospinal output (Bergmann et al., 2019; Hussain et al., 2019; Suresh & Hussain, 2023; Wischnewski et al., 2022; Zrenner et al., 2018) and interhemispheric communication between homologous M1 subregions (Stefanou et al., 2018) are also increased during sensorimotor mu rhythm trough relative to peak phases.

In addition to single-neuron spiking rates, corticospinal output, and inter-hemispheric communication, sensorimotor mu rhythm phase covaries with the likelihood of inducing long-term potentiation-like (LTP-like) corticospinal plasticity in humans. For example, transcranial magnetic stimulation (TMS) interventions applied to M1 during mu rhythm trough phases preferentially induce LTP-like corticospinal plasticity, measured as increases in corticospinal output (Baur et al., 2020; Zrenner et al., 2018). In contrast, identical TMS interventions applied during mu rhythm peak phases induce weak long-term depression-like (LTD-like) corticospinal plasticity, measured as decreases in corticospinal output (Baur et al., 2020; Zrenner et al., 2018). These findings suggest that corticospinal plasticity induction is causally related to sensorimotor mu rhythm phase, with LTP-like and weak LTD-like corticospinal plasticity occurring preferentially during mu rhythm trough and peak phases, respectively. Given that LTP-like plasticity processes have been proposed to support human motor learning (Cantarero et al., 2013a; Cantarero et al., 2013b), and TMS interventions preferentially enhance motor learning when applied during mu rhythm trough phases (Hussain et al., 2021), the neurophysiological mechanisms underlying human motor learning may thus be most active during sensorimotor mu rhythm trough phases.

Outside of the motor system, brain oscillatory activity rhythmically controls various

attentional (Busch & VanRullen, 2010; VanRullen et al., 2011), perceptual (Baumgarten et

al., 2015; Busch et al., 2009; Dugué et al., 2011), and memory processes (Kerrén et al.,

2018; Siegel et al., 2009). For example, the formation and retrieval of declarative memories

has been proposed to occur during opposing hippocampal oscillatory phases (Hasselmo et

al., 2002; Kerrén et al., 2018), object memory formation preferentially occurs during distinct

prefrontal oscillatory phases (Siegel et al., 2009), and replay of whole-brain neuronal activity

patterns associated with working memory are phase-locked to the theta rhythm (Fuentemilla

et al., 2010). These findings suggest that oscillatory phase-dependent memory processing

may be a widespread learning mechanism present across neural domains (Watrous et al.,

2015). If so, phase-dependent neurophysiological signatures of motor learning should also

be detectable within the human motor system.

Here, we reasoned that, if motor learning relies on a sensorimotor mu phase-dependent

mechanism, then learning-related corticospinal plasticity should be more evident during

some mu rhythm phases than others. Based on previous studies showing that corticospinal

output (Hussain et al., 2019; Suresh & Hussain, 2023; Zrenner et al., 2018),

interhemispheric communication (Stefanou et al., 2018), corticospinal plasticity (Baur et al.,

2020; Zrenner et al., 2018), and the effects of TMS interventions on motor sequence

learning (Hussain et al., 2021) are all enhanced during mu rhythm trough relative to peak

phases, we predicted that motor sequence learning-related changes in corticospinal output

would be more evident when measured during mu rhythm trough than peak phases.

To test this possibility, we measured sensorimotor mu phase-independent and phase-

dependent corticospinal output using single-pulse, real-time EEG-informed TMS immediately

before and after healthy adults practiced a serial reaction time task (SRTT) that included

either an embedded, repeating 12-item motor sequence (sequence group) or a variation of

the same task that included no sequence (no-sequence group). We predicted that mu

phase-dependent corticospinal plasticity would be more prominent for the sequence group

than the no-sequence group, with motor sequence-specific learning-related corticospinal

plasticity being stronger when measured during mu trough than peak phases. As expected,

our results show that participants in the sequence group exhibited greater sequence-specific

learning than those in the no-sequence group. While both groups showed increased mu

phase-independent corticospinal output following SRTT performance, each group showed a

distinct pattern of mu phase-dependent corticospinal plasticity. Relative to the no-sequence

group, the sequence group showed greater increases in *peak*-specific corticospinal output.

Further, peak-specific increases in corticospinal output were negatively associated with

sequence-specific learning. Surprisingly, these findings indicate that corticospinal plasticity

induced by motor sequence-specific learning is more prominent during mu rhythm peak than

trough phases. Findings establish the mechanistic role of mu rhythm phase in behaviorally-

relevant corticospinal plasticity within the human motor system.

**Methods**

*Participants*

Thirty-six naïve right-handed healthy adult participants (age = 20.9 ± 0.52, 25 F and 11 M)

participated in a single-session study that involved single-pulse transcranial magnetic

stimulation (TMS), electroencephalography (EEG), electromyography (EMG), and behavioral

testing. Right-handedness was confirmed using the Edinburgh Handedness Inventory

(Oldfield, 1971). This study was approved by the Institutional Review Board at the University

of Texas at Austin and all participants provided their written informed consent. Participants

were randomly assigned to either the sequence group (N = 18, age = 20 ± 0.37 years, 13 F

and 5 M) or the no-sequence group (N=18, age = 21.8 ± 0.94 years, 12 F and 6 M) prior to

beginning experimental procedures. This sample size was chosen based on previous

studies that identified significant mu phase-dependent variation in corticospinal output and

corticospinal plasticity, as well as phase-dependent enhancement of motor sequence

learning (N = 12-23, Zrenner et al., 2018; Bergmann et al., 2019; Hussain et al., 2019;

Suresh & Hussain, 2023; Hussain et al., 2021).

*Experimental design and timeline*

After EEG and EMG preparation, the scalp hotspot for the right first dorsal interosseous

muscle (FDI) and the stimulation intensity needed to elicit motor-evoked potentials (MEPs)

within this muscle were empirically determined for each participant. Participants then

performed a choice reaction time task (cRT), after which parameters for real-time targeting

of sensorimotor mu rhythm phases were individually optimized. Phase targeting accuracy

was assessed during EEG recordings while participants rested quietly with their eyes open.

Then, single-pulse TMS was delivered over the scalp hotspot during sensorimotor mu

rhythm peak, trough, and random phases. Participants subsequently practiced one of two

versions of the serial reaction time task (SRTT; see below for details). Immediately and 30

minutes after task completion, single-pulse TMS was again delivered during mu peak, trough

and random phases. One hour after completing the SRTT, both groups performed a brief

retention test that involved the same SRTT version performed during practice. After retention

testing, all participants completed the Process Dissociation Procedure (PDP; Wilkinson &

Shanks, 2004). Given that corticospinal plasticity following motor sequence learning under

implicit and explicit task conditions may differ (Tunovic et al., 2014), the PDP was used to

evaluate the development of explicit sequence awareness. Data from the cRT are not

reported in this manuscript. See Figure 1 for experimental design and timeline.

154
155
**Figure 1. Experimental timeline and serial reaction time task. A)** Experimental timeline.
Real-time EEG analysis algorithm parameters were individually optimized during resting
EEG recordings. After acquisition of phase-independent (random) and phase-dependent
(peak and trough) MEPs at baseline, participants in both groups performed the SRTT. Post-
SRTT phase-independent and phase-dependent MEPs were recorded immediately and 30
minutes after SRTT acquisition. SRTT retention was evaluated 1 hour after SRTT
acquisition. **B)** Depiction of implicit SRTT. During acquisition, participants completed 14
blocks (120 trials per block). During retention, participants completed 7 blocks (120 trials per
block). During sequence blocks, the sequence group practiced the SRTT with an embedded
12-item sequence, while during these blocks the no-sequence group practiced the task with
no sequence (all random blocks).
167

*Data acquisition*

<u>EEG and EMG recording.</u> 64-channel EEG and bipolar EMG signals were recorded using

TMS-compatible amplifiers (NeurOne Tesla with real-time digital out option, Bittium

Biosignals, Finland). EEG impedances were maintained below 10 kΩ. EMG signals were

recorded from the right FDI using disposable adhesive electrodes placed in a belly-tendon

montage. EEG and EMG were recorded at a sampling rate of 5 kHz, with low-pass hardware

filtering at 1250 Hz and a resolution of 0.001 µV.

Transcranial magnetic stimulation. We empirically determined the motor hotspot as the scalp

site at which suprathreshold, single-pulse TMS elicited a reliable, focal twitch within the right

FDI as well as a peak-to-peak motor evoked potential (MEP) amplitude of at least 50 µV.

After hotspot identification, resting motor threshold (RMT) was determined using an

automatic threshold-tracking algorithm (Borckardt et al., 2006). On average, RMT was 64.3 ±

3.8% of maximum stimulator output for the sequence group and 61.2 ± 3.1% for the no-

sequence group. All TMS procedures were performed using a figure-of-eight coil held at

~45° relative to the midsagittal line (DuoMAG XT-100, Deymed Diagnostic, biphasic pulse

shape). Coil position accuracy was monitored online using frameless neuronavigation

(BrainSight, Rogue Research) during hotspot identification, RMT determination, and mu

phase-independent and mu phase-dependent single-pulse TMS.

Real-time EEG analysis. Real-time EEG-triggered TMS was performed using an established

method (Zrenner et al., 2018; Shirinpour et al., 2020; Hussain et al., 2021; Suresh &

Hussain, 2023). The EEG system was configured to stream data from 5 pre-selected EEG

channels overlying the left sensorimotor cortex (C3, FC3, FC5, CP3 and CP5) using a local

wired network connection (packet rate = 1 kHz, 5 packets per sample) between the EEG

system and a dedicated PC for signal processing (Microsoft Windows 10, 10 cores, Intel i9

processor, 16 GB RAM) using Lab Streaming Layer software (Kothe, 2014) in real-time.

Custom MATLAB scripts were used to receive EEG data, process it, and send TTL triggers

to the TMS device upon detection of the desired mu rhythm phase. During data streaming,

EEG data was first downsampled to 1 kHz. To attenuate the effects of volume conduction,

the Hjorth transformation was applied by subtracting the mean of the surrounding electrodes

(FC3, FC5, CP3 and CP5) from the center electrode (C3; Hjorth 1975). The resulting signal

was then buffered to create overlapping 500 ms windows for real-time analysis. At each time

step, the current data window was zero-phase bandpass filtered using a Finite Impulse

Response (FIR) filter (8-13 Hz, order = 128) and 64 ms data segments at the beginning and

end of the 500 ms window were removed to eliminate edge effects. The remaining 372 ms

data window was then used to fit an autoregressive model (Yule-Walker method, order: 30)

that forward predicted the bandpass filtered signal 128 ms. After forward prediction, the

Hilbert transform was used to generate an estimated phase angle timeseries signal. The

phase estimation algorithm was also configured to trigger TMS only when each participant's

sensorimotor mu power exceeded a predefined threshold. To calculate this threshold, resting

state EEG data was divided into non-overlapping 500 ms segments, bandpass filtered using

a Finite Impulse Response (FIR) filter (8-13 Hz, order = 128), Hjorth transformed,

downsampled to 1 kHz, and then spectrally decomposed (Welch's method, 4-100 Hz with

0.25 Hz resolution excluding frequencies between 57.75 – 62 Hz to reduce the influence of

line noise). For each participant, the median of the resulting frequency-domain signal

between 8 and 13 Hz was defined as the power threshold. When the estimated

instantaneous phase angle at the time point of interest matched the pre-determined mu

phase condition and mu power exceeded each participant's threshold, a single TTL pulse

was sent to the TMS system and also marked the time of TMS delivery in the EEG

recording.

Phase targeting accuracy and real-time EEG-informed single-pulse TMS. Prior to delivering

real-time EEG-informed TMS, we first evaluated the accuracy of phase targeting in the

absence of TMS delivery in each participant. 25 TTL pulses were sent to the EEG system at

mu peak, trough and random phases, but no TMS was delivered (minimum inter-stimulus

interval = 5 s + random jitter). During mu phase-independent single-pulse TMS, blocks of 25

TMS pulses were delivered to the scalp hotspot for the right FDI during random mu phases

(120% RMT, inter-stimulus interval = 5 s + random jitter). During mu phase-dependent

single-pulse TMS, blocks of 25 TMS pulses were delivered during mu peak and trough

phases (120% RMT, minimum inter-stimulus interval = 5 s + random jitter). Real-time EEG-

informed single-pulse TMS was delivered at baseline, immediately after SRTT acquisition,

and 30 minutes after SRTT acquisition. The order of phases targeted per block (i.e., random,

peak, and trough) was counterbalanced across participants in both groups.

<u>Serial Reaction Time Task</u>. Participants in both the sequence and no-sequence groups were

seated in front of a computer monitor with their right arm resting on a tabletop and their right

hand positioned atop a serial response box (Chronos Response Device, Psychology

Software Tools, see Figure 1). Participants performed one of two versions (sequence or no-

sequence) of the SRTT (Nissen & Bullemer, 1987; Perez et al., 2007) programmed using E-

prime 3 (Psychology Software Tools). The no-sequence group was included in this study to

experimentally account for non-specific effects of motor task engagement (including

repeated keypresses) on MEP amplitude changes. During the SRTT, the monitor displayed

four equally-spaced green dashes against a black background. On each trial, one of these

dashes was replaced by a circle. Prior to beginning the task, participants were instructed to

respond to the presentation of the circle as fast and accurately as possible by pressing one

of the four keys with the corresponding finger of their right hand, as shown in Figure 1. When

the participant pressed the correct key, the circle disappeared, and the trial was complete. If

an incorrect key was pressed, the circle remained on the monitor and the trial continued until

the correct key was pressed. 500 ms elapsed between trials. No additional feedback on

keypress accuracy or keypress latency was provided.

Participants became familiarized with the task during a practice block that included 10 trials.

During SRTT acquisition, participants in the sequence group performed 14 blocks of

keypresses, with one minute of rest between blocks. Each block included 10 repetitions of a

12-item sequence, amounting to a total of 120 keypresses per block. Blocks 3-7 and 9-13

included a pre-determined, embedded 12-item sequence (2-3-1-4-3-2-4-1-3-4-2-1). Blocks 1,

2, 8 and 14 were random blocks that contained no embedded sequence. During SRTT

retention, participants in the sequence group performed 7 blocks of 120 keypresses each,

again resting for one minute between blocks. Blocks 1 and 7 were random blocks while

blocks 2-6 were sequence blocks.

Random blocks contained no consecutive item repeats and each item had approximately the

same frequency of appearance. Participants in the no-sequence group performed

randomized keypresses for all blocks. Participants were not informed about the presence or

absence of the sequence or its length.

<u>Process Dissociation Procedure.</u> After SRTT retention, participants completed the PDP. The

PDP assesses conscious control over sequence knowledge that may develop during implicit

SRTT performance (Destrebecqz et al., 2005; Wilkinson et al., 2009, 2015; Wilkinson &

Jahanshahi, 2007; Wilkinson & Shanks, 2004) and is based on the rationale that conscious

knowledge is both reportable upon request and under intentional control. The PDP consists

of an inclusion and exclusion test, each of which involve 12 unique three-item fragments of

the 12-item sequence that were embedded within the sequence version of the SRTT (e.g., 2-

3-1, 3-1-4, 1-4-3, etc.).

The inclusion test evaluates reportable, consciously accessible knowledge. During the

inclusion test, participants were presented with two elements of each of 12 three-item

fragments and were asked to produce the next element in the fragment by pressing the

corresponding button. The exclusion test evaluates intentional control. During the exclusion

test, participants were again presented with two elements of each of 12 three-item fragments

and were asked to *avoid* producing the next element in the fragment by pressing any button

*other than* the one that corresponded to the next element in the fragment. At the beginning

of the inclusion and exclusion tests, participants were informed that repeated element

sequences (i.e., a 2 immediately after a 2) were not allowed. For each trial of the inclusion

test, there was a 1/3 probability of pressing the correct key by chance. For each trial of the

exclusion test, there was a 2/3 probability of pressing the correct key by chance. The order

of inclusion and exclusion tests was counterbalanced across participants in both groups.

*Data analysis*

<u>Serial Reaction Time Task.</u> Reaction times (RTs) were averaged within each block. We

defined acquisition and retention skill as the difference between the mean RT of the last

random block and last sequence block during SRTT acquisition and the difference between

the mean RT of the first random block and first sequence block during SRTT retention,

respectively. We also defined offline changes in skill as the difference between acquisition

skill and retention skill. Here, positive values reflect offline losses while negative values

reflect offline gains.

$Acquisition\ skill = RT_{Acquisition\ block\ 14} - RT_{Acquisition\ block\ 13}$

$Retention\ skill = RT_{Retention\ block\ 1} - RT_{Retention\ block\ 2}$

$Offline\ skill\ changes = Acquisition\ skill - Retention\ skill$

<u>Process Dissociation Procedure.</u> Keypresses during inclusion trials were considered

accurate when the participant correctly completed the 3-element fragment by pressing the

correct key in the sequence. Keypresses during exclusion trials were considered accurate

when the participant correctly avoided the next element in the fragment by pressing one of

the two other key choices. The total number of accurate trials in the inclusion and exclusion

conditions were compared to the corresponding chance levels (i.e., 4/12 for the inclusion test

and 8/12 for the exclusion test) to test for explicit awareness of the embedded sequence.

<u>EEG analysis.</u> EEG signals were processed using a combination of Fieldtrip (Oostenveld et

al., 2010) and custom-written MATLAB scripts. To evaluate phase targeting accuracy, we

segmented EEG data into 6 s segments (± 3 s relative to the targeted phase), and then

demeaned and detrended these segments. To ensure that all offline data analysis

procedures mimicked those applied during real-time phase targeting, we applied the Hjorth

transformation to the EEG data by subtracting the mean of four surrounding sensorimotor

electrodes (FC3, FC5, CP3 and CP5) from the center sensorimotor electrode (C3). Resulting

data segments were zero-phase bandpass filtered using an FIR filter (8-13 Hz). The

instantaneous phase of the filtered signal at the time of phase targeting was identified using

the Hilbert transformation. We then combined phase angles across all participants per

group.

Given that mu phase-power interactions shape MEP amplitude (Hussain et al., 2019;

Ozdemir et al., 2022; Suresh & Hussain, 2023), we also calculated pre-stimulus

sensorimotor mu rhythm power. Continuous EEG data obtained during TMS delivery were

segmented into 500 ms segments (0.51 to 0.01 seconds prior to each TMS pulse),

demeaned, linearly detrended and Hjorth transformed (center = C3, surround = FC1, FC5,

CP1, CP5; Hjorth, 1975; Zrenner et al., 2018; Hussain et al., 2021; Suresh & Hussain,

2023). The resulting Hjorth-transformed signals were then spectrally decomposed (Welch's

method, 4-50 Hz with 0.25 Hz resolution, Suresh & Hussain, 2023). Each trial's power

values were averaged in the mu (8-13 Hz) range to generate a mixed mu power estimate

(i.e., the combination of aperiodic and periodic components of mu power; Donoghue et al.,

2020). Mixed mu power was calculated instead of periodic mu power as our previous work

demonstrated that statistical models including mu rhythm phase and mixed mu power

explain MEP amplitude variation better than those including mu phase and periodic mu

power (Suresh & Hussain, 2023).

MEP analysis. Pre-stimulus EMG activity was filtered with a discrete Fourier transform filter

to reduce line noise. All trials were visually inspected, and those that precluded reliable

calculation of peak-to-peak MEP amplitudes were rejected. We also calculated background

EMG as the root-mean-square of pre-stimulus EMG activity between 100 and 25 ms before

each TMS pulse; trials with values greater than 10 µV were rejected. Peak-to-peak MEP

amplitudes were calculated as the maximum voltage deflection between 20 and 40 ms after

each TMS pulse. Overall, 5.5% of all trials were rejected. MEP amplitudes were log-

transformed to reduce positive skew and improve statistical model fits (Hussain et al., 2019;

Suresh & Hussain, 2023).

*Statistical analysis*

Phase targeting accuracy. Phase angle distributions at the time of phase targeting were

evaluated for deviations from a circular uniform distribution in the 90° (peak phases) and

270° (trough phases) directions using separate V-tests for each group. For random phases,

general deviations from circular uniformity were tested using Rayleigh's test. For each group,

mu peak and trough phase angle distributions were compared using the Watson-Williams

test. Phase angle distributions for each condition (peak, trough, and random) were also

compared across groups using the Watson-Williams test.

SRTT acquisition and retention. To evaluate changes in mean RT during SRTT acquisition

and retention, we fit separate linear mixed-effects models (LMMs) for data acquired during

SRTT acquisition and retention periods. These included mean RT as the response variable,

BLOCK (categorical variable: acquisition blocks 1-14, retention blocks 1-7), GROUP

(categorical variable: sequence, no-sequence), and BLOCK x GROUP interactions as fixed

effects. PARTICIPANT was included as random intercept. To evaluate changes in skill from

SRTT acquisition to retention, we fit an LMM that included the change in mean RT as the

response variable, TASK (categorical variable: acquisition, retention, offline change),

GROUP (categorical variable: sequence, no-sequence), and TASK x GROUP interactions as

fixed effects. Post hoc pairwise comparisons were used to compare SRTT acquisition and

retention between groups. To evaluate offline skill, we compared offline changes to zero

using separate single-sample t-tests for the sequence and no-sequence groups.

Process dissociation procedure (PDP). To assess explicit sequence awareness, we

compared fragment completion accuracy to chance using separate single-sample right-tailed

t-tests for the inclusion and exclusion tests. One participant in the no-sequence group did not

complete the PDP.

MEP analysis. We first examined the influence of SRTT acquisition on phase-independent

MEP amplitudes using an LMM with GROUP (categorical variable: sequence, no-sequence),

TIME (categorical variable: baseline, immediately after SRTT acquisition, 30 minutes after

SRTT acquisition), and the GROUP x TIME interaction as fixed factors and natural log-

transformed POWER (continuous variable) as a covariate, and natural log-transformed MEP

amplitudes elicited during random mu phases as the response variable. The random

intercept of PARTICIPANT was also included (Hussain et al., 2019; Hussain et al., 2021;

Suresh & Hussain, 2023). Post hoc pairwise comparisons were used to compare phase-

independent MEP amplitudes across time points.

Next, we evaluated the influence of SRTT task performance on phase-dependent MEP

amplitudes by fitting an LMM that included fixed effects of GROUP (categorical variable:

sequence, no-sequence), PHASE (categorical variable: peak, trough), TIME (categorical

variable: baseline, immediately after SRTT acquisition, 30 minutes after SRTT acquisition),

as well as their two- and three-way interactions as fixed factors, the PHASE x natural log-

transformed POWER interaction as a covariate, and natural log-transformed MEP

amplitudes as the response variable. PARTICIPANT was included as the random intercept.

This LMM revealed a significant three-way interaction (see *Results*). To further characterize

this three-way interaction, we fit separate LMMs for each group. These post hoc LMMs

included PHASE (categorical variable: peak, trough), TIME (categorical variable: baseline,

30 minutes after SRTT acquisition), and the PHASE x TIME interaction as fixed factors, as

well as the PHASE x log-transformed POWER interaction as a covariate. Natural log-

transformed mu MEP amplitudes were the response variable, and PARTICIPANT was

included as the random intercept. Post hoc pairwise comparisons were used to compare

MEP amplitudes across time points and phases. Given the mu PHASE x POWER interaction

identified in our previous work and that of others (Bigoni et al., 2024; Hussain et al., 2019;

Ozdemir et al., 2022; Suresh & Hussain, 2023) and our primary interest in evaluating mu

phase-dependent MEP amplitude changes, we included the PHASE x natural log-

transformed POWER interaction as a covariate in all our LMMs to control for this known

interaction effect (Hussain et al., 2019; Suresh & Hussain, 2023).

Finally, we examined relationships between phase-dependent MEP amplitude changes and

SRTT performance changes. We first used the LMM that evaluated phase-dependent

changes in MEP amplitudes across groups to obtain trial wise peak- and trough-specific

model-fitted MEP amplitudes at baseline and 30 minutes after SRTT acquisition. These time

points were chosen because our phase-dependent MEP analysis revealed a significant

three-way GROUP x PHASE x TIME interaction across these two time points (see *Results*),

Then, we calculated the magnitude of phase-dependent MEP amplitude changes for each

participant by dividing the fitted mean phase-dependent MEP amplitudes 30 minutes after

SRTT acquisition by the fitted mean phase-dependent MEP amplitudes at baseline.

Separate LMMs were then used to evaluate relationships between (1) mu peak-specific MEP

changes and SRTT performance and (2) mu trough-specific MEP changes and SRTT

performance. Each LMM included GROUP (categorical variable: sequence, no-sequence),

TASK (categorical variable: acquisition, retention), MEP ratio (continuous variable), and their

two- and three-way interactions as fixed factors. Each LMM included also SKILL (i.e.,

change in mean RT during SRTT acquisition and retention, see *Data analysis*) as the

response variable and PARTICIPANT as the random intercept. Post hoc pairwise

comparisons were used to statistically evaluate relationships between peak-specific MEP

amplitude changes and SRTT performance for each group.

All statistical analyses were performed in R (version 4.3.1). Model fits were visually

inspected using histograms of residuals and quantile-quantile plots. The significance of fixed

effects was determined using likelihood ratio tests. All reported β-coefficients represent the

difference in estimates between groups and phases. Alpha was equal to 0.05 for all

comparisons and p-values were corrected using the Tukey's Honestly Significant Difference

test for multiple comparisons where appropriate. Data are expressed as mean ± standard

error of the mean.

**Results**

*Phase targeting accuracy*

Real-time EEG analysis accurately identified and targeted mu peak (Figure 2, target phase =

90°, observed mean phase for sequence group = 85.7° [95% confidence interval {CI} =

80.27°– 91.12°], V = 237.9, p < 0.001; observed mean phase for no-sequence group =

86.93° [95% CI = 80.77°– 93.10°], V = 217.52, p < 0.001) and mu trough phases (target

phase = 270°, observed mean phase for sequence group = 271.72° [95% CI = 266.17° –

277.25°], V = 56.01, p < 0.001; observed mean phase for no-sequence group = 269.66°

[95% CI = 263.99° – 275.31°], V = 45.71, p = 0.001). As expected, targeted phases differed

significantly between peak and trough conditions in both the sequence group (F = 1687.0, p

< 0.001) and the no-sequence group (F = 1495.8, p < 0.001). For random phases, we

observed a deviation from circular uniformity in the 20° direction in the sequence group (R =

5.56, p = 0.003, observed mean phase = 19.05° [95% CI = 343.15° – 54.95°]) and no

significant deviation from uniformity in the no-sequence group (R = 2.47, p = 0.08, observed

mean phase = 12.93° [95% CI = 311.2° – 74.66°]). However, we observed no significant

differences in phase targeting accuracy between the sequence and no-sequence groups for

any phase (peak: F = 0.11, p = 0.741; trough: F = 0.341, p = 0.559; random: F = 0.188, p =

0.665).

**Figure 2. Phase targeting accuracy. A, B)** Group-averaged raw and filtered (8-13 Hz) EEG
signals recorded over left sensorimotor cortex during evaluation of mu phase targeting
accuracy in the sequence (A) and no-sequence (B) groups. Black vertical lines reflect the
time point at which the real-time EEG analysis algorithm identified mu peak, trough, and
random phases. Shading represents mean ± 1 SEM. **C, D)** Circular histograms indicate the
phase angle of the sensorimotor mu (8-13 Hz) rhythm at the time of phase targeting for
peak, trough, and random phases in the sequence (C) and no-sequence (D) groups. **E, F)**
Scalp distribution of C3-Hjorth transformed and band-pass filtered EEG activity during mu
peak, trough and random phases at the time of phase targeting for each phase condition in
the sequence (E) and no-sequence (F) groups. Topographical plots reflect average scalp-
recorded EEG signals (filtered between 8-13 Hz) within a 5 ms window centered on the time
of phase targeting. Bolded channels indicate those used to calculate C3-Hjorth transformed
EEG data. Black asterisks reflect statistical significance.

*SRTT*

During SRTT acquisition, participants in both the sequence and no-sequence groups

decreased their mean RTs across blocks (Figure 3, main effect of BLOCK: F = 12.96 and p

< 0.001). RTs did not differ between groups for random blocks, but participants in the

sequence group tended to show faster mean RTs during the sequence blocks than the no-

sequence group during the corresponding blocks (BLOCK x GROUP interaction: F = 9.37

and p < 0.001; no main effect of GROUP: F = 0.63 and p = 0.42). However, post hoc

pairwise comparisons did not identify any significant differences in mean RTs between

groups during these blocks (i.e., blocks 3-7 or 9-13, see Figure 3). For both groups, mean

RTs were not statistically distinguishable between the first and last random blocks (blocks 1

vs. 14, $t < 0.165$, $p = 0.99$). For the sequence group, mean RTs were faster during the last

sequence block than the last random block (block 13 vs. 14, t ratio = -8.51, $p < 0.001$), but

this was not the case for the no-sequence group (t ratio = 0.16, $p = 0.99$). Finally, acquisition

skill was significantly greater in the sequence than no-sequence group (Figure 3, t ratio =

5.94, $p < 0.001$). Thus, as expected, participants in the sequence group experienced

sequence-specific learning.

During SRTT retention, participants in both groups decreased their mean RTs across blocks

(Figure 3, main effect of BLOCK: $F = 18.91$ and $p < 0.001$) and mean RTs were overall

faster for the sequence group than the no-sequence group (main effect of GROUP: $F = 6.04$,

$p = 0.01$). Mean RTs were not statistically distinguishable between groups during random

blocks but were significantly faster for the sequence group than the no-sequence group

during sequence blocks (GROUP x BLOCK interaction: $F = 21.14$, $p < 0.001$). Post hoc

pairwise comparisons revealed that mean RTs were faster for the sequence group than for

the no-sequence group during blocks 4-6 ($t > 3.59$, p values $< 0.03$). In addition, retention

skill was significantly greater in the sequence group than the no-sequence group (Figure 3, t

ratio = 5.17, $p < 0.001$). Thus, participants in the sequence group retained sequence-specific

skill obtained during SRTT acquisition. Finally, participants in the sequence group showed

significant offline losses ($t = 5.23$ $p < 0.0001$, mean = 98.17 ms [95% CI = 58.61 ms - 137.72

ms] while those in the no-sequence group did not ($t = -0.51$, $p = 0.616$, mean = -1.95 ms

[95% CI = -10.01 ms - 6.11ms]). However, direct comparison of the magnitude of offline

changes revealed no significant difference between groups (Figure 3, $t = 0.76$, $p = 0.97$).

**Figure 3**. **Serial reaction time task performance. A)** Participants in the sequence group showed sequence-specific decreases in mean RTs during SRTT acquisition and retention. Vertical dotted lines indicate random blocks during the SRTT (practice block not shown). Green asterisks reflect significant post hoc pairwise comparisons between blocks 13 and 14 during SRTT acquisition and blocks 1 and 2 during retention for the sequence group. Shading represents the mean ± 1 SEM. **B)** Sequence-specific skill acquisition and retention were greater in the sequence group than in the no-sequence group. Black asterisks reflect significant post hoc comparisons for acquisition, retention and offline skill changes between groups. Note that for offline changes, positive values reflect offline losses while negative values reflect offline gains. For boxplots, black lines represent median, white dots represent mean, black dots represent individual participants, and error bars represent upper and lower inter-quartile ranges.

*Explicit sequence awareness*

In the sequence group, mean fragment completion accuracies exceeded chance for the

inclusion test (6 ± 0.66 [SEM], t = 3.02, p = 0.007, chance = 4) but not the exclusion test

(7.72 ± 0.54, t = -0.51, p = 0.61, chance = 8). In the no-sequence group, fragment

completion accuracies did not exceed chance for either the inclusion or exclusion tests

(Figure 4, inclusion = 3.88 ± 0.45, exclusion = 7.7 ± 0.54, t < -0.52 for both, p > 0.59 for

both). Given that explicit awareness requires both conscious knowledge of the sequence

and the ability to intentionally control sequence production, there was no strong evidence of

explicit sequence awareness in either group.

**Figure 4. Process Dissociation Procedure.** Mean number of fragments completed or avoided during the inclusion and exclusion portions of the PDP, respectively. The dashed lines reflect chance performance (4/12 for inclusion and 8/12 for exclusion). Black asterisks reflect significant difference between mean fragment completion accuracy and inclusion chance level in the sequence group. For boxplots, black lines represent group median, white dots represent group mean, black dots represent individual participants, and error bars represent upper and lower inter-quartile ranges.

*Phase-independent changes in MEP amplitudes*

We first characterized the effects of SRTT acquisition on phase-independent MEP amplitudes elicited during random mu phases. In both groups, MEP amplitudes increased following SRTT acquisition (significant main effect of TIME; F = 30.01, p < 0.001, no main effect of GROUP; F = 0.68, p = 0.41, no GROUP x TIME interaction, F = 0.88, p = 0.41). Post hoc pairwise comparisons revealed that random, phase-independent MEP amplitudes increased from baseline to immediately after SRTT acquisition (β = -0.08, t ratio = -2.38, p = 0.04), from baseline to 30 minutes after SRTT acquisition (β = -0.25, t ratio = -7.58, p < 0.0001), and from immediately to 30 minutes after SRTT acquisition (β = -0.17, t ratio = -5.21, p < 0.001). Thus, phase-independent MEP amplitudes increased similarly following SRTT acquisition in both groups.

*Phase-dependent changes in MEP amplitudes*

We next evaluated the effects of SRTT acquisition on phase-dependent MEP amplitudes elicited during mu peak and trough phases. We found a significant three-way GROUP x

TIME x PHASE interaction (Figure 5, likelihood ratio test: F = 4.95, p = 0.007). However, this

three-way interaction was only present when comparing changes in MEP amplitudes from

baseline to 30 minutes after SRTT acquisition (no GROUP x TIME x PHASE interaction for

baseline versus immediately after SRTT: t = -0.9, p = 0.36, significant GROUP x TIME x

PHASE interaction for baseline versus 30 minutes after SRTT: t = 2.19, p = 0.02), indicating

that the magnitude of MEP amplitude changes from baseline to 30 minutes after SRTT

acquisition more strongly differed across mu phases in one group than the other. Given the

significant three-way GROUP x PHASE x TIME interaction present from baseline to 30

minutes after SRTT acquisition, we evaluated mu phase-dependent changes in MEP

amplitudes across these two time points separately for each group. For the sequence group,

this post hoc analysis revealed a significant effect of PHASE (F = 5.32, p = 0.02) and TIME

(F = 68.59, p < 0.0001), but no PHASE x TIME interaction (F = 1.29, p = 0.25). Additional

post hoc pairwise comparisons further revealed that MEP amplitudes were significantly

larger during mu trough than peak phases (β = -0.08, t ratio = -2.35, p = 0.01), while MEP

amplitudes elicited during both phases were significantly larger 30 minutes after SRTT

acquisition than baseline (β = -0.29, t ratio = -8.28, p < 0.0001). For the no-sequence group,

post hoc analysis revealed a significant effect of PHASE (F = 24.9, p < 0.0001), TIME (F =

104.89, p < 0.0001) and a significant PHASE x TIME interaction (F = 4.06, p = 0.04).

Additional post hoc pairwise comparisons showed no differences between peak and trough

MEP amplitudes at baseline (β = -0.08, t ratio = -1.68, p= 0.33), but MEP amplitudes were

larger during trough than peak phases 30 minutes after SRTT acquisition (β = -0.22, t ratio =

-4.48, p < 0.0001).

**Figure 5. Mu phase-independent and phase-dependent changes in MEP amplitudes.**
**A)** Group-level linear mixed-effects model coefficients depicting relationships between mu phase-independent MEP amplitudes and group at baseline, immediately and 30 minutes after SRTT acquisition. **B)** Group-level linear mixed-effects model coefficients depicting relationships between mu phase-dependent MEP amplitudes and group at baseline, immediately, and 30 minutes after SRTT acquisition. For A and B, dots represent model coefficients and error bars indicate SEM. For A, black asterisks reflect significant post hoc pairwise comparisons across time points. For B, text above each panel indicates results of the PHASE x TIME interaction, and the black asterisk reflects significant GROUP x PHASE x TIME interaction for the no-sequence group.

To visualize differences in mu phase-dependent MEP amplitude changes between groups

(i.e., in the sequence group relative to the no-sequence group) identified by the GROUP x

PHASE x TIME interaction, we subtracted no-sequence group model-fitted MEP amplitudes

from sequence group model-fitted MEP amplitudes (Figure 6). This subtraction illustrated

that sequence-specific learning-related MEP amplitude changes were significantly greater

during mu rhythm peak than trough phases 30 minutes after SRTT acquisition.

**Figure 6. Effects of motor sequence-specific learning on mu phase-dependent MEP amplitude changes.** At each time point, group-level model fitted MEP amplitudes obtained during mu peak and trough phases in the no-sequence group were subtracted from corresponding group-level model fitted MEP amplitudes in the sequence group. Black asterisk reflects significant GROUP x PHASE x TIME interaction.

*Relationships between phase-dependent MEP amplitude changes and SRTT performance*

Finally*,* we examined relationships between mu phase-dependent MEP amplitude changes and SRTT performance. When evaluating relationships between peak-specific MEP amplitude changes and SRTT performance, there was a significant effect of GROUP (F = 9.07, p = .005), MEP ratio (F = 8.32, p = .006), and a significant GROUP x MEP ratio interaction (F = 8.82, p = .005), but no main effect of TASK (F = 0.06, p = .8) nor a GROUP x MEP ratio x TASK interaction (F = 0.01, p = .9). The GROUP x MEP ratio interaction was driven by a significantly stronger negative relationship between peak-specific MEP amplitude changes and SRTT performance for the sequence group relative to the no-sequence group (Figure 7A). In contrast, when evaluating relationships between trough-specific MEP amplitude changes and SRTT performance, there was no main effect of GROUP (F = 1.69, p = .2), MEP ratio (F = 1.7, p = .2), or TASK (F = 1.86, p = .18), nor a GROUP x MEP ratio interaction (F = 1.58, p = .21) or a GROUP x MEP ratio x TASK interaction (F = 0.55, p = .46). That is, the relationship between trough-specific MEP amplitude changes and SRTT performance did not significantly differ between groups (Figure 7B). Overall, these findings demonstrate that peak- but not trough-specific MEP amplitude changes were significantly

609 more negatively associated with SRTT performance in the sequence group than in the no-

610 sequence group.

**Figure 7. Relationships between mu phase-dependent MEP amplitude changes and**
**SRTT performance. (A)** Relationship between mu peak-specific MEP amplitude changes
and SRTT acquisition skill (left) and retention skill (right). **(B)** Relationships between trough-
specific MEP amplitude changes and SRTT acquisition skill (left) and retention skill (right).
Phase-specific MEP amplitude changes are represented as the ratio between model-fitted
phase-specific MEPs obtained 30 minutes after SRTT acquisition and those obtained at
baseline. Larger MEP ratio values indicate greater peak-specific (A) or trough-specific (B)
MEP amplitude changes. For both A and B, larger skill values indicate greater reaction time
improvements during SRTT acquisition (left) and retention (right). Dots represent individual
participant data and lines represent the best fit line. For A, black asterisks reflect significant
GROUP x MEP ratio interaction.

**Discussion**

Based on our previous work showing that mu phase-dependent TMS interventions coupled

to sensorimotor mu rhythm trough but not peak phases improve motor sequence learning,

we predicted that motor sequence learning-related corticospinal plasticity would be more

evident when measured during sensorimotor mu rhythm trough than peak phases. To test

this hypothesis, we measured MEP amplitudes during mu rhythm peak, trough, and random

phases immediately before and after healthy adults performed a version of the implicit SRTT

that included either an embedded, repeating motor sequence (sequence group) or no

sequence (no-sequence group). Behaviorally, participants in the sequence group acquired

greater sequence-specific knowledge than participants in the no-sequence group and

retained this knowledge over time. Physiologically, participants in both groups showed mu

phase-independent increases in corticospinal output following SRTT acquisition, reflecting

the occurrence of significant mu phase-independent corticospinal plasticity that did not differ

between participants who did and did not acquire sequence-specific knowledge. However,

we identified differences in mu phase-dependent changes in corticospinal output between

groups. In the sequence group, SRTT acquisition elicited corticospinal plasticity that was

similar in magnitude across mu peak and trough phases. In contrast, in the no-sequence

group, SRTT acquisition elicited corticospinal plasticity that was greater during mu trough

than peak phases. Direct comparisons between groups showed that participants in the

sequence group exhibited greater peak-specific corticospinal plasticity than those in the no-

sequence group. Further, the magnitude of peak- but not trough-specific corticospinal

plasticity was significantly more negatively associated with the magnitude of sequence-

specific learning for participants in the sequence relative to the no-sequence group. These

findings strongly suggest that performance of each SRTT variant elicits phase-independent

corticospinal plasticity, while sequence-specific learning elicits stronger corticospinal

plasticity during mu rhythm peak than trough phases.

Motor learning potentiates corticospinal output (Pascual-Leone et al., 1994; Perez et al.,

2007) and occludes subsequent induction of LTP-like corticospinal plasticity (Cantarero et

al., 2013a; Cantarero et al., 2013b). This occlusion is thought to support retention of newly-

acquired motor skills, such that its experimental reversal significantly diminishes retention

(Cantarero et al., 2013a). Thus, LTP-like corticospinal plasticity and its occlusion both play

key mechanistic roles in motor learning. Like motor learning, repetitive TMS interventions

can also induce corticospinal plasticity. Specifically, recent studies using real-time EEG-TMS

have shown that both low- and high-frequency repetitive TMS interventions preferentially

induce LTP-like corticospinal plasticity when applied during sensorimotor mu trough but not

peak phases (Baur et al., 2020; Zrenner et al., 2018). Such mu phase-dependent

corticospinal plasticity could therefore reflect a common neurophysiological mechanism

underlying the effects of repetitive TMS interventions and motor learning. If mu phase-

dependent corticospinal plasticity does support motor learning, then TMS interventions

should more strongly affect motor learning when delivered during specific mu phases. We

previously evaluated this possibility by measuring the effects of a mu phase-dependent

repetitive TMS intervention on offline changes in explicit motor sequence task performance

(Hussain et al., 2021). In that study, a repetitive TMS intervention delivered during mu trough

but not peak phases enhanced offline gains in motor sequence performance. These findings

raised the intriguing possibility that the neurophysiological mechanisms underlying motor

learning may be most active and thus most susceptible to modification during sensorimotor

mu trough phases. However, in the current study, sequence-specific learning elicited mu

phase-dependent corticospinal plasticity that was stronger during mu peak than trough

phases.

What could explain these surprising findings? Our experimental design directly compared

changes in MEP amplitudes following performance of one of two related motor tasks. In the

sequence group, participants performed repeated keypresses while also acquiring

knowledge of an embedded sequence, while in the no-sequence group, participants

performed repeated keypresses in the absence of any embedded sequence. To isolate the

effects of sequence learning on mu phase-dependent corticospinal plasticity, we therefore

subtracted no-sequence group model-fitted MEP amplitudes from sequence group model-

fitted MEP amplitudes. This direct contrast revealed that sequence learning, after accounting

for the effects of repeated keypresses, induced mu phase-dependent corticospinal plasticity

that was stronger during mu peak than trough phases. Consistent with this finding, we also

observed a significant negative relationship between peak-specific corticospinal plasticity

and sequence-specific learning. We previously proposed that mu rhythm peak phases reflect

a neutral excitability state that is not strongly affected by synchronization of corticothalamic

inputs to layer V pyramidal cells (Hussain et al., 2019). Specifically, M1 single neuron spiking

rates are lower during mu rhythm peak than trough phases in non-human primates

(Haegens et al., 2011), which may correspond to the decreased corticospinal output

observed during mu peak relative to trough phases in humans (Bergmann et al., 2019;

Hussain et al., 2019; Suresh & Hussain, 2023). Indeed, Bergmann et al. (2019) showed that

mu peak phases reflect a lack of pulsed excitation rather than pulsed inhibition of

corticospinal output. Together, these previous findings suggest that mu peaks reflect brief

quiescent windows during which corticospinal and sensorimotor network communication is

diminished relative to mu troughs. As a result, neural activity should only weakly propagate

through an extended sensorimotor network during mu peak phases. We and others

previously provided empirical support for this possibility by showing that (1) TMS-related

whole-brain broadband power, an indirect marker of population-level neuronal spiking (Gao

et al., 2017; Manning et al., 2009), is weaker when M1 TMS delivery coincides with mu

rhythm peak than trough phases (Hussain et al., 2021), and (2) interhemispheric

communication between bilateral motor cortices is lower during mu peak than trough phases

(Stefanou et al., 2018). The decreased network communication that occurs during mu peak

phases may prevent motor sequence memories from being disrupted by task-irrelevant

sensory information carried by thalamocortical projections, such that sequence memories

can be replayed and consolidated by sensorimotor and hippocampal networks without

interference (Buch et al., 2021; Jacobacci et al., 2020; Robertson & Genzel, 2020). This idea

aligns with previous work showing that wakeful rest after motor sequence learning supports

consolidation and retention of the learned sequence (Bönstrup et al., 2019, 2020; Buch et

al., 2021; Robertson, 2019; Robertson et al., 2004; Walker et al., 2003). Mu peak phases

could therefore provide a protected time window during which replay and consolidation of

motor sequence memories may occur. Such a scenario could explain why motor sequence-

specific learning-related corticospinal plasticity was more pronounced during mu peak

phases in the current study. Further, we also observed a negative relationship between

peak-specific MEP amplitude changes and SRTT performance that was significantly

stronger in the sequence group than the no-sequence group. Although this result may at first

appear counterintuitive, it is possible that poorer motor sequence-specific learning more

strongly recruits peak-specific corticospinal plasticity processes in order to protect newly-

formed yet fragile sequence memories from interference during consolidation. Thus, peak-

specific corticospinal plasticity changes could reflect a compensatory, protective mechanism

that is most prominent in participants who initially experience the least sequence-specific

learning. Future work could directly test the causal nature of the correlative relationship

observed in the current study by evaluating the behavioral consequences of experimentally

altering mu peak phase properties immediately after motor sequence learning using either

neurostimulation or closed-loop neurofeedback paradigms.

Although we previously suggested that the neurophysiological mechanisms underlying motor

sequence learning may be most active during sensorimotor mu trough phases (Hussain et

al., 2021), here we report that corticospinal plasticity following motor sequence-specific

learning is more strongly expressed during mu peak than trough phases. Two key

differences between our previous work and the current study could explain these diverging

results. First, the current study used the implicit SRTT, while our previous work used the

explicit motor sequence learning task. Although both tasks involve acquisition of sequential

knowledge, they rely on dissimilar sensorimotor networks. Specifically, the implicit SRTT

likely more heavily recruits corticostriatal circuits (Grafton et al., 1995; Hazeltine et al., 1997),

while the explicit motor sequence learning task may more strongly engage prefrontal and

premotor areas (Pinsard et al., 2019; Yokoi & Diedrichsen, 2019). Given the fundamental

role of oscillatory phase in inter-regional neural communication (Fries, 2005, 2015;

Womelsdorf et al., 2007), these two brain networks may optimally interact with sensorimotor

cortical neuronal populations during different mu rhythm phases. Second, the current study

directly assessed within-session phase-dependent neurophysiological changes induced by

motor sequence-specific learning, whereas our previous work examined the influence of a

phase-dependent TMS intervention on between-session behavioral changes in motor

sequence performance. Thus, corticospinal plasticity supporting motor sequence-specific

learning may be most evident during mu peak phases, while TMS interventions may best

improve motor sequence learning during mu trough phases. Along these lines, it is possible

that the neurophysiological mechanisms which protect newly-formed motor sequence

memories from interference during consolidation are maximally active during mu peak

phases, such that peak-specific TMS interventions cannot further upregulate them. If so, the

same protective mechanisms may be only partially active during mu trough phases, allowing

them to be further enhanced by trough-specific TMS interventions. Regardless, our current

findings and previous work both support the notion that the neurophysiological mechanisms

underlying motor sequence learning are sensorimotor mu phase-dependent.

Several strengths of this study exist. First, our experimental design included two groups: one

that experienced sequence learning and one that did not. This design allowed us to

dissociate the effects of sequence-specific and non-sequence-specific learning on mu

phase-independent and mu phase-dependent corticospinal plasticity. Of note, non-specific

effects of motor task engagement are often not considered in motor sequence learning

studies (Bönstrup et al., 2019, 2020; Perez et al., 2007; Walker et al., 2003), including our

own previous work (Hussain et al., 2021). However, without directly contrasting phase-

dependent corticospinal plasticity between the sequence and no-sequence groups, the

current study would have failed to identify any role of mu phase-dependent corticospinal

plasticity in sequence learning. Second, we quantified both mu phase-independent and

phase-dependent corticospinal plasticity. This quantification showed that mu phase-

independent corticospinal plasticity can be elicited by repeated keypresses even in the

absence of sequence learning itself. Third, participants were not required to meet any pre-

defined neurophysiological criteria to participate in this study; our results are likely to be

more generalizable across diverse groups of participants than previous studies using similar

techniques (Baur et al., 2020; Zrenner et al., 2018). Finally, rather than inferring the

existence of a phase-dependent learning mechanism from an interventional study (Hussain

et al., 2021), here we specifically quantified phase-dependent neurophysiological changes

induced by sequence-specific and non-sequence-specific learning. Thus, the results of the

current study can be directly attributed to motor sequence-specific learning rather than the

combined effects of motor sequence learning and a phase-dependent TMS intervention.

While our study provides novel insights into phase-dependent mechanisms of motor

learning, it is not without limitations. First, previous studies have shown that mu rhythm

phase and power interdependently influence corticospinal output at rest (Hussain et al.,

2019; Ozdemir et al., 2022; Suresh & Hussain, 2023), but here we evaluated only mu phase-

dependent changes in corticospinal output. Although this was achieved by introducing a

covariate into our statistical models that controlled for mu phase-power interactions, this

approach does not capture more complex learning-related changes in mu phase-power

interactions. We also did not measure changes in corticospinal output during other mu

phases (Wischnewski et al., 2022; Zrenner et al., 2023), other sensorimotor rhythms

(Hussain et al., 2019; Wischnewski et al., 2022) or broadband activity (Davis et al., 2022).

Experimental designs using phase-independent measures of corticospinal excitability and

post hoc trial sorting approaches may be better suited to explore the contribution of other

brain rhythms and phases to learning-related corticospinal plasticity.

In conclusion, we report that motor sequence-specific learning elicits mu peak-specific

corticospinal plasticity, and that the magnitude of this peak-specific corticospinal plasticity is

significantly negatively associated with sequence-specific learning. These findings provide

first direct evidence that motor learning recruits mu phase-dependent neurophysiological

mechanisms in humans and establish the mechanistic role of sensorimotor mu rhythm phase

in behaviorally-relevant corticospinal plasticity.

**Data availability:** Data used for analysis are available from the corresponding author upon

reasonable request.

**Acknowledgements**: We thank Kristen Pulliam for assistance with data collection.

**Grants**: SJH was supported by K12HD093427, R21NS133605, and R03HD114188 and TS

was supported by R25HD105583 and R21NS133605.

**Disclosures**: The authors declare no conflicts of interest, financial or otherwise, to disclose.

**Author contributions**: TS, FI, MZ, MVF and SJH conceived and designed research; TS,

MMc, and MMa performed experiments; TS and SJH analyzed data; TS, MVF, and SJH

interpreted results of experiments; TS and SJH prepared figures; TS and SJH drafted

manuscript; TS, MVF, and SJH edited and revised manuscript, TS, FI, MZ, MMc, MMa, MVF

and SJH approved final version of manuscript.

**References**

Baumgarten, T. J., Schnitzler, A., & Lange, J. (2015). Beta oscillations define discrete

   perceptual cycles in the somatosensory domain. *Proceedings of the National*

   *Academy of Sciences*, *112*(39), 12187–12192.

   https://doi.org/10.1073/pnas.1501438112

Baur, D., Galevska, D., Hussain, S., Cohen, L. G., Ziemann, U., & Zrenner, C. (2020).

   Induction of LTD-like corticospinal plasticity by low-frequency rTMS depends on pre-

   stimulus phase of sensorimotor μ-rhythm. *Brain Stimulation*, *13*(6), 1580–1587.

   https://doi.org/10.1016/j.brs.2020.09.005

Bergmann, T. O., Lieb, A., Zrenner, C., & Ziemann, U. (2019). Pulsed Facilitation of

   Corticospinal Excitability by the Sensorimotor μ-Alpha Rhythm. *Journal of*

   *Neuroscience*, *39*(50), 10034–10043. https://doi.org/10.1523/JNEUROSCI.1730-

   19.2019

Bigoni, C., Pagnamenta, S., Cadic-Melchior, A., Bevilacqua, M., Harquel, S., Raffin, E., &

   Hummel, F. C. (2024). MEP and TEP features variability: Is it just the brain-state?

   *Journal of Neural Engineering*. https://doi.org/10.1088/1741-2552/ad1dc2

Bönstrup, M., Iturrate, I., Hebart, M. N., Censor, N., & Cohen, L. G. (2020). Mechanisms of

   offline motor learning at a microscale of seconds in large-scale crowdsourced data.

   *Npj Science of Learning*, *5*(1), Article 1. https://doi.org/10.1038/s41539-020-0066-9

Bönstrup, M., Iturrate, I., Thompson, R., Cruciani, G., Censor, N., & Cohen, L. G. (2019). A

   Rapid Form of Offline Consolidation in Skill Learning. *Current Biology*, *29*(8), 1346-

   1351.e4. https://doi.org/10.1016/j.cub.2019.02.049

Borckardt, J. J., Nahas, Z., Koola, J., & George, M. S. (2006). Estimating Resting Motor

   Thresholds in Transcranial Magnetic Stimulation Research and Practice: A computer

   Simulation Evaluation of Best Methods. *The Journal of ECT*, *22*(3), 169.

   https://doi.org/10.1097/01.yct.0000235923.52741.72

Buch, E. R., Claudino, L., Quentin, R., Bönstrup, M., & Cohen, L. G. (2021). Consolidation of

human skill linked to waking hippocampo-neocortical replay. *Cell Reports*, *35*(10).

https://doi.org/10.1016/j.celrep.2021.109193

Busch, N. A., Dubois, J., & VanRullen, R. (2009). The Phase of Ongoing EEG Oscillations

Predicts Visual Perception. *Journal of Neuroscience*, *29*(24), 7869–7876.

https://doi.org/10.1523/JNEUROSCI.0113-09.2009

Busch, N. A., & VanRullen, R. (2010). Spontaneous EEG oscillations reveal periodic

sampling of visual attention. *Proceedings of the National Academy of Sciences*,

*107*(37), 16048–16053. https://doi.org/10.1073/pnas.1004801107

Cantarero, G., Lloyd, A., & Celnik, P. (2013). Reversal of Long-Term Potentiation-Like

Plasticity Processes after Motor Learning Disrupts Skill Retention. *Journal of*

*Neuroscience*, *33*(31), 12862–12869. https://doi.org/10.1523/JNEUROSCI.1399-

13.2013

Cantarero, G., Tang, B., O'Malley, R., Salas, R., & Celnik, P. (2013). Motor Learning

Interference Is Proportional to Occlusion of LTP-Like Plasticity. *Journal of*

*Neuroscience*, *33*(11), 4634–4641. https://doi.org/10.1523/JNEUROSCI.4706-

12.2013

Davis, Z. W., Muller, L., & Reynolds, J. H. (2022). Spontaneous Spiking Is Governed by

Broadband Fluctuations. *The Journal of Neuroscience*, *42*(26), 5159–5172.

https://doi.org/10.1523/JNEUROSCI.1899-21.2022

Destrebecqz, A., Peigneux, P., Laureys, S., Degueldre, C., Fiore, G. D., Aerts, J., Luxen, A.,

Linden, M. V. D., Cleeremans, A., & Maquet, P. (2005). The neural correlates of

implicit and explicit sequence learning: Interacting networks revealed by the process

dissociation procedure. *Learning & Memory*, *12*(5), 480–490.

https://doi.org/10.1101/lm.95605

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara,

A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing

neural power spectra into periodic and aperiodic components. *Nature Neuroscience*,

*23*(12), Article 12. https://doi.org/10.1038/s41593-020-00744-x

Dugué, L., Marque, P., & VanRullen, R. (2011). The Phase of Ongoing Oscillations Mediates

the Causal Relation between Brain Excitation and Visual Perception. *Journal of*

*Neuroscience*, *31*(33), 11889–11893. https://doi.org/10.1523/JNEUROSCI.1161-

11.2011

Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through

neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480.

https://doi.org/10.1016/j.tics.2005.08.011

Fries, P. (2015). Rhythms for Cognition: Communication through Coherence. *Neuron*, *88*(1),

220–235. https://doi.org/10.1016/j.neuron.2015.09.034

Fuentemilla, L., Penny, W. D., Cashdollar, N., Bunzeck, N., & Düzel, E. (2010). Theta-

Coupled Periodic Replay in Working Memory. *Current Biology*, *20*(7), 606–612.

https://doi.org/10.1016/j.cub.2010.01.057

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance

from field potentials. *NeuroImage*, *158*, 70–78.

https://doi.org/10.1016/j.neuroimage.2017.06.078

Grafton, S. T., Hazeltine, E., & Ivry, R. (1995). Functional Mapping of Sequence Learning in

Normal Humans. *Journal of Cognitive Neuroscience*, *7*(4), 497–510.

https://doi.org/10.1162/jocn.1995.7.4.497

Haegens, S., Nácher, V., Luna, R., Romo, R., & Jensen, O. (2011). α-Oscillations in the

monkey sensorimotor network influence discrimination performance by rhythmical

inhibition of neuronal spiking. *Proceedings of the National Academy of Sciences*,

*108*(48), 19377–19382. https://doi.org/10.1073/pnas.1117190108

Hasselmo, M. E., Bodelón, C., & Wyble, B. P. (2002). A Proposed Function for Hippocampal

Theta Rhythm: Separate Phases of Encoding and Retrieval Enhance Reversal of

Prior Learning. *Neural Computation*, *14*(4), 793–817.

https://doi.org/10.1162/089976602317318965

Hazeltine, E., Grafton, S. T., & Ivry, R. (1997). Attention and stimulus characteristics

determine the locus of motor-sequence encoding. A PET study. *Brain*, *120*(1), 123–

140. https://doi.org/10.1093/brain/120.1.123

Hjorth, B. (1975). An on-line transformation of EEG scalp potentials into orthogonal source

derivations. *Electroencephalography and Clinical Neurophysiology*, *39*(5), 526–530.

https://doi.org/10.1016/0013-4694(75)90056-5

Hussain, S. J., Claudino, L., Bönstrup, M., Norato, G., Cruciani, G., Thompson, R., Zrenner,

C., Ziemann, U., Buch, E., & Cohen, L. G. (2019). Sensorimotor Oscillatory Phase–

Power Interaction Gates Resting Human Corticospinal Output. *Cerebral Cortex*,

*29*(9), 3766–3777. https://doi.org/10.1093/cercor/bhy255

Hussain, S. J., Cohen, L. G., & Bönstrup, M. (2019). Beta rhythm events predict corticospinal

motor output. *Scientific Reports*, *9*(1), 18305. https://doi.org/10.1038/s41598-019-

54706-w

Hussain, S. J., Vollmer, M. K., Stimely, J., Norato, G., Zrenner, C., Ziemann, U., Buch, E. R.,

& Cohen, L. G. (2021). Phase-dependent offline enhancement of human motor

memory. *Brain Stimulation*, *14*(4), 873–883. https://doi.org/10.1016/j.brs.2021.05.009

Jacobacci, F., Armony, J. L., Yeffal, A., Lerner, G., Amaro, E., Jovicich, J., Doyon, J., &

Della-Maggiore, V. (2020). Rapid hippocampal plasticity supports motor sequence

learning. *Proceedings of the National Academy of Sciences*, *117*(38), 23898–23903.

https://doi.org/10.1073/pnas.2009576117

Kerrén, C., Linde-Domingo, J., Hanslmayr, S., & Wimber, M. (2018). An Optimal Oscillatory

Phase for Pattern Reactivation during Memory Retrieval. *Current Biology*, *28*(21),

3383-3392.e6. https://doi.org/10.1016/j.cub.2018.08.065

Kothe, C. (2014). *Lab streaming layer (LSL)*.

Manning, J. R., Jacobs, J., Fried, I., & Kahana, M. J. (2009). Broadband Shifts in Local Field

Potential Power Spectra Are Correlated with Single-Neuron Spiking in Humans.

*Journal of Neuroscience*, *29*(43), 13613–13620.

https://doi.org/10.1523/JNEUROSCI.2041-09.2009

Nissen, M. J., & Bullemer, P. (1987). Attentional requirements of learning: Evidence from

performance measures. *Cognitive Psychology*, *19*(1), 1–32.

https://doi.org/10.1016/0010-0285(87)90002-8

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh

inventory. *Neuropsychologia*, *9*(1), 97–113. https://doi.org/10.1016/0028-

3932(71)90067-4

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: Open Source

Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological

Data. *Computational Intelligence and Neuroscience*, *2011*, e156869.

https://doi.org/10.1155/2011/156869

Ozdemir, R. A., Kirkman, S., Magnuson, J. R., Fried, P. J., Pascual-Leone, A., & Shafi, M.

M. (2022). Phase matters when there is power: Phasic modulation of corticospinal

excitability occurs at high amplitude sensorimotor mu-oscillations. *Neuroimage:*

*Reports*, *2*(4), 100132. https://doi.org/10.1016/j.ynirp.2022.100132

Pascual-Leone, A., Grafman, J., & Hallett, M. (1994). Modulation of cortical motor output

maps during development of implicit and explicit knowledge. *Science*, *263*(5151),

1287–1289. https://doi.org/10.1126/science.8122113

Perez, M. A., Wise, S. P., Willingham, D. T., & Cohen, L. G. (2007). Neurophysiological

Mechanisms Involved in Transfer of Procedural Knowledge. *Journal of Neuroscience*,

*27*(5), 1045–1053. https://doi.org/10.1523/JNEUROSCI.4128-06.2007

Pinsard, B., Boutin, A., Gabitov, E., Lungu, O., Benali, H., & Doyon, J. (2019). Consolidation

alters motor sequence-specific distributed representations. *eLife*, *8*, e39324.

https://doi.org/10.7554/eLife.39324

Robertson, E. M. (2019). Skill Memory: Mind the Ever-Decreasing Gap for Offline

Processing. *Current Biology*, *29*(8), R287–R289.

https://doi.org/10.1016/j.cub.2019.03.007

Robertson, E. M., & Genzel, L. (2020). Memories replayed: Reactivating past successes and

new dilemmas. *Philosophical Transactions of the Royal Society B: Biological*

*Sciences*, *375*(1799), 20190226. https://doi.org/10.1098/rstb.2019.0226

Robertson, E. M., Pascual-Leone, A., & Miall, R. C. (2004). Current concepts in procedural

consolidation. *Nature Reviews Neuroscience*, *5*(7), 576–582.

https://doi.org/10.1038/nrn1426

Shirinpour, S., Alekseichuk, I., Mantell, K., & Opitz, A. (2020). Experimental evaluation of

methods for real-time EEG phase-specific transcranial magnetic stimulation. *Journal*

*of Neural Engineering*, *17*(4), 046002. https://doi.org/10.1088/1741-2552/ab9dba

Siegel, M., Warden, M. R., & Miller, E. K. (2009). Phase-dependent neuronal coding of

objects in short-term memory. *Proceedings of the National Academy of Sciences*,

*106*(50), 21341–21346. https://doi.org/10.1073/pnas.0908193106

Stefanou, M.-I., Desideri, D., Belardinelli, P., Zrenner, C., & Ziemann, U. (2018). Phase

Synchronicity of μ-Rhythm Determines Efficacy of Interhemispheric Communication

Between Human Motor Cortices. *The Journal of Neuroscience*, *38*(49), 10525–

10534. https://doi.org/10.1523/JNEUROSCI.1470-18.2018

Suresh, T., & Hussain, S. J. (2023a). Re-evaluating the contribution of sensorimotor mu

rhythm phase and power to human corticospinal output: A replication study. *Brain*

*Stimulation*. https://doi.org/10.1016/j.brs.2023.05.022

Suresh, T., & Hussain, S. J. (2023b). Re-evaluating the contribution of sensorimotor mu

rhythm phase and power to human corticospinal output: A replication study. *Brain*

*Stimulation: Basic, Translational, and Clinical Research in Neuromodulation*, *16*(3),

936–938. https://doi.org/10.1016/j.brs.2023.05.022

Tunovic, S., Press, D. Z., & Robertson, E. M. (2014). A Physiological Signal That Prevents

Motor Skill Improvements during Consolidation. *Journal of Neuroscience*, *34*(15),

5302–5310. https://doi.org/10.1523/JNEUROSCI.3497-13.2014

VanRullen, R., Busch, N. A., Drewes, J., & Dubois, J. (2011). Ongoing EEG Phase as a

Trial-by-Trial Predictor of Perceptual and Attentional Variability. *Frontiers in*

*Psychology*, *2*, 60. https://doi.org/10.3389/fpsyg.2011.00060

Walker, M. P., Brakefield, T., Allan Hobson, J., & Stickgold, R. (2003). Dissociable stages of

human memory consolidation and reconsolidation. *Nature*, *425*(6958), 616–620.

https://doi.org/10.1038/nature01930

Watrous, A. J., Fell, J., Ekstrom, A. D., & Axmacher, N. (2015). More than spikes: Common

oscillatory mechanisms for content specific neural representations during perception

and memory. *Current Opinion in Neurobiology*, *31*, 33–39.

https://doi.org/10.1016/j.conb.2014.07.024

Wilkinson, L., & Jahanshahi, M. (2007). The striatum and probabilistic implicit sequence

learning. *Brain Research*, *1137*, 117–130.

https://doi.org/10.1016/j.brainres.2006.12.051

Wilkinson, L., Khan, Z., & Jahanshahi, M. (2009). The role of the basal ganglia and its

cortical connections in sequence learning: Evidence from implicit and explicit

sequence learning in Parkinson's disease. *Neuropsychologia*, *47*(12), 2564–2573.

https://doi.org/10.1016/j.neuropsychologia.2009.05.003

Wilkinson, L., & Shanks, D. R. (2004). Intentional Control and Implicit Sequence Learning.

*Journal of Experimental Psychology: Learning, Memory, and Cognition*, *30*(2), 354–

369. https://doi.org/10.1037/0278-7393.30.2.354

Wilkinson, L., Steel, A., Mooshagian, E., Zimmermann, T., Keisler, A., Lewis, J. D., &

Wassermann, E. M. (2015). Online feedback enhances early consolidation of motor

sequence learning and reverses recall deficit from transcranial stimulation of motor

cortex. *Cortex*, *71*, 134–147. https://doi.org/10.1016/j.cortex.2015.06.012

Wischnewski, M., Haigh, Z. J., Shirinpour, S., Alekseichuk, I., & Opitz, A. (2022). The phase

of sensorimotor mu and beta oscillations has the opposite effect on corticospinal

excitability. *Brain Stimulation: Basic, Translational, and Clinical Research in*

*Neuromodulation*, *15*(5), 1093–1100. https://doi.org/10.1016/j.brs.2022.08.005

Womelsdorf, T., Schoffelen, J.-M., Oostenveld, R., Singer, W., Desimone, R., Engel, A. K., &

Fries, P. (2007). Modulation of Neuronal Interactions Through Neuronal

Synchronization. *Science*, *316*(5831), 1609–1612.

https://doi.org/10.1126/science.1139597

Yokoi, A., & Diedrichsen, J. (2019). Neural Organization of Hierarchical Motor Sequence

Representations in the Human Neocortex. *Neuron*, *103*(6), 1178-1190.e7.

https://doi.org/10.1016/j.neuron.2019.06.017

Zrenner, C., Desideri, D., Belardinelli, P., & Ziemann, U. (2018). Real-time EEG-defined

excitability states determine efficacy of TMS-induced plasticity in human motor

cortex. *Brain Stimulation*, *11*(2), 374–389. https://doi.org/10.1016/j.brs.2017.11.016

Zrenner, C., Kozák, G., Schaworonkow, N., Metsomaa, J., Baur, D., Vetter, D., Blumberger,

D. M., Ziemann, U., & Belardinelli, P. (2023). Corticospinal excitability is highest at

the early rising phase of sensorimotor μ-rhythm. *NeuroImage*, *266*, 119805.

https://doi.org/10.1016/j.neuroimage.2022.119805

1018