A Subset of Cortical Areas Exhibit Adult-like Functional Network Patterns in Early Childhood

Authors:

Jiaxin Cindy Tu[1], Yu Wang[2], Xintian Wang[1], Donna Dierker[1], Chloe M. Sobolewski[1,3], Trevor K. M. Day[4,5,6], Omid Kardan[7], Óscar Miranda-Domínguez[4], Lucille A. Moore[4], Eric Feczko[4], Damien A. Fair[4,5], Jed T. Elison[4,5], Evan M. Gordon[1], Timothy O. Laumann[1], Adam T. Eggebrecht[1], Muriah D. Wheelock[1]

Affiliations:

[1]Department of Radiology, Washington University in St. Louis.
[2]Department of Mathematics and Statistics, Washington University in St. Louis.
[3]Department of Psychology, Virginia Commonwealth University.
[4]Masonic Institute for the Developing Brain, University of Minnesota.
[5]Institute of Child Development, University of Minnesota.
[6]Center for Brain Plasticity and Recovery, Georgetown University.
[7]Department of Psychiatry, University of Michigan.

Corresponding Author

Muriah D. Wheelock
Mallinckrodt Institute of Radiology
4525 Scott Ave
St. Louis, MO 63110
mdwheelock@wustl.edu

Highlights

- Previous studies primarily investigated age-specific networks in infants, with limited focus on how well adult networks describe infant functional connectivity (FC).
- Our analysis identified a subset of areas in infants showing adult-like network organization, where within-network FC shows less age-related variation and higher scan-to-scan reliability.
- These areas are positioned near locations with low variability in functional network identity in adults, indicating a potential link between developmental sequence and interindividual variability in functional network organization.

Keywords

fMRI, functional connectivity, system, network, infant, resting state

Abstract

The human cerebral cortex contains groups of areas that support sensory, motor, cognitive, and affective functions, often categorized into functional networks. These

networks show stronger internal and weaker external functional connectivity (FC), with
FC profiles more similar within the same network. Previous studies have shown these
networks develop from nascent forms before birth to their mature, adult-like structures in
childhood. However, these analyses often rely on adult functional network definitions.
This study assesses the potential misidentification of infant functional networks when
using adult models and explores the consequences and possible solutions to this
problem.
Our findings suggest that although adult networks only marginally describe infant
FC organization better than chance, misidentification is primarily driven by specific
areas. Restricting functional networks to areas with adult-like network clustering
revealed consistent within-network FC across scans and throughout development.
These areas are also near locations with low network identity variability. Our results
highlight the implications of using adult networks for infants and offer guidance for
selecting and utilizing functional network models based on research questions and
scenarios.
58

1. Introduction

The human cerebral cortex comprises specialized, large-scale, functional
networks (Power et al., 2011; Yeo et al., 2011), supporting sensory, motor, higher-
cognitive, and affective functions (Petersen & Sporns, 2015; Wig, 2017). These
networks help segregate information processing across different sensory modalities and
cognitive domains (Grayson & Fair, 2017; Petersen & Sporns, 2015). In adults, these
networks consistently exhibit similar spatial topographies across various acquisition
paradigms (task and resting states), and individuals (Gratton et al., 2018). They can
also be disrupted by disease (Fornito et al., 2015; Fox & Greicius, 2010). Research
indicates that functional networks develop from infancy through old age (Grayson &
Fair, 2017; Sun et al., 2023; Wig, 2017), mirroring the progression of complex behavior
functions (Grayson & Fair, 2017; Petersen & Sporns, 2015). Preliminary forms of adult
functional networks are detectable in utero (Thomason et al., 2013; Turk et al., 2019).
Robust, sometimes bilateral, segregated networks for somatomotor, primary auditory,
primary visual, and extrastriate visual cortex are present in infants, whereas higher-
order networks appear less mature in early infancy (Eyre et al., 2021; Fransson et al.,
2007; Gao, Alcauter, Elton, et al., 2015, 2015; Moore et al., 2024; Myers et al., 2024;
Smyser et al., 2010; Sylvester et al., 2022). By 1-2 years, the default mode network
becomes more adult-like in some studies (Gao, Alcauter, Elton, et al., 2015; Gao,
Alcauter, Smith, et al., 2015; Gao et al., 2009) but remains localized in others
(Eggebrecht et al., 2017; Kardan et al., 2022; Marrus et al., 2018; F. Wang et al., 2023).
Researchers analyzing infant neuroimaging data often face a dilemma in
choosing an appropriate representation model. Some use the adult network models to
describe the relationships between functional connectivity (FC) and behavioral
phenotypes in infants (Nielsen et al., 2022; Rudolph et al., 2018; Tooley et al., 2024) or
to compare infants and adults (Yates et al., 2023). This choice is sometimes justified by
the need to enhance biological interpretability and facilitate communication using
consistent terminology across different research groups. However, applying exact adult
topography to define functional networks in infants is prone to inaccuracies and may
cause the mixing of fMRI BOLD signals (Smith et al., 2011) across different functional

networks, reducing statistical power. Additionally, differences in FC may stem from
variations in network topography or identity (Bijsterbosch et al., 2018, 2019). Prior-
based precision functional mapping using an adult network atlas is also commonly
employed in pediatric cohorts (Hermosillo et al., 2024; Moore et al., 2024; Sun et al.,
2023). However, these methods assume that infant network organization closely
resembles that of adults, which may not be valid.
An alternative approach involves deriving data-driven putative functional
networks specific to each developmental stage (Eggebrecht et al., 2017; Kardan et al.,
2022; Marrus et al., 2018; Tu et al., 2024; Wheelock et al., 2019). This method could
potentially address the issue of poor FC representation within functional networks and
improve reproducibility. However, the utility and interpretability of these putative
functional networks are less clear, as these putative functional networks have not been
validated through task activation studies. Ultimately, the choice of network model should
depend on the research goal. It is also crucial to understand the extent to which adult
network topography fits infant data. If the adult functional network topography differs
significantly from that of infants, using adult functional networks in infant studies could
lead to low reliability (Marek et al., 2022). Here, we aim to investigate this issue further
and examine the similarities and differences between adult and infant networks in
describing the modular structure of FC.
Converging evidence from various modalities indicates that the human cortex
develops non-uniformly across different areas. Primary sensory and motor cortex areas
mature earlier than higher-order association cortex areas (Ahmad et al., 2023; Flechsig,
1901; Garcia et al., 2018; Grayson & Fair, 2017; Hill et al., 2010; Sydnor et al., 2021;
Truzzi & Cusack, 2023). Early maturation in some areas results in limited future
plasticity (Hill et al., 2010), which may lead to lower interindividual variability and
reduced susceptibility to environmental influences (Gao et al., 2017) and
psychopathological factors (Sydnor et al., 2021). We hypothesize that some areas
would exhibit early signs of adult-like organization, especially in sensorimotor areas
(Gao, Alcauter, Elton, et al., 2015; Sydnor et al., 2021). Additionally, we hypothesize
that areas with adult-like organization would overlap with regions showing low
interindividual variability in functional network assignments (Dworetsky et al., 2021;
Gordon, Laumann, Adeyemo, et al., 2017; Gratton et al., 2018; Hermosillo et al., 2024;
Kong et al., 2019; Langs et al., 2016; Seitzman et al., 2019).
In the present work, we used fMRI datasets from two groups: 120 adults (aged
19-32 years) and 181 typically developing infants (aged 8-60 months). We aimed to
quantify how well the adult and infant networks describe the modular structure in both
adult and infant FC. We also examined if adult networks fit some areas better than
others by mapping the spatial distribution of this fit. We identified a subset of areas
where infant FC was more similar within their assigned adult networks than within
alternative adult networks. We demonstrate the consequences of network model choice
by comparing the relationship between chronological age and within-network FC in our
area subset to that of all areas. Lastly, we compared the spatial distribution of our area
subset to locations with low variance in functional network identity across individuals
and demonstrate their proximity to each other. Our findings will help researchers
working with infant neuroimaging data understand the pros and cons of using adult and

infant functional network models, interpret current results in the literature, and select appropriate models for future research.

2. Materials and Methods

2.1. Data Collection

2.1.1. Washington University 120 (WU 120)

This dataset has been previously described in detail (Power et al., 2017). Briefly, data were collected from 120 healthy young adult subjects during relaxed eyes–open fixation (60 females, mean age = 25 years, age range = 19–32 years). All subjects were native speakers of English and right-handed. Subjects were recruited from the Washington University community and were screened with a self-report questionnaire to ensure that they had no current or previous history of neurological or psychiatric diagnosis, as well as no head injuries resulting in a loss of consciousness for more than 5 minutes. Informed consent was obtained from all subjects. The study was approved by the Washington University School of Medicine Human Studies Committee and Institutional Review Board.

Structural and functional MRI data were obtained with a Siemens MAGNETOM Trio Tim 3.0-T Scanner (Erlangen, Germany) and a Siemens 12-channel Head Matrix Coil. A T1-weighted sagittal magnetization-prepared rapid acquisition gradient-echo (MP-RAGE) structural image was obtained [time echo (TE) = 3.08 ms, time repetition, TR (partition) = 2.4 s, time to inversion (TI) = 1000 ms, flip angle = 8°, 176 slices with 1 × 1 × 1 mm voxels]. An auto-align pulse sequence protocol provided in the Siemens software was used to align the acquisition slices of the functional scans parallel to the anterior commissure–posterior commissure plane of the MP-RAGE and centered on the brain. This plane is parallel to the slices in the Talairach atlas (Talairach & Tournoux, 1988).

During functional MRI data acquisition, subjects were instructed to relax while fixating on a black crosshair that was presented against a white background. Functional imaging was performed using a BOLD contrast-sensitive gradient-echo echo-planar imaging (EPI) sequence (TE = 27 ms, flip angle = 90°, in-plane resolution = 4 × 4 mm). Whole-brain EPI volumes (MR frames) of 32 contiguous, 4-mm-thick axial slices were obtained every 2.5 s. A T2-weighted turbo spin-echo structural image (TE = 84 ms, TR = 6.8 s, 32 slices with 1 × 1 × 4 mm voxels) in the same anatomical planes as the BOLD images was also obtained to improve alignment to an atlas. Anterior→Posterior (AP) phase encoding was used for fMRI acquisition. The number of volumes collected from subjects ranged from 184 to 724 (mean = 336 frames, 14.0 min).

2.1.2. Baby Connectome Project (BCP)

Full-term (gestational age of 37-42 weeks) infants free of any major pregnancy and delivery complications were recruited as part of the Baby Connectome Project (Howell et al., 2019). All procedures were approved by the University of North Carolina at Chapel Hill and the University of Minnesota Institutional Review Boards. Informed consent was obtained from the parents of all participants. In the final cohort used following fMRI data quality control (described below), we retained 313 fMRI sessions

from 181 individuals (95 females, 8-60 months, mean 19.1 months, and standard
deviation 8.3 months) during natural sleep (Supplementary Figure 1A). The number of
longitudinal points range from 1 to 6: 90 individuals with 1 time point, 60 individuals with
2 time points, 24 individuals with 3 time points, 5 individuals with 4 time points, 1
individual with 5 time points and 1 individual with 6 time points (Supplementary Figure
1B). In a supplementary analysis, we used an additional 15 fMRI sessions collected
while participants were watching a movie clip.
All MRI images were acquired on a Siemens 3T Prisma scanner with a 32-
channel head coil at the University of Minnesota and the University of North Carolina at
Chapel Hill during natural sleep without the use of sedating medications. T1-weighted
(TR=2400 ms, TE=2.24 ms, 0.8 mm isotropic; flip angle = 8°), T2-weighted images
(TR=3200 ms, TE=564 ms, 0.8 mm isotropic), spin echo field maps (SEFM) (TR=8000
ms, TE=66 ms, 2 mm isotropic, MB=1), and fMRI data (TR=800 ms, TE=37 ms, 2 mm
isotropic, MB=8) were collected. A mixture of Anterior→Posterior (AP) and
Posterior→Anterior (PA) phase encoding directions was used for fMRI acquisition in
each session, but they were concatenated into one time series. A subset of data had a
720-ms TR (N = 95 out of 313 sessions). The number of low-motion volumes collected
from subjects ranged from 840 to 2100 (mean = 1306 frames, 16.9 min).
198
199 **2.2. fMRI analysis**
200
201 **2.2.1. MRI data preprocessing**
202   While both datasets were acquired on Siemens 3T scanners, acquisition
203 parameters and preprocessing of the structural and functional MRI data were optimized
204 for each specific cohort. For a detailed comparison of the acquisition, processing, and
205 quality control procedures, please refer to Supplementary Table 1.
206
2.2.1.1. MRI data preprocessing – WU120
Functional images were first processed to reduce artifacts including (1)
Correction of odd versus even slice intensity differences attributable to interleaved
acquisition without gaps, (2) correction for head movement within and across runs, and
(3) across-run intensity normalization to a whole-brain mode value of 1000. Atlas
transformation of the functional data was computed for each individual using the MP-
RAGE scan. Each run was then resampled to an isotropic 3-mm atlas space (Talairach
& Tournoux, 1988), combining movement correction and atlas transformation in a single
cubic spline interpolation (Lancaster et al., 1995).
Additional preprocessing steps were applied to the functional data to reduce the
effect of high-motion frames. This was performed in two iterations. In the first iteration,
the processing steps were (1) demeaning and detrending, (2), multiple regression
including whole-brain, ventricular cerebrospinal fluid (CSF), and white matter signals,
and motion regressors derived by Volterra expansion and (3) a band-pass filter (0.009
Hz < f < 0.08 Hz). Following the initial FC preprocessing iteration, temporal masks were
created to flag motion-contaminated frames. Motion-contaminated volumes were
identified by framewise displacement (FD), defined as the squared sum of the motion
vectors (Power et al., 2012). Volumes with FD > 0.2 mm and data segments lasting
fewer than 5 contiguous volumes were censored.

The data were then reprocessed in a second iteration, incorporating the temporal
masks described above. This reprocessing was identical to the initial processing stream
but ignored censored data. Data were interpolated across censored frames using least
squares spectral estimation (Power et al., 2014) of the values at censored frames, so
that continuous data could be passed through the band-pass filter (0.009 Hz < f < 0.08
Hz) without contaminating frames near high motion frames. Censored frames were
ultimately ignored during functional connectivity matrix generation.
Individual surfaces were generated from the structural images and the functional
data was sampled to surface space (Glasser et al., 2013). First, following volumetric
registration, anatomical surfaces for the left and right hemispheres were generated from
each subject's MP-RAGE image using FreeSurfer's default recon-all processing pipeline
(v5.0)(Fischl, 2012). This pipeline included brain extraction, segmentation, generation of
white matter and pial surfaces, inflation of the surfaces to a sphere, and surface shape-
based spherical registration of the subject's "native" surface to the fsaverage surface.
The fsaverage-registered left and right hemisphere surfaces were then brought into
register with each other (Van Essen et al., 2012), resampled to a resolution of 164000
vertices using Caret tools (Van Essen et al., 2001) and subsequently down-sampled to
a 32492 vertex surface (32k fs_LR). The BOLD volumes were sampled to each
subject's individual "native" midthickness surface (generated as the average of the white
and pial surfaces) using the ribbon-constrained sampling procedure available in
Connectome Workbench (v0.84) and then deformed and resampled from the
individual's "native" surface to the 32k fs_LR surface. Finally, the time courses were
smoothed along the 32k fs_LR surface using a Gaussian smoothing kernel (σ = 2.55
mm).
250
2.2.1.2. MRI data preprocessing – BCP
MRI data were processed using the DCAN-Labs infant-abcd-bids-pipeline
(v0.0.22) largely following steps described previously (Feczko et al., 2021). Structural
MRI data underwent HCP-style processing (Feczko et al., 2021; Glasser et al., 2013),
including ANTS N4 bias correction, ANTS denoising, T1/T2 distortion
correction/registration, and finally ANTS SyN algorithm deformation alignment to an
infant MNI template. In addition, a refined brain mask was generated from data that was
segmented using in-house age-specific templates via Joint Label Fusion (JLF). The
toddler-specific mask and segmentation were substituted into the FreeSurfer (Fischl,
2012) pipeline and used to refine the white matter segmentation and guide the
FreeSurfer surface delineation. The native surface data were then deformed to the 32k
fs_LR template via a spherical registration.
A scout image (frame 16 in each run) was selected from the fMRI time series for
functional MRI preprocessing. The scout was distortion-corrected via spin-echo field
maps, served as the reference for motion correction via rigid-body realignment (Feczko
et al., 2021), and was registered to the native T1. Across-run intensity normalization to a
whole-brain mode value of 10,000 was then performed. These steps were combined in
a single resampling with the MNI template transformation from the previous step, such
that all fMRI frames were registered to the infant MNI template. Manual inspection of
image quality of structural and functional data was conducted to exclude sessions with
bad data quality.

To prepare the functional data for FC analysis, further processing steps were
applied after sampling the BOLD data to the 32k fs_LR surface space using steps
described in 2.2.1.1. First, functional data were demeaned and detrended in time.
Denoising was then performed using a general linear model with regressors including
signal and motion variables. Signal regressors included mean CIFTI gray-ordinate time
series, Joint Label Fusion (JLF)-defined white matter, and JLF-defined CSF. Motion
regressors included volume-based translational and rotational components and their 24-
parameter Volterra expansion. The movement of the head was measured by FD and an
age-specific respiratory notch filter (0.28-0.48 Hz) was applied to the FD traces and
motion parameter estimates to mitigate the effects of factitious head motion due to
infant respiration (Fair, 2020; Kaplan et al., 2022). Frames were censored during
demeaning/detrending if their post-respiratory filtering FD value exceeded 0.3 mm to
generate the denoised beta values in the general linear model. Bandpass filtering was
applied using a second-order Butterworth filter (0.008–0.09 Hz). To preserve the
temporal sequence and avoid aliasing caused by missing time points during bandpass
filtering, interpolation was used to replace missing frames, and residuals were acquired
from the denoising general linear model. In addition, zero-padding was applied to both
ends of the BOLD data before filtering to minimize the distortions in the edges of the
time series. The data were originally minimally spatially smoothed with a geodesic 2D
Gaussian kernel (σ = 0.85 mm). A further smoothing with a geodesic 2D Gaussian
kernel (σ = 2.40 mm) was applied to give a final effective smoothing of σ = 2.55 mm to
match the smoothing used in the adult dataset (WU 120). Finally, the time series were
concatenated across all complete and partially completed scan runs with good data
quality. The first 7 frames from each run, frames with > 0.2 mm FD post-respiratory
filtering (Kaplan et al., 2022), and outlier frames whose across-vertex standard deviation
was more than 3 median absolute deviations from the median of the low FD frames
were censored and ignored for functional connectivity matrix construction.
299
300   **2.2.2. Functional Connectivity Matrix Construction**
301   The preprocessed BOLD time series data of each session were parcellated into
333 non-overlapping areas using the Gordon parcellation (Gordon et al., 2016). This
choice of parcellation was justified by recent work by our group that demonstrated that
the Gordon parcellation had the best fit among a set of adult parcellations and
performed comparably to most available infant parcellations in data from infants aged
around 8-30 months (Tu et al., 2024). After that, a total number of frames equivalent to
7.2 minutes of data (560 frames for TR = 0.72 and 600 frames for TR = 0.8) were
randomly sampled from the full censored time series in each fMRI session. Pearson's
correlation between the parcellated time series was computed to create a 333 x 333
functional connectivity (FC) matrix. This matrix was then Fisher-Z-transformed. The
group-average FC matrix was calculated as the mean FC across fMRI sessions.
312
313   **2.3. Infant and Adult Functional Network Schemes**
314   We used the Gordon network assignments (Gordon et al., 2016) for "Adult
Networks" (Figure 1A) and Kardan network assignments (Kardan et al., 2022) for "Infant
Networks" (Figure 1B). These network assignments were derived from group-average
adult and infant FC on the 333 areas using an Infomap community detection algorithm

(Rosvall & Bergstrom, 2010) optimized for identifying networks in FC data (Power et al.,
2011). Among the 333 areas, some were originally assigned in communities with fewer
than 5 areas and considered unassigned (named "None" and "Unspecified"). These
areas commonly fall under locations subjected to the biggest susceptibility artifact
(Ojemann et al., 1997). We removed them from all analyses and had 286 areas left for
the adult networks ("Gordon") and 328 areas left for the infant networks ("Kardan").
The 12 Gordon networks include the auditory (Aud), cingulo-opercular (CON),
parietal memory (PMN), default mode (DMN), dorsal attention (DAN), fronto-parietal
(FPN), retrosplenial temporal (RTN), somatomotor hand (SMN hand), somatomotor
mouth (SMN mouth), salience (Sal), ventral attention (VAN), and visual (Vis) networks.
The 10 Kardan networks include somatomotor (SMN), temporal (Tem), posterior
frontoparietal (pFPN), posterior default mode (pDMN), lateral visual (lVis), medial visual
(mVis), dorsal attention (DAN), anterior fronto-parietal (aFPN), anterior default mode
(aDMN).
332
2.4. Functional Network Overlap
The overlap between a network in the Gordon networks and a network in the
Kardan networks can be measured with the Dice coefficient, with 0 indicating no overlap
and 1 indicating complete overlap. For this analysis, each network is represented with a
333 x 1 vector with 1 for the areas in the network and 0 for the areas outside the
network.
339
340 **2.5. Silhouette Index Calculation**
341   Following prior procedures in the literature (Rousseeuw, 1987; Yeo et al., 2011),
342 we calculated the silhouette index (SI) for each area with the correlation distance (i.e. 1-
343 Pearson's correlation) in the FC profiles:
344
345
$$SI = \frac{b - a}{\max{(a, b)}} \ (Equation \ 1)$$

346
where *b* is the mean between-network correlation distance of the FC profiles, and *a* is
the mean within-network correlation distance of the FC profiles. FC profiles here refer to
the FC from each area to all other areas (i.e., one row in the FC matrix).
Intuitively, the SI ranges from +1 to -1 with the sign indicating whether the area
has a more similar FC profile to areas in its own network (+) or to areas in an alternative
network (-). The magnitude indicates the confidence of this assignment, with a higher
magnitude suggestive of strong confidence. The average SI for the FC in a network
scheme was defined as the average SI across all areas.
We also calculated the silhouette index with the average of all networks rather
than just the alternative network in the Supplementary Materials.
To obtain a confidence interval for the average SI across individual sessions, we
used a bootstrap 95% confidence interval estimate in 1000 random draws of individual
sessions from the full sample (N = 120 for WU120 and N = 313 for BCP) with
replacement. The p-value for the average SI different from zero was calculated using
the bootstrap distribution of the average SI. The p-value for the difference in the

average SI across two samples was calculated using the bootstrapped distribution of
the difference in the average SI. Both assumed a two-tailed test.
364
2.6. Identify the Subset of Areas with Similar Network Organization to the Gordon
Networks in Infants
A positive SI indicates that the area has a more similar FC profile to areas in its
own network. We obtained the subset of areas that had a similar network organization
to other areas defined in the Gordon network scheme in infants by only retaining the
areas with a positive SI in the group-average infant FC. We refer to this set of positive
SI areas as our "area subset".
372
2.7. Distance Between High Consensus Regions of Interests (ROIs) and the
"Gordon Subset" Areas
To quantify the spatial distribution similarity between the locations of low
interindividual variability and our area subset, we calculated the Euclidean distance
between the high consensus cortical ROIs and the centers of those areas. We used
published coordinates of 153 high consensus ROIs calculated previously by identifying
locations that demonstrated consistent network assignment across a large majority (i.e.
$\geq 75\%$) of subjects in the Dartmouth dataset (N = 69 subjects, 56 female, average age
20.2 years)(Gordon et al., 2016) when a template-matching procedure (Gordon,
Laumann, Adeyemo, et al., 2017) was applied to identify individual network
assignments (Dworetsky et al., 2021). The distances in mm were obtained from the ROI
locations on a standard adult brain template (a.k.a. MNI coordinates).
For each of the "high consensus" ROIs, we found its distances to the nearest
area within and outside our area subset. An average of this difference was recorded
and named "distance difference". A negative distance difference indicates that on
average, the high consensus regions were closer to the areas within compared to
outside our area subset. To account for the potential effect of differences in the number
of areas within (N = 166) and outside (N = 120) our area subset, we repeated the
analysis but with the labels of within and outside area subset randomly assigned 1000
times to obtain a null distribution.
393
2.8. Moving Average Analysis Across Age
To examine the FC fit to different network schemes across infancy, we used a
moving average analysis across age. For this analysis, we limited our data to the 281
sessions collected at the age of 8-27 months because the data became very sparse and
less evenly distributed after 27 months (Supplementary Figure 1A). We first sorted the
fMRI sessions by age at scan. Sessions were arranged chronologically by age, and FC
averages were computed for consecutive windows of 20 sessions, with each window
representing the mean age within it. This window was then shifted by one session at a
time until all 281 sessions were accounted for. Subsequently, we calculated the average
similarity index (SI) using the same method.
404
2.9. Intraclass Correlation Coefficient
To assess the differences in the reliability of within-network edges for using the
Gordon networks with our area subset versus all areas. We quantified the test-retest

reliability of FC with the intraclass correlation coefficient (ICC). We assessed the
consistency among measurements under the fixed levels of the session factor (Tozzi et
al., 2020), referred to as ICC 'C-1' (McGraw & Wong, 1996) or ICC (3,1) (Shrout &
Fleiss, 1979).
For this analysis, we re-calculated the FC matrices for each individual with two
non-overlapping time windows of data from each session. "Test" and "re-test" were
defined as the first 6 min and last 6 min of low-motion data, separated by at least 1.2
min low motion data in between to reduce the impact of temporal autocorrelation (i.e.
total > 13.2 min low-motion data). Only 167 sessions had enough low-motion data for
this analysis. First, the FC values in the upper triangle of each subject's connectivity
matrix were entered as rows in two large matrices (one matrix for "test" and another for
"re-test", one row per subject in each matrix). Then, the corresponding columns of these
matrices were compared to obtain an ICC value for each edge. The mean and standard
error of the mean of the ICCs within each of the Gordon networks were calculated for
our area subset and all areas.
423
424 **3. Results**
425
3.1. Adult and Infant functional connectivity clustering was best described by the
adult and infant network assignments respectively
Adult networks ("Gordon") (Figure 1A) and Infant networks ("Kardan") (Figure 1B)
assignments derived using data-driven methods demonstrate a moderate degree of
agreement: Normalized Mutual Information (NMI) = 0.5 for the overlapping 281 areas
after excluding the "None"/ "Unspecified" network in both adult and infant network
assignments. The CON, pDMN, aDMN, SMN, mVis and lVis networks in the Kardan
networks tend to have a large dice overlap with a single Gordon network, but Tem,
DAN, pFPN, aFPN have a match to multiple Gordon networks (Supplementary Figure 2,
network full names defined in section 2.3).
Next, we asked how closely the network assignments matched the similarity of
FC profiles within and between different networks and quantified it with the silhouette
index (SI; Rousseeuw, 1987; Yeo et al., 2011). We used the average FC across 120
adult sessions and the average FC across 313 infant sessions. We found that adult FC
had a more modular organization (Figure 1C) when grouping into adult networks (Figure
1A) than infant networks (Figure 1E). The average SI for areas assigned to adult
networks in adult FC (0.333, 95% bootstrap CI = [0.3088, 0.3417], $p_{bootsrap}$ < 0.001) was
much higher ($p_{bootsrap}$ < 0.001) than the average SI for areas assigned to infant networks
in adult FC (0.009, 95% bootstrap CI = [-0.0015, 0.0168], $p_{bootsrap}$ = 0.042). In contrast,
the opposite was observed for infant FC, with a higher average SI ($p_{bootsrap}$ < 0.001) for
areas assigned to infant networks in infant FC (0.336, 95% bootstrap CI = [0.3280,
0.3397], $p_{bootsrap}$ < 0.001; Figure 1F) than the average SI for areas assigned to adult
networks in infant FC (0.049, 95% bootstrap CI = [0.0406,0.0560], $p_{bootsrap}$ < 0.001;
Figure 1D). Furthermore, the results were also qualitatively validated across individual
sessions (details in Supplementary Materials), with a much higher SI of adult networks
than infant networks on adult FC (Cohen's d = 1.215, p < 0.001) (Supplementary Figure
3C), and a much higher SI of infant networks than adult networks on infant FC (Cohen's
d = 2.744, p < 0.001) (Supplementary Figure 4C). Taken together, the adult networks

better describe the modular organization in adult FC than infant FC, and the infant
networks better describe the modular organization in infant FC than adult FC. However,
the SI is comparable for adult FC and infant FC using the best network model,
suggesting the presence of modular organization in both cohorts. We also repeated the
same analysis for 15 infant sessions scanned while the participants were awake and
watching movies and 14 infant sessions scanned during natural sleep with similar age
range (Supplementary Figure 5). The difference in average SI between the adult
("Gordon") networks and infant ("Kardan") networks on the awake sessions was not
significant ($p_{bootsrap}$ = 0.258, paired t-test on individual sessions = 0.71), unlike the
difference in asleep sessions with matching age range ($p_{bootsrap}$ <0.001, paired t-test on
individual sessions < 0.001). The average SI for areas assigned to adult networks in
awake infant FC (0.174, 95% bootstrap CI = [0.1048, 0.1895], $p_{bootsrap}$ <0.001) was
much higher ($p_{bootsrap}$ = 0.002) than that in sleeping infant FC (0.084, 95% bootstrap CI
= [0.0355, 0.0962], $p_{bootsrap}$ = 0.001), but still lower ($p_{bootsrap}$ < 0.001) than that in adult
FC.
Notably, some areas tend to have a positive SI for areas assigned to adult
networks regardless of the FC age group (Figure 1G & I). Since the spatial distribution
of SI across sessions (Supplementary 3A-B & 4A-B) was relatively consistent, it is
unlikely that the low SI magnitude in infants was purely driven by high interindividual
variability.
Because the Kardan networks were derived from a subset of BCP sessions used
in the present analysis and this might inflate the SI in the infant FC data, we also
replicated our analysis with networks derived from an independent 2-year-old dataset
with similar acquisition and preprocessing protocols (Tu et al., 2024) (Supplementary
Figure 6). The SI of the independently derived infant 2-yearl-old networks was lower
than the Kardan networks. This may be partially attributed to the wider age range of the
BCP data relative to the toddler data used to derive the 2-year networks. However, it
may also suggest that the high average SI with Kardan networks is partially attributable
to data leakage. However, importantly, the independently derived infant networks
demonstrate significantly higher SI compared to Gordon adult networks when applied to
infant FC ($p_{bootsrap}$ < 0.001 for the Tu-12networks and Tu-19networks).

Figure 1. *Adult and Infant FC ordered by the adult and infant networks.* A) 12 adult networks, B) 10 infant networks, C) average adult FC sorted by adult networks, D) average infant FC sorted by adult networks, E) average adult FC sorted by infant networks, F) average infant FC sorted by infant networks, G) SI of parcels with adult network assignments in adults, H) SI of parcels with adult network assignments in infants, I) SI of parcels with infant network assignments in adults, J) SI of parcels with infant network assignments in infants. Network abbreviations: auditory (Aud), cingulo-opercular (CON), parietal memory (PMN), default mode (DMN), dorsal attention (DAN), fronto-parietal (FPN), retrosplenial temporal (RTN), somatomotor hand (SMN hand), somatomotor mouth (SMN mouth), salience (Sal), and ventral attention (VAN), visual (Vis), somatomotor (SMN), temporal (Tem), posterior frontoparietal (pFPN), posterior default mode (pDMN), lateral visual (lVis), medial visual (mVis), anterior fronto-parietal (aFPN), anterior default mode (aDMN).

3.2. A subset of areas demonstrates adult-like network organization throughout development

An SI above zero for an area indicates that its FC profile is more similar to the FC profiles in areas from the same network than areas from any alternative network within a given network scheme (e.g., adult Gordon networks). Therefore, we selected the subset of areas with an SI above zero when the adult networks were applied to the infant FC (166 in total, Figure 2A, Supplementary Table 2). These areas fell into all 11 out of the 12 Gordon networks (i.e., all except for PMN), with the whole RTN, SMN mouth, and Sal networks retained, and the remaining eight networks partially retained (Figure 2B). We validated that the areas with SI above zero are highly consistent across bootstrap samples, with 156 out of the 166 areas having SI above zero in at least 950 out of 1000 bootstraps (Supplementary Figure 7).

Figure 2. A subset of areas in infants demonstrates adult-like network organization.
A) A subset of areas with SI > 0 to adult networks in infants (Figure 1H). B) The number of areas in each adult network with all areas ("All") or area subset ("Subset"). C) The sorted average FC in infants with our area subset. D) The percentage representation of each network in all areas versus. E) The average within-network FC with all areas (left) versus area subset (right) across individual BCP sessions. FDR-corrected p for paired t-tests. *** p < 0.001.

As expected, the average SI for areas assigned to adult networks in infant FC
was much higher ($p_{bootstrap}$ < 0.001) using our area subset (0.388, 95% bootstrap CI =
[0.3792, 0.3925], $p_{bootstrap}$ < 0.001; Figure 2C) than using all areas. In addition, the
average SI for areas assigned to adult networks in adult FC was also marginally higher
($p_{bootstrap}$ < 0.001) using our area subset (0.419, 95% bootstrap CI = [0.3925,0.4300],
$p_{bootstrap}$ < 0.001) (Supplementary Figure 8A) than using all areas. Our results suggest
that this subset captures the part of the adult networks with more within-network
consistency in both infants and adults. Compared with all areas, our area subset was
disproportionally enriched in the SMN networks (SMN hand and SMN mouth) (Figure
2D). As expected, the within-network FC was significantly higher across infant sessions
(paired t-test, FDR-corrected p < 0.05) for all eight partially retained networks when our
area subset was used instead of all areas, with little change in variability (Figure 2E).
Similarly, within-network FC was significantly higher across adult sessions (paired t-test,
FDR-corrected p < 0.05) for all seven out of eight partially retained networks
(Supplementary Figure 8B). Because our data spanned a wide developmental window,
we further investigated whether our observation was influenced by chronological age.
We found that within-network FC for all eight partially retained networks was higher in
our area subset in most sessions and that there was no correlation (FDR corrected p >
0.05) between the within-network FC differences and age (Supplementary Figure 9). In
general, the within-network FC differences between "All" and "Subset" were larger in
infants than in adults (Table 1).
520

521   **Table 1.** *Cohen's d of the within-network FC differences in All V.S. Subset.*

	Aud	CON	DMN	DAN	FPN	SMN hand	VAN	Vis
Infant FC	-2.41	-0.29	-1.80	-2.23	-1.54	-2.81	-2.64	-2.91
Adult FC	-1.13	-0.31	-0.46	-1.47	0.70	-1.09	-1.53	-2.10

Specifically, for seven out of the eight partially retained Gordon networks the
adult FC was significantly higher (FDR-corrected *p* < 0.05) than the infant FC with all
areas (Figure 3A, Table 2). On the other hand, only five out of the eight still
demonstrated significantly higher within-network FC when using the subset (FDR-
corrected *p* < 0.05) in adults compared to infants, and two out of the eight demonstrated
significantly lower within-network FC (FDR-corrected *p* < 0.05) (Figure 3B, Table 2).

Table 2. *Cohen's d of the within-network FC differences in adults V.S. infants*

	Aud	CON	DMN	DAN	FPN	SMN hand	VAN	Vis
All	2.35	2.35	0.80	2.11	0.23	1.04	0.03	1.22
Subset	1.04	1.88	0.13	1.04	-0.26	0.43	-0.34	0.94

Additionally, we found that the effect of chronological age on within-network FC
within the infant cohort was also reduced when our area subset was used in place of all
areas (Supplementary Figure 10).

Taken together, our results suggest that while the within-network FC within our
area subset was higher than that within all areas in both infant and adult datasets, using
our area subset would reduce the difference across age.

Figure 3. *Violin plot of within-network average FC in the Gordon networks.* A) within-network average FC using all areas in adults and infants. B) Within-network average FC using our area subset in adults and infants. FDR-corrected *p* for two-sample t-tests. * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001.

538
3.3 Adult versus infant FC organizations across 1-2-year-olds

Next, we investigated whether there was any variation between how the different
network schemes fit the infant FC at various stages between 1 to 2 years (Gordon adult
networks and Kardan infant networks). In addition, we also included a comparison when
using our area subset (166 areas, Figure 2A). Using a moving average approach across
infant ages, we found a consistent order of the network schemes across 1 to 2 years,
with the Gordon (Subset), Kardan (All), and Kardan (Subset) having a similar average
SI, and Gordon (All) having a much lower average SI (Figure 4A). Nevertheless, we saw
a subtle decrease in difference between adult and infant networks on the infant FC with
increasing age. We further investigated this trend using average SI in individual

Figure 4. *Moving average analysis with the adult networks ("Gordon") and the infant networks ("Kardan") using all areas and our area subset.* A) Average SI of the average FC in a window for different network assignments. B) Average SI of the window around 1 year old sorted by different network assignments. C) Average SI of the window around 2 years old sorted by different network assignments.

sessions spanning 8 to 60 months (Supplementary Figure 4). We found a weak to
moderate negative correlation (r = -0.21, p < 0.001, Supplementary Figure 11A) in infant
networks ("Kardan") fit (average SI) with age, and a weak positive correlation (r = 0.13,
p = 0.020, Supplementary Figure 11B) in adult networks ("Gordon") fit (average SI) with
age. Together, this amounted to a moderate negative correlation in the difference in
average SI between infant and adult networks with age (Supplementary Figure 11C).
We also replicated Figure 4A with network parcellations derived from an
independent dataset to eliminate potential circularity in using the infant networks derived
from a subsample of our infant FC data (Supplementary Figure 12).
558
3.4 Our area subset with adult-like network organization is in spatial proximity to
the high consensus regions across adult individuals
To quantify the spatial distribution similarity between the locations of low
interindividual variability in network identity ("high consensus cortical ROIs") (Dworetsky
et al., 2021) and our area subset, we calculated the Euclidean distance between the
centers of the areas within our area subset or alternative areas to the "high consensus
cortical ROIs" (Figure 5). Areas within our area subset were 9.5 ± 4.5 mm to the closest

high consensus ROIs, whereas areas outside our area subset were 13.5 ± 6.6 mm to
the closest high consensus ROIs (Supplementary Figure 13). On average, areas within
our area subset were 3.9 mm closer to the "high consensus cortical ROIs". To rule out
the possibility that this difference was driven by the differences in the number of areas
within and outside our area subset, we repeated the same analysis by permuting the
binary within and outside area subset labels 1000 times to generate a null distribution.
We found that the actual difference (3.9 mm) was significantly higher than the null (p <
0.001, permutation testing) (Figure 5C).

Figure 5. *Comparison of area centers to high-consensus regions.* A) Our area subset with adult-like network organization (colored surface patches) overlayed with high-consensus regions (rosy brown spheres). B) The areas outside the subset (colored surface patches) overlayed with high-consensus regions (rosy brown spheres). C) The average Euclidean distance between the high-consensus regions and the closest parcel center. Dashed line: the actual difference between the distances in panel A and panel B. Histogram: the difference between the distances with parcels randomly assigned to be in the adult-like (panel A) and not adult-like (panel B) groups 1000 times.

3.6. Within-network FC edges in our area subset have a higher test-retest
reliability and a higher consistency across subjects
Comparing FC computed from non-overlapping time windows in the same
session demonstrated that our area subset had significantly higher within-session
reliability than all areas. In particular, four out of eight networks partially retained (Figure
2B) exhibited higher average ICC with our area subset than all areas (two-sample t-test,
FDR-corrected $p < 0.05$): Aud (Cohen's $d = 0.68$), DMN (Cohen's $d = 0.13$), DAN
(Cohen's $d = 0.55$), and VAN (Cohen's $d = 0.43$) (Figure 6). To examine whether the
contribution of FC edges to individual identification varied across the within- and
between-network blocks by the three network schemes, we also quantified the FC
group consistency (ϕ) and differential power (DP) (Finn et al., 2015). We found the FC-
edges connecting our area subset tend to be the ones more consistent across scans
and subjects (Supplementary Materials S4, Supplementary Tables 3-6, Supplementary

Figure 14).

Figure 6. *Reliability of within-network edges using our area subset versus all areas.* Reliability of within-network edges across two non-overlapping 6-minute windows. Bars show the mean of the ICC and error bars show the standard error of the mean. FDR-corrected *p* for two-sample t-tests. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

588
589
4. Discussion
4.1. Similarity and differences between adult and infant network organization
We observed that for infants at 8-60 months, adult networks captured some of the
features in the infant FC (average SI>0), although much weaker than that in adult FC.
Furthermore, it seemed that instead of being less modular and more random, the infant
FC data were better described with notably different but related modular organization,
including fragmented anterior and posterior segments of higher-order association
networks (Eggebrecht et al., 2017; Eyre et al., 2021; Kardan et al., 2022; Marrus et al.,
2018).

599
4.2. Identification of a subset of areas that are stable across development
We found that a subset of the areas tended to exhibit more of an adult-like network
FC clustering pattern even in infants, forming the cores of adult networks. While
substantial attention has been devoted to the differences in infant network organization
throughout development (F. Wang et al., 2023; Wen et al., 2019, 2020), it is also desirable
to note their similarities to older children and adults (Fransson et al., 2007; Gao, Alcauter,
Elton, et al., 2015). Here we observed that the difference in within-network FC across
ages was reduced when an area subset was used instead of all areas. The FC within our
area subset was also more consistent both within the session and across sessions. These
areas likely form the early scaffold for what will eventually become the adult networks
(Grayson & Fair, 2017).

611
612   ### 4.3. The possible role of childhood experience in shaping the interindividual
613   differences in functional networks
614   Our results suggest that interindividual variability in functional network topography
615   might have a developmental origin. To a first approximation, the spatial topography of our

area subset closely matches the locations with low interindividual variability in network identity (Dworetsky et al., 2021; Gordon, Laumann, Gilmore, et al., 2017; Hermosillo et al., 2024). On the other hand, the other areas approximately correspond to previously identified hub nodes (Power et al., 2013) and integration zones (Hermosillo et al., 2024).

One potential explanation for this observation is that some parts of the brain mature earlier than others in terms of functional network organization. There might be biological or evolutionary reasons that parts of the adult networks mature later, such as to devote prenatal resources to regions most important for early survival (Hill et al., 2010). In addition, the regional variability of network stability might also be linked to variability in the expression of excitatory and inhibitory features across the cortex (Sydnor et al., 2021). The idea that areas with higher FC variability have more significance in interindividual differences is further reinforced by research demonstrating that FC in cortical areas with high FC variability has more predictive power in behavioral and cognitive domain features (Mueller et al., 2013). It is important that future researchers recognize the regional variability in functional network stability. This variability may serve as a useful biomarker for psychopathology (Sydnor et al., 2021), or may guide targeted brain stimulation interventions.

We did not observe a strong bias in the over-representation of sensorimotor networks compared to association networks in our area subset, despite the literature suggesting that sensorimotor networks mature earlier than association networks (Gao, Alcauter, Elton, et al., 2015; Sydnor et al., 2021). Our area subset spanned both sensorimotor and association networks along the functional hierarchy of the neocortex (Flechsig, 1901; Mesulam, 1998; Sydnor et al., 2021). One potential limitation is that our infant cohort was older than eight months and significant earlier neurodevelopmental changes along the sensorimotor-association hierarchy might have happened before eight months (Bethlehem et al., 2022; Flechsig, 1901). Another possibility is that the sensorimotor functional networks definition was inaccurate, e.g. the auditory network might have incorporated parts of secondary somatosensory regions (Raju & Tadi, 2024), leading to heterogeneous FC profiles within the auditory network

4.4. Using an area subset to improve statistical power and interpretability

There are pros and cons of using a pre-existing functional network model and a data-driven functional network model for infant neuroimaging research. Studies of functional networks in infants have often implemented unsupervised methods (i.e. clustering or similar types of community detection algorithms) to find age-specific modules and called them "functional networks" (Eggebrecht et al., 2017; Kardan et al., 2022; Marrus et al., 2018; Molloy & Saygin, 2022; Myers et al., 2024; Sylvester et al., 2022; F. Wang et al., 2023; Wen et al., 2019, 2020). These identified modules were by definition a good representation of the organizational structure in the data and may help alleviate the problem of reproducibility in brain-wise association studies (Hermosillo et al., 2024; Marek et al., 2022). However, unlike the adult networks that have been extensively validated with task fMRI data to corroborate their "functional" roles (Power et al., 2011; Wig, 2017; Yeo et al., 2011), those age-specific modules often lack biological support for their functions, reducing their relevance to a broader developmental context. On the other hand, using the adult-network topography directly on infants neglects the infant-specific organizational features and risks including spurious variability in neuroimaging

measurements (be they from fMRI, PET, or fNIRS), leading to reduced effect size and statistical power (Hermosillo et al., 2024), or an exaggerated difference across development. For example, our results in section 3.2 suggest that differences in within-network FC across age groups might be partially attributed to the misspecification of functional network identity.

Here, we propose an alternative strategy that uses an area subset that is relatively stable throughout development. This approach strikes a balance between interpretability/comparability across cohorts and reliability/reproducibility. This idea of using a subset of the brain areas to define ROIs as an approach to improve statistical power has been mentioned in prior literature (Dworetsky et al., 2021; Hermosillo et al., 2024). However, instead of focusing on the subset of brain areas with interindividual variability, we focus on excluding the subset of brain areas that had a potential network misidentification in the infant cohort. Alternatively, one might be interested in focusing on the areas that are unstable across development depending on the research question at hand, which may have behavioral or clinical significance.

4.5. Implications for Precision Functional Mapping in Developmental Cohorts Using Adult Group Priors

As demonstrated in our results, on average the adult functional networks did not well represent the organization of infant FC into internally similar clusters. This observation might have important implications for research using an adult functional network model to generate individual-specific functional networks in pediatric cohorts (Hermosillo et al., 2024; Moore et al., 2024; Sun et al., 2023). Motivated by recent research findings showing that functional network topography across human individuals qualitatively differs from group-average estimates (Gordon, Laumann, Gilmore, et al., 2017; Gratton et al., 2018; Laumann et al., 2015), researchers have emphasized the importance of precision functional mapping (Gratton et al., 2022). However, reliable identification of individualized functional networks with unsupervised clustering or community detection procedures requires extended data acquisition. For example, with the Infomap algorithm (Power et al., 2011; Rosvall & Bergstrom, 2008), more than 90 minutes of data are required to achieve an average network overlap dice coefficient of > 0.75 (Gordon, Laumann, Gilmore, et al., 2017). Therefore, researchers have developed several semi-supervised methods using adult networks as priors to derive individual functional networks in data with shorter acquisition time (Cui et al., 2020; Gordon, Laumann, Adeyemo, et al., 2017; Hacker et al., 2013; Kong et al., 2019; D. Wang et al., 2015). However, these approaches generally assume that the individual functional networks are highly similar to the adult group average template. This assumption might not be suitable for developmental cohorts. As we demonstrate in this paper, the adult functional networks poorly represent the organization of the infant FC into internally coherent clusters. Two unwanted consequences might arise from this observation. First, the network templates generated by averaging the FC profiles within a poorly defined network might be noisy and inaccurate. Second, the algorithms may incorrectly force a categorical label for locations that poorly match all available networks. Future studies using adult-based priors in developmental cohorts should keep those limitations in mind and develop strategies to mitigate them.

4.6. Limitations and Future Directions

The differences in state (asleep in infants and awake in adults) may contribute to the worse fit of adult networks to infant FC. Sleep and the level of arousal are known to modify the FC structure in adults (Chang et al., 2016; Mitra et al., 2017; Tagliazucchi et al., 2012), and FC patterns in sleeping 6- and 12-month-old infants more closely resemble FC patterns in sleeping adults (Mitra et al., 2017). Based on our results from a small sample of awake fMRI in 36-to-60-month-olds, the difference in the quality of clustering between awake adult and awake infant FC using the adult Gordon networks is smaller than the difference in sleeping infant FC. Taken together with the existing literature, these results tentatively support the hypothesis that infant network clustering quality is driven by the interaction of both brain development and sleeping state.

Other differences in the acquisition and processing of the two datasets might introduce further confounds. Additionally, while we observed minimal effects of age on within-network FC, this could be due to the narrow age range of our sample (mostly between 1 and 3 years). The mixing of fMRI signals within the ill-defined areas using an adult parcellation might contribute to the low SI observed in some of those areas which could be improved with the use of infant specific areas (Tu et al., 2024).

Future studies could examine the cellular, molecular, and genetic properties of the areas that have already developed an adult-like organization in infancy to fully understand the biological underpinning of our observation. Furthermore, future studies with larger samples and well-defined behavior measures can explicitly test our deduction that the use of our area subset could improve statistical power and reproducibility for brain-wide association studies. Moreover, it is worth investigating whether the topography and diversity of thalamocortical projections would relate to the variability of functional network stability across the neocortex.

4.7. Conclusions

We found that there exists a subset of cortical areas whose FC profiles demonstrate adult-like network organization even in infants, despite the noticeable differences in FC organization between infants and adults. These areas were spatially closer than alternative areas to previously described locations of high network identity consensus in adults. Additionally, within-network FC defined with our area subset was higher in magnitude and more reliable across scans, individuals, and chronological age. We propose the use of adult networks defined by our area subset as a complementary approach of studying infant FC than using age-specific functional networks derived from data-driven methods. This would strengthen reliability, yet at the same time encourage interpretability and comparability across developmental stages. The biological basis of our observations as well as their psychopathological and behavioral impacts may become interesting topics for future research.

Author Contributions

JCT conceptualized the project. JCT and YW conducted a formal analysis. OK, OM, TKMD, CMS, DD, XW, and YW processed/curated the data. MDW, JCT, JTE were responsible for funding acquisition. JCT and MDW wrote the original draft. Everyone contributed to the review and editing of the final manuscript.

Funding

This work is supported by the CCSN fellowship from the McDonnell Center for Systems Neuroscience at Washington University School of Medicine in St. Louis to JCT and from NIH grants including EB029343 to MDW. The Baby Connectome Project was supported by NIMH R01 MH104324 and NIMH U01 MH110274.

Declaration of Competing Interests

The authors declared that they have no competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data and Code Availability

The WU 120 data can be downloaded from https://legacy.openfmri.org/dataset/ds000243/.
The BCP data can be downloaded from the National Institute of Mental Health Data Archive (NDA) at https://nda.nih.gov/edit_collection.html?id= 2848. The preprocessing scripts are available at https://github.com/DCAN-Labs/dcan-infant-pipeline. The analysis scripts used to generate the results and figures are available on https://github.com/cindyhfls/Tu-2024-GordonSubset-DCN.
The test for the difference between correlations were implemented from the function corr_rtest downloaded from MATLAB central. The Intraclass Correlation Coefficient (ICC) was calculated from the function ICC downloaded from MATLAB central.

Acknowledgement

The authors would like to thank Ari Segel for assistance in statistical analysis. The authors would also like to thank all families participated and technicians who have helped with the collection, processing, and curation of the data.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used ChatGPT in order to improve sentence structure and language precision. After using this tool/service, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

References

Ahmad, S., Wu, Y., Wu, Z., Thung, K.-H., Liu, S., Lin, W., Li, G., Wang, L., & Yap, P.-T. (2023).

Multifaceted atlases of the human brain in its infancy. *Nature Methods*, *20*(1), Article 1.

https://doi.org/10.1038/s41592-022-01703-z

Bethlehem, R. a. I., Seidlitz, J., White, S. R., Vogel, J. W., Anderson, K. M., Adamson, C., Adler, S.,

Alexopoulos, G. S., Anagnostou, E., Areces-Gonzalez, A., Astle, D. E., Auyeung, B., Ayub,

M., Bae, J., Ball, G., Baron-Cohen, S., Beare, R., Bedford, S. A., Benegal, V., … Alexander-

Bloch, A. F. (2022). Brain charts for the human lifespan. *Nature*, *604*(7906), Article 7906.

https://doi.org/10.1038/s41586-022-04554-y

Bijsterbosch, J. D., Beckmann, C. F., Woolrich, M. W., Smith, S. M., & Harrison, S. J. (2019). The

relationship between spatial configuration and functional connectivity of brain regions

revisited. *ELife*, *8*, e44890. https://doi.org/10.7554/eLife.44890

Bijsterbosch, J. D., Woolrich, M. W., Glasser, M. F., Robinson, E. C., Beckmann, C. F., Van Essen,

D. C., Harrison, S. J., & Smith, S. M. (2018). The relationship between spatial

configuration and functional connectivity of brain regions. *ELife*, *7*, e32992.

https://doi.org/10.7554/eLife.32992

Chang, C., Leopold, D. A., Schölvinck, M. L., Mandelkow, H., Picchioni, D., Liu, X., Ye, F. Q.,

Turchi, J. N., & Duyn, J. H. (2016). Tracking brain arousal fluctuations with fMRI.

Proceedings of the National Academy of Sciences, *113*(16), 4518–4523.

https://doi.org/10.1073/pnas.1520613113

Cui, Z., Li, H., Xia, C. H., Larsen, B., Adebimpe, A., Baum, G. L., Cieslak, M., Gur, R. E., Gur, R. C.,

Moore, T. M., Oathes, D. J., Alexander-Bloch, A. F., Raznahan, A., Roalf, D. R., Shinohara,

R. T., Wolf, D. H., Davatzikos, C., Bassett, D. S., Fair, D. A., … Satterthwaite, T. D. (2020).

Individual Variation in Functional Topography of Association Networks in Youth. *Neuron*,

106(2), 340-353.e8. https://doi.org/10.1016/j.neuron.2020.01.029

Dworetsky, A., Seitzman, B. A., Adeyemo, B., Neta, M., Coalson, R. S., Petersen, S. E., & Gratton,

C. (2021). Probabilistic mapping of human functional brain networks identifies regions of

high group consensus. *NeuroImage*, *237*, 118164.

https://doi.org/10.1016/j.neuroimage.2021.118164

Eggebrecht, A. T., Elison, J. T., Feczko, E., Todorov, A., Wolff, J. J., Kandala, S., Adams, C. M.,

Snyder, A. Z., Lewis, J. D., Estes, A. M., Zwaigenbaum, L., Botteron, K. N., McKinstry, R.

C., Constantino, J. N., Evans, A., Hazlett, H. C., Dager, S., Paterson, S. J., Schultz, R. T., …

Pruett, J. R., Jr. (2017). Joint Attention and Brain Functional Connectivity in Infants and

Toddlers. *Cerebral Cortex*, *27*(3), 1709–1720. https://doi.org/10.1093/cercor/bhw403

Eyre, M., Fitzgibbon, S. P., Ciarrusta, J., Cordero-Grande, L., Price, A. N., Poppe, T., Schuh, A.,

Hughes, E., O'Keeffe, C., Brandon, J., Cromb, D., Vecchiato, K., Andersson, J., Duff, E. P.,

Counsell, S. J., Smith, S. M., Rueckert, D., Hajnal, J. V., Arichi, T., … Edwards, A. D. (2021).

The Developing Human Connectome Project: Typical and disrupted perinatal functional

connectivity. *Brain*, *144*(7), 2199–2213. https://doi.org/10.1093/brain/awab118

Fair, D. A. (2020). Correction of respiratory artifacts in MRI head motion estimates.

NeuroImage, 17.

Feczko, E., Conan, G., Marek, S., Tervo-Clemmens, B., Cordova, M., Doyle, O., Earl, E., Perrone,

A., Sturgeon, D., Klein, R., Harman, G., Kilamovich, D., Hermosillo, R., Miranda-

Dominguez, O., Adebimpe, A., Bertolero, M., Cieslak, M., Covitz, S., Hendrickson, T., …

Fair, D. A. (2021). *Adolescent Brain Cognitive Development (ABCD) Community MRI*

Collection and Utilities (p. 2021.07.09.451638). bioRxiv.

https://doi.org/10.1101/2021.07.09.451638

Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., Papademetris, X., &

Constable, R. T. (2015). Functional connectome fingerprinting: Identifying individuals

using patterns of brain connectivity. *Nature Neuroscience*, *18*(11), Article 11.

https://doi.org/10.1038/nn.4135

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Flechsig, P. (1901). DEVELOPMENTAL (MYELOGENETIC) LOCALISATION OF THE CEREBRAL

CORTEX IN THE HUMAN SUBJECT. *The Lancet*, *158*(4077), 1027–1030.

https://doi.org/10.1016/S0140-6736(01)01429-5

Fornito, A., Zalesky, A., & Breakspear, M. (2015). The connectomics of brain disorders. *Nature*

Reviews Neuroscience, *16*(3), 159–172. https://doi.org/10.1038/nrn3901

Fox, M., & Greicius, M. (2010). Clinical applications of resting state functional connectivity.

Frontiers in Systems Neuroscience, *4*.

https://www.frontiersin.org/articles/10.3389/fnsys.2010.00019

Fransson, P., Skiöld, B., Horsch, S., Nordell, A., Blennow, M., Lagercrantz, H., & Åden, U. (2007).

Resting-state networks in the infant brain. *Proceedings of the National Academy of*

Sciences, *104*(39), 15531–15536. https://doi.org/10.1073/pnas.0704380104

Gao, W., Alcauter, S., Elton, A., Hernandez-Castillo, C. R., Smith, J. K., Ramirez, J., & Lin, W.

(2015). Functional Network Development During the First Year: Relative Sequence and

Socioeconomic Correlations. *Cerebral Cortex (New York, N.Y.: 1991)*, *25*(9), 2919–2928.

https://doi.org/10.1093/cercor/bhu088

Gao, W., Alcauter, S., Smith, J. K., Gilmore, J. H., & Lin, W. (2015). Development of human brain

cortical network architecture during infancy. *Brain Structure and Function*, *220*(2),

1173–1186. https://doi.org/10.1007/s00429-014-0710-3

Gao, W., Lin, W., Grewen, K., & Gilmore, J. H. (2017). Functional Connectivity of the Infant

Human Brain: Plastic and Modifiable. *The Neuroscientist*, *23*(2), 169–184.

https://doi.org/10.1177/1073858416635986

Gao, W., Zhu, H., Giovanello, K. S., Smith, J. K., Shen, D., Gilmore, J. H., & Lin, W. (2009).

Evidence on the emergence of the brain's default network from 2-week-old to 2-year-

old healthy pediatric subjects. *Proceedings of the National Academy of Sciences*,

106(16), 6790–6795. https://doi.org/10.1073/pnas.0811221106

Garcia, K. E., Robinson, E. C., Alexopoulos, D., Dierker, D. L., Glasser, M. F., Coalson, T. S.,

Ortinau, C. M., Rueckert, D., Taber, L. A., Van Essen, D. C., Rogers, C. E., Smyser, C. D., &

Bayly, P. V. (2018). Dynamic patterns of cortical expansion during folding of the preterm

human brain. *Proceedings of the National Academy of Sciences of the United States of*

America, *115*(12), 3156–3161. https://doi.org/10.1073/pnas.1715451115

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J.,

Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., & Jenkinson, M. (2013). The

minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*, *80*,

105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Gordon, E. M., Laumann, T. O., Adeyemo, B., Huckins, J. F., Kelley, W. M., & Petersen, S. E.

(2016). Generation and Evaluation of a Cortical Area Parcellation from Resting-State

Correlations. *Cerebral Cortex*, *26*(1), 288–303. https://doi.org/10.1093/cercor/bhu239

Gordon, E. M., Laumann, T. O., Adeyemo, B., & Petersen, S. E. (2017). Individual Variability of

the System-Level Organization of the Human Brain. *Cerebral Cortex*, *27*(1), 386–399.

https://doi.org/10.1093/cercor/bhv239

Gordon, E. M., Laumann, T. O., Gilmore, A. W., Newbold, D. J., Greene, D. J., Berg, J. J., Ortega,

M., Hoyt-Drazen, C., Gratton, C., Sun, H., Hampton, J. M., Coalson, R. S., Nguyen, A. L.,

McDermott, K. B., Shimony, J. S., Snyder, A. Z., Schlaggar, B. L., Petersen, S. E., Nelson, S.

M., & Dosenbach, N. U. F. (2017). Precision Functional Mapping of Individual Human

Brains. *Neuron*, *95*(4). https://doi.org/10.1016/j.neuron.2017.07.011

Gratton, C., Laumann, T. O., Nielsen, A. N., Greene, D. J., Gordon, E. M., Gilmore, A. W., Nelson,

S. M., Coalson, R. S., Snyder, A. Z., Schlaggar, B. L., Dosenbach, N. U. F., & Petersen, S. E.

(2018). Functional Brain Networks Are Dominated by Stable Group and Individual

Factors, Not Cognitive or Daily Variation. *Neuron*, *98*(2), 439-452.e5.

https://doi.org/10.1016/j.neuron.2018.03.035

Gratton, C., Nelson, S. M., & Gordon, E. M. (2022). Brain-behavior correlations: Two paths

toward reliability. *Neuron*, *110*(9), 1446–1449.

https://doi.org/10.1016/j.neuron.2022.04.018

Grayson, D. S., & Fair, D. A. (2017). Development of large-scale functional networks from birth

to adulthood: A guide to the neuroimaging literature. *NeuroImage*, *160*, 15–31.

https://doi.org/10.1016/j.neuroimage.2017.01.079

Hacker, C. D., Laumann, T. O., Szrama, N. P., Baldassarre, A., Snyder, A. Z., Leuthardt, E. C., &

Corbetta, M. (2013). Resting state network estimation in individual subjects.

NeuroImage, *82*, 616–633. https://doi.org/10.1016/j.neuroimage.2013.05.108

Hermosillo, R. J. M., Moore, L. A., Feczko, E., Miranda-Domínguez, Ó., Pines, A., Dworetsky, A.,

Conan, G., Mooney, M. A., Randolph, A., Graham, A., Adeyemo, B., Earl, E., Perrone, A.,

Carrasco, C. M., Uriarte-Lopez, J., Snider, K., Doyle, O., Cordova, M., Koirala, S., … Fair, D.

A. (2024). A precision functional atlas of personalized network topography and

probabilities. *Nature Neuroscience*, *27*(5), 1000–1013. https://doi.org/10.1038/s41593-

024-01596-5

Hill, J., Inder, T., Neil, J., Dierker, D., Harwell, J., & Van Essen, D. (2010). Similar patterns of

cortical expansion during human development and evolution. *Proceedings of the*

National Academy of Sciences, *107*(29), 13135–13140.

https://doi.org/10.1073/pnas.1001229107

Howell, B. R., Styner, M. A., Gao, W., Yap, P.-T., Wang, L., Baluyot, K., Yacoub, E., Chen, G.,

Potts, T., Salzwedel, A., Li, G., Gilmore, J. H., Piven, J., Smith, J. K., Shen, D., Ugurbil, K.,

Zhu, H., Lin, W., & Elison, J. T. (2019). The UNC/UMN Baby Connectome Project (BCP):

An overview of the study design and protocol development. *NeuroImage*, *185*, 891–905.

https://doi.org/10.1016/j.neuroimage.2018.03.049

Hu, D., Wang, F., Zhang, H., Wu, Z., Zhou, Z., Li, G., Wang, L., Lin, W., Li, G., & Consortium, U. B.

C. P. (2022). Existence of Functional Connectome Fingerprint during Infancy and Its

Stability over Months. *Journal of Neuroscience*, *42*(3), 377–389.

https://doi.org/10.1523/JNEUROSCI.0480-21.2021

Ji, J. L., Spronk, M., Kulkarni, K., Repovš, G., Anticevic, A., & Cole, M. W. (2019). Mapping the

human brain's cortical-subcortical functional network organization. *NeuroImage*, *185*,

35–57. https://doi.org/10.1016/j.neuroimage.2018.10.006

Kaplan, S., Meyer, D., Miranda-Dominguez, O., Perrone, A., Earl, E., Alexopoulos, D., Barch, D.

M., Day, T. K. M., Dust, J., Eggebrecht, A. T., Feczko, E., Kardan, O., Kenley, J. K., Rogers,

C. E., Wheelock, M. D., Yacoub, E., Rosenberg, M., Elison, J. T., Fair, D. A., & Smyser, C. D.

(2022). Filtering respiratory motion artifact from resting state fMRI data in infant and

toddler populations. *NeuroImage*, *247*, 118838.

https://doi.org/10.1016/j.neuroimage.2021.118838

Kardan, O., Kaplan, S., Wheelock, M. D., Feczko, E., Day, T. K. M., Miranda-Domínguez, Ó.,

Meyer, D., Eggebrecht, A. T., Moore, L. A., Sung, S., Chamberlain, T. A., Earl, E., Snider,

K., Graham, A., Berman, M. G., Uğurbil, K., Yacoub, E., Elison, J. T., Smyser, C. D., …

Rosenberg, M. D. (2022). Resting-state functional connectivity identifies individuals and

predicts age in 8-to-26-month-olds. *Developmental Cognitive Neuroscience*, *56*, 101123.

https://doi.org/10.1016/j.dcn.2022.101123

Kong, R., Li, J., Orban, C., Sabuncu, M. R., Liu, H., Schaefer, A., Sun, N., Zuo, X.-N., Holmes, A. J.,

Eickhoff, S. B., & Yeo, B. T. T. (2019). Spatial Topography of Individual-Specific Cortical

Networks Predicts Human Cognition, Personality, and Emotion. *Cerebral Cortex*, *29*(6),

2533–2551. https://doi.org/10.1093/cercor/bhy123

Lancaster, J. L., Glass, T. G., Lankipalli, B. R., Downs, H., Mayberg, H., & Fox, P. T. (1995). A

modality-independent approach to spatial normalization of tomographic images of the

human brain. *Human Brain Mapping*, *3*(3), 209–223.

https://doi.org/10.1002/hbm.460030305

Langs, G., Wang, D., Golland, P., Mueller, S., Pan, R., Sabuncu, M. R., Sun, W., Li, K., & Liu, H.

(2016). Identifying Shared Brain Networks in Individuals by Decoupling Functional and

Anatomical Variability. *Cerebral Cortex*, *26*(10), 4004–4014.

https://doi.org/10.1093/cercor/bhv189

Laumann, T. O., Gordon, E. M., Adeyemo, B., Snyder, A. Z., Joo, S. J., Chen, M.-Y., Gilmore, A.

W., McDermott, K. B., Nelson, S. M., Dosenbach, N. U. F., Schlaggar, B. L., Mumford, J.

A., Poldrack, R. A., & Petersen, S. E. (2015). Functional System and Areal Organization of

a Highly Sampled Individual Human Brain. *Neuron*, *87*(3), 657–670.

https://doi.org/10.1016/j.neuron.2015.06.037

Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S., Donohue,

M. R., Foran, W., Miller, R. L., Hendrickson, T. J., Malone, S. M., Kandala, S., Feczko, E.,

Miranda-Dominguez, O., Graham, A. M., Earl, E. A., Perrone, A. J., Cordova, M., Doyle,

O., … Dosenbach, N. U. F. (2022). Reproducible brain-wide association studies require

thousands of individuals. *Nature*, 1–7. https://doi.org/10.1038/s41586-022-04492-9

Marrus, N., Eggebrecht, A. T., Todorov, A., Elison, J. T., Wolff, J. J., Cole, L., Gao, W., Pandey, J.,

Shen, M. D., Swanson, M. R., Emerson, R. W., Klohr, C. L., Adams, C. M., Estes, A. M.,

Zwaigenbaum, L., Botteron, K. N., McKinstry, R. C., Constantino, J. N., Evans, A. C., …

Pruett, J. R., Jr. (2018). Walking, Gross Motor Development, and Brain Functional

Connectivity in Infants and Toddlers. *Cerebral Cortex*, *28*(2), 750–763.

https://doi.org/10.1093/cercor/bhx313

Mesulam, M. (1998). From sensation to cognition. *Brain*, *121*(6), 1013–1052.

https://doi.org/10.1093/brain/121.6.1013

Mitra, A., Snyder, A. Z., Tagliazucchi, E., Laufs, H., Elison, J., Emerson, R. W., Shen, M. D., Wolff,

J. J., Botteron, K. N., Dager, S., Estes, A. M., Evans, A., Gerig, G., Hazlett, H. C., Paterson,

S. J., Schultz, R. T., Styner, M. A., Zwaigenbaum, L., Network, T. I., … Raichle, M. (2017).

Resting-state fMRI in sleeping infants more closely resembles adult sleep than adult

wakefulness. *PLOS ONE*, *12*(11), e0188122.

https://doi.org/10.1371/journal.pone.0188122

Molloy, M. F., & Saygin, Z. M. (2022). Individual variability in functional organization of the

neonatal brain. *NeuroImage*, *253*, 119101.

https://doi.org/10.1016/j.neuroimage.2022.119101

Moore, L. A., Hermosillo, R. J. M., Feczko, E., Moser, J., Koirala, S., Allen, M. C., Buss, C., Conan,

G., Juliano, A. C., Marr, M., Miranda-Dominguez, O., Mooney, M., Myers, M.,

Rasmussen, J., Rogers, C. E., Smyser, C. D., Snider, K., Sylvester, C., Thomas, E., …

Graham, A. M. (2024). Towards personalized precision functional mapping in infancy.

Imaging Neuroscience, *2*, 1–20. https://doi.org/10.1162/imag_a_00165

Mueller, S., Wang, D., Fox, M. D., Yeo, B. T. T., Sepulcre, J., Sabuncu, M. R., Shafee, R., Lu, J., &

Liu, H. (2013). Individual Variability in Functional Connectivity Architecture of the

Human Brain. *Neuron*, *77*(3), 586–595. https://doi.org/10.1016/j.neuron.2012.12.028

Myers, M. J., Labonte, A. K., Gordon, E. M., Laumann, T. O., Tu, J. C., Wheelock, M. D., Nielsen,

A. N., Schwarzlose, R. F., Camacho, M. C., Alexopoulos, D., Warner, B. B., Raghuraman,

N., Luby, J. L., Barch, D. M., Fair, D. A., Petersen, S. E., Rogers, C. E., Smyser, C. D., &

Sylvester, C. M. (2024). Functional parcellation of the neonatal cortical surface. *Cerebral*

Cortex, *34*(2), bhae047. https://doi.org/10.1093/cercor/bhae047

Nielsen, A. N., Kaplan, S., Meyer, D., Alexopoulos, D., Kenley, J. K., Smyser, T. A., Wakschlag, L.

S., Norton, E. S., Raghuraman, N., Warner, B. B., Shimony, J. S., Luby, J. L., Neil, J. J.,

990   Petersen, S. E., Barch, D. M., Rogers, C. E., Sylvester, C. M., & Smyser, C. D. (2022).

991   Maturation of large-scale brain systems over the first month of life. *Cerebral Cortex*,

992   bhac242. https://doi.org/10.1093/cercor/bhac242

993 Ojemann, J. G., Akbudak, E., Snyder, A. Z., McKinstry, R. C., Raichle, M. E., & Conturo, T. E.

994   (1997). Anatomic Localization and Quantitative Analysis of Gradient Refocused Echo-

995   Planar fMRI Susceptibility Artifacts. *NeuroImage*, *6*(3), 156–167.

996   https://doi.org/10.1006/nimg.1997.0289

997 Petersen, S. E., & Sporns, O. (2015). Brain Networks and Cognitive Architectures. *Neuron*, *88*(1),

998   207–219. https://doi.org/10.1016/j.neuron.2015.09.027

999 Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2012). Spurious but

1000   systematic correlations in functional connectivity MRI networks arise from subject

1001   motion. *NeuroImage*, *59*(3), 2142–2154.

1002   https://doi.org/10.1016/j.neuroimage.2011.10.018

1003 Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C.,

1004   Laumann, T. O., Miezin, F. M., Schlaggar, B. L., & Petersen, S. E. (2011). Functional

1005   Network Organization of the Human Brain. *Neuron*, *72*(4), 665–678.

1006   https://doi.org/10.1016/j.neuron.2011.09.006

1007 Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014).

1008   Methods to detect, characterize, and remove motion artifact in resting state fMRI.

1009   *NeuroImage*, *84*, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Power, J. D., Plitt, M., Kundu, P., Bandettini, P. A., & Martin, A. (2017). Temporal interpolation

alters motion in fMRI scans: Magnitudes and consequences for artifact detection. *PLoS*

ONE, *12*(9), e0182939. https://doi.org/10.1371/journal.pone.0182939

Power, J. D., Schlaggar, B. L., Lessov-Schlaggar, C. N., & Petersen, S. E. (2013). Evidence for Hubs

in Human Functional Brain Networks. *Neuron*, *79*(4), 798–813.

https://doi.org/10.1016/j.neuron.2013.07.035

Raju, H., & Tadi, P. (2024). Neuroanatomy, Somatosensory Cortex. In *StatPearls*. StatPearls

Publishing. http://www.ncbi.nlm.nih.gov/books/NBK555915/

Rosvall, M., & Bergstrom, C. T. (2008). Maps of random walks on complex networks reveal

community structure. *Proceedings of the National Academy of Sciences*, *105*(4), 1118–

1123. https://doi.org/10.1073/pnas.0706851105

Rosvall, M., & Bergstrom, C. T. (2010). Mapping Change in Large Networks. *PLoS ONE*, *5*(1),

e8694. https://doi.org/10.1371/journal.pone.0008694

Rousseeuw, P. J. (1987). Silhouettes: A graphical aid to the interpretation and validation of

cluster analysis. *Journal of Computational and Applied Mathematics*, *20*, 53–65.

https://doi.org/10.1016/0377-0427(87)90125-7

Rudolph, M. D., Graham, A. M., Feczko, E., Miranda-Dominguez, O., Rasmussen, J. M., Nardos,

R., Entringer, S., Wadhwa, P. D., Buss, C., & Fair, D. A. (2018). Maternal IL-6 during

pregnancy can be estimated from newborn brain connectivity and predicts future

working memory in offspring. *Nature Neuroscience*, *21*(5), 765–772.

https://doi.org/10.1038/s41593-018-0128-y

Seitzman, B. A., Gratton, C., Laumann, T. O., Gordon, E. M., Adeyemo, B., Dworetsky, A., Kraus,

B. T., Gilmore, A. W., Berg, J. J., Ortega, M., Nguyen, A., Greene, D. J., McDermott, K. B.,

Nelson, S. M., Lessov-Schlaggar, C. N., Schlaggar, B. L., Dosenbach, N. U. F., & Petersen,

S. E. (2019). Trait-like variants in human functional brain networks. *Proceedings of the*

National Academy of Sciences, *116*(45), 22851–22861.

https://doi.org/10.1073/pnas.1902932116

Smith, S. M., Miller, K. L., Salimi-Khorshidi, G., Webster, M., Beckmann, C. F., Nichols, T. E.,

Ramsey, J. D., & Woolrich, M. W. (2011). Network modelling methods for FMRI.

NeuroImage, *54*(2), 875–891. https://doi.org/10.1016/j.neuroimage.2010.08.063

Smyser, C. D., Inder, T. E., Shimony, J. S., Hill, J. E., Degnan, A. J., Snyder, A. Z., & Neil, J. J.

(2010). Longitudinal Analysis of Neural Network Development in Preterm Infants.

Cerebral Cortex, *20*(12), 2852–2862. https://doi.org/10.1093/cercor/bhq035

Sun, L., Zhao, T., Liang, X., Xia, M., Li, Q., Liao, X., Gong, G., Wang, Q., Pang, C., Yu, Q., Bi, Y.,

Chen, P., Chen, R., Chen, Y., Chen, T., Cheng, J., Cheng, Y., Cui, Z., Dai, Z., … He, Y. (2023).

Functional connectome through the human life span (p. 2023.09.12.557193). bioRxiv.

https://doi.org/10.1101/2023.09.12.557193

Sydnor, V. J., Larsen, B., Bassett, D. S., Alexander-Bloch, A., Fair, D. A., Liston, C., Mackey, A. P.,

Milham, M. P., Pines, A., Roalf, D. R., Seidlitz, J., Xu, T., Raznahan, A., & Satterthwaite, T.

D. (2021). Neurodevelopment of the association cortices: Patterns, mechanisms, and

implications for psychopathology. *Neuron*, *109*(18), 2820–2846.

https://doi.org/10.1016/j.neuron.2021.06.016

Sylvester, C. M., Kaplan, S., Myers, M. J., Gordon, E. M., Schwarzlose, R. F., Alexopoulos, D.,

    Nielsen, A. N., Kenley, J. K., Meyer, D., Yu, Q., Graham, A. M., Fair, D. A., Warner, B. B.,

    Barch, D. M., Rogers, C. E., Luby, J. L., Petersen, S. E., & Smyser, C. D. (2022). Network-

    specific selectivity of functional connections in the neonatal brain. *Cerebral Cortex*,

    bhac202. https://doi.org/10.1093/cercor/bhac202

Tagliazucchi, E., Von Wegner, F., Morzelewski, A., Brodbeck, V., & Laufs, H. (2012). Dynamic

    BOLD functional connectivity in humans and its electrophysiological correlates. *Frontiers*

    in Human Neuroscience, *6*.

    https://www.frontiersin.org/articles/10.3389/fnhum.2012.00339

Talairach, J., & Tournoux, P. (1988). *Co-planar Stereotaxic Atlas of the Human Brain: 3-*

    dimensional Proportional System : an Approach to Cerebral Imaging. G. Thieme.

Thomason, M. E., Dassanayake, M. T., Shen, S., Katkuri, Y., Alexis, M., Anderson, A. L., Yeo, L.,

    Mody, S., Hernandez-Andrade, E., Hassan, S. S., Studholme, C., Jeong, J.-W., & Romero,

    R. (2013). Cross-Hemispheric Functional Connectivity in the Human Fetal Brain. *Science*

    Translational Medicine, *5*(173). https://doi.org/10.1126/scitranslmed.3004978

Tooley, U. A., Latham, A., Kenley, J. K., Alexopoulos, D., Smyser, T. A., Nielsen, A. N., Gorham, L.,

    Warner, B. B., Shimony, J. S., & Neil, J. J. (2024). Prenatal environment is associated with

    the pace of cortical network development over the first three years of life. *Nature*

    Communications, *15*(1), 7932.

Truzzi, A., & Cusack, R. (2023). The development of intrinsic timescales: A comparison between

    the neonate and adult brain. *NeuroImage*, *275*, 120155.

    https://doi.org/10.1016/j.neuroimage.2023.120155

Tu, J., Myers, M., Li, W., Li, J., Wang, X., Dierker, D., Day, T., Snyder, A. Z., Latham, A., Kenley, J.

K., & others. (2024). Early life neuroimaging: The generalizability of cortical area

parcellations across development. *BioRxiv : The Preprint Server for Biology*, 2024–09.

Turk, E., Heuvel, M. I. van den, Benders, M. J., Heus, R. de, Franx, A., Manning, J. H., Hect, J. L.,

Hernandez-Andrade, E., Hassan, S. S., Romero, R., Kahn, R. S., Thomason, M. E., &

Heuvel, M. P. van den. (2019). Functional Connectome of the Fetal Brain. *Journal of*

Neuroscience, *39*(49), 9716–9724. https://doi.org/10.1523/JNEUROSCI.2891-18.2019

Van Essen, D. C., Drury, H. A., Dickson, J., Harwell, J., Hanlon, D., & Anderson, C. H. (2001). An

Integrated Software Suite for Surface-based Analyses of Cerebral Cortex. *Journal of the*

American Medical Informatics Association, *8*(5), 443–459.

https://doi.org/10.1136/jamia.2001.0080443

Van Essen, D. C., Glasser, M. F., Dierker, D. L., Harwell, J., & Coalson, T. (2012). Parcellations and

Hemispheric Asymmetries of Human Cerebral Cortex Analyzed on Surface-Based

Atlases. *Cerebral Cortex*, *22*(10), 2241–2262. https://doi.org/10.1093/cercor/bhr291

Wang, D., Buckner, R. L., Fox, M. D., Holt, D. J., Holmes, A. J., Stoecklein, S., Langs, G., Pan, R.,

Qian, T., Li, K., Baker, J. T., Stufflebeam, S. M., Wang, K., Wang, X., Hong, B., & Liu, H.

(2015). Parcellating cortical functional networks in individuals. *Nature Neuroscience*,

18(12), Article 12. https://doi.org/10.1038/nn.4164

Wang, F., Zhang, H., Wu, Z., Hu, D., Zhou, Z., Girault, J. B., Wang, L., Lin, W., & Li, G. (2023). Fine-

grained functional parcellation maps of the infant cerebral cortex. *ELife*, *12*, e75401.

https://doi.org/10.7554/eLife.75401

Wen, X., Hsu, L., Lin, W., Zhang, H., & Shen, D. (2020). Co-evolution of Functional Brain Network

at Multiple Scales during Early Infancy. *ArXiv:2009.06899 [Cs, q-Bio]*.

http://arxiv.org/abs/2009.06899

Wen, X., Zhang, H., Li, G., Liu, M., Yin, W., Lin, W., Zhang, J., & Shen, D. (2019). First-year

development of modules and hubs in infant brain functional networks. *NeuroImage*,

185, 222–235. https://doi.org/10.1016/j.neuroimage.2018.10.019

Wheelock, M. D., Hect, J. L., Hernandez-Andrade, E., Hassan, S. S., Romero, R., Eggebrecht, A. T.,

& Thomason, M. E. (2019). Sex differences in functional connectivity during fetal brain

development. *Developmental Cognitive Neuroscience*, *36*, 100632.

https://doi.org/10.1016/j.dcn.2019.100632

Wig, G. S. (2017). Segregated Systems of Human Brain Networks. *Trends in Cognitive Sciences*,

21(12), 981–996. https://doi.org/10.1016/j.tics.2017.09.006

Yates, T. S., Ellis, C. T., & Turk-Browne, N. B. (2023). Functional networks in the infant brain

during sleep and wake states. *Cerebral Cortex*, bhad327.

https://doi.org/10.1093/cercor/bhad327

Yeo, B. T. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman,

J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., & Buckner, R. L. (2011).

The organization of the human cerebral cortex estimated by intrinsic functional

connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165.

https://doi.org/10.1152/jn.00338.2011

1115
1116

Supplementary Materials

Supplementary Results
S1. Silhouette index in individual sessions

We calculated the silhouette index in group-average data because individual sessions are noisy and have a relatively short acquisition time. However, our group level results were consistent with results obtained by calculating silhouette index in individual sessions (Supplementary Figures 3-4). Specifically, the average SI (-0.0054 $\pm$ 0.0714) was not different from zero in individual adult sessions (two-tailed t-test, p = 0.41) for adult ("Gordon") networks. On the other hand, the average SI (-0.0859 $\pm$ 0.0318) was significantly smaller than zero in individual adult sessions for infant ("Kardan") networks (two-tailed t-test, p<0.001). In addition, the average SI (-0.0636 $\pm$ 0.0492) was smaller than zero in individual infant sessions (two-tailed t-test, p < 0.001) for adult("Gordon") networks. On the other hand, the average SI (0.0492 $\pm$ 0.0472) was significantly greater than zero in individual adult sessions for infant ("Kardan") networks (two-tailed t-test, p<0.001).

S2. Silhouette index of adult networks in infant FC with the mean in all alternative networks

By default, the silhouette index compares the current network to the best alternative network, which also depends on the quality of alternatives. However, other researchers have chosen to use a similar metric that compares the average within-network similarity to the average between-network similarity across all alternative networks, rather than just the best alternative (Ji et al., 2019). This approach tends to be less conservative and generally results in a higher silhouette index when calculated in this manner.

When the SI was calculated using the mean in all alternative networks rather than the mean of the best alternative network, they were still moderately correlated with the SI reported in the main results (Pearson's r = 0.74, p <0.001). However, since the mean of similarity to all alternative networks (especially to the ones spatially distant from the area in question) would tend to be lower than the best alternative, the SI is positively shifted with almost all parcels having SI > 0 (Supplementary Figure 15).

S3. Age effect on within-network (Gordon networks) FC is smaller in magnitude with our area subset than with all areas

To test the hypothesis that our area subset has relatively stable within-network FC across chronological age in infants, we compared the age effect on within-network FC when the networks include only our area subset versus all areas. The age effect of within-network FC was quantified with a Spearman's correlation (ρ). The significance of the difference between the correlation between chronological age and within-network FC in our area subset versus all areas is calculated with a Z-test on Fisher-Z-transformed r values.

We additionally examined the within-network FC in infants across chronological age. We computed within-network FC across age using full versus our area subset. For the eight networks that were partially retained, five networks demonstrated a significant correlation between within-network FC and age (p < 0.05, Spearman's ρ): the within-

network Aud, SMN hand and Vis networks were negatively correlated with age and the
within-network FC in DAN and the FPN were positively correlated with age. The age
effect was greater in magnitude with the full set of areas (Figure 3A) than with only the
partially retained areas (Figure 3B) for the SMN hand network, although not significant
when comparing the Fisher-Z-transformed ρ values (Z = 1.588, one-sided p = 0.056).
Similar results were found for other networks, where the age effect was less negative
for Aud, SMN hand and Vis networks, and less positive for DAN and FPN, but none of
them had a significant (p < 0.05) Z-test. To examine the robustness of our result to the
selection of data samples, we generated 1000 bootstrapped samples of the infant
sessions. We found that the sign of the difference was consistent across bootstrap
samples (i.e., on average the networks using our area subset was less correlated with
age than all areas) (Figure 3C). The mean and 95% confidence interval for the
bootstrap showed a mean difference in Fisher-Z-transformed ρ values for full versus
subset was -0.1139 [-0.1721,0.0129] for Aud, -0.1386 [-0.1684, -0.0814] for SMN hand,
0.0020 [-0.0887, 0.0348] for Vis, -0.0205 [-0.0120, 0.1439] for DAN and 0.0089 [0.0071,
0.0511] for FPN (Figure 3C).
1179
S4. Group consistency and differential power of FC edges
Prior studies suggested that it was possible to identify individuals using FC in
infants from the BCP dataset (Hu et al., 2022; Kardan et al., 2022). To assess which FC
edges (i.e. connections between a pair of areas) are more consistent across individuals
versus distinct across individuals, we calculated the group consistency (ϕ) and
differential power (DP) measures (Finn et al., 2015). We aim to describe the distribution
of highly consistent edges and highly differentiating edges with respect to adult and
infant network models. For this analysis, we only use the one session from each of the
115 unique subjects with at least 13.2 min low-motion data. Given two sets of
connectivity $[X_i^{R1}]$, $[X_i^{R2}]$ obtained from the two resting scan windows ($R1$ and $R2$) after
z-score normalization, the edgewise product vector φ_i was computed as
1191
1192 $$\varphi_i\ (e) =\ X_i^{R1}(e) * X_i^{R2}(e), e = 1, \dots, M\ (Equation\ 2)$$
1193
where *i* indexed the subject, *e* indexed the edge, and *M* indexed the total number of FC
edges. The sum of φ_i over all edges is the correlation between $[X_i^{R1}]$, $[X_i^{R2}]$. The group
consistency ϕ was computed as the mean of φ_i across all subjects. We defined the
edges with the top 10% ϕ values to be "highly consistent".
Similarly, the edgewise product vector φ_{ij} was calculated between patterns from
different subjects, for example:
$$\varphi_{ij}\ (e) =\ X_i^{R1}(e) * X_j^{R2}(e), e = 1, \dots, M, i \neq j\ (Equation\ 3)$$
$$P_i(e) =\ P\big|\varphi_{ij}\ (e) > \varphi_{ii}\ (e)\ or\ \varphi_{ji}\ (e) > \varphi_{ii}\ (e)\big|\ (Equation\ 4)$$
$$DP\ (e) = \sum_i\{-\ln\ (P_i(e))\}\ (Equation\ 5)$$
1203
We defined the edges with the top 10% DP values as "highly differentiating".
Consistent with the findings in previous literature, we observed that a large
percentage (~50%) of FC edges in the within-network blocks tend to be highly
consistent. On the other hand, much fewer FC edges in between-network blocks (~6%)

were highly consistent (Supplementary Figure 14; Supplementary Table 3). The
sensorimotor networks especially had a large proportion of highly consistent within-
network FC edges (Supplementary Table 4). Moreover, using adult networks defined by
our area subset, the percentage of highly consistent edges within networks increased
substantially for all eight partially retained networks (Supplementary Table 4), indicating
that the adult network spanned by our area subset over-represented areas with highly
consistent FC between them.
1215   On the other hand, within-network blocks tend to have only a slightly larger
1216   percentage of highly differentiating FC edges (~15%) than between-network blocks
1217   (~10%) (Supplementary Table 5-6), with both increased and decreased proportion of
1218   highly differentiating edges when using our area subset instead of all areas.
1219

Supplementary Figure 1. *Distribution of age and sex of individual infants in the BCP dataset.*
A) The age time points of 181 infants ordered by sex. B) The count of number of individuals with 1-6 longitudinal points.

Supplementary Figure 2. *Dice overlap between Adult Networks ("Gordon") and Infant Networks ("Kardan").* Network abbreviations: auditory (Aud), cingulo-opercular (CON), parietal memory (PMN), default mode (DMN), dorsal attention (DAN), fronto-parietal (FPN), retrosplenial temporal (RTN), somatomotor hand (SMN hand), somatomotor mouth (SMN mouth), salience (Sal), and ventral attention (VAN), visual (Vis), somatomotor (SMN), temporal (Tem), posterior frontoparietal (pFPN), posterior default mode (pDMN), lateral visual (lVis), medial visual (mVis), anterior fronto-parietal (aFPN), anterior default mode (aDMN).

Supplementary Figure 3. *Silhouette index (SI) of adult and infant networks on individual adults' FC.* A) SI across adult networks ("Gordon", 286 areas). B) SI across infant networks ("Kardan", 328 areas). C) average SI of adult and infant networks across areas on individual adults' FC. *** p < 0.001 in paired t-test. D) Pearson's correlation of SI of adult networks on group average FC and the mean of SI on individual FC across 286 areas. E) Pearson's correlation of SI of infant networks on group average FC and the mean of SI on individual FC across 328 areas.

Supplementary Figure 4. *Silhouette index (SI) of adult and infant networks on individual infants' FC.* A) SI across adult networks ("Gordon", 286 areas). B) SI across infant networks ("Kardan", 328 areas). C) average SI of adult and infant networks across areas on individual infants' FC. *** *p* < 0.001 in paired t-test. D) Pearson's correlation of SI of adult networks on group average FC and the mean of SI on individual FC across 286 areas. E) Pearson's correlation of SI of infant networks on group average FC and the mean of SI on individual FC across 328 areas. Sessions in A and B are sorted by increasing age from left to right.

BCP awake 36-60 months,
N = 15 (49.9 ± 7.4 months)

BCP asleep 36-60 months,
N = 14 (43.5 ± 8.8 months)

1225
1226

Supplementary Figure 5. *Awake V.S. sleeping infant FC organized by adult ("Gordon") and infant ("Kardan") networks.* A) The average FC for 15 BCP sessions sorted by adult networks ("Gordon"). B) The average FC for 15 BCP sessions sorted by infant networks ("Kardan"). C) Average silhouette index for individual sessions. D-F) Same as A-C, but for 14 BCP sessions also in approximately the same age range (36-60 months). *** *p* < 0.001 in paired t-test.

Supplementary Figure 6. *Infant FC sorted by the Tu (326) 12 and 19 networks.* A) The average FC for 313 BCP sessions sorted by Tu (326) 12 networks. B) The average FC for 313 BCP sessions sorted by Tu (326) 19 networks. C-D) Silhouette index for each area parcel for A-B. E-F) Average silhouette index for individual sessions. *** *p* < 0.001 in paired t-test.

frequency

0   0.5   1

Supplementary Figure 7. *Frequency of SI > 0 across 1000 bootstraps.*

Supplementary Figure 8. *Adult FC using our area subset.* A) The sorted average FC in adults with our area subset. B) The average within-network FC with all areas (left) versus our area subset subset (right) across sessions.

1233

Supplementary Figure 9. *Within-network FC difference (All – Subset) across eight partially-retained networks.*

Supplementary Figure 10. *Correlation between age and within-network FC in using our area subset versus all areas.* A) Scatter plot of within-network FC versus age for SMN hand network using all areas. B) Scatter plot of within-network FC versus age for SMN hand network using our area subset. C) The within-network FC for three networks is negatively correlated with age (Aud, SMN hand, Vis), and the within-network FC for two networks is positively correlated with age (DAN, FPN). The x-axis is the Fisher-Z-transformed Spearman's correlation (ρ) between within-network FC using all areas and age. The y-axis is the Fisher-Z-transformed Spearman's correlation (ρ) within-network FC using our area subset and age). Each data point represents a bootstrap sample of sessions (N = 1000). Red line shows the line of least-squared fit in A-B and the line of identity in C.

Supplementary Figure 11. *A scatter plot between chronological age in years and average SI for individual BCP sessions.* A) infant networks ("Kardan"). B) adult networks ("Gordon"). C) Difference in infant networks ("Kardan") and adult networks ("Gordon").

Supplementary Figure 12. *Moving average analysis (Figure 4A) adding the results using infant networks from an independent dataset.*

Supplementary Figure 13. *Distance between our area subset and alternative areas to the 153 Dworestky high consensus ROI.*

Supplementary Figure 14. *Fraction of high consistency (φ) and high differential power (DP) edges (top 10%) across (A-B) adult networks ("Gordon"), (C-D) adult networks ("Gordon" Subset), (E-F) infant networks ("Kardan").*

1257

Supplementary Figure 15. *Correlation between silhouette index calculated with the best network or with all alternative networks.*

Supplementary Table 1			
	Cohort	BCP	WashU 120
Acquisition	Location	University of Minnesota	Washington University in St. Louis
	Scanners	Siemens Prisma 3T Scanner	Siemens 3T Trio Tim
	Headcoil	32-channel	12-channel
	Sequence type	Gradient-echo EPI	Gradient-echo EPI
	Resolution (BOLD)	2mm isotropic	4 mm isotropic
	Phase encoding direction	AP+PA	AP
	TR (s)	UMN - 0.72 (N = 70), 0.8 (N = 107)	2.5
	TE (ms)	37	27
	Resolution (T1)	0.8 mm	1 mm
	Multi-band factor	8	N/A
	State	Natural Sleep	Fixation on crosshair
	Average clean data frames		14.0 min
	Frames per BOLD run	420	120
	BOLD runs	2-4	2
Processing	Processing pipeline version	DCAN-Infant (0.0.22)	adult EPI (BOLD) preprocessing pipeline using the 4dfp tool suite
	Distortion correction	ANTs SyN registration	None
	Bias field correction	N4 method	None
	Denoising	Respiratory notch filter, demean, detrend, 24 parameters nuisance regression, remove FD>0.3mm to apply bandpass filtering (0.008-0.09 Hz) then interpolate the missing frames	Demeaning and detrending, , multiple regression including: whole-brain, ventricular and white matter signals, and motion regressors derived by Volterra expansion (Friston et al. 1996), and a band-pass filter (0.009 Hz < f < 0.08Hz).
	Scrubbing threshold	filtered FD<0.2mm, outlier (across-vertex STD on low FD frames >3MAD of the median of all frames)	FD<0.2mm, at least 5 consecutive low-motion frames
	Tissue Segmentation	ANTs Joint Label Fusion was performed using a set of ALBERT atlases with segmentations generated and manually corrected by DCAN for ages 0-5 months old. For older ages, the atlases used for JLF were a set of 10 ABCD subjects for which we generated segmentations. Manual	FreeSurfer's default recon-all processing pipeline (version 5.0)

		curation of tissue segmentation was performed where necessary.	
	Surface reconstruction & registration	Modified FreeSurfer reconstruction (no hires, aseg from JLF, adjusted class means of tissue to fit T1w contrasts), registered to fs_LR32k using spherical registration (with MSMsulc).	FreeSurfer's default recon-all processing pipeline (version 5.0)
	BOLD data geodesic smoothing	σ = 2.55 mm	σ = 2.55 mm
Quality Control			
		BrainSwipes crowdsource ratings with average aggregated passing rate >75% for anatomical or functional images + manual screening.	N/A
	Data availability	https://nda.nih.gov/edit_collection.html?id=2848	https://legacy.openfmri.org/dataset/ds000243/

1258

Supplementary Table 2

x	y	z
-18.8	-48.7	65
-51.8	-7.8	38.5
-18.4	-85.5	21.6
-47.2	-58	30.8
-38.1	48.8	10.5
-55.9	-47.7	-9.3
-14.4	-57.8	18.4
-8.8	-49.8	4.2
-11.3	-83.2	3.9
-1.7	-17.7	39.1
-10	33.9	21.5
-10.7	-47.5	60.3
-15.6	-33.1	66.1
-10.9	-29.3	69.5
-6.6	-20.4	74.2
-10.8	-41.1	64.9
-5	-28.2	60.4
-5.4	-15.9	48.8
-35.8	-29.7	54.5
-41.5	-12.5	50.4
-42.1	-4.5	47.3
-27.3	1.9	52.9
-19.8	6.4	55.7
-19.5	30.1	45.5
-36.8	-22.8	61.9
-20.5	-24.9	64.5
-23.4	-13.8	64.2
-17.2	-8.6	67.9
-28.6	-44.7	61.7
-31.1	-48.9	47.1
-42.9	-45	43
-51.5	-11.9	29.7
-51.7	-30.9	39.9
-27.5	-37.2	61.4
-47.2	-31.4	54.8
-46.1	-17.8	52.7
-44.8	-54	14.6
-51.6	-55.9	11.4
-48.1	-40	2.4

-46.3	-41.4	25.9
-52.7	-20.6	5.4
-58.7	-29.9	11.1
-40.6	-38.3	14.5
-38.7	-16	-5.3
-50	20.8	10.6
-37.7	2.9	11.7
-40.3	50.4	-4.8
-32.5	17.2	-7.8
-44.3	33.2	-7.2
-45.4	28.8	0.8
-20.4	-64.6	51.4
-34.1	-61	42.4
-31.3	-84.2	9
-34.2	-86.6	-0.5
-46.2	-57.7	-7.9
-55.1	-32.3	23
-43	19.4	33.5
-40.2	23.6	23.3
-48.6	7.5	11.1
-5.9	54.8	-11.3
-6.8	38.2	-9.4
-33.8	-33.2	-15.4
-28.8	-58.8	-9.1
-34.4	-63.9	-15.7
-34.3	-43.8	-21.6
-5.4	-88	18.6
-8.6	-77.5	-3.5
-22.6	-81.7	-11.7
-22.5	-37.1	-15
-15.9	48.6	37.2
-19.5	56.3	27.5
-21.3	63.1	1.9
-28.6	50.9	10.1
-6.5	54.7	18.1
-15.7	64.7	13.7
-26.2	26.6	38.8
-29.3	16.8	50.7
-41.7	16.1	47.5
-54.4	-1.4	-0.7
-59	-18	-3

20.8	-48.2	66.1
49.6	-7.4	36.1
22	-84.6	23.7
47.9	-42.5	41.5
38.1	45.9	7.7
59.7	-41	-10.9
13.8	-54.1	10.9
15.5	-74.1	9.4
6.7	5	55.9
8.4	34.7	22.6
3	-19.6	37.9
8.8	10.8	45.9
16.5	-32.8	67.7
4.8	-27.1	64.8
11.9	-40.7	67
5.1	-17.1	51.6
6.8	-8.1	50.9
42.3	-11	47.3
42.5	-2.3	47.2
29.2	1.9	52.4
21.9	21	46.2
38.1	-22.4	60.3
19.7	-25	65.2
12.4	-28.3	69.6
29.2	-13.5	64.2
17	-16.9	70.9
20.9	-6.4	65
29.5	-42.5	60.4
38.8	-42.6	40.4
53.9	-8.3	26.1
28	-34.8	63.1
39.2	-34.6	57.5
37.3	-25.9	50.9
47.8	-15.1	49.3
48.9	-53	28.6
57.5	-45.3	9
60.9	-38.7	1.7
54.9	-27	29.6
57.1	-17	-2.6
53.8	-15.8	5.2
47.4	-39.6	13.2

45.5	-37.3	3.4
48.5	-26.5	-0.1
60	-25.2	10.2
38.8	-14.4	-5
42.8	48.3	-5.1
45.2	30.7	-5.6
30.6	22.8	-4.7
7.7	-85.6	31.6
35.4	-77.1	21.1
41.5	-53.5	44
33.5	-48.2	49.4
31.7	-85.7	2.4
43.8	-67.2	2
57	-53.8	-1.1
37.8	28.7	35.6
41.8	29.1	21.6
50.1	3	3.9
38.6	18.8	25.5
28.4	57	-5.1
4.8	65.1	-7.1
7.2	48.4	-10.1
34.6	-35.6	-12.3
34.6	-23.9	-20.4
26.9	-69.1	-6.6
34.9	-44	-20
13.8	-92.3	14.7
10.5	-73.8	-1.5
20.4	-87.3	-6.6
5.1	-80.2	23.1
24.5	-36.2	-13.2
21	32.8	42.1
21.4	42.8	35.1
23.5	59.1	4.9
30.9	52.2	9.9
8.2	53.8	14
5.9	54.9	29.4
13.8	46.7	42.1
6.8	44.5	34.8
30.6	18.9	48.7
42.4	19.5	48.2
38.9	9.6	42.7

39.7	-22.5	2.6
55.8	2	-2
57.1	-6.3	-7.7
46.6	-21.5	-8.5

1259

Supplementary Table 3. Percentage of highly consistent (ϕ) edges across different network assignment schemes

	Gordon All	Gordon Subset	Kardan
within-network	44.1%	63.1%	49.5%
between-network	8.3%	6.5%	5.4%

Supplementary Table 4. Percentage of highly consistent (ϕ) edges across different network assignment schemes. The eight partially retained networks were bolded and had an asterisk.

within-network	Gordon All	Gordon Subset
Aud*	41.30%	84.21%
CON*	35.64%	61.73%
PMN	40.00%	/
DMN*	43.17%	59.07%
DAN*	31.45%	65.17%
FPN*	43.12%	50.84%
RTN	100%	100%
SMN hand*	61.45%	72.82%
SMN mouth	89.29%	89.29%
Sal	33.33%	33.33%
VAN*	38.34%	60.94%
Vis*	45.89%	65.17%

Supplementary Table 5. Percentage of high differential power (*DP*) edges across different network assignment schemes.

	Gordon All	Gordon Subset	Kardan
within-network	16.50%	14.73%	12.79%
between-network	10.81%	10.98%	9.93%

Supplementary Table 6. Percentage of high differential power (*DP*) edges across different network assignment schemes. The eight partially retained networks were bolded and had an asterisk.

within-network	Gordon All	Gordon Subset
Aud*	7.97%	10.53%
CON*	8.33%	4.94%
PMN	50.00%	/
DMN*	10.73%	11.59%
DAN*	30.24%	47.19%
FPN*	15.58%	13.91%
RTN	7.14%	7.14%
SMN hand*	17.92%	15.82%
SMN mouth	21.43%	21.43%
Sal	0%	0%
VAN*	14.62%	12.02%
Vis*	24.97%	18.43%