Protein degradation shapes developmental tempo in mouse and human neural progenitors.

Shota Nakanoh[1*], Despina Stamataki[2*], Lorena Garcia-Perez[2], Chiara Azzi[1], Hayley L Carr[3], Alexandra Pokhilko[1], Lu Yu[4], Steven Howell [2], Mark Skehel[2], David Oxley [4], Simon Andrews[3], James Briscoe[2#], Teresa Rayon[1,5#]

[1] Epigenetics Programme, Babraham Institute, Cambridge, UK.
[2] The Francis Crick Institute, London, UK.
[3] Bioinformatics, Babraham Institute, Cambridge, UK
[4] Mass Spectrometry, Babraham Institute, Cambridge, UK
[5] Centre for Trophoblast Research; Cambridge Stem Cell Institute, University of Cambridge, Cambridge, UK.
[*] These authors contributed equally.
[#] Authors for correspondence: james.briscoe@crick.ac.uk; teresa.rayon@babraham.ac.uk

ABSTRACT

The speed of embryonic development varies considerably between mammalian species, yet the underlying molecular mechanisms remain poorly understood. To investigate the basis for species-specific developmental tempo, we performed a comprehensive comparative analysis of protein dynamics in mouse and human neural progenitors (NPs). Through a combination of targeted protein labelling, quantitative mass spectrometry, and protein depletion with self-labeling tags, we demonstrate that protein degradation is a key driver of tempo differences between mouse and human NPs. We observe a systematic 1.5-fold increase in protein half-lives in human NPs compared to mouse, independent of cellular compartment or protein function. This difference persists in post-mitotic neurons, indicating active degradation as the primary mechanism. Proteasomal activity is also ~1.5-fold higher in mouse NPs, consistent with upregulation of proteasome-associated proteins. Importantly, increasing the rate of proteolytic degradation of a key transcriptional repressor in neural progenitors accelerates the expression of its target gene. Despite differences in degradation rates, protein synthesis rates are similar between species, resulting in higher protein content in human NPs. Our findings highlight the central role of protein degradation in controlling developmental tempo and provide insight into the molecular basis of evolutionary changes in developmental timing across species.

INTRODUCTION

The developmental mechanisms governing embryogenesis are remarkably conserved across mammalian species, yet the pace at which these programmes unfold is species-specific[1–4]. These species differences in developmental timing are recapitulated in vitro using stem cell models, which has allowed investigations into the mechanisms controlling tempo[2,3,5–8]. Previous work implicated protein turnover, the balance of protein synthesis and degradation, as an important contributor to developmental tempo[5,6]. Specifically, reduced rates of protein degradation corresponded to slower tempos in human neural progenitors compared to mouse. Similarly, the pace of the segmentation clock during the elongation of the vertebrate axis correlates with the rate of protein turnover in a variety of mammalian species[6,9].

While these findings suggest protein turnover is crucial for tempo control, several questions remain unresolved. Regulation of the production and degradation machineries and their

dynamics during development are poorly understood. More precisely, a systematic
characterization of the similarities and differences in protein half-lives between species is
lacking, and it remains to be determined whether overall abundance of proteins in embryos
differs between species due to species-specific protein turnover dynamics, or compensatory
mechanisms adjust protein synthesis rates to achieve equivalent steady-state protein levels.
Finally, whether manipulating the stability of individual proteins affects the pace of
development is unclear.
56
Here, we address these questions by quantifying protein production, degradation, and
abundance in equivalent mouse and human neural progenitors (NPs). Combining targeted
protein labelling, high-resolution quantitative mass spectrometry, and protein depletion with
self-labeling tags, we systematically identify species differences in protein turnover rates and
establish the underlying mechanisms. Specifically, we elucidate a role for active protein
degradation independent of dilution via cell division to tempo divergence. We also provide
evidence to support a role for proteolytic degradation in dictating developmental tempo by
decreasing the stability of a key regulatory protein within the gene regulatory network
governing motor neuron formation in the neural tube.
66
67 Collectively, our work provides a comprehensive comparative analysis of protein dynamics
68 between two mammalian species. Our findings highlight the centrality of protein degradation
69 in controlling developmental tempo. This has broad implications for understanding
70 phenotypic diversity across evolution.
71
RESULTS
SNAP Tagged OLIG2 demonstrates differences in protein turnover between mouse
and human neural progenitors
The balance between protein synthesis and degradation (protein turnover) determines
protein levels in cells. Having observed global differences in protein decay between mouse
and human neural progenitors (NPs)[5], we set out to test if similar differences were observed
for specific proteins. To this end we focused on OLIG2, an essential transcription factor for
motor neuron differentiation[10,11]. To obtain data on the turnover for OLIG2 in mouse and
human NPs, we tagged the C-terminus of endogenous OLIG2 in mouse and human
embryonic stem cells with a HA tag and a SNAP tag, a self-labelling protein tag that enables
the covalent labelling by substrates[12].
83
We confirmed that the differentiation dynamics of OLIG2::HA::SNAP (OLIG2-SNAP) targeted
cells followed the expected pattern for OLIG2 in mouse and human NPs, with OLIG2
increasing upon Retinoic Acid (RA) and Smoothened Agonist addition (SAG)[5]. We detected
OLIG2 positive cells at the expected days of differentiation, and there was a good correlation
between OLIG2 antibody and HA tag (Fig. S1a). This indicated that tagging does not affect
OLIG2 dynamics or function in mouse or human NPs. Next, we tested the specificity of the
SNAP fluorescent substrate SNAP 647-SiR. We incubated human NPs from day 4 or day 6
of differentiation with SNAP or HALO ligands conjugated to a 647 fluorophore. We detected
specific labelling with the SNAP 647-SiR ligand; the HALO-JF647 substrate did not cross-
react with the hOLIG2-SNAP line (Fig. S1 b).
94
95 Having established ligand specificity, we then determined the half-life of OLIG2 in mouse
96 and human NPs. First, we cultured NPs for 1 hour in media containing SNAP 647- SiR

ligand. This was then replaced with media containing SNAP-ligands without 647 fluorophore or SNAP-cell block, a cell permeable and non-fluorescent compound that blocks the reactivity of the SNAP-tag. We observed similar trends in the decay of OLIG2 intensity in human NPs at day 4 and day 6 of differentiation (Fig. S1c), suggesting similar protein decay rates for OLIG2 during differentiation. Comparative SNAP pulse-chase analysis indicated the OLIG2 half-life was ~1.4-fold longer in human compared to mouse (mOLIG2 t1/2 = 2.31 ± 0.43h; hOLIG2 t1/2 = 3.21 ± 0.30h) (Fig. 1a). These data corroborate an increased OLIG2 stability in human compared to mouse[5].

We next determined OLIG2 production rates for mouse and human. SNAP ligands bind to mature proteins (Fig. 1b). To distinguish between old OLIG2 and newly synthesized molecules, we cultured mouse and human neural progenitors with SNAP-cell block to obstruct all pre-existing OLIG2 molecules present in cells. NPs were then switched into media containing SNAP 647-ligand and samples collected at intervals over the following two hours (Fig. 1b). We observed similar accumulation of labelled OLIG2 in mouse and human NPs (Fig. 1c). Quantifications of the geometric mean intensities for the ligand at various time points indicated that OLIG2 production rates in human NPs were slightly lower in human in comparison to mouse at 15min and 30min, but not markedly different between species over the course of 120 minutes (Fig. 1b).

No differences in global protein translation rates between mouse and human neural progenitors

We next set out to compare global protein production rates and abundances in mouse and human NPs. We performed metabolic labelling of nascent proteins by adding the methionine analog L-azidohomoalanine (AHA)[13] to media, and determined the rate of incorporation over one hour. We treated cells with AHA and collected mouse and human NPs after 15, 30 or 60 minutes of treatment. The production rate appeared similar in mouse and human NPs (Fig. 1d). To confirm equivalent production rates between mouse and human NPs, we measured puromycin incorporation rates[14]. While both AHA and puromycin incorporate in nascent polypeptides, puromycin blocks protein synthesis and causes premature termination, with the resulting truncated polypeptides often failing to fold properly[15,16]. We treated mouse and human NPs with various concentrations of puromycin to identify the lowest concentration that allowed sufficient labelling for quantification. Puromycin concentrations of 0.1 mg/ml were sufficient to detect nascent protein in mouse and human NPs without triggering a stress response, as measured by levels of EIF2A phosphorylated at Ser51[17] (Fig. S1d-f). Side-by-side puromycin incorporation assays in mouse and human NPs showed similar rates of protein production (Fig. 1e). The puromycin results were consistent with the measurements of AHA and confirmed that the rates of production of newly synthesized proteins are similar in mouse and human NPs.

Similar rates of protein production but faster degradation rates in mouse compared to human implies that the amount of protein in mouse NPs should be lower than in human NPs. To test this, we measured protein content using bicinchoninic acid assay (BCA)[18] and CellTrace[19] as an orthogonal approach. The BCA assay confirmed a difference of ~1.6-fold in protein content between mouse and human cells (mouse 37.7 ± 8.53 pg/cell; human 61.3 ± 12.4 pg/cell) (Fig. 1f). Similarly, CellTrace treatment, which covalently attaches succinimidyl ester groups to amine groups in proteins, indicated increased protein levels in human NPs compared to mouse (Fig. S1g). Together, these results suggest little if any differences in

145 protein synthesis, yet a ~1.6- fold increase in protein content, in human NPs compared to
146 mouse.
147
Global proteomic differences associated to protein degradation in mouse and human
neural progenitors
Given the observed differences in protein abundance, we asked if specific sets of proteins
showed a greater species difference than others. We performed data-independent
acquisition mass spectrometry (DIA-MS) of mouse and human NPs, and calculated protein
copy number per cell using the 'proteomic ruler' approach[20] (see Methods) (Fig. S2a-e). We
identified 5718 orthologous proteins between mouse and human (Table S1), which we used
for differential expression analysis. We found 1791 differentially expressed proteins with ≥ 2-
fold change and FDR ≤ 0.05 (Table S2): 997 proteins upregulated in human and 794 in
mouse (Fig. 2a). The differentially expressed proteins included lowly and highly abundant
proteins (Fig. S2f).
159
Many of the differentially expressed proteins were related to the ubiquitin–proteasome
system and ubiquitination (Fig. 2d, Table S2). There were species-specific differences in
subunits of the PA28 proteasome regulatory particles, with human NPs having increased
levels of PA28 $\alpha\beta$ (PSME1/E2), and mouse NPs containing higher PA28γ/Psme3 levels (Fig.
2e). Both PA28$\alpha\beta$ and PA28 γ particles have been shown to be less efficient than the 19S
canonical regulatory particle at activating the 20S proteolytic activity[21], and could compete
with one another for binding to this 20S core. Proteasome-related proteins such as Ttc3,
Psma3, Psmd5, Psmd12; ubiquitin ligases Smurf1, Rnf34 and Cul7; F box protein Tbl1x;
proteasome accessory protein Ubqln1, Ubqln2, Ubr1, Psmd5, Ubr2, Psmd12, and Psmg3
and zinc finger protein Zfand1 were upregulated in mouse NPs, supporting a potential
increase in protein degradation in mouse compared to human NPs. Proteins related to
sumoylation (Ube2i, Sumo2), and enzyme transferases (Prmt9, Nsd, Hcfc1) were detected
as differentially increased in mouse NPs as well (Fig. 2a). By contrast, the chaperones and
regulatory subunits PSMD10, NGLY1, PSMD4, TRIP12, and PPP1CC were significantly
upregulated in human NPs (Fig. 2b, c). The expression of deubiquitinases USP7, USP11,
and ABRAXAS2 , implicated in the stabilization of proteins[22,23], were increased in human
NPs.
By contrast with these differences in degradative machineries between the two species, the
proteomic analyses did not identify differential expression of proteins associated with protein
synthesis, supporting the observation of equivalent production rates in mouse and human
NPs.
181
To explore further the differences between mouse and human cells, we performed pathway
enrichment analysis[24] (Table S3). The pathways enriched in mouse were the anaphase-
promoting ubiquitination complex, a major E3 ubiquitin ligase complex with roles in the
control of cell cycle and neurogenesis[25–27], the mitochondrial electron transport chain,
specifically complex III/ Ubiquinol-Cytochrome C Reductase, and enzymes of the TCA cycle
(e.g. isocitrate dehydrogenase subunits IDH3G and IDH3A) (Fig. 2b,c). The pathways
enriched in human NPs represented upregulation of cell-cell communication (e.g., NCAM1,
log2FC=6; PODXL), Ca2+ signalling, immunity and secretion (e.g., GSN, SRP14), as well as
Rho GTPase-mediated signal transduction (e.g., GNA13, GNAI1). Specific human-
upregulated pathways included semaphorins, involved in the axonal development (e.g.,

DPYSL5 hydrolase and dihydropyrimidinase CRMP1), as well as sphingolipid metabolism
(e.g., SMPD4), involved in the neuronal development[28,29]. The Pentose/Glucoronate
interconversion pathway was upregulated in human cells. Localized in the cytoplasm, this
pathway is a key route to carbohydrate synthesis. It also plays a central role in recycling
NADP to NADPH, a cellular reducing agent for reductive biosynthesis and protection against
oxidative stress[30].
198
199   Together, the comparison of the mouse and human proteome in NPs indicated selective
200   upregulation of proteins related to degradation and mitochondrial metabolism in the mouse.
201   Proteins overrepresented in human were associated with neural phenotypes, the
202   biosynthesis of carbohydrates, and the stabilization of the proteome.
203
Proteins in all cellular components display similar fold-change differences in stability
Having established a genome-wide difference in protein turnover between mouse and
human NPs[5], we proceeded to profile protein turnover in mouse and human to determine the
identity of specific proteins and pathways that exhibit similar or divergent protein dynamics
between species. We performed dynamic SILAC (Stable Isotope Labeling by Amino acids in
cell Culture) coupled to mass spectrometry in mouse and human NPs to estimate half-lives
of individual proteins at a proteome-wide scale[31]. The labelling was centered on equivalent
time points in mouse and human NPs based on our previous stability analysis[5], and the
collection of five time points per species was distributed according to the expected stability
differences between mouse and human. The media was switched to heavy lysine and
arginine isotopes 24 hours before day 2 in mouse. For human, we started labeling on day 4,
48 hours before human NPs reached the stage assayed (Fig. 3a). The samples were
analyzed by LC-MS/MS at each time-point, and the degradation rate constants for individual
proteins calculated (see Methods). We determined the half-lives of 4990 and 4640 proteins
in mouse and human, respectively (Table S5, S6). Their median half-lives were 18.2 hours
and 27.9 hours, respectively (Fig. 3b). To obtain more detailed insights into protein features
associated with differences in stability in the mouse and human datasets, we performed
gene set enrichment analysis (GSEA) across all GO terms (Fig.S3a, b). Structural proteins
were enriched for long half-lives in both mouse and human, as expected. Furthermore,
proteins with metabolism- and translation-related functions, and proteins located in the
mitochondrial and ribosomes were more stable on average, especially in mouse (Table S7,
S8).
226
We further compared the half-lives of 3402 homologous proteins detected in both mouse
and human datasets (Fig.3c, Table S9). Half-lives of human homologues were generally
longer than those of mouse homologues, showing a median fold difference of 1.48 (Fig. 3c),
consistent with the measurement for OLIG2 (Fig. 1a). To investigate whether specific
features of proteins determine their stabilities, we first split the 3402 homologous proteins
into quartiles based on their size (amino acid sequence length) or abundance (average total
expression level). Each quartile exhibited similar species-specific trends of half-lives and
interspecies fold differences (Fig. 3d, Fig. S3c, d). We observed a mild correlation between
protein abundance and half-lives, with more abundant proteins tending to be more stable;
however, the half-life fold differences between species were similar irrespective of
expression levels (Fig. 3e, Fig. S3c, e). These results indicated that the overall differences in
protein stability between mouse and human NPs are independent of protein size and
abundance.

To examine if differences in the subcellular localization of proteins affected half-life, we
analysed the dataset categorised by cellular compartments (see Methods). Proteins
assigned to the nucleus, cytoplasm, membrane or extracellular compartment showed similar
median half-lives in each species (Fig. 3f). Consequently, the observed stabilisation of the
human proteome compared to the mouse occurs in all cellular compartments. We then
categorised proteins based on their molecular functions by referring to GO terms (Fig. S3g).
Although most of the groups followed the overall trend, proteins classified as "Structural
molecule activity", such as TUBB, RPL32 and LMNA, and "ATP-dependent activity"
appeared longer-lived, while the groups "Transcription regulator activity", including
transcription factors like POU3F2, YAP1 and ZEB2, and "Molecular adaptor activity" were
short-lived in comparison with the overall median (Table S10). Nevertheless, the fold
difference in half-life between the species was preserved for the proteins. Together, the
results indicate that proteins in diverse biophysical properties, functional categories, and
subcelullar locations have similar stability differences between mouse and human NPs.

Given that our earlier comparative proteomics analysis had indicated several differences in
the proteasome between the two species, we reasoned it was possible that the observed
differences in proteome half-lives could be explained by differences in the stability of specific
components of the proteasome, as it accounts for ~70% of the cellular protein degradation[32].
We grouped proteins associated with the proteasome into core proteasomal components,
proteasomal regulatory components, or proteasomal activators and inhibitors from a curated
list generated by the proteostasis consortium[33]. We compared the median half-lives of
proteins in these categories to the median half-lives of the proteome in mouse and human.
Whereas core particles followed the median trend for mice and human (Fig. S3h), regulatory
particles were more stable than average in both species, especially in mice (Fig. S3h). For
proteins in the "activator or inhibitor" category, we observed that human proteins were more
unstable compared to the human median, but this was not the case for mouse (Fig. S3h, i).
For example, PSME3 half-life was shorter in human compared to the median protein stability
(Fig. S3i). In mouse, Psme3 half-life followed the median trend for mouse. The increased
degradation rate in human PSME3 may contribute to the significant upregulation of Psme3 in
mouse in the global dataset (Fig. 2e). By contrast, the proteasome regulator Psmf1[34] was
more unstable than the median in mouse, and this was not the case in human. Global
proteomic quantification indicated that Psmf1 was expressed at almost twice the level in
mouse compared to human (Fig. 2e). Together, these results identify differences in the
stability of activators and inhibitors of the proteasome between mouse and human and
suggest a stabilization of the mouse proteasome and/or lower proteasomal activity in human
compared to mouse.

A role for active protein degradation in mouse and human neural progenitors
The faster cell cycle in mouse compared to human[5] raised the possibility that the lower rate
of protein turnover in human NPs could arise from lower cell division rates resulting in a
lower rate of passive dilution. To test this, we set out to compare protein stability in non-
dividing cells by inducing the generation of post-mitotic neurons *in vitro* from mouse and
human NPs (Fig. 4a, b). Treatment with the gamma-secretase inhibitor Dibenzazepine
(DBZ), which inhibits the activation of Notch pathway, increased the production of TUBB3+
neurons at the expense of SOX2 expressing NPs[35] (Fig. S4a, b). Measurements of protein
stability in these conditions by AHA pulse-chase labeling[5] showed that the half-life in mouse

post-mitotic neurons was ~10h (t1/2 = 11.45 ± 1.8 hours), and ~20h in human neurons (t1/2
= 22.93 ± 4.58 hours) (Fig. 4c, d, Fig. S4c). These rates were similar to those measured in
NPs (mouse NPs t1/2 = 9.3h ± 1.99 hours; human NPs t1/2 = 17.57h ± 3.6 hours). These
results indicate that protein degradation is slower in human neurons than mouse neurons
and support the conclusion that active protein degradation mechanisms drive differences in
protein stability between mouse and human.
294
A role for active protein degradation in the species differences in protein turnover predicts
higher proteasome activity in mouse NPs than human NPs. This is supported by the global
and SILAC proteomic analyses that indicated differences associated with protein
degradation (Fig.2b, Fig. S3h, i). To test this directly, we used a luminogenic assay to
measure chymotrypsin-like proteasomal activity[36,37] in mouse and human NPs at equivalent
developmental stages (Fig. 4e). We observed 1.43-fold higher activity in the mouse
compared to human (Fig. 4f). This suggests that higher proteasomal activity in mouse NPs
compared to human NPs contributes to the higher rate of protein degradation of proteins and
is consistent with the idea that active protein degradation mechanisms are responsible for
the differences in protein stability between mouse and human.
305
Irx3 depletion accelerates Olig2 appearance
Given the findings suggesting a role for protein degradation in the differing developmental
tempos between mouse and human NPs, we sought to test whether modulating the stability
of a specific regulatory protein affected the tempo in NPs. We focused on IRX3, which is
expressed early in spinal cord development, prior to cells receiving sonic hedgehog
signaling, and represses OLIG2 and motor neuron progenitor formation[10,38]. To this end we
generated an Irx3::HA::HALO mouse ESC line (Irx3-HALO) by inserting the HALO tag at the
C-terminus of the Irx3 gene. An HA epitope was inserted as a linker between the Irx3 coding
region and HaloTag[39].
315
We tested Irx3-HALO expression and functionality in spinal cord motor neuron
differentiation. We differentiated Irx3-HALO cells and we exposed cells to RA only on day 3
to promote IRX3 expression (Fig. S5c). We then performed a pulse-chase experiment with
HALO-JF549 ligand (Fig. S5a). Cells were pulsed for 1h, ligand was then removed, and
samples collected 0h, 4h and 24h later. We observed decreasing levels of HALO-JF549
signal during the chase, while the level of IRX3 expression, assayed by HA
immunofluorescence, remained constant across the time course (Fig. S5b, c). This
confirmed that the Irx3-HALO line reproduces the expected expression pattern of IRX3 and
validates the use of the HA::HALO tag as a multimodal tag to track IRX3 levels and
dynamics of turnover.
326
The HALOtag allowed us to use a small molecule degrader, HaloPROTAC, to induce IRX3
degradation. HaloPROTAC binds covalently to the HaloTagged protein and recruits specific
E3 ligases, which promote ubiquitination of the protein and subsequent degradation by the
26S proteasome[39,40]. To test whether HaloPROTAC decreased IRX3 protein stability, IRX3
expression was induced by addition of RA for 24h as before. HaloPROTAC was added to the
Irx3-HALO at the time of RA addition. We collected samples at 0h, 12h, and 24h after
HaloPROTAC treatment as well as in a control Irx3-/- cell line. While IRX3 was detected
across time points in the untreated control condition (- PROTAC), IRX3 was reduced 24h
after HaloPROTAC addition (0h time point), and HA intensity was reduced almost to the

levels found in the Irx3-/- line 48h after HaloPROTAC addition (Fig. S5d). These results
confirm protein depletion of by addition of HaloPROTAC.
338
We next tested how acute IRX3 degradation impacted the dynamics of OLIG2 expression, a
target of IRX3 repression[11,41]. We cultured differentiating Irx3-HALO cells for 24h in RA to
induce Irx3 in neural progenitors. Cells were then switched to media containing RA and SAG
either in the presence or absence of HaloPROTAC degrader. This culture regime allowed the
maximal expression of IRX3 prior to OLIG2 expression induced by SAG induction. We
assayed OLIG2 expression 12-48h after SAG and HaloPROTAC addition (Fig. 5b, c). In cells
that had not been exposed to HaloPROTAC, SAG-dependent repression of IRX3 and
induction of Olig2 took more than 18h. By contrast, in cells treated with HaloPROTAC, IRX3
was largely absent by 18h and an increase in the percentage of OLIG2+ cells was apparent
(Fig. 5d). Quantifications of the IRX3-;OLIG2+ population revealed a higher proportion of
OLIG2+ cells and lower variability in the HaloPROTAC condition compared to control (Fig.
5d). Moreover, fewer cells co-expressed IRX3 and OLIG2 with HaloPROTAC, indicating a
more rapid transition from the IRX3+;OLIG2- to an IRX3-;OLIG2+ state (Fig. S5e). Together,
these results demonstrate that acute degradation of the transcriptional repressor IRX3
accelerates the appearance of OLIG2 expressing neural progenitor cells, providing evidence
that modulating protein stability can influence the tempo of developmental gene regulatory
networks.
356
DISCUSSION
Here we provide a systematic comparison of protein dynamics between mouse and human
neural progenitors. Through a combination of targeted protein labelling, quantitative mass
spectrometry, and genetic manipulation of endogenous proteins, we have characterised
protein abundance, production, and degradation rates in equivalent mouse and human NPs.
The data provide insight into the mechanisms underlying the differences in developmental
tempo between these species.
364
Our findings show a ~1.5 fold-difference in the stability of homologous proteins between
mouse and human. The trends we observe are consistent with previous assays that
measured the overall degradation rate of the proteome by global metabolic labeling[42–46].
Despite the differences in protein degradation rates between species, we find no marked
differences in the rate of production between mouse and human NPs. The similarity in
protein production rates between species suggests that differential translation rates is not a
primary driver of the observed tempo differences. While different from the findings in
presomitic cells[6,7], our results are reminiscent of findings in mouse spinal cord progenitors *in*
vivo which showed that embryos with reduced protein synthesis rates (Rpl24(Bst)/+) are
noticeably smaller in size, but the patterning of progenitors occurs at the same ratio and
pace[47].
376
The similarity in protein translation rates coupled with differences in degradation rates
between species results in higher protein content in human NPs in comparison to mouse.
This could contribute to increased protein stability by increasing interactions or decreasing
diffusivity[48,49] and lead to a decrease in protein-proteasome encounters. However, the net
effect of crowding on protein stability depends on the specific properties of individual
proteins and their interactions in the cellular environment. Future studies investigating
protein concentration in mouse and human NPs, and the relationship between molecular

crowding and protein stability could provide valuable insights into how these biophysical
properties contribute to species-specific developmental timing.
386
Our findings support a role for active protein degradation in controlling developmental tempo.
Unbiased global proteomic analysis suggests an increased abundance of proteins
associated with degradative pathways and metabolic functions in mouse. Importantly, we
found that the differences in protein stability persisted in post-mitotic neurons, indicating that
active degradation, rather than passive dilution through cell division, is the primary
mechanism driving the differences in species-specific protein turnover rates. The increased
stability of human proteins in NPs was observed across most cellular compartments and was
independent of aminoacid sequence, protein size or abundance. At least part of this
differential protein degradation rates between mouse and human NPs can be explained by
differences in proteasomal activity. We observed 1.43-fold higher chymotrypsin-like
proteasomal activity in mouse NPs compared to human. There did not appear to be clear
differences in the abundance of core particle proteasome subunits between species
(including that of its chymotrypsin-like Psmb5 subunit), suggesting that the observed
differences in proteasomal activity may be regulated either by differences in the activity of
regulatory complexes and/or the proteasome inhibitory proteins, through the availability of
proteasome targets, or through other post-translational regulatory mechanisms. In line with
this, our comparative proteomics identified differences between the two species in ubiquitin-
proteasome pathway components that could have roles in each of these facets of
proteasomal regulation. For example, we identify differences in the abundance and stability
of regulatory particles of the PA28αβ and γ complexes. While the PA28αβ is thought to be
characteristic of the immune system, PA28γ is expressed in all cell types. Also, PA28
complexes promote protein degradation in an ATP and ubiquitin-independent manner[21,50–52].
In the future, dissecting the functional outcomes of the different proteasomal regulatory
complexes in mouse and human NPs will be key to understand how homologous proteins
are degraded at different rates.
412
The importance of protein degradation in regulating developmental pace was highlighted by
the targeted manipulation of IRX3, a key transcriptional repressor in neural progenitor
pattern[10,41]. Increasing the degradation rate of IRX3 using a HaloPROTAC system led to
accelerated expression of its target gene, OLIG2, and suggests an accelerated transition of
intermediate NPs to ventral motor neuron progenitors. This demonstrates that modulating
the stability of individual regulatory proteins can have a significant effect on gene expression
dynamics and cell fate transitions during development. Moreover, these experiments offer
proof-of-concept for the use of the HaloPROTAC system for targeted perturbation of
components of a gene regulatory network that could be broadly applied to dissect gene
regulatory mechanisms.
423
Consistent with our findings, Matsuda et al. report a pervasive ~1.5-fold slower rate of
protein degradation in presomitic mesoderm[53], and Swovick et al. describe differences in
protein turnover across twelve different mammalian species with divergent lifespan[54]. The
conservation in different tissues of species-specific differences in protein degradation rates
suggests that this may be a fundamental mechanism for evolutionary adaptation of timing in
development and homeostasis. Our work measures differences in abundance and stability of
regulatory proteasomal subunits that may have a global impact on tempo. Future studies

investigating the molecular basis for the increased proteasomal activity in mouse cells, as
well as potential differences in other degradation pathways such as the autophagy-lysosome
system, will be important for fully understanding these species-specific differences.
434
While much attention for the basis of interspecies differences has focused on the regulatory
logic of gene networks through gene duplication or de novo acquisition of enhancers[2,3], our
results emphasise that post-translational processes, particularly protein degradation, play a
crucial role in shaping the temporal dynamics of development. Modulating protein stability
could offer a flexible and precise method for globally controlling developmental tempo,
without requiring extensive changes to gene regulatory networks or protein sequences in
individual developmental processes. Whether differences in protein degradation rates also
account for heterochronic changes in developmental pace, such as those associated with
the expansion of the primate cortex or the protracted maturation of cortical neurons remains
to be determined[3,4,55,56]. Our work highlights the importance of considering post-translational
regulation when studying developmental systems and evolutionary divergence between
species. Little is known about the mechanisms that drive and maintain the homeostasis of
proteins (proteostasis) across species and how they affect gene regulatory programs in
development. Exploring protein turnover in different evolutionary contexts could provide
insight into the constraints and selective pressures shaping developmental timing across
species.
451
In conclusion, our study provides a comparison of protein dynamics between mouse and
human NPs, revealing that differential protein degradation appears to be a key driver of
species-specific developmental timing. These findings raise questions about the molecular
mechanisms underlying evolutionary changes in developmental tempo and the potential role
of protein turnover in other aspects of phenotypic diversity between species. Future studies
exploring how protein degradation is controlled globally and its impact across developmental
processes in diverse organisms will be essential for deepening our understanding of the
molecular basis of evolutionary change. Understanding these mechanisms will not only
contribute to our knowledge of evolutionary and developmental biology but may also inform
strategies for manipulating cellular timing in contexts such as stem cell differentiation,
regenerative medicine, and homeostasis.
463
MATERIALS AND METHODS
Cell culture and neural progenitor differentiation
Mouse ESCs (HM1, OLIG2::HA::SNAP and IRX3::HA::HALO mouse ESC line) were
propagated on mitotically inactivated mouse embryonic fibroblasts (feeders) in DMEM
knockout medium supplemented with 1000U/ml ESGRO-LIF (ESG117 Sigma Aldrich), 10%
cell-culture validated fetal bovine serum, penicillin/streptomycin, 2 mM L-glutamine (GIBCO)
or on 0.1% gelatine in ES medium supplemented with LIF/2i, consisting of 1 µM PD0325901
and 3 µM CHIR99021P (Stem Cell Institute).
WA09/H9 ESC line (WiCell), and OLIG2::HA::SNAP H9 ESC lines were routinely cultured in
Essential 8 medium (Thermo Fisher A1517001) or StemFlex medium (Thermo Fisher
A3349401) on 0.5 µg/cm^2 laminin-coated plates (Thermo Fisher A29249) or 0.5 µg/cm^2 of
Vitronectin (Thermo Fisher A14700), and split using Versene (Gibco 15040066). The use of
human ESCs was carried out in accordance with approvals from the UK Stem Cell Bank
Steering Committee (SCSC14-18 to J.B. and SCSC21-46 to T.R.). All cell lines used in this
study were confirmed to be mycoplasma negative.

Mouse and human NP differentiation was performed following Rayon et al.[5]. For cells grown
on ES medium with LIF/2i, 2i was removed at the split prior to differentiation. The γ-
secretase inhibitor DBZ was used in experiments to fully differentiate neurons. Cells were
treated with 10 ng/uL DBZ (Tocris Biosciences Cat. No. 4489) for 72h starting at day 2 for
mouse and day 5 in human.
For IRX3::HA:: HALO mESC differentiation, N2B27 was supplemented with 100 nM RA
(R2625 Sigma Aldrich) for neural induction at day 0 and supplemented with RA and 500 nM
SAG (566660 Calbiochem) thereafter. To achieve IRX3 depletion, from day 1 onwards, cells
were treated with 1 μM HaloPROTAC3 (Promega GA3110) or DMSO as control. Samples
were collected after 12h, 18h, 24h, 33h, 42h, 48h of HaloPROTAC or DMSO treatment and
processed for intracellular flow cytometry (see below). Medium was changed daily.
491

Generation of mouse and human ESC lines by CRISPR
All transgenic cell lines were generated by CRISPR-Cas9–mediated homologous
recombination, as in Gouti et al. 2017[57] and Rayon et al. 2020[5]. Briefly, 2×10^6 cells were
electroporated with 2 μg of each plasmid using program A23 of Nucleofector II (Amaxa) and
mouse stem cell Nucleofector kit (Lonza DPH-1001) or human stem cell Nucleofector I kit
(Lonza VPH-5012). For selection, colonies were first treated with 0.5 mg/ml Puromycin
(Sigma P9620) for 2 days followed by 50 μg/ml Geneticin (Thermo Fisher 10131027)
selection. Individual colonies were picked and replated to allow expansion and a second
round of Geneticin selection. Correct integration of the T2A-HA-SNAPTag transgene was
verified using long-range PCRs and Sanger sequencing.
502

Generation of the pNTKV-HA-HaloTag and pNTKV-HA-SnapTag vectors
The coding sequence for the HaloTag was extracted by 2-step PCR amplification with
Phusion High-Fidelity DNA Polymerase (NEB, M0530) from the pFC27A HaloTag CMV-neo
Flexi vector (Promega, G842A), while, at the same time, the coding sequence for the HA
epitope (tatccctatgacgtcccggactatgca) and two stop codons were introduced as part of the 5'
primer used for the amplification process.
For the HA-SnapTag cell line, the coding sequence for the SnapTag was extracted equally
by PCR amplification from the pSNAPf commercialvector (NEB, N9183S), while the HA
coding sequence and two stop codons were introduced as part of the 5' primer.
The resulting purified DNA elements were separately cloned, by restriction digestion and
inserted with in-Fusion cloning into the pNTKV-T2A-mKate vector[58].
514

Generation of Olig2::HA::SNAP mouse and human ESC lines by CRISPR
For CRISPR-Cas9–mediated homologous recombination, short guide RNA (sgRNA)
sequences (mouse: CGGCCAGCGGGGGTGCGTCC, human: CTGTCGCCAGAACGCGC)
were cloned into pSpCas9(BB)-2A-Puro (Addgene pX459 plasmid no. 62988). As donor
vector, the HA-SnapTag cassette was inserted at the 3′ end of the Olig2 open reading frame,
using 2.83-Kb upstream and 5.04-Kb downstream arms for mouse and 2.78-Kb upstream
and 1.98-Kb downstream homology arms for human.
522

Generation of mouse Irx3::HA::HALO ESC line
For CRISPR-Cas9–mediated homologous recombination, short guide RNA (sgRNA)
sequence: ATGGTTGAAAAGTTAAGACG was cloned into pSpCas9(BB)-2A-Puro (Addgene
pX459 plasmid no. 62988). As donor vector, the HA-HaloTag cassette was inserted at the 3′

end of the Irx3 open reading frame, using 2.32-Kb upstream and 0.34-Kb downstream
homology arms.
529
530 Generation of mouse Irx3 knockout line
531 For CRISPR-Cas9–mediated non-homologous end joining to generate loss of function by
532 frameshift two short guide RNA (sgRNA) sequences: TCTCTACCCACCCGAACGCC and
533 GGATGTACTGGTATCCGAGC were cloned into pSpCas9(BB)-2A-Puro (Addgene pX459
534 plasmid no. 62988).
535
SNAP/HALO labelling for pulse or pulse-chase experiments
For OLIG2:HA::SNAP experiments, mouse and human ESCs were differentiated to NP day 2
and day 6, respectively, as described above. To estimate OLIG2 degradation rates,
OLIG2::HA::SNAP cells were incubated for one hour in RA and SAG medium containing 100
nM SNAP-Cell 647-SiR ligand (NEB S9102S). After two PBS washes, cells were incubated
in RA and SAG medium containing 1μM SNAP-Cell Block (NEB S9106S) or 100nM SNAP-
Cell TMR-Star ligand (NEB S9105S) for the indicated chase periods. To estimate OLIG2
production rates, cells were incubated for one hour in RA and SAG medium containing
SNAP-Cell Block ligand. After two PBS washes, cells were incubated in RA and SAG
medium containing SNAP 647-SiR ligand for the indicated chase periods.
For IRX3::HA::SNAP mouse pulse-chase experiments, cells were labeled for 1h in medium
supplemented with 100nM Janelia Fluor HaloTag Ligand (549, Promega GA111) at day 0 on
RA as described above. After two PBS washes, cells were incubated in RA and SAG
medium for the indicated chase periods.
After SNAP/HALO ligand treatments, cells were washed once or twice with PBS for
degradation or production measurements, respectively, and processed for intracellular flow
cytometry or imaging. At least 3 biological replicates per species per time point from
independent experiments were used. For OLIG2 protein dynamics, geometrical mean was
measured on OLIG2 positive and negative populations. Half-life estimation was based on the
non-linear model fitting in GraphPad Prism version 10.0.0 for MacOS using values
normalized by samples not treated with SNAP-cell 647-SiR ligand. Error intervals reported
correspond with 95% confidence intervals. IRX3-HALO Images were taken on a Leica SP8
confocal microscope and analysed in FIJI.
559
Intracellular Flow Cytometry
Cells were dissociated with 0.5ml accutase (GIBCO) and fixed in 4% paraformaldehyde in
PBS for 5min. For stainings, 0.5 - 1x10^6 cells were used. Cells were incubated overnight with
an antibody mix on PBST with 1% BSA at 4^{o}C. The following day, and when HA (Cell
Signaling C29F4, 1:1000) or OLIG2 (R&D AF2418, 1:800) primary antibody were used, cells
were pelleted and incubated with donkey anti goat/rabbit Alexa Fluor secondary antibodies
(1:1000) at room temperature for 1h. Cells were resuspended in 0.5 mL PBS and filtered for
data acquisition on LSR Fortessa or Biorad ZE5 and MacsQuant Analysers. For OLIG2
measurements, 10,000 ~ 50,000 total events were recorded. For other experiments 10,000
events gated on SOX2 were recorded. Analysis was performed using FlowJo software. The
following fluorophore-conjugated antibodies were used: SOX2-V450 (BD 561610, 1:100),
TUBB3-AF488 (BioLegend 801203, 1:100), TUBB3-AF647 (BioLegend 560394, 1:100).
For Cell Trace Violet (Thermo Fisher C34557), 1x10^6 cells were incubated at the stated
concentrations and counterstained with LIVE/DEAD nearIR marker (Thermo Fisher L34975)
following manufacturer's intructions. Cells were quantified live.

Global protein production measurements
AHA uptake experiment

Experiments were performed on differentiated neural progenitors (mouse day 2 and human day 6), as described above. The experiments were performed in the presence of methionine as we reasoned that methionine deprivation might trigger a reduction in protein production and impact protein translation. Spent medium was replaced by 1 mM AHA in differentiation medium supplemented with RA and SAG at 0h timepoint. Samples were collected at the indicated timepoints for intracellular staining with Click It Cell Reaction buffer kit (Invitrogen C10269) and Alkyne Alexa Fluor 488 (Invitrogen A10267) followed by SOX2-V450.

Puromycin uptake experiment

We tested a range of puromycin concentrations and timepoints to identify conditions that do not trigger integrated stress response to the neural progenitors. 0.1 mg/ml puromycin for up to 2 hours gave sufficient staining intensity without inducing a cell stress response. Puromycin was applied to the differentiating NPs in differentiation medium supplemented with RA and SAG at 0h timepoint. Samples were collected at the indicated timepoints for intracellular staining with anti-puromycin Alexa Fluor 488 (EDM Millipore MABE343-AF488) and SOX2-V450 conjugated antibodies.

Global proteomics
Sample prep

Cells were treated with accutase to detach from cell culture plate, washed with PBS, counted and centrifuged. Cell pellets were frozen in an ethanol/dry ice bath and stored in -80oC until ready to process. Protein extracts were prepared with lysis buffer containing: 50mMTris pH8, 1% SDS, complete protease inhibitors (one tablet per 3.5 ml buffer, ROCHE 05056489001) and benzonase (250 U/ml, Pierce 88701). Protein concentration was determined using the BCA assay according to manufacturer's instructions (Pierce 23227). To prepare peptides 100 µg of proteins per sample were acetone precipitated, washed twice with 80% acetone and solubilised in 100 mM Hepes pH 8 at 1 mg/ml protein concentration. To reduce disulfide bonds and alkylate free cysteine residues samples were treated with 10 mM DTT for 30 min followed by 20 mM Iodoacetamide for 30 min, followed again by 10 mM DTT for 30 min. Proteins were digested with LysC (1 µg LysC per 100 µg of proteins) overnight (O/N) followed by Trypsin (1.33 µg Trypsin per 100 µg of proteins) O/N. Samples were stored at -80oC until ready to proceed with mass spec.

Data Independent Acquisition (DIA) analysis on timsTOF Pro2

Peptides were analysed using an Evosep One LC system (EvoSep) coupled to a timsTOF Pro 2 mass spectrometer (Bruker Daltonik GmbH) using a commercial 150-mm analytical column (EV1113 ENDURANCE COLUMN, Evosep Biosystems) and an integrated Captive Spray Emitter (IonOpticks). Buffer A was 0.1% formic acid in water, Buffer B was 0.1% formic acid in acetonitrile. Data was collected using diaPASEF with 1 MS frame and 9 diaPASEF frames per cycle with an accumulation and ramp time of 100 ms, for a total cycle time of 1.07 seconds. The diaPASEF frames were separated into 3 ion mobility windows, in total covering the 400 – 1000 m/z mass range with 25 m/z-wide windows between an ion mobility range of 0.64–1.4 Vs/cm^2. The collision energy was ramped linearly over the ion mobility range, with 20 eV applied at 0.6 Vs/cm^2 to 59 eV at 1.6 Vs/cm^2.

DIA Data Processing and Analysis
Data were analyzed using DIA-NN (version 1.8) with all settings as default except we
omitted normalisation. Specifically, a maximum of one tryptic missed cleavage was allowed,
with fixed modifications of N-term methionine excision and carbamidomethylation of cysteine
residues. No variable modifications were selected. A *Homo sapiens and a Mus musculus*
UniProt database was used for the analysis of the relevant samples. A default threshold of
1% false discovery rate was used at both the peptide and protein level. At least two unique
peptides were required to identify a protein. The library-free mode of DIA-NN was used to
generate precursor and fragment ions *in silico* from the UniProt database. Additionally, the
programme generates a library of decoy precursors (negative controls). Retention time
alignment was performed using endogenous and iRT peptides, and peak scores were
calculated by comparison of peak properties between observed and reference spectra.
635
Data Analysis
The data were analysed in three samples (biological replicates) derived from three bulked
samples of mouse or human NPs. Each sample was run three times on the mass-spec
(three technical replicates). Protein copy numbers per cell were estimated using the
proteomic ruler method[20], and 1:1 homologs between mouse and human were retained for
downstream analysis. The table with homologs was extracted from ensemble using
BioMart[59] and then manually curated to include missing gene names (Table S4). The
proteins that were identified in at least 2 samples in each group (human or mouse) were
used for downstream analysis. To reduce potential technical biases between mouse and
human samples, we performed a global correction of protein abundances by dividing them to
the size factors of the samples (total copy number of each sample per the mean of all
samples). The average copy number between the non-zero technical replicates were used
for differential expression, with zeros being replaced by medians intensities in each group
(human and mouse). The abundances were transformed to log2 scale and differential
expression was performed with R package limma[60], using empirical Bayes method for 2
group comparison with standard t-test of the eb.fit function output[61]. The differentially
expressed proteins (DEPs) were chosen based on the criteria of ≥ 2 fold change (FC) and
the false discovery rate (FDR) adjusted P values for multiple-hypothesis testing with the
Benjamini–Hochberg method FDR ≤ 0.05.
The pathway enrichment analysis was performed separately on mouse- or human-
upregulated DEPs. We used David functional enrichment analysis[24] with annotations from
GO:BP, GO:MF, GO:CC, KEGG and Reactome databases. The background was
represented by orthologous human or mouse proteins identified in mouse or human
samples, respectively (Table S1). The DAVID functional annotation clustering was used to
group similar and redundant terms and the top representative terms of each cluster were
combined for plotting.
662
Dynamic SILAC
Cell Culture and Stable Isotope Labeling
Mouse and human ESCs were differentiated as described above until 1st and 4th day of RA
and SAG addition, respectively. Then, cells were collected as 0h time point, or continued the
differentiation in media containing heavy arginine and lysine until collected after 3h, 8h, 24h
and 48h for mouse NPs, or after 6h, 24h, 48h and 96h for human NPs in three biological
replicates. For the "heavy" condition, DMEM/F12 and Neurobasal were replaced by SILAC
DMEM/F12 and SILAC Neural Basal (AthenaES 0423 and 0428) supplemented with

methionine and leucine. L-arginine hydrochloride (13C6, 15N4) and L-lysine hydrochloride
(13C6, 15N2) (Cambridge Isotope Laboratories CNLM-539-H and CNLM-291-H) were
dissolved in PBS and kept at -80°C and were used at the final concentrations of 0.549 mM
and 0.648 mM, respectively, based on the concentrations in the original medium. Medium
change intervals, seeding densities, coating methods and supplements of inhibitors and
growth factors followed the differentiation protocols described above.
677
Sample prep
The cell pellets were lysed in 5% SDS/100mM triethylammonium bicarbonate buffer (Sigma
TEAB) by probe sonication and heating. Protein concentration was measured by Pierce BCA
Assay. 40 µg proteins per sample were taken, reduced by tris(2-carboxyethyl)phosphine
hydrochloride solution (Sigma) and alkylated by iodoacetamide (Sigma) then purified by
acetone precipitation. Proteins were digested with trypsin (0.8 µg; Sequence modified,
Promega) overnight at 30°C in 50 mM ammonium bicarbonate.
685
Data Independent Acquisition (DIA) analysis on Orbitrap Eclipse Tribrid
The LC-MS/MS analysis was on the Orbitrap Eclipse Tribrid mass spectrometer coupled with
a U3000 RSLCnano system (both from ThermoFisher Scientific) using DIA mode. 200 ng
peptides were first loaded to the trap column (100 µm i.d. x 5 cm) and then separated on an
analytical column (75 µm id x 50 cm) using a 120-min gradient from 2 - 30%
acetonitrile/0.1% formic acid at a flow rate of 300 nl/min. Both trap column and analytical
column were packed in-house with C18 (3 µm, Reprosil C18AQ, Dr. Maisch). MS1 scans
were acquired from m/z 390 - 865 using a resolution setting at 120,000, AGC at 400,000 and
maximum injection time at 50 msec. MS/MS scans were acquired in the Orbitrap in DIA
mode, where the precursor mass range was set at m/z 400 – 855 with isolation windows of
16 m/z and 1 m/z overlap. The scan range was set at m/z 200 – 1800, with a resolution at
30,000, maximum injection time at 54 msec, and relative HCD collision energy at 30.
698
Data Analysis
DIA raw files were analysed using Spectronaut version 18.0 using the spectral library-free
directDIA approach, searching against a human or mouse UniProtKB-proteome database
(version June 2022). The Pulsar search settings were: Trypsin (specific) with maximum 2-
missed cleavage sites; peptide length from 7 to 52; Carbamidomethyl (C) as fixed
modification; toggle N-terminal M is true; Oxidation (M) and Acetyl (Protein N-terminus) as
variable modifications. FDRs at PSM, peptide and protein level were all set to 0.01.
Quantitation was based on peak areas from MS2 data, with channel 1 set for light (label-
free), and channel 2 set for heavy peptides (Arg(10) and Lys(8)). All other settings were
default.
709
Data were processed in R (v4.4.1) and RStudio (v2023.06.0+421) and mouse and human
datasets were processed in parallel. The fraction of light (channel 1) compared to heavy
(channel 2) chain was calculated for each replicate at each time point (L/(L+H)). Starting
from 8127 and 8509 detected proteins from mouse and human NPs, respectively. Quality
check filters were applied to remove proteins that had missing data, <0.85 fraction light at
the 0h timepoint, <1000 total expression level over all timepoints, or unstable total
expression levels across the timepoints. Instability in expression levels was defined as
differences in total expression across the timepoints (Kruskal Wallis p<0.01).

Fraction of light was transformed to a log2 scale, and a linear model was fitted for each
protein across the time points using all three replicates, with the y-intercept constrained to 0.
Those with R2 >0.85 and sigma <0.7 were kept, and the slope of the model was used to
calculate protein half-life using the following equation: $t1/2 = \frac{\log2(0.5)}{slope}$

Gene set enrichment analysis on the half-lives was conducted using GSEAPreranked
[v4.3.3,][62,63] to identify GO:BP, GP:MF, and GO:CC terms enriched in short or long half-lives.
Terms were deemed significant if they had a False Discovery Rate (FDR) less than 5%
(q<0.05). Revigo[64] was used to collapse similar terms, and those with a dispensability score
<0.5 were retained. Terms with fewer than 25 proteins in the dataset or that had greater than
80% overlapping genes with a term with larger normalised effect size (NES) were removed.
728
729 Homologs between mouse and human were extracted from Ensembl v112 using
730 BioMart[59,65], and those with 1:1 homologs were retained. Gene homologs that matched to
731 multiple Uniprot IDs were removed. Half-life fold difference between mouse and human was
732 calculated (human/mouse) to visualise differences in half-life between the species (Table
733 S9).
734
To group the proteins by GO terms, at least one of the homologous proteins annotated with
descendant terms of following were counted: CC: cytoplasm (GO:0005737), nucleus
(GO:0005634), membrane (GO:0016020) and extracellular region (GO:0005576); MF:
catalytic activity (GO:0003824), structural molecule activity (GO:0005198), transporter
activity (GO:0005215), binding (GO:0005488), molecular adaptor activity (GO:0060090),
molecular function regulator activity (GO:0098772), transcription regulator activity
(GO:0140110). ATP-dependent activity (GO:0140657); For MF, only the terms with more
than 100 proteins were shown, and terms with complete overlap were excluded (Table S10).
743
Proteome stability measurements in neurons
AHA pulse and chase experiments were performed as in Rayon et. al 2020[5]. Cells were
starved by replacing complete differentiation medium with methionine-free medium for 30
min. Next, 100μM AHA was added to the methionine-free medium for 1h. To measure protein
stability, AHA pulse was removed by washing the cells once with PBS and growing the cells
on complete differentiation medium for the course of the experiment. On the indicated days
of differentiation and at specific time points after AHA removal, cells were processed for
intracellular flow cytometry.
AHA-incorporated proteins were labeled using Click-iT™ Cell Reaction Buffer Kit (Thermo
Fisher Scientific C10269) on a volume of 150 μl per sample reaction. Estimations of global
proteome stability were calculated as in Rayon et al. 2020[5]. Briefly, estimations of proteome
stability were obtained by normalizing each individual replicate using an initial exponential fit
$y(t) = B + C \cdot \exp(kt)$ to determine the baseline B and initial fluorescence intensity C,
allowing a comparison of all the replicates together. Exponential fits of the bootstraps of the
normalized ensemble were used to construct confidence intervals of the degradation rates.
Error intervals reported correspond with 95% confidence intervals. At least 2 biological
replicates in technical duplicates per species per time point from independent experiments
were used.
762
Proteasome activity measurements

Mouse and human NPs were washed with PBS once, and then scraped in proteasome
assay buffer (50 mM Tris-HCl, pH7.5, 5 mM MgCl2, 0.5 mM EDTA and 10% Glycerol). Then,
lysate was homogenized by passing through 27-gauge needles ten times. Lysate was
centrifuged at 10,000 x *g* for 10 min at 4°C and supernatant was aliquoted for storage at -
80°C. 20 µg of total proteins per well of 96-well plate with black walls (Greiner, 655090) were
diluted in the proteasome assay buffer containing 2 mM ATP (Thermo Fisher Scientific), 1
mM dithiothreitol (Sigma) and 0.333 mg/ml Z-GGL-AMC (Cambridge Bioscience).
Fluorescence was measured by PHERA star plate reader (360 nm excitation, 430 nm
emission, BMG Labtech) every 2 min for 60 cycles at 37°C. Kinetic slopes were measured
for the stable 60 min. Protein amount was estimated by BCA assay.

Data and code availability
Scripts to analyse global and SILAC proteomics are both found at
https://github.com/POTsnake2/Nakanoh-Stamataki_2024.
777
778 Global proteomics datasets have been deposited to the PRIDE database[66] with the project
779 accession identifier: PXD054152. SILAC datasets have been deposited with the project
780 accession identifier: PXD053697.
781

REFERENCES
1. Ebisuya, M. & Briscoe, J. What does time mean in development? *Dev. Camb. Engl.* **145**,
dev164368 (2018).
2. Rayon, T. Cell time: How cells control developmental timetables. *Sci. Adv.* **9**, eadh1849
(2023).
3. Wallace, J. L. & Pollen, A. A. Human neuronal maturation comes of age: cellular
mechanisms and species differences. *Nat. Rev. Neurosci.* **25**, 7–29 (2024).
4. Iwata, R. & Vanderhaeghen, P. Metabolic mechanisms of species-specific developmental
tempo. *Dev. Cell* **59**, 1628–1639 (2024).
5. Rayon, T. *et al.* Species-specific pace of development is associated with differences in
protein stability. *Science* **369**, eaba7667 (2020).
6. Matsuda, M. *et al.* Species-specific segmentation clock periods are due to differential
biochemical reaction speeds. *Science* **369**, 1450–1455 (2020).
7. Diaz-Cuadros, M. *et al.* Metabolic regulation of species-specific developmental rates.
Nature **613**, 550–557 (2023).
8. Iwata, R., Casimir, P. & Vanderhaeghen, P. Mitochondrial dynamics in postmitotic cells
regulate neurogenesis. *Science* **369**, 858–862 (2020).
9. Lázaro, J. *et al.* A stem cell zoo uncovers intracellular scaling of developmental tempo
across mammals. *Cell Stem Cell* **30**, 938-949.e7 (2023).
10. Novitch, B. G., Chen, A. I. & Jessell, T. M. Coordinate regulation of motor neuron
subtype identity and pan-neuronal properties by the bHLH repressor Olig2. *Neuron* **31**,
773–789 (2001).
11. Frith, T. J. R., Briscoe, J. & Boezio, G. L. M. Chapter Five - From signalling to form: the
coordination of neural tube patterning. in *Current Topics in Developmental Biology* (ed.
Mallo, M.) vol. 159 168–231 (Academic Press, 2024).
12. Gautier, A. *et al.* An Engineered Protein Tag for Multiprotein Labeling in Living Cells.
Chem. Biol. **15**, 128–136 (2008).
13. Dieterich, D. C., Link, A. J., Graumann, J., Tirrell, D. A. & Schuman, E. M. Selective
identification of newly synthesized proteins in mammalian cells using bioorthogonal
noncanonical amino acid tagging (BONCAT). *Proc. Natl. Acad. Sci. U. S. A.* **103**, 9482–
9487 (2006).
14. Schmidt, E. K., Clavarino, G., Ceppi, M. & Pierre, P. SUnSET, a nonradioactive method
to monitor protein synthesis. *Nat. Methods* **6**, 275–277 (2009).

15. Marciano, R., Leprivier, G. & Rotblat, B. Puromycin labeling does not allow protein
synthesis to be measured in energy-starved cells. *Cell Death Dis.* **9**, 1–3 (2018).
16. Aviner, R. The science of puromycin: From studies of ribosome function to applications
in biotechnology. *Comput. Struct. Biotechnol. J.* **18**, 1074–1083 (2020).
17. Pakos-Zebrucka, K. *et al.* The integrated stress response. *EMBO Rep.* **17**, 1374–1395
(2016).
18. Smith, P. K. *et al.* Measurement of protein using bicinchoninic acid. *Anal. Biochem.* **150**,
76–85 (1985).
19. Berenson, D. F., Zatulovskiy, E., Xie, S. & Skotheim, J. M. Constitutive expression of a
fluorescent protein reports the size of live human cells. *Mol. Biol. Cell* **30**, 2985–2995
(2019).
20. Wiśniewski, J. R., Hein, M. Y., Cox, J. & Mann, M. A "Proteomic Ruler" for Protein Copy
Number and Concentration Estimation without Spike-in Standards. *Mol. Cell. Proteomics*
MCP **13**, 3497 (2014).
21. Mao, I., Liu, J., Li, X. & Luo, H. REGgamma, a proteasome activator and beyond? *Cell.*
Mol. Life Sci. CMLS **65**, 3971–3980 (2008).
22. Clague, M. J., Coulson, J. M. & Urbé, S. Cellular functions of the DUBs. *J. Cell Sci.* **125**,
277–286 (2012).
23. Raule, M. *et al.* PA28αβ Reduces Size and Increases Hydrophilicity of 20S
Immunoproteasome Peptide Products. *Chem. Biol.* **21**, 470–480 (2014).
24. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and
functional annotation of gene lists (2021 update). *Nucleic Acids Res.* **50**, W216–W221
(2022).
25. Peters, J.-M. The anaphase promoting complex/cyclosome: a machine designed to
destroy. *Nat. Rev. Mol. Cell Biol.* **7**, 644–656 (2006).
26. Ferguson, C. J. *et al.* APC7 mediates ubiquitin signaling in constitutive heterochromatin
in the developing mammalian brain. *Mol. Cell* **82**, 90-105.e13 (2022).
27. Huang, J. & Bonni, A. A decade of the anaphase-promoting complex in the nervous
system. *Genes Dev.* **30**, 622–638 (2016).
28. Inskeep, K. A., Crase, B. & Stottmann, R. W. SMPD4 mediated sphingolipid metabolism
regulates brain and primary cilia development. *bioRxiv* 2023.12.15.571873 (2023)
doi:10.1101/2023.12.15.571873.
29. Koncina, E., Roth, L., Gonthier, B. & Bagnard, D. Role of Semaphorins during Axon
Growth and Guidance. in *Madame Curie Bioscience Database [Internet]* (Landes
Bioscience, 2013).
30. Kovářová, J. & Barrett, M. P. The Pentose Phosphate Pathway in Parasitic
Trypanosomatids. *Trends Parasitol.* **32**, 622–634 (2016).
31. Doherty, M. K., Hammond, D. E., Clague, M. J., Gaskell, S. J. & Beynon, R. J. Turnover
of the Human Proteome: Determination of Protein Intracellular Stability by Dynamic
SILAC. doi:10.1021/pr800641v.
32. Collins, G. A. & Goldberg, A. L. The Logic of the 26S Proteasome. *Cell* **169**, 792–806
(2017).
33. Consortium, T. P. *et al.* A Comprehensive Enumeration of the Human Proteostasis
Network. 1. Components of Translation, Protein Folding, and Organelle-Specific
Systems. Preprint at https://doi.org/10.1101/2022.08.30.505920 (2022).
34. Minis, A. *et al.* The proteasome regulator PI31 is required for protein homeostasis,
synapse maintenance, and neuronal survival in mice. *Proc. Natl. Acad. Sci. U. S. A.* **116**,
24639–24650 (2019).
35. Delás, M. J. *et al.* Developmental cell fate choice in neural tube progenitors employs two
distinct cis-regulatory strategies. *Dev. Cell* **58**, 3-17.e8 (2023).
36. Vilchez, D. *et al.* Increased proteasome activity in human embryonic stem cells is
regulated by PSMD11. *Nature* **489**, 304–308 (2012).
37. Gaczynska, M. & Osmulski, P. A. Characterization of noncompetitive regulators of
proteasome activity. *Methods Enzymol.* **398**, 425–438 (2005).

38. Chen, J. A. *et al.* Mir-17-3p controls spinal neural progenitor patterning by regulating
Olig2/Irx3 cross-repressive loop. *Neuron* **69**, 721–735 (2011).
39. Los, G. V. *et al.* HaloTag: A novel protein labeling technology for cell imaging and protein
analysis. *ACS Chem. Biol.* **3**, 373–382 (2008).
40. Buckley, D. L. *et al.* HaloPROTACS: Use of Small Molecule PROTACs to Induce
Degradation of HaloTag Fusion Proteins. *ACS Chem. Biol.* **10**, 1831–1837 (2015).
41. Novitch, B. G., Wichterle, H., Jessell, T. M. & Sockanathan, S. A requirement for retinoic
acid-mediated transcriptional activation in ventral neural patterning and motor neuron
specification. *Neuron* **40**, 81–95 (2003).
42. Swovick, K. *et al.* Cross-species Comparison of Proteome Turnover Kinetics. *Mol. Cell.*
Proteomics **17**, 580–591 (2018).
43. Harasimov, K. *et al.* The maintenance of oocytes in the mammalian ovary involves
extreme protein longevity. *Nat. Cell Biol.* **26**, 1124–1138 (2024).
44. Dörrbaum, A. R., Alvarez-Castelao, B., Nassim-Assir, B., Langer, J. D. & Schuman, E. M.
Proteome dynamics during homeostatic scaling in cultured neurons. *eLife* **9**, e52939
(2020).
45. Dörrbaum, A. R., Kochen, L., Langer, J. D. & Schuman, E. M. Local and global
influences on protein turnover in neurons and glia. *eLife* **7**, e34202 (2018).
46. Schwanhäusser, B. *et al.* Global quantification of mammalian gene expression control.
Nature **473**, 337–42 (2011).
47. Kicheva, A. *et al.* Coordination of progenitor specification and growth in mouse and chick
spinal cord. *Science* **345**, 1254927 (2014).
48. Speer, S. L., Stewart, C. J., Sapir, L., Harries, D. & Pielak, G. J. Macromolecular
Crowding Is More than Hard-Core Repulsions. *Annu. Rev. Biophys.* **51**, 267–300 (2022).
49. Macromolecular Crowding Tunes Protein Stability by Manipulating Solvent Accessibility -
Köhn - 2019 - ChemBioChem - Wiley Online Library. https://chemistry-
europe.onlinelibrary.wiley.com/doi/full/10.1002/cbic.201800679.
50. Rechsteiner, M. & Hill, C. P. Mobilizing the proteolytic machine: cell biological roles of
proteasome activators and inhibitors. *Trends Cell Biol.* **15**, 27–33 (2005).
51. Stadtmueller, B. M. & Hill, C. P. Proteasome Activators. *Mol. Cell* **41**, 8–19 (2011).
52. Lee, H. J. *et al.* Cold temperature extends longevity and prevents disease-related protein
aggregation through PA28γ-induced proteasomes. *Nat. Aging* 1–21 (2023)
doi:10.1038/s43587-023-00383-4.
53. Matsuda, M., Hammaren, H. M., Lazaro, J., Savitski, M. M. & Ebisuya, M. Systematic
analysis of protein stability associated with species-specific developmental tempo.
bioRxiv 2024.06.07.597977 (2024) doi:10.1101/2024.06.07.597977.
54. Swovick, K. *et al.* Interspecies Differences in Proteome Turnover Kinetics Are Correlated
With Life Spans and Energetic Demands. *Mol. Cell. Proteomics* **20**, 100041 (2021).
55. Espinós, A., Fernández-Ortuño, E., Negri, E. & Borrell, V. Evolution of genetic
mechanisms regulating cortical neurogenesis. *Dev. Neurobiol.* **82**, 428–453 (2022).
56. Lindhout, F. W., Krienen, F. M., Pollard, K. S. & Lancaster, M. A. A molecular and cellular
perspective on human brain evolution and tempo. *Nature* **630**, 596–608 (2024).
57. Gouti, M. *et al.* A Gene Regulatory Network Balances Neural and Mesoderm
Specification during Vertebrate Trunk Development. *Dev. Cell* **41**, 243--261.e7 (2017).
58. Sagner, A. *et al.* Olig2 and Hes regulatory dynamics during motor neuron differentiation
revealed by single cell transcriptomics. *PLOS Biol.* **16**, e2003127 (2018).
59. Smedley, D. *et al.* BioMart – biological queries made easy. *BMC Genomics* **10**, 22
(2009).
60. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).
61. Kammers, K., Cole, R. N., Tiengwe, C. & Ruczinski, I. Detecting Significant Changes in
Protein Abundance. *EuPA Open Proteomics* **7**, 11–19 (2015).
62. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.* **102**, 15545–15550
(2005).

63. Mootha, V. K. *et al.* PGC-1α-responsive genes involved in oxidative phosphorylation are
coordinately downregulated in human diabetes. *Nat. Genet.* **34**, 267–273 (2003).
64. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes
Long Lists of Gene Ontology Terms. *PLOS ONE* **6**, e21800 (2011).
65. Harrison, P. W. *et al.* Ensembl 2024. *Nucleic Acids Res.* **52**, D891–D899 (2024).
66. Martens, L. *et al.* PRIDE: the proteomics identifications database. *Proteomics* **5**, 3537–
3545 (2005).

FIGURE LEGENDS
Figure 1. Dynamics of protein production and degradation in mouse and human
neural progenitors
(a) OLIG2-SNAP pulse and chase experiments in mouse (orange) and human (blue) NPs.
Line and shaded areas show exponential fit with 95% confidence intervals (bottom); (mouse
n=3, human n=5 except for t_{3h} n=1; t_{4h}=4; t_{8h} n=4).
(b) OLIG2-SNAP production measurements in mouse (orange) and human (blue) NPs.
Geometric mean ± standard deviation of SNAP 647-SiR ligand incorporation at the indicated
time points in mouse and human NPs. No significant differences were observed using
multiple sample paired t-tests (n=3).
(c) Representative flow cytometry analysis of 647-SiR ligand incorporation in mouse and
human NPs for OLIG2-SNAP production measurements.
(d) Quantification of AHA incorporation at the indicated time points in mouse and human
NPs. Values correspond to geometric mean intensity and standard deviation (n=3).
(e) Quantification of puromycin incorporation at the indicated time points in mouse and
human NPs (0.1 µg/ml puromycin). Geometric mean ± standard deviation assayed with
fluorescently conjugated puromycin antibody (n=2).
(f) Protein content of mouse and human NPs measured by BCA quantification. Mean ±
standard deviation (n=3, including at least 20 wells; p-value < 0.0001, paired sample t-test).
951

Figure S1. Assessment of SNAP-Tag suitability to determine protein dynamics
(a) Representative FACS plot showing co-expression of OLIG2 and HA tag in OLIG2-SNAP
mouse and human NPs.
(b) Representative flow cytometry analysis of 647-SiR ligand incorporation in pulse-chase
experiments of mouse and human NPs at the indicated days of differentiation with HALO
ligands as controls.
(c) Representative flow cytometry analysis of 647-SiR ligand incorporation on day 4 (red) or
day 6 (blue) in human NPs pulse-chase experiment.
(d) Geometric mean ± standard deviation of fluorescent intensity of conjugated puromycin
antibody in mouse NPs treated with the indicated concentrations of puromycin (n=2).
(e) Geometric mean ± standard deviation of fluorescent intensity of conjugated puromycin
antibody in human NPs treated with the indicated concentrations of puromycin (n=2).
(f) Quantitation of phospho-EIF2A in mouse and human NPs treated with (mouse, red;
human, dark blue) or without puromycin (mouse, orange; human, blue) at the indicated time
points after puromycin incorporation. Data represent mean ± standard deviation (n=2).
(g) Quantification of Cell Trace incorporation of mouse and human NPs as a proxy for
protein content. (n=3)
969

Figure 2. Global proteomic differences in mouse and human neural progenitors
(a) Volcano plots showing 1791 Differentially Expressed Proteins (DEPs) in at least 2
biological replicates, with ≥ 2 FC and FDR ≤ 0.05. DEPs enriched in mouse or human are

depicted in orange (794) and blue (997), respectively. The full list of DEPs is presented in
Table S2.
(b) Bubble plot depicting the top pathways identified in pathway analysis with DAVID (Table
S3). The pathways enriched amongst human- or mouse-upregulated DEPs are shown in
blue and orange, respectively. Bubble sizes indicate the number of DEPs in each pathway.
The axes show the fold enrichment of each term compared to the background, and -log of P
value from a Fisher's test.
(c) Heatmap of the normalized scaled intensities of DEPs assigned to Gene Ontology
categories (DAVID) in mouse and human NPs. Each pathway is color coded and ordered by
p value.
(d) Abundance of proteins associated to the ubiquitin-proteasome pathway in mouse and
human NPs. DEPs are highlighted in darker color and labeled.
(e) Mean abundance of DEPs associated with the ubiquitin-proteasome pathway in mouse
and human NPs.
987
Figure S2. Quality control and differential expression of human and mouse proteins
identified by DIA-MS in NPs
(a-b) Percentage of SOX2 (a) and TUBB3 (b) positive cells in mouse and human NP
differentiations. Data represent mean ± standard deviation (mouse n=3, human n=2).
(c) Principal Component Analysis plot of the mouse and human NPs used for differential
protein expression. Each point represents a replicate ran in triplicate for DIA-MS.
(d) MA plot displaying the logged intensity ratio (M) versus the mean logged intensities (A) of
changes in protein abundances of the mouse and human NPs. The log2 of fold changes in
mean human/mouse protein abundance is shown against log10 of the averaged protein
abundance. Red dots indicate differentially expressed proteins.
(e) Boxplots of the normalised values per technical replicate for the DIA-MS in mouse and
human NPs. Each biological replicate was run in triplicate for DIA-MS.
(f) Mean abundance of DEPs in mouse and human NPs across the replicates. Proteins
mentioned in the main text and shown on the volcano plot are labeled.
(g) Mean abundance of core particles of the proteasome in mouse (orange) and human
(blue) NPs as determined by DIA-MS. Data represent mean ± standard deviation (n=3, in
triplicates).
(h) Mean abundance of regulatory particle subunits of the proteasome in mouse (orange)
and human (blue) NPs as determined by DIA-MS. Data represent mean ± standard deviation
(n=3, in triplicates).
1008
Figure 3. Dynamic SILAC proteomics identifies global differences in protein stability
(a) Schematic of dynamic SILAC combined with quantitative proteomics.
(b) Violin plot showing the estimated half-lives of mouse and human proteins in NPs. Median
half-lives for the mouse and human datasets are indicated. Quartile lines in the violin plots
are the 25th, 50th (median) and 75th percentiles.□
(c) Violin plot illustrating the fold change in half-lives between human and mouse proteins
(left), and dot plot showing the half-lives of 3402 homologous proteins in mouse and human
(right). The blue line represents a 1:1 relationship, indicating identical half-lives in both
species. The red line marks the median fold-difference in half-life between human and
mouse proteins. Violin plot as in (b).
(d) Half-life measurements of homologous proteins divided into quartiles based on protein
size (aminoacid sequence length), from smallest to largest. Orange and blue box plots

represent the values for mouse and human proteins, respectively. Grey box plots indicate
the fold difference in half-lives in human versus mouse. Box-plot: Centre line is the median;
box represents upper and lower quartiles; whiskers are1.5x interquartile range from the
upper and lower quartiles; dots are outliers.
(e) Quartiles of measured half-lives homologous proteins partitioned according to protein
abundance (average total expression level), from low to high. Orange and blue box plots
correspond to mouse and human values selected for partitioning, respectively. Grey box
plots indicate the fold difference in half-lives in human versus mouse. Box-plot as (d).
(f) Half-lives of homologous proteins associated with four GO cellular component terms.
Dashed lines indicate median half-lives of mouse and human homologues. Box-plot as in
(d).
1032
Figure S3. Additional data analysis of the SILAC dataset
(a, b) Bubble plot showing top enriched GO terms in mouse (a) and human (b) half-lives
from GSEA Preranked analysis. Positive normalized enrichment scores (NES) indicate the
enrichment in long-lived proteins, and negative NES values indicate enrichment in short-
lived proteins. Size of the points depict size of the GO term, colour indicates -log10 p-value.
(c) Human to mouse fold difference in half-lives in human homologous proteins partitioned in
quartiles according to protein size in human, from small to large. Box-plot: Centre line is the
median; box represents upper and lower quartiles; whiskers are1.5x interquartile range from
the upper and lower quartiles; dots are outliers.
(d) Dot plot showing amino acid sequence length of mouse to human homologues.
(e) Quartiles of measured half-lives in human homologous proteins partitioned according to
abundance of human proteins, from low to high. Box-plot as in (c).
(f) Dot plot showing average total expression levels of mouse and human homologues.
(g) Half-lives of homologous proteins associated with 8 GO molecular function terms. Box-
plot as in (c).
(h) Median fraction of light over time for proteins related to proteasomal categories in the
mouse (orange) and human (blue) datasets. Core proteasomal particles (n=14), regulatory
subunits (n= 16), proteasomal activators and inhibitors (n= 5). Line and shadowed areas
show best linear fit with 95% confidence intervals. Dashed lines represent median protein
half-lives in mouse (orange) and human (blue).
(i) Fraction of light over time for specific activators and inhibitors of the proteasome in the
mouse and human NP SILAC datasets. Line and shadowed areas show best linear fit with
95% confidence intervals. Dashed lines represent median protein half-lives in mouse
(orange) and human (blue) (n=3).
1057
Figure 4. Proteasomal activity is higher in mouse neural progenitors
(a-b) Schema of the experimental design for neuronal differentiation from ESCs. A Notch
inhibitor (10mM DBZ, gamma secretase inhibitor) was added to increase the differentiation
of NPs into post-mitotic neurons. In mouse, treatment was administered from day 2 to day 4
of the differentiation. On day 4, AHA pulse and chase experiments were performed to
measure protein stability in progenitors and differentiated neurons in mouse (a). Similarly,
human NPs were treated for 48 hours with the Notch inhibitor and protein stability measured
after treatment starting at day 4 (b).
(c) Normalized intensity of AHA levels in mouse and human NPs (mouse, orange; human,
blue) and post-mitotic neurons (mouse, red; human, dark blue) across timepoints. Line and

shadowed areas show best exponential fit with 95% confidence intervals (bottom); (mouse n
= 2, human n = 2).
(d) Half-live estimation of the proteome in mouse and human neural progenitors (mouse,
orange; human, blue) and post-mitotic neurons (mouse, red; human, dark blue) for datasets
in (c). Quartile lines in the violin plots are the 25th, 50th (median) and 75th
percentiles. Statistical significance (*) corresponds with < 5% of overlap between the
distributions of parameter xestimations (mouse n = 2, human n = 2).
(e) Measurements of proteasomal activity over time in mouse and human NPs. A line
connects the mean values at the measured timepoints, and the shaded areas represent the
standard deviation (n=4).
(f) Relative slope of proteasomal activity in mouse and human NPs. Data are presented as
mean ± standard deviation; unpaired sample t-test (n=4; p-value = 0.0046, unpaired sample
t-test).

1081

Figure S4. Differences in protein stability between mouse and human neural cells
(a-b) Quantifications of the proportion of neural progenitors (SOX2+) and post-mitotic
neurons (TUBB3+) after treatment compared to controls in mouse (a) and human (b)
differentiations. Data represent mean ± standard deviation (mouse n=2, human n=2).
(c) Representative flow cytometry analysis of AHA incorporation in pulse-chase experiments
of mouse and human NPs and neurons at the indicated time points.

1088

Figure 5. HALO-PROTAC addition depletes IRX3 and increases the proportion of
OLIG2 expressing neural progenitors.
(a) Schematics of HaloPROTAC targeted degradation and treatment regimes. HaloPROTAC
covalently binds to HALO and induces the 26S proteasomal degradation of IRX3-HALO
through the recruitment of the von Hippel-Lindau (VHL) E3 ubiquitin ligase.
(b) Representative FACS plots of OLIG2 expression measured in SOX2 expressing mouse
NPs in controls (DMSO, turquoise) and PROTAC (+PROTAC, violet) treated samples at 12h
and 18h post addition.
(c) Percentage of total OLIG2 positive mouse NPs at 0h, 12h and 18h post SAG addition in
controls and PROTAC treated samples (12h n=2; 18h n=3).
(d) Percentage of IRX3+;OLIG2- and OLIG2+;IRX3-* NPs during differentiation in DMSO or
+PROTAC samples. A line connects the mean values (dots), and the shaded areas
represent the standard deviation. (12h n=2; 18h n=3).

1102

Figure S5. Characterization of the IRX3-HALO mESC line.
(a) Diagram of pulse-chase treatment for IRX3::HALO ligand. 100nM of HALO JF549 ligand
was added for 1h. Media was then replaced, and mNPs collected at the indicated time
points.
(b) Confocal images of IRX3 detection and decay from the pulse-chase experiments. IRX3 is
detected with an antibody against the HA tag in blue throughout the time course, while the
fully labelled IRX3 with the HALO JF549 ligand (magenta) at 0h shows the decay of IRX3
protein at 4h and 24h post-labelling (n=2).
(c) Schema indicating medium conditions in the pulse-chase labelling experiment and
representative flow cytometry histograms of IRX3 expression and decay from the pulse-
chase experiments. The histogram on the top corresponds to total IRX3 levels detected with
an HA antibody. The lower panel corresponds to the decay of IRX3-JF549 labeled proteins
at 4h and 24h post-labelling (n=2). Scale bar 25 μm.

(d) Validation of IRX3 depletion by HALO PROTAC as compared to an Irx3-/- (IRX3[KO]) cell line. Representative histograms of IRX3 expression and depletion over time upon PROTAC addition (n=2).

(e) Representative contour plots of IRX3 depletion in 12h and 18h time courses. X-axis represents OLIG2 intensity and y-axis IRX3 intensity. Numbers in each quadrant indicate percentage of NPs quantified in DMSO (turquoise) or PROTAC (violet) conditions.

ACKNOWLEDGEMENTS

We thank Miki Ebisuya for productive discussions; Rahul Samant, Ian McGough, and all members of the Briscoe lab and Rayon lab for advice and feedback. We are grateful to David Vilchez for advice and sharing the proteasome activity protocol. Thanks to Florence Wood for performing some experiments on the regimes of treatment for the IRX3-HALO line. We thank the Babraham Institute Facilities in particular the Flow Facility, Mass Spec and Bioinformatics as well as Stores, BICS, and Tech Services. The latter were instrumental when setting up the lab. We thank the Crick Science Technology Platforms for their expertise and assistance, particularly the Genomics, the Flow Cytometry, and the Proteomics STPs.

AUTHOR CONTRIBUTIONS

T.R. and J.B. conceived the project, interpreted the data, and wrote the manuscript with input from all authors. T.R. designed and performed experiments and data analysis. S.N. designed and performed Olig2-SNAP turnover measurements, proteasome activity, and SILAC experiments and data analysis, including bioinformatic analysis. D.S. generated and characterised the OLIG2-SNAP hESC line and designed, performed and analyzed global production measurements, protein content, and global proteomic experiments. L.G.P. generated and characterized the OLIG2-SNAP and IRX3-HALO mESCs lines and designed, performed and analyzed experiments. C.A. designed and performed experiments and data analysis of IRX3-HALO mESCs. A.P. performed statistical analysis of global proteomics. S.H and M.S performed global DIA mass spec. L.Y and D.O performed SILAC mass spec analysis. H.C. and S.A. performed bioinformatic analysis of the SILAC datasets.

FUNDING

Work in JB laboratory was supported by the Francis Crick Institute which receives its core funding from Cancer Research UK (CC001051), the UK Medical Research Council (CC001051), and Wellcome (CC001051); by the European Research Council under European Union (EU) Horizon 2020 research and innovation program grant 742138; and by UK Human Developmental Biology Initiative (Wellcome, 215116_Z_18_Z). Work in the laboratory of TR is supported by the Babraham Institute's BBSRC Institute Strategic Programmes Grant (ISPG) [BB/Y006909/1]; the Engineering and Physical Sciences Research Council [EP/X021521/1]; the Babraham Institute's BBSRC Core Capability Grant (CCG) [BB/CCG2210/1], and the Institute Development Grant [BB/IDG2210/1].

COMPETING INTERESTS

The authors declare no competing or financial interests.

Figure S5