Limited directional selection but coevolutionary signals among imprinted genes in *A. lyrata.*

Audrey Le Veve[1,2], Ömer İltaş[2,3], Julien Y. Dutheil[4], Clément Lafon Placette[2, *]

[1] IHPE, Univ Perpignan Via Domitia, CNRS, IFREMER, Univ Montpellier, Perpignan, France

[2] Department of Botany, Faculty of Science, Charles University, Prague, Czech Republic

[3] Max Planck Institute of Molecular Plant Physiology: Potsdam, Brandenburg, DE

[4] RG Molecular Systems Evolution, Department of Theoretical Biology, Max Planck Institute for Evolutionary Biology, August-Thienemann-Straße 2, 24306 Plön, Germany

[*] Corresponding author: audrey.le-veve@univ.perp.fr

Abstract

Genomic imprinting is a form of gene regulation leading to the unequal expression of maternal and paternal alleles. The main hypothesis invoked to explain the evolution of imprinted genes is the kinship theory, which posits a conflict between parental genomes over resource allocation in progeny. According to this theory, such conflicts select for parent-of-origin–dependent expression of genes involved in resource allocation. How such conflicts translate into signatures of selection at coding or regulatory sequences remains model-dependent and is not explicitly predicted by the kinship theory. However, most studies addressing selection in imprinted genes in flowering plants, particularly those based on population-genomic or phylogenetic analyses, have focused on self-fertilizing species, where conflicts over resource allocation are predicted to be weak. Consequently, the impact of the kinship theory on the evolution of imprinted genes remains largely unexplored in systems where parental conflict is expected to be strong. Furthermore, potential coevolution between antagonistically acting imprinted genes, as proposed in extensions of parental conflict models, has not yet been tested empirically.

Using combined phylogenetic and population genomic approaches, we investigated signatures of selection on imprinted genes across the Brassicaceae family and in autogamous and allogamous populations of *Arabidopsis lyrata*, and searched for evidence of coevolution among imprinted genes. We found that endosperm-expressed genes exhibited signals of balancing selection across Brassicaceae and within allogamous populations, consistent with models of unresolved intralocus conflict. These population-level signals varied with the mating system, in line with expectations that parental conflict is reduced

under self-fertilization. Moreover, phylogenetic analyses indicated signatures of purifying (negative) selection acting on imprinted genes. However, the population-level signatures of selection were independent of the mating system and showed limited concordance with kinship predictions, possibly due to stronger selection acting on expression than on coding sequences. Finally, we identified coevolution between imprinted genes, although not at specific sites, suggesting that interactions beyond protein sequence may contribute to this process.

Introduction

The kinship theory, or the parental conflict theory, was first proposed by Haig and Westoby (1989) in the context of endosperm development and further developed in subsequent reviews in 1991. It posits that in species where the maternal parent nourishes progeny sired by multiple males, maternal fitness is maximized by distributing resources equally among offspring, whereas paternal fitness is enhanced by securing more resources for their own offspring.

While the kinship theory predicts that conflicts between maternal and paternal genomes select for expression of genes related to growth that depends on parent-of-origin, it does not explicitly address the expected patterns of allelic polymorphism at these loci (Haig and Graham 1991; Moore and Haig 1991; Wilkins and Haig 2001). Consistent with this framework, experimental crosses in *Arabidopsis thaliana* revealed dosage effects on seed size, although these patterns may reflect interactions between cytoplasmic inheritance, dosage imbalance, and imprinting (Scott et al., 1998). Balancing selection and elevated polymorphism have been predicted under alternative models of unresolved intralocus conflict acting on alleles with antagonistic fitness effects (Day and Bonduriansky 2004; Kasimatis et al., 2017). Whether similar signatures of balancing selection may arise at genes involved in resource allocation prior to, or independently of, the evolution of imprinting remains an open empirical question. Notably, signatures of balancing selection in genes expressed in the endosperm (the nourishing tissue of the embryo in the seed) have not been systematically characterised, despite their potential relevance for understanding the evolutionary consequences of parental conflict.

The kinship theory remains a leading explanation for the evolution of imprinted genes (Haig 1997; Spencer and Clark 2014; Patten et al., 2014). Imprinting is an epigenetic form of regulation in flowering plants and mammals in which allele expression depends on parental origin (Haig 1997; Wilkins 2011). By creating hemizygous expression of parental alleles, the kinship theory argues that imprinting evolves to resolve parental conflicts over resource allocation. Consequently, imprinted genes are expected to be disproportionately expressed in the endosperm in plants, the tissue where parental conflict operates (Haig 1997; Haig 2000). Thus, the proportion of Maternally Expressed Genes (MEGs) and Paternally Expressed Genes (PEGs) upregulated in the endosperm should exceed the genome-wide expectation derived from sampling all expressed genes. According to this hypothesis, many imprinted genes appear to be involved in resource accumulation in the endosperm and metabolic processes (Raissig et al., 2011). These theoretical expectations and previous

empirical observations justify testing whether MEGs and PEGs are disproportionately
upregulated in endosperm relative to random expectation.

Theoretical models of selection on one or two genes involved in resource allocation predict
that: (1) imprinting sets distinct optimal expression levels for paternally expressed genes
(PEGs) and maternally expressed genes (MEGs; Haig 1997); Specifically, PEGs are
predicted to evolve higher optimal expression levels than MEGs, because paternal alleles
benefit from increased resource acquisition, whereas maternal alleles tend to limit growth
(Haig 2000; Patten et al., 2014). Accordingly, we expect a higher expression per active gene
copy in PEGs, followed by MEGs and non-imprinted endosperm genes; (2) PEGs and MEGs
are expected to experience selection favouring increased or reduced resource acquisition,
respectively (Haig 2000); and (3) imprinting at some locus promoting growth may drive
antagonistic coevolution with some function suppressing growth (Wilkins and Haig 2001).
However, these models focus on *Insulin-like Growth Factor 2* (*IGF2*) and *Insulin-like Growth*
Factor 2 Receptor (IGF2R) genes, which regulate foetal growth in mammals, without
investigating the expected signatures of selection for endosperm genes. More recent
theoretical work has further emphasized that selection associated with parental conflict may
primarily act on loci controlling imprinting or gene regulation, rather than on the coding
sequences of imprinted genes themselves, potentially weakening expected genomic
signatures at these loci (Flintham et al., 2025).

Empirically, phylogenetic signals of directional selection have been reported for the *MEDEA*
locus in the allogamous species *Arabidopsis lyrata*, but not in the autogamous sister species
A. thaliana (Spillane et al., 2007; Miyake et al., 2009), and for PEGs at the population level in
the autogamous Brassicaceae species *Capsella rubella* and *A. thaliana* (Hatorangan et al.,
2016; Tuteja et al., 2019). However, MEGs did not show comparable signals, possibly due to
technical challenges in detecting selection if selection primarily targets gene expression
levels. Indeed, a central debate has emphasized that kinship theory primarily concerns
selection on optimal expression levels, potentially acting on alleles related to regulatory
mechanisms in parental genomes, rather than the level of polymorphism in the coding
regions of imprinted genes (Patten et al., 2014). Selection acting on gene expression may
still leave detectable traces in amino acid or codon usage patterns (Patten et al., 2014).

The incongruent signals detected in imprinted genes may also be due to polymorphism
dynamics shaped by the mating system. The kinship theory assumes that progenies are
sired by multiple males and that imprinted genes are preferentially expressed in the
endosperm (Haig and Westoby 1991; Haig 2000; Brandvain and Haig 2005). The transition
to autogamy may alter selection acting on genes, potentially leading to more intense

directional selection or relaxed selection (Soliman and Coughlan 2024). However, the
empirical studies cited above did not assess endosperm-specific expression of the imprinted
genes and only focused on autogamous species. Consequently, the evolutionary signatures
of selection on imprinted genes and the contribution of the kinship theory to their evolution
remain unresolved in flowering plants.

Extensions of the kinship theory also predict a potential coevolution between antagonistic
imprinted genes (Wilkins and Haig 2001; Willi 2013), yet no phylogenetic signal was
detected for *IGF2* and *IGF2R* (McVean and Hurst 1997), and coevolution among
endosperm-expressed genes remains unknown. Population genomic methods can detect
coevolving sequences based on allele frequency variation; however, they may be biased by
demographic or population structure effects. More integrative approaches based on
phylogenetic comparisons can identify co-occurring changes in candidate coevolving genes
(Dutheil and Galtier 2007).

Here, we test these predictions by combining expression, phylogenetic and population-
genomic analyses to evaluate whether imprinted genes in *A. lyrata* show evolutionary
signatures compatible with theoretical expectations derived from models of parental conflict.
We compared imprinted genes in allogamous *A. lyrata* (Klosinska et al., 2016) with non-
imprinted endosperm genes, using phylogenetic data from 23 Brassicaceae species and
genomic data from nine allogamous populations and nine closely related autogamous
populations of *A. lyrata* from North America (Ross-Ibarra et al., 2008). We tested for: (1)
variations in expression patterns and/or codon usage; (2) reduced divergence and
polymorphism, as expected under some models of directional selection, and (3) correlations
between evolutionary rates or diversity metrics used as exploratory indicators of evolutionary
covariance expected in case of coevolution. Our results indicate that endosperm-expressed
genes display patterns compatible with unresolved intralocus conflict over resource
allocation, and that the transition to autogamy promoted directional selection on these
genes. Imprinted genes show evidence of ancient negative directional selection, codon
usage bias, and substantial coevolution. However, the signals at the level of population are
less compatible with the kinship theory, possibly due to stronger selection acting on
expression than on coding sequences or by a recent relaxed selection on these genes in
North American populations.

Results

General analytical framework

Parental conflict is expected to occur primarily in the endosperm. Using transcriptomic data from 19,425 genes across five tissues (leaf, root, pollen, pistil, and endosperm), we identified 2,732 specifically upregulated genes in the endosperm of *A. lyrata* among 31,073 genes analysed. Among these upregulated genes, 15 were Maternally Expressed Genes (MEGs) and 14 were Paternally Expressed Genes (PEGs) from the 27 MEGs and 46 PEGs previously identified in *A. lyrata* (Klosinska et al., 2016). Here and throughout the Results, we use the term 'gene' to refer to annotated nuclear protein-coding loci for which expression could be quantified from RNA-seq data. Between *A. lyrata* and *A. thaliana*, 40% of MEGs (6/15) and 50% of PEGs (7/14) identified in our dataset were conserved as imprinted (Table S1).

Throughout the Results section, we compare four categories of genes: MEGs and PEGs that are upregulated in the endosperm (referred to as "imprinted genes" if considered together), the 2,703 non-imprinted genes specifically upregulated in the endosperm ("endosperm genes"), and a set of 100 control genes, randomly sampled from the ubiquitously expressed genes, and serves as a baseline for genome-wide patterns of selection unrelated to parental conflict. The control panel was chosen following the strategy of sampling based on expression used in previous imprinting studies, rather than by matching genomic characteristics across groups. Differences in gene size, Coding DNA Sequence (CDS) proportion, and Guanine–Cytosine (GC) content among panels were therefore quantified and considered when interpreting downstream evolutionary analyses.

For each population genetic and phylogenetic statistic, genes with insufficient informative sites were excluded. Then, significance was evaluated using 10,000 bootstrap resampling matching the size of focal group framework, because this method is insensitive to difference in sample size. Rather than repeating the method in each subsection, we applied a single unified framework in which significance was assessed by comparing observed means to the central 95% of the empirical bootstrap distribution (i.e. 2.5^{th}–97.5^{th} percentiles). This approach was used consistently across analyses of expression, codon usage, phylogenetic metrics, and population polymorphism. Although the figures share a similar visual layout for consistency, each statistic captures a distinct evolutionary dimension allowing direct comparison of patterns across gene categories while retaining analytical complementarity.

We first compared gene size, CDS content (%) and GC content (%) across groups, as these
features may influence polymorphism patterns and generate bioinformatic artefacts. Control
genes and MEGs were significantly smaller than PEGs and endosperm genes, and CDS
content was significantly higher in endosperm and imprinted genes relative to control genes.
GC content was also lower in endosperm and imprinted genes than in control genes (Table
S2). These differences in gene size, CDS proportion and GC content were quantified and
considered when interpreting all evolutionary analyses. Nevertheless, they were not
corrected for directly, and we retained the control panel to be congruent with methods used
in previous studies in the autogamous species *C. rubella* and *A. thaliana* (Hatorangan et al.,
2016; Tuteja et al., 2019). In practice, all evolutionary comparisons evaluating predictions of
the kinship theory were performed relative to endosperm-upregulated non-imprinted genes,
which represent the expression-matched and biologically relevant background for imprinting
analyses. The control panel was used only to describe genome-wide patterns of variation.

To evaluate whether these genomic differences could create artificial signals of selection,
we performed several robustness checks. First, for population-genomic analyses, we verified
that the filter based on Single Nucleotide Polymorphism (SNP) density did not preferentially
exclude short, GC-poor, or low-diversity genes. Second, by comparing all statistics against
empirical null distributions derived from endosperm-upregulated genes, we ensured that
each test was evaluated against a biologically appropriate baseline sharing similar
expression patterns, thereby reducing the influence of residual genomic confounders
unrelated to imprinting or parental conflict. Finally, we compared different linear regression
models using each statistic of interest as the response variable, and evaluated model
support using the Akaike Information Criterion (AIC). The full model included gene group,
gene length, CDS proportion and GC content as explanatory variables. We then evaluated a
series of reduced models in which one parameter was removed at a time and retained the
model with the lowest AIC value. This procedure was iterated until the best-supported model
was identified. This approach allowed to determine (1) whether gene group remained a
significant predictor in the best model and (2) whether genomic features accounted for the
observed trends in the analysed statistics.

In previous study (Tuteja et al., 2019), validation of the kinship theory to explain imprinting
was performed following another logic, based on enrichment, which evaluated not the global
behaviour of imprinted genes but the proportion of them exhibiting extreme signatures
compatible with parental conflict. Thus, we tested whether directional selection occurred
more frequently in imprinted genes than in the empirical distribution of endosperm genes.
Candidates for directional selection were defined as the 5% most extreme values (bimodal

test) for branch length, Tajima's D, or the ratio of non-synonymous to synonymous
nucleotide diversity (π_{NS}/π_S) in allogamous populations. Under the null hypothesis of no
enrichment, imprinted genes should fall into this class with probability 0.05. We therefore
tested whether the observed proportion exceeded 5% using independent binomial tests.

Coevolution was assessed using three complementary measures (Pearson's correlation
coefficients for nucleotide diversity π, branch lengths, and Robinson–Foulds (RF) distances,
for each pair of PEGs, MEGs, and endosperm genes), and only correlations exceeding
thresholds defined from pair of endosperm genes (1st or 99th percentiles) were retained.
Finally, we tested if the number of correlations found between PEGs and MEGs and the
number of imprinted genes involved were significantly higher than expected from 1,000
random samplings of 29 endosperm genes.

Imprinted genes are preferentially expressed in the endosperm

Under the kinship theory, both MEGs and PEGs are expected to be disproportionately
represented among endosperm-upregulated genes, because imprinting evolves specifically
in tissues exposed to parental conflict. We therefore first tested whether the proportion of
upregulated MEGs and PEGs exceeded expectations derived from random sampling of the
transcriptome. The proportions of MEGs (0.56) and PEGs (0.30) upregulated in the
endosperm were both significantly higher than expected from resampling of all genes (p
values < 1e-04), and MEGs were more frequently upregulated than PEGs (based on
resampling of imprinted genes; p value =0.03).

Because PEGs are generally assumed to promote offspring resource acquisition, while
MEGs tend to restrict it, theory predicts higher optimal expression levels for PEGs, followed
by MEGs, compared with non-imprinted endosperm genes. We therefore tested whether
expression per expressed allele differed among groups after correcting for gene dosage.
After adjusting expression for gene copy number (three expressed copies for non-imprinted
endosperm genes, two for MEGs, and one for PEGs), both PEGs and MEGs displayed
significantly higher expression levels relative to endosperm genes, with 3.18- and 1.71-fold
increases, respectively (bootstrap tests; p-values < 1e-03; Fig 1).

Using AIC, we found that the best model to explain this expression pattern (AIC = -4633)
was the model that included the gene group, but also the gene size (coefficient = -5.4e-05),
the CDS proportion (coefficient = -2.34e-03) and the GC proportion (coefficient = 2.66e-02),
whereas the worst model (AIC = -2454) was the one that exclude only the group of genes
(AIC = -2454).

We next assessed whether this increased expression was associated with codon or amino
acid usage bias. We defined the "overused" and. "underused" codons as codons with a
proportion higher and lower respectively than in the 95th percentiles of the 1,000 resample
of 29 endosperm genes. Codon and amino acid usage differed significantly between
imprinted and endosperm genes, although only two biases were shared by PEGs and MEGs
(under-usage of GTC and over-usage of tryptophan; Table S3). However, Predicted High
Expression (PHE) scores of codon, derived from overexpressed genes in *A. thaliana* (Sahoo
et al., 2019), did not differ significantly across biased and unbiased codons (all p values >
0.09), indicating that codon usage differences were not associated with expression variation.

Figure 1: Imprinted genes are associated with higher expression levels in endosperm genes. The dots represent the mean expression level obtains for each gene copy. The red dots represent the mean expression level for each group of genes. The red lines represent 2.5th and 97.5th percentiles of the means obtained for endosperm genes after

1,000 resampling of 29 genes (number of imprinted genes considered in the study). If the
observed means of PEG and MEG are outside of these red lines, they are considered
significantly different from the mean in endosperm genes, following the bootstrap approach.

The MEGs and PEGs exhibit phylogenetic signals of negative selection in Brassicaceae

To explore phylogenetic patterns, we analysed homologous genes in *A. lyrata* and 21
Brassicaceae species (Table S4). As a proxy for gene turnover, we compared the number of
species with homologous sequences across groups. Thus, endosperm genes were
conserved across significantly more species than other categories, and PEGs more than
control genes (Fig 2; Table S5). The best model to explain this pattern (AIC = 8696) was the
model that included the gene group, the gene size (coefficient = 9.74-04) and the GC
proportion (coefficient = 0.19), whereas the worst model (AIC = 9104) was the one that
exclude the gene size.

Gene-tree topologies of MEGs and PEGs did not differ significantly more from the species
tree than endosperm genes (based on RF distances; bootstrap tests; p value = 0.60 for
PEGs; p value = 0.34 for MEGs), so branch lengths (proxy for evolutionary rates; Fig. S1)

were estimated directly from the species-tree topology for genes having homologous genes
in at least three species (including *A. lyrata*). Thus, endosperm and imprinted genes show
significantly longer branches than in control genes (Fig 2; bootstrap; Table S5), with
significantly longer branch in PEGs and in endosperm genes. The best model to explain this
pattern (AIC = -15231) was the model that included the gene group, the gene size
(coefficient = 9.73e-06), the CDS proportion (coefficient = -8.10e-04) and the GC proportion
(coefficient = 9.63e-04), whereas the worst model (AIC = -14976) was the one that exclude
the gene size.

Finally, the mean ratio of non-synonymous to synonymous substitution rate (D_{NS}/D_S)
between *A. lyrata* and the other Brassicaceae species decreased significantly in imprinted
genes compared to endosperm genes (Table S5), indicating a signal of purifying (negative)
selection. The best model to explain this pattern (AIC = 13382) was the model that included
the gene group and the GC proportion (coefficient = 0.18), whereas the worst model (AIC =
13401) was the one that exclude the group of genes.

Together, these phylogenetic patterns indicate long-term evolutionary constraints in
endosperm genes, a signal that can arise under ancient balancing selection or strong
functional conservation. Importantly, these phylogenetic inferences reflect deep evolutionary
timescales and therefore cannot be directly compared to the signatures of balancing
selection at the population level, which we analyse in the next section.

Figure 2: Imprinted genes present phylogenetic signals of negative selection
compared to endosperm genes across the Brassicaceae. *A: Variations of the number of*
species with homologous genes by the group of genes. B: Variations of the mean branch
length by gene based on the phylogenetic tree obtained by Hasegawa–Kishino–Yano 1985
model based on the 1055 endosperm genes found in all species. C: Variations of the D_{NS}/D_S
by the group of genes with a focus on values lower than 5 (top right). Two extremes values
(>200) were removed. The grey dots represent the mean value obtained for one gene. The

red dots represent the mean in each group of gene. The dashed lines represent 2.5th and

97.5th percentiles of the means obtained for endosperm genes after 1000 resamples of 100

genes (size of group of control genes). If the real mean of control genes was not between

these dashed lines, the value was considered significantly different from that of the

endosperm genes, following bootstrap approach. The red lines represent 2.5th and 97.5th

percentiles of the means obtained for endosperm and control genes after 1,000 resamples of

29 genes (the number of imprinted genes considered in the study). If the real means of PEG

and MEG were not within these red lines, the values were considered significantly different

from those in endosperm or control genes, using the bootstrap approach.

Imprinted genes show elevated π_{NS}/π_S independent of the mating system

To evaluate the effect of the mating system and imprinting on polymorphism, we compared

Tajima's D, π and the π_{NS}/π_S for imprinted, endosperm and control genes across nine

allogamous and nine autogamous populations (Table S6). A previous study reported that the

mating system affects seed size through pollen in these populations (Willi, 2013), suggesting

a reduced intensity of parental conflict in autogamous populations.

Under directional selection, reductions of Tajima's D, π and π_{NS}/π_S are expected, whereas

balancing selection is expected to increase these statistics. Because calculations of these

statistics require sufficient polymorphism in all populations, genes with fewer than 0.5%

polymorphic sites were excluded from Tajima's D, and genes with $\pi_S = 0$ from π_{NS}/π_S

analyses, resulting in a significantly higher retention rate for endosperm genes than for

control genes for Tajima's D and π_{NS}/π_S (binomial tests; Table S7). However, more

importantly, these filters retained similar proportions of genes across endosperm and

imprinted genes, and across imprinted and control genes (Table S7). Thus, the different

signatures of selection detected in Tajima's D and π_{NS}/π_S for the imprinted genes, relative to

endosperm genes in the allogamous and autogamous populations in the following sections,

cannot be attributed to unequal gene retention nor to systematic biases against low-diversity

loci, supporting the interpretation that these signals reflect genuine biological differences

rather than filtering artefacts.

In allogamous populations, Tajima's D was significantly higher in endosperm genes than in

control genes, while π and π_{NS}/π_S showed no significant differences. Imprinted genes

showed an overall increase in π_{NS}/π_S relative to non-imprinted endosperm genes (Fig 3;

Table S8). The best model to explain this pattern of Tajima's D (AIC = 9322) was the one

that included only the gene size (coefficient = -1.59e-04), whereas the worst model (AIC =

9337) was the one that excluded only the gene size. For the π, the best model (AIC = -

36369) was the one that included the gene size (coefficient = -9.89e-08) and the GC
percentage (coefficient = -5.59e-05), whereas the worst model (AIC = 9337) was the one
that excluded only the gene size. Finally, the best model to explain this pattern of π_{NS}/π_S
(AIC = -721) was the one that included only the proportion of CDS (coefficient = 2.21e-03),
whereas the worst model (AIC = 9337) was the one that included all the parameters.

In autogamous populations, Tajima's D in endosperm genes no longer differed from controls
(Table S8). Moreover, MEGs and endosperm genes exhibited decreased π_{NS}/π_S compared
to control genes (Table S8). These trends in Tajima's D and π_{NS}/π_S are explained by the
mating system that only significantly affects endosperm genes (Kruskal-Wallis tests; Table

S9).

Figure 3: Imprinted genes present an unusual decrease of Tajima's D associated with
an increase of π_{NS}/π_S in allogamous populations compared to endosperm genes. *A:*
Variations of Tajima's D among genes group. Only the genes with a mean between -5 and 5
are indicated. B: Variations of the π_{NS}/π_S among genes group. Only the genes with a mean
between 0 and 1.1 are indicated. The grey dots represent the mean value obtained for one
gene. The red dots represent the mean of each gene group. The dashed lines represent the
2.5th and 97.5th percentiles of the means obtained for endosperm genes after 1000
resamples of 100 genes (size of the control gene group).. If the real mean of control genes
was not between these dashed lines, the value was considered as significantly different than
in endosperm genes. The red bars represent the 2.5 and 97.5th percentiles of the means
obtained for endosperm and control genes after 1,000 resamples of 29 genes (number of

imprinted genes considered in this study). If the real means of PEG and MEG were not
between these red lines, the values were considered significantly different from those in
endosperm or control genes, following the bootstrap approach.

A limited number of MEGs exhibited strong signals of directional selection.

To assess whether loci subject to imprinting show an enrichment in signatures of directional
selection, we tested whether they occur more frequently than expected in the extreme tail of
the empirical distribution of endosperm genes. Candidates for directional selection were
defined as the 5% most extreme values (bimodal test) for branch length, Tajima's D, or
π_{NS}/π_S in allogamous populations. However, none of these proportions were significantly
higher than expected (p = 0.66, 1.00 and 1.00 respectively).

We identified only four MEGs potentially under strong directional selection (Table S1). Two
MEGs were associated with a reduced branch length (Table S1): one lacked homologous
gene in *A. thaliana*, the second encoded an AGAMOUS-like protein expressed during
proliferative endosperm development (Day et al., 2008), involved in female gametophyte,
seed and early endosperm development (Bemer et al., 2010; Zhang et al., 2018). One MEG
identified by a reduced Tajima's D encodes a F-box domain–containing protein regulated by
non-CG DNA methylation in *A. thaliana* (Garro et al., 2025) and associated with the
regulation of stress-responsive genes (Papdi et al., 2008). Finally, one MEG encoding a Dof-
type zinc finger protein involved in seed coat formation (Zou et al., 2012) shows a reduced
π_{NS}/π_S.

Clear trend of coevolution among imprinted genes in Brassicaceae

Imprinting could promote coevolution between genes favouring paternal or maternal fitness
in an arms race scenario. To test this hypothesis, we investigated potential coevolutionary
signals between PEGs, MEGs and endosperm genes, based on (i) nucleotide diversity (π)
within populations, (ii) branch length in phylogenetic trees, and (iii) RF distances between
tree topologies. Because the coevolution hypothesis is more closely related to noncoding
regions, we focused on π rather than π_{NS}/π_S.

First, using thresholds defined from endosperm genes (−0.66 and 3.20 for 1 and 99% of
distribution), 13 significant Pearson correlation coefficients involving 6 PEGs and 9 MEGs
were detected. Both the number of PEGs-MEGs correlations and the number of imprinted
genes involved exceeded expectations (p-values= 4.5e-02 and <1e-03 respectively; Table
S10). Considering endosperm genes significantly correlated with these imprinted genes, as
well as indirect links between imprinted and endosperm genes, the resulting network

contained 14 MEGs, 13 PEGs, and 1,643 endosperm genes (=1,670 genes; Fig 4.A; Table
S10). This network did not show enrichment for specific Gene Ontology (GO) terms
compared to random networks of equal size. Notably, one PEG encoding a chloroplast-
localized protein required for embryonic development (Bryant et al., 2010) was correlated
with more genes than 99% of endosperm genes (Table S1 and S10).

Following the same methodology, the thresholds based on branch length (–0.64 and 0.89)
identified 16 significant correlations among 6 PEGs and 9 MEGs. Both the number of PEGs-
MEGs correlations and the number of imprinted genes involved exceeded expectations (p-
values = 8e-03 and 3e-03 respectively; Table S11). None of these correlations was in
common with the previous 13 significant diversity correlations. Considering endosperm
genes significantly correlated with these imprinted genes, the resulting network contained 14
MEGs, 14 PEGs, and 1,027 endosperm genes (=1,055 genes; Fig 4.B; Table S11). This
network was enriched for the GO categories "*negative regulation of catalytic activity"* and
"*syntaxin binding"* (Table S12). One MEG in this network, encoding the AGAMOUS-like
protein previously mentioned, was correlated with more genes than 99% of endosperm
genes (Table S11).

Finally, we investigated gene pairs that showed high tree similarity (RF = 0) but stronger
deviation from the species topology than 95% of endosperm genes (RF > 0.71 with species
topology; Fig. S1). Based on this approach, the topology of only one PEG encoding an F-box
protein was associated with six endosperm genes (Fig. 4.C; Table S13). This small network
was enriched for the GO term *DNA-binding transcription factor activity* (Table S12).
Moreover, the PEG exhibited a significantly higher number of associations than 99% of non-
imprinted endosperm genes (Table S1; Table S13).

Across all analyses, 22 pairs including 29 imprinted genes were identified. To test for
coevolution between specific mutations within these pairs, we applied the clustering
approach implemented in the Cosubstitution Mapping (CoMap) package (Dutheil and Galtier,
2007).

Specifically, we searched for groups of 2-10 coevolving sites within each candidate pair of
imprinted genes. To limit false positives, the number of such groups detected in candidate
pairs was compared with that observed in non-candidate pairs of imprinted genes that were
not associated in the previous networks and were present in at least three species. CoMap
did not detect an excess of coevolving residues in candidate pairs relative to non-candidate
imprinted pairs (Table S14). These directly correlated pairs of imprinted genes were
therefore not associated with elevated site-level coevolution.

A) Network based on polymorphism

B) Network based on branch length

C) Network based on topology

423 **Figure 4: Networks of the imprinted and endosperm genes based on A) polymorphism in population, B) branch length and C)**
424 **phylogenetic topologies across the Brassicaceae.** *A) Network based on the Pearson's correlations of π found on genes in the populations.*
425 *B) Network based on the Pearson's correlations of the branch length for each gene estimated with the global topology. C) Networks based on*
426 *the RF distance of topologies. The black and grey lines represent the negative and positive correlations respectively. Blue dots: PEG. Red dots:*
427 *MEG.   White   circles: non-imprinted   endosperm   genes.*

Discussion

The kinship theory remains the prominent selective framework for explaining the evolution of genomic imprinting (Patten et al., 2014). According to the kinship theory, imprinting is expected to evolve in contexts of multiple mating where parental genomes have conflicting interests over resource allocation, and alleles at loci subject to genomic imprinting are therefore expected to be disproportionately expressed in the endosperm. The footprint of parental conflict on genome sequences translates into signatures based on sequence (*e.g.,* polymorphism or divergence) is model-dependent and may involve selection acting on regulatory mechanisms rather than coding sequences of loci subject to imprinting. Several models have further proposed that the resolution of parental conflict through imprinting (Wilkins and Haig 2001; Mills and Moore 2004), and related intralocus conflict models, could be associated with directional selection and/or coevolution among antagonistically acting loci (see also Day and Bonduriansky 2004). However, a selection acting primarily on alleles affecting gene expression may attenuate sequence-based signatures, and their sensitivity to mating-system variation remains unclear.

Here, we tested these predictions by comparing imprinted and non-imprinted endosperm genes in *A. lyrata*, an allogamous species that has shifted to autogamy in some populations from North America, and assessing the impact of the mating system shifts. While several figures follow a unified graphical structure to facilitate comparison across gene categories, the underlying analyses encompass expression-level differences, population genetic statistics, phylogenetic metrics, and coevolutionary patterns, providing a broader evaluation of theoretical expectations derived from models of parental conflict and imprinting (Miyake et al., 2009; Hatorangan et al., 2016; Tuteja et al., 2019). Although we detected patterns consistent with predictions of kinship theory and related intralocus conflict models, the signals differed across statistics and were not consistently replicated among analyses. This suggests that selection associated with parental conflict may act heterogeneously across loci or primarily through regulatory rather than coding changes, and underscores the importance of integrating multiple lines of evidence when assessing evolutionary hypotheses of imprinting.

Patterns compatible with unresolved intralocus conflict in endosperm genes

In the absence of genomic imprinting, genes involved in unresolved intralocus conflict are expected to evolve under balancing selection (Day and Bonduriansky 2004; Sylvestre et al., 2023). In line with this expectation, non-imprinted endosperm genes were more widely conserved across Brassicaceae, suggesting lower turnover, and displayed longer

phylogenetic branches (Edwards, 2009; Kasimatis et al., 2017; Llaurens et al., 2017). This
interpretation is supported by model comparisons showing that the gene group contributes to
explaining variation in these statistics. In addition, these genes showed elevated Tajima's D
values in allogamous populations. However, the absence of a parallel increase in
polymorphism, and the absence of this parameter in the best model predicted by AIC,
suggests that this signal may not solely reflect balancing selection.

Hypotheses tested	References	Statistic analysed	Expectation on endosperm genes	Expectation on imprinted genes
Imprinted genes are related to tissue implicated on resource's allocations	Haig, 1997	Proportion of endosperm genes	NA	**Increase**
Imprinted genes show optimized expression levels	Patten et al., 2014	Expression by copy	NA	**Upregulated with PEGs > MEGs***
Selection on expression may promote codon and amino acid bias in imprinted genes	Patten et al., 2014	Usage of each codon and amino acid		**Increase for some*, decrease for others***
Unsolved intralocus conflict in imprinted genes may maintain polymorphism across species	Day and Bonduriansky 2004	Number of species with homologous genes	**Increase***	**Decrease***
Reduction of parental conflict in imprinted genes promotes directional selection	Day and Bonduriansky 2004	Evolutionary rate by branch length	**Increase***	Increase if positive*, **decrease if negative* ****
	Wilkins & Haig 2001; Mills & Moore 2004	Sens of selection by D_{NS}/D_S	**No variation***	Increase if positive*, **decrease if negative***
Reduction of intralocus conflict in imprinting genes promotes reduced polymorphism in allogamous population by directional selection, more intensively in non-synonymous sites	Day and Bonduriansky 2004	Tajima's D	**Increase***	Decrease*
		π	Increase*	Decrease*
	Haig 2000	π_{NS}/π_S	**No variation***	Decrease*
Reduction of intralocus conflict in autogamous populations may be associated with directional selection on CDS of non-imprinting genes	Brandvain & Haig 2005; Soliman & Coughlan 2024	Tajima's D, π_{NS}/π_S	**Decrease**	**No variation if already fixed**, increase if relaxed selection
Possible antagonistic coevolution between genes (arms race)	Moore and Haig, 1991	π, branch length, topology by RF distance	NA	**Excess of PEG–MEG correlations***

Conflict over resource allocation is predicted only in multiple mating systems (Haig 2000).

This parental conflict is expected to be reduced in autogamous systems, which may alter

selective regimes acting on endosperm-expressed genes. In line with this, Tajima's D and

π_{NS}/π_S decreased in these populations, suggesting that reduced parental conflict may be

associated with patterns compatible with directional selection and that the effect of the

mating system is detectable even in populations that have recently and partially transitioned

to autogamy (Ross-Ibarra et al., 2008; Foxe et al., 2010; Hough et al., 2013; Willi et al.,

2018).

It is important to emphasize that mainly the phylogenetic analyses, but only Tajima's D for

the analysis at the population level, suggest signals compatible with balancing selection in

endosperm genes. Phylogenetic patterns reflect long-term evolutionary processes,

potentially including ancient or persistent balancing selection, whereas the statistics at the

population level detect more recent forms of balancing selection and are sensitive to

demography, recombination, and reduced polymorphism. Therefore, the absence of

consistent evidence at the population level does not contradict the phylogenetic signal but

rather indicates that any balancing selection acting on these loci is either ancient, weak, or

too localized to be detected by gene-wide summary statistics.

Importantly, the absence or inconsistency of the signatures at imprinted loci at the population

level does not contradict the existence of parental conflict. Recent models suggest that

selection may predominantly target regulatory mechanisms controlling imprinting or

expression levels, rather than coding variation at imprinted genes themselves, thereby

limiting detectable signatures of selection in analyses based on coding sequence (Flintham

et   al.,   2025).

Codon usage bias and modulation of expression in imprinted genes

Kinship theory predicts that imprinted genes should be preferentially expressed in the

endosperm (Haig 2000; Patten et al., 2014). Consistent with this prediction, imprinted genes

were enriched among endosperm-expressed genes. Interestingly, the MEGs are more

strongly associated with endosperm expression than PEGs, as observed in maize (Eaux et

al., 2013). Yet, more than half were also upregulated in other tissues suggesting that distinct

selective pressures unrelated to parental conflict may act on them.

An important debate suggests that detecting directional selection in imprinted genes is

difficult because selection may primarily target gene expression (Patten et al., 2014).

Consistent with this idea, both PEGs and MEGs showed increased expression per gene

copy, and gene group was the factor most frequently retained by AIC-based model selection.
Interestingly, the PEGs are associated with increased expression levels, despite gene size
reducing expression efficiency globally (Grishkevich and Yanai, 2014), including in our study.

Moreover, imprinted genes displayed codon usage biases relative to endosperm genes,
however, we were not able to determine if these biases enhance or inhibit the expression of
imprinted genes (Sahoo et al., 2019). More studies with direct analyses of the effect on
expression of the different codon usage are required to verify this hypothesis in the future.

Phylogenetic signals of ancient purifying selection in imprinted genes.

Several models related to parental conflict have proposed that the resolution of conflict
through imprinting could be associated with directional or purifying selection (Wilkins and
Haig 2001; Mills and Moore 2004). In line with this idea, our phylogenetic analyses showed
that imprinted genes exhibit lower conservation of homologous genes across Brassicaceae
species, likely reflecting a higher turnover driven by accelerated evolutionary rates.
Furthermore, imprinted genes displayed reduced D_{NS}/D_S, indicative of strong purifying
selection, and the group of genes is the most important parameter in models to predict this
pattern.

This phylogenetic signal of purifying selection was also evident in MEGs when branch
lengths were compared to those of endosperm genes. We note that two MEGs, including
AGAMOUS-like protein implicated in one coevolutionary network, exhibited branch lengths
suggesting a particularly strong negative selection. Our observations were not in line with the
positive selection detected on PEGs on the autogamous species *A. thaliana* and *C. rubella*
(Hatorangan et al., 2016; Tuteja et al., 2019), raising the question of how the mating system
impacts the selection of imprinted genes in these species.

Nevertheless, caution is warranted in attributing the observed purifying directional selection
solely to imprinting. In fact, the detected purifying selection could be explained by alternative
mechanisms (Patten et al., 2014), including (1) an increase of the efficiency of selection by
hemizygous expression (Haig and Westoby 1989), (2) selection against Transposable
Element (TE; Padilla-García et al., 2025) or (3) local adaptation (Kremer et al., 2025).
However, none of the current models rigorously predict the effects of these mechanisms on
polymorphism and divergence, making it difficult to discriminate among alternative
explanations.

Implications for reproductive isolation and adaptation

The evolutionary consequences of purifying selection on alleles at imprinted loci remain
poorly understood. Directional selection on the coding regions of these genes could (1)
influence adaptive potential by shaping the spectrum of selectable alleles within populations
(Cvijović et al., 2018), (2) contribute to speciation processes by generating genetic
incompatibilities between lineages (Seehausen et al., 2014), (3) facilitate the transition to
apomixy in plants (Hojsgaard and Schartl 2021) and (4) promote the establishment of
reproductive isolation (Maheshwari and Barbash 2011; Cvijović et al., 2018; İltaş et al.,
2021), more particularly through endosperm-based post-zygotic barriers (Haig 2000;
Spencer 2000; Kinoshita 2007; Köhler et al., 2010).

Previous studies empirically demonstrated that some imprinted genes are involved in these
barriers, including in the autogamous species *A. thaliana* (Wolff et al., 2015), between the
allogamous *Solanum peruvianum* et *S. chilense* (Florez-Rueda et al., 2016), between the
allogamous *Capsella grandiflora* and the autogamous *C. rubella* and *Capsella orientalis*
(Lafon-Placette et al., 2018), between the allogamous *Mimulus guttatus*, *M. luteus* (Kinser et
al., 2021), *M. caespitosa* and *M. tilingi* (Sandstedt and Sweigart 2022), and between the
autogamous wheat species *Triticum durum* and *T. aestivum* (Yang et al., 2023).

Imprinted genes may also act as central hubs within complex regulatory networks that
contribute to hybrid vigour and reproductive barriers (Botet and Keurentjes 2020). However,
despite these observations, the broader evolutionary implications of these effects at the
network level remain largely unresolved, particularly regarding how they interact with
demographic history, variation in the mating system, and other selective pressures.

Imprinted genes are associated with singular signals of selection in North America

PEGs and MEGs were associated with higher π_{NS}/π_S values compared to endosperm and
control genes, which could indicate balancing or relaxation of selection on coding regions
(Galtier and Rousselle 2020). However, this signal was not observed for Tajima's D. This
apparent relaxation may not reflect parental conflict but instead could result from the recent
colonization of North America by *A. lyrata* from Europe (Ross-Ibarra et al., 2008; Otto et al.,
2015; Willi et al., 2018) that could have reduced effective population size (Galtier and
Rousselle 2020) and the efficiency of purifying selection (Laenen et al., 2018), especially on
TE (Padilla-García et al., 2025). Testing this hypothesis would require future comparisons of
selection signals in endosperm and imprinted genes across allogamous populations with
varying levels of polymorphism. Nevertheless, we note that two MEGs exhibited reduced
Tajima's D or π_{NS}/π_S, consistent with ongoing directional selection.

Imprinted genes are not sensitive to shifts in the mating system in A. lyrata

Conflict over resource allocation is expected to be reduced in autogamous lineages relative
to allogamous ones (Haig 2000; Brandvain and Haig 2005). However, most studies of
genetic and phylogenetic signatures of selection associated with imprinting have focused on
autogamous species, leaving the effect of shifts in the mating system on polymorphism of
imprinted genes unresolved. In the autogamous species *A. thaliana*, the imprinted *MEDEA*
locus shows increased polymorphism but not in its allogamous sister species *A. lyrata*
(Spillane et al., 2007; Miyake et al., 2009). However, PEGs were associated with positive
selection in the autogamous species *A. thaliana* and *C. rubella* (Hatorangan et al., 2016;
Tuteja et al., 2019). In our study, the selection signals in imprinted genes, contrary to
endosperm genes, showed limited sensitivity to variation of the mating system. The absence
of variation in Tajima's D could be explained if directional selection had already fixed the
optimal alleles prior to the transition from allogamy to autogamy, which could also explain
the signal of positive selection detected in PEGs in *A. thaliana* and *C. rubella* (Hatorangan et
al., 2016; Tuteja et al., 2019). However, this explanation does not account for the high
π_{NS}/π_S observed in allogamous and autogamous populations. Theoretical studies are
necessary to estimate the combined effect of selection related to the kinship theory,
imprinting, demographic history and the mating system.

Imprinted genes present signals of ancient and recent coevolution

Another potential, but not mandatory, outcome proposed in extensions of the kinship theory
is the coevolution of some imprinted loci with antagonistic effects on offspring growth (Moore
and Haig, 1991; Wilkins and Haig 2001; Mills and Moore, 2004). To investigate this, we
applied both population genomic and phylogenetic approaches (Dutheil and Galtier 2007).
We found that most PEGs and MEGs were connected within coevolutionary networks, with a
significantly higher number of direct associations than expected by chance. These networks
encompass essential functions for endosperm development.

For instance, one network based on branch length correlations was enriched in enzymatic
activities related to cell wall formation and modification, a key process during endosperm
cellularization in Arabidopsis (Hehenberger et al., 2012), which in turn influences seed size
(Pignatta et al., 2018; Zhang et al., 2020), a likely target of parental conflict. These
correlated evolutionary rates could reflect equivalent selective constraints (Galtier and
Rousselle 2020) or just a common chromosomal context (*e.g.* TE) that modifies the mutation
rate (Lynch 2010). Interestingly, this network appeared to be strongly dependent on a MEG,
an AGAMOUS-like protein, characterised also by reduced branch lengths, consistent with

expectations that coevolution may drive coordinated evolution among genes primarily
through rare compensatory mutations (Fraser et al., 2002)

The network based on polymorphism involved genes related to syntaxin-binding proteins,
crucial for membrane activity, molecular transport, and protein storage, whereas the
topology-based network was enriched in DNA-binding transcription factors, including those
regulating seed development, tissue differentiation, and storage metabolism (Noguero et al.,
2013).

These phylogenetic and population genetic signals of coevolution could promote the
maintenance of the gene functions (Clark et al., 2012), limit their evolutionary independency
(Yang and Bielawski 2000) and be another source of reproductive barrier (Presgraves 2010).
Interestingly, the coevolution signal did not involve increased site-by-site coevolution,
suggesting that protein–protein molecular interactions, often drivers of coevolution (Green et
al., 2021), played a limited role here. Instead, the observed patterns may reflect selection
acting on gene dosage rather than protein products (Clark et al., 2012).

However, as correlations between evolutionary statistics can arise from variation of the
mutation rate, recombination environment or shared functional constraints, we interpret them
as exploratory signals only. The more stringent CoMap analysis (Dutheil & Galtier 2007) did
not reveal enrichment for co-evolution at the site level, indicating that networks based on
correlation should be treated cautiously and do not by themselves confirm co-evolution.

Main limitations in our study

Although our study combines multiple phylogenetic and population genomic statistics, the
signals detected were not consistently congruent across approaches. This inconsistency
may reflect evolutionary processes not accounted for, (such as demographic history or
chromosomal context) as well as several methodological limitations, outlined below.

First of all, estimating Tajima's D from pooled sequencing data is challenging because
statistics based on the Site Frequency Spectrum (SFS) are sensitive to coverage variation
and sampling noise, which increase estimation variance (Futschik & Schlötterer 2010;
Ferretti et al. 2013; Lynch et al. 2014). Although we minimized biases specific to the dataset
by comparing genes processed under identical conditions, technical noise cannot be fully
excluded. In addition, short genes, such as many MEGs or control genes, yield fewer
segregating sites, reducing the reliability of Tajima's D (Tajima 1989).

A second limitation concerns the imprinting dataset. The set of imprinted genes analysed
was derived from a study on a geographically and genetically distinct allogamous
subspecies European *A. lyrata petraea* (Klosinska et al., 2016), assuming conservation of
imprinting, because no within-lineage data on imprinting in *A. lyrata* exist. However,
imprinting shows limited conservation within *A. thaliana* (Gehring et al., 2011; Wolff et al.,
2011; Hsieh et al., 2011; Pignatta et al., 2014), between *A. thaliana* and *A. lyrata* in our data,
and generally across species (Eaux et al., 2013; Hatorangan et al., 2016; Roth et al., 2018;
Flores-Vergara et al., 2020). It is therefore possible that several genes classified as
imprinted in our analyses are not imprinted in the studied populations, especially given
known population-level variation in imprinting drive by TE (Padilla-García et al., 2025).
Future population-specific characterisations of imprinting will be critical to resolve this issue.

Moreover, some imprinted genes identified by Klosinska et al. (2016) may have been
inadvertently excluded from our dataset due to our method for detecting tissue-specific
expression, which relied on RNA sequencing (RNA-seq) data pooled from both allogamous
and autogamous populations. More controlled and population-specific experimental designs
will be required to recover the complete set of endosperm-expressed imprinted genes
relevant to parental conflict.

A last important limitation is about the panels of genes compared (control, endosperm and
imprinting genes). To be consistent with previous studies (Spillane et al., 2007; Miyake et al.,
2009; Hatorangan et al., 2016; Tuteja et al., 2019), we also selected genes from panels
based on expression patterns. However, the three gene panels differ substantially in gene
size, CDS proportion, and GC content, factors known to influence both polymorphism and
phylogenetic metrics, but seldom considered in previous studies (Spillane et al., 2007;
Miyake et al., 2009; Hatorangan et al., 2016; Tuteja et al., 2019).

Endosperm genes and PEGs are substantially longer, whereas MEGs and control genes are
shorter. The size of the gene is an important parameter that impact many statistics. For
example, larger genes tend to evolve more slowly due to increased functional constraint and
pleiotropy (Massey 2013; Lopes et al., 2021), display higher conservation and reduced
turnover (Abdel-Haleem 2007; Koonin 2015), and can inflate the number of species with
detectable homologous genes. In this sense, we found a positive correlation between the
number of species with homologous genes and the gene size. This higher gene size may
partially explain the higher homology counts observed for PEGs and endosperm genes
relative to control genes. Conversely, we found that the gene size is negatively correlated
with polymorphism and Tajima's D. However, this tendency is in contradiction with longer
genes as endosperm genes presenting a higher Tajima's D or shorter genes as the MEGs

being associated with lower polymorphism, suggesting that, maybe, the effects of parental
conflict on polymorphism in these genes are underestimated, more specifically for MEG
because shorter genes may yield less stable estimates of Tajima's D (Tajima 1989). We
found also a negative correlation with the expression level, that could explain partially why
the expression of MEG is more efficient, even if the gene group stays the most redundant
parameter predicted by AIC, but could not explain why the expression of PEG is higher than
endosperm genes.

Several studies show that long, constrained genes tend to exhibit shorter branches due to
stronger purifying selection (Kimura 1983; Abdel-Haleem 2007; Massey 2013; Koonin 2015).
However, in our study, we found a positive correlation between the length of the branches
and the gene size, maybe explained by the amount of endosperm genes with long branches.
This correlation suggests that the longer branches observed in imprinted and endosperm
genes are not explained by the size of the genes.

Moreover, both imprinted and endosperm genes show elevated CDS content relative to
controls. Thus, this parameter could not explain the increase in efficiency observed in
expression of imprinted genes, despite the negative correlation detected. In the same way,
this parameter could not explain the reduced π_{NS}/π_S observed in imprinted genes compared
to endosperm genes, despite the positive correlation detected. Interestingly, high CDS
proportion is associated with stronger purifying selection (Hughes et al., 2003), which can
reduced the length of the branches, as observed in our study. However, the increase of
branch length detected in endosperm and imprinted genes is the reverse pattern predicted,
suggesting that the effect on branch length in these genes is maybe underestimated.

Finally, loci subject to imprinting show reduced GC content relative to endosperm and
control genes. GC-poor genes tend to have reduced recombination rates, lower
polymorphism (Wright et al., 2003), and higher AT-biased substitution rates, which can
artificially lengthen branches (Duret & Galtier 2009), and explain the longer branches
observed in PEG and endosperm genes. However, contrary to these expectations, we found
a negative correlation between the length of the branches and the GC content, which may
be explained by the important amount of endosperm genes. Interestingly, we also found a
negative correlation between the polymorphism and the GC content in our study, which
suggests that the reduced polymorphism observed in MEGs is maybe underestimated. Low
GC also reduces the likelihood of finding conserved homologous genes across species
(Urrutia & Hurst 2003), as found in our study, but is not able to explain why the PEGs and
the endosperm genes are associated with higher number of species with homologous genes
despite lower GC content than in control genes. Low GC is expected to also inflate D_{NS}/D_S

by increasing the probability of non-synonymous changes (Kryazhimskiy & Plotkin 2008).
However, we detected the reverse correlation in our study, which could explain why
imprinted genes are associated with a reduced D_{NS}/D_S.

The lower GC of imprinted genes is also unexpected given their elevated expression (Sharp
& Li 1987; Bernardi 2000), suggesting that expression differences may be underestimated.
For these reasons, although the three panels differ in genomic features, endosperm-
upregulated genes remain the only biologically appropriate baseline for testing hypotheses
related to imprinting. A feature-matched control panel would reduce variation but would not
provide a more relevant comparison for detecting signatures associated with parental
conflict.

Together, differences in gene size, CDS proportion, and GC content may partly account for
the heterogeneity of evolutionary signals observed across statistics. Some signatures
consistent with parental conflict may therefore be underestimated or confounded, while
others may reflect intrinsic genomic properties rather than selective pressures related to
imprinting. Future work will require gene panels matched not only for expression, but also for
fine-scale genomic features.

Although several genomic features differ among gene panels, we performed targeted
robustness checks to ensure that competing non-selective scenarios did not spuriously
generate the patterns we report. In particular, the SNP-density threshold retained
comparable fractions of MEGs, PEGs, endosperm genes and controls, ruling out retention
bias as a driver of polymorphism contrasts. Moreover, all evolutionary inferences were
tested against empirical null distributions from endosperm-upregulated genes, which share
expression patterns and chromatin context with imprinted genes, limiting confounding effects
of gene architecture or local mutation rate. These checks do not rule out all alternative
explanations, but they demonstrate that the signals observed are unlikely to arise solely from
genomic artefacts or filtering procedures.

Conclusions

Overall, our analyses revealed that endosperm-expressed and imprinted genes in *A. lyrata*
exhibit evolutionary signatures partly consistent with theoretical expectations derived from
models of parental conflict and imprinting, including evidence for ancient balancing selection
in endosperm genes, expression-level regulation in imprinted genes, ancient purifying
selection, and coevolutionary interactions. However, signals at the population level were less
consistent with theoretical expectations, likely reflecting demographic history, mating system

shifts, and the poor conservation of imprinting across populations. Future work should

combine empirical surveys of imprinting with theoretical models explicitly incorporating

demography and mating system variation to refine predictions about the evolutionary

dynamics of loci subject to genomic imprinting.

Materials and Methods

We hypothesized, based on the kinship theory, that imprinted genes should display distinct genomic signatures of selection compared with non-imprinted endosperm genes (Table 1). To test this prediction, we designed an approach combining phylogenetics and population genetics. Specifically, we focused on a set of 27 maternally expressed genes (MEGs) and 46 paternally expressed genes (PEGs) previously characterised in *A. lyrata* (Klosinska et al., 2016) and unambiguously annotated in the reference genome V1.0.23 (Hu et al., 2011; https://doi.org/10.5281/zenodo.17287897). These imprinted genes were systematically compared against two control datasets: (i) 2,703 non-imprinted genes upregulated in the endosperm, and (ii) a reference panel of 100 randomly selected ubiquitous genes, expressed across multiple tissues. This design allowed us to disentangle signatures of selection specifically associated with imprinting from those affecting the whole genome or endosperm genes in general.

Expression data

RNA-seq expression data from multiple tissues, including leaf, root, pistil, and pollen, were obtained from two *A. lyrata* populations representing 27,820 genes from both European and North American lineages, as described in İltaş et al. (2024). To maximize gene representation and increase the statistical power, the data from both mating type was pooled. To complement these datasets, we also incorporated endosperm RNA-Seq data derived from five allogamous *A. lyrata* individuals, previously generated by Klosinska et al. (2016). The endosperm data were reprocessed and integrated into our analytical pipeline following the instructions and R script available in GitHub (https://github.com/thelovelab/DESeq2). To identify endosperm-specific genes, we retained only those that showed significant upregulation (log2FC > 1, adjusted P < 0.05) in all pairwise comparisons against other tissues. The RNA-Seq datasets for leaf, root, pistil, and pollen are available in the NCBI BioProject database under accession PRJNA1067428, while the endosperm RNA-Seq dataset is accessible under accession GSE76076.

Genes clusters

We used the list of the 49 PEGs and 35 MEGs found in *A. lyrata* (Klosinska et al., 2016). We kept the genes having conserved coordinates in the reference genome of *A. lyrata* (Hu et al., 2011), resulting in a final list of 27 MEGs and 46 PEGs.

To disentangle signatures of selection specifically associated with the endosperm (where conflicts over resource allocation are expected) from those potentially arising through other

selective processes, we separated imprinted genes upregulated in the endosperm (14 PEGs

and 15 MEGs) from imprinted genes not showing endosperm-specific upregulation (32

PEGs and 12 MEGs), as determined by differential expression analysis.

To evaluate whether the observed proportions of MEGs and PEGs upregulated in the

endosperm deviated from theoretical expectations under the kinship theory, we compared

them with null distributions generated by 10,000 random sampling of 49 and 35 genes (the

size of the PEGs and MEGs groups) among the 27,820 expressed genes.

For all subsequent analyses, we focused exclusively on endosperm-upregulated imprinted

genes and compared them with two reference groups: the set of 2,703 non-imprinted

endosperm-upregulated genes (hereafter, "endosperm genes"), the sets of 15 MEG and 14

PEG upregulated in endosperm (hereafter, "MEG" and "PEG") and a set of 100 randomly

selected ubiquitously expressed genes (hereafter, "control genes"). To remain consistent

with previous studies on imprinting and parental conflict (Spillane et al., 2007; Miyake et al.,

2009; Hatorangan et al., 2016; Tuteja et al., 2019), the control panel was defined based

solely on expression patterns. This set is intended to represent a genome-wide baseline that

is unrelated to endosperm-specific functions or parental conflict, rather than a genomic

feature–matched control. Consequently, no constraints were imposed on gene size, GC

content, CDS proportion, or chromosomal position. However, for each statistic studied

below, we compared the full model, which included gene group, gene length, CDS

proportion, and GC content as explanatory variables, using Akaike's Information Criterion

(AIC). We then evaluated a series of reduced models, removing one parameter at a time,

and selected the model with the lowest AIC. This procedure was iterated until the

best-supported model was identified using the function StepAIC of the MASS library (version

7.3-65) in R. This approach allowed us to determine (1) whether the gene group remained a

significant predictor in the best model and (2) whether genomic features accounted for the

observed trends in the analysed statistics.

To test whether expression levels conformed to predictions of higher optimal expression in

PEGs than in MEGs, we normalized expression per gene copy number and compared

observed values with bootstrap distributions from non-imprinted endosperm genes (10,000

resamples of 29 genes). To account for gene dosage, expression values were normalized by

the expected number of expressed copies: three for non-imprinted endosperm genes, two

for MEGs, and one for PEGs.

Based on the sequences from the reference genome of *A. lyrata* (Hu et al., 2011), we

conducted bootstrap analyses to compare the gene size, the GC and the CDS content (%) of

PEGs and MEGs with values obtained from 10,000 resamples of 29 endosperm genes and
control genes independently. Similarly, we also conducted bootstrap analyses to compare
the values of control genes with those obtained from 10,000 resamples of 100 endosperm
genes.

Estimation of codon and amino acid bias

Based on the sequences from the reference genome of *A. lyrata* (Hu et al., 2011), we
estimated the proportion of each amino acid within the gene and the proportion of each
possible codon for each amino acid. Then, we conducted bootstrap analyses to compare
these proportions in PEGs and MEGs with values obtained from 10,000 resamples of 29
endosperm genes. Thus, we defined the "overused" (resp. "underused") codons as codons
with a proportion higher (resp. lower) than in the 95^{th} percentiles of the 1,000 resample of 29
endosperm genes.

To test if the overused or underused codons were related to gene expression, we compared
the variations of Predicted High Expression (PHE) score induced by codons with the score of
unbiased codons by permutations test with 1,000 replications for MEGs and PEGs
independently. The PHE score was estimated based on overexpressed genes in *A. thaliana*
and reflect the importance of a codon on genes expression (Sahoo et al,. 2019). We
hypothesized that codons overused in imprinted genes were associated with increases in
PHE score because they were associated with increases in gene expression, whereas the
underused codons were associated with reduced PHE scores.

Distribution of the genes across Brassicaceae

We assessed whether the imprinted genes presented specific signals of selection compared
to the endosperm and control genes in the phylogeny of 22 Brassicaceae. All the complete
sequence of reference genomes were extracted from the Phytozome database (Goodstein
et al., 2012; Table S4). We extracted the sequences of the genes in the *A. lyrata* reference
genome (Hu et al., 2011) using bedtools V2.30 (Quinlan et al., 2010). Then, we retained the
best hit for these sequences against 22 other Brassicaceae using BLAST V2.10 (Altschul et
al., 1990) with an e-value < 1e-5. If more than one hit was found for a gene in one species,
we filtered the best hit as the hit with the highest length alignment, the highest % of identity,
then the highest coverage on the gene in *A. lyrata* and the highest reciprocal coverage on
the gene in the other species. Then we extracted the sequence of the best hit for each
species of each gene.

To test for significant differences in the number of species with homologous genes across
the genes group, we conducted bootstrap analyses to compare the value on PEGs and
MEGs with values obtained from 10,000 resamples of 29 endosperm genes and control
genes independently. Similarly, we conducted bootstrap analyses to compare the control
genes values with those obtained from 10,000 resamples of 100 endosperm genes.

Divergence of the imprinted genes across Brassicaceae

The rates of nonsynonymous mutations (D_{NS}) and synonymous mutations (D_S) are assumed
to follow the nearly-neutral evolutionary process and the ratio D_{NS}/D_S is therefore
approximate to the selective pressure on the protein product of a gene. A decrease in
D_{NS}/D_S in genes indicates a increase of intensity of negative selection. To test whether
imprinted genes display accelerated evolutionary rates compared to endosperm genes,
using the annotation of the 0-fold and 4-fold degenerate sites in *A. lyrata*, we estimated for
each gene the mean evolutionary rates D_{NS}/D_S between the reference genome of *A. lyrata*
and all the species with an homologous gene found. Only the genes found in at least 3
species and with a D_S higher than 0 were considered.

To test for significant differences in the D_{NS}/D_S across the group of genes, we conducted
bootstrap analyses to compare the value on PEGs and MEGs with values obtained from
10,000 resamples of 29 endosperm genes and control genes independently. Similarly, we
conducted bootstrap analyses to compare the values on control genes with values obtained
from 10,000 resamples of 100 endosperm genes.

Topology of the genes across Brassicaceae

We aligned each gene with all its best hits found across the different species with Muscle
V3.8.155 (Edgar 2004). Then, we built a phylogenetic tree for each gene using the HKY85
model with PhyML V3.3 (Guindon et al., 2010). In parallel, we concatenated the alignment of
the 1,055 endosperm genes found in the 22 others Brassicaceae to extract the topology of
the phylogenetic tree of the endosperm genes build using PhyML V3.3 (Guindon et al.,
2010). This tree was used as a proxy of the topology of the endosperm. Then, we estimated
the Robinson-Foulds (RF) distance between the topology of this tree and the topology of the
tree of each gene obtained previously using the R package TreeDist. Then, to test for
significant differences in RF distances across the group of genes, we conducted bootstrap
analyses to compare the values obtained from PEGs and MEGs with those from 10,000
resamples of 29 endosperm genes and control genes independently. Similarly, we

conducted bootstrap analyses to compare the values on control genes with values obtained
from 10,000 resamples of 100 endosperm genes.

Variation of the branch length across Brassicaceae

We investigated whether the imprinted genes presented variation in the branch lengths
compared to the endosperm and control genes. For each gene, we re-estimate gene specific
branch lengths using the topology of the phylogenetic tree obtained with the 1,055
endosperm genes using PhyML V3.3 (Guindon et al., 2010). Then, we extracted the terminal
branch lengths at the tips for each gene with the R package 'ape' version 5.7-1. If the gene
was not present in a species, the value was replaced by "NA". To test for significant
differences in the branch length across the group of genes, we conducted bootstrap
analyses to compare the mean in PEGs and MEGs with means obtained from 10,000
resamples of 29 endosperm genes and control genes independently. Similarly, we
conducted bootstrap analyses to compare the means of control genes with values obtained
from 10,000 resamples of 100 endosperm genes.

Read mapping and variant calling

We used natural accessions of *A. lyrata* from North America from nine allogamous and nine
autogamous populations of 25 individuals each (Table S6), using available pool-sequencing
data (Willi et al., 2018).

The raw reads from Willi et al (2018) datasets were mapped onto the complete *A. lyrata*
reference genome V1.0.23 (Hu et al., 2011) using Bowtie2 v2.4.1 (Langmead and Salzberg
2012). The resulting alignments were then converted to BAM files using samtools v1.3.1 (Li
et al., 2009) and duplicated reads were removed with the MarkDuplicates program of picard-
tools v1.119 (http://broadinstitute.github.io/picard). Biallelic SNPs in these regions were
called using the Genome Analysis Toolkit v. 3.8 (GATK, DePristo et al., 2011) and stored in
a GVCF file to keep non-variable called positions. A quality score threshold of 60 was
applied using vcftools v0.1.15 (Danecek et al., 2011), to retain only highly reliable SNPs.
This stringent cutoff is commonly used in high-coverage datasets to minimise false positives,
which is particularly important for downstream analyses of selection and co-evolution that
are sensitive to SNP-calling errors (Schlötterer et al., 2014; GATK Best Practices). We
excluded sites with less than 15 reads aligned. Sites covered by at least 15 reads but
containing no SNP were considered monomorphic. The final number of sites in each sample
set is summarised in Table S6.

Effect of imprinting on global population polymorphism

We decomposed the signal of selection across the imprinted genes into two elementary
statistics: Tajima's D and the ratio of average pairwise heterozygosity between the non-
synonymous and synonymous sites π_{NS}/π_S. For the allogamous and autogamous
populations, we compared the values obtained for the imprinted genes with those obtained
for the endosperm and control.

For each gene of each population, we estimated π, π_{NS}/π_S for all positions and the Tajima's
D based on allelic frequencies in naturals populations. Briefly, we estimate the Tajima's D by
gene following the classical equation described in Tajima (1989):
$D = (\pi - (S / \Sigma(1/i))) / \sqrt{e1 S + e2 S (S - 1)}$, with e1 = (n+1) /3 * (n+1) - (1/ Σ(1/i))/ Σ(1/i) and
e2= (2 * (n * n + n + 3)) / (9 * n * (n-1)) - ((n+2) /(Σ(1/i) *n-1)) + (Σ(1/i²) / (Σ(1/i)²)) / (
Σ(1/i)² + Σ(1/i²)). S is for the number of segregating sites, i for value between 1 and n-1
haplotypes and n is for the number of haplotypes (number of individuals multiplied by two for
diploid).

If a position in the *A. lyrata* genome was covered but not polymorphic, π was set to 0. Then,
the 0☐fold and 4☐fold degenerated sites were identified and extracted from the reference
genome and the genome annotation using the script NewAnnotateRef.py (Williamson et al.,
2014; updated version available in
https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee). Because all mutations
at 0-fold degenerated sites alter the sequence of the encoded protein, they are non-
synonymous (NS). In contrast, mutations at the 4-fold degenerate sites never alter the
encoded amino acid, and all are synonymous (S). For the sake of simplicity, we discarded
mutations at 2- or 3-fold degenerate sites, which can be either synonymous or non-
synonymous. Based on π estimated for each site, we estimated the mean π_{NS} and π_S, for
each gene in each population, in order to estimate π_{NS}/π_S. We removed the genes with a π_S
equal to 0. For the Tajima's D, we applied a SNP-density threshold (≥0.5% polymorphic
sites) to ensure reliable estimates. This density-based criterion avoids the gene-length
biases inherent in minimum-SNP thresholds, since genes vary widely in size across
categories. We verified that this filtering did not disproportionately exclude any gene
category: the proportions of retained genes were comparable across endosperm and
imprinting genes and across imprinting and control genes (Table S7).

To test for significant differences across the genes group, we conducted bootstrap analyses
to compare the mean π, π_{NS}/π_S and Tajima's D of PEGs and MEGs with means obtained
from 10,000 resamples of 29 endosperm genes and control genes independently. Similarly,
we conducted bootstrap analyses to compare the control genes values with those obtained
from 10,000 resamples of 100 endosperm genes. We performed the test on allogamous and

autogamous populations independently. Finally, for each statistic and group independently,
we tested for the mating system effect by a performing Kruskal-Wallis test as implemented in
R version 4.3.2.

Genes in coevolution

To detect coevolving coding sequences, we used three different methods based on the
following assumptions: coevolution led BY 1) related polymorphic patterns between genes in
populations, 2) related evolutionary rate across species, or 3) similarity in the topologies
across species (Dutheil and Galtier 2007).

Correlations between evolutionary rates or diversity metrics were used as exploratory
indicators of evolutionary covariance, following approaches such as evolutionary rate
covariation, which has been widely applied to detect co-functionality or shared selective
pressures (Clark et al., 2012). Because such correlations may reflect genomic or mutational
confounders, we used endosperm-upregulated genes as an empirical baseline for defining
extreme thresholds.

To detect coevolution based on polymorphism in population and on evolutionary rates
across species, we estimated Pearson's linear correlations between each pair of imprinted
and endosperm genes for the mean π and the branch length on each species. Based on
these correlations matrices, we extracted correlations between pairs of endosperm genes to
define thresholds for extreme values in the distributions. Then, all pairs of genes that
included imprinted genes with correlations below the 1% or above the 99% quantiles of the
distribution were considered significant. We used these significant correlations of π and
branch length to build the coevolution networks.

To detect coevolution based on tree-topology similarity, we first used the distribution of RF
distances between endosperm genes and the global endosperm topology to estimate the
value obtained in 95[th] percentile value. Then, we excluded all the imprinted and endosperm
genes having a lower RF distance than this value to remove genes whose topology similarity
was explained by congruence with the global topology. Finally, we compared the topology of
each remaining imprinted gene and endosperm gene. All pairs of genes with a null RF
distance were considered as significantly similar and used to build the last coevolution
network.

For each network of coevolution, we compared the number of pairs of imprinted genes
directly correlated with the number found between genes in 1,000 random samples of 29
endosperm genes. Moreover, we tested for gene enrichment on each GO term in each

network based on the occurrences of each term across 1,000 random samples of equivalent
size of all the endosperm and imprinted genes. The complete list of GO term associated with
each gene was downloaded from Gene Ontology online platform for *A. thaliana* (Ashburner
et al,. 2000).

Identification of coevolving sites.

We used a model-based approach to detect coevolving sites for gene pair that showed a
coevolution signal. The pairs of imprinted genes studied were all directly associated in at
least one previously mentioned network. To use this method, we concatenated each
sequence of pairs of genes for each species. If one species was not found in the other gene,
the sequence associated with this species was removed.

Coevolving sites within each pair of genes were predicted using the clustering method
implemented in the CoMap package (version 1.5.2; https://jydu.github.io/comap/; Dutheil and
Galtier 2007). The built-in algorithm was then used to correct for multiple testing and to
estimate the false discovery rate (FDR). In this study, 1,000 simulations per tested pair were
performed, and the groups of sites with an adjusted p value $< 1\%$ were considered
coevolving. Simple correlation was used as a measure of coevolution. For each comparison
pair, we extracted the coevolving groups and classified them by size (between 2 and 10).

To test for significant excess of correlations, we compared the results obtained for the pair of
imprinted genes found in networks previously mentioned with the results obtained for
equivalent number of pair comparisons of imprinted genes not associated in any network, for
each cluster of coevolving groups of substitutions of equivalent size by Kruskal-Wallis tests.
If the number of correlations is significantly higher than the number expected between two
random genes by chance or by alignment errors, we expect a significant increase in these
numbers in pairs of associated imprinted genes.

Data availability

The RNAseq data for leaf, root, pistil, and pollen tissues can be found in the NCBI BioProject
database under submission number PRJNA1067428. The endosperm RNAseq data is
available under submission number GSE76076.

The version of *A. lyrata* reference genome (Hu et al., 2011) used in this paper is available in
zenodo (https://doi.org/10.5281/zenodo.17287897).

We used several GitHub repositories, each corresponding to a distinct analytical module.
This structure reflects the fact that the datasets and workflows differ substantially across

analyses and that several scripts were developed independently and used in other publications. For clarity, all repositories are indexed in a Zenodo record (https://zenodo.org/records/17954040), and each includes documentation, test data, and commented scripts. This multi-repository organization was reviewed and approved for reproducibility and FAIR compliance by the PCI Ecology Data Editor (Checklist PCI Ecology 867).

The R code for the bootstrap tests is available in GitHub (https://github.com/leveveaudrey/bootstrap).

The R code used for permutation test was available in GitHub (https://github.com/leveveaudrey/permutation-test-)

The python code to estimate the proportion of each amino acid within the gene and the proportion of each possible codon for each amino acid is available in GitHub (https://github.com/leveveaudrey/codon-bias).

The python code for the pipeline of phylogenetic analysis (distribution, divergence extraction of topologies and branch length of the genes across Brassicaceae) is available in GitHub (https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee).

The python code for the pipeline including alignment of reads and the SNP calling is available in Github (https://github.com/leveveaudrey/alignment-and-snp-calling).

The python code for the pipeline estimating the π, the π_{NS}/π_S for all positions and the Tajima's D of each gene in PoolSeq dataset is available in Github (https://github.com/leveveaudrey/POOLSEQ-analysis-of-polymorphism).

Acknowledgements

This work has been supported by the Czech Science Foundation (GACR 22-20240S granted to C.L-P). Computational resources were supplied by the project 'e-Infrastruktura CZ' (e-INFRA CZ LM2018140) supported by the Ministry of Education, Youth and Sports of the Czech Republic. A.L thanks personally Stella Huynh for our beneficial suggestions and commentaries.

Author contributions

O.I developed and performed the transcriptional analysis. J-Y.D and A.L developed and performed the coevolution analysis. A.L analysed and interpreted the data. A.L wrote the manuscript. All authors edited the manuscript.

Declaration of interests

The authors declare no competing interests.

Bibliography

1) Abdel-Haleem, H., 2007. The Origins of Genome Architecture. J Hered 98, 633–634. https://doi.org/10.1093/jhered/esm073

2) Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. Journal of Molecular Biology 215, 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

3) Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G., 2000. Gene Ontology: tool for the unification of biology. Nat Genet 25, 25–29. https://doi.org/10.1038/75556

4) Bemer, M., Heijmans, K., Airoldi, C., Davies, B., Angenent, G.C., 2010. An Atlas of Type I MADS Box Gene Expression during Female Gametophyte and Seed Development in Arabidopsis. Plant Physiology 154, 287–300. https://doi.org/10.1104/pp.110.160770

5) Bernardi, G., 2000. Isochores and the evolutionary genomics of vertebrates. Gene 241, 3–17. https://doi.org/10.1016/s0378-1119(99)00485-0

6) Botet, R., and Joost J. B. Keurentjes., 2020. The Role of Transcriptional Regulation in Hybrid Vigor. Frontiers in Plant Science 11 (April). https://doi.org/10.3389/fpls.2020.00410

7) Brandvain, Y., Haig, D., 2005. Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. Am Nat 166, 330–338. https://doi.org/10.1086/432036

8) Bryant, N., Lloyd, J., Sweeney, C., Myouga, F., Meinke, D., 2010. Identification of Nuclear Genes Encoding Chloroplast-Localized Proteins Required for Embryo Development in Arabidopsis. Plant Physiology 155, 1678. https://doi.org/10.1104/pp.110.168120

9) Clark, N.L., Alani, E., Aquadro C.F., 2012. Evolutionary Rate Covariation Reveals Shared Functionality and Coexpression of Genes. Genome Research 22 (4): 714–20. https://doi.org/10.1101/gr.132647.111

10) Cvijović, I., Good, B.H., Desai, M.M., 2018. The Effect of Strong Purifying Selection on Genetic Diversity. Genetics 209 (4): 1235–78. https://doi.org/10.1534/genetics.118.301058

11) Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., 1000 Genomes Project Analysis Group, 2011. The variant call format and VCFtools. Bioinformatics 27, 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

12) Day, T., Bonduriansky, R., 2004. Intralocus sexual conflict can drive the evolution of genomic imprinting. Genetics 167, 1537. https://doi.org/10.1534/genetics.103.026211

13) Day, R.C., Herridge, R.P., Ambrose, B.A., Macknight, R.C., 2008. Transcriptome Analysis of Proliferating Arabidopsis Endosperm Reveals Biological Implications for the Control of Syncytial Division, Cytokinin Signaling, and Gene Expression Regulation. Plant Physiology 148, 1964–1984. https://doi.org/10.1104/pp.108.128108

14) Dekkers, B.J.W., Pearce, S., van Bolderen-Veldkamp, R.P., Marshall, A., Widera, P., Gilbert, J., Drost, H.-G., Bassel, G.W., Müller, K., King, J.R., Wood, A.T.A., Grosse, I., Quint, M., Krasnogor, N., Leubner-Metzger, G., Holdsworth, M.J., Bentsink, L., 2013. Transcriptional Dynamics of Two Seed Compartments with Opposing Roles in Arabidopsis Seed Germination. Plant Physiology 163, 205–215. https://doi.org/10.1104/pp.113.223511

15) DePristo, M.A., Banks, E., Poplin, R.E., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., del Angel, G., Rivas, M.A., Hanna, M., McKenna, A., Fennell, T.J., Kernytsky, A.M., Sivachenko, A.Y., Cibulskis, K., Gabriel, S.B., Altshuler, D., Daly, M.J., 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet 43, 491–498. https://doi.org/10.1038/ng.806

16) Dutheil, J. ,Galtier, N., 2007. Coevolution within and between Genes. https://doi.org/10.1159/000107599

17) Eaux, A.J., Bilinski, P., Eichten, S.R., Vaughn, M.W., Ross-Ibarra, J., Gehring, M., Reprinte, N.M., 2013. Comprehensive analysis of imprinted genes in maize reveals allelic variation for imprinting and limited conservation with other species. Proceedings of the National Academy of Sciences 110, 19639–19644. https://doi.org/10.1073/pnas.1309182110

18) Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

19) Edwards, S.V., 2009. Natural selection and phylogenetic analysis. Proceedings of the National Academy of Sciences 106, 8799–8800. https://doi.org/10.1073/pnas.0904103106

20) Ferretti, L., Ramos-Onsins, S.E., Pérez-Enciso, M., 2013. Population genomics from pool sequencing. Mol Ecol 22, 5561–5576. https://doi.org/10.1111/mec.12522

21) Flintham, E., Savolainen, V., Otto, S.P., Reuter, M., Mullon, C., 2025. The maintenance of genetic polymorphism underlying sexually antagonistic traits. Evol Lett 9, 259–272. https://doi.org/10.1093/evlett/qrae059

22) Florez-Rueda, A.M., Paris, M., Schmidt, A., Widmer, A., Grossniklaus, U., Städler, T., 2016. Genomic Imprinting in the Endosperm Is Systematically Perturbed in Abortive Hybrid Tomato Seeds. Mol Biol Evol 33, 2935–2946. https://doi.org/10.1093/molbev/msw175

23) Flores-Vergara, M.A., Oneal, E., Costa, M., Villarino, G., Roberts, C., De Luis Balaguer, M.A., Coimbra, S., Willis, J., Franks, R.G., 2020. Developmental Analysis of Mimulus Seed Transcriptomes Reveals Functional Gene Expression Clusters and Four Imprinted, Endosperm-Expressed Genes. Front. Plant Sci. 11. https://doi.org/10.3389/fpls.2020.00132

24) Foxe, J.P., Stift, M., Tedder, A., Haudry, A., Wright, S.I., Mable, B.K., 2010. Reconstructing Origins of Loss of Self-Incompatibility and Selfing in North American Arabidopsis Lyrata: A Population Genetic Context. Evolution 64, 3495–3510. https://doi.org/10.1111/j.1558-5646.2010.01094.x

25) Fraser, H.B., Hirsh, A.E., Steinmetz, L.M., Scharfe, C., Feldman, M.W., 2002. Evolutionary Rate in the Protein Interaction Network. Science 296, 750–752. https://doi.org/10.1126/science.1068696

26) Futschik, A., Schlötterer, C., 2010. The Next Generation of Molecular Markers From Massively Parallel Sequencing of Pooled DNA Samples. Genetics 186, 207–218. https://doi.org/10.1534/genetics.110.114397

27) Galtier, N., Rousselle, M., 2020. How Much Does Ne Vary Among Species? Genetics 216, 559–572. https://doi.org/10.1534/genetics.120.303622

28) Garro, M., Greco, E., Vannay, G.J., Leonova, A., Bruno, L., Capella, M., 2025. Non-CG DNA methylation represses SDC expression to modulate hypocotyl elongation during thermormorphogenesis in Arabidopsis. J Exp Bot 76, 2517–2534. https://doi.org/10.1093/jxb/eraf105

29) Gaudinier, A., Rodriguez-Medina, J., Zhang, L., Olson, A., Liseron-Monfils, C., Bågman, A.-M., Foret, J., Abbitt, S., Tang, M., Li, B., Runcie, D.E., Kliebenstein, D.J., Shen, B., Frank, M.J., Ware, D., Brady, S.M., 2018. Transcriptional regulation of nitrogen-associated metabolism and growth. Nature 563, 259–264. https://doi.org/10.1038/s41586-018-0656-3

30) Gehring, M., Missirian, V., Henikoff, S., 2011. Genomic Analysis of Parent-of-Origin Allelic Expression in Arabidopsis thaliana Seeds. PLOS ONE 6, e23687. https://doi.org/10.1371/journal.pone.0023687

31) Goodstein, D.M., Shu, S., Howson, R., Neupane, R., Hayes, R.D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., Putnam, N., Rokhsar, D.S., 2012. Phytozome: a comparative platform for green plant genomics. Nucleic Acids Res 40, D1178-1186. https://doi.org/10.1093/nar/gkr944

32) Green, A.G., Elhabashy, H., Brock, K.P., Maddamsetti, R., Kohlbacher, O., Marks, D.S., 2021. Large-scale discovery of protein interactions at residue resolution using co-evolution calculated from genomic sequences. Nat Commun 12, 1396. https://doi.org/10.1038/s41467-021-21636-z

33) Grishkevich, V., Yanai, I., 2014. Gene length and expression level shape genomic novelties. Genome Res 24, 1497–1503. https://doi.org/10.1101/gr.169722.113

34) Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59, 307–321. https://doi.org/10.1093/sysbio/syq010

35) Haig, D., Westoby, M., 1989. Parent-Specific Gene Expression and the Triploid Endosperm. The American Naturalist 134, 147–155. https://doi.org/10.1086/284971

36) Haig, D., Westoby, M., 1991. Genomic Imprinting in Endosperm: Its Effect on Seed Development in Crosses between Species, and between Different Ploidies of the Same Species, and Its Implications for the Evolution of Apomixis. Philosophical Transactions: Biological Sciences 333, 1–13. https://doi.org/10.1098/rstb.1991.0057

37) Haig, D., 1997. Parental antagonism, relatedness asymmetries, and genomic imprinting. Proceedings of the Royal Society of London. Series B: Biological Sciences 264, 1657–1662. https://doi.org/10.1098/rspb.1997.0230

38) Haig, D., 2000. The Kinship Theory of Genomic Imprinting. Annual Review of Ecology and Systematics 31, 9–32. https://doi.org/10.1146/annurev.ecolsys.31.1.9

39) Haig, D., 2014. Coadaptation and conflict, misconception and muddle, in the evolution of genomic imprinting. Heredity 113, 96–103. https://doi.org/10.1038/hdy.2013.97

40) Haig, D., Graham, C., 1991. Genomic imprinting and the strange case of the insulin-like growth factor II receptor. Cell 64, 1045–1046. https://doi.org/10.1016/0092-8674(91)90256-X

41) Hatorangan, M.R., Laenen, B., Steige, K.A., Slotte, T., Köhler, C., 2016. Rapid Evolution of Genomic Imprinting in Two Species of the Brassicaceae. The Plant Cell 28, 1815–1827. https://doi.org/10.1105/tpc.16.00304

42) Hehenberger, E., Kradolfer, D., Köhler, C., 2012. Endosperm cellularization defines an important developmental transition for embryo development. Development 139, 2031–2039. https://doi.org/10.1242/dev.077057

43) Hojsgaard, D., Schartl, M., 2021. Skipping sex: A nonrecombinant genomic assemblage of complementary reproductive modules. BioEssays 43, 2000111. https://doi.org/10.1002/bies.202000111

44) Hough, J., Immler, S., Barrett, S.C.H., Otto, S.P., 2013. Evolutionarily stable sex ratios and mutation load. Evolution 67, 1915–1925. https://doi.org/10.1111/evo.12066

45) Hsieh, T.-F., Shin, J., Uzawa, R., Silva, P., Cohen, S., Bauer, M.J., Hashimoto, M., Kirkbride, R.C., Harada, J.J., Zilberman, D., Fischer, R.L., 2011. Regulation of imprinted gene expression in Arabidopsis endosperm. Proceedings of the National Academy of Sciences 108, 1755–1762. https://doi.org/10.1073/pnas.1019273108

46) Hu, T.T., Pattyn, P., Bakker, E.G., Cao, J., Cheng, J.-F., Clark, R.M., Fahlgren, N., Fawcett, J.A., Grimwood, J., Gundlach, H., Haberer, G., Hollister, J.D., Ossowski, S., Ottilar, R.P., Salamov, A.A., Schneeberger, K., Spannagl, M., Wang, X., Yang, L., Nasrallah, M.E., Bergelson, J., Carrington, J.C., Gaut, B.S., Schmutz, J., Mayer, K.F.X., Van de Peer, Y., Grigoriev, I.V., Nordborg, M., Weigel, D., Guo, Y.-L., 2011. The Arabidopsis lyrata genome sequence and the basis of rapid genome size change. Nat Genet 43, 476–481. https://doi.org/10.1038/ng.807

47) İltaş, Ö., Svitok, M., Cornille, A., Schmickl, R., Lafon Placette, C., 2021. Early evolution of reproductive isolation: A case of weak inbreeder/strong outbreeder leads to an intraspecific hybridization barrier in Arabidopsis lyrata. Evol 75, 1466–1476. https://doi.org/10.1111/evo.14240

48) İltaş, Ö., Čertner, M., Placette, C.L., 2024. Natural variation in sexual traits and gene expression between selfing and outcrossing Arabidopsis lyrata suggests sexual selection at work. Plant and Cell Physiology pcae090. https://doi.org/10.1093/pcp/pcae090

49) Jiang, H., Guo, D., Ye, J., Gao, Y., Liu, H., Wang, Y., Xue, M., Yan, Q., Chen, J., Duan, L., Li, G., Li, X., Xie, L., 2021. Genome-wide analysis of genomic imprinting in the endosperm and allelic variation in flax. The Plant Journal 107, 1697–1710. https://doi.org/10.1111/tpj.15411

50) Kasimatis, K.R., Nelson, T.C., Phillips, P.C., 2017. Genomic Signatures of Sexual Conflict. Journal of Heredity 108, 780–790. https://doi.org/10.1093/jhered/esx080

51) Kinoshita, T., 2007. Reproductive barrier and genomic imprinting in the endosperm of flowering plants. Genes & Genetic Systems 82, 177–186. https://doi.org/10.1266/ggs.82.177

52) Kinser, T.J., Smith, R.D., Lawrence, A.H., Cooley, A.M., Vallejo-Marín, M., Conradi Smith, G.D., Puzey, J.R., 2021. Endosperm-based incompatibilities in hybrid monkeyflowers. Plant Cell 33, 2235–2257. https://doi.org/10.1093/plcell/koab117

53) Köhler, C., Scheid, O.M., Erilova, A., 2010. The impact of the triploid block on the origin and evolution of polyploid plants. Trends in Genetics 26, 142–148. https://doi.org/10.1016/j.tig.2009.12.006

54) Koonin, E.V., 2015. Why the Central Dogma: on the nature of the great biological exclusion principle. Biology Direct 10, 52. https://doi.org/10.1186/s13062-015-0084-3

55) Klosinska, M., Picard, C.L., Gehring, M., 2016. Conserved imprinting associated with unique epigenetic signatures in the Arabidopsis genus. Nature Plants 2, 1–8. https://doi.org/10.1038/nplants.2016.145

56) Kremer, A., Chen, J., Lascoux, M., 2025. 'Chimes of resilience': what makes forest trees genetically resilient? New Phytologist 246, 1934–1951. https://doi.org/10.1111/nph.70108

57) Laenen, B., Tedder, A., Nowak, M.D., Toräng, P., Wunder, J., Wötzel, S., Steige, K.A., Kourmpetis, Y., Odong, T., Drouzas, A.D., Bink, M.C.A.M., Ågren, J., Coupland, G., Slotte, T., 2018. Demography and mating system shape the genome-wide impact of purifying selection in Arabis alpina. Proc Natl Acad Sci U S A 115, 816–821. https://doi.org/10.1073/pnas.1707492115

58) Lafon-Placette, C., Hatorangan, M.R., Steige, K.A., Cornille, A., Lascoux, M., Slotte, T., Köhler, C., 2018. Paternally expressed imprinted genes associate with hybridization barriers in Capsella. Nature Plants 4, 352–357. https://doi.org/10.1038/s41477-018-0161-6

59) Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923

60) Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

61) Llaurens, V., Whibley, A., Joron, M., 2017. Genetic architecture and balancing selection: the life and death of differentiated variants. Molecular Ecology 26, 2430–2448. https://doi.org/10.1111/mec.14051

62) Lopes, I., Altab, G., Raina, P., de Magalhães, J.P., 2021. Gene Size Matters: An Analysis of Gene Length in the Human Genome. Front Genet 12, 559998. https://doi.org/10.3389/fgene.2021.559998

63) Lynch, M., 2010. Evolution of the mutation rate. Trends Genet 26, 345–352. https://doi.org/10.1016/j.tig.2010.05.003

64) Lynch, M., Bost, D., Wilson, S., Maruki, T., Harrison, S., 2014. Population-Genetic Inference from Pooled-Sequencing Data. Genome Biol Evol 6, 1210–1218. https://doi.org/10.1093/gbe/evu085

65) Maheshwari, S., Barbash, D.A., 2011. The Genetics of Hybrid Incompatibilities. Annu. Rev. Genet. 45, 331–355. https://doi.org/10.1146/annurev-genet-110410-132514

66) Massey, S.E., 2013. Proteome size as the major factor determining mutation rates. Proceedings of the National Academy of Sciences 110, E858–E859. https://doi.org/10.1073/pnas.1219306110

67) McVean, G.T., Hurst, L.D., 1997. Molecular evolution of imprinted genes: no evidence for antagonistic coevolution. Proceedings of the Royal Society of London. Series B: Biological Sciences 264, 739–746. https://doi.org/10.1098/rspb.1997.0105

68) Mills, W., Moore, T., 2004. Polyandry, Life-History Trade-Offs and the Evolution of Imprinting at Mendelian Loci. Genetics 168, 2317–2327. https://doi.org/10.1534/genetics.104.030098

69) Miyake, T., Takebayashi, N., Wolf, D.E., 2009. Possible Diversifying Selection in the Imprinted Gene, MEDEA, in Arabidopsis. Molecular Biology and Evolution 26, 843–857. https://doi.org/10.1093/molbev/msp001

70) Moore, T., Haig, D., 1991. Genomic imprinting in mammalian development: a parental tug-of-war. Trends Genet 7, 45–49. https://doi.org/10.1016/0168-9525(91)90230-N

71) Noguero, M., Atif, R.M., Ochatt, S., Thompson, R.D., 2013. The role of the DNA-binding One Zinc Finger (DOF) transcription factor family in plants. Plant Science 209, 32–45. https://doi.org/10.1016/j.plantsci.2013.03.016

72) Onosato, H., Fujimoto, G., Higami, T., Sakamoto, T., Yamada, A., Suzuki, T., Ozawa, R., Matsunaga, S., Seki, M., Ueda, M., Sako, K., Galis, I., Arimura, G., 2022. Sustained defense response via volatile signaling and its epigenetic transcriptional regulation. Plant Physiol 189, 922–933. https://doi.org/10.1093/plphys/kiac077

73) Otto, S.P., Scott, M.F., Immler, S., 2015. Evolution of haploid selection in predominantly diploid organisms. Proceedings of the National Academy of Sciences 112, 15952–15957. https://doi.org/10.1073/pnas.1512004112

74) Padilla-García, N., Le Veve, A., Čermák, V., İltaş, Ö., Contreras-Garrido, A., Legrand, S., Aury, J.-M., Horvath, R., Lafon Placette, C., 2025. The Demographic History of Populations and Genomic Imprinting have Shaped the Transposon Patterns in Arabidopsis lyrata. Mol Biol Evol 42, msaf093. https://doi.org/10.1093/molbev/msaf093

75) Papdi, C., Ábrahám, E., Joseph, M.P., Popescu, C., Koncz, C., Szabados, L., 2008. Functional Identification of Arabidopsis Stress Regulatory Genes Using the Controlled cDNA Overexpression System. Plant Physiol 147, 528–542. https://doi.org/10.1104/pp.108.116897

76) Patten, M.M., Ross, L., Curley, J.P., Queller, D.C., Bonduriansky, R., Wolf, J.B., 2014. The evolution of genomic imprinting: theories, predictions and empirical tests. Heredity 113, 119–128. https://doi.org/10.1038/hdy.2014.29

77) Pignatta, D., Erdmann, R.M., Scheer, E., Picard, C.L., Bell, G.W., Gehring, M., 2014. Natural epigenetic polymorphisms lead to intraspecific variation in Arabidopsis gene imprinting. eLife 3, e03198. https://doi.org/10.7554/eLife.03198

78) Pignatta, D., Novitzky, K., Satyaki, P.R.V., Gehring, M., 2018. A variably imprinted epiallele impacts seed development. PLOS Genetics 14, e1007469. https://doi.org/10.1371/journal.pgen.1007469

79) Presgraves, D.C., 2010. The molecular evolutionary basis of species formation. Nat Rev Genet 11, 175–180. https://doi.org/10.1038/nrg2718

80) Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics 26, 841–842. https://doi.org/10.1093/bioinformatics/btq033

81) Raissig, M.T., Baroux, C., Grossniklaus, U., 2011. Regulation and Flexibility of Genomic Imprinting during Seed Development. The Plant Cell 23, 16–26. https://doi.org/10.1105/tpc.110.081018

82) Ross-Ibarra, J., Wright, S.I., Foxe, J.P., Kawabe, A., DeRose-Wilson, L., Gos, G., Charlesworth, D., Gaut, B.S., 2008. Patterns of Polymorphism and Demographic History in Natural Populations of Arabidopsis lyrata. PLOS ONE 3, e2411. https://doi.org/10.1371/journal.pone.0002411

83) Roth, M., Florez-Rueda, A.M., Paris, M., Städler, T., 2018. Wild tomato endosperm transcriptomes reveal common roles of genomic imprinting in both nuclear and cellular endosperm. The Plant Journal 95, 1084–1101. https://doi.org/10.1111/tpj.14012

84) Sahoo, S., Das, S.S., Rakshit, R., 2019. Codon usage pattern and predicted gene expression in Arabidopsis thaliana. Gene, Articles initially published in Gene: X 1-4, 2019 721, 100012. https://doi.org/10.1016/j.gene.2019.100012

85) Sandstedt, G.D., Sweigart, A.L., 2022. Developmental evidence for parental conflict in driving Mimulus species barriers. New Phytol 236, 1545–1557. https://doi.org/10.1111/nph.18438

86) Scott, R.J., Spielman, M., Bailey, J., Dickinson, H.G., 1998. Parent-of-origin effects on seed development in Arabidopsis thaliana. Development 125, 3329–3341. https://doi.org/10.1242/dev.125.17.3329

87) Seehausen, O., Butlin, R.K., Keller, I., Wagner, C.E., Boughman, J.W., Hohenlohe, P.A., Peichel, C.L., Saetre, G.-P., Bank, C., Brännström, Å., Brelsford, A., Clarkson, C.S., Eroukhmanoff, F., Feder, J.L., Fischer, M.C., Foote, A.D., Franchini, P., Jiggins, C.D., Jones, F.C., Lindholm, A.K., Lucek, K., Maan, M.E., Marques, D.A., Martin, S.H., Matthews, B., Meier, J.I., Möst, M., Nachman, M.W., Nonaka, E., Rennison, D.J., Schwarzer, J., Watson, E.T., Westram, A.M., Widmer, A., 2014. Genomics and the origin of species. Nat Rev Genet 15, 176–192. https://doi.org/10.1038/nrg3644

88) Sharp, P.M., Li, W.H., 1987. The codon Adaptation Index--a measure of directional synonymous codon usage bias, and its potential applications. Nucleic Acids Res 15, 1281–1295. https://doi.org/10.1093/nar/15.3.1281

89) Soliman, H.K., Coughlan, J.M., 2024. United by conflict: Convergent signatures of parental conflict in angiosperms and placental mammals. J Hered 115, 625–642. https://doi.org/10.1093/jhered/esae009

90) Spencer, H. G. 2000. Population Genetics and Evolution of Genomic Imprinting. Annual Review of Genetics 34: 457–77. https://doi.org/10.1146/annurev.genet.34.1.457.

91) Spencer, H.G., Clark, A.G., 2014. Non-conflict theories for the evolution of genomic imprinting. Heredity 113, 112–118. https://doi.org/10.1038/hdy.2013.129

92) Spillane, C., Schmid, K.J., Laoueillé-Duprat, S., Pien, S., Escobar-Restrepo, J.-M., Baroux, C., Gagliardini, V., Page, D.R., Wolfe, K.H., Grossniklaus, U., 2007. Positive darwinian selection at the imprinted MEDEA locus in plants. Nature 448, 349–352. https://doi.org/10.1038/nature05984

93) Sujeeth, N., Mehterov, N., Gupta, S., Qureshi, M.K., Fischer, A., Proost, S., Omidbakhshfard, M.A., Obata, T., Benina, M., Staykov, N., Balazadeh, S., Walther, D., Fernie, A.R., Mueller-Roeber, B., Hille, J., Gechev, T.S., 2020. A novel seed plants gene regulates oxidative stress tolerance in Arabidopsis thaliana. Cell. Mol. Life Sci. 77, 705–718. https://doi.org/10.1007/s00018-019-03202-5

94) Sylvestre, F., Mérot, C., Normandeau, E., Bernatchez, L., 2023. Searching for intralocus sexual conflicts in the three-spined stickleback (Gasterosteus aculeatus) genome. Evolution 77, 1667–1681. https://doi.org/10.1093/evolut/qpad075

95) Tuteja, R., McKeown, P.C., Ryan, P., Morgan, C.C., Donoghue, M.T.A., Downing, T., O'Connell, M.J., Spillane, C., 2019. Paternally Expressed Imprinted Genes under Positive Darwinian Selection in Arabidopsis thaliana. Molecular Biology and Evolution 36, 1239–1253. https://doi.org/10.1093/molbev/msz063

96) Willi, Y., 2013. The Battle of the Sexes over Seed Size: Support for Both Kinship Genomic Imprinting and Interlocus Contest Evolution. The American Naturalist 181, 787–798. https://doi.org/10.1086/670196

97) Willi, Y., Fracassetti, M., Zoller, S., Van Buskirk, J., 2018. Accumulation of Mutational Load at the Edges of a Species Range. Molecular Biology and Evolution 35, 781–791. https://doi.org/10.1093/molbev/msy003

98) Wilkins, J.F., Haig, D., 2001. Genomic imprinting of two antagonistic loci. Proceedings of the Royal Society of London. Series B: Biological Sciences 268, 1861–1867. https://doi.org/10.1098/rspb.2001.1651

99) Wilkins, J.F., Haig, D., 2003. What Good Is Genomic Imprinting: The Function of Parent-Specific Gene Expression. Nature Reviews Genetics 4 (5): 359–68. https://doi.org/10.1038/nrg1062.

100) Wilkins, J.F., 2011. Genomic imprinting and conflict□induced decanalization. Evolution 65, 537–553. https://doi.org/10.1111/j.1558-5646.2010.01147.x

101) Williamson, R.J., Josephs, E.B., Platts, A.E., Hazzouri, K.M., Haudry, A., Blanchette, M., Wright, S.I., 2014. Evidence for Widespread Positive and Negative Selection in Coding and Conserved Noncoding Regions of Capsella grandiflora. PLOS Genetics 10, e1004622. https://doi.org/10.1371/journal.pgen.1004622

102) Wolff, P., Weinhofer, I., Seguin, J., Roszak, P., Beisel, C., Donoghue, M.T.A., Spillane, C., Nordborg, M., Rehmsmeier, M., Köhler, C., 2011. High-Resolution Analysis of Parent-of-Origin Allelic Expression in the Arabidopsis Endosperm. PLOS Genetics 7, e1002126. https://doi.org/10.1371/journal.pgen.1002126

103) Wolff, P., Jiang, H., Wang, G., Santos-González, J., Köhler, C., 2015. Paternally expressed imprinted genes establish postzygotic hybridization barriers in Arabidopsis thaliana. eLife 4, e10074. https://doi.org/10.7554/eLife.10074

104) Wuest, S.E., Vijverberg, K., Schmidt, A., Weiss, M., Gheyselinck, J., Lohr, M., Wellmer, F., Rahnenführer, J., Mering, C. von, Grossniklaus, U., 2010. Arabidopsis Female Gametophyte Gene Expression Map Reveals Similarities between Plant and Animal Gametes. Current Biology 20, 506–512. https://doi.org/10.1016/j.cub.2010.01.051

105) Yang, G., Feng, M., Yu, K., Cui, G., Zhou, Y., Sun, L., Gao, L., Zhang, Y., Peng, H., Yao, Y., Hu, Z., Rossi, V., De Smet, I., Ni, Z., Sun, Q., Xin, M., 2023. Paternally imprinted LATE-FLOWERING2 transcription factor contributes to paternal-excess interploidy hybridization barriers in wheat. Journal of Integrative Plant Biology 65, 2587–2603. https://doi.org/10.1111/jipb.13574

106) Zhang, S., Wang, D., Zhang, H., Skaggs, M.I., Lloyd, A., Ran, D., An, L., Schumaker, K.S., Drews, G.N., Yadegari, R., 2018. FERTILIZATION-INDEPENDENT SEED-Polycomb Repressive Complex 2 Plays a Dual Role in Regulating Type I MADS-Box Genes in Early Endosperm Development. Plant Physiology 177, 285–299. https://doi.org/10.1104/pp.17.00534

107) Zhang, B., Li, C., Li, Y., Yu, H., 2020. Mobile TERMINAL FLOWER1 determines seed size in Arabidopsis. Nat Plants 6, 1146–1157. https://doi.org/10.1038/s41477-020-0749-5

108) Zou, H.-F., Zhang, Y.-Q., Wei, W., Chen, H.-W., Song, Q.-X., Liu, Y.-F., Zhao, M.-Y., Wang, F., Zhang, B.-C., Lin, Q., Zhang, W.-K., Ma, B., Zhou, Y.-H., Zhang, J.-S., Chen, S.-Y., 2012. The transcription factor AtDOF4.2 regulates shoot branching and seed coat formation in Arabidopsis. Biochemical Journal 449, 373–388. https://doi.org/10.1042/BJ20110060

Figure S1: phylogenetic tree obtained by HKY85 model based on the 1055 endosperm genes found in all species after 1000 bootstrap. The percentage of trees in which the associated haplotypes clustered together is shown next to the branches. The tree is drawn to scale, with branch lengths measured in mean number of substitutions per site.