Limited directional selection but coevolutionary signals among imprinted genes in *A. lyrata.*

Audrey Le Veve[1,2], Ömer İltaş[2,3], Julien Y. Dutheil[4], Clément Lafon Placette[2, *]

[1] IHPE, Univ Perpignan Via Domitia, CNRS, IFREMER, Univ Montpellier, Perpignan, France

[2] Department of Botany, Faculty of Science, Charles University, Prague, Czech Republic

[3] Max Planck Institute of Molecular Plant Physiology: Potsdam, Brandenburg, DE

[4] RG Molecular Systems Evolution, Department of Theoretical Biology, Max Planck Institute for Evolutionary Biology, August-Thienemann-Straße 2, 24306 Plön, Germany

[*] Corresponding author: audrey.le-veve@univ.perp.fr

Abstract

Genomic imprinting is a form of gene regulation leading to the unequal expression of maternal and paternal alleles. The main hypothesis invoked to explain the evolution of imprinted genes is the kinship theory, which posits a conflict between parental genomes over resource allocation in progeny. According to this theory, imprinted genes are expected to undergo a shift from balancing to directional selection, in coding or regulatory regions. However, most studies addressing selection in imprinted genes have focused on self-fertilizing species, where conflicts over resource allocation are predicted to be weak. Consequently, the impact of kinship theory on the evolution of imprinted genes remains largely untested. Furthermore, the predicted coevolution between certain imprinted genes under kinship theory has not been tested yet empirically.

Combined phylogenetic and population genomic approaches, we investigated signatures of selection on imprinted genes across the Brassicaceae family and in autogamous and allogamous populations of *Arabidopsis lyrata*, and searched for evidence of coevolution among imprinted genes. We found that endosperm-expressed genes exhibited signals of balancing selection across Brassicaceae and within allogamous populations, consistent with unresolved intralocus conflicts. Supporting the kinship theory, these signals varied with mating system. Moreover, phylogenetic analyses further indicated negative directional selection acting on imprinted genes. However, population-level signals were independent of mating system and showed limited concordance with kinship predictions, possibly due to stronger selection acting on expression than on coding sequences. Finally, we identified coevolution

31   between imprinted genes, although not at specific sites, suggesting that interactions beyond

32   protein sequence drive this process.

34   **Key words:** Genomic imprinting, Kinship theory, Endosperm, Coevolution, Mating system.

Introduction

The kinship theory, or the parental conflict theory, was first proposed by Haig and Westoby (1989) in the context of endosperm development and elaborated in a review in 1991. It posits that in species where the maternal parent nourishes progeny sired by multiple males, maternal fitness is maximized by distributing resources equally among offspring, whereas paternal fitness is enhanced by securing more resources for their own offspring. This conflict is expected to drive intralocus conflict between growth-promoting and growth-limiting genes (Haig and Graham 1991; Moore and Haig 1991; Wilkins and Haig 2001), consequently leaving signatures of balancing selection, such as reduced divergence across species and increased polymorphism within populations (Day and Bonduriansky 2004; Kasimatis et al., 2017). This has been shown in *Arabidopsis thaliana,* whose crosses showed dosage effects on seed size, although these may reflect interactions between cytoplasmic inheritance, dosage imbalance, and imprinting (Scott et al., 1998). Notably, balancing selection in genes expressed in endosperm (embryo's nourishing tissue in the seed) has not been systematically studied to validate the kinship theory.

The kinship theory remains the leading explanation for the evolution of imprinted genes (Haig 1997; Spencer and Clark 2014; Patten et al., 2014). Imprinting is an epigenetic regulation in flowering plants and mammals in which allele expression depends on the parental origin (Haig 1997; Wilkins 2011). By creating hemizygous expression of parental alleles, the kinship theory argues that imprinting evolves to resolve parental conflicts over resource allocation. Consequently, imprinted genes are expected to be disproportionately expressed in the endosperm in plants, the tissue where parental conflict operates (Haig 1997; Haig 2000). Thus, the proportion of maternally expressed genes (MEGs) and paternally expressed genes (PEGs) upregulated in the endosperm should exceed the genome-wide expectation derived from sampling all expressed genes. According to this hypothesis, many imprinted genes appear to be involved in resource accumulation in the endosperm and metabolic processes (Raissig et al., 2011). These theoretical expectations and previous empirical observations justify testing whether MEGs and PEGs are disproportionately upregulated in endosperm relative to random expectation.

Theoretical models using models of selection on one or two genes involved in resource allocation predicted that: (1) imprinting set distinct optimal expression levels for paternally expressed genes (PEGs) and maternally expressed genes (MEGs; Haig 1997); Exactly, PEGs are predicted to evolve higher optimal expression levels than MEGs, because paternal alleles benefit from increased resource acquisition, whereas maternal alleles tend to limit growth (Haig 2000; Patten et al., 2014). Accordingly, we expect an higher expression per active gene

copy in PEGs than in MEGs and finally in non-imprinted endosperm genes; (2) PEGs and
MEGs are under directional selection to increase and limit resource acquisition by offspring
respectively (Haig 2000); and (3) imprinting at some growth-promoting locus may drive
antagonistic coevolution with some growth-suppressing function (Wilkins and Haig 2001).
However, these models focus on *IGF2* and *IGF2R* genes, which regulate foetal growth in
mammals, and without investigating the expected signatures of selection for endosperm
genes.

Empirically, phylogenetic signals of directional selection have been reported for *MEDEA* locus
in the allogamous species *Arabidopsis lyrata*, but not in the autogamous sister species *A.*
thaliana (Spillane et al., 2007; Miyake et al., 2009), and for PEGs at the population levels in
the autogamous Brassicaceae species *Capsella rubella* and *A. thaliana* (Hatorangan et al.,
2016; Tuteja et al., 2019). However, MEGs did not show comparable signals, possibly due to
technical challenges in detecting selection if selection primarily targets gene expression levels.
Indeed, a central debate has emphasized that kinship theory may concern selection on the
optimal expression levels of genes between maternal and paternal genomes, and not on
alleles (Patten et al., 2014). Selection acting on gene expression may still leave detectable
traces in amino acid or codon usage patterns (Patten et al., 2014).

The incongruent signals detected in imprinted genes may also be due to polymorphism
dynamics shaped by the mating system. The kinship theory assumes that progenies are sired
by multiple males and that imprinted genes are preferentially expressed in the endosperm
(Haig and Westoby 1991; Haig 2000; Brandvain and Haig 2005). The transition to autogamy
may alter selection acting on genes, potentially leading to more intense directional selection
or relaxed selection (Soliman and Coughlan 2024). However, the empirical studies cited above
did not assess endosperm-specific expression of the imprinted genes and only focused on
autogamous species. Consequently, the evolutionary signatures of selection on imprinted
genes and the contribution of the kinship theory to their evolution remain unresolved in plants.

Kinship theory also predicts a potential coevolution between antagonistic imprinted genes
(Wilkins and Haig 2001; Willi 2013), yet no phylogenetic signal was detected for *IGF2* and
IGF2R (McVean and Hurst 1997), and coevolution among endosperm-expressed genes
remains unknown. Population genomic methods can detect coevolving sequences based on
allele frequency variation; however, they may be biased by demographic or population
structure effects. More integrative approaches based on phylogenetic comparisons can
identify co-occurring changes in candidate coevolving genes (Dutheil and Galtier 2007).

Here, we test these predictions by combining expression, phylogenetic and population-
genomic analyses to evaluate whether imprinted genes in *A. lyrata* show evolutionary

signatures consistent with the kinship theory. We compared imprinted genes in allogamous *A.*
lyrata (Klosinska et al., 2016) with non-imprinted endosperm genes, using phylogenetic data
from 23 Brassicaceae species and genomic data from nine allogamous populations and nine
closely related autogamous populations of *A. lyrata* from North America (Ross-Ibarra et al.,
2008). We tested for : (1) variations in expression patterns and/or codon usage; (2) reduced
divergence and polymorphism, as expected under directional selection, and (3) correlations
between evolutionary rates or diversity metrics used as exploratory indicators of evolutionary
covariance expected in case of coevolution. Our results indicate that endosperm-expressed
genes display patterns consistent with intralocus conflict and that transition to autogamy
promoted directional selection on these genes. Imprinted genes show evidence of ancient
negative directional selection, codon usage bias, and substantial coevolution. However,
population-level signals are less consistent with kinship theory, possibly due to stronger
selection acting on expression than on coding sequences or by a recent relaxed selection on
these genes in North American populations.

Results

General analytical framework

Parental conflict is expected to occur primarily in the endosperm. Using transcriptomic data from 19,425 genes across five tissues (leaf, root, pollen, pistil, and endosperm), we identified 2,732 specifically upregulated genes in the endosperm of *A. lyrata* among 31,073 genes analysed. Among these upregulated genes, 15 were MEGs and 14 were PEGs from the 27 MEGs and 46 PEGs previously identified in *A. lyrata* (Klosinska et al., 2016). Between *A. lyrata* and *A. thaliana*, 40% of MEGs (6/15) and 50% of PEGs (7/14) identified in our dataset were conserved as imprinted (Table S1).

Throughout the Results section, we compare four categories of genes: maternally expressed genes (MEGs) and paternally expressed genes (PEGs) that are upregulated in the endosperm (referred "imprinted genes" if considered together), the 2,703 non-imprinted genes specifically upregulated in the endosperm ("endosperm genes"), and a set of 100 control genes. The control panel was randomly sampled from the set of 126 ubiquitously expressed genes and serves as a baseline for genome-wide patterns of selection unrelated to parental conflict. The control panel was chosen following the expression-based sampling strategy used in previous imprinting studies, rather than by matching genomic characteristics across groups. Differences in gene size, CDS proportion, and GC content among panels were therefore quantified and considered when interpreting downstream evolutionary analyses.

For each statistic, genes with insufficient informative sites were excluded. Then, significance was evaluated using 10,000 bootstrap resampling matching the size of focal group. framework because this method is unsensitive to difference of samples sizes. Rather than repeating the method in each subsection, we applied a single unified framework in which observed means are compared to the 5th–95th percentiles of bootstrap distributions. This approach was used consistently across analyses of expression, codon usage, phylogenetic metrics, and population polymorphism. Although the figures share a similar visual layout for consistency, each statistic captures a distinct evolutionary dimension allowing direct comparison of patterns across gene categories while retaining analytical complementarity.

We first compared gene size, CDS content (%) and GC content (%) across groups, as these features may influence polymorphism patterns and generate bioinformatic artefacts. Control genes and MEGs were significantly smaller than PEGs and endosperm genes, and CDS content was significantly higher in endosperm and imprinted genes relative to control genes. GC content was also lower in imprinted genes than in control genes (Table S2). These

differences were quantified and considered when interpreting all evolutionary analyses. Nevertheless, differences in gene size, CDS proportion and GC content were not corrected for directly, and we retained the control panel to be congruent with methods used in previous studies in the autogamous species *C. rubella* and *A. thaliana* (Hatorangan et al., 2016; Tuteja et al., 2019). In practice, all evolutionary comparisons evaluating predictions of the kinship theory were performed relative to endosperm-upregulated non-imprinted genes, which represent the expression-matched and biologically relevant background for imprinting analyses. The control panel was used only to describe genome-wide patterns of variation.

To evaluate whether these genomic differences could create artificial signals of selection, we performed several robustness checks. First, for population-genomic analyses, we verified that the SNP-density filter did not preferentially exclude short, GC-poor or low-diversity genes. Second, by comparing all statistics against empirical null distributions derived from endosperm-upregulated genes, we ensured that each test was evaluated against a biologically appropriate baseline sharing similar expression patterns, thereby reducing the influence of residual genomic confounders unrelated to imprinting or parental conflict. Finally, we compared models using the Akaike Information Criterion (AIC). The full model included gene group, gene length, CDS proportion and GC content as explanatory variables. We then evaluated a series of reduced models in which one parameter was removed at a time and retained the model with the lowest AIC value. This procedure was iterated until the best-supported model was identified. This approach allowed to determine (1) whether gene group remained a significant predictor in the best model and (2) whether genomic features accounted for the observed trends in the analysed statistics.

In previous study (Tuteja et al., 2019), validation of kinship theory to explain imprinting was performed following another enrichment-based logic, which evaluated not the global behaviour of imprinted genes but the proportion of them exhibiting extreme signatures compatible with parental conflict. Thus, we tested whether directional selection occurred more frequently in imprinted genes than in the empirical distribution of endosperm genes. Candidates for directional selection were defined as the 5% most extreme values for branch length, Tajima's D, or π_{NS}/π_S in allogamous populations. Under the null hypothesis of no enrichment, imprinted genes should fall into this class with probability 0.05. We therefore tested whether the observed proportion exceeded 5% using independent binomial tests.

Coevolution was assessed using three complementary measures (Pearson's correlation coefficients for π, branch lengths, and RF distances, for each pair of PEGs, MEGs, and endosperm genes), and only correlations exceeding thresholds defined from pair of endosperm genes (1st or 99th percentiles) were retained. Finally, we tested if the number of

187 correlations found between PEGs and MEGs and the number of imprinted genes involved
188 were significantly higher than expected from 1,000 random samplings of 29 endosperm genes.

Imprinted genes are preferentially expressed in the endosperm

Under kinship theory, both MEGs and PEGs are expected to be disproportionately
represented among endosperm-upregulated genes, because imprinting evolves specifically in
tissues exposed to parental conflict. We therefore first tested whether the proportion of
upregulated MEGs and PEGs exceeded expectations derived from random sampling of the
transcriptome. The proportions of MEGs (0.56) and PEGs (0.30) upregulated in the
endosperm were both significantly higher than expected from resampling of all genes (p values
< 1e-04), and MEGs were more frequently upregulated than PEGs (based on resampling of
imprinted genes; p value =0.03).

Because PEGs are predicted to increase offspring resource acquisition, while MEGs tend to
restrict it, theory predicts higher optimal expression levels for PEGs, followed by MEGs,
compared with non-imprinted endosperm genes. We therefore tested whether expression per
expressed allele differed among groups after correcting for gene dosage. After adjusting
expression for gene copy number (three expressed copies for non-imprinted endosperm
genes, two for MEGs, and one for PEGs), both PEGs and MEGs displayed significantly higher
expression levels relative to endosperm genes, with 3.18- and 1.71-fold increases,
respectively (bootstrap tests; p-values < 1e-03; Fig 1).

Using AIC, we found that the best model to explain this expression pattern (AIC = -4633) was
this that include the group of genes, but also the gene size (coefficient = -5.4e-05), the CDS
proportion (coefficient = -2.34e-03) and the GC proportion (coefficient = 2.66e-02), whereas
the worst model (AIC = -2454) was this that exclude only the group of genes (AIC = -2454).

We next assessed whether this increased expression was associated with codon or amino
acid usage bias. To avoid repeating the same test structure, results for codon usage are
presented concisely: codon and amino acid usage differed significantly between imprinted and
endosperm genes, although only two biases were shared by PEGs and MEGs (under-usage
of GTC and over-usage of tryptophan; Table S3). However, PHE scores (an index of codon
impact on expression, derived from overexpressed genes in *A. thaliana*; Sahoo et al., 2019)
did not differ significantly across biased and unbiased codons (all p values > 0.09), indicating
that codon usage differences were not associated with expression variation.

Figure 1: Imprinted genes are associated to higher expression level by gene copies than in endosperm genes. The grey dots represent the mean expression level obtained for one gene. The red dots represent the mean expression level for each group of gene. The red lines represent 5% and 95% of the means obtained for endosperm genes after 1,000 resampling of 29 genes (number of imprinted genes considered in study). If the observed means of PEG and MEG are outside of these red lines, they are considered significantly different from the mean in endosperm genes, following the bootstrap approach.

The MEGs and PEGs exhibit phylogenetic signals of negative selection in Brassicaceae

To explore phylogenetic patterns, we analysed genes homologs in *A. lyrata* and 21
Brassicaceae species (Table S4). As a proxy for gene turnover, we compared the number of
species with homologous sequences across groups. Thus, endosperm genes were conserved
across significantly more species than other categories, and PEGs more than control genes
(Fig 2; Table S5). The best model to explain this pattern (AIC = 8696) was this that include
the group of genes, the gene size (coefficient = 9.74-04) and the GC proportion (coefficient =
0.19), whereas the worst model (AIC = 9104) was this that exclude the genes size.

Gene-tree topologies of MEGs and PEGs did not differ significantly in imprinted genes from
the species tree (based on Robinson–Foulds RF distances; bootstrap tests; p value = 0.60 for
PEGs; p value = 0.34 for MEGs), so branch lengths (proxy for evolutionary rates; Fig. S1)
were estimated directly from the species-tree topology to avoid repeating similar topological
tests for genes having homologs in at least three species (including *A. lyrata*). Thus,
endosperm and imprinted genes show significantly longer branches than in control genes (Fig
2; bootstrap; Table S5), even if MEGs exhibited significantly shorter branch lengths than
endosperm genes. The best model to explain this pattern (AIC = -15231) was this that include
the group of genes, the gene size (coefficient = 9.73e-06), the CDS proportion (coefficient = -
8.10e-04) and the GC proportion (coefficient = 9.63e-04), whereas the worst model (AIC = -
14976) was this that exclude the genes size.

Finally, mean D_{NS}/D_S between *A. lyrata* and the other Brassicaceae species decreased
significantly in imprinted genes (Table S5), indicating a signal of purifying (negative) selection.

The best model to explain this pattern (AIC = 13382) was this that include the group of genes
and the GC proportion (coefficient = 0.18), whereas the worst model (AIC = 13401) was this
that exclude the group of genes.

Together, these phylogenetic patterns indicate long-term evolutionary constraints in
endosperm genes, a signal that can arise under ancient balancing selection or strong
functional conservation. Importantly, these inferences reflect deep evolutionary timescales
and therefore cannot be directly compared to population-level signatures of balancing
selection, which we analyse in the next section.

Figure 2: Imprinted genes present phylogenetic signal of negative selection compared
to endosperm genes across the Brassicaceae. A: Variations of the number of species with
homologous genes by group of genes. B: Variations of the mean branch length by gene based
on the phylogenetic tree obtained by HKY85 model based on the 1055 endosperm genes
found in all species. C: Variations of the D_{NS}/D_S by group of genes with a focus on values lower
than 5 (top right). Two extremes values (>200) were removed. The grey dots represent the
mean value obtained for one gene. The red dots represent the mean in each group of gene.
The dashed lines represent 5 and 95% of the means obtained for endosperm genes after 1000
resamples of 100 genes (size of group of control genes). If the real mean of control genes was
not between these dashed lines, the value was considered as significantly different than in
endosperm genes, following bootstrap approach. The red lines represent 5 and 95% of the
means obtained for endosperm and control genes after 1000 resamples of 29 genes (number
of imprinted genes considered in study). If the real means of PEG and MEG were not between
these red lines, the values were considered as significantly different than in endosperm or
control genes, following bootstrap approach.

Imprinted genes show elevated π_{NS}/π_S independent of mating system

To evaluate the effect of mating system and imprinting on polymorphism, we compared
Tajima's D, π and the π_{NS}/π_S for imprinted, endosperm and control genes across nine
allogamous and nine autogamous populations (Table S6). A previous study reported an effect
of mating system on seed size mediated by pollen in these populations (Willi, 2013),
suggesting decreased intensity of parental conflict in autogamous populations.

Under directional selection, reductions of Tajima's D, π and π_{NS}/π_S are expected, whereas
balancing selection is expected to increase these statistics. Because the calculations these
statistic required sufficient polymorphism in all populations, genes with fewer than 0.5%
polymorphic sites were excluded from Tajima's D, and genes with $\pi_S = 0$ from π_{NS}/π_S
analyses, resulting in a significantly higher retention rates for endosperm genes than for
control genes for Tajima's D and π_{NS}/π_S (binomial tests; Table S7). However, more
importantly, these filters retained similar proportions of genes across endosperm and
imprinted genes, and across imprinted and control genes (Table S7). Thus, the observed
contrasts in Tajima's D or π_{NS}/π_S for the imprinted genes cannot be attributed to unequal gene
retention nor to systematic biases against low-diversity loci, supporting that the signals reflect
genuine biological differences rather than filtering artefacts.

In allogamous populations, the Tajima's D was significantly higher in endosperm genes than
in control genes and PEGs, while the π and the π_{NS}/π_S showed no significant differences.
Imprinted genes showed an overall increase in π_{NS}/π_S, while MEGs displayed a significantly
lower π relative to control genes (Fig 3; Table S8). The best model to explain this pattern of
Tajima's D (AIC = 9322) was this that include only the gene size (coefficient = -1.59e-04),
whereas the worst model (AIC = 9337) was this that exclude only the gene size. For the π,
the best model (AIC = -36369) was this that include the gene size (coefficient = -9.89e-08)
and the GC percentage (coefficient = -5.59e-05) whereas the worst model (AIC = 9337) was
this that exclude only the gene size. Finally, the best model to explain this pattern of T π_{NS}/π_S
(AIC = -721) was this that include only the proportion of cds (coefficient = 2.21e-03), whereas
the worst model (AIC = 9337) was this that include all the parameters.

In autogamous populations, Tajima's D in endosperm genes no longer differed from controls
(Table S8). At the opposite, MEGs showed a significantly higher Tajima's D than endosperm
genes. Moreover, MEGs and endosperm genes exhibited decreased π and π_{NS}/π_S compared
to control genes (Table S8). These trends are explained by mating-system that only
significantly affects endosperm genes (Kruskal-Wallis tests; Table S9).

Figure 3: Imprinted genes present an unusual decrease of Tajima's D associated to an increase of π_{NS}/π_S in allogamous populations compared to endosperm genes. A: Variations of the Tajima's D by group of genes. Only the genes with a mean between -5 and 5 are indicated. B: Variations of the π_{NS}/π_S by group of genes. Only the genes with a mean between 0 and 1.1 are indicated. The grey dots represent the mean value obtained for one gene. The red dots represent the mean in each group of gene. The dashed lines represent 5 and 95% of the means obtained for endosperm genes after 1000 resamples of 100 genes (size of group of control genes). If the real mean of control genes was not between these dashed lines, the value was considered as significantly different than in endosperm genes, following bootstrap approach. The red lines represent 5 and 95% of the means obtained for endosperm and control genes after 1,000 resamples of 29 genes (number of imprinted genes considered in study). If the real means of PEG and MEG were not between these red lines, the values were considered as significantly different than in endosperm or control genes, following bootstrap approach.

A limited number of MEGs exhibited strong signals of directional selection.

To assess whether imprinted genes show an enrichment in signatures of directional selection, we tested whether they occur more frequently than expected in the extreme tail of the empirical distribution of endosperm genes. Candidates for directional selection were defined as the 5% most extreme values for branch length, Tajima's D, or π_{NS}/π_S in allogamous populations. However, none of these proportions were significantly higher than expected (p = 0.66, 1 and 1 respectively).

We identified only four MEGs potentially under strong directional selection (Table S1). Two MEGs were associated with a reduced branches length (Table S1): one lacked homologous gene in *A. thaliana*, the second encoded an AGAMOUS-like protein expressed during proliferative endosperm development (Day et al., 2008), involved in female gametophyte, seed and early endosperm development (Bemer et al., 2010; Zhang et al., 2018). One MEG identifing by a reduced Tajima's D encodes a F-box domain–containing protein regulated by non-CG DNA methylation in *A. thaliana* (Garro et al., 2025) and associated with the regulation

of stress-responsive genes (Papdi et al., 2008). Finally, one MEG encoding a Dof-type zinc

finger protein involved in seed coat formation (Zou et al., 2012) shows a reduced π_{NS}/π_S.

Clear trend of coevolution among imprinted genes in Brassicaceae

Imprinting could promote coevolution between genes favouring paternal or maternal fitness in

an arms race scenario. To test this hypothesis, we investigated potential coevolutionary

signals between PEGs, MEGs and endosperm genes, based on (i) nucleotide diversity (π)

within populations, (ii) branch length in phylogenetic trees, and (iii) Robinson–Foulds (RF)

distances between tree topologies. Because the coevolution hypothesis is more closely

related to noncoding regions, we focused on π rather than π_{NS}/π_S.

First, using thresholds defined from endosperm genes (−0.66 and 3.20 for 1 and 99% of

distribution), 13 significant Pearson's correlation coefficient correlations involving 6 PEGs and

9 MEGs were detected. Both the number of PEGs-MEGs correlations and the number of

imprinted genes involved exceeded expectations (p-values= 4.5e-02 and <1e-03 respectively;

Table S10). Considering endosperm genes significantly correlated with these imprinted genes,

as well as indirect links between imprinted and endosperm genes, the resulting network

contained 14 MEGs, 13 PEGs, and 1,643 endosperm genes (=1,670 genes; Fig 4.A; Table

S10). This network did not show enrichment for specific GO terms compared to random

networks of equal size. Notably, one PEG encoding a chloroplast-localized protein required

for embryonic development (Bryant et al., 2010) was correlated with more genes than 99% of

endosperm genes (Table S1 and S10).

Following the same methodology, the thresholds based on branches length(−0.64 and 0.89)

identified 16 significant correlations among 6 PEGs and 9 MEGs. Both the number of PEGs-

MEGs correlations and the number of imprinted genes involved exceeded expectations (p-

values = 8e-03 and 3e-03 respectively; Table S11). None of these correlations was in common

with the previous 13 significant correlations of polymorphism. Considering endosperm genes

significantly correlated with these imprinted genes, the resulting network contained 14 MEGs,

14 PEGs, and 1,027 endosperm genes (=1,055 genes; Fig 4.B; Table S11). This network was

enriched for the GO categories "*negative regulation of catalytic activity*" and "*syntaxin binding*"

(Table S12). One MEG in this network, encoding the AGAMOUS-like protein previously

mentioned, was correlated with more genes than 99% of endosperm genes (Table S11).

Finally, we investigated gene pairs that showed high similarity together (RF = 0) but higher

deviation from the species topology than 95% of endosperm genes (RF > 0.71 with species

topology; Fig. S1). Based on this approach, the topology of only one PEG encoding an F-box

protein was associated with six endosperm genes (Fig. 4.C; Table S13). This small network

was enriched for the GO term *DNA-binding transcription factor activity* (Table S12). Moreover,

the PEG exhibited a significantly higher number of associations than 99% of non-imprinted

endosperm genes (Table S1; Table S13).

Across all analyses, 22 imprinted genes pairs (29 genes) were identified. To test for

coevolution between specific mutations within these pairs, we applied the clustering approach

implemented in the CoMap package (Dutheil and Galtier, 2007).

Specifically, we searched for groups of 2 to 10 coevolving sites within each candidate pair of

imprinted genes. To limit false positives, the number of such groups detected in candidate

pairs was compared to the number observed in non-candidate pairs of imprinted genes that

were not associated in the previous networks and were present in at least three species.

CoMap did not detect an excess of coevolving residues in candidate pairs relative to non-

candidate imprinted pairs (Table S14). These directly correlated pairs of imprinted genes were

therefore not associated with elevated levels of site-level coevolution.

A) Network based on polymorphism

B) Network based on lenght of branches

C) Network based on topology

Figure 4: Networks of the imprinted and endosperm genes based on A) polymorphism in population , B) length of branches and C) phylogenetic topologies across the Brassicaceae. A) Network based on the Pearson's correlations of π found on genes in the populations. B) Network based on the Pearson's correlations of the length of branches for each gene estimated with the global topology. C) Networks based on the RF distance of topologies. The black and grey lines represent the negative and positive correlations respectively. Blue dots: PEG. Red dots: MEG. White circles: non-imprinted endosperm genes.

Discussion

The kinship theory remains the prominent selective framework for explaining the evolution of genomic imprinting (Patten et al., 2014). According to this theory, imprinted genes are should (I) be involved in resource allocation within systems with multiple mating (Haig 2000), (II) exhibit signatures of directional selection relative to non-imprinted endosperm genes (Day and Bonduriansky 2004; Kasimatis et al., 2017; Table 1), and could present some signals of coevolution among genes (Wilkins and Haig 2001; Willi 2013). However, a selection acting primarily on gene expression may attenuated these genomic signatures, and their sensitivity to mating-system variation remains unclear in this case.

Here, we tested these predictions by comparing imprinted and non-imprinted endosperm genes in *A. lyrata*, an allogamous species that shift to autogamy in some population from North America, and assessing the impact of mating system shifts. While several figures follow a unified graphical structure to facilitate comparison across gene categories, the underlying analyses encompass expression-level differences, population genetic statistics, phylogenetic metrics, and coevolutionary patterns, providing a broader and complementary evaluation of kinship-theory predictions than previous work (Miyake et al., 2009; Hatorangan et al., 2016; Tuteja et al., 2019). Although we detected patterns consistent with predictions, the signals differed across statistics and were not consistently replicated among analyses. This suggests that selection associated with parental conflict may act heterogeneously across loci or primarily through regulatory rather than coding changes, and underscores the importance of integrating multiple lines of evidence when assessing evolutionary hypotheses of imprinting.

Signals of ancient balancing selection in endosperm genes

In the absence of genomic imprinting, genes involved in resource allocation conflict are expected to evolve under balancing selection (Day and Bonduriansky 2004; Sylvestre et al., 2023). Consistently, non-imprinted endosperm genes were more widely conserved across Brassicaceae, suggesting lower turnover, and displayed longer phylogenetic branches (Edwards, 2009; Kasimatis et al., 2017; Llaurens et al., 2017). These assumptions are comforted by the presence of the group of genes in best models predicted by AIC for these statistics. In addition, these genes showed elevated Tajima's D values in allogamous populations. However, the absence of a parallel increase in polymorphism, and the absence of this parameter in best model predicted by AIC, suggests that this signal may not solely reflect balancing selection.

Hypotheses tested	References	Statistic analysed	Expectation on endosperm genes	Expectation on imprinted genes
Imprinted genes are related to tissue implicated on resource's allocations	Haig, 1997	Proportion of endosperm genes	NA	**Increase**
Imprinted genes are selected to optimize gene expression	Patten et al., 2014	Expression by copy		**Upregulated with PEGs > MEGs***
Selection on expression promotes codon and amino acid bias in imprinted genes		Usage of each codon and amino acid		**Increase for some*, decrease for others***
Reducing of intralocus conflict in imprinted genes promotes lower conservation of genes across species and higher turnover	Haig 2000; Spencer 2000	Number of species with homologous genes	**Increase***	**Decrease***
Reducing of intralocus conflict in imprinted genes promotes directional selection		Evolutionary rate by length of branches	**Increase***	Increase if positive*, **decrease if negative* ****
	Spencer 2000	Sens of selection by D_{NS}/D_S	**No variation***	Increase if positive*, **decrease if negative***
Reducing of intralocus conflict in imprinting genes promotes reduced polymorphism in allogamous population by directional selection, more intensively in non-synonymous sites	Spencer 2000; Day and Bonduriansky 2004	Tajima's D	**Increase***	Decrease*
		π	Increase*	Decrease*
	Haig 2000	π_{NS}/π_S	**No variation***	Decrease*
Reducing of intralocus conflict in autogamous populations promotes directional selection in non-imprinting genes		Tajima's D, π_{NS}/π_S	**Decrease**	**No variation if already fixed**, increase if relaxed selection
Possible antagonist coevolution between genes (arms race)	Moore and Haig, 1991	π, length of branches, topology by RF distance	NA	**Excess of PEG–MEG correlations***

Conflict over resource allocation is predicted only in multiple mating systems (Haig 2000).
This conflict is resolved in autogamous systems, potentially allowing directional selection on
endosperm genes. In line with this, Tajima's D and π_{NS}/π_S decreased in these populations,
indicating that relaxation of conflict facilitates directional selection and that the effect of mating
system is detectable even in populations that have recently and partially transitioned to
autogamy (Ross-Ibarra et al., 2008; Foxe et al., 2010; Hough et al., 2013; Willi et al., 2018).

It is important to emphasize that mainly the phylogenetic analyses, but only Tajima's D for the
population-genetic ones, suggest signals compatible with balancing selection in endosperm
or imprinted genes. Phylogenetic patterns reflect long-term evolutionary processes, potentially
including ancient or persistent balancing selection, whereas population-level statistics detect
more recent forms of balancing selection and are sensitive to demography, recombination,
and reduced polymorphism. The absence of consistent population-level evidence therefore
does not contradict the phylogenetic signal but rather indicates that any balancing selection
acting on these loci is either ancient, weak, or too localized to be detected with gene-wide
summary statistics.

Codon usage bias and modulation of expression in imprinted genes

Kinship theory predicts that imprinted genes should be preferentially expressed in the
endosperm (Haig 2000; Patten et al., 2014). Consistent with this prediction, imprinted genes
were enriched among endosperm-expressed genes. Interestingly, the MEGs are more related
to endosperm than PEGs, as observed in maize (Eaux et al., 2013). Yet, more than half were
also upregulated in other tissues suggesting that distinct selective pressures unrelated to
parental conflict may act on them.

An important debate suggests that detecting directional selection in imprinted genes is difficult
because selection may primarily target gene expression (Patten et al., 2014). Consistent with
this idea, both PEGs and MEGs showed increased expression per gene copy, and the group
of genes is the more essential parameter in models to predict expression pattern by AIC.
Interestingly, the PEGs are associated to an increase of expression level, despite that gene
size reduced globally the efficiency of expression in some study(Grishkevich and Yanai, 2014),
including in our study.

Moreover, imprinted genes displayed codon usage biases relative to endosperm genes,
however, we were not able to determine if these biases enhance or inhibit the expression of
imprinted genes (Sahoo et al., 2019). More studies with direct analyses of effect on expression
of the different codon usage are required to verify this hypothesis in future.

Phylogenetic signals of ancient purifying selection in imprinted genes.

The kinship theory is expected to promote directional selection on imprinted genes (Wilkins

and Haig 2001; Mills and Moore 2004; Day and Bonduriansky 2004; Kasimatis et al., 2017).

Consistently, our phylogenetic analyses showed that imprinted genes exhibit lower

conservation of homologs across Brassicaceae species, likely reflecting a higher turnover

driven by accelerated evolutionary rates. Furthermore, imprinted genes displayed reduced

D_{NS}/D_S, indicative of strong negative directional selection, and the group of genes is the more

essential parameter in models to predict this pattern by AIC.

This signal of purifying selection was also evident in MEGs when branch lengths were

compared to those of endosperm genes. We note that two MEGs, including AGAMOUS-like

protein implicated in one coevolutionary network, exhibited branch lengths suggesting a

particularly strong negative selection. Our observations were not in line with the positive

selection detected on PEGs on the autogamous species *A. thaliana* and *C. rubella*

(Hatorangan et al., 2016; Tuteja et al., 2019), raising the question of the effect of the mating

system on the selection of imprinted genes in these species.

Nevertheless, caution is warranted in attributing the observed purifying directional selection

solely to imprinting. In fact, the detected purifying selection could be explained by alternative

explanations (Patten et al., 2014), including (1) an increase of the efficiency of selection by

hemizygous expression (Haig and Westoby 1989), (2) selection against transposable element

(Padilla-García et al., 2025) or (3) local adaptation (Kremer et al., 2025). However, none of

the current models rigorously predict the effects of these mechanisms on polymorphism and

divergence, making it difficult to discriminate among alternative explanations.

Implications for reproductive isolation and adaptation

The evolutionary consequences of purifying selection on imprinted genes remains poorly

understood. Directional selection on imprinted genes could (1) influence adaptive potential by

shaping the spectrum of selectable alleles within populations (Cvijović et al., 2018), (2)

contribute to speciation processes by generating genetic incompatibilities between lineages

(Seehausen et al., 2014), (3) facilitate the transition to apomixy in plants (Hojsgaard and

Schartl 2021) and (4) promote the establishment of reproductive isolation (Maheshwari and

Barbash 2011; Cvijović et al., 2018; İltaş et al., 2021), more particularly through endosperm-

based post-zygotic barriers (Haig 2000; Spencer 2000; Kinoshita 2007; Köhler et al., 2010).

Previous studies empirically demonstrated that some imprinted genes are involved in these

barriers, including in the autogamous species *A. thaliana* (Wolff et al., 2015), between the

allogamous *Solanum peruvianum* et *S. chilense* (Florez-Rueda et al., 2016), between the allogamous *C. grandiflora* and the autogamous *C. rubella* and *C. orientalis* (Lafon-Placette et al., 2018), between the allogamous *Mimulus guttatus*, *M. luteus* (Kinser et al., 2021), *M. caespitosa* and *M. tilingi* (Sandstedt and Sweigart 2022), and between the autogamous wheat species *Triticum durum* and *T. aestivum* (Yang et al., 2023).

Imprinted genes may also act as central hubs within complex regulatory networks that contribute to hybrid vigour and reproductive barriers (Botet and Keurentjes 2020). However, despite these observations, the broader evolutionary implications of these network-level effects remain largely unresolved, particularly regarding how they interact with demographic history, mating system variation, and other selective pressures.

Imprinted genes are associated with singular signals of selection in North America

Consistent with the expectations under the kinship theory, PEGs were associated with reduced Tajima's D (Haig 2014). However, this signal was not observed for π_{NS}/π_S. More interestingly, π_{NS}/π_S values were higher in both PEGs and MEGs compared to endosperm and control genes, which could indicate a relaxation of selection on coding regions (Galtier and Rousselle 2020). This apparent relaxation may not reflect parental conflict but instead could result from the recent colonization of North America by *A. lyrata* from Europe (Ross-Ibarra et al., 2008; Otto et al., 2015; Willi et al., 2018) that could have reduced effective population size (Galtier and Rousselle 2020) and the efficiency of purifying selection (Laenen et al., 2018), especially on transposable element (Padilla-García et al., 2025). Testing this hypothesis would require future comparisons of selection signals in endosperm and imprinted genes across allogamous populations with varying levels of polymorphism. Nevertheless, we note that two MEGs exhibited reduced Tajima's D or π_{NS}/π_S, consistent with ongoing directional selection.

Imprinted genes are not sensitive to shifting mating system in A. lyrata

Conflict over resource allocation is expected to be reduced in autogamous lineages relative to allogamous ones (Haig 2000; Brandvain and Haig 2005). However, most studies of genetic and phylogenetic signatures of selection associated with imprinting have focused on autogamous species, leaving the effect of shifting mating system on polymorphism of imprinted genes unresolved. In the autogamous species *A. thaliana*, the imprinted *MEDEA* locus shows increased polymorphism but not in its allogamous sister species *A. lyrata* (Spillane et al., 2007; Miyake et al., 2009). However, PEGs were associated to positive selection in the autogamous species *A. thaliana* and *C. rubella* (Hatorangan et al., 2016;

Tuteja et al., 2019). In our study, the selection signals in imprinted genes, contrary to
endosperm genes, were insensitive to the mating system. The absence of variation in Tajima's
D could be explained if directional selection had already fixed the optimal alleles prior to the
transition from allogamy to autogamy, which could also explain the signal of positive selection
detected in PEGs in *A. thaliana* and *C. rubella* (Hatorangan et al., 2016; Tuteja et al., 2019).
However, this explanation does not account for the high π_{NS}/π_S observed in allogamous and
autogamous populations. Theoretical studies are necessary to estimate the cumulated effect
of selection related to kinship theory, imprinting, demographic history and mating system.

Imprinted genes present signal of ancient and recent coevolution

Another potential, but not mandatory, predicted outcome of the kinship theory is the
coevolution of some imprinted genes with antagonistic effects on offspring growth (Moore and
Haig, 1991; Wilkins and Haig 2001; Mills and Moore, 2004). To investigate this, we applied
both population genomic and phylogenetic approaches (Dutheil and Galtier 2007). We found
that most PEGs and MEGs were connected within coevolutionary networks, with a significantly
higher number of direct associations than expected by chance. These networks encompass
essential functions for endosperm development.

For instance, one network based on branch-length correlations was enriched in enzymatic
activities related to cell wall formation and modification, a key process during endosperm
cellularization in Arabidopsis (Hehenberger et al., 2012), which in turn influences seed size
(Pignatta et al., 2018; Zhang et al., 2020), a likely target of parental conflict. These correlated
evolutionary rates could reflect equivalent selective constraints (Galtier and Rousselle 2020)
or just a common chromosomic context (e.g. transposable element) that modify mutation rate
(Lynch 2010). Interestingly, this network appeared to be strongly dependent on a MEG, an
AGAMOUS-like protein, characterized also by reduced branch lengths, consistent with
expectations that coevolution may drive coordinated evolution among genes primarily through
rare compensatory mutations (Fraser et al., 2002)

The polymorphism-based network involved genes related to syntaxin-binding proteins, crucial
for membrane activity, molecular transport, and protein storage, whereas the topology-based
network was enriched in DNA-binding transcription factors, including those regulating seed
development, tissue differentiation, and storage metabolism (Noguero et al., 2013).

These signals of coevolution could promote the maintenance of the gene functions (Clark et
al., 2012), limit their evolutionary independency (Yang and Bielawski 2000) and be another
source of reproductive barrier (Presgraves 2010). Interestingly, the coevolution signal did not

involve increased site-by-site coevolution, suggesting that protein–protein molecular interactions, often drivers of coevolution (Green et al., 2021), played a limited role here. Instead, the observed patterns may reflect selection acting on gene dosage rather than protein products (Clark et al., 2012).

However, as correlations between evolutionary statistics can arise from mutation-rate variation, recombination environment or shared functional constraints, we interpret them as exploratory signals only. The more stringent CoMap analysis (Dutheil & Galtier 2007) did not reveal enrichment for residue-level co-evolution, indicating that correlation-based networks should be treated cautiously and do not by themselves confirm co-evolution.

Main limitations in our study

Although our study combines multiple phylogenetic and population genomic statistics, the signals detected were not consistently congruent across approaches. This inconsistency may reflect evolutionary processes not accounted for, (such as demographic history or chromosomal context) as well as several methodological limitations, outlined below.

First of all, estimating Tajima's D from pooled sequencing data is challenging because SFS-based statistics are sensitive to coverage variation and sampling noise, which increase estimation variance (Futschik & Schlötterer 2010; Ferretti et al. 2013; Lynch et al. 2014). Although we minimized dataset-specific biases by comparing genes processed under identical conditions, technical noise cannot be fully excluded. In addition, short genes, such as many MEGs or control genes, yield fewer segregating sites, reducing the reliability of Tajima's D (Tajima 1989).

A second limitation concerns the imprinting dataset. The set of imprinted genes analysed was derived from a study on a geographically and genetically distinct allogamous subspecies European *A. lyrata petraea* (Klosinska et al., 2016), assuming conservation of imprinting, because no within-lineage data on imprinting in *A. lyrata* exist. However, imprinting shows limited conservation within *A. thaliana* (Gehring et al., 2011; Wolff et al., 2011; Hsieh et al., 2011; Pignatta et al., 2014), between *A. thaliana* and *A. lyrata* in our data, and generally across species (Eaux et al., 2013; Hatorangan et al., 2016; Roth et al., 2018; Flores-Vergara et al., 2020). It is therefore possible that several genes classified as imprinted in our analyses are not imprinted in the studied populations , especially given known population-level variation in TE-driven imprinting (Padilla-García et al., 2025). Future population-specific characterizations of imprinting will be critical to resolve this issue..

Moreover, some imprinted genes identified by Klosinska et al. (2016) may have been
inadvertently excluded from our dataset due to our method for detecting tissue-specific
expression, which relied on RNA-seq data pooled from both allogamous and autogamous
populations. More controlled and population-specific experimental designs will be required to
recover the complete set of endosperm-expressed imprinted genes relevant to parental
conflict.

A last important limitation is about the panels of genes compared (control, endosperm and
imprinting genes). To be congruent with previous studies (Spillane et al., 2007; Miyake et al.,
2009; Hatorangan et al., 2016; Tuteja et al., 2019), we also choose genes in panels based on
expression patterns. However, the three gene panels differ substantially in gene size, CDS
proportion, and GC content, factors known to influence both polymorphism and phylogenetic
metrics, but seldom considered in previous studies (Spillane et al., 2007; Miyake et al., 2009;
Hatorangan et al., 2016; Tuteja et al., 2019).

Endosperm genes and PEGs are substantially longer, whereas MEGs and control genes are
shorter. The size of the gene is an important parameter that could modify an important amount
of statistic. For example, larger genes tend to evolve more slowly due to increased functional
constraint and pleiotropy (Massey 2013; Lopes et al., 2021), display higher conservation and
reduced turnover (Abdel-Haleem 2007; Koonin 2015), and can inflate the number of species
with detectable homologues. In this sense, we found a positive correlation between the
number of species with homologous genes and the gene size. This may explain partially the
higher homology counts observed for PEGs and endosperm genes relative to control genes.
Conversely, we found that the gene size is negatively correlated with polymorphism and
Tajima's D. However, this tendency contradict that longer genes as endosperm genes
presented higher Tajima's D or that the shorter genes as the MEGs were associated to lower
polymorphism, suggesting that, maybe, the effects of parental conflict on polymorphism in
these genes are underestimated, more specifically for MEG because shorter genes may yield
less stable estimates of Tajima's D (Tajima 1989). We found also a negative correlation with
the expression level, that could explain partially why the expression of MEG is more efficient,
even if the group of gene stays the more important parameter predicted by AIC, but could not
explain why the expression of PEG is higher than endosperm genes.

The effect of gene size on branch length is more debated: while longer genes trivially
encompass more substitutions (Graur & Li 2000), several studies show that long, constrained
genes tend to exhibit shorter branches due to stronger purifying selection (Kimura 1983;
Abdel-Haleem 2007; Massey 2013; Koonin 2015). These opposing forces complicate
interpretation of branch-length differences between MEGs, PEGs and endosperm genes. In

our study, we found a positive correlation between the length of the branches and the gene
size, maybe explained by the amount of endosperm genes with long branches. This correlation
suggests that the longer branches observed in PEGs and endosperm genes are partially
explained by the size of the genes, but could not explain the longer branches observed in
MEGs.

Moreover, both imprinted and endosperm genes show elevated CDS content relative to
controls. Thus, this parameter could not explain the increase of efficiency observed in
expression of imprinted genes, despite the negative correlation detected. In the same way,
this parameter could not explain the reduced π_{NS}/π_S observed in imprinted genes compared
to endosperm genes, despite the positive correlation detected. Interestingly, high CDS
proportion is associated with stronger purifying selection (Hughes et al., 2003), which can
reduced the length of the branches, as observed in our study. However, the increase of length
of branches detected in endosperm and imprinted genes is the reverse pattern predicted,
suggesting that the effect on length of branches in these genes is maybe underestimated.

Finally, imprinted genes show reduced GC content relative to endosperm and control genes.
GC-poor genes tend to have reduced recombination rates, lower polymorphism (Wright et al.,
2003), and higher AT-biased substitution rates, which can artificially lengthen branches (Duret
& Galtier 2009), and explain the longer branches observed in PEG and endosperm genes.
However, contrary to these expectations, we found a negative collation between the length of
the branches and the GC content, maybe explain by the important amount of endosperm
genes. Interestingly, we found also a negative correlation between the polymorphism and the
GC proportion in our study, that suggest that the reduced polymorphism observed in MEGs is
maybe underestimated. Low GC also reduces the likelihood of finding conserved homologues
across species (Urrutia & Hurst 2003), as found in our study, but is not able to explain why the
PEGs and the endosperm genes are associated to higher number of species with homologous
genes despite lower GC content than in control genes. Low GC is expected to also inflate
D_{NS}/D_S by increasing the probability of non-synonymous changes (Kryazhimskiy & Plotkin
2008). However, we detected the reverse correlation in our study, that could explain why
imprinted genes are associated to a reduced D_{NS}/D_S.

The lower GC of imprinted genes is also unexpected given their elevated expression (Sharp
& Li 1987; Bernardi 2000), suggesting that expression differences may be underestimated.
For these reasons, although the three panels differ in genomic features, endosperm-
upregulated genes remain the only biologically appropriate baseline for testing imprinting-
related hypotheses. A feature-matched control panel would reduce variation but would not
provide a more relevant comparison for detecting signatures associated with parental conflict.

Together, these factors indicate that differences in gene size, CDS proportion, and GC content
may partly account for the heterogeneity of evolutionary signals observed across statistics.
Some signatures consistent with parental conflict may therefore be underestimated or
confounded, while others may reflect intrinsic genomic properties rather than imprinting-
related selective pressures. Future work will require gene panels matched not only for
expression, but also for fine-scale genomic features.

Although several genomic features differ among gene panels, we performed targeted
robustness checks to ensure that competing non-selective scenarios did not spuriously
generate the patterns we report. In particular, the SNP-density threshold retained comparable
fractions of MEGs, PEGs, endosperm genes and controls, ruling out retention bias as a driver
of polymorphism contrasts. Moreover, all evolutionary inferences were tested against
empirical null distributions from endosperm-upregulated genes, which share expression
patterns and chromatin context with imprinted genes, limiting confounding effects of gene
architecture or local mutation rate. These checks do not eliminate all alternative explanations,
but they demonstrate that the signals observed are unlikely to arise solely from genomic
artefacts or filtering procedures.

Conclusions

Overall, our analyses revealed that endosperm-expressed and imprinted genes in *A. lyrata*
exhibit evolutionary signatures partly consistent with kinship theory, including evidence for
ancient balancing selection in endosperm genes, expression-level regulation in imprinted
genes, ancient purifying selection, and coevolutionary interactions. However, signals at the
population level were less congruent with theoretical expectations, likely reflecting
demographic history, mating system shifts, and the poor conservation of imprinting across
populations. Future work should combine empirical surveys of imprinting with theoretical
models explicitly incorporating demography and mating system variation to refine predictions
about the evolutionary dynamics of imprinted genes.

Materials and Methods

We hypothesized, based on kinship theory, that imprinted genes should display distinct
genomic signatures of selection compared to non-imprinted endosperm genes (Table 1). To
test this prediction, we designed an approach combining phylogenetic analyses and
polymorphism data. Specifically, we focused on a set of 27 maternally expressed genes
(MEGs) and 46 paternally expressed genes (PEGs) previously characterized in *A. lyrata*
(Klosinska et al., 2016) and unambiguously annotated in the reference genome (Hu et al.,
2011; https://doi.org/10.5281/zenodo.17287897). These imprinted genes were systematically
compared against two control datasets: (i) 2,703 non-imprinted genes upregulated in the
endosperm, and (ii) a reference panel of 100 randomly selected ubiquitous genes, expressed
across multiple tissues. This design allowed us to disentangle signatures of selection
specifically associated with imprinting from those affecting endosperm genes more generally
or the whole genome.

We used several GitHub repositories, each corresponding to a distinct analytical module. This
structure reflects the fact that the datasets and workflows differ substantially across analyses
and that several scripts were developed independently and used in other publications. For
clarity, all repositories are indexed in a Zenodo record (https://zenodo.org/records/17954040),
and each includes documentation, test data, and commented scripts. This multi-repository
organization was reviewed and approved for reproducibility and FAIR compliance by the PCI
Ecology Data Editor (Checklist PCI Ecology 867).

Expression data

RNA-seq expression data from multiple tissues, including leaf, root, pistil, and pollen, were
obtained from two *A. lyrata* populations representing 27,820 genes from both European and
North American lineages, as described in İltaş et al. (2024). To maximize gene representation
and increase the statistical power, the data from both mating systems was pooled. To
complement these datasets, we also incorporated endosperm RNA-seq data derived from five
allogamous *A. lyrata* individuals, previously generated by Klosinska et al. (2016). The
endosperm data were reprocessed and integrated into our analytical pipeline following the
instructions and R script available in GitHub (https://github.com/thelovelab/DESeq2). To
identifying endosperm-specific genes, we retained only those that showed significant
upregulation (log2FC > 1, adjusted P < 0.05) in all pairwise comparisons against other tissues.
The RNA-seq datasets for leaf, root, pistil, and pollen are available in the NCBI BioProject
database under accession PRJNA1067428, while the endosperm RNA-seq dataset is
accessible under accession GSE76076.

Genes clusters

We used the list of the 49 PEGs and 35 MEGs found in *A. lyrata* (Klosinska et al., 2016). We kept the genes having conserved coordinates in the reference genome of *A. lyrata* V1.0.23 (Hu et al., 2011; https://doi.org/10.5281/zenodo.17287897), resulting in a final list of study 27 MEGs and 46 PEGs.

To disentangle signatures of selection associated specifically with the endosperm, where conflicts over resource allocation are expected, from those potentially arising through other selective processes, we separated imprinted genes upregulated in the endosperm (14 PEGs and 15 MEGs) from imprinted genes not showing endosperm-specific upregulation (32 PEGs and 12 MEGs), as determined by differential expression analysis.

To evaluate whether the observed proportions of MEGs and PEGs upregulated in endosperm deviated from theoretical expectations under kinship theory, we compared them with null distributions generated by 10,000 random sampling of 49 and 35 genes (size of PEGs and MEGs groups) between the 27,820 genes expressed genes.

For all subsequent analyses, we focused exclusively on endosperm-upregulated imprinted genes and compared them with two reference groups: the set of 2,703 non-imprinted endosperm-upregulated genes (hereafter, "endosperm genes"), the sets of 15 MEG and 14 PEG upregulated in endosperm (hereafter, "MEG" and "PEG") and a set of 100 randomly selected ubiquitously expressed genes (hereafter, "control genes"). To remain consistent with previous studies of imprinting and parental conflict (Spillane et al., 2007; Miyake et al., 2009; Hatorangan et al., 2016; Tuteja et al., 2019), the control panel was defined on the basis of expression patterns only. This set is intended to represent a genome-wide baseline unrelated to endosperm-specific functions or parental conflict, rather than a genomic feature–matched control. Consequently, no constraints were imposed on gene size, GC content, CDS proportion, or chromosomal position. However, for each statistics studied after, we compared the full model included gene group, gene length, CDS proportion and GC content as explanatory variables using the Akaike Information Criterion (AIC). We then evaluated a series of reduced models in which one parameter was removed at a time and retained the model with the lowest AIC value. This procedure was iterated until the best-supported model was identified using the function StepAIC of the MASS library (version 7.3-65) in R. This approach allowed to determine (1) whether gene group remained a significant predictor in the best model and (2) whether genomic features accounted for the observed trends in the analysed statistics.

To test whether expression levels conformed to predictions of higher optimal expression in
PEGs than in MEGs, we normalized expression by gene copy number and compared
observed values with bootstrap distributions from non-imprinted endosperm genes (10,000
resamples of 29 genes). To account for gene dosage, expression values were normalized by
the expected number of expressed copies: three for non-imprinted endosperm genes, two for
MEGs, and one for PEGs.

Based on the sequences from the reference genome of *A. lyrata* (Hu et al., 2011;
https://doi.org/10.5281/zenodo.17287897), we conducted bootstrap analyses to compare the
gene size, the GC and the CDS content (%) of PEGs and MEGs with values obtained from
10,000 resamples of 29 endosperm genes and control genes independently. Similarly, we also
conducted bootstrap analyses to compare the values on control genes with values obtained
from 10,000 resamples of 100 endosperm genes.

The　　R　　code　　for　　the　　bootstrap　　tests　　is　　available　　in　　GitHub
(https://github.com/leveveaudrey/bootstrap).

Estimation of codon and amino acid bias

Based on the sequences from the reference genome of *A. lyrata* (Hu et al., 2011;
https://doi.org/10.5281/zenodo.17287897)　　and　　a　　custom　　python　　code
(https://github.com/leveveaudrey/codon-bias), we estimated the proportion of each amino acid
within the gene and the proportion of each possible codon for each amino acid. Then, we
conducted bootstrap analyses to compare these proportions in PEGs and MEGs with values
obtained from 10,000 resamples of 29 endosperm genes. The R code for the bootstrap test is
available in GitHub (https://github.com/leveveaudrey/bootstrap). Thus, we defined the codons
"overused" as codons more used in imprinted genes than in endosperm genes, the codons
"underused" as codons less used in imprinted genes than in endosperm genes and the
"unbiased" codons as codons equivalently used in imprinted genes than in endosperm genes.

To test if the overused or underused codons were related to gene expression, we compared
the variations of PHE score induced by codons with the score of unbiased codons by
permutations test with 1,000 replications for MEGs and PEGs independently. The PHE score
was estimated based on overexpressed genes in *A. thaliana* and reflect the importance of a
codon on genes expression (Sahoo et al,. 2019). We hypothesized that codons overused in
imprinted genes were associated to increases of PHE score because associated to increases
of gene expression whereas the underused codon were associated to reduce PHE scores.

The   code   used   for   permutation   test   was   available   in   GitHub
(https://github.com/leveveaudrey/permutation-test-)

Distribution of the genes across Brassicaceae

We assessed whether the imprinted genes presented specific signals of selection compared
to the endosperm and control genes in the phylogeny of 22 Brassicaceae. All the complete
sequence of reference genomes (fasta files) were extracted from the Phytozome database
(Goodstein et al., 2012; Table S4). We extracted the sequences of the genes in the *A. lyrata*
reference genome (Hu et al., 2011) using bedtools V2.30 (Quinlan et al., 2010). Then, we
retained the best hit of these sequences on 22 other Brassicaceae using BLAST V2.10
(Altschul et al., 1990) with e-value < 1e-5. If more of one hit was found for a gene in one
specie, we filtered the best hit as the hit with: the higher length alignment, the higher % of
identity, then the higher coverage on the gene in *A. lyrata* and the higher reciprocal coverage
on the gene in the other species. Then we extracted the sequence of the best hit for each
species of each gene. These steps were integrated into a pipeline of analysis available in
GitHub (https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee).

To test for significant differences in the number of species with homologous genes across the
group of genes, we conducted bootstrap analyses to compare the value on PEGs and MEGs
with values obtained from 10,000 resamples of 29 endosperm genes and control genes
independently. Similarly, we conducted bootstrap analyses to compare the values on control
genes with values obtained from 10,000 resamples of 100 endosperm genes. The R code for
the bootstrap test is available in GitHub (https://github.com/leveveaudrey/bootstrap).

Divergence of the imprinted genes across Brassicaceae

The rates of nonsynonymous mutations (D_{NS}) and synonymous mutations (D_S) are assumed
to follow the neutral evolutionary process and the ratio D_{NS}/D_S is therefore approximate to the
selective pressure on the protein product of a gene. An increase of the D_{NS}/D_S in genes
indicates positive selection, while a decrease indicates negative selection. To test whether
imprinted genes display accelerated evolutionary rates compared to endosperm genes, using
the annotation of the 0-fold and 4-fold degenerate sites in *A. lyrata*, we estimated for each
gene the mean evolutionary rates D_{NS}/D_S between the reference genome of *A. lyrata* and all
the species with an homologue found. Only the genes found in at least 3 species and with a
D_S higher than 0 were considered. These steps were integrated into a pipeline of analysis
available in GitHub (https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee).

To test for significant differences in the D_{NS}/D_S across the group of genes, we conducted
bootstrap analyses to compare the value on PEGs and MEGs with values obtained from
10,000 resamples of 29 endosperm genes and control genes independently. Similarly, we
conducted bootstrap analyses to compare the values on control genes with values obtained
from 10,000 resamples of 100 endosperm genes. The R code for the bootstrap test is available
in GitHub (https://github.com/leveveaudrey/bootstrap).

Topology of the genes across Brassicaceae

For each gene, we aligned the gene and all its best hits found across the different species in
with Muscle V3.8.155 (Edgar 2004). Then, we built a phylogenetic tree following the HKY85
model for each gene with phyml V3.3 (Guindon et al., 2010). In parallel, we concatenated the
alignment of the 1,055 endosperm genes found in the 22 others Brassicaceae to build a
phylogenetic tree of the endosperm, without branch length, using phyml V3.3 (Guindon et al.,
2010). This tree was used as a proxy of the topology of the endosperm. These steps were
integrated   into   a   pipeline   of   analysis   available   in   GitHub
(https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee). Then, we estimated
the Robinson-Foulds distance between the topology of this tree with the topology of the tree
of each gene obtained previously using the R package TreeDist. Then, to test for significant
differences between the RF distances across the group of genes, we conducted bootstrap
analyses to compare the value on PEGs and MEGs with values obtained from 10,000
resamples of 29 endosperm genes and control genes independently. Similarly, we conducted
bootstrap analyses to compare the values on control genes with values obtained from 10,000
resamples of 100 endosperm genes. The R code for the bootstrap test is available in GitHub
(https://github.com/leveveaudrey/bootstrap).

Variation of the length of branches across Brassicaceae

We investigated whether the imprinted genes presented variation of the branch length
compared to the endosperm and control genes. For each gene, we estimated the length of
branches for each species with the topology of the phylogenetic tree obtained with the 1,055
endosperm genes using phyml V3.3 (Guindon et al., 2010). These steps were integrated into
a   pipeline   of   analysis   available   in   GitHub
(https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee). We extracted the
branches length for each gene with the R package 'ape' version 5.7-1. If the gene was not
present in a species, the value was replaced by "NA". To test for significant differences in the
branch length across the group of genes, we conducted bootstrap analyses to compare the
mean in PEGs and MEGs with means obtained from 10,000 resamples of 29 endosperm

genes and control genes independently. Similarly, we conducted bootstrap analyses to
compare the means on control genes with values obtained from 10,000 resamples of 100
endosperm genes. The R code for the bootstrap test is available in GitHub
(https://github.com/leveveaudrey/bootstrap).

Read mapping and variant calling

We used natural accessions of *A. lyrata* from North America from nine allogamous and nine
autogamous populations of 25 individuals each (Table S6), using available pool-sequencing
data (Willi et al., 2018).

The raw reads from Willi et al (2018) datasets were mapped onto the complete *A. lyrata*
reference genome V1.0.23 (Hu et al., 2011; https://doi.org/10.5281/zenodo.17287897) using
Bowtie2 v2.4.1 (Langmead and Salzberg 2012). File formats were then converted to BAM
using samtools v1.3.1 (Li et al., 2009) and duplicated reads were removed with the
MarkDuplicates program of picard-tools v1.119 (http://broadinstitute.github.io/picard). Biallelic
SNPs in these regions were called using the Genome Analysis Toolkit v. 3.8 (GATK, DePristo
et al., 2011) with the option GVCF and a quality score threshold of 60 using vcftools v0.1.15
(Danecek et al., 2011), to retain only highly reliable SNPs. This stringent cutoff is commonly
used in high-coverage datasets to minimise false positives, which is particularly important for
downstream analyses of selection and co-evolution that are sensitive to SNP-calling errors
(Schlötterer et al., 2014; GATK Best Practices). We excluded sites with less than 15 reads
aligned. Sites covered by at least 15 reads but containing no SNP were considered
monomorphic. The final number of sites in each sample set is summarised in Table S6. The
command lines used for all these steps are available in Github
(https://github.com/leveveaudrey/alignment-and-snp-calling).

Effect of imprinting on global population polymorphism

We decomposed the signal of selection across the imprinted genes into two elementary
statistics: the Tajima's D and the ratio of average polymorphism between the non-synonymous
and synonymous sites π_{NS}/π_S. For the allogamous and autogamous populations, we
compared the values obtained for the imprinted genes with the values obtained across the
endosperm genes and the control genes.

For each gene of each population, we estimated the π, the π_{NS}/π_S for all positions and the
Tajima's D of each gene using a custom python code
(https://github.com/leveveaudrey/POOLSEQ-analysis-of-polymorphism). Briefly, we estimate
the Tajima's D by gene following the classical equation described in Tajima (1989):
D = (π − (S / $\Sigma(1/i)$)) / $\sqrt{e1S + e2S\,(S-1)}$, with e1 = (n+1) /3 * (n+1) - (1/ $\Sigma(1/i)$)/ $\Sigma(1/i)$ and

e2= (2 * (n * n + n + 3)) / (9 * n * (n-1)) - ((n+2) /(Σ(1/i) *n-1)) + (Σ(1/i²) / (Σ(1/i)²)) / (Σ(1/i)²

+ Σ(1/i²)). S is for the number of segregating sites, i for value between 1 and n-1 haplotypes

and n is for the number of haplotypes (number of individuals multiplied by two for diploid).

If a position in the *A. lyrata* genome was covered but not polymorphic, the π was set to 0.

Then, the 0-fold and 4-fold degenerated sites were identified and extracted from the reference

genome and the genome annotation using the script NewAnnotateRef.py (Williamson et al.,

2014;   updated   version   available   in

https://github.com/leveveaudrey/phylogenetic_analyse_brassicacee)*. Because all mutations

at 0-fold degenerated sites alter the sequence of the encoded protein, we assumed that these

mutations are non-synonymous (NS). In contrast, mutations at the 4-fold degenerate sites

never alter the encoded amino acid, thus we assumed that these mutations are synonymous

(S). For the sake of simplicity, we discarded mutations at 2- or 3-fold degenerate sites. Based

on the π estimated for each site, we estimated the mean π_{NS} and π_S, for each gene in each

population, in order to estimate the π_{NS}/π_S. We removed the genes with a π_S to 0. For the

Tajima's D, we applied a SNP-density threshold (≥0.5% polymorphic sites) to ensure reliable

estimates. This density-based criterion avoids gene-length biases inherent to minimum-SNP

thresholds, as genes vary widely in size across categories. We verified that this filtering did

not disproportionately exclude any gene category: the proportions of retained genes were

comparable across endosperm and imprinting genes and across imprinting and control genes

(Table S7).

To test for significant differences across the group of genes, we conducted bootstrap analyses

to compare the mean π, π_{NS}/π_S and Tajima's D of PEGs and MEGs with means obtained from

10,000 resamples of 29 endosperm genes and control genes independently. Similarly, we

conducted bootstrap analyses to compare the values on control genes with values obtained

from 10,000 resamples of 100 endosperm genes. The R code for the bootstrap test is available

in GitHub (https://github.com/leveveaudrey/bootstrap). We performed the test on allogamous

and   autogamous   populations   independently.   Finally,   for   each   statistic   and   group

independently, we tested for the mating system effect by a performing Kruskal-Wallis test as

implemented in R version 4.3.2.

Genes in coevolution

To detect coevolving coding sequences, we used three different methods based on the

following assumptions: coevolution lead to 1) related polymorphic patterns between genes in

populations, 2) related evolutionary rate across species and/or 3) similarity in the topologies

across species (Dutheil and Galtier 2007).

Correlations between evolutionary rates or diversity metrics were used as exploratory
indicators of evolutionary covariance, following approaches such as evolutionary rate
covariation (ERC), which has been widely applied to detect co-functionality or shared selective
pressures (Clark et al., 2012). Because such correlations may reflect genomic or mutational
confounders, we used endosperm-upregulated genes as an empirical baseline for defining
extreme thresholds.

To detect coevolution based on polymorphism in population and on evolutionary rates across
species, we estimated Pearson's linear correlations between each pair of imprinted and
endosperm genes for the mean π and the length of branches on each species. Based on
these correlations matrices, we extracted the correlations between pairs of endosperm genes
to define the thresholds of extreme values in the distributions. Then, all pairs of genes that
included imprinted genes with a correlation lower than the value at 1% of the distributions or
higher than the value at 99% of the distributions were considered as significant. We used
these significant correlations of π and length of branches to build the networks of coevolution.

To detect coevolution based on topology with other genes, we used first the distribution of RF
distances between endosperm genes and the global endosperm topology to estimate the
value obtained in 95% of the distribution. Then, we excluded all the imprinted and endosperm
genes having lower RF distance than this value to remove genes whose topology was
explained by the global topology. Finally, we compared the topology of each remaining
imprinted genes and endosperm genes. All pairs of genes with a null RF distance were
considered as significantly similar and used to build the last coevolution network.

For each network of coevolution, we compared the number of pairs of imprinted genes directly
correlated with the number found between genes in 1,000 random samples of 29 endosperm
genes. Moreover, we tested for gene enrichment on each GO term in each network based on
the occurrences of each term across 1,000 random samples of equivalent size of all the
endosperm and imprinted genes. The complete list of GO term associated to each gene was
downloaded from Gene Ontology online platform for *A. thaliana* (Ashburner et al,. 2000).

Identification of sites in coevolution

We used a model-based approach to detect coevolving sites for each pair of genes showing
a coevolution signal. The pairs of imprinted genes studied were all the pairs directly associated
within at least one network previously mentioned. To use this method, we concatenated each
sequence of pairs of genes for each species. If one species was not found in the other gene,
the sequence associated with this species was removed.

Coevolving sites within each pair of genes were predicted using the clustering method implemented in the CoMap package (version 1.5.2; https://jydu.github.io/comap/; Dutheil and Galtier 2007). The built-in algorithm was then used to correct for multiple testing and to estimate the false discovery rate (FDR). In this study, 1,000 simulations per tested pair were performed, and the groups of sites with an adjusted p value < 1% were considered coevolving. Simple correlation was used as a measure of coevolution. For each comparison pair, we extracted the coevolving groups of substitutions and classified them by size (between 2 and 10).

To test for significant excess of correlations, we compared the results obtained for the pair of imprinted genes found in networks previously mentioned with the results obtained for equivalent number of pair comparisons of imprinted genes not associated in any network, for each cluster of coevolving groups of substitutions of equivalent size by Kruskal-Wallis tests. If the number of correlations is significantly higher than the number expected between two random genes by chance or by alignment errors, we expect a significant increase of these numbers in pairs of associated imprinted genes.

Data availability

The RNAseq data for leaf, root, pistil, and pollen tissues can be found in the NCBI BioProject database under submission number PRJNA1067428. The endosperm RNAseq data is available under submission number GSE76076.

Acknowledgements

This work has been supported by the Czech Science Foundation (GACR 22-20240S granted to C.L-P). Computational resources were supplied by the project 'e-Infrastruktura CZ' (e-INFRA CZ LM2018140) supported by the Ministry of Education, Youth and Sports of the Czech Republic. A.L thanks personally Stella Huynh for our beneficial suggestions and commentaries.

Author contributions

O.I developed and performed the transcriptional analysis. J-Y.D and A.L developed and performed the coevolution analysis. A.L analysed and interpreted the data. A.L wrote the manuscript. All authors edited the manuscript.

Declaration of interests

The authors declare no competing interests.

Bibliography

1) Abdel-Haleem, H., 2007. The Origins of Genome Architecture. J Hered 98, 633–634. https://doi.org/10.1093/jhered/esm073

2) Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. Journal of Molecular Biology 215, 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

3) Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G., 2000. Gene Ontology: tool for the unification of biology. Nat Genet 25, 25–29. https://doi.org/10.1038/75556

4) Bemer, M., Heijmans, K., Airoldi, C., Davies, B., Angenent, G.C., 2010. An Atlas of Type I MADS Box Gene Expression during Female Gametophyte and Seed Development in Arabidopsis. Plant Physiology 154, 287–300. https://doi.org/10.1104/pp.110.160770

5) Bernardi, G., 2000. Isochores and the evolutionary genomics of vertebrates. Gene 241, 3–17. https://doi.org/10.1016/s0378-1119(99)00485-0

6) Botet, R., and Joost J. B. Keurentjes., 2020. The Role of Transcriptional Regulation in Hybrid Vigor. Frontiers in Plant Science 11 (April). https://doi.org/10.3389/fpls.2020.00410

7) Brandvain, Y., Haig, D., 2005. Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. Am Nat 166, 330–338. https://doi.org/10.1086/432036

8) Bryant, N., Lloyd, J., Sweeney, C., Myouga, F., Meinke, D., 2010. Identification of Nuclear Genes Encoding Chloroplast-Localized Proteins Required for Embryo Development in Arabidopsis. Plant Physiology 155, 1678. https://doi.org/10.1104/pp.110.168120

9) Clark, N.L., Alani, E., Aquadro C.F., 2012. Evolutionary Rate Covariation Reveals Shared Functionality and Coexpression of Genes. Genome Research 22 (4): 714–20. https://doi.org/10.1101/gr.132647.111

10) Cvijović, I., Good, B.H., Desai, M.M., 2018. The Effect of Strong Purifying Selection on Genetic Diversity. Genetics 209 (4): 1235–78. https://doi.org/10.1534/genetics.118.301058

11) Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., 1000 Genomes Project Analysis Group, 2011. The variant call format and VCFtools. Bioinformatics 27, 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

12) Day, T., Bondurianský, R., 2004. Intralocus sexual conflict can drive the evolution of genomic imprinting. Genetics 167, 1537. https://doi.org/10.1534/genetics.103.026211

13) Day, R.C., Herridge, R.P., Ambrose, B.A., Macknight, R.C., 2008. Transcriptome Analysis of Proliferating Arabidopsis Endosperm Reveals Biological Implications for the Control of Syncytial Division, Cytokinin Signaling, and Gene Expression Regulation. Plant Physiology 148, 1964–1984. https://doi.org/10.1104/pp.108.128108

14) Dekkers, B.J.W., Pearce, S., van Bolderen-Veldkamp, R.P., Marshall, A., Widera, P., Gilbert, J., Drost, H.-G., Bassel, G.W., Müller, K., King, J.R., Wood, A.T.A., Grosse, I.,

Quint, M., Krasnogor, N., Leubner-Metzger, G., Holdsworth, M.J., Bentsink, L., 2013. Transcriptional Dynamics of Two Seed Compartments with Opposing Roles in Arabidopsis Seed Germination. Plant Physiology 163, 205–215. https://doi.org/10.1104/pp.113.223511

15) DePristo, M.A., Banks, E., Poplin, R.E., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., del Angel, G., Rivas, M.A., Hanna, M., McKenna, A., Fennell, T.J., Kernytsky, A.M., Sivachenko, A.Y., Cibulskis, K., Gabriel, S.B., Altshuler, D., Daly, M.J., 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet 43, 491–498. https://doi.org/10.1038/ng.806

16) Dutheil, J. ,Galtier, N., 2007. Coevolution within and between Genes. https://doi.org/10.1159/000107599

17) Eaux, A.J., Bilinski, P., Eichten, S.R., Vaughn, M.W., Ross-Ibarra, J., Gehring, M., Reprinte, N.M., 2013. Comprehensive analysis of imprinted genes in maize reveals allelic variation for imprinting and limited conservation with other species. Proceedings of the National Academy of Sciences 110, 19639–19644. https://doi.org/10.1073/pnas.1309182110

18) Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res 32, 1792–1797. https://doi.org/10.1093/nar/gkh340

19) Edwards, S.V., 2009. Natural selection and phylogenetic analysis. Proceedings of the National Academy of Sciences 106, 8799–8800. https://doi.org/10.1073/pnas.0904103106

20) Ferretti, L., Ramos-Onsins, S.E., Pérez-Enciso, M., 2013. Population genomics from pool sequencing. Mol Ecol 22, 5561–5576. https://doi.org/10.1111/mec.12522

21) Florez-Rueda, A.M., Paris, M., Schmidt, A., Widmer, A., Grossniklaus, U., Städler, T., 2016. Genomic Imprinting in the Endosperm Is Systematically Perturbed in Abortive Hybrid Tomato Seeds. Mol Biol Evol 33, 2935–2946. https://doi.org/10.1093/molbev/msw175

22) Flores-Vergara, M.A., Oneal, E., Costa, M., Villarino, G., Roberts, C., De Luis Balaguer, M.A., Coimbra, S., Willis, J., Franks, R.G., 2020. Developmental Analysis of Mimulus Seed Transcriptomes Reveals Functional Gene Expression Clusters and Four Imprinted, Endosperm-Expressed Genes. Front. Plant Sci. 11. https://doi.org/10.3389/fpls.2020.00132

23) Foxe, J.P., Stift, M., Tedder, A., Haudry, A., Wright, S.I., Mable, B.K., 2010. Reconstructing Origins of Loss of Self-Incompatibility and Selfing in North American Arabidopsis Lyrata: A Population Genetic Context. Evolution 64, 3495–3510. https://doi.org/10.1111/j.1558-5646.2010.01094.x

24) Fraser, H.B., Hirsh, A.E., Steinmetz, L.M., Scharfe, C., Feldman, M.W., 2002. Evolutionary Rate in the Protein Interaction Network. Science 296, 750–752. https://doi.org/10.1126/science.1068696

25) Futschik, A., Schlötterer, C., 2010. The Next Generation of Molecular Markers From Massively Parallel Sequencing of Pooled DNA Samples. Genetics 186, 207–218. https://doi.org/10.1534/genetics.110.114397

26) Galtier, N., Rousselle, M., 2020. How Much Does Ne Vary Among Species? Genetics 216, 559–572. https://doi.org/10.1534/genetics.120.303622

27) Garro, M., Greco, E., Vannay, G.J., Leonova, A., Bruno, L., Capella, M., 2025. Non-CG DNA methylation represses SDC expression to modulate hypocotyl elongation during thermormorphogenesis in Arabidopsis. J Exp Bot 76, 2517–2534. https://doi.org/10.1093/jxb/eraf105

28) Gaudinier, A., Rodriguez-Medina, J., Zhang, L., Olson, A., Liseron-Monfils, C., Bågman, A.-M., Foret, J., Abbitt, S., Tang, M., Li, B., Runcie, D.E., Kliebenstein, D.J., Shen, B., Frank, M.J., Ware, D., Brady, S.M., 2018. Transcriptional regulation of nitrogen-associated metabolism and growth. Nature 563, 259–264. https://doi.org/10.1038/s41586-018-0656-3

29) Gehring, M., Missirian, V., Henikoff, S., 2011. Genomic Analysis of Parent-of-Origin Allelic Expression in Arabidopsis thaliana Seeds. PLOS ONE 6, e23687. https://doi.org/10.1371/journal.pone.0023687

30) Goodstein, D.M., Shu, S., Howson, R., Neupane, R., Hayes, R.D., Fazo, J., Mitros, T., Dirks, W., Hellsten, U., Putnam, N., Rokhsar, D.S., 2012. Phytozome: a comparative platform for green plant genomics. Nucleic Acids Res 40, D1178-1186. https://doi.org/10.1093/nar/gkr944

31) Green, A.G., Elhabashy, H., Brock, K.P., Maddamsetti, R., Kohlbacher, O., Marks, D.S., 2021. Large-scale discovery of protein interactions at residue resolution using co-evolution calculated from genomic sequences. Nat Commun 12, 1396. https://doi.org/10.1038/s41467-021-21636-z

32) Grishkevich, V., Yanai, I., 2014. Gene length and expression level shape genomic novelties. Genome Res 24, 1497–1503. https://doi.org/10.1101/gr.169722.113

33) Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59, 307–321. https://doi.org/10.1093/sysbio/syq010

34) Haig, D., Westoby, M., 1989. Parent-Specific Gene Expression and the Triploid Endosperm. The American Naturalist 134, 147–155. https://doi.org/10.1086/284971

35) Haig, D., Westoby, M., 1991. Genomic Imprinting in Endosperm: Its Effect on Seed Development in Crosses between Species, and between Different Ploidies of the Same Species, and Its Implications for the Evolution of Apomixis. Philosophical Transactions: Biological Sciences 333, 1–13. https://doi.org/10.1098/rstb.1991.0057

36) Haig, D., 1997. Parental antagonism, relatedness asymmetries, and genomic imprinting. Proceedings of the Royal Society of London. Series B: Biological Sciences 264, 1657–1662. https://doi.org/10.1098/rspb.1997.0230

37) Haig, D., 2000. The Kinship Theory of Genomic Imprinting. Annual Review of Ecology and Systematics 31, 9–32. https://doi.org/10.1146/annurev.ecolsys.31.1.9

38) Haig, D., 2014. Coadaptation and conflict, misconception and muddle, in the evolution of genomic imprinting. Heredity 113, 96–103. https://doi.org/10.1038/hdy.2013.97

39) Halg, D., Graham, C., 1991. Genomic imprinting and the strange case of the insulin-like growth factor II receptor. Cell 64, 1045–1046. https://doi.org/10.1016/0092-8674(91)90256-X

40) Hatorangan, M.R., Laenen, B., Steige, K.A., Slotte, T., Köhler, C., 2016. Rapid Evolution of Genomic Imprinting in Two Species of the Brassicaceae. The Plant Cell 28, 1815–1827. https://doi.org/10.1105/tpc.16.00304

41) Hehenberger, E., Kradolfer, D., Köhler, C., 2012. Endosperm cellularization defines an important developmental transition for embryo development. Development 139, 2031–2039. https://doi.org/10.1242/dev.077057

42) Hojsgaard, D., Schartl, M., 2021. Skipping sex: A nonrecombinant genomic assemblage of complementary reproductive modules. BioEssays 43, 2000111. https://doi.org/10.1002/bies.202000111

43) Hough, J., Immler, S., Barrett, S.C.H., Otto, S.P., 2013. Evolutionarily stable sex ratios and mutation load. Evolution 67, 1915–1925. https://doi.org/10.1111/evo.12066

44) Hsieh, T.-F., Shin, J., Uzawa, R., Silva, P., Cohen, S., Bauer, M.J., Hashimoto, M., Kirkbride, R.C., Harada, J.J., Zilberman, D., Fischer, R.L., 2011. Regulation of imprinted gene expression in Arabidopsis endosperm. Proceedings of the National Academy of Sciences 108, 1755–1762. https://doi.org/10.1073/pnas.1019273108

45) Hu, T.T., Pattyn, P., Bakker, E.G., Cao, J., Cheng, J.-F., Clark, R.M., Fahlgren, N., Fawcett, J.A., Grimwood, J., Gundlach, H., Haberer, G., Hollister, J.D., Ossowski, S., Ottilar, R.P., Salamov, A.A., Schneeberger, K., Spannagl, M., Wang, X., Yang, L., Nasrallah, M.E., Bergelson, J., Carrington, J.C., Gaut, B.S., Schmutz, J., Mayer, K.F.X., Van de Peer, Y., Grigoriev, I.V., Nordborg, M., Weigel, D., Guo, Y.-L., 2011. The Arabidopsis lyrata genome sequence and the basis of rapid genome size change. Nat Genet 43, 476–481. https://doi.org/10.1038/ng.807

46) İltaş, Ö., Svitok, M., Cornille, A., Schmickl, R., Lafon Placette, C., 2021. Early evolution of reproductive isolation: A case of weak inbreeder/strong outbreeder leads to an intraspecific hybridization barrier in Arabidopsis lyrata. Evol 75, 1466–1476. https://doi.org/10.1111/evo.14240

47) İltaş, Ö., Čertner, M., Placette, C.L., 2024. Natural variation in sexual traits and gene expression between selfing and outcrossing Arabidopsis lyrata suggests sexual selection at work. Plant and Cell Physiology pcae090. https://doi.org/10.1093/pcp/pcae090

48) Jiang, H., Guo, D., Ye, J., Gao, Y., Liu, H., Wang, Y., Xue, M., Yan, Q., Chen, J., Duan, L., Li, G., Li, X., Xie, L., 2021. Genome-wide analysis of genomic imprinting in the endosperm and allelic variation in flax. The Plant Journal 107, 1697–1710. https://doi.org/10.1111/tpj.15411

49) Kasimatis, K.R., Nelson, T.C., Phillips, P.C., 2017. Genomic Signatures of Sexual Conflict. Journal of Heredity 108, 780–790. https://doi.org/10.1093/jhered/esx080

50) Kinoshita, T., 2007. Reproductive barrier and genomic imprinting in the endosperm of flowering plants. Genes & Genetic Systems 82, 177–186. https://doi.org/10.1266/ggs.82.177

51) Kinser, T.J., Smith, R.D., Lawrence, A.H., Cooley, A.M., Vallejo-Marín, M., Conradi Smith, G.D., Puzey, J.R., 2021. Endosperm-based incompatibilities in hybrid monkeyflowers. Plant Cell 33, 2235–2257. https://doi.org/10.1093/plcell/koab117

52) Köhler, C., Scheid, O.M., Erilova, A., 2010. The impact of the triploid block on the origin and evolution of polyploid plants. Trends in Genetics 26, 142–148. https://doi.org/10.1016/j.tig.2009.12.006

53) Koonin, E.V., 2015. Why the Central Dogma: on the nature of the great biological exclusion principle. Biology Direct 10, 52. https://doi.org/10.1186/s13062-015-0084-3

54) Klosinska, M., Picard, C.L., Gehring, M., 2016. Conserved imprinting associated with unique epigenetic signatures in the Arabidopsis genus. Nature Plants 2, 1–8. https://doi.org/10.1038/nplants.2016.145

55) Kremer, A., Chen, J., Lascoux, M., 2025. 'Chimes of resilience': what makes forest trees genetically resilient? New Phytologist 246, 1934–1951. https://doi.org/10.1111/nph.70108

56) Laenen, B., Tedder, A., Nowak, M.D., Toräng, P., Wunder, J., Wötzel, S., Steige, K.A., Kourmpetis, Y., Odong, T., Drouzas, A.D., Bink, M.C.A.M., Ågren, J., Coupland, G., Slotte, T., 2018. Demography and mating system shape the genome-wide impact of purifying selection in Arabis alpina. Proc Natl Acad Sci U S A 115, 816–821. https://doi.org/10.1073/pnas.1707492115

57) Lafon-Placette, C., Hatorangan, M.R., Steige, K.A., Cornille, A., Lascoux, M., Slotte, T., Köhler, C., 2018. Paternally expressed imprinted genes associate with hybridization barriers in Capsella. Nature Plants 4, 352–357. https://doi.org/10.1038/s41477-018-0161-6

58) Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods 9, 357–359. https://doi.org/10.1038/nmeth.1923

59) Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

60) Llaurens, V., Whibley, A., Joron, M., 2017. Genetic architecture and balancing selection: the life and death of differentiated variants. Molecular Ecology 26, 2430–2448. https://doi.org/10.1111/mec.14051

61) Lopes, I., Altab, G., Raina, P., de Magalhães, J.P., 2021. Gene Size Matters: An Analysis of Gene Length in the Human Genome. Front Genet 12, 559998. https://doi.org/10.3389/fgene.2021.559998

62) Lynch, M., 2010. Evolution of the mutation rate. Trends Genet 26, 345–352. https://doi.org/10.1016/j.tig.2010.05.003

63) Lynch, M., Bost, D., Wilson, S., Maruki, T., Harrison, S., 2014. Population-Genetic Inference from Pooled-Sequencing Data. Genome Biol Evol 6, 1210–1218. https://doi.org/10.1093/gbe/evu085

64) Maheshwari, S., Barbash, D.A., 2011. The Genetics of Hybrid Incompatibilities. Annu. Rev. Genet. 45, 331–355. https://doi.org/10.1146/annurev-genet-110410-132514

65) Massey, S.E., 2013. Proteome size as the major factor determining mutation rates. Proceedings of the National Academy of Sciences 110, E858–E859. https://doi.org/10.1073/pnas.1219306110

66) McVean, G.T., Hurst, L.D., 1997. Molecular evolution of imprinted genes: no evidence for antagonistic coevolution. Proceedings of the Royal Society of London. Series B: Biological Sciences 264, 739–746. https://doi.org/10.1098/rspb.1997.0105

67) Mills, W., Moore, T., 2004. Polyandry, Life-History Trade-Offs and the Evolution of Imprinting at Mendelian Loci. Genetics 168, 2317–2327. https://doi.org/10.1534/genetics.104.030098

68) Miyake, T., Takebayashi, N., Wolf, D.E., 2009. Possible Diversifying Selection in the Imprinted Gene, MEDEA, in Arabidopsis. Molecular Biology and Evolution 26, 843–857. https://doi.org/10.1093/molbev/msp001

69) Moore, T., Haig, D., 1991. Genomic imprinting in mammalian development: a parental
tug-of-war. Trends Genet 7, 45–49. https://doi.org/10.1016/0168-9525(91)90230-N

70) Noguero, M., Atif, R.M., Ochatt, S., Thompson, R.D., 2013. The role of the DNA-
binding One Zinc Finger (DOF) transcription factor family in plants. Plant Science 209,
32–45. https://doi.org/10.1016/j.plantsci.2013.03.016

71) Onosato, H., Fujimoto, G., Higami, T., Sakamoto, T., Yamada, A., Suzuki, T., Ozawa,
R., Matsunaga, S., Seki, M., Ueda, M., Sako, K., Galis, I., Arimura, G., 2022. Sustained
defense response via volatile signaling and its epigenetic transcriptional regulation.
Plant Physiol 189, 922–933. https://doi.org/10.1093/plphys/kiac077

72) Otto, S.P., Scott, M.F., Immler, S., 2015. Evolution of haploid selection in
predominantly diploid organisms. Proceedings of the National Academy of Sciences
112, 15952–15957. https://doi.org/10.1073/pnas.1512004112

73) Padilla-García, N., Le Veve, A., Čermák, V., İltaş, Ö., Contreras-Garrido, A., Legrand,
S., Aury, J.-M., Horvath, R., Lafon Placette, C., 2025. The Demographic History of
Populations and Genomic Imprinting have Shaped the Transposon Patterns in
Arabidopsis lyrata. Mol Biol Evol 42, msaf093. https://doi.org/10.1093/molbev/msaf093

74) Papdi, C., Ábrahám, E., Joseph, M.P., Popescu, C., Koncz, C., Szabados, L., 2008.
Functional Identification of Arabidopsis Stress Regulatory Genes Using the Controlled
cDNA Overexpression System. Plant Physiol 147, 528–542.
https://doi.org/10.1104/pp.108.116897

75) Patten, M.M., Ross, L., Curley, J.P., Queller, D.C., Bonduriansky, R., Wolf, J.B., 2014.
The evolution of genomic imprinting: theories, predictions and empirical tests. Heredity
113, 119–128. https://doi.org/10.1038/hdy.2014.29

76) Pignatta, D., Erdmann, R.M., Scheer, E., Picard, C.L., Bell, G.W., Gehring, M., 2014.
Natural epigenetic polymorphisms lead to intraspecific variation in Arabidopsis gene
imprinting. eLife 3, e03198. https://doi.org/10.7554/eLife.03198

77) Pignatta, D., Novitzky, K., Satyaki, P.R.V., Gehring, M., 2018. A variably imprinted
epiallele impacts seed development. PLOS Genetics 14, e1007469.
https://doi.org/10.1371/journal.pgen.1007469

78) Presgraves, D.C., 2010. The molecular evolutionary basis of species formation. Nat
Rev Genet 11, 175–180. https://doi.org/10.1038/nrg2718

79) Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics 26, 841–842.
https://doi.org/10.1093/bioinformatics/btq033

80) Raissig, M.T., Baroux, C., Grossniklaus, U., 2011. Regulation and Flexibility of
Genomic Imprinting during Seed Development. The Plant Cell 23, 16–26.
https://doi.org/10.1105/tpc.110.081018

81) Ross-Ibarra, J., Wright, S.I., Foxe, J.P., Kawabe, A., DeRose-Wilson, L., Gos, G.,
Charlesworth, D., Gaut, B.S., 2008. Patterns of Polymorphism and Demographic
History in Natural Populations of Arabidopsis lyrata. PLOS ONE 3, e2411.
https://doi.org/10.1371/journal.pone.0002411

82) Roth, M., Florez-Rueda, A.M., Paris, M., Städler, T., 2018. Wild tomato endosperm
transcriptomes reveal common roles of genomic imprinting in both nuclear and cellular
endosperm. The Plant Journal 95, 1084–1101. https://doi.org/10.1111/tpj.14012

83) Sahoo, S., Das, S.S., Rakshit, R., 2019. Codon usage pattern and predicted gene expression in Arabidopsis thaliana. Gene, Articles initially published in Gene: X 1-4, 2019 721, 100012. https://doi.org/10.1016/j.gene.2019.100012

84) Sandstedt, G.D., Sweigart, A.L., 2022. Developmental evidence for parental conflict in driving Mimulus species barriers. New Phytol 236, 1545–1557. https://doi.org/10.1111/nph.18438

85) Scott, R.J., Spielman, M., Bailey, J., Dickinson, H.G., 1998. Parent-of-origin effects on seed development in Arabidopsis thaliana. Development 125, 3329–3341. https://doi.org/10.1242/dev.125.17.3329

86) Seehausen, O., Butlin, R.K., Keller, I., Wagner, C.E., Boughman, J.W., Hohenlohe, P.A., Peichel, C.L., Saetre, G.-P., Bank, C., Brännström, Å., Brelsford, A., Clarkson, C.S., Eroukhmanoff, F., Feder, J.L., Fischer, M.C., Foote, A.D., Franchini, P., Jiggins, C.D., Jones, F.C., Lindholm, A.K., Lucek, K., Maan, M.E., Marques, D.A., Martin, S.H., Matthews, B., Meier, J.I., Möst, M., Nachman, M.W., Nonaka, E., Rennison, D.J., Schwarzer, J., Watson, E.T., Westram, A.M., Widmer, A., 2014. Genomics and the origin of species. Nat Rev Genet 15, 176–192. https://doi.org/10.1038/nrg3644

87) Sharp, P.M., Li, W.H., 1987. The codon Adaptation Index--a measure of directional synonymous codon usage bias, and its potential applications. Nucleic Acids Res 15, 1281–1295. https://doi.org/10.1093/nar/15.3.1281

88) Soliman, H.K., Coughlan, J.M., 2024. United by conflict: Convergent signatures of parental conflict in angiosperms and placental mammals. J Hered 115, 625–642. https://doi.org/10.1093/jhered/esae009

89) Spencer, H. G. 2000. Population Genetics and Evolution of Genomic Imprinting. Annual Review of Genetics 34: 457–77. https://doi.org/10.1146/annurev.genet.34.1.457.

90) Spencer, H.G., Clark, A.G., 2014. Non-conflict theories for the evolution of genomic imprinting. Heredity 113, 112–118. https://doi.org/10.1038/hdy.2013.129

91) Spillane, C., Schmid, K.J., Laoueillé-Duprat, S., Pien, S., Escobar-Restrepo, J.-M., Baroux, C., Gagliardini, V., Page, D.R., Wolfe, K.H., Grossniklaus, U., 2007. Positive darwinian selection at the imprinted MEDEA locus in plants. Nature 448, 349–352. https://doi.org/10.1038/nature05984

92) Sujeeth, N., Mehterov, N., Gupta, S., Qureshi, M.K., Fischer, A., Proost, S., Omidbakhshfard, M.A., Obata, T., Benina, M., Staykov, N., Balazadeh, S., Walther, D., Fernie, A.R., Mueller-Roeber, B., Hille, J., Gechev, T.S., 2020. A novel seed plants gene regulates oxidative stress tolerance in Arabidopsis thaliana. Cell. Mol. Life Sci. 77, 705–718. https://doi.org/10.1007/s00018-019-03202-5

93) Sylvestre, F., Mérot, C., Normandeau, E., Bernatchez, L., 2023. Searching for intralocus sexual conflicts in the three-spined stickleback (Gasterosteus aculeatus) genome. Evolution 77, 1667–1681. https://doi.org/10.1093/evolut/qpad075

94) Tuteja, R., McKeown, P.C., Ryan, P., Morgan, C.C., Donoghue, M.T.A., Downing, T., O'Connell, M.J., Spillane, C., 2019. Paternally Expressed Imprinted Genes under Positive Darwinian Selection in Arabidopsis thaliana. Molecular Biology and Evolution 36, 1239–1253. https://doi.org/10.1093/molbev/msz063

95) Willi, Y., 2013. The Battle of the Sexes over Seed Size: Support for Both Kinship Genomic Imprinting and Interlocus Contest Evolution. The American Naturalist 181, 787–798. https://doi.org/10.1086/670196

96) Willi, Y., Fracassetti, M., Zoller, S., Van Buskirk, J., 2018. Accumulation of Mutational Load at the Edges of a Species Range. Molecular Biology and Evolution 35, 781–791. https://doi.org/10.1093/molbev/msy003

97) Wilkins, J.F., Haig, D., 2001. Genomic imprinting of two antagonistic loci. Proceedings of the Royal Society of London. Series B: Biological Sciences 268, 1861–1867. https://doi.org/10.1098/rspb.2001.1651

98) Wilkins, J.F., Haig, D., 2003. What Good Is Genomic Imprinting: The Function of Parent-Specific Gene Expression. Nature Reviews Genetics 4 (5): 359–68. https://doi.org/10.1038/nrg1062.

99) Wilkins, J.F., 2011. Genomic imprinting and conflict-induced decanalization. Evolution 65, 537–553. https://doi.org/10.1111/j.1558-5646.2010.01147.x

100) Williamson, R.J., Josephs, E.B., Platts, A.E., Hazzouri, K.M., Haudry, A., Blanchette, M., Wright, S.I., 2014. Evidence for Widespread Positive and Negative Selection in Coding and Conserved Noncoding Regions of Capsella grandiflora. PLOS Genetics 10, e1004622. https://doi.org/10.1371/journal.pgen.1004622

101) Wolff, P., Weinhofer, I., Seguin, J., Roszak, P., Beisel, C., Donoghue, M.T.A., Spillane, C., Nordborg, M., Rehmsmeier, M., Köhler, C., 2011. High-Resolution Analysis of Parent-of-Origin Allelic Expression in the Arabidopsis Endosperm. PLOS Genetics 7, e1002126. https://doi.org/10.1371/journal.pgen.1002126

102) Wolff, P., Jiang, H., Wang, G., Santos-González, J., Köhler, C., 2015. Paternally expressed imprinted genes establish postzygotic hybridization barriers in Arabidopsis thaliana. eLife 4, e10074. https://doi.org/10.7554/eLife.10074

103) Wuest, S.E., Vijverberg, K., Schmidt, A., Weiss, M., Gheyselinck, J., Lohr, M., Wellmer, F., Rahnenführer, J., Mering, C. von, Grossniklaus, U., 2010. Arabidopsis Female Gametophyte Gene Expression Map Reveals Similarities between Plant and Animal Gametes. Current Biology 20, 506–512. https://doi.org/10.1016/j.cub.2010.01.051

104) Yang, G., Feng, M., Yu, K., Cui, G., Zhou, Y., Sun, L., Gao, L., Zhang, Y., Peng, H., Yao, Y., Hu, Z., Rossi, V., De Smet, I., Ni, Z., Sun, Q., Xin, M., 2023. Paternally imprinted LATE-FLOWERING2 transcription factor contributes to paternal-excess interploidy hybridization barriers in wheat. Journal of Integrative Plant Biology 65, 2587–2603. https://doi.org/10.1111/jipb.13574

105) Zhang, S., Wang, D., Zhang, H., Skaggs, M.I., Lloyd, A., Ran, D., An, L., Schumaker, K.S., Drews, G.N., Yadegari, R., 2018. FERTILIZATION-INDEPENDENT SEED-Polycomb Repressive Complex 2 Plays a Dual Role in Regulating Type I MADS-Box Genes in Early Endosperm Development. Plant Physiology 177, 285–299. https://doi.org/10.1104/pp.17.00534

106) Zhang, B., Li, C., Li, Y., Yu, H., 2020. Mobile TERMINAL FLOWER1 determines seed size in Arabidopsis. Nat Plants 6, 1146–1157. https://doi.org/10.1038/s41477-020-0749-5

107) Zou, H.-F., Zhang, Y.-Q., Wei, W., Chen, H.-W., Song, Q.-X., Liu, Y.-F., Zhao, M.-Y., Wang, F., Zhang, B.-C., Lin, Q., Zhang, W.-K., Ma, B., Zhou, Y.-H., Zhang, J.-S., Chen, S.-Y., 2012. The transcription factor AtDOF4.2 regulates shoot branching and seed coat formation in Arabidopsis. Biochemical Journal 449, 373–388. https://doi.org/10.1042/BJ20110060

Figure S1: phylogenetic tree obtained by HKY85 model based on the 1055 endosperm genes found in all species after 1000 bootstrap. The percentage of trees in which the associated haplotypes clustered together is shown next to the branches. The tree is drawn to scale, with branch lengths measured in mean number of substitutions per site.