Activation and regulation of a p38-MAPK by its downstream MAPKAP kinase
through feedback phosphorylation and LLPS-driven condensate formation.

Pranita Ray[a], Chakshu Mangal[b], Pallavi Rao T[c,d], Mintu Nandi[e,f], Afreen Haque[a], Vibha
Viswanath[b], Samrat Mitra[a], Swasti Raychaudhuri[c,d], Anindya Ghosh-Roy[b] and Smarajit
Polley[a,1]

a Department of Biological Sciences, Bose Institute, Sector V, Bidhan Nagar, Kolkata
700091, India; b Department of Cellular and Molecular Neuroscience, BRIC-National Brain
Research Centre, Manesar 122052, India; c CSIR-Centre for Cellular and Molecular Biology,
Uppal Road, Hyderabad 500007, India; d Academy of Scientific and Innovative Research
(AcSIR), Ghaziabad- 201002, India; e Department of Chemistry, Indian Institute of
Engineering Science and Technology, Shibpur, Howrah 711103, India; current address:
Universal Biology Institute, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-
0033, Japan.

1 corresponding author: smarajit.polley@gmail.com, spolley@jcbose.ac.in

Abstract:

MAP kinases (MAPKs) represent a class of evolutionarily conserved stress and
extracellular stimuli responsive signaling molecules. p38 group of MAPKs known to
be activated by dual-specific upstream MAPK kinases and also by
autophosphorylation. They activate MAPK activated protein kinases (MAPKAPKs) in a
context dependent manner by specific phosphorylation, and together, they play
crucial biological roles. One such pair is PMK3, p38α-MAPK and its cognate
MAPKAPK, MAK2, downstream of DLK1 (MAPK kinase kinase) and MKK4 (MAPK
kinase) in *C. elegans*. They are implicated in axonal regeneration, degeneration and
synaptic pruning in response to neuronal injury. Here, we report that PMK3 and MAK2
engage in a feedback relationship, leading to phosphorylation-mediated activation of
both kinases. Interaction of PMK3 with MAK2 through the canonical docking site
leads to negligible *de novo* autophosphorylation of PMK3. Phosphorylation of both
Thr and Tyr residues on the TxY-motif necessary for the full activation of PMK3
requires catalytic activity of MAK2 as confirmed by mass spectrometry. Distribution
of phosphorylation sites on either kinase when incubated together, and presence of
long intrinsically disordered regions in each kinase indicate that PMK3:MAK2

complex is conformationally plastic in nature. MAK2 increases bioavailability of aggregation-prone PMK3 by virtue of its ability to form LLPS-driven condensates *in vitro*, wherein they retain enzymatic activities and phosphorylate each other. Furthermore, experiments with transgenic *C. elegans* reveal that MAK2 controls stability of PMK3 and localization of PMK3 puncta in touch neuron. Our observations offer an unreported activation mechanism of a p38-MAPK by its downstream MAPKAPK.

Significance Statement:

MAPK kinases are evolutionarily conserved, and are key players in stress response, cell survival, differentiation, metabolic processes and neuronal response to injury. MAPK pathways are primarily activated through unidirectional flow of phosphorylation-signal MAP3K to MAP2K to MAPK, and in some cases to a downstream MAPKAPK. We found that a *C. elegans* MAPK (PMK3) can also be activated by its downstream MAPKAPK (MAK2) through feedback phosphorylation that ensures robust activation of PMK3 by MAK2 without requiring the MAP2K. Furthermore, MAK2 increases bioavailability of activation-competent and active PMK3 by preventing its aggregation through LLPS-driven condensate formation *in vitro* and in the worm neuron. This feedback relationship might ensure rapid activation of such MAPK pathways in response to nervous system injury or stress.

Introduction:

Protein kinases are implicated in almost, if not all, signaling events in eukaryotes (1). Mitogen activated protein kinases (MAPKs) are activated by a variety of extracellular stimuli that regulate a plethora of cellular processes including but not limited to gene expression, cell cycle, cell survival, infection, inflammation, apoptosis, differentiation and metabolism (2-6). They are one of the most ancient signal transduction molecules found throughout evolution. Multiple MAPK pathways are found to operate in concert in eukaryotes. A canonical MAPK module consists of two upstream kinases, MAPK kinase (MAPKK or MAP2K) and MAPKK kinase (MAPKKK or MAP3K)(7). In response to extracellular stimuli, a MAP3K activates its cognate MAP2K that in turn activates the downstream MAPK – making it a three-kinase signaling module where the signal transmits from the MAP3K to MAPK via the MAP2K. Activation of p38α by autophosphorylation has also been documented (8) apart from the canonical mode of MAPK activation by MAP2K, suggesting multiple modes of MAPK-activation in play. MAPKs are Ser/Thr kinases that phosphorylate different kinds of protein substrates in cells including transcription factors and specific protein kinases called the MAPK activated protein kinases (MAPKAPKs) (9).

p38 group of MAPKs are perhaps the most widely implicated and most well-studied pathways among the four distinct subgroups of the MAPK family (2, 10). In mammals, there are four isoforms of p38: α, β, γ and δ. Among these p38α is the most well studied one. Among many substrates of p38α, MAPKAPK2 or MK2 have been implicated in many cellular processes that phosphorylate substrates including small heat shock protein 27 (Hsp27), cAMP response element-binding protein (CREB), lymphocyte-specific protein 1 (LSP1), transcription factor ATF1 etc. (11). p38 MAPKs are present both in the cytoplasm and nuclei of cells and are sequestered mostly in the nuclei in certain stress conditions. Nucleocytoplasmic shuttling of the p38 MAPKs is also regulated by their association with the MAPKAPKs (12).

C. elegans genome consists of three genes, *pmk1, pmk2* and *pmk3* that code for three p38 MAPK isoforms: PMK1, PMK2 and PMK3, respectively. These three genes constitute an operon with a polarity from *pmk-1* to *pmk-3*, and are expressed under the influence of a single promoter (13). Among these three p38 MAPK isoforms, PMK3 has been implicated in axonal regeneration acting as the MAPK downstream of MAP2K, MKK4 that is activated by the MAP3K, DLK1 in both *C. elegans* and other organisms (14-17) This signaling mechanism is also implicated in synaptic maturation in multiple organisms, which is under the tight regulation of E3 ubiquitin ligase (18-20). In the axonal regeneration pathway involving DLK1, the terminal kinase however is not the MAPK but a cognate MAPKAPK of PMK3 known as MAK2 akin to human MK2 or MK3 (Fig. 1A). MAK2 is

involved in the maintenance of the neuronal synapses and axon morphology. It has also
been shown that MAK2 regulates touch neuron axon regeneration by modulating *cebp-1*
mRNA stability in response to axon injury in nematode (21). A search for the protein-protein
interaction networks related to PMK3 and MAK2 on the STRING database revealed strong
interactions between the two proteins (Fig. 1B), both in the full STRING network as well as
physical subnetwork (22). We asked the question, whether activation, regulation and mutual
relationship of p38α and its MAPKAPK established in mammalian systems also holds true in
the PMK3:MAK2 system, which is involved in multiple cellular responses such as axon
regeneration (23), degeneration (24), and synaptic pruning (25) following neuronal injury.
We used the nematode specific counterparts of MAPK (PMK3) and MAPKAPK (MAK2) as
our model systems. Our biochemical, computational, mass spectrometric and organismal
analyses revealed that PMK3 participates in a hitherto unknown feedback relationship with
MAK2. PMK3 activates its downstream kinase MAK2 that leads to the double-
phosphorylation of the activating TxY motif in PMK3 leading to its full activation. We also
found that, MAK2 but not PMK3 forms condensates through liquid-liquid phase separation
(LLPS) *in vitro* and prevents aggregation of PMK3. Presence of MAK2 increases level of
PMK3 in neurons under stress conditions in *C. elegans*. Further, the feedback-
phosphorylation relationship remains functional in the condensate state. These observations
point towards an unknown regulatory aspect of a MAP kinase (PMK3), wherein it is
regulated by its cognate downstream kinase (MAK2), not only by means of its catalytic
activity to chemically modify (activating phosphorylation) but also through change of physico-
chemical microenvironment. Overall, our study revealed a new mechanistic understanding of
the regulation of the injury sensing MAPK, PMK3 via its downstream kinase MAPKAPK,
MAK2.
119
Results:
In vitro reconstitution of a MAPK-MAPKAPK module implicated in axonal regeneration
in C. elegans:
PMK3 (Uniprot id: O44514, M.W. 54.9kDa) is a p38 MAPK in *C. elegans*. It
participates in the DLK1-dependent signaling cascade in axonal regeneration pathway
downstream of the MAP2K, MKK4 (KEGG pathway cel04361) (21, 23). PMK3
phosphorylates a downstream kinase MAK2 (Uniprot id: Q965G5, M.W. 41.5kDa), a MAP
kinase activated protein kinase (MAPKAPK). Involvement of MAK2 in this cascade turns the
canonical three-kinase MAPK-module into a four-kinase module (Fig. 1A). Schematic
representation of the domain architectures and different constructs used in this study are
shown in Fig. 1C. PMK3 shares highest sequence similarity with p38α among the human

p38-isoforms (Fig. S1A), while MAK2 shares more than 50% identity with two human
MAPKAPKs, MK2 and MK3 (Fig. S1B). To understand the mutual relationship between
these two terminal kinases in further detail, we purified PMK3 and MAK2 recombinantly from
E. coli, that were active in an *in vitro* radioactive kinase assay towards a generic substrate,
myelin basic protein (MBP) (26) (Fig. S1C).

We employed *in vitro* kinase assay using γ-^{32}P-ATP to check phosphorylation of
MAK2 by PMK3. Purified PMK3 and MAK2, mixed at different ratios, were incubated for
different time intervals at 25°C and *de novo* phosphorylation and autophosphorylation were
monitored by autoradiography. It was observed that, both PMK3 and MAK2 were heavily
phosphorylated in a time dependent manner when incubated together (lanes 9-12), but did
not show appreciable autophosphorylation when incubated alone in the same assay
conditions (lanes 1-8) (Fig. 1D). Enhancement of phosphorylation of MAK2 in presence of
PMK3 was expected. However, profound increase in phosphorylation of PMK3 in presence
of MAK2 may result from the direct catalytic activity of MAK2 or by virtue of physical
interaction that leads to autophosphorylation of PMK3 (27, 28). This interdependence was
independent of the relative stoichiometry of the two kinases as similar observations were
made at two different molar ratios of the kinases used in Fig1D (2μM of each) and 1E (1μM
PMK3 and 6μM MAK2).

It is known that, active MKK4 is able to activate PMK3 by phosphorylation (19),
similar to human p38α activation by MKK6 (29-31). PMK3 did not enhance *de novo*
phosphorylation of its upstream kinase MKK4, in a similar assay involving PMK3 and MKK4,
where MKK4 efficiently trans-phosphorylated PMK3 as expected (Fig. 1F).

PMK3 and MAK2 engage in a feedback phosphorylation loop:

Next, we wanted to further analyze the enhancement of phosphorylation of PMK3
and MAK2 in presence of each other (Fig. 2A). We took advantage of a mutant PMK3,
where the invariant ATP binding Lys, K150 was replaced by Met, giving rise to K150M
mutant or kinase dead version of PMK3 (PMK3-KD). This ATP-anchoring Lys, conserved in
all eukaryotic protein kinases, helps position ATP in the kinase, resulting in a productive
phospho-transfer reaction. Mutation of this Lys leads to abrogation of kinase activity; such
mutants, in essence, are either kinase dead or retain minimal activity, if any. In a radioactive
in vitro phosphorylation assay using PMK3-KD and MAK2, phosphorylation signal for PMK3-
KD in the presence or absence of MAK2 (Fig. 2A, lanes 1-8), was hardly detectable. MAK2,
in presence or absence of PMK3-KD, was faintly phosphorylated (Fig. 2A, lanes 5-12), but
this phosphorylated form of MAK2 failed to enhance phosphorylation of PMK3-KD. Next, we
used PMK3-activated phospho-MAK2 (P-MAK2) and monitored phosphorylation of PMK3-

KD and P-MAK2 using the same experimental scheme. Phosphorylation of PMK3-KD in
absence of P-MAK2 (i.e., auto-) was barely detectable, while significant phosphorylation of
PMK3-KD was observed in presence of P-MAK2 (Fig. 2B, compare lanes 1-4 and 5-8).
PMK3-activated P-MAK2, on the other hand, was capable of high level of self-
phosphorylation as well (Fig. 2B, lanes 9-12). We also created a kinase dead version of
MAK2 (K42M mutant, MAK2-KD) and found that, while it failed to undergo any self-
phosphorylation (Fig. 2C, lanes 9-12), it was mildly trans-phosphorylated by PMK3 that
showed very faint phosphorylation signal in these reactions (Fig. 2C, lanes 5-8). In
comparison to MAK2-KD, MAK2 underwent significantly higher-level phosphorylation in
presence of PMK3 (Fig. 2D). Likewise, PMK3 displayed much more efficient *de novo*
phosphorylation signal in presence of MAK2 than MAK2-KD (Fig. 2D).

In order to understand the effect of such feedback relationship, we modeled the
system as depicted in Fig. 2E; wherein X denotes MKK4 (MAP2K), Y denotes PMK3
(MAPK) and Z denotes MAK2 (MAPKAPK). Y_p and Z_p represent the phosphorylated
versions of the respective kinases and are considered active. The model captures the effect
of feedback on the phosphorylation kinetics and relative amounts of phosphorylated kinases
(with respect to the total kinase amount) (Fig. 2E). The effect of feedback is accounted by
considering a dimensionless parameter A_{Z_p}, called feedback strength and sensitivity.
Clearly, such feedback is advantageous to generate active pools of those kinases than
without feedback.

Mass spectrometric analysis of PMK3 and MAK2 phosphorylation:
p38-MAPKs require phosphorylation of both T and Y in the TxY motif in their
activation loops to usher maximum activity, while phosphorylation of only T show much
reduced activity and that of only Y renders the kinase inactive (32). In the previous sections,
we observed that, phosphorylation on both PMK3 and MAK2 increased many folds when
they were both present in the reaction compared to the lone-kinases. However, identities of
phosphosites are unknown. We performed high resolution LC-MS/MS experiments to
identify phosphorylation sites after incubating PMK3, MAK2, and a mixture of PMK3 and
MAK2 with ATP. We considered only those phosphorylated residues with probability of
phosphorylation greater than 0.85 and that were detected at least twice (MS/MS count ≥2).
List of those sites are presented in Table 1; for clarity the phosphosite-identification data are
collated in Venn diagrams (Fig. 3A) and in individual cartoon representation of PMK3 and
MAK2 (Fig. S2A and B, respectively). Evidently, more phosphorylated residues were
identified in both PMK3 and MAK2 when they were incubated together, than individually with
ATP. Most relevant phosphorylation event in the context of PMK3 activation is double

phosphorylation of T285 and Y287 (pT285pY287). Singly phosphorylated peptides bearing pT285 or pY287 were detected in PMK3 alone treated with ATP, without the detection of any peptide containing pT285pY287, although peptides containing pT282pY287 or pS283pY287 were successfully detected in the same sample (Fig. 3A and B). No singly phosphorylated pT285 bearing PMK3 peptide was detected when incubated with ATP in presence of MAK2 although peptides bearing pY287 were observed (Fig. 3A and B). PMK3-peptides containing MAPK-activating pT285pY287 were detected exclusively in presence of MAK2 (Fig. 3B and C, S3C) along with multiple other doubly phosphorylated sites including those observed in the PMK3-only sample. Distribution of phosphorylation sites in the activation segments of PMK3 (Fig. 3D) and in MAK2 (Fig. S2D) are also shown. We also found, using a context-independent phospho-tyrosine specific antibody, that a small fraction of PMK3 was already tyrosine-phosphorylated that enhanced significantly when treated with ATP (Fig. 3E, lanes 1-4) and treatment with MAK2 further enhanced it (Fig. 3E, lanes 5-8). MAK2, on the other hand, was tyrosine-phosphorylated to a very small extent by PMK3 (Fig. 3E, lanes 5-8) while that on MAK2 in absence of PMK3 with or without ATP was barely detectable (Fig. 3E, lanes 9-12).

In order to look at the spatial distribution of phosphorylation sites in three dimensions, two different structural models of the PMK3:MAK2 complex were generated using AlphaFold-predicted structures (33), due to the lack of any experimentally derived structure for either kinase or their complex. Firstly, we superimposed AlphaFold-derived structures of PMK3 and MAK2 on to the experimentally derived structure of the human p38α:MK2 complex (pdb id: 2oza), and also generated a model of the PMK3:MAK2 complex using AlphaFold2. When phosphorylation sites on both PMK3 and MAK2 were highlighted in these two structural models of the said complex, it became clear that it's unlikely for either kinase to access all the phosphorylation sites on the other kinase from either pose (Fig. S2E and F, left and right panels). Neither could these kinases autophosphorylate themselves at sites far from the respective active sites. We speculate that PMK3:MAK2 complex is dynamic in nature giving rise to an ensemble of conformations, and the experimentally derived pose of the human p38α:MK2 complex or AlphaFold2-derived PMK3:MAK2 complex are insufficient to capture the dynamicity and/or plasticity of the said complex.

Sequence and structural analysis of PMK3 and MAK2:

To glean into the complex between PMK3 and MAK2, we used the above-mentioned superimposed-complex structure (pdb id: 2oza) and analyzed the CD and ED domains on PMK3 that are major contributors in recognizing the MAPKAPKs (34, 35). Residues in the ED domain are largely conserved with the exception of N159 of p38α changed to another

polar residue T261 in PMK3, and C162 of p38α changed to S264 in PMK3, that changes -

SH to -OH (Fig. S3A). In the CD domain, however, changes are more drastic; acidic D313 of

p38α changed to hydrophobic L424 in PMK3 and polar, phosphorylatable Y131 of p38α

changed to hydrophobic, phospho-ablative F422 in PMK3 (Fig. S3B). When we turned our

attention to the p38-docking site (R/K–R/K–X_{10}–R/K–R/K–R/K–R/K–R/K), we realized that

this region of MAK2 is not fully conserved compared to its human counterparts MK2 and

MK3 (Fig. S3C). For example, 2[nd] R/K (K374 of MK2) and last R/K (K389 of MK2) have been

changed to S325 and G340, respectively in MAK2 (Fig. S3C). Such changes in the

interacting residues among the partner kinases of human and nematode origins raise the

possibility that the ensuing PMK3:MAK2 and p38α:MK2 complexes may differ in spatial

arrangements as well as affinity.

We also noticed that PMK3 contains a long (~90aa) insert at the N-terminus before

the kinase domain that is absent in the human p38 isoforms (Fig. S1A). On the other hand,

MK2 and MK3 contain long N-terminal insertion before the beginning of the kinase domain

that is absent in MK5 and MAK2 (Fig. S1B). MAK2, however contains a highly charged C-

terminal region beyond the MAPK-docking sequence that is absent in MK2 or MK3 (Fig.

S1B). Both the N-terminal extension in PMK3 and C-terminal extension in MAK2 are largely

unstructured in AlphaFold2-predicted structures (Fig. 4A). Next, we analyzed those

sequences using web-based tools, like MolPhase, FuzDrop, IUPred3, PONDR and

FoldIndex, to gain further insights on their physico-chemical properties (36-39). For PMK3,

pLDDT score was lowest at the N-terminus prior to the kinase domain and the same region

was predicted to be intrinsically disordered by each of the predictive tools employed (Fig.

4A, left panel and S4D); the same was true for the MAK2 C-terminus beyond the kinase

domain (Fig. 4A, right panel and S4E). Presence of intrinsically disordered regions (IDRs)

bestow their parent proteins with multivalency of interaction (40, 41), and presence of long

IDRs in both PMK3 and MAK2 may posit each other in ways to result in an ensemble of

conformation making phosphorylation at different surfaces possible. Notably, MAPK-docking

site on MAK2 resides in its C-terminal IDR (42).

MAK2 undergoes liquid-liquid phase separation and prevents aggregation of PMK3:

Presence of IDRs also gives rise to the possibility of formation of molecular

condensates through liquid-liquid phase separation (LLPS). MolPhase scores of 0.995 and

0.939 for PMK3 and MAK2, respectively, prompted us to investigate this possibility. We

found that Fluorescein-labeled MAK2 (F-MAK2) forms condensates through LLPS at PEG

concentrations of 8% and above (Fig. 4B), and both the number and size of the condensates

increased with the increase in PEG concentration. At 12% PEG, the condensates seemed to

be transitioning from a liquid-like state to an arrested state. To further confirm that LLPS in

MAK2 was not due to labeling of the protein with Fluorescein, we generated and purified a

MAK2 construct containing GFP at its C-terminus (MAK2-GFP) that also underwent LLPS to

form condensate in a similar range of PEG concentrations (Fig. S3F). Size of these

condensates increased as protein concentration was increased from 6uM to 10uM or more

(Fig. S3G). To assess the liquid like properties of the condensates, we performed FRAP

experiments at different concentrations of PEG (43). It was observed that the fluorescence

fully recovered within 60s of bleaching at 8% PEG, while it was gradually arrested as the

PEG concentration was increased to 10% and 12% (Fig. 4C). We also found that LLPS of

MAK2 was severely compromised at higher concentration of NaCl (Fig. S3H). It suggests

that MAK2 LLPS is a reversible process, energetics of which is likely dominated by

electrostatic interactions (44, 45).

Unlike MAK2, Fluorescein-labeled PMK3 (F-PMK3) formed aggregates instead of

phase-separated droplets in the same conditions that lead to LLPS-mediated droplet

formation in MAK2 (Fig. 4D). Furthermore, this aggregation of F-PMK3 could not be

prevented by the chaperone HSP90 or a non-interacting non-specific protein, BSA (Fig. S3I).

Of note, low MW fraction of PMK3 displayed very high activity towards MAK2 (lane 6), while

large MW fraction of PMK3 did not show any detectable activity (lane 3), indicating that

PMK3 present in large self-assembly was inactive (Fig. S3J).

Next, we sought to investigate if MAK2 could prevent the aggregation of PMK3 and

lead to LLPS-mediated droplet formation in PMK3. To this end, we took a fixed

concentration F-PMK3 and incubated it with 10% PEG 8K in absence or presence of

increasing concentration of unlabeled MAK2, and monitored F-PMK3 droplet formation at

different time interval. Interestingly, instead of aggregating, F-PMK3 formed droplets through

LLPS in presence of MAK2 (Fig. 4E). We also observed that, Alexa Fluor 594-labeled PMK3

(AF-PMK3) and MAK2-GFP colocalized in the same droplets as early as 10 mins post

incubation (Fig. 4F), when MAK2-GFP and AF-PMK3 were used in a similar assay. At 3

hours post incubation, droplets showed tendency to aggregate with increasing concentration

of PEG (Fig. 4F). Interestingly, AF-PMK3 and MAK2-GFP complex formed droplets through

LLPS at PEG concentration as low as 1-2% and very few droplets were seen even when no

PEG was present (Fig. 4G). These assays strongly bolstered the idea that MAK2 specifically

engages with PMK3 to prevent its aggregation and it's not a non-specific bystander effect.

MAK2 regulates stability and localization of PMK3 in neuron:

Our findings suggested that PMK3 is prone to aggregation and presence of MAK2

may prevent it to a large extent through LLPS *in vitro*. We wanted to investigate whether this

MAK2 dependent regulation of PMK3 holds true within *C. elegans* neuron. Transgenic

expression of PMK3-TagRFP reporter in touch neuron showed that PMK3 protein majorly
localized to the cell body (Fig. 5A "WT" panel; named WT because it is Wild-Type for *mak-2*;
arrow pointing PLM cell soma) and its nuclear localization was obvious. However, the
intensity of PMK3-TagRFP was dim in the cell body, and it was hardly visible in both anterior
and posterior neurites of PLM neuron (Fig. 5A "WT" panel). In the deletion mutant for *mak-2*,
the intensity of the PMK3-TagRFP reporter gets further dimmed within PLM soma (Fig. 5A
"*mak-2(lf)*" panel). Next, we co-injected PMK3-TagRFP and TagBFP-MAK2 in 1:1 and 1:6
ratios for co-expressing them in touch neurons, named "*mak-2(+)*" and "*mak-2(++)*",
respectively. When MAK2 was co-expressed, the intensity of PMK3-TagRFP was elevated
in the cell body (arrow, Fig. 5A "*mak-2(+)*" panel). Interestingly, PMK3-TagRFP was visible
within the neurites in presence of MAK2 (arrowheads, Fig. A "*mak-2(+)*" panel). Average
intensity of PMK3-TagRFP was significantly higher in *mak-2* overexpression backgrounds as
compared to wildtype in both the cell body and neurites, although diffusible GFP intensity
remained unchanged (Fig. 5B and 5C, Fig. S4A). Of note, PMK3 intensity was slightly
higher in the 1:6 expression line as compared to the 1:1 line (Fig. 5B and 5C). Same trend
was observed in both L4 (larval) and aged animals (Fig. S4B and C). This suggested that
MAK2 helps increase the stability and thereby intensity of PMK3 in both cell body and
neurites of PLM neuron. Similar phenomenon was also noticed in ALM neuron.

We further noticed that when TagBFP-MAK2 was expressed with PMK3-TagRFP, it
resulted in infrequent puncta (condensed structures) formation in the PLM soma of L4 stage
animals (Fig. 5D, "*mak-2(++)*" panel), although these puncta did not appear in transgenic
line which exclusively expresses the PMK3-TagRFP reporter (Fig. 5D, "WT" panel).This
observation was made in both 1:1 and 1:6 lines (arrows, Fig. 5D "*mak-2(++)*" panel, Fig.
S4D). Also, the percentage of worms showing puncta phenotype increased with age (Day3
vs L4 data, Fig. 5E). This indicated that age related stress might be the factor that promotes
these puncta formation, which requires MAK2.

We speculated that MAK2 could be helping in the formation of such condensed
structures in response to stress to accommodate the aggregation-prone PMK3 (46, 47). In
order to test this hypothesis, we used temperature as a stress factor and checked the effect
of heat-shock on PMK3 in presence and absence of MAK2. Of note, heat-shock was shown
to enhance LLPS conditions in other studies (48, 49). After heat shock at 34^0C for two-hours,
the condensed puncta of PMK3-TagRFP in cytoplasm of PLM soma increased significantly
(Fig. 5F and 5G). These puncta were reversible as they disappear after animal was allowed
to recover at 20^0C for same time-period. Whereas in *mak-2* mutant background, such puncta
formation was not seen for PMK3::TagRFP upon heat shock (Fig. 5F and 5G); however,
PMK3::TagRFP intensity permanently decreased in those animals even after recovery at
20^0C (Fig. 5F).

These results indicate that stress condition (like heat shock) causes aggregation of
(possibly misfolded) PMK3 which can be reversed by endogenous MAK2 by preventing
aggregation of PMK3 at favorable conditions. We speculate that PMK3 could be getting
degraded by one of the degradation pathways in absence of MAK2. Similar results were
obtained upon heat-shocking the animals at the Day1 and Day3 adult stage and with *mak-*
2(++) line (Fig. 5G, Fig. S4E). Overall, these experiments demonstrate that MAK-2 regulates
PMK-3 localization and stability *in vivo.*
356

Effect of IDRs and phosphorylation on the feedback relationship of MAK2 and PMK3:

It was evident from our biochemical analyses that MAK2 and PMK3 participate in a
feedback phosphorylation (trans-) loop, whereas they undergo very little, if any, self-
phosphorylation when incubated alone with ATP. To investigate how these self- vs. trans-
phosphorylation modulate LLPS-mediated droplet formation and *vice-versa*, we designed
three different reaction conditions, sets A, B and C as outlined in Fig. S5A. In set A, PMK3
was incubated alone in presence of 10% PEG 8K, where PMK3 aggregates, and then MAK2
was added. In this condition, only the fraction of PMK3 that is available in solution or
condensate would serve as the kinase to phosphorylate MAK2 and as the substrate to be
phosphorylated by MAK2. In set B, MAK2 was incubated alone in presence of 10% PEG 8K
that is favorable for MAK2 condensate formation, and then PMK3 was added. In set C,
PMK3 and MAK2 were incubated together in 10% PEG 8K, where the PMK3:MAK2 complex
should form condensate. Both in sets B and C, larger fraction of PMK3 should be
biochemically available to serve as the kinase or the substrate compared to set A. As
expected, it was observed that in case of set A, phosphorylation signal for PMK3 was much
less as not much of PMK3 was available in solution/condensate, and only a catalytic amount
of PMK3 was available to phosphorylate MAK2 (Fig. 6A, left panel, lanes 1-3). In sets B and
C, where PMK3:MAK2 complex was allowed to form condensates, both proteins underwent
significant phosphorylation (Fig. 6A, left panel, lanes 4-9). We also confirmed that MAK2
was able to phosphorylate PMK3 on tyrosine(s), when both were present in the condensate
(set C) (Fig. 6A, right panel) as was observed when both were present in solution (Fig. 3E).
Next, we checked the effect of ATP-treatment on LLPS of MAK2-GFP and AF-PMK3 by
incubating individual proteins along with 200µM of ATP at 8% PEG-concentration for
different time intervals. They behaved similarly as to form droplets through LLPS for MAK2
and form aggregates for PMK3 when compared with the same protein in absence of ATP
(Fig. S5B and C). Together these results inform us that autophosphorylation, if any, or in
other words, mere presence of ATP is not sufficient to change PMK3's tendency to
aggregate or MAK2's ability to from condensates through LLPS. MAK2:PMK3 complex
underwent droplet formation through LLPS both in absence (Fig. S5D), as was already seen,

and presence of ATP (Fig. 6B). Interestingly, we noticed that after prolonged incubation, the
complex also showed tendency to aggregate in absence of ATP but that tendency to
aggregate was not apparent in presence of ATP as the droplets remained rounded even
after 3hrs post-incubation. These results show that MAK2 is able to form condensates and
incorporate PMK3 into those condensates both in presence and absence of ATP, which
might increase the bio-availability of PMK3 and activate it in presence of ATP.

Next, we wanted to check the effect of the predicted intrinsically disordered regions
(IDRs) in each kinase on their interdependence. In MAK2, the major IDR was predicted to be
between residues 325-366 region; we created and purified two deletion constructs,
MAK2ΔC1 and MAK2ΔC2 encompassing residues 1-317 and 1-341, respectively (Fig. 1C).
MAK2ΔC2 behaved similarly as the full length MAK2 that displayed enhancement of PMK3
phosphorylation but MAK2ΔC1 was unable to do so, wherein only faint phosphorylation
signal of PMK3 was visible (Fig. 6C). We also created two deletion constructs for PMK3,
PMK3ΔN1 and PMK3ΔN2 encompassing residues 45-474 and 95-474, respectively as the
predicted IDRs were located between residues 1-92 (Fig. 1C). Of note, we could
successfully express and purify PMK3ΔN1 but not PMK3ΔN2. Similar observations were
made for PMK3ΔN1 that displayed very little autophosphorylation, and full length MAK2 and
MAK2ΔC2 efficiently phosphorylated it while MAK2ΔC1 couldn't (Fig. 6D). Indeed, with
gradual shortening of the IDR of MAK2 its ability to undergo LLPS went down and shorter
constructs formed progressively smaller droplets compared to the full length MAK2 in similar
assay conditions (Fig. S5E).

Discussion:
MAPK signaling modules have remained in the forefront of signal transduction
research for many years. Many research articles employing biochemical, structural, genetics
and other approaches helped the field to understand the importance of these kinases and
detailed regulatory mechanisms (9, 50, 51). Studies of MAPK signaling in multicellular
(mammalian) and unicellular (yeast) organisms show that the signal flows in a unidirectional
path starting from a MAP3K that activates a cognate MAP2K to activate its downstream
MAPK upon encountering extracellular stimuli. Basic structure and skeleton of this three-
kinase module is preserved wherein MAP2Ks are dual specificity kinases required to
phosphorylate the conserved TxY motif on the activation loop of the MAPKs, essential for
the MAPKs to realize their full scale of activity (32). The three-kinase MAPK modules exhibit
their roles either by directly phosphorylating effector-substrates (like transcription factors) or
by activating a downstream kinase, MAPKAPK, by phosphorylation that ultimately carry out
the phosphorylation of the effector-substrates (9). This makes the respective module a four-

kinase MAPK module. Role of such MAPK-MAPKAPK duo are well established in the literature and have been implicated in mammalian biology, especially in the realm of immune signaling (52-54). Heterodimeric p38α:MK2 complex is known to orchestrate modulation of proinflammatory cytokine productions (55). The complex is resident in the nucleus in resting cells but migrates to the cytoplasm in response to stimuli (12). p38α lacks NES; however, it is transported to the cytoplasm in complex with MK2 that bears an NES. MK2-NES remains masked in its inactive form, phosphorylation (Thr334 in human) of NES by p38α unmasks it (12, 56, 57).

As mentioned earlier, MAPK pathways represent one of the most ancient eukaryotic signal transduction pathways throughout evolution. In *C. elegans* PMK3 and MAK2 are implicated in the axon regeneration process downstream of DLK1 (MAP3K) and MKK4 (MAP2K) (21, 23). The DLK1 pathway is negatively regulated by the E3 ubiquitin ligase RPM-1/Highwire/Phr in worm, fly and mice in order to stabilize the synaptic connections (18-20, 58). PMK3 and MAK2 were shown to interact by yeast two hybrid assay and in a co-immunoprecipitation assay performed in HEK293T cells (21). As expected PMK3 was able to phosphorylate MAK2 *in vitro.* We found that, PMK3 tends to form large oligomeric forms in solution that is catalytically inactive along with catalytically active low molecular weight fractions. PMK3 phosphorylates MAK2 at T173 and Thr285 in addition to other residues. A phospho-ablative double mutant of MAK2 (T173A T285A) failed to exert its role in axonal regeneration in nematode (15). Thr285 in MAK2 is equivalent to Thr334 in human MK2 located in the NES region, phosphorylation of which has been well established in the activation and nuclear export of MK2 and p38α:MK2 complex in human to exert their signaling roles. We found that the activated MAK2 phosphorylates its upstream kinase PMK3 at multiple locations including the TxY (T285-x-Y287) motif essential for PMK3 to attain its highest-activity state. This is a new insight which is unreported in the literature. Identification of doubly phosphorylated pT285pY287 bearing peptide exclusively in presence of MAK2 provokes us to propose this newly found feedback loop as a mechanism for fully activating PMK3 by its downstream kinase. However, allosteric autophosphorylation of PMK3 at T285 by virtue of its interaction with MAK2 through the docking site also contribute to its activation to a minor extent (Fig. 2C & 6C). Interaction of PMK3 with MAK2 through the canonical docking site on MAK2 enables trans-phosphorylation of the substrate kinase as expected (59) but does not enhance PMK3-autophosphorylation that requires catalytic activity of MAK2 as well (Fig. 2C & D, 6C) In addition, PMK3 specifically engages in such a feedback relationship with its downstream MAK2 but not with its upstream MKK4 (MAP2K) (Fig. 1F). Of note, MK2 (mammalian counterpart of MAK2) has been reported to participate in a feedback relationship with RIPK1 that downregulates RIPK1 activity (53, 54). We

speculate that the feedback loop either may act as a fail-safe mechanism to activate PMK3
in a situation when MKK4 is either unavailable or inactivated; or it may help gain quicker and
more robust response of this pathway as and when required (Fig. 2E). Since DLK1 signaling
is used in multiple phenomena such as axon regeneration (23), degeneration (24, 60), and
synaptic pruning (25) in response to nervous system injury, it might need a seamless
mechanisms for robust activation and faithful transmission inside the cell. A critical
dependence on two different p38 MAPK cascades involving DLK1 and MLK1 for axonal
regeneration, indicates a robust cross-talk between parallel signaling kinases in neuronal
injury response (61). Future course of studies will shed light on the scenario in the cellular
context in presence and absence of MKK4.

Phosphorylation mediated regulation of kinases is a common theme in kinase
biology. Distribution of phosphorylation sites on either kinase in this study suggested that the
existing structural models with similar kinases from other organisms were inadequate to
capture the possible modes of binding between PMK3 and MAK2 to enable such distribution
of phosphorylation sites. In addition, both these kinases contained IDRs and showed
theoretical propensity to form droplets. These observations led us to hypothesize that
droplet/condensate formation through LLPS might be a plausible mechanism to
accommodate the ensemble conformational space for this complex. In many cases, kinases
have been found to depend upon scaffolding proteins to undergo phosphorylation by a trans-
kinase or to attain a critical local concentration for self-phosphorylation. Activation of IKK-
complex is a classic example of such mechanism wherein IKK-complex undergo activation
by means of condensate formation through LLPS in the signaling context in a NEMO-
dependent manner (62). Regulation of kinase function through LLPS-driven condensate
formation have been reported in other cases as well (63-68). However, to the best of our
knowledge, LLPS-mediated formation of kinase-condensate that relies upon the presence of
a trans-phosphorylating kinase is not known. Here, we report that an aggregation-prone p38
MAPK (PMK3) is kept bio-available by its activating downstream kinase MAPKAPK (MAK2)
by formation of condensates through LLPS. Furthermore, this LLPS-mediated condensate
formation leads to productive association between the two kinases where a feedback
phosphorylation relationship is functional. Our studies with transgenic *C. elegans* strongly
suggest such functional association between these kinases in the organism, wherein MAK2
regulates the amount of PMK3 under normal and stress conditions. Interestingly, it has
been shown recently that human MKK6:p38α complex is also highly dynamic in nature and
the disordered N-terminal linker of the MAP2K (MKK6) plays critical role in determining
specificity of the MAPK to be activated by phosphorylation (69). This report and our
observations provoke us to speculate that MAPK activation across species might occur
through a highly dynamical engagement of the MAPK with its upstream (MAP2K) or

downstream (MAPKAP) kinases with ensuing conformational plasticity. Species specific
regulatory nuances maybe required for fine tuning of the process.

Together, we propose a hitherto unreported regulatory mechanism for p38-MAP
kinases wherein a critical signaling p38-MAP kinase is kept bio-available through physical
interaction and condensate formation by its downstream substrate kinase that in turn
activate the upstream kinase through a feedback phosphorylation loop (Fig. 6E). Interaction
through the canonical docking-motif on MAK2 is necessary but not sufficient for the full
activation of PMK3 through allosteric means. Full activation of PMK3 defined by double
phosphorylation of the TxY-motif requires catalytic activity of MAK2. Future course of studies
will shed light whether this is a rare, kinase-specific mechanism active in a particular
organism or a more wide-spread aspect in kinase activation and regulation.

Acknowledgements and funding sources: This study was primarily supported by DBT-
Wellcome Trust India Alliance Intermediate Fellowship to SP (IA/I/15/1/501852) and Senior
Fellowship to AGR (IA/S/22/1/506243), and intramural core funding from Bose Institute and
BRIC-NBRC to SP and AGR, respectively. PR and AH thanks the University Grants
Commission, Government of India, and SM thanks Bose Institute for graduate research
fellowship. MN thanks Science and Engineering Research Board (GoI) for the National Post-
doctoral fellowship (PDF/2022/001807). Authors thank Dr. Amitabha Majumdar (NCCS,
Pune) for helpful discussion and the proteomics facility at CSIR-CCMB for mass
spectrometric data collection.

Author contribution:
Conceptualization: SP; Experimental design: SP, AGR and PR; Investigation: PR, CM, PRT,
AH, VB, SM (structural modeling) and MN (kinetic modeling); Data analysis: SP, PR, AGR,
SRC, CM, PRT; Figure: SP, PR, CM, PRT, MN, SM; First draft: SP; Editing: PR, AGR, PRT,
MN

Competing interests: Authors declare no competing financial interests.

References
525

1.   G. Manning, G. D. Plowman, T. Hunter, S. Sudarsanam, Evolution of protein kinase
signaling from yeast to man. *Trends Biochem Sci* **27**, 514-520 (2002).
2.   A. Cuadrado, A. R. Nebreda, Mechanisms and functions of p38 MAPK signalling.
Biochem J **429**, 403-417 (2010).
3.   C. Widmann, S. Gibson, M. B. Jarpe, G. L. Johnson, Mitogen-activated protein
kinase: conservation of a three-kinase module from yeast to human. *Physiol Rev* **79**,
143-180 (1999).
4.   M. Bouhaddou *et al.*, The Global Phosphorylation Landscape of SARS-CoV-2
Infection. *Cell* **182**, 685-712 e619 (2020).

5. S. Luecke *et al.*, Dynamical and combinatorial coding by MAPK p38 and NFkappaB
in the inflammatory response of macrophages. *Mol Syst Biol* 10.1038/s44320-024-
00047-4 (2024).
6. D. S. Battagello *et al.*, Unpuzzling COVID-19: tissue-related signaling pathways
associated with SARS-CoV-2 infection and transmission. *Clin Sci (Lond)* **134**, 2137-
2160 (2020).
7. G. Pearson *et al.*, Mitogen-activated protein (MAP) kinase pathways: regulation and
physiological functions. *Endocr Rev* **22**, 153-183 (2001).
8. B. Ge *et al.*, MAPKK-independent activation of p38alpha mediated by TAB1-
dependent autophosphorylation of p38alpha. *Science* **295**, 1291-1294 (2002).
9. M. Cargnello, P. P. Roux, Activation and function of the MAPKs and their substrates,
the MAPK-activated protein kinases. *Microbiol Mol Biol Rev* **75**, 50-83 (2011).
10. B. Canovas, A. R. Nebreda, Diversity and versatility of p38 kinase signalling in health
and disease. *Nat Rev Mol Cell Biol* **22**, 346-366 (2021).
11. T. Zarubin, J. Han, Activation and signaling of the p38 MAP kinase pathway. *Cell*
Res **15**, 11-18 (2005).
12. R. Ben-Levy, S. Hooper, R. Wilson, H. F. Paterson, C. J. Marshall, Nuclear export of
the stress-activated protein kinase p38 mediated by its substrate MAPKAP kinase-2.
Curr Biol **8**, 1049-1057 (1998).
13. K. Berman, J. McKay, L. Avery, M. Cobb, Isolation and characterization of pmk-(1-3):
three p38 homologs in Caenorhabditis elegans. *Mol Cell Biol Res Commun* **4**, 337-
344 (2001).
14. M. Hammarlund, P. Nix, L. Hauth, E. M. Jorgensen, M. Bastiani, Axon regeneration
requires a conserved MAP kinase pathway. *Science* **323**, 802-806 (2009).
15. D. Yan, Z. Wu, A. D. Chisholm, Y. Jin, The DLK-1 kinase promotes mRNA stability
and local translation in C. elegans synapses and axon regeneration. *Cell* **138**, 1005-
1018 (2009).
16. Y. Hao *et al.*, An evolutionarily conserved mechanism for cAMP elicited axonal
regeneration involves direct activation of the dual leucine zipper kinase DLK. *Elife* **5**
(2016).
17. J. E. Shin *et al.*, Dual leucine zipper kinase is required for retrograde injury signaling
and axonal regeneration. *Neuron* **74**, 1015-1022 (2012).
18. A. M. Schaefer, G. D. Hadwiger, M. L. Nonet, rpm-1, a conserved neuronal gene that
regulates targeting and synaptogenesis in C. elegans. *Neuron* **26**, 345-356 (2000).
19. K. Nakata *et al.*, Regulation of a DLK-1 and p38 MAP kinase pathway by the
ubiquitin ligase RPM-1 is required for presynaptic development. *Cell* **120**, 407-420
(2005).
20. J. W. Lewcock, N. Genoud, K. Lettieri, S. L. Pfaff, The ubiquitin ligase Phr1 regulates
axon outgrowth through modulation of microtubule dynamics. *Neuron* **56**, 604-620
(2007).
21. D. Yan, Z. Wu, A. D. Chisholm, Y. Jin, The DLK-1 kinase promotes mRNA stability
and local translation in C. elegans synapses and axon regeneration. *Cell* **138**, 1005-
1018 (2009).
22. D. Szklarczyk *et al.*, The STRING database in 2023: protein–protein association
networks and functional enrichment analyses for any sequenced genome of interest.
Nucleic acids research **51**, D638-D646 (2023).
23. M. Hammarlund, P. Nix, L. Hauth, E. M. Jorgensen, M. Bastiani, Axon regeneration
requires a conserved MAP kinase pathway. *Science* **323**, 802-806 (2009).
24. E. Asghari Adib, L. J. Smithson, C. A. Collins, An axonal stress response pathway:
degenerative and regenerative signaling by DLK. *Curr Opin Neurobiol* **53**, 110-119
(2018).
25. E. Asghari Adib *et al.*, DLK signaling in axotomized neurons triggers complement
activation and loss of upstream synapses. *Cell Rep* **43**, 113801 (2024).

26.   M. F. Cicirelli, S. L. Pelech, E. G. Krebs, Activation of multiple protein kinases during
the burst in protein phosphorylation that precedes the first meiotic cell division in
Xenopus oocytes. *J Biol Chem* **263**, 2009-2019 (1988).
27.   J. Beenstock *et al.*, The p38β mitogen-activated protein kinase possesses an intrinsic
autophosphorylation activity, generated by a short region composed of the α-G helix
and MAPK insert. *Journal of Biological Chemistry* **289**, 23546-23556 (2014).
28.   J. Beenstock, N. Mooshayef, D. Engelberg, How do protein kinases take a selfie
(autophosphorylate)? *Trends in biochemical sciences* **41**, 938-953 (2016).
29.   J. Han *et al.*, Characterization of the structure and function of a novel MAP kinase
kinase (MKK6). *J Biol Chem* **271**, 2886-2891 (1996).
30.   T. Moriguchi *et al.*, A novel kinase cascade mediated by mitogen-activated protein
kinase kinase 6 and MKK3. *J Biol Chem* **271**, 13675-13679 (1996).
31.   T. Moriguchi *et al.*, Purification and identification of a major activator for p38 from
osmotically shocked cells. Activation of mitogen-activated protein kinase kinase 6 by
osmotic shock, tumor necrosis factor-alpha, and H2O2. *J Biol Chem* **271**, 26981-
26988 (1996).
32.   Y. Y. Zhang, Z. Q. Mei, J. W. Wu, Z. X. Wang, Enzymatic activity and substrate
specificity of mitogen-activated protein kinase p38alpha in different phosphorylation
states. *J Biol Chem* **283**, 26591-26601 (2008).
33.   J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583-589 (2021).
34.   T. Tanoue, M. Adachi, T. Moriguchi, E. Nishida, A conserved docking motif in MAP
kinases common to substrates, activators and regulators. *Nat Cell Biol* **2**, 110-116
(2000).
35.   T. Tanoue, R. Maeda, M. Adachi, E. Nishida, Identification of a docking groove on
ERK and p38 MAP kinases that regulates the specificity of docking interactions.
EMBO J **20**, 466-479 (2001).
36.   Q. Liang *et al.*, MolPhase, an advanced prediction algorithm for protein phase
separation. *EMBO J* **43**, 1898-1918 (2024).
37.   G. Erdos, M. Pajkos, Z. Dosztanyi, IUPred3: prediction of protein disorder enhanced
with unambiguous experimental annotation and visualization of evolutionary
conservation. *Nucleic Acids Res* **49**, W297-W303 (2021).
38.   A. Hatos, S. C. E. Tosatto, M. Vendruscolo, M. Fuxreiter, FuzDrop on AlphaFold:
visualizing the sequence-dependent propensity of liquid-liquid phase separation and
aggregation of proteins. *Nucleic Acids Res* **50**, W337-W344 (2022).
39.   J. Prilusky *et al.*, FoldIndex: a simple tool to predict whether a given protein
sequence is intrinsically unfolded. *Bioinformatics* **21**, 3435-3438 (2005).
40.   M. M. Babu, R. van der Lee, N. S. de Groot, J. Gsponer, Intrinsically disordered
proteins: regulation and disease. *Curr Opin Struct Biol* **21**, 432-440 (2011).
41.   P. E. Wright, H. J. Dyson, Intrinsically disordered proteins in cellular signalling and
regulation. *Nat Rev Mol Cell Biol* **16**, 18-29 (2015).
42.   J. Torres Robles, H. J. Lou, G. Shi, P. L. Pan, B. E. Turk, Linear motif specificity in
signaling through p38α and ERK2 mitogen–activated protein kinases. *Proceedings of*
the National Academy of Sciences **120**, e2316599120 (2023).
43.   N. O. Taylor, M. T. Wei, H. A. Stone, C. P. Brangwynne, Quantifying Dynamics in
Phase-Separated Condensates Using Fluorescence Recovery after Photobleaching.
Biophys J **117**, 1285-1300 (2019).
44.   A. E. Posey, A. S. Holehouse, R. V. Pappu, Phase Separation of Intrinsically
Disordered Proteins. *Methods Enzymol* **611**, 1-30 (2018).
45.   S. Sanulli, G. J. Narlikar, Generation and Biochemical Characterization of Phase-
Separated Droplets Formed by Nucleic Acid Binding Proteins: Using HP1 as a Model
System. *Curr Protoc* **1**, e109 (2021).
46.   P. Panda, V. Sarohi, T. Basak, P. Kasturi, Elucidation of site-specific ubiquitination
on chaperones in response to mutant huntingtin. *Cellular and Molecular*
Neurobiology **44**, 3 (2024).

643 47. B. S. Rutledge, W.-Y. Choy, M. L. Duennwald, Folding or holding?—Hsp70 and
644   Hsp90 chaperoning of misfolded proteins in neurodegenerative disease. *Journal of*
645   *Biological Chemistry* **298** (2022).
646 48. A. Sandhu *et al.*, The microtubule regulator EFA-6 forms cortical foci dependent on
647   its intrinsically disordered region and interactions with tubulins. *Cell reports* **43**
648   (2024).
649 49. L. Thomas, B. Taleb Ismail, P. Askjaer, G. Seydoux, Nucleoporin foci are
650   stress-sensitive condensates dispensable for C. elegans nuclear pore assembly. *The*
651   *EMBO Journal* **42**, e112987 (2023).
652 50. L. Chang, M. Karin, Mammalian MAP kinase signalling cascades. *Nature* **410**, 37-40
653   (2001).
654 51. E. J. Goldsmith, X. Min, H. He, T. Zhou, Structural studies of MAP Kinase cascade
655   components. *Methods Mol Biol* **661**, 223-237 (2010).
656 52. M. Gaestel, MAPKAP kinases - MKs - two's company, three's a crowd. *Nat Rev Mol*
657   *Cell Biol* **7**, 120-130 (2006).
658 53. M. B. Menon *et al.*, p38(MAPK)/MK2-dependent phosphorylation controls cytotoxic
659   RIPK1 signalling in inflammation and infection. *Nat Cell Biol* **19**, 1248-1259 (2017).
660 54. M. B. Menon, J. Gropengiesser, K. Ruckdeschel, M. Gaestel, To die or not to die:
661   Regulatory feedback phosphorylation circuits determine receptor-interacting protein
662   kinase-1 (RIPK1) function. *Mol Cell Oncol* **5**, e1396389 (2018).
663 55. M. B. Menon, M. Gaestel, MK2-TNF-Signaling Comes Full Circle. *Trends Biochem*
664   *Sci* **43**, 170-179 (2018).
665 56. K. Engel, A. Kotlyarov, M. Gaestel, Leptomycin B-sensitive nuclear export of
666   MAPKAP kinase 2 is regulated by phosphorylation. *EMBO J* **17**, 3363-3371 (1998).
667 57. W. Meng *et al.*, Structure of mitogen-activated protein kinase-activated protein
668   (MAPKAP) kinase 2 suggests a bifunctional switch that couples kinase activation
669   with nuclear export. *J Biol Chem* **277**, 37401-37405 (2002).
670 58. Q. Chang, R. J. Balice-Gordon, Highwire, rpm-1, and futsch: balancing synaptic
671   growth and stability. *Neuron* **26**, 287-290 (2000).
672 59. Y. Tokunaga, K. Takeuchi, H. Takahashi, I. Shimada, Allosteric enhancement of
673   MAP kinase p38α's activity and substrate selectivity by docking interactions. *Nature*
674   *Structural & Molecular Biology* **21**, 704-711 (2014).
675 60. J. Gerdts, D. W. Summers, J. Milbrandt, A. DiAntonio, Axon Self-Destruction: New
676   Links among SARM1, MAPKs, and NAD+ Metabolism. *Neuron* **89**, 449-460 (2016).
677 61. P. Nix, N. Hisamoto, K. Matsumoto, M. Bastiani, Axon regeneration requires
678   coordinate activation of p38 and JNK MAPK pathways. *Proc Natl Acad Sci U S A*
679   **108**, 10738-10743 (2011).
680 62. M. Du, C. K. Ea, Y. Fang, Z. J. Chen, Liquid phase separation of NEMO induced by
681   polyubiquitin chains activates NF-kappaB. *Mol Cell* **82**, 2415-2426 e2415 (2022).
682 63. C. C. Lin *et al.*, Receptor tyrosine kinases regulate signal transduction through a
683   liquid-liquid phase separated state. *Mol Cell* **82**, 1089-1106 e1012 (2022).
684 64. T. P. Lopez-Palacios, J. L. Andersen, Kinase regulation by liquid-liquid phase
685   separation. *Trends Cell Biol* **33**, 649-666 (2023).
686 65. D. Schlutermann *et al.*, FIP200 controls the TBK1 activation threshold at
687   SQSTM1/p62-positive condensates. *Sci Rep* **11**, 13863 (2021).
688 66. J. Z. Zhang *et al.*, Phase Separation of a PKA Regulatory Subunit Controls cAMP
689   Compartmentation and Oncogenic Signaling. *Cell* **182**, 1531-1544 e1515 (2020).
690 67. T. Zhu *et al.*, Phase separation underlies signaling activation of oncogenic NTRK
691   fusions. *Proc Natl Acad Sci U S A* **120**, e2219589120 (2023).
692 68. D. Sang *et al.*, Condensed-phase signaling can expand kinase specificity and
693   respond to macromolecular crowding. *Mol Cell* **82**, 3693-3711 e3610 (2022).
694 69. P. Juyoux *et al.*, Architecture of the MKK6-p38alpha complex defines the basis of
695   MAPK specificity and activation. *Science* **381**, 1217-1225 (2023).
696

 Figure Legends

Figure 1: *In vitro* reconstruction of a MAPK-MAPKAPK module involved in
axonal regeneration in *C. elegans.*

(A) Schematic representation of the four-kinase DLK-1 MAPK signaling pathway in *C.*
elegans (blue) along with their human counterparts (dark). PMK3 (MAPK) and MAK2
(MAPKAPK) act as the two terminal kinases in this module. **(B)** Analysis of the
protein-protein interaction network between PMK3 and MAK2 using the STRING
database. Full STRING network as well as physical interaction subnetwork are
shown. **(C)** Top: Domain architecture of PMK3 depicting the kinase domain (Cyan)
and the TxY motif within the activation loop. Lower panel: Domain architecture of
MAK2 depicting the kinase domain (Blue), the nuclear export signal (NES, Orange)
and the nuclear localization signal (NLS, purple). The domain borders have been
numbered. For each kinase, location of the ATP-anchoring invariant lysine residues
and predicted IDR regions are marked. Design of deletion constructs used in this
study are also depicted. **(D-F)** *In vitro* radioactive phosphorylation assays with
recombinant PMK3 (MAPK), MAK2 (MAPKAPK) and MKK4 (MAP2K). (D) *De novo*
phosphorylation of PMK3 and MAK2 were monitored either alone or in presence of
each other. (E) Same as (A) at a different relative stoichiometry to check reliance on
the relative stoichiometry of the two constituent kinases. (F) Similar assay was
performed to monitor directionality of phosphorylation signal transmission between
PMK3 and its upstream kinase MKK4.

Figure 2: PMK3 engages in feedback phosphorylation with the downstream
Kinase MAK2, but not the upstream kinase MKK4.

(A) *de novo* phosphorylation status of a kinase dead PMK3 mutant (PMK3 K150M or
PMK3-KD) and wild type MAK2 were monitored in an *in vitro* radioactive kinase assay
at different time intervals. **(B)** Similar assay was performed with PMK3-KD and
activated MAK2 (P-MAK2, pre-phosphorylated by PMK3) and their *de novo*
phosphorylation status were monitored as a function of time. **(C)** Same as (A), except
a kinase dead mutant of MAK2 (MAK2 K42M or MAK2-KD) and wild type PMK3 was
used in this case. **(D)** *De novo* phosphorylation of wild type PMK3, MAK2 and MAK2-
KD were monitored at different time points in two separate reactions where PMK3
was incubated with MAK2 or MAK2-KD. **(E)** Computational modelling of the effect of
feedback loop on the overall phosphorylation kinetics of the two kinases. The
temporal evolution of relative concentrations of Y_p (phosphorylated PMK3) and Z_p
(phosphorylated MAK2) are presented with respect to total kinase amount of each

kinase for different values of feedback strengths (A_{Z_p}), which is a dimensionless

measure of feedback strength and sensitivity.

Figure 3: Identification of phosphorylation sites on PMK3 and MAK2.

(A) Venn diagrams showing identities of singly and doubly-phosphorylated residues

in PMK3 and MAK2 identified by LC-MS/MS analyses when incubated with ATP

either individually or together. PMK3-residues phosphorylated in PMK3-only and with

MAK2 samples are shown in blue and purple, respectively while residues common in

both samples are depicted in dark letters (left panel). MAK2-residues phosphorylated

in MAK2-only and with PMK3 samples are shown in green and purple, respectively

and those common in both samples are depicted in dark letters (right panel). In each

case, doubly-phosphorylated entities are shaded with respective colors. **(B)**

Percentage occurrence of individual PMK3-phosphosite in presence (purple) and

absence (blue) of MAK2 are compared. Percentage occurrence of phosphorylation

for each phosphorylation site in PMK3 is calculated from MS/MS counts of each

phospho-peptide in total phospho-peptide pool. Only those phosphorylation sites are

considered having probability > 0.85 and MS/MS counts ≥ 2. **(C)** MS/MS spectrum of

a PMK3-peptide containing doubly phosphorylated TxY motif (-KDTSLpTQpYVQTR-)

is shown. **(D)** Ribbon representation of the activation segment of PMK3 from DFG- to

APE-motif. Phosphorylation sites identified in this study are shown in sticks; wherein

purple color depicts residues phosphorylated only in presence of MAK2 while those in

dark gray are found both in presence and absence of MAK2. **(E)** PMK3 and MAK2

were incubated with non-radioactive 'cold' ATP and subjected to immunoblot

analyses using a context-independent anti p-tyrosine antibody. The blot is a

representative of three independent experiments.

Figure 4: PMK3 and MAK2 harbor unstructured, disordered regions and

propensity to undergo phase separation

(A) Left panel: FuzDrop prediction shows the droplet promoting regions (DPRs) along

with the aggregation hotspots in the PMK3 sequence, AlphaFold predicted structure

of PMK3 colored according to the pLDDT score (signifies confidence of prediction) is

also shown. Right panel: Shows FuzDrop analysis of MAK2 and its AlphaFold

predicted structure colored in the same manner as PMK3 structure. The number of

aggregation hot spots in the MAK2 sequence is distinctly lower than those found in

PMK3. Region spanning amino acids 325-341 contains the MAPK-docking site. Color

coding of the pLDDT score is shown at the bottom. **(B)** Fluorescein labelled MAK2 (F-

MAK2) was incubated with varying concentrations of PEG-8000 for different time
periods and imaged under the microscope. MAK2 undergoes phase separation to
form the round droplets under both the conditions. Images are representative of three
independent experiments. Scale bar is 10µm. **(C)** FRAP analysis on the phase
separated MAK2-GFP condensates at different PEG concentrations (left). Scale bar
is 10µm.The fluorescence recovery in each case was followed up to 60 seconds post
bleaching and the normalized fluorescence intensity was plotted against time (right).
The data values are shown as mean±s.d. of normalized fluorescence intensity values
from three independent measurements (n=3). **(D)** *In vitro* droplet formation assays
with fluorescein labelled PMK3 show that PMK3 forms aggregates at different protein
concentrations in the presence of 10% PEG-8K. Images are representative of three
independent experiments. Scale bar is 10µm. **(E)** F-PMK3 was mixed with unlabeled
MAK2 at two different concentrations in presence of 10% PEG and imaged at 10
minutes and 2 hours respectively. In presence of MAK2, PMK3 undergoes liquid-
liquid phase separation to form globular droplets instead of aggregates. Images are
representative of two independent experiments. Scale bar is 10µm. **(F)** MAK2 and
PMK3 co-localize within the same phase separated droplets. MAK-GFP (green) and
Alexa 594 labelled PMK3 (red) were mixed together in presence of varying
concentrations of PEG-8K (8-12%) and imaged at two time points. Merged (yellow)
images show the co-localization of the two kinases within a single droplet. At 3 hours,
the droplets exhibit a tendency to form aggregates with an increase in PEG
concentration. Images are representative of two independent experiments. Scale bar
is 8µm. **(G)** In presence of MAK2-GFP, PMK3 undergoes LLPS even in the absence
of any crowding agent or as low as 1% PEG-8K. Scale bar is 8µm.

Figure 5: MAK2 regulates stability and localization of PMK3 in neuron.
(A) The expression of P*mec4-*PMK3-TagRFP using a transgenic extra-chromosomal
array (*shrEx522*) shows that PMK-3 localizes to the cell body (including nucleus) of
posterior gentle-touch (PLM) neuron of C. elegans ("WT" panel, named WT because
it is Wild-Type for *mak-2*), arrow pointing PLM cell soma). PMK3-TagRFP intensity
gets dimmer in *mak-2*(ok2394) background. Co-expression of P*mec4*-PMK3-TagRFP
and P*mec4*-TagBFP-MAK2 in a transgenic extra-chromosomal array (*shrEx526*) of
1:1 ratio, named "*mak-2*(+)", shows that PMK3-TagRFP intensity is brighter both in
the cell body (arrow, "*mak-2*(+)" panel) and axon (arrowheads, "mak-2(+)" panel)
(Scale bar - 10 µm). (B) Scatter-plot showing the ratio of the mean intensity of PMK3-
TagRFP vs diffusible GFP in region of interest (ROIs) in cell body (elliptical ROI-

major axis=6 μm, minor axis=4 μm) in different genetic backgrounds. Increasing trend
of mean-intensity ratio between PMK3-TagRFP and diffusible GFP in *mak-2*
overexpression backgrounds as compared to wildtype in PLM cell body and axon
(anterior neurite). (C) Scatter-plot showing the ratio of the mean intensity of PMK3-
TagRFP vs diffusible GFP in region of interest (ROIs) (~ 60-μm line ROI of 5-pixel
thickness) in PLM axon in different genetic backgrounds. (D) High-resolution images
of the cell body region of PLM neuron in wild type and in a transgenic
extrachromosomal array (*shrEx527*) co-expressing P*mec4*-PMK3-TagRFP and
P*mec4*-TagBFP-MAK2 in 1:6 ratio, named "*mak-2*(++)" (Scale bar - 2 μm). (E)
Quantification of number of PMK3-TagRFP/TagBFP-MAK2 puncta in PLM soma
upon aging. The violin plot shows that the penetrance of puncta phenotype among
animals increases with age. N=1, n~15 for each genotype. (F) The high-resolution
images of the PLM cell body region for Pmec4-PMK3-TagRFP localization before and
after heat shock. Upon heat-shock at 34deg for 2 hours, PMK3-TagRFP puncta can
be seen in PLM soma. After recovery at 20 degrees, the puncta disappear. This
puncta phenomenon is perturbed in *mak-2*(ok2394) (Scale bar - 2 micron). (G)
Quantification of puncta number in L4 (larval) PLM soma pre- and post- heat shock
and recovery. N=2, n~30 for each genotype except for mak-2(++) where N=1, n~15.
Each dot in the graphs represent number of puncta per worm. Statistics: One-way
Anova with Tukey's multiple comparison test, P<0.001(***), <0.002(**), <0.033(*),
<0.12(ns).

Figure 6: Effect of auto- and feedback phosphorylation on LLPS of the MAK2-
PMK3 complex.
(A) Left panel: PMK3 and MAK2 were incubated with radioactive ATP as per the
reaction scheme shown in Figure S6, resolved by SDS-PAGE and *de novo*
phosphorylation was monitored by autoradiography. Right panel: Reaction scheme C
was performed with cold ATP and immunoblotting was performed using anti p-
tyrosine antibodies to detect context-independent phosphorylation at the tyrosine
residues of the two kinases. The blot is a representative of three independent
experiments. **(B)** Effect of trans-phosphorylation on PMK3: MAK2 condensates. Scale
bar is 8μm. In presence of ATP, the PMK3: MAK2 droplets maintained their globular
look even at 3 hours post incubation. **(C)** *In vitro* radioactive kinase assay was
performed using full length PMK3 incubated separately with full length MAK2 and two
deletion constructs of MAK2, and *de novo* phosphorylation of each protein was
monitored at different time points using autoradiography. **(D)** Similar assay, as

described in *C* was performed using the same constructs of MAK2 with an N-terminal

deletion construct of PMK3. **(E)** A graphical model depicting mutual relationship and

signal transmission through PMK3 and MAK2. PMK3, a neuronal p38-MAPK is

activated by its downstream MAPKAPK MAK2 without requiring the presence of

canonical activator MAP2K, MKK4. MAK2 increases bioavailability of aggregation

prone PMK3 by LLPS-driven condensate formation.

Supplementary Figures:

Figure S1: (A) Left panel: Multiple sequence alignment of C. elegans PMK3

(CePMK3) with the four human p38 (Hsp38) MAPK isoforms α, β, γ and δ. Right

panel: The percent identity matrix obtained from the MSA of the p38 MAPK

sequences. Darker shade in the box implies higher sequence identity. PMK3 shares

highest sequence similarity with human p38α. **(B)** Left panel: Multiple sequence

alignment (MSA) of MAK2 with the human MAPKAPKs MK2, MK3 and MK5. Right

panel: The percent identity matrix shows that MAK2 shares more than 50% sequence

similarity with MK2 and MK3. **(C)** *In vitro* radioactive kinase assay with purified,

recombinant PMK3 and MAK2 using myelin basic protein (MBP) as a generic

substrate (n=2).

Figure S2: (A-B) Phosphorylation sites identified by LC-MS/MS experiments are

shown on cartoon representations of PMK3 and MAK2 in *A* and *B*, respectively.

Residues colored in blue were exclusively identified in individual kinases when the

other kinase was absent, while those in purple were absent in the sample containing

both kinases, and those in dark gray were identified in both types of samples.

Residues connected with lines signify identification of those residues in doubly

(dotted lines) or triply (solid lines) phosphorylated peptides. **(C)** MS/MS spectrum of a

PMK3-peptide containing doubly phosphorylated TxY motif (DTSLpTQpYVQTR). **(D)**

Ribbon representation of the activation segment of MAK2 from DFG- to APE-motif.

Phosphorylation sites identified in this study are shown in sticks where in purple

shows residues phosphorylated only in presence of PMK3 while those in dark gray

are found both in presence and absence of PMK3. **(E)** Ribbon representation of the

PMK3(cyan):MAK2(blue) complex obtained by superimposing on a p38α:MK2 (pdb

id: 2oza). Phosphorylated residues identified by mass-spectrometric analyses have

been highlighted (orange and purple) on both the kinases. **(F)** Ribbon representation

of the AlphaFold2-predicted PMK3(cyan):MAK2(blue) complex. Phosphorylated

residues identified by mass-spectrometric analyses have been highlighted (orange

and purple) on those kinases.

Figure S3: (A-B) Superimposition of the ED and CD domains of PMK3 and human
p38α. The respective sequence alignments have been shown in the lower panel.
Changes in the nature of amino acids have been highlighted in boxes. **(C)**
Superimposition of the docking sites of MAK2 and MK2 notable changes are boxed.
(D-E) IUPred3, PONDR, FoldIndex and MolPhase webserver-based predictions of
PMK3 (D) and MAK2 (E) primary sequences. **(F)** A MAK2 construct with a GFP tag at
its C-terminus (MAK2-GFP) could also successfully form these condensates through
LLPS at the different PEG concentrations used. Images are representative of three
independent experiments. Scale bar is 10μm. **(G)** Concentration dependent phase
separation of MAK2-GFP in presence of 6% PEG 8K. MAK2-GFP was mixed with 6%
PEG at different protein concentrations ranging from 1 to 40 μM and imaged under
the microscope to monitor droplet formation. Scale bar is 8μm. **(H)** F-MAK2 and
MAK2-GFP were subjected to phase separation in presence of increased amount of
NaCl, wherein the droplet formation was significantly inhibited in both the cases at
600mM NaCl. Images are representative of two independent experiments. Scale bar
is 8μm (upper panel) and 10μm (lower panel) respectively. **(I)** PMK3 forms
aggregates even in the presence of chaperone Hsp90 or a non-specific protein, BSA.
Scale bar is 10μm. **(J)** Left panel: Comparison of the size exclusion profiles of 6X His-
PMK3 and a molecular weight standard resolved on a preparative gel filtration
column. The absorbance at 280nm has been plotted against the elution volume. The
larger molecular weight fractions constitute Peak 1 and the lower molecular weight
fractions constitute Peak2. Right panel: *In vitro* radioactive kinase assay with purified,
recombinant PMK3 and MAK2. PMK3 from two peaks in SEC was used for the assay
in different reactions.

Figure S4: (A) Comparison of mean intensity of PMK3-TagRFP (Red-scatter plot)
and diffusible GFP (Green-scatter plot) in region of interest in cell body and PLM
axon in different genetic backgrounds at Day1 adult stage. **(B)** Comparison of mean
intensity of PMK3-TagRFP, diffusible GFP and their ratio in region of interest in cell
body and PLM axon in different genetic backgrounds at L4 stage. **(C)** Comparison of
mean intensity of PMK3-TagRFP, diffusible GFP and their ratio in region of interest in
cell body and PLM axon in different genetic backgrounds at Day3 adult stage. **(D)**
High-resolution images of the cell body region of PLM neuron in mak-2(ok2394)
background and in transgenic extrachromosomal array (shrEx526) co-expressing
Pmec4-PMK3-TagRFP and Pmec4-TagBFP-MAK2 in 1:1 ratio (Scale bar - 2 μm). **(E)**
High-resolution images of the PLM soma before and after heat shock in the

923 transgenic line (shrEx527) co-expressing Pmec4-PMK3-TagRFP and Pmec4-
924 TagBFP-MAK2 in 1:6 ratio.

926 **Figure S5: (A)** Outline of the reaction schemes followed for the *in vitro* LLPS and
927 subsequent kinase assays. **(B-C)** Condensate formation of MAK2 (B) and
928 aggregation of PMK3 (C) in presence and absence of ATP. **(D)** Condensate
929 formation of the PMK3:MAK2 complex through LLPS in absence of ATP. **(E)**
930 Decrease in droplet size with the shortening of the C-terminal IDR of MAK2.

Figure 1

A) Axonal Injury

DLK1 *(MAP3K)* **MAP3K12**
↓
MKK4 *(MAP2K)* **MKK6**
↓
PMK3 *(MAPK)* **p38α**
↓
MAK2 *(MAPKAPK)* **MK2**

B)

supr-1
pmk-2
pmk-1
oxbp-1
pmk-3
pmk-3
mak-1
sek-1
oxpp-6
mak-2

— from curated databases
— experimentally determined
— gene neighborhood
— gene fusions
— gene co-occurrence
— textmining
— co-expression
— protein homology

Full STRING Network   Physical Subnetwork

C)

IDR — Kinase Domain — PMK3
1   92   113 K150M   416

T285 TxY Y207

PMK3-KD
PMK3ΔN1   45 — 474
PMK3ΔN2   95 — 474

T173 T285 IDR
MAK2   1 13   K42M   276 301 318 353 366
MAK2-KD   366
MAK2ΔC1   317
MAK2ΔC2   341

D) γP³²-ATP

PMK3   PMK3

Time (mins.): 0 15 30 60   0 15 30 60   0 15 30 60
            PMK3   MAK2

Autoradiograph
75—
PMK3► 50—
MAK2► 37—

Coomassie
75—
PMK3►
50—
MAK2► 37—

Lanes: 1 2 3 4   5 6 7 8   9 10 11 12

E) γP³²-ATP

PMK3   MAK2

Time (mins.): 0 15 30 60   0 15 30 60   0 15 30 60
            PMK3   MAK2

Autoradiograph
75—
PMK3► 50—
MAK2► 37—

Coomassie
75—
PMK3►
50—
MAK2► 37—

Lanes: 1 2 3 4   5 6 7 8   9 10 11 12

F) ATP/ATP*

MKK4   PMK3

Time (mins.): 15 30   15 30 60   15 30 60
            MKK4   PMK3

Autoradiograph
75—
PMK3► 50—
MKK4► 37—

Coomassie
75—
PMK3►
50—
MKK4► 37—

Lanes: 1 2 3   4 5 6   7 8 9

Figure 2

Figure 3

A)

Phosphorylation sites on PMK3

Phosphorylation sites on MAK2

B)

C)

D)

PMK3 Activation Segment

E)

Figure S2

Figure S3

A) Comparison of ED domain

p38α: N159 E160 D161 C162 E163
PMK3: T261 D262 D263 S264 D265

B) Comparison of CD domain

p38α: D313 D316 E178 F231 Y129
PMK3: L424 D427 D178 F231 F422

C) Comparison of docking sites of MAK2 and MK2

MK2:   K373 K374 X10 K R R K 385
MAK2: K324 S325 X10 K336 R337 R338 K340

D) PMK3 — MolPhase parameters of PMK3; score 0.995

E) MAK2 — MolPhase parameters of MAK2; 0.939

F)
MAK2—GFP — 8% PEG 8K, 10% PEG 8K, 12% PEG 8K (1 hour, 2 hours)

G) 6% PEG 8K, 10 minutes — 1μM, 4μM, 8μM, 12μM, 16μM, 20μM, 30μM, 40μM — MAK2—GFP / MAK2

H) 100mM NaCl / 600mM NaCl — 10% PEG 8K, 12% PEG 8K — MAK2—GFP / MAK2 — 10 mins

I) PMK3 — BSA / HSP90 — 10% PEG 8K — 10 minutes, 2 hours, 3 hours

J) PMK3 Peak 1 / PMK3 Peak 2 — M.W. (kDa) Std. (670, 158, 44, 17, 1.35) — A280 (mAU) vs Elution Volume (ml)

PMK3 / MAK2 — Peak 1 / Peak 2 — Autoradiograph — Lanes 1–8

Figure S4

Figure S5

A)

Incubation at 25°C for 45 minutes in 10% PEG

Add **MAK2**   Add **PMK3**

@Room Temperature

Centrifuge @16K rpm, 30mins

Discard supernatant | Resuspend the pellet

In vitro kinase assay   γ-P^{32}-ATP

Reaction stopped at indicated time

SDS-PAGE & autoradiography

(Cold ATP was used for the immunoblot analysis)

B) MAK2 MAK2-GFP MAK2

30 minutes / 3 hours — 8% PEG 8K

No ATP | + ATP

C) PMK3 PMK3 PMK3

30 minutes / 3 hours — 8% PEG 8K

No ATP | + ATP

D) PMK3 PMK3 MAK2-GFP MAK2 MAK2

30 minutes / 3 hours — 8% PEG8K

MAK2-GFP | PMK3 | Merged

E)

Average Feret diameter (µm)

MAK2ΔC1   MAK2ΔC2   MAK2WT