Establishment of a rapid method to assemble and transfer DNA fragments into the JCVI-syn3B minimal synthetic bacterial genome

Atsuko Uenoyama[1], Hana Kiyama[1], Mone Mimura[1], and Makoto Miyata[1,2]*

[1]Graduate School of Science, Osaka Metropolitan University, Osaka, 558-8585, Japan
[2]The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka City University, Osaka, 558-8585, Japan

*Corresponding author:
Makoto Miyata
Graduate School of Science, Osaka Metropolitan University, Osaka, 558-8585, Japan
Tel: +81-6-6605-3157
E-mail: miyata@omu.ac.jp
ORCID ID: https://orcid.org/0000-0002-7478-7390

Abstract

JCVI-syn3B (syn3B), a minimal synthetic bacterium that only possesses essential

genes, facilitates the examination of heterogeneous gene functions in minimal life.

Conventionally, *Escherichia coli* is used to construct DNA fragments for gene transfer

into the syn3B genome. However, the construction process is challenging and time-

consuming due to various issues, including the inhibition of *E. coli* growth and

unexpected recombination, especially with AT-rich DNA sequences such as those found

in *Mycoplasma* genes. Therefore, in this study, we aimed to develop a new

transformation method to overcome these issues. We assembled the vector and target

DNA fragments using an in vitro homologous recombination system and subsequently

transferred the products into the syn3B genome. We obtained approximately 5,000

recombinant colonies per milliliter of the original culture in eight days, which is four

days shorter than the conventional period, without any recombination issues, even for

AT-rich DNA.

Key words

Homologous recombination, *loxP*, Transformation

Significance

The minimal synthetic bacterium JCVI-Syn3B is a useful tool for basic and applied

biology. Plasmids for transformation are generally constructed using *Escherichia coli*;

however, unexpected recombination occurs in some cases. To prevent such issues, this

study developed a new method without any in vivo *E. coli* system.

Introduction

JCVI-syn3B (syn3B) is a minimal synthetic bacterium whose genome only consists of essential genes derived from the small bacterium, *Mycoplasma mycoides* [1-3]. Its genome contains a locus of *loxP* sites to which external genes are transferred via the Cre recombinase activity (Fig. 1) [2]. It is useful in both basic and applied biology [4-8]. Transformation requires a plasmid containing the target DNA, Cre recombinase, *loxP* sites, and puromycin resistance gene [2]. We previously constructed a plasmid using *Escherichia coli* (Fig. 2) as described below. Day 1: Polymerase chain reaction (PCR)-amplified vector and target DNA fragment were assembled via homologous recombination-based methods using the NEBuilder® and In-Fusion® systems. The constructs were then used to transform *E. coli* cells. Day 2: Transformed colonies were inoculated into a liquid medium. Days 3–5: Plasmids were isolated and sequenced. Days 6–9: Syn3B was transformed with the plasmid. Days 10–12: Colonies were cultured in a liquid medium and target gene transfer was confirmed via direct PCR. This method exhibited some drawbacks, including poor *E. coli* growth and unexpected recombination. Moreover, unexpected gene recombination was observed in *Mycoplasma* and related species, possibly due to their high AT content (67–76%) [9,10]. Therefore, in this study, we aimed to explore four transfer methods without *E. coli* to develop a new *E. coli*-free approach that is four days faster than the conventional method for gene transfer.

Materials and Methods

Bacterial strains and culture conditions

JCVI-syn3B (GenBank, CP069345.1) and *E. coli* (DH5α) were cultured in the SP4 [1,2] and Luria–Bertani media at 37°C, respectively.

Preparation of DNA fragments

Mycoplasma mobile genome was isolated as previously described [11]. Vector- and focused *M. mobile* DNA fragments were amplified from pSD128 DNA [4] and *M. mobile* genomic DNA using primers P1–P2 and P3–P4 (Table 1), respectively. The DNA fragments were assembled using the NEBuilder® HiFi DNA Assembly Master Mix (New England Biolabs, Inc., MA, USA) and In-Fusion HD® Cloning Kit (Takara Bio Inc., Kusatsu, Japan). Then, the assembled product was amplified via PCR using

primers P5 and P6 and circularized via self-assembly with NEBuilder®.

Transformation and direct PCR

Transformation of JCVI-syn3B was performed as previously described

[5]. Transformed colonies were picked, inoculated into 100 μL of SP4 medium

containing puromycin (3 μg/mL), cultured for 18–24 h, and confirmed via PCR using

primers P7 and P8.

Results and Discussion

Transformation using assembled and amplified linear DNA

For transformation, we assembled the plasmid using an in vitro homologous

recombination system and amplified it using PCR. The amplified DNA fragments (Fig.

3(i)) were transferred into the syn3B genome. As our target DNA, we used a 1,916-bp

fragment containing the 1,536-bp gene encoding phosphoglycerate kinase from *M.*

mobile (MMOB4530) and its flanking region (at 565,247–567,161 nt on the genome)

[12-16]. Initially, we performed assembly using the In-Fusion® system, but the

resulting target fragments could not be successfully amplified via PCR, possibly due to

the absence of DNA polymerase and ligase for the gap-filling activity of the In-Fusion®

system that prevents mutagenesis during filling [17,18]. Subsequently, we used

NEBuilder® with an in-built gap-filling activity [19]. Using NEBuilder®, we

successfully amplified the DNA fragments (Fig. 4A). Upon transforming syn3B with

100 ng of the PCR product, 392 ± 124 colonies (n=6) were obtained. However, the

efficiency was approximately 1/20 of that achieved with plasmid pSD128 of *E. coli*

(Fig. 5(i)). We investigated the cause for this low efficiency by performing

transformation using circularized DNA.

Transformation using assembled, amplified, and circularized DNA

PCR products were circularized by NEBuilder® (Fig. 3(ii)). The amount of DNA

converted to a form other than the original linear form was estimated from the agarose

gel electrophoresis band intensity (Fig. 4B), which indicated a 94.6% conversion. The

DNA mixture was used for transformation (Fig. 5(ii)). Notably, transformation

efficiency was similar to that achieved with the *E. coli* plasmid. These results suggest

that circularized DNA achieves a transformation efficiency similar to that achieved with

the supercoiled plasmids of *E. coli*. Next, we examined the products assembled without

PCR amplification.

Transformation using assembled circular DNA

The product assembled using NEBuilder®, which was expected to be circular, was

directly integrated into the syn3B genome (Fig. 3(iii)). The DNA product assembled

using 30 ng vector and 28 ng target DNA was transferred into syn3B, resulting in 2,067

$\pm$ 262 colonies from 425 µL cultures (n=6; Fig. 5(iii)). This shows the advantage of

circular DNA in the uptake process. Next, we assembled the product using In-Fusion®

(Fig. 3(iv)) and obtained 1,282 $\pm$ 418 colonies from 425 µL cultures (n=6; Fig. 5(iv)),

indicating plasmid instability caused by the gap region ($P = 0.005$ via Student's *t*-test).

Here, the transformation efficiency could not be directly compared with that of the

conventional method as the precise amount of assembled DNA was unknown. However,

sufficient number of colonies were obtained for gene transfer using the NEBuilder® and

In-Fusion® products.

Unexpected recombination in the transformants

We examined the transfer of the target DNA into *loxP* via direct PCR using primers

P7 and P8 (Fig. 1) for 18 transformants, and the process was repeated six times (Fig. 6).

For the colonies obtained using the conventional method, we observed PCR bands at the

correct positions. However, compared to that of the conventional method, success rates

(74, 75, 93, and 92%) were low for the linear (Fig. 3(i)), circular (Fig. 3(ii)),

NEBuilder®-assembled (Fig. 3(iii)), and In-Fusion®-assembled (Fig. 3(iv)) DNAs,

respectively. This decrease in success rate was due to unexpected amplification during

PCR. Additionally, transformation efficiency of our direct transformation method was

lower than that of conventional transformation using *E. coli* plasmids. This was

expected, as the isolated plasmids were cloned into *E. coli* colonies.

To date, we have constructed 58 syn3B constructs with AT content of 65–76%.

Moreover, using the established method, we successfully transferred a 14.5-kb fragment

consisting of eight genes of *M. mobile* into the syn3B genome. We also transferred the

genes of *Haloplasma contractile* into syn3B using the new method [20,21], which is

impossible with *E. coli*-based gene manipulation systems. Approximately 58 kb of the

transferred DNA region was sequenced from 18 transformants. Two clones each with

single-nucleotide deletions and substitutions were observed. These mutations were
possibly caused by PCR, as similar mutations have been reported in *E. coli*-based
recombination PCR products.
139

Conclusion
141   Overall, we developed a direct rapid method for gene transfer to the JCVI-syn3B
142 strain, particularly useful for DNA sequences unstable or harmful to *E. coli*.
143

Conflict of Interest
145   The authors declare no conflicts of interest.
146

Author Contributions
148   AU performed all of the experiments. AU and MM wrote an early version of the
149 manuscript. All authors contributed to the project and manuscript.
150

Data Availability Statement
152   All data generated/analyzed in this study are included in this article.
153

Acknowledgements
155   This study was supported by the Grant-in-Aid for Scientific Research (A) (grant
156 number JP17H01544), JST CREST grant (grant number JPMJCR19S5 to MM), and
157 Grant-in-Aid for JSPS Fellows (grant number JP24KJ0189 to HK).

Figure Legends
160 **Fig. 1** Schematic of DNA transfer into the JCVI-syn3B (syn3B) genome. Plasmid
161 pSD128 used for gene transfer contained a Cre/*loxP*-related region and puromycin
162 resistance gene. Target DNA was transferred into the syn3B genome via the Cre
163 recombinase activity. Blue arrows indicate the primers used for direct polymerase chain
164 reaction (PCR).
165

Fig. 2 Comparison of the conventional method with the new established method. The
new method was four days shorter than the conventional method.

Fig. 3 DNA preparation for transformation. (i) Linear DNA, including the target DNA,

Cre/*loxP* system, and puromycin resistance gene. (ii) Circular DNA prepared using

NEBuilder®. (iii) DNA assembled from two fragments using NEBuilder®. (iv) DNA

assembled from two fragments using In-Fusion®. Four types of DNA were used for

syn3B transformation.

Fig. 4 DNA fragments used for transformation. Size markers are shown in the left lanes.

(A) Three PCR products used in this study. (B) DNA fragments used for transformation.

(i) Linear and (ii) circular DNA fragments, as indicated in Fig. 3, are shown.

Fig. 5 Transformation efficiencies with various DNA fragments. Amount of circular

DNA was estimated based on the electrophoresis band intensity. DNA amounts

obtained with (iii) NEBuilder® and (iv) In-Fusion® accounted for the total DNA

amount used for assembly.

Fig. 6 Direct PCR analysis of the transformants. PCR was performed on 18 colonies

obtained using each DNA plasmid.

Table 1. Primers used in this study

Name	Sequence
P1	5'-GTCAAAGATCATACCGTTCG-3'
P2	5'-AGCTAGCTCAGTCCTTATTT-3'
P3	5'-AAATAAGGACTGAGCTAGCTTTGAATCGAACTGAATATGC-3'
P4	5'-CGAACGGTATGATCTTTGACTACTCATAATATGCTACCTC-3'
P5	5'-GGTTGGACTCAAGACGATAG-3'
P6	5'-GTCTTGAGTCCAACCGGCCGGCCCGGAGGCGTCCC-3'
P7	5'-GTAAATTCCCAAATTATTCCATCA-3'
P8	5'-TATGTGATAATGCCAATCGCTAAG-3'

References

[1] Hutchison, C.A., 3rd, Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Ellisman, M.H., et al. Design and synthesis of a minimal bacterial genome. Science 351, aad6253 (2016). https://doi.org/10.1126/science.aad6253

[2] Pelletier, J.F., Sun, L., Wise, K.S., Assad-Garcia, N., Karas, B.J., Deerinck, T.J., et al. Genetic requirements for cell division in a genomically minimal cell. Cell 184, 2430-2440 e2416 (2021). https://doi.org/10.1016/j.cell.2021.03.008

[3] Pelletier, J.F., Glass, J.I., Strychalski, E.A. Cellular mechanics during division of a genomically minimal cell. Trends Cell Biol 32, 900-907 (2022). https://doi.org/10.1016/j.tcb.2022.06.009

[4] Nishiumi, F., Kawai, Y., Nakura, Y., Yoshimura, M., Wu, H.N., Hamaguchi, M., et al. Blockade of endoplasmic reticulum stress-induced cell death by *Ureaplasma parvum* vacuolating factor. Cell Microbiol, e13392 (2021). https://doi.org/10.1111/cmi.13392

[5] Kiyama, H., Kakizawa, S., Sasajima, Y., Tahara, Y.O., Miyata, M. Reconstitution of a minimal motility system based on *Spiroplasma* swimming by two bacterial actins in a synthetic minimal bacterium. Sci Adv 8, eabo7490 (2022). https://doi.org/10.1126/sciadv.abo7490

[6] Moger-Reischer, R.Z., Glass, J.I., Wise, K.S., Sun, L., Bittencourt, D.M.C., Lehmkuhl, B.K., et al. Evolution of a minimal cell. Nature 620, 122-127 (2023). https://doi.org/10.1038/s41586-023-06288-x

[7] Sandberg, T.E., Wise, K.S., Dalldorf, C., Szubin, R., Feist, A.M., Glass, J.I., et al. Adaptive evolution of a minimal organism with a synthetic genome. iScience 26, 107500 (2023). https://doi.org/10.1016/j.isci.2023.107500

[8] Bittencourt, D.M.C., Brown, D.M., Assad-Garcia, N., Romero, M.R., Sun, L., Palhares de Melo, L.A.M., et al. Minimal bacterial cell JCVI-syn3B as a chassis to investigate interactions between bacteria and mammalian cells. ACS Synth Biol 13, 1128-1141 (2024). https://doi.org/10.1021/acssynbio.3c00513

[9] Razin, S. The genus *Mycoplasma* and related genera (class Mollicutes). in The Prokaryotes: Volume 4: Bacteria: Firmicutes, Cyanobacteria. (Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., Stackebrandt, E. eds.), pp. 836-904 Springer US, New York, NY (2006).

[10] Wang, Y., Huang, J.M., Zhou, Y.L., Almeida, A., Finn, R.D., Danchin, A., et al. Phylogenomics of expanding uncultured environmental Tenericutes provides insights into their pathogenicity and evolutionary relationship with *Bacilli*. BMC Genomics 21, 408 (2020). https://doi.org/10.1186/s12864-020-06807-4

[11] Uenoyama, A., Kusumoto, A., Miyata, M. Identification of a 349-kilodalton
protein (Gli349) responsible for cytadherence and glass binding during gliding
of *Mycoplasma mobile*. J Bacteriol 186, 1537-1545 (2004).
https://doi.org/https://doi.org/10.1128%2FJB.186.5.1537-1545.2004

[12] Toyonaga, T., Kato, T., Kawamoto, A., Kodera, N., Hamaguchi, T., Tahara,
Y.O., et al. Chained structure of dimeric F_1-like ATPase in *Mycoplasma mobile*
gliding machinery. mBio 12, e0141421 (2021).
https://doi.org/10.1128/mBio.01414-21

[13] Tulum, I., Kimura, K., Miyata, M. Identification and sequence analyses of the
gliding machinery proteins from *Mycoplasma mobile*. Sci Rep 10, 3792 (2020).
https://doi.org/10.1038/s41598-020-60535-z

[14] Nishikawa, M., Nakane, D., Toyonaga, T., Kawamoto, A., Kato, T., Namba, K.,
et al. Refined mechanism of *Mycoplasma mobile* gliding based on structure,
ATPase activity, and sialic acid binding of machinery. mBio 10, e02846-02819
(2019). https://doi.org/10.1128/mBio.02846-19

[15] Nakane, D., Miyata, M. Cytoskeletal "jellyfish" structure of *Mycoplasma*
mobile. Proc Natl Acad Sci U S A 104, 19518-19523 (2007).
https://doi.org/https://doi.org/10.1073/pnas.0704280104

[16] Jaffe, J.D., Stange-Thomann, N., Smith, C., DeCaprio, D., Fisher, S., Butler, J.,
et al. The complete genome and proteome of *Mycoplasma mobile*. Genome Res
14, 1447-1461 (2004). https://doi.org/https://doi.org/10.1101%2Fgr.2674004

[17] Chen, C.G., Fabri, L.J., Wilson, M.J., Panousis, C. One-step zero-background
IgG reformatting of phage-displayed antibody fragments enabling rapid and
high-throughput lead identification. Nucleic Acids Res 42, e26 (2014).
https://doi.org/10.1093/nar/gkt1142

[18] Oh, S., Jokhadze, G., Duong, T., Bostick, M., Farmer, A.A. Optimized In-
Fusion cloningsystem demonstrates high efficiency and accuracy of multi-
fragment cloning. *Synthetic Biology: Engineering, Evolution & Design*, (2014).

[19] Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A., 3rd,
Smith, H.O. Enzymatic assembly of DNA molecules up to several hundred
kilobases. Nat Methods 6, 343-345 (2009). https://doi.org/10.1038/nmeth.1318

[20] Antunes, A., Alam, I., El Dorry, H., Siam, R., Robertson, A., Bajic, V.B., et al.
Genome sequence of Haloplasma contractile, an unusual contractile bacterium
from a deep-sea anoxic brine lake. J Bacteriol 193, 4551-4552 (2011).
https://doi.org/10.1128/JB.05461-11

[21]   Antunes, A., Rainey, F.A., Wanner, G., Taborda, M., Patzold, J., Nobre, M.F., et

261   al. A new lineage of halophilic, wall-less, contractile bacteria from a brine-filled

262   deep of the Red Sea. J Bacteriol 190, 3580-3587 (2008).

263   https://doi.org/10.1128/JB.01860-07

Fig. 1

Fig. 2

Conventional method	Day	New method
· Assembly of vector and target gene · Transformation into *E. coli*	1	· Assembly of vector and target DNA · Transfer into JCVI-syn3B cell
· Colony pick up and cultivation	2	
· Plasmid isolation · Sequencing	3	
	4	
	5	
· Transfer into JCVI-syn3B cell	6	· Colony pick up and directPCR · Sequencing
	7	
	8	
	9	
· Colony pick up and directPCR	10	
	11	
	12	

Fig. 3

Fig. 4

Fig. 5

Fig. 6