Rapid *in vitro* method to assemble and transfer DNA fragments into the JCVI-syn3B minimal synthetic bacterial genome through Cre/*loxP* system

Atsuko Uenoyama[1], Hana Kiyama[1], Mone Mimura[1], and Makoto Miyata[1,2]*

[1]Graduate School of Science, Osaka Metropolitan University, Osaka, 558-8585, Japan
[2]The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka City University, Osaka, 558-8585, Japan

*Corresponding author:
Makoto Miyata
Graduate School of Science, Osaka Metropolitan University, Osaka, 558-8585, Japan
Tel: +81-6-6605-3157
E-mail: miyata@omu.ac.jp
ORCID ID: https://orcid.org/0000-0002-7478-7390

Abstract

JCVI-syn3B (syn3B), a minimal synthetic bacterium that only possesses essential

genes, facilitates the examination of heterogeneous gene functions in minimal life.

Conventionally, *Escherichia coli* is used to construct DNA fragments for gene transfer

into the syn3B genome through Cre/*loxP* system. However, the construction process is

challenging and time-consuming due to various issues, including the inhibition of *E.*

coli growth and unexpected recombination, especially with AT-rich DNA sequences

such as those found in *Mycoplasma* genes. Therefore, in this study, we aimed to develop

a new transformation method to overcome these issues. We assembled the vector and

target DNA fragments using an in vitro homologous recombination system and

subsequently transferred the products into the syn3B genome. We obtained

approximately $10^3 \sim 10^4$ recombinant colonies per milliliter of the original culture in

eight days, which is four days shorter than the conventional period, without any

recombination issues, even for AT-rich DNA. This method may be applicable to other

gene manipulation systems based on Cre/*loxP* system.

Key words

Homologous recombination, Transformation

Significance

A rapid and trouble free method was developed to transfer genes to the genome of

minimal synthetic bacterium JCVI-Syn3B through Cre/*loxP* system. This method can be

applied to Cre/*loxP*-based gene manipulation system in various research fields.

Introduction

JCVI-syn3B (syn3B) is a "synthetic" bacterium based on a small bacterium, *Mycoplasma mycoides*, which was established to answer the question what is the minimal life? The necessity of individual *M. mycoides* genes were judged by bioinformatics and experiments. The syn3B genome was designed by a computer, synthesized chemically, assembled in *E. coli* and yeast, and then transplanted to a related species, *Mycoplasma capricolum* [1]. The syn3B genome contains a locus of *loxP* site to which external genes can be efficiently transferred via the Cre recombinase activity (Fig. 1) [2-4]. Therefore, syn3B is a useful platform for both basic and applied researches [3,5-8]. Our group is discussing the origin of cell motility, based on syn3B experiments [5,9]. In the transformation of syn3B by external genes requires a plasmid containing the target DNA, Cre recombinase, *loxP* sites, and puromycin resistance gene [2]. We construct these plasmids using *Escherichia coli* (Fig. 1) as described below. (a, day 1) Polymerase chain reaction (PCR)-amplified vector and target DNA fragment were assembled via homologous recombination-based methods using the NEBuilder® and In-Fusion® systems. The constructs were then used to transform *E. coli* cells. (b, day 2) Transformed colonies were inoculated into a liquid medium. (c, days 3–5) Plasmids were isolated and sequenced. (d, days 6–9) Syn3B was transformed with the plasmid. (e, days 10–12) Colonies were cultured in a liquid medium and target gene transfer was confirmed via direct PCR. This method exhibited some drawbacks, including poor *E. coli* growth and unexpected recombination. Moreover, we suffered from unexpected gene recombination in transfer of some *Mycoplasma* and related species, possibly due to their high AT content (67–76%) [10,11]. Therefore, in this study, we aimed to explore four transfer methods without *E. coli* to develop a new *E. coli*-free approach that is four days faster than the conventional method for gene transfer.

Materials and Methods

Bacterial strains and culture conditions

JCVI-syn3B (GenBank, CP069345.1) and *E. coli* (DH5α) were cultured in the SP4 [1,2] and Luria–Bertani media at 37°C, respectively.

Preparation of DNA fragments

Mycoplasma mobile genome was isolated as previously described [12]. Vector and

focused *M. mobile* DNA fragments were amplified from pSD128 DNA [3] and *M.*

mobile genomic DNA using primers P1–P2 and P3–P4 (Table 1), respectively. The

DNA fragments were assembled using the NEBuilder® HiFi DNA Assembly Master

Mix (New England Biolabs, Inc., MA, USA) and In-Fusion HD® Cloning Kit (Takara

Bio Inc., Kusatsu, Japan). Then, the assembled product was amplified via PCR using

primers P5 and P6 and circularized via self-assembly with NEBuilder®.

Transformation and direct PCR

Transformation of JCVI-syn3B was performed as previously described

[5]. In short, competent cells were incubated with respective DNA for transfer for 15

min on ice, treated with 70% PEG_{6000}, and then incubated in SP4 medium at 37°C for 3

h. Cells were then plated on a SP4 (containing horse serum instead of FBS) plate

supplemented by 3 μg/mL puromycin. After incubation at 37°C for four days, the

number of transformed colonies was counted. The transformation efficiency is shown as

values relative to those of 100 ng pSD128 plasmid. This is because the efficiency

largely depended on serum. For further studies, the transformed colonies were picked

and inoculated into 100 μL of SP4 medium containing puromycin (3 μg/mL) for 18–24

h to make stocks, and confirmed via PCR using primers P7 and P8.

Results and Discussion

Rationale of the transformation protocol

To skip DNA manipulation in *E. coli*, we assembled the DNA fragments using an

in vitro homologous recombination system and amplified them using PCR to obtain

large amount of uniform DNA fragments, beneficial for transformation. Four types of

DNAs were tried as shown in Fig. 2. As our target DNA, we used a 1,916-bp fragment

containing the 1,536-bp gene encoding phosphoglycerate kinase from *M. mobile*

(MMOB4530) and its flanking region (at 565,247–567,161 nt on the genome) [13-17].

Transformation using assembled and amplified linear DNA

First, we tried an assembled and amplified DNA fragments (Fig. 2(i)). Initially, we

performed assembly using the In-Fusion® system, but the size of the resulting target

fragments was shorter than the full length (Fig. 2B). In-Fusion® system has no DNA

polymerase and ligase for the gap-filling activity that prevents mutagenesis during

filling [18,19]. Therefore, the target DNA with the region between P5 or P6 should be

amplified by the PCR (Fig. 2B right). This assumption was supported by observations

that the amplified fragment changed with the length of target fragment. Subsequently,

we used NEBuilder® with an in-built gap-filling activity [20]. Using NEBuilder®, we

successfully amplified the DNA fragments (Fig. 2B). Upon transforming syn3B with

100 ng of the PCR product, 392 ± 124 colonies (n = 6) were obtained. However, the

efficiency was approximately 1/50 of that achieved with a plasmid (pSD128 including

target DNA) isolated from *E. coli* (Fig. 3A(i)). We investigated the cause for this low

efficiency by performing transformation using circularized DNA.

Transformation using assembled, amplified, and circularized DNA

PCR products were circularized by NEBuilder® (Fig. 2(ii)). The amount of DNA

converted to a form other than the original linear form was estimated from the agarose

gel electrophoresis band intensity (Fig. 2B), which indicated a 94.6% conversion. The

DNA mixture was used for transformation (Fig. 3A(ii)). Notably, 6,458 ± 1428 colonies

from 425 μL cultures (n = 6) were obtained. The transformation efficiency was

approximately 3/10 that achieved with the plasmid isolated from *E. coli*. These results

suggest that circularized DNA achieves a transformation efficiency similar to that

achieved with the supercoiled plasmids isolated from *E. coli*. Next, we examined the

products assembled without PCR amplification.

Transformation using assembled circular DNA

The product assembled using NEBuilder®, which was expected to be circular, was

directly transferred into the syn3B genome (Fig. 2(iii)). The DNA product assembled

using 30 ng vector and 28 ng target DNA was transferred into syn3B, resulting in 2,067

± 262 colonies from 425 μL cultures (n = 6; Fig. 3A(iii)). This shows the advantage of

circular DNA in the uptake process. Next, we assembled the product using In-Fusion®

(Fig. 2(iv)) and obtained 1,282 ± 418 colonies from 425 μL cultures (n = 6; Fig.

3A(iv)), indicating plasmid instability caused by the gap region ($P = 0.005$ via Student's

t-test). Here, the transformation efficiency could not be directly compared with that of

the conventional method as the precise amount of assembled DNA was unknown.

However, sufficient number of colonies were obtained for gene transfer using the

NEBuilder® and In-Fusion® products.

Unexpected recombination in the transformants

We examined the transfer of the target DNA into *loxP* via direct PCR using primers

P7 and P8 (Fig. 1) for 18 transformants, and repeated the whole process of this

examination for six times (Fig. 3B). For the colonies obtained using the conventional

method, we observed PCR bands at the correct positions in 97% of colonies. However,

compared to that of the conventional method, success rates (74, 75, 93, and 92%, n =

108) were low for the linear (Fig. 2(i)), circular (Fig. 2(ii)), NEBuilder®-assembled

(Fig. 2(iii)), and In-Fusion®-assembled (Fig. 2(iv)) DNAs, respectively. This decrease

in success rate may be due to unexpected amplification during PCR, because the

decrease depended on number of PCR application and the success rate was 97% when

the plasmid cloned in *E. coli* was used. Additionally, transformation efficiency of our

direct transformation method was lower than that of conventional transformation using

plasmids isolated from *E. coli*.

Originally, we started this method to overcome problems in transferring AT-rich

Mollicutes DNA into the syn3B genome. To date, we have constructed 58 syn3B

transformants with AT content of 65–76%. Moreover, using the established method, we

successfully transferred a 14.5-kb fragment consisting of eight genes of *M. mobile* into

the syn3B genome. We also transferred the genes of *Haloplasma contractile* into syn3B

using the new method [21,22], which is impossible with *E. coli*-based gene

manipulation systems. Approximately 58 kb of the transferred DNA region was

sequenced from 18 transformants. Two clones each with single-nucleotide deletions and

substitutions were observed. These mutations were possibly caused by PCR, as similar

mutations have been reported in *E. coli*-based recombination of PCR products.

Cre/*loxP* system is used for genome and gene manipulation widely for microorganisms,

animals, plants, and human [23-25]. Although our study focused on basic research of

syn3B, the direct assembly method may be applicable for other targets.

Conclusion

165 Overall, we developed an *in vitro* direct rapid method for gene transfer to the JCVI-

166 syn3B strain appreciable for Cre/*loxP* recombination. Our method is particularly useful

167   for DNA sequences unstable or harmful to *E. coli*, and also appreciable for various

168   systems based on Cre/*loxP* recombination.

Conflict of Interest

171   The authors declare no conflicts of interest.

Author Contributions

174   AU performed all of the experiments. AU and MM wrote an early version of the

175   manuscript. All authors contributed to the project and manuscript.

Data Availability Statement

178   All data generated/analyzed in this study are included in this article.

Acknowledgements

181   This study was supported by the Grant-in-Aid for Scientific Research (A) (grant

182   number JP17H01544), JST CREST grant (grant number JPMJCR19S5 to MM), and

183   Grant-in-Aid for JSPS Fellows (grant number JP24KJ0189 to HK).

Figure Legends

Fig. 1 DNA transfer into the JCVI-syn3B (syn3B) genome. (A) Schematic illustration.

Plasmid pSD128 used for gene transfer contains a Cre/*loxP*-related region and

puromycin resistance gene. Target DNA is transferred into the syn3B genome via the

Cre recombinase activity. Blue arrows indicate the primers used for direct polymerase

chain reaction (PCR). The conventional (red) and new (green) processes are shown as

five (a–e) and two (f–g) steps, respectively. (B) Gantt chart for the conventional (red)

and new (green) processes.

Fig. 2 DNAs used for transformation. (A) Schematic illustration for preparation of four

DNAs. (i) Linear DNA, including the target DNA, Cre/*loxP* system, and puromycin

resistance gene. (ii) Circular DNA prepared using NEBuilder®. (iii) DNA assembled

from two fragments using NEBuilder®. (iv) DNA assembled from two fragments using

In-Fusion®. Four types of DNA were used for syn3B transformation. (B) Left:

Electrophoresis of DNAs used for transformation. Size markers are shown in the left

lane. pSD128 and Target DNA are the fragments to be assembled. The fourth lane is the

result of PCR amplified from In-Fusion® product. Four DNAs used for transformation

are shown as numbered (i) – (iv). 200 and 150 ng DNA loaded on lanes (ii) – (iii) and

(iv), respectively. The other lanes were loaded with 100 ng DNA. Right: Schematic

illustration of amplified In-Fusion® product. The outermost arcs present DNA

fragments amplified by this PCR.

Fig. 3 Transformation of syn3B. (A) Efficiencies with various DNAs, shown as values

relative to the results of 100 ng pSD128, ranging 1,800~23,000 colonies per mL.

Amount of circular DNA was estimated based on the electrophoresis band intensity.

DNA amounts obtained with (iii) NEBuilder® and (iv) In-Fusion® accounted for the

total DNA amount used for assembly. (B) Direct PCR analysis of the transformants.

PCR was performed using P7 and P8 primers as shown in Fig. 1A on 18 colonies

obtained using individual DNAs. The band position for expected transfer is marked by

red triangles.

Table 1. Primers used in this study

Name	Sequence
P1	5'-GTCAAAGATCATACCGTTCG-3'
P2	5'-AGCTAGCTCAGTCCTTATTT-3'
P3	5'-AAATAAGGACTGAGCTAGCTTTGAATCGAACTGAATATGC-3'
P4	5'-CGAACGGTATGATCTTTGACTACTCATAATATGCTACCTC-3'
P5	5'-GGTTGGACTCAAGACGATAG-3'
P6	5'-GTCTTGAGTCCAACCGGCCGGCCCGGAGGCGTCCC-3'
P7	5'-GTAAATTCCCAAATTATTCCATCA-3'
P8	5'-TATGTGATAATGCCAATCGCTAAG-3'

References

[1]   Hutchison, C.A., 3rd, Chuang, R.Y., Noskov, V.N., Assad-Garcia, N., Deerinck, T.J., Ellisman, M.H., et al. Design and synthesis of a minimal bacterial genome. Science 351, aad6253 (2016). https://doi.org/10.1126/science.aad6253

[2]   Pelletier, J.F., Sun, L., Wise, K.S., Assad-Garcia, N., Karas, B.J., Deerinck, T.J., et al. Genetic requirements for cell division in a genomically minimal cell. Cell 184, 2430-2440 e2416 (2021). https://doi.org/10.1016/j.cell.2021.03.008

[3]   Nishiumi, F., Kawai, Y., Nakura, Y., Yoshimura, M., Wu, H.N., Hamaguchi, M., et al. Blockade of endoplasmic reticulum stress-induced cell death by *Ureaplasma parvum* vacuolating factor. Cell Microbiol, e13392 (2021). https://doi.org/10.1111/cmi.13392

[4]   Mariscal, A.M., Kakizawa, S., Hsu, J.Y., Tanaka, K., Gonzalez-Gonzalez, L., Broto, A., et al. Tuning gene activity by inducible and targeted regulation of gene expression in minimal bacterial cells. ACS Synth Biol 7, 1538-1552 (2018). https://doi.org/10.1021/acssynbio.8b00028

[5]   Kiyama, H., Kakizawa, S., Sasajima, Y., Tahara, Y.O., Miyata, M. Reconstitution of a minimal motility system based on *Spiroplasma* swimming by two bacterial actins in a synthetic minimal bacterium. Sci Adv 8, eabo7490 (2022). https://doi.org/10.1126/sciadv.abo7490

[6]   Moger-Reischer, R.Z., Glass, J.I., Wise, K.S., Sun, L., Bittencourt, D.M.C., Lehmkuhl, B.K., et al. Evolution of a minimal cell. Nature 620, 122-127 (2023). https://doi.org/10.1038/s41586-023-06288-x

[7]   Sandberg, T.E., Wise, K.S., Dalldorf, C., Szubin, R., Feist, A.M., Glass, J.I., et al. Adaptive evolution of a minimal organism with a synthetic genome. iScience 26, 107500 (2023). https://doi.org/10.1016/j.isci.2023.107500

[8]   Bittencourt, D.M.C., Brown, D.M., Assad-Garcia, N., Romero, M.R., Sun, L., Palhares de Melo, L.A.M., et al. Minimal bacterial cell JCVI-syn3B as a chassis to investigate interactions between bacteria and mammalian cells. ACS Synth Biol 13, 1128-1141 (2024). https://doi.org/10.1021/acssynbio.3c00513

[9]   Miyata, M., Robinson, R.C., Uyeda, T.Q.P., Fukumori, Y., Fukushima, S.I., Haruta, S., et al. Tree of motility - A proposed history of motility

systems in the tree of life. Genes Cells 25, 6-21 (2020).

https://doi.org/10.1111/gtc.12737

[10] Razin, S. The genus *Mycoplasma* and related genera (class Mollicutes).

in The Prokaryotes: Volume 4: Bacteria: Firmicutes, Cyanobacteria.

(Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., Stackebrandt,

E. eds.), pp. 836-904 Springer US, New York, NY (2006).

[11] Wang, Y., Huang, J.M., Zhou, Y.L., Almeida, A., Finn, R.D., Danchin, A.,

et al. Phylogenomics of expanding uncultured environmental Tenericutes

provides insights into their pathogenicity and evolutionary relationship

with *Bacilli*. BMC Genomics 21, 408 (2020).

https://doi.org/10.1186/s12864-020-06807-4

[12] Uenoyama, A., Kusumoto, A., Miyata, M. Identification of a 349-kilodalton

protein (Gli349) responsible for cytadherence and glass binding during

gliding of *Mycoplasma mobile*. J Bacteriol 186, 1537-1545 (2004).

https://doi.org/https://doi.org/10.1128%2FJB.186.5.1537-1545.2004

[13] Toyonaga, T., Kato, T., Kawamoto, A., Kodera, N., Hamaguchi, T.,

Tahara, Y.O., et al. Chained structure of dimeric F_1-like ATPase in

Mycoplasma mobile gliding machinery. mBio 12, e0141421 (2021).

https://doi.org/10.1128/mBio.01414-21

[14] Tulum, I., Kimura, K., Miyata, M. Identification and sequence analyses of

the gliding machinery proteins from *Mycoplasma mobile*. Sci Rep 10,

3792 (2020). https://doi.org/10.1038/s41598-020-60535-z

[15] Nishikawa, M., Nakane, D., Toyonaga, T., Kawamoto, A., Kato, T.,

Namba, K., et al. Refined mechanism of *Mycoplasma mobile* gliding

based on structure, ATPase activity, and sialic acid binding of machinery.

mBio 10, e02846-02819 (2019). https://doi.org/10.1128/mBio.02846-19

[16] Nakane, D., Miyata, M. Cytoskeletal "jellyfish" structure of *Mycoplasma*

mobile. Proc Natl Acad Sci U S A 104, 19518-19523 (2007).

https://doi.org/https://doi.org/10.1073/pnas.0704280104

[17] Jaffe, J.D., Stange-Thomann, N., Smith, C., DeCaprio, D., Fisher, S.,

Butler, J., et al. The complete genome and proteome of *Mycoplasma*

mobile. Genome Res 14, 1447-1461 (2004).

https://doi.org/https://doi.org/10.1101%2Fgr.2674004

[18] Chen, C.G., Fabri, L.J., Wilson, M.J., Panousis, C. One-step zero-

background IgG reformatting of phage-displayed antibody fragments

enabling rapid and high-throughput lead identification. Nucleic Acids Res
42, e26 (2014). https://doi.org/10.1093/nar/gkt1142

[19]   Oh, S., Jokhadze, G., Duong, T., Bostick, M., Farmer, A.A. Optimized In-
Fusion cloningsystem demonstrates high efficiency and accuracy of
multi-fragment cloning. *Synthetic Biology: Engineering, Evolution &*
Design, (2014).

[20]   Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A.,
3rd, Smith, H.O. Enzymatic assembly of DNA molecules up to several
hundred kilobases. Nat Methods 6, 343-345 (2009).
https://doi.org/10.1038/nmeth.1318

[21]   Antunes, A., Alam, I., El Dorry, H., Siam, R., Robertson, A., Bajic, V.B.,
et al. Genome sequence of Haloplasma contractile, an unusual
contractile bacterium from a deep-sea anoxic brine lake. J Bacteriol 193,
4551-4552 (2011). https://doi.org/10.1128/JB.05461-11

[22]   Antunes, A., Rainey, F.A., Wanner, G., Taborda, M., Patzold, J., Nobre,
M.F., et al. A new lineage of halophilic, wall-less, contractile bacteria
from a brine-filled deep of the Red Sea. J Bacteriol 190, 3580-3587
(2008). https://doi.org/10.1128/JB.01860-07

[23]   Foster, M.P., Benedek, M.J., Billings, T.D., Montgomery, J.S. Dynamics
in Cre-*loxP* site-specific recombination. Curr Opin Struct Biol 88, 102878
(2024). https://doi.org/10.1016/j.sbi.2024.102878

[24]   Maizels, N. Genome engineering with Cre-*loxP*. J Immunol 191, 5-6
(2013). https://doi.org/10.4049/jimmunol.1301241

[25]   Sternberg, N., Hamilton, D. Bacteriophage P1 site-specific
recombination. I. Recombination between *loxP* sites. J Mol Biol 150, 467-
486 (1981). https://doi.org/10.1016/0022-2836(81)90375-2

Fig. 1

A

B

Fig. 2

Fig. 3