Title: Plant functional groups and root traits are linked to exudation rates of mature temperate trees.

Authors: Young E. Oh[1], Meghan G. Midgley[2], M. Luke McCormack[2], Katilyn V. Beidler[3],
Marshall McCall[4], Savannah Henderson[2], Renato K. Braghiere[5,6], and Richard P. Phillips[1]

1. Department of Biology, Indiana University, Bloomington IN, USA
2. Center for Tree Science, The Morton Arboretum, Lisle, IL, United States,
3. Department of Plant and Microbial Biology, University of Minnesota, Saint Paul MN, USA
4. School of the Environment, Yale University, New Haven, USA
5. Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA
6. California Institute of Technology, Pasadena, CA, US

Keywords: fine root, root traits, root economic space, rhizodeposition, belowground carbon
allocation, rhizosphere carbon flux

Abstract.

While root exudation has the potential to affect soil biogeochemistry profoundly, the process is
rarely quantified in mature, field-grown trees. We measured rates of carbon (C) exudation in 11
trees species that exhibit divergent root traits, including gymnosperms and angiosperms that
associate with either arbuscular mycorrhizal (AM) or ectomycorrhizal (EcM) fungi. Our goal was
to explore how tree species, plant functional groups and root traits collectively influence exudation
patterns. Intraspecific variation in exudation rates was larger than interspecific variation, and
neither functional groups nor morphological traits alone could sufficiently explain variation in this
flux. EcM-associated gymnosperms exuded 2.4 times more C than EcM angiosperms and 1.5 times
more than AM gymnosperms. Exudation rates correlated positively with specific root length (SRL)
and specific root area (SRA), and were correlated with root tissue density and root diameter in
EcM-associated species. Mixed-effect models revealed that exudation rates were best determined
by a combination of phylogenetic group, tree-mycorrhizal type and SRA, though a large portion
of unexplained variation suggests that contemporary environmental and local edaphic conditions
are likely important. Collectively, our results reveal that exudation is a complex physiological
process governed by multiple factors and cannot be fully explained by functional groups or root
traits alone. Instead, a combined consideration of these factors and new experimental approaches
may be needed before exudation patterns can be linked to plant trait frameworks and incorporated
into large-scale models.

Introduction

Global environmental changes are altering plant community composition, with poorly understood impacts on belowground processes and biogeochemical cycling (Fei et al., 2017; Jo et al., 2019). Root carbon (C) exudation is a physiological process that links aboveground-belowground interactions (Bardgett, 2014; McCormack et al., 2017; Weemstra et al., 2022; Wen et al., 2022) and, in many cases, mediates ecosystem responses to global change (Norby et al., 2024; Phillips et al., 2011). Root exudates represent 5–20% of photosynthetically-fixed C (Chari et al., 2024), and much of this C fuels rhizosphere microbes that, in turn, determine soil organic matter dynamics (Chari & Taylor, 2022) and nutrient availability (Brzostek et al., 2014; Finzi et al., 2015; Meier et al., 2017; Yin et al., 2014). In this way, exudation rates affect ecosystem C balance through their effects on both nutrient uptake/primary production and microbial decomposition. Given this role, a deeper understanding of factors that mediate exudate fluxes should enhance our understanding of the ecosystem effects of plant community change (Freschet et al., 2021; Jo et al., 2019; McCormack et al., 2017).

Root traits impact many ecosystem processes (Bardgett, 2014), yet there is little consensus about which traits, if any, align with exudation rates. Exudation rates have been reported to be associated positively with both specific root length (SRL; Meier et al., 2020; Tückmantel et al., 2017; Wang et al., 2021) and negatively with root tissue density (RTD; Sun et al., 2017, 2021) - traits that capture different dimensions of root economic space (RES; Bergmann et al., 2020; McCormack & Iversen, 2019; Weemstra et al., 2016; Weigelt et al., 2021). In the RES, RTD and root N represent a 'conservation' gradient (e.g., 'fast' *vs.* 'slow'), whereby long-lived, tissue-dense roots (high RTD) slowly provision N to hosts relative to fast-growing acquisitive roots with low RTD and high root N (Bergmann et al., 2020; Weigelt et al., 2021). Orthogonal to this axis is the 'collaboration' gradient (e.g., 'outsourcing' *vs.* 'do-it-yourself') defined by SRL and root diameter (Bergmann et al., 2020; McCormack & Iversen, 2019; Weemstra et al., 2016; Weigelt et al., 2021; Wen et al., 2022; Yaffar et al., 2022). Here, large diameter roots with low SRL are colonized by mycorrhizal fungi to a greater extent than thin, high SRL roots (Bergmann et al., 2020; Weigelt et al., 2021). However, links between exudation and both axes of the RES remain unclear, indicating that the relationship may depend on site factors (climate, soils and nutrient availability) and the traits of

the species under consideration. As such, investigations of multiple tree species (with divergent

traits) growing in a common soil may help resolve this apparent paradox.

Tree species also exhibit varying degrees of plasticity in terms of their root traits (Weemstra &

Valverde-Barrantes, 2022), which could influence exudation dynamics. Traits like SRL and

branching intensity (BI) are associated with nutrient acquisition (Comas & Eissenstat, 2009), and

tend to be more plastic than traits linked to structural stability and longevity, such as RTD and root

diameter (Comas et al., 2012, 2014; Comas & Eissenstat, 2009; Sun et al., 2021). Whether

exudation rates display greater intraspecific variation than morphological root traits remains

unresolved (Sun et al., 2021), posing a key challenge for detecting the exudation-trait relationships.

Many root traits show a strong phylogenetic signal (Brundrett, 2002; Comas et al., 2012, 2014)

suggesting that exudation rates may differ among tree species with divergent evolutionary histories

(e.g., angiosperms *vs.* gymnosperms) and distinct mycorrhizal associations (e.g., arbuscular *vs.*

ectomycorrhizal associations; AM *vs.* EcM). Moreover, if exudation patterns evolved as a means

for dealing with nutrient limitations, links between tree species' evolutionary history, root traits

and exudation might be expected. Early gymnosperms had thick, dense, long-lived roots that

associated with 'ancestral' AM fungi (Brundrett, 2002; Comas et al., 2012). As greater water and

nutrient limitations emerged and selected for gymnosperms with highly-branched roots colonized

by fungi derived from saprotrophs (i.e., EcM fungi), high exudation rates may have represented

an additional strategy for nutrient acquisition (Brundrett, 2002; Read & Perez-Moreno, 2003).

When angiosperms arose in the early Cretaceous, some species evolved thin diameter, highly

proliferative roots (Brundrett, 2002; Comas et al., 2012; Guo et al., 2008) whereas others - often

in nutrient-poor soils - developed EcM associations (Comas et al., 2012; Read & Perez-Moreno,

2003). Whether exudation rates relate to tree species' belowground C allocation and nutrient

acquisition strategies is unknown, yet there are reasons to suspect that the evolutionary processes

that shape root trait syndromes and tree-mycorrhizal associations also affect exudation.

There has been little consensus over whether exudation rates differ among tree species from

different functional groups (Brzostek et al., 2013; Liese et al., 2018; Wang et al., 2021). Exudation

could be greater in EcM trees (relative to AM trees) if exudation is a reflection of the C sink

strength of roots, which is typically greater in EcM root systems (Hobbie, 2006). Alternatively, if

exudation rates reflect C allocation tradeoffs within the root system (e.g., C exuded by roots comes

at a cost to C used to support mycorrhizal fungi; Wen et al., 2019), one might expect higher

exudation in AM trees where the C costs of supporting AM hyphae are low relative to EcM

mycelium (Hawkins et al., 2023). To date, support for both hypotheses is apparent. In temperate

forests, EcM trees have been shown to exude more C than AM trees (Brzostek et al., 2013; Phillips

& Fahey, 2006; Yin et al., 2014), though this effect is not apparent in young trees (Liese et al.,

2018) and the opposite pattern has been reported in sub-tropical forests (Sun et al., 2021). Likewise,

deciduous trees have been shown to have higher exudation rates than evergreen trees in some

temperate forests (Sun, et al., 2017; Wang et al., 2021) but not others (Brzostek et al. 2013). In a

recent synthesis of dozens of studies, Chari et al. (2024) found no evidence of exudation

differences between angiosperms and gymnosperms or between AM and EcM trees. These mixed

findings highlight the need to investigate tree functional group effects on exudation at a common

site where other factors (climate, tree age, soil characteristics, etc.) can be controlled for.

In this study, we assessed the effects of tree species, functional groups, and root traits on exudation

rates in mature trees grown in monodominant plots in a common soil. Importantly, to disentangle

the effects of functional groups (e.g., AM and EcM *vs.* angiosperms and gymnosperms) from root

traits across the RES, we selected tree species from each functional group that spanned a range of

root trait space. Our objectives were to (1) characterize the extent to which exudation rates vary

among tree species and across functional groups, (2) determine which root traits in the RES, if any,

are closely related to exudation rates, and (3) build a framework for predicting exudation using

readily-measurable root traits and tree functional groups. We hypothesized that exudation rates

would differ among tree species and functional groups due to differences in root traits, leading to

the prediction that considering both root traits and functional groups would better predict exudation

rates (H1). Additionally, we hypothesized that exudation is linked to one or more axes of the RES

(H2): (a) congruent to the collaboration gradient (leading to the prediction that exudation correlates

positively with SRL or SRA) or (b) congruent to the conservation gradient (leading to the

prediction that exudation correlates negatively to RTD).

Materials and methods

Site description

This study was conducted in monoculture plots at the Morton Arboretum, Lisle, Illinois (41.81N,

88.05W). The plots were established between 1922 and 1948 to test and study "all the timber trees

of the world which might come under consideration for reforestation purposes in this part of the

country" (Morton Arboretum Staff, 1929). Soils in the plots are poorly drained Alfisols that form

from a thin layer of loess (0.31 m) underlain by glacial till and Mollisols that formed from alluvium

(Soil Survey Staff, NRCS, USDA, 2024). The soil series in the plots are primarily Ozaukee silt

loams and Sawmill silty clay loam (Midgley & Sims, 2020). The area has a continental climate

with temperatures ranging from -6°C in January to 22°C in July and 800-1,000 mm mean annual

precipitation.

Eleven tree species were selected to capture the heterogeneity in root traits among species from

distinct functional groups: phylogenetic group (angiosperm *vs.* gymnosperm), tree-mycorrhizal

association (AM *vs.* EcM), and leaf habit (deciduous *vs.* evergreen). Within each group, species

were chosen based on mean SRL and root tissue N concentration (root N) - the traits that were

found to correlate positively with exudation rates in previous studies (Meier et al., 2020; Sun et

al., 2021; Wang et al., 2021). As such, selected eleven species spanned a wide range of SRL and

root N for each group, ensuring that all selected species captured diverse trait space (Table 1). This

allowed for minimizing phylogenetic covariations among traits while maximizing species trait

dissimilarities. Out of the eight combinations, only two combinations were absent: evergreen-AM-

angiosperms and evergreen-EcM-angiosperms (Table 1).

Root exudation rates

Fine-root exudates were collected during the growing season of 2022 (i.e., from May to July 2022)

using an *in-situ* culture-based cuvette system (Phillips et al., 2008). To mitigate the impact of

variable weather, sampling campaigns were conducted under sunny and clear conditions, to the

extent possible, and each plot was visited twice: once in late May/early June to collect exudates

from three individuals and once in late June/early July to collect exudates from 3-4 additional

individuals. The terminal roots were excavated carefully from the mineral topsoil below the

organic layer. The excavated root segments were examined to ensure that the fine-root system

consisted of the first three branching orders with an intact absorptive function. Organic matter and

soil particles adhering to the root system were removed with DDI water with extreme caution while

keeping the roots moist with wet paper towels. In cases where the distal fine roots were damaged

or broken off, samples were discarded, and a new sample was prepared. The intact root systems

were placed in cuvettes (30mL syringe) filled with sterile, C-free glass beads (>1mm diameter).

The root systems with glass beads were flushed three times with C-free nutrient solution (0.5 mM

NH_4NO_3, 0.1 mM KH_2PO_4, 0.2 mM K_2SO_4, 0.15 mM $MgSO_4$, 0.3 mM $CaCl_2$) to ensure the root

segments and glass beads were well-mixed and to remove any C adhering to the root surface. To

ensure the same amount of solution was added to the cuvettes, we added 15mL of nutrient trap

solution in the field using a bottle-top dispenser. The cuvette was covered in aluminum foil to

allow the root system to equilibrate with the cuvette environment. The same procedure was applied

to the control (i.e., no root) cuvette with the same glass beads and nutrient solution. The cuvettes

were placed at the exact excavated area and covered with soils and organic matter and incubated

for approximately 24 hrs.

After the one-day incubation period, the sampled roots with the cuvette were clipped with care and

brought to the laboratory for analysis. Within one hour of clipping, each cuvette was flushed with

15mL of the working nutrient solution three times to remove accumulated exudates in the cuvette.

All solutions were filtered immediately through a sterile 0.22 μm syringe filter (Millex-GV 0.22μm

PVDF 33mm Gamma Sterilized 50/Pk, Millipore Co., Billerica, MA) and refrigerated at 4°C until

analyses (<24 h). All samples were analyzed for non-particulate organic C on a TOC analyzer

(Shimadzu Scientific Instruments, Columbia, MD) within a day of sample collection. The total

mass-specific exudation rate was calculated with the total C captured from the trap solution minus

the total C flushed from the root-free control cuvettes divided by the dry root biomass and day (mg

$C * g_{root}^{-1} * day^{-1}$).

Root morphological and chemical traits

Roots originally placed in the cuvette were carefully collected from the cuvette, washed, and stored

at 4°C until processing. Fine-root morphology was analyzed for all the fine roots with a transparent

flat-bed scanner and the WinRHIZO program (Regent Instruments, Quebec, QC, Canada). Scans

were collected at a resolution of 600 dpi. All root samples were dried at 65°C for at least 48 h, and

the dried root biomass was used for root trait calculations. Specific root length (SRL, in m g^{-1} : the

length of the fine roots divided by the corresponding root dry weight), specific root area (SRA, in

cm^2 g^{-1} : the area of the fine roots divided by the corresponding root dry weight), root tissue density

(RTD, in g cm^{-3} : root dry weight divided by root volume), root branching intensity (BI, in the

number of tips per total fine-root length), and root diameter (diameter, in cm) were calculated from

WinRHIZO. Root N concentration (per dry weight) was measured independently in the lab using

an elemental combustion system (Costech Analytical Technologies).

Statistical analyses

We used an analysis of variance (ANOVA), mixed linear models, and variance partitioning to

characterize the extent to which root exudation rates vary among tree species and across functional

groups. To test for differences in exudation rates among tree species, we conducted pairwise

comparisons after an ANOVA using a Tukey's Honest Significant Difference (HSD) test. To test

for differences in exudation rates among tree functional groups, we built a mixed linear model with

mycorrhizal type, phylogeny, and their interaction as fixed effects and species-plot as a random

effect using restricted maximum likelihood ('lme4::lmer' via REML). To evaluate the significance

of each nested group in the model after accounting for all other groups, Type III ANOVA with

Satterthwaite's Method using the 'lme4::anova' was performed to summarize the results of each

model. To control the likelihood of false positives in all linear mixed effects models, adjusted p-

values from BH Correction (Benjamini-Hochberg) test were performed using the p.adjust function.

To quantify the contributions of inter- vs. intraspecific variation to exudation rates in mixed effects

models, a variation partitioning analysis was performed using the 'VEGAN::varpart'. To show co-

variations among root traits, a pairwise trait relationships between exudation rates and root traits

were also performed using Pearson's correlations at the individual tree level using 'corr.test'

function. Root traits and exudation rates were natural-log-transformed prior to analyses to meet

model assumptions of residual normality and homogeneity of variance.

To assess how and the extent to which root exudation rates are associated with root trait

coordination, we used principal components analysis (PCA) (Weigelt et al., 2023) and Redundancy

Analysis (RDA). To examine how exudation rates aligns with major dimensions in the PCA, we

created an ordination of RTD, root N, SRL, Diameter, SRA, and BI along with exudation rates

using princomp () with standardized PCA. To examine the significance of linear relationships
between exudation rates and the first four axes, we created a PCA without exudation rates and
preformed Pearson's product-moment correlation test between PCs and exudation using cor.test
function. To select the best predicting root trait or subset of predictors, we built a PCA with four
core variables (RTD, root N, SRL, and Diameter) and evaluated the relationship between root
exudation rates as a trait and the traits that comprise the PCA. We used RDA models for PCs to
explain exudation using 'VEGAN::rda' and selected the best predicting trait using
'VEGAN::ordistep' with both forward and backward stepwise model selection.

To identify the functional groups and root traits that collectively best predict exudation rates, we
used a stepwise model selection approach using linear mixed-effects models by 'lme4::lmer' via
REML. The fixed effects included six root traits (RTD, root N, SRL, Diameter, SRA, and BI) along
with mycorrhizal type or phylogenetic group. Monodominant plot identity (i.e., species-plot) was
treated as a random effect. Model selection was based on improvements in Akaike Information
Criterion (AIC) and likelihood ratio tests comparing full and reduced models. Building on the best-
performing model, we further tested interactions between traits and functional groups (e.g.,
Exudation ~ Trait × Functional Group). We examined the explanatory power of each model by
calculating marginal (R²m) and conditional (R²c) R-squared values, where R²m represents variance
explained by fixed effects and R²c includes both fixed and random effects (Nakagawa & Schielzeth,
2013). Model assumptions for selected models were verified via checks for residual normality,
homoscedasticity, and unbiasedness. All statistical analyses were performed using R v.3.5.3 (R
Core Team, 2017).

Results

Hypothesis 1: Exudation differences among species and functional groups

Variation in exudation rates among tree species and functional groups
We found partial support for H1, as species explained 22% of the total variation in exudation rates
in our model (Adj. R^2 = 0.22; p = 0.01). While not all species differed in their exudation rates
(ANOVA using Tukey's HSD test), *Larix gmelinii* exhibited significantly higher rates of root

exudation compared to *Chamaecyparis pisifera* (p=0.03) and *Carya ovata* (p<0.01) (Fig. 1a; Table 2). The mean exudation rate of *L. gmelinii* (3.82 mgC g_{root}^{-1} * day^{-1}) was more than twice that of the second-highest species, *Picea abies* (1.52 mgC g_{root}^{-1} * day^{-1}) (Fig. 1a: Table 2).

We also found a significant interaction between mycorrhizal type and phylogenetic group in exudation rates (Chisq = 9.27, Adj. p < 0.01; Table 3). On average, EcM gymnosperms exuded 2.4 times more C than EcM angiosperms and 1.5 times more C than AM gymnosperms (Fig. 1b). Despite these notable differences, tree mycorrhizal type, phylogenetic group, and leaf habit alone did not significantly explain variation in exudation (Chisq $\leq$ 0.98, Adj. p $\geq$ 0.48; Table 3). Additionally, we did not detect a significant interaction between mycorrhizal association and leaf habit (Chisq = 1.90, Adj. p =0.34; Table 3).

Exudation rates exhibited over twice the variability of most root traits with a coefficient of variation (CV%) of 119% compared to lower variability across root traits (Table S1). This higher CV% corresponded to a high intraspecific variation (i.e., 77%) (Fig. S9; Table S3). Most root traits had CV%s below 40%, except for BI at 44% (Table S1), suggesting morphological traits are generally more conservative (i.e., less plastic) than exudation. High interspecific variation was observed in most root traits (RTD, root N, root diameter, and BI), whereas 'composite' and 'acquisitive' root traits such as SRL and SRA showed high intraspecific variability (>50%) and intermediate CV%s (38% and 30%, respectively) (Fig S9; Table S3). Together, unlike most root traits, exudation rates in our study can only be partially explained by species.

Hypothesis 2: Exudation and the root economic space (RES)

The first two principal axes of the PCA generated with seven core root traits (RTD, root N, SRL, Diameter, SRA, BI, and exudation rates) explained 73% of the total variation in root traits (Fig. 2a; Table S5). The axis generated by RTD-root N was closely mapped onto PC1 reflecting the conservation gradient, while the axis generated by SRL-Diameter was loaded closely onto PC2 representing the collaboration gradient (Bergmann et al., 2020; Fig. 2a). The first five PCs demonstrated eigenvalues exceeding those predicted by random chance (as determined by Broken Stick analysis; Fig. S6), indicating that these axes accounted for more variance than would be

expected under random conditions. In addition, the first three PCs showed eigenvalues greater than

1.0, indicating the significant contribution of three PCs to the ordination (Table S5; Tabachnik &

Fidell, 1996).

Redundancy analysis (RDA) with permutation tests indicated that exudation rates were not

significantly associated with variation along PC1 and PC2, which were generated with six root

traits excluding exudation rates (p>0.15; Fig. S7). However, exudation rates were significantly

correlated with PC3 ($R^2 = 0.08$, p=0.03) and PC4 ($R^2 = 0.10$, p=0.02), while the two PCs accounted

for 10% and 6% of the total variation, respectively (Fig. S7). PC3 and PC4 together explained a

greater proportion of variance in exudation (exudation ~ PC3 + PC4; $R^2 = 0.18$, p=0.004). Notably,

stepwise model selection based on RDA of the PCA derived from four core root traits (SRL–

Diameter, RTD–Root N; Fig. S4) identified RTD as the only significant predictor of exudation

(AIC = 3.56, F = 5.05, p=0.015). These results suggest that exudation likely correlates with the

conservation gradient, but also suggest that root exudation is more strongly associated with trait

variation captured by more than just the first two principal components in the RES.

The effects of tree functional groups on trait-exudation relationships

Mixed-effect model predictions

Consistent with the prediction derived from the first hypothesis, incorporating both root traits and

tree functional groups enhanced model predictions of exudation. As such, the best-performing

mixed-effects model included SRA and the interaction term between the phylogenetic group and

mycorrhizal types (exudation~ SRA + mycorrhizal-type:phylogeny + (1|species); $R^2c = 0.36$;

p<0.01; Fig. 3; Table 4). That is, tree species with higher SRA - indicative of more acquisitive root

strategies - exhibited significantly higher exudation rates (Std β = 1.13 ± 0.39, Adj. p=0.013; Fig.

3; Table 4). However, the interaction between mycorrhizal-type and phylogeny modified this

relationship: compared to the baseline group (EcM-Gymnosperms), all other combinations (AM-

Angiosperms, EcM-Angiosperms, and AM-Gymnosperms) showed significantly lower exudation

rates (Adj. p=0.01; Fig. 1b). A larger model that additionally included root N also significantly

predicted exudation rates (p<0.01; Table S4), albeit with a slightly reduced fit. These results

supported our second hypothesis, suggesting that exudation rates might be influenced by both acquisitive and conservative root traits, as captured by variation in SRA and root N.

Exudation-trait correlations in mixed effects models

Linear mixed effects models that included significant effects of mycorrhizal association on exudation provide partial support for exudation as both an acquisitive resource exploitation strategy and a physiological process governed by C allocation tradeoffs. Across all species, variation in SRL, and SRA, and partly in root diameter, significantly accounted for variation in root exudation (Fig. 4; Table 4). The negative relationship of exudation with RTD and positive relationship with root diameter were modulated by mycorrhizal association (Fig. 4; Table 4), while phylogenetic group showed little effects on trait-exudation relationships (Table S2).

Specifically, high SRL and SRA were positively correlated with exudation rates across species (SRL: Std β = 1.5, Adj. p<0.001; SRA: Std β = 1.8, Adj. p=0.03; Table 4). Unlike SRA, this positive relationship between exudation and SRL appeared stronger among AM trees, as indicated by a marginally significant linear relationship in a simple linear model nested in mycorrhizal association (R^2 = 0.13, p=0.051; Fig 4b). The marginally significant interaction term (SRL × Mycorrhizal Type, Std β = -1.09, Adj. p=0.11; Table 4) suggests that EcM association may modulate this relationship, potentially exhibiting a weaker or even negative association between SRL and exudation compared to AM trees. However, further study is needed to confirm these effects, especially since the relationship was not detected in the bi-variate analysis (Pearson's correlation 'r'; Fig S1).

While no trend between exudation and RTD across species was detectable, RTD and exudation showed a significant interaction (Std β = -2.0, Adj. p=0.001, Table 4), showing that in ECM trees, increasing RTD was associated with a large decrease in exudation rates (Fig. 4d), with this relationship being more pronounced in gymnosperms (Table 3; Fig 1b). Exudation generally decreased with root diameter across species with marginal significance (Std β = -1.5, Adj. p=0.06; Fig. 4a; Table 4). Notably, EcM trees exhibited higher exudation rates than AM trees after controlling for diameter (Std β = 2.9, adjusted p=0.01; Fig 4a; Table 4), leading to a significant interaction between tree mycorrhizal association and diameter. Together, exploitative root traits

such as SRA and SRL associated positively with exudation across species while EcM association

significantly influenced the relationships of exudation with RTD, SRA and root diameter.

Discussion

Our study aimed to identify key drivers of mass-specific root exudation in mature field-grown

trees. We hypothesized that exudation rates would differ among tree species and functional groups

(H1). Further, we hypothesized that exudation rates would be associated with at least one of the

axes of the RES (H2): along the collaboration gradient of the RES owing to exudation's functional

role as a nutrient acquisition strategy and/or along the conservation gradient of the RES owing to

exudation's role as a competing sink for C (e.g., root tissue construction costs). We found partial

support for H1, as exudation rates varied partly among tree species, and EcM gymnosperms

exhibited greater exudation rates than other tree functional groups. In partial support of our second

hypothesis, exudation rates correlated positively with SRL and SRA (across all species) and in

EcM trees, correlated with conserved traits such as root diameter and RTD. However, while

exudation rates were loaded weakly onto the 'fast' side of the conservative gradient in the RES,

exudation was not correlated with the first two axes. Rather, exudation was better predicted by

independent third and fourth (i.e., non-RES) axes. Finally, we found that the best model to predict

exudation rates across all 11 species contained SRA (with a significant correlation with SRL)

coupled with a strong influence from mycorrhizal type on phylogeny. Collectively, our study

indicates that root exudation is a complex physiological process and cannot be fully explained by

species identity or root traits alone. Instead, a combined consideration of these factors offers a

more accurate prediction of fine-root physiological functioning.

Tree functional groups partially account for variation in exudation

EcM gymnosperms generally had the highest exudation rates (two-fold higher than other groups),

though the reasons remain elusive. High exudation rates have been reported for other EcM

gymnosperms in temperate forests (Abramoff & Finzi, 2016; Brunn et al., 2022; Jiang et al., 2021;

Yang et al., 2020; Zhang et al., 2016) and under controlled conditions (Sell et al., 2022; Yin et al.,

2013). One of the highest exuders in our study, *Larix gmelinii*, has been reported to have elevated

exudation rates in a boreal forest as well (Yin et al., 2023). Although the influence of EcM

associations on root traits is well documented, their differential impacts on root traits and exudation

between angiosperms and gymnosperms remain unclear.

Three potential mechanisms driving this pattern include tradeoffs between C allocation to EcM

fungi and root exudates, leakiness of highly colonized gymnosperm roots, and high exudation rates

of EcM fungi themselves. First, EcM gymnosperms might allocate more C to the rhizosphere and

lesser to their mycorrhizal partners. For instance, a global synthesis by Hawkins et al. (2023)

reported that C allocation to EcM mycelium biomass was ~2.5 times greater in broad-leaved trees

(mostly angiosperms) than needle-leaf trees (mostly gymnosperms). However, the Hawkins et al.

(2023) synthesis did not account for biome differences, suggesting that reports of lesser C

allocation to mycelium in EcM gymnosperms may have been confounded by climate (e.g., if boreal

forest needle-leaf trees were compared to temperate/tropical forest broad-leafed trees). In fact,

some of the most striking examples of prodigious EcM mycelium come from studies of EcM

gymnosperms like *Pinus spp.* (Anderson & Cairney, 2007), and EcM gymnosperms tend to possess

short, thick roots (Fig S8) that are well-colonized by mycorrhizal fungi (Cheng et al., 2016; Comas

& Eissenstat, 2009; Ma et al., 2018). As such, a second factor that could explain greater exudation

in EcM gymnosperms is mycorrhizal-root leakiness. Little is known about whether the mycorrhizal

roots of EcM gymnosperms differ from those of EcM angiosperms in leakiness (Farrar et al., 2003).

If the magnitude of leakiness is determined by the strength of the C sink - as described by the "hole

in the pipe model" for nitrous oxide gas in Firestone and Davidson (1989) - greater exudation in

EcM gymnosperms would result from greater C allocation to mycelium. Finally, while EcM fungi

are a sink for plant-derived C (Prescott et al., 2020), they too can exude organic C (e.g., oxalic

acid), resulting in an additional source of C in the cuvettes (Ahonen-Jonnarth et al., 2000; Sun et

al., 1999; Van Schöll et al., 2008). Further studies are required to elucidate the mechanisms that

drive high exudation rates of EcM gymnosperms.

The relationships between exudation and root traits that comprise the RES

Exudation rates were aligned weakly with the fast side of the 'conservation' gradient defined by

RTD and root N in the RES while also showing a strong association with SRA (Sun et al., 2021).

These results suggest that exudation may be linked to acquisitive root traits (e.g., high SRA and

root N) to optimize soil resource uptake (Hypothesis 2a; Eissenstat, 1991; Eissenstat & Yanai,

1997; Lv et al., 2023; Weemstra et al., 2016). These patterns may also reflect a crucial root-soil process where rapid exudation increases soil N availability (Meier et al., 2017), thereby enhancing root metabolic activity characterized by low RTD and high root N (Sun et al., 2021; Wen et al., 2022). We also found that RTD was the single significant predictor for exudation in the RES, partially supporting the prediction that exudation could correlate with root traits related to the conservation gradient reflecting structural stability, longevity, and metabolic activity such as RTD and root N (Hypothesis 2b). The observed significant relationships of exudation with RTD (from RDA analysis) and SRA (from mixed-effects models) likely result from the interplay of root area, tissue permeability, and diffusion strength - all of which can influence exudation rates (Akatsuki & Makita, 2020; Farrar et al., 2003; Jones et al., 2009; Lv et al., 2023; Sun et al., 2021). Since higher RTD often is association with reduced root leakiness and metabolic activity (Farrar et al., 2003), roots with less dense tissues, high root N, and larger surface areas are more likely to exude C. Consistent with our findings, recent studies in cool-temperate and subtropical forests have also reported significant relationships of exudation rates with RTD (-), root N (+), and SRA (+), but not SRL (Akatsuki & Makita, 2020; Sun et al., 2021).

The alignment of exudation within the RES suggests that exudation may function as an alternative soil resource exploitation strategy (Sun et al., 2021; Wen et al., 2022), modulated by the interplay between mycorrhizal associations (Brzostek et al., 2013) and root traits under phylogenetic constraints (Williams et al., 2022). However, exudation rates were not associated with the first two PC axes, and additional PC axes (PC3 and PC4; Fig. 2b) were needed to explain variation in exudation. This suggests that exudation may be more strongly associated with traits not considered or other factors. The RES simplifies trait space by collapsing trait variation into 'collaboration' and 'conservation' gradients (Bergmann et al., 2020), yet other dimensions of plant trait variation (e.g., plant height and rooting depth) were found to exist (Weigelt et al., 2021). Whether exudation aligns with this third axis of plant variation is unknown (Freschet et al., 2021; McCormack et al., 2017; Weemstra et al., 2022) or links to other unmeasured processes (e.g., C assimilation and allocation) is not well-understood, but this would be a fruitful line of inquiry in future studies.

Trait-to-exudation relationships constrained by functional group interactions

Exudation rates were significantly linked to acquisitive root traits such as high SRA, high SRL,

low RTD, while their variability was strongly influenced by EcM association, root morphology,

and C construction costs. Notably, the model that included the interaction between mycorrhizal

association and phylogenetic lineage in predicting exudation–SRA relationships outperformed all

others (Table 4). These findings collectively suggest that exudation may be linked to acquisitive

strategies (e.g., high SRL and SRA, narrow diameters, and low RTD) that favor foraging efficiency

under the control of both mycorrhizal association and evolutionary history (Hypothesis 2a;

Eissenstat, 1991; Eissenstat & Yanai, 1997; Lv et al., 2023; Weemstra et al., 2016).

Interestingly, we observed that EcM association influenced the relationship between exudation and

diameter, where exudation rates increased with root diameter. This trend was predominantly driven

by EcM gymnosperms, which exhibited short, thick roots (Fig. S8) and high exudation rates (Fig.

1b) - highlighting their reliance on symbiotic partners for nutrient acquisition. However, a previous

study measuring exudation rates of EcM gymnosperms (*Pinus* and *Larix spp.*) found a significant

negative relationship between exudation rates and root diameter (Akatsuki & Makita, 2020). These

divergent results suggest that while the associations with EcM do broadly affect plant exudation

rates, the direction of those relationships likely depends on several factors, including species

identity, trait variations and methodological differences (Williams et al., 2021). For example,

Akatsuki & Makita (2020) used a glass-fiber filter method that targeted smaller root segments (<

3rd order), potentially capturing higher mass-specific exudation rates than the cuvette-base

methods of Phillips et al. (2008). This suggests that EcM gymnosperms exude root carbon mostly

from lower-order roots and that our cuvette-base methods may underestimate mass-specific

exudation rates owing to inclusion of larger root segments (Akatsuki & Makita, 2020). Together,

this highlights the complexity of trait-to-exudation relationships and the need for further studies to

uncover generalizable patterns across diverse ecosystems.

Unlike EcM trees, which exhibit differences in some root traits across phylogenies, AM association

did not significantly predict most root traits and trait-exudation relationships in our study (Fig. S8)

(Akatsuki & Makita, 2020). Instead, AM angiosperms demonstrated greater within-group

variability, including exudation rates (Fig. S8), suggesting a flexible, 'do-it-yourself' resource

acquisition strategy with high plasticity (Bergmann et al., 2020). Together, these findings indicate

that elevated exudation rates can result from (1) the interplay of EcM-associated root traits and

phylogenetic constraints or (2) enhanced intraspecific variability in acquisitive root traits, allowing

for efficient soil resource exploitation with moderate exudation rates.

Challenges and opportunities

Exudation rates for mature, field-grown trees are rare, as all methods used to measure exudation

introduce some artifacts; our study is no exception. While the root cuvette method of Phillips et al.

(2008) has been used in dozens of forest studies (Chari et al., 2024), important methodological

questions remain about how to minimize damage to root systems before placement into cuvettes,

how to normalize the rates measured in different scales, the chemistry and sterility of the trapping

solution, the duration of the equilibrium period, the duration of the collection period, and the

contribution of mycorrhizal fungi to the total exudates (Oburger & Jones, 2018). Moreover, the

high degree of spatio-temporal variation in exudation (Jacoby et al., 2017; Jiang et al., 2021; Yang

et al., 2020) affects how many samples are needed to draw inferences about species-specific

patterns and trait-exudation correlations. While we attempted to minimize variability by selecting

tree species grown in a common soil, applying the same methodology to all trees, etc., the

substantial amount of variability in exudation within species suggests that this flux may be better

linked to dynamic physiological processes such as C assimilation and allocation than to (relatively)

static morphological traits.

Conclusion

We revealed that root exudation was positively associated with acquisitive root traits (SRL and

SRA), while exudation-trait relationships were modulated by mycorrhizal association and

phylogenetic lineage. EcM trees appeared to influence trait–exudation relationships, especially for

RTD and root diameter, with these effects being more pronounced in gymnosperms. As such,

constraining trait-exudation relationships with tree functional groups did improve model

predictability. Importantly, we demonstrate that root exudation may be a complex physiological

process that cannot be explained by species identity, functional groups or individual root traits

alone. High intraspecific variation in root exudation (unlike stable morphological traits) likely

contributed to the weaker alignment of exudation with the RES. Instead, exudation was linked to

additional functional axes beyond the first two axes in the RES. Given the emerging interest in including root physiological traits into the RES (which is based mostly on morphological and chemical traits), our findings suggest that incorporating such dynamic processes into the RES may pose significant challenges and require to identify additional drivers of such dynamics other than root traits. As more mechanistic studies are developed (e.g. tracking exudate responses to tree girdling or nutrient solution culture alterations) and methods for capturing exudates are improved, our ability to understand the role and function of exudates in the environment should come into greater focus.

Acknowledgements

This study was supported by the Center for Tree Science fellowship from The Morton Arboretum and Indiana University-Bloomington. Funding to YO and RPP was provided by NSF, DEB, MacroSysBIO & NEON-Enabled Science (Award# 2106096). We thank Elizabeth Huenupi for facilitating lab work in Indiana University and The Morton Arboretum Soils Lab REU students for lab assistance. We are grateful for constructive comments provided by Phillips Lab members.

Competing interests

The authors declare that they have no conflict of interest.

Author contributions

M.G.M., K.V.B., M.L.M., Y.E.O. and R.P.P. designed the study; M.G.M., K.V.B., M.L.M., Y.E.O., M.M., R.K.B., and S.H. performed the research; M.G.M., M.M., and Y.E.O. analyzed the data; Y.E.O., M.G.M. and R.P.P. wrote the paper with input from all authors.

References

Abramoff, R. Z., & Finzi, A. C. (2016). Seasonality and partitioning of root allocation to rhizosphere soils in a midlatitude forest. *Ecosphere*, *7*(11), e01547. https://doi.org/10.1002/ecs2.1547

Ahonen-Jonnarth, U., Van Hees, P. A. W., Lundström, U. S., & Finlay, R. D. (2000). Organic acids produced by mycorrhizal *Pinus sylvestris* exposed to elevated aluminium and heavy metal concentrations. *New Phytologist*, *146*(3), 557–567. https://doi.org/10.1046/j.1469-8137.2000.00653.x

Akatsuki, M., & Makita, N. (2020). Influence of fine root traits on in situ exudation rates in four conifers from different mycorrhizal associations. *Tree Physiology*, *40*(8), 1071–1079. https://doi.org/10.1093/treephys/tpaa051

Anderson, I. C., & Cairney, J. W. G. (2007). Ectomycorrhizal fungi: Exploring the mycelial frontier. *FEMS Microbiology Reviews*, *31*(4), 388–406. https://doi.org/10.1111/j.1574-6976.2007.00073.x

Bardgett, R. D. (2014). *Going underground: Root traits as drivers of ecosystem processes*. *29*(12), 8.

Bergmann, J., Weigelt, A., & Laughlin, D. C. (2020). The fungal collaboration gradient dominates the root economics space in plants. *SCIENCE ADVANCES*, 10.

Brundrett, M. C. (2002). Coevolution of roots and mycorrhizas of land plants. *New Phytologist*, *154*(2), 275–304. https://doi.org/10.1046/j.1469-8137.2002.00397.x

Brunn, M., Hafner, B. D., Zwetsloot, M. J., Weikl, F., Pritsch, K., Hikino, K., Ruehr, N. K., Sayer, E. J., & Bauerle, T. L. (2022). Carbon allocation to root exudates is maintained in mature temperate tree species under drought. *New Phytologist*, *235*(3), 965–977. https://doi.org/10.1111/nph.18157

Brzostek, E. R., Fisher, J. B., & Phillips, R. P. (2014). Modeling the carbon cost of plant nitrogen acquisition: Mycorrhizal trade-offs and multipath resistance uptake improve predictions of retranslocation: Carbon cost of mycorrhizae. *Journal of Geophysical Research: Biogeosciences*, *119*(8), 1684–1697. https://doi.org/10.1002/2014JG002660

Brzostek, E. R., Greco, A., Drake, J. E., & Finzi, A. C. (2013). Root carbon inputs to the rhizosphere stimulate extracellular enzyme activity and increase nitrogen availability in temperate forest soils. *Biogeochemistry*, *115*(1–3), 65–76. https://doi.org/10.1007/s10533-012-9818-9

Canarini, A., Kaiser, C., Merchant, A., Richter, A., & Wanek, W. (2019). Root Exudation of Primary Metabolites: Mechanisms and Their Roles in Plant Responses to Environmental Stimuli. *Frontiers in Plant Science*, *10*, 157. https://doi.org/10.3389/fpls.2019.00157

Chari, N. R., & Taylor, B. N. (2022). Soil organic matter formation and loss are mediated by root exudates in a temperate forest. *Nature Geoscience,* 15(12), 1011-1016.

Chari, N. R., Tumber-Dávila, S. J., Phillips, R. P., Bauerle, T. L., Brunn, M., Hafner, B. D., Klein, T., Obersteiner, S., Reay, M. K., Ullah, S., & Taylor, B. N. (2024). Estimating the global root exudate carbon flux. *Biogeochemistry*. https://doi.org/10.1007/s10533-024-01161-z

Cheng, L., Chen, W., Adams, T. S., Wei, X., Li, L., McCormack, M. L., DeForest, J. L., Koide, R. T., & Eissenstat, D. M. (2016). Mycorrhizal fungi and roots are complementary in foraging within nutrient patches. *Ecology*, *97*(10), 2815–2823. https://doi.org/10.1002/ecy.1514

Comas, L. H., Callahan, H. S., & Midford, P. E. (2014). Patterns in root traits of woody species hosting arbuscular and ectomycorrhizas: Implications for the evolution of belowground strategies. *Ecology and Evolution*, *4*(15), 2979–2990. https://doi.org/10.1002/ece3.1147

Comas, L. H., & Eissenstat, D. M. (2009). Patterns in root trait variation among 25 co-existing North American forest species. *New Phytologist*, *182*(4), 919–928. https://doi.org/10.1111/j.1469-8137.2009.02799.x

Comas, L. H., Mueller, K. E., Taylor, L. L., Midford, P. E., Callahan, H. S., & Beerling, D. J. (2012). Evolutionary Patterns and Biogeochemical Significance of Angiosperm Root Traits. *International Journal of Plant Sciences*, *173*(6), 584–595. https://doi.org/10.1086/665823

Eissenstat, D. M. (1991). On the relationship between specific root length and the rate of root proliferation: A field study using citrus rootstocks. *New Phytologist*, *118*(1), 63–68. https://doi.org/10.1111/j.1469-8137.1991.tb00565.x

Eissenstat, D. M., & Yanai, R. D. (1997). The Ecology of Root Lifespan. In *Advances in Ecological Research* (Vol. 27, pp. 1–60). Elsevier. https://doi.org/10.1016/S0065-2504(08)60005-7

Farrar, J., Hawes, M., Jones, D., & Lindow, S. (2003). How roots control the flux of carbon to the rhizosphere. *Ecology*, *84*(4), 827–837. https://doi.org/10.1890/0012-9658(2003)084[0827:HRCTFO]2.0.CO;2

Fei, S., Desprez, J. M., Potter, K. M., Jo, I., Knott, J. A., & Oswalt, C. M. (2017). Divergence of species responses to climate change. *Science Advances*, *3*(5), e1603055. https://doi.org/10.1126/sciadv.1603055

Finzi, A. C., Abramoff, R. Z., Spiller, K. S., Brzostek, E. R., Darby, B. A., Kramer, M. A., & Phillips, R. P. (2015). Rhizosphere processes are quantitatively important components of terrestrial carbon and nutrient cycles. *Global Change Biology*, *21*(5), 2082–2094. https://doi.org/10.1111/gcb.12816

Firestone, M. K., & Davidson, E. A. (1989). Microbiological basis of NO and N 2 O production and consumption in soil. Exchange of trace gases between terrestrial ecosystems and the *atmosphere*, 47, 7-21.

Freschet, G. T., Roumet, C., Comas, L. H., Weemstra, M., Bengough, A. G., Rewald, B., Bardgett, R. D., De Deyn, G. B., Johnson, D., Klimešová, J., Lukac, M., McCormack, M. L., Meier, I. C., Pagès, L., Poorter, H., Prieto, I., Wurzburger, N., Zadworny, M., Bagniewska-Zadworna, A., … Stokes, A. (2021). Root traits as drivers of plant and ecosystem functioning: Current understanding, pitfalls and future research needs. *New Phytologist*, *232*(3), 1123–1158. https://doi.org/10.1111/nph.17072

Guo, D., Xia, M., Wei, X., Chang, W., Liu, Y., & Wang, Z. (2008). Anatomical traits associated with absorption and mycorrhizal colonization are linked to root branch order in twenty-three Chinese temperate tree species. *New Phytologist*, *180*(3), 673–683. https://doi.org/10.1111/j.1469-8137.2008.02573.x

Hawkins, H.-J., Cargill, R. I. M., Van Nuland, M. E., Hagen, S. C., Field, K. J., Sheldrake, M., Soudzilovskaia, N. A., & Kiers, E. T. (2023). Mycorrhizal mycelium as a global carbon pool. *Current Biology*, *33*(11), R560–R573. https://doi.org/10.1016/j.cub.2023.02.027

Hobbie, E. A. (2006). Carbon allocation to ectomycorrhizal fungi correlates with belowground allocation in culture studies. *Ecology*, *87*(3), 563–569. https://doi.org/10.1890/05-0755

Jacoby, R., Peukert, M., Succurro, A., Koprivova, A., & Kopriva, S. (2017). The Role of Soil Microorganisms in Plant Mineral Nutrition—Current Knowledge and Future Directions. *Frontiers in Plant Science*, *8*, 1617. https://doi.org/10.3389/fpls.2017.01617

Jiang, Z., Wang, Q., Xiao, J., Zhang, Z., & Yin, H. (2021). Differential responses of N benefit mediated by root exudate inputs to N addition between two subalpine forests. *Rhizosphere*, *19*, 100404. https://doi.org/10.1016/j.rhisph.2021.100404

Jo, I., Fei, S., Oswalt, C. M., Domke, G. M., & Phillips, R. P. (2019). Shifts in dominant tree mycorrhizal associations in response to anthropogenic impacts. *Science Advances*, *5*(4), eaav6358. https://doi.org/10.1126/sciadv.aav6358

Jones, D. L., Nguyen, C., & Finlay, R. D. (2009). Carbon flow in the rhizosphere: Carbon trading at the soil–root interface. *Plant and Soil*, *321*(1–2), 5–33. https://doi.org/10.1007/s11104-009-9925-0

Kumordzi, B. B., Aubin, I., Cardou, F., Shipley, B., Violle, C., Johnstone, J., Anand, M., Arsenault, A., Bell, F. W., Bergeron, Y., Boulangeat, I., Brousseau, M., De Grandpré, L., Delagrange, S., Fenton, N. J., Gravel, D., Macdonald, S. E., Hamel, B., Higelin, M., … Munson, A. D. (2019). Geographic scale and disturbance influence intraspecific trait variability in leaves and roots of North American understorey plants. *Functional Ecology*, *33*(9), 1771–1784. https://doi.org/10.1111/1365-2435.13402

Liese, R., Lübbe, T., Albers, N. W., & Meier, I. C. (2018). The mycorrhizal type governs root exudation and nitrogen uptake of temperate tree species. *Tree Physiology*, *38*(1), 83–95. https://doi.org/10.1093/treephys/tpx131

Lv, C., Wang, C., Li, Y., & Zhou, Z. (2023). Coordination among root exudation C, mycorrhizal colonization, and functional traits and their responses to drought in five temperate tree species. *Forest Ecology and Management*, *546*, 121316. https://doi.org/10.1016/j.foreco.2023.121316

Ma, Z., Guo, D., Xu, X., Lu, M., Bardgett, R. D., Eissenstat, D. M., McCormack, M. L., & Hedin, L. O. (2018). Evolutionary history resolves global organization of root functional traits. *Nature*, *555*(7694), 94–97. https://doi.org/10.1038/nature25783

McCormack, M. L., Guo, D., Iversen, C. M., Chen, W., Eissenstat, D. M., Fernandez, C. W., Li, L., Ma, C., Ma, Z., Poorter, H., Reich, P. B., Zadworny, M., & Zanne, A. (2017). Building a better foundation: Improving root-trait measurements to understand and model plant and ecosystem processes. *New Phytologist*, *215*(1), 27–37. https://doi.org/10.1111/nph.14459

McCormack, M. L., & Iversen, C. M. (2019). Physical and Functional Constraints on Viable Belowground Acquisition Strategies. *Frontiers in Plant Science*, *10*, 1215. https://doi.org/10.3389/fpls.2019.01215

Meier, I. C., Finzi, A. C., & Phillips, R. P. (2017). Root exudates increase N availability by stimulating microbial turnover of fast-cycling N pools. *Soil Biology and Biochemistry*, *106*, 119–128. https://doi.org/10.1016/j.soilbio.2016.12.004

Meier, I. C., Tückmantel, T., Heitkötter, J., Müller, K., Preusser, S., Wrobel, T. J., Kandeler, E., Marschner, B., & Leuschner, C. (2020). Root exudation of mature beech forests across a nutrient availability gradient: The role of root morphology and fungal activity. *New Phytologist*, *226*(2), 583–594. https://doi.org/10.1111/nph.16389

Midgley, M. G., & Sims, R. S. (2020). Mycorrhizal Association Better Predicts Tree Effects on Soil Than Leaf Habit. *Frontiers in Forests and Global Change*, *3*, 74. https://doi.org/10.3389/ffgc.2020.00074

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2 from generalized linear mixed-effects models. *Methods in ecology and evolution*, *4*(2), 133-142.

Niu, K., Zhang, S., & Lechowicz, M. J. (2020). Harsh environmental regimes increase the functional significance of intraspecific variation in plant communities. *Functional Ecology*, *34*(8), 1666–1677. https://doi.org/10.1111/1365-2435.13582

Norby, R. J., Loader, N. J., Mayoral, C., Ullah, S., Curioni, G., Smith, A. R., Reay, M. K., Van Wijngaarden, K., Amjad, M. S., Brettle, D., Crockatt, M. E., Denny, G., Grzesik, R. T., Hamilton, R. L., Hart, K. M., Hartley, I. P., Jones, A. G., Kourmouli, A., Larsen, J. R., … MacKenzie, A. R. (2024). Enhanced woody biomass production in a mature temperate forest under elevated CO2. *Nature Climate Change*, *14*(9), 983–988. https://doi.org/10.1038/s41558-024-02090-3

Oburger, E., & Jones, D. L. (2018). Sampling root exudates – Mission impossible? *Rhizosphere*, *6*, 116–133. https://doi.org/10.1016/j.rhisph.2018.06.004

Phillips, R. P., Erlitz, Y., Bier, R., & Bernhardt, E. S. (2008). New approach for capturing soluble root exudates in forest soils. *Functional Ecology*, *22*(6), 990–999. https://doi.org/10.1111/j.1365-2435.2008.01495.x

Phillips, R. P., & Fahey, T. J. (2006). TREE SPECIES AND MYCORRHIZAL ASSOCIATIONS INFLUENCE THE MAGNITUDE OF RHIZOSPHERE EFFECTS. *Ecology*, *87*(5), 1302–1313. https://doi.org/10.1890/0012-9658(2006)87[1302:TSAMAI]2.0.CO;2

Phillips, R. P., Finzi, A. C., & Bernhardt, E. S. (2011). Enhanced root exudation induces microbial feedbacks to N cycling in a pine forest under long-term CO2 fumigation: Rhizosphere feedbacks in CO2-enriched forests. *Ecology Letters*, *14*(2), 187–194. https://doi.org/10.1111/j.1461-0248.2010.01570.x

Read, D. J., & Perez-Moreno, J. (2003). Mycorrhizas and nutrient cycling in ecosystems – a journey towards relevance? *New Phytologist*, *157*(3), 475–492. https://doi.org/10.1046/j.1469-8137.2003.00704.x

Sell, M., Ostonen, I., Rohula-Okunev, G., Rusalepp, L., Rezapour, A., & Kupper, P. (2022). Responses of fine root exudation, respiration and morphology in three early successional tree species to increased air humidity and different soil nitrogen sources. *Tree Physiology*, *42*(3), 557–569. https://doi.org/10.1093/treephys/tpab118

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. (2024). *Web Soil Survey*. Retrieved April 22, 2025, from https://websoilsurvey.nrcs.usda.gov/

Sun, L., Ataka, M., Han, M., Han, Y., Gan, D., Xu, T., Guo, Y., & Zhu, B. (2021). Root exudation as a major competitive fine-root functional trait of 18 coexisting species in a subtropical forest. *New Phytologist*, *229*(1), 259–271. https://doi.org/10.1111/nph.16865

Sun, L., Ataka, M., Kominami, Y., & Yoshimura, K. (2017). Relationship between fine-root exudation and respiration of two Quercus species in a Japanese temperate forest. *Tree Physiology*, *37*(8), 1011–1020. https://doi.org/10.1093/treephys/tpx026

Sun, Y.-P., Unestam, T., Lucas, S. D., Johanson, K. J., Kenne, L., & Finlay, R. (1999). Exudation-reabsorption in a mycorrhizal fungus, the dynamic interface for interaction with soil and soil microorganisms. *Mycorrhiza*, *9*(3), 137–144. https://doi.org/10.1007/s005720050298

Tabachnik, B.G. and Fidell, L.S. (1996). Using multivariate statistics. 3rd ed. Harper Collins College. Publ. New York.

Tückmantel, T., Leuschner, C., Preusser, S., Kandeler, E., Angst, G., Mueller, C. W., & Meier, I. C. (2017). Root exudation patterns in a beech forest: Dependence on soil depth, root

morphology, and environment. *Soil Biology and Biochemistry*, *107*, 188–197. https://doi.org/10.1016/j.soilbio.2017.01.006

Van Schöll, L., Kuyper, T. W., Smits, M. M., Landeweert, R., Hoffland, E., & Breemen, N. V. (2008). Rock-eating mycorrhizas: Their role in plant nutrition and biogeochemical cycles. *Plant and Soil*, *303*(1–2), 35–47. https://doi.org/10.1007/s11104-007-9513-0

Wang, Q., Xiao, J., Ding, J., Zou, T., Zhang, Z., Liu, Q., & Yin, H. (2021). Differences in root exudate inputs and rhizosphere effects on soil N transformation between deciduous and evergreen trees. *Plant and Soil*, *458*(1–2), 277–289. https://doi.org/10.1007/s11104-019-04156-0

Weemstra, M., Freschet, G. T., Stokes, A., & Roumet, C. (2021). Patterns in intraspecific variation in root traits are species-specific along an elevation gradient. *Functional Ecology*, *35*(2), 342–356. https://doi.org/10.1111/1365-2435.13723

Weemstra, M., Kuyper, T. W., Sterck, F. J., & Umaña, M. N. (2022). Incorporating belowground traits: Avenues towards a whole-tree perspective on performance. *Oikos*. https://doi.org/10.1111/oik.08827

Weemstra, M. and Valverde-Barrantes, O. J. 2022. Above- and below-ground trait responses to environmental variation: the need to distinguish inter- and intraspecific variability. A commentary on 'Above and below-ground plant traits are not consistent in response to drought and competition treatments.' Ann. Bot.: mcac135. doi: 10.1093/aob/mcac135.

Weemstra, M., Mommer, L., Visser, E. J. W., Ruijven, J., Kuyper, T. W., Mohren, G. M. J., & Sterck, F. J. (2016). Towards a multidimensional root trait framework: A tree root review. *New Phytologist*, *211*(4), 1159–1169. https://doi.org/10.1111/nph.14003

Weigelt, A., Mommer, L., Andraczek, K., Iversen, C. M., Bergmann, J., Bruelheide, H., Fan, Y., Freschet, G. T., Guerrero-Ramírez, N. R., Kattge, J., Kuyper, T. W., Laughlin, D. C., Meier, I. C., Plas, F., Poorter, H., Roumet, C., Ruijven, J., Sabatini, F. M., Semchenko, M., … McCormack, M. L. (2021). An integrated framework of plant form and function: The belowground perspective. *New Phytologist*, *232*(1), 42–59. https://doi.org/10.1111/nph.17590

Weigelt, A., Mommer, L., Andraczek, K., Iversen, C. M., Bergmann, J., Bruelheide, H., Freschet, G. T., Guerrero-Ramírez, N. R., Kattge, J., Kuyper, T. W., Laughlin, D. C., Meier, I. C., Van Der Plas, F., Poorter, H., Roumet, C., Van Ruijven, J., Sabatini, F. M., Semchenko, M., Sweeney, C. J., … McCormack, M. L. (2023). The importance of trait selection in ecology. *Nature*, *618*(7967), E29–E30. https://doi.org/10.1038/s41586-023-06148-8

Wen, Z., Li, H., Shen, Q., Tang, X., Xiong, C., Li, H., Pang, J., Ryan, M. H., Lambers, H., & Shen, J. (2019). Tradeoffs among root morphology, exudation and mycorrhizal symbioses for phosphorus-acquisition strategies of 16 crop species. *New Phytologist*, *223*(2), 882–895. https://doi.org/10.1111/nph.15833

Wen, Z., White, P. J., Shen, J., & Lambers, H. (2022). Linking root exudation to belowground economic traits for resource acquisition. *New Phytologist*, *233*(4), 1620–1635. https://doi.org/10.1111/nph.17854

Williams, A., Langridge, H., Straathof, A. L., Fox, G., Muhammadali, H., Hollywood, K. A., Xu, Y., Goodacre, R., & de Vries, F. T. (2021). Comparing root exudate collection techniques: An improved hybrid method. *Soil Biology and Biochemistry*, *161*, 108391. https://doi.org/10.1016/j.soilbio.2021.108391

Williams, A., Langridge, H., Straathof, A. L., Muhamadali, H., Hollywood, K. A., Goodacre, R., & De Vries, F. T. (2022). Root functional traits explain root exudation rate and

748   composition across a range of grassland species. *Journal of Ecology*, *110*(1), 21–33.
749   https://doi.org/10.1111/1365-2745.13630
750 Yaffar, D., Cabugao, K. G., & Meier, I. C. (2022). Representing root physiological traits in the
751   root economic space framework. *New Phytologist*, *234*(3), 773–775.
752   https://doi.org/10.1111/nph.18070
753 Yang, L., Wang, X., Mao, Z., Jiang, Z., Gao, Y., Chen, X., & Aubrey, D. P. (2020). Root
754   Exudation Rates Decrease with Increasing Latitude in Some Tree Species. *Forests*, *11*(10),
755   1045. https://doi.org/10.3390/f11101045
756 Yin, H., Li, Y., Xiao, J., Xu, Z., Cheng, X., & Liu, Q. (2013). Enhanced root exudation stimulates
757   soil nitrogen transformations in a subalpine coniferous forest under experimental
758   warming. *Global Change Biology*, *19*(7), 2158–2167. https://doi.org/10.1111/gcb.12161
759 Yin, H., Wheeler, E., & Phillips, R. P. (2014). Root-induced changes in nutrient cycling in forests
760   depend on exudation rates. *Soil Biology and Biochemistry*, *78*, 213–221.
761   https://doi.org/10.1016/j.soilbio.2014.07.022
762 Yin, S., Wang, C., Lv, C., & Zhou, Z. (2023). Short-term responses of root traits and carbon
763   exudation to drought in a Larix gmelinii plantation. *Plant and Soil*, *484*(1–2), 393–405.
764   https://doi.org/10.1007/s11104-022-05800-y
765 Zhang, Z., Qiao, M., Li, D., Yin, H., & Liu, Q. (2016). Do warming-induced changes in quantity
766   and stoichiometry of root exudation promote soil N transformations via stimulation of soil
767   nitrifiers, denitrifiers and ammonifiers? *European Journal of Soil Biology*, *74*, 60–68.
768   https://doi.org/10.1016/j.ejsobi.2016.03.007
769

 Figures and tables

772
Fig 1) Boxplot of root exudation variation among: (a) 11 tree species and (b) combinations of mycorrhizal
type (AM vs. EcM) and phylogenetic group (Angiosperm vs. gymnosperm) at The Morton Arboretum,
Lisle, IL, USA. The central box in each boxplot represents the median and the interquartile range. The
whiskers extend to the minimum and maximum value. In (a), different lowercase letters denote significant
differences, as determined by a post hoc TukeyHSD test. In (b), different lowercase letters denote
significant differences (Linear mixed model fit by REML; Std β = 9.6, p=0.03). Closed circles denote
outliers.

(a) PCA of seven fine-root traits on an individual tree level

(b) PCA of seven fine-root traits (PC3 & PC4)

Fig 2) Standardized principal components analysis (PCA) of six fine-root functional traits and exudation rates on an individual tree level. 'cos2' values indicate that the quality of the representation of the variable on that principal component as a cos2 value close to 1 indicates that the variable is very well represented by the PC(s) in question. (a) PCA generated by six core root functional traits. PC1 and PC2 likely correspond to conservation and collaboration gradient respectively (Bergmann et al., 2020). (b) Two additional axes (PC 3 & 4) from the PCA.

Fig 3) Interaction effects of tree functional groups on the relationship between log-transformed specific root area (SRA) and mass-specific exudation rates. The model depicts the interaction effects described in the best-predicting model selected by a stepwise model selection approach: Exudation ~ SRA + myco.type:phylogeny + (1|species) (Table 4). Linear regression lines are shown for each group combination defined by mycorrhizal association (AM or EcM) and phylogeny (Angiosperm or Gymnosperm). Each equation displays the linear fit, R^2, and p-value. Among groups, EcM Angiosperms showed the steepest positive relationship between SRA and exudation rates ($R^2 = 0.33$, $P = 0.064$), while other groups exhibited weaker or non-significant trends. This interaction reflects differential trait-based exudation patterns across mycorrhizal and phylogenetic strategies.

Correlation of root exudation with four root Traits

Fig 4) Linear relationships of log-transformed root exudation with (a) root diameter, (b) SRL, (c) SRA, and (d) RTD measured at The Morton Arboretum, Lisle, IL, USA. The fitted lines indicate statistically significant relationships of root exudation with root traits (p<0.1) that are nested with AM and EcM mycorrhizal association.

Table 1) List of eleven tree species in monodominant forestry plots at The Morton Arboretum (Lisle, IL, USA); All species are ≥70 years old except *Asimina triloba*

Species	Soil series	Soil order	Mycorrhizal type	Phylogenetic group	Leaf habit	Clade
Acer saccharum	Ozaukee	Alfisol	AM	Angiosperm	Deciduous	Rosids
Asimina triloba	Ozaukee	Alfisol	AM	Angiosperm	Deciduous	Magnoliids
Platanus occidentalis	Ashkum	Mollisol	AM	Angiosperm	Deciduous	Eudicots
Chamaecyparis pisifera	Ozaukee	Alfisol	AM	Gymnosperm	Evergreen	Gymnosperms
Juniperus chinensis	Ozaukee	Alfisol	AM	Gymnosperm	Evergreen	Gymnosperms
Taxodium distichum	Sawmill	Mollisol	AM	Gymnosperm	Deciduous	Gymnosperms
Carya ovata	Ozaukee	Alfisol	EcM	Angiosperm	Deciduous	Rosids
Quercus bicolor	Sawmill	Mollisol	EcM	Angiosperm	Deciduous	Rosids
Larix gmelinii	Ozaukee	Alfisol	EcM	Gymnosperm	Deciduous	Gymnosperms
Picea abies	Ozaukee	Alfisol	EcM	Gymnosperm	Evergreen	Gymnosperms
Tsuga canadensis	Ozaukee	Alfisol	EcM	Gymnosperm	Evergreen	Gymnosperms

Note: AM, arbuscular mycorrhizal type; EcM, ectomycorrhizal type

Table 2) The species-specific means ± SDs of six core root traits at The Morton Arboretum, Lisle, IL, USA. n denotes the number of tree individuals sampled. Exudation: mass-specific exudation rate.

Species (plot)	n	Exudation rate (mg C g$_{root}^{-1}$ * day^{-1})	Specific root length (SRL (cm g^{-1}))	Root diameter (Diameter (mm))	Root tissue density (RTD (g cm^{-3}))	N concentration (root N (%))	Specific root area (SRA (cm^2 g^{-1}))	Branching intensity (BI (Tips cm^{-1}))
Acer saccharum	5	0.91 ± 0.87	1636.36 ± 529.41	0.34 ± 0.03	0.73 ± 0.14	1.23 ± 0.09	169.90 ± 42.69	3.19 ± 0.50
Asimina triloba	4	0.60 ± 0.63	1110.21 ± 104.32	0.64 ± 0.02	0.29 ± 0.05	2.91 ± 0.23	222.05 ± 28.12	0.72 ± 0.06
Platanus occidentalis	5	1.19 ± 0.88	1045.53 ± 540.38	0.52 ± 0.10	0.52 ± 0.07	1.74 ± 0.23	157.53 ± 52.86	2.95 ± 0.53
Chamaecyparis pisifer	7	0.75 ± 0.71	1528.93 ± 460.45	0.42 ± 0.06	0.51 ± 0.09	1.87 ± 0.18	195.13 ± 39.51	2.18 ± 0.56
Juniperus chinensis	6	1.22 ± 1.78	863.63 ± 273.33	0.55 ± 0.09	0.54 ± 0.11	1.39 ± 0.11	142.73 ± 25.23	2.40 ± 0.90
Taxodium distichum	4	0.58 ± 0.16	1696.22 ± 374.94	0.51 ± 0.08	0.31 ± 0.04	1.89 ± 0.26	264.06 ± 26.82	2.02 ± 0.42
Carya ovata	5	0.52 ± 0.69	1160.74 ± 209.51	0.37 ± 0.05	0.85 ± 0.17	1.25 ± 0.26	133.43 ± 25.23	5.32 ± 1.15
Quercus bicolor	6	0.73 ± 0.54	1439.00 ± 397.30	0.33 ± 0.02	0.86 ± 0.14	1.29 ± 0.14	146.67 ± 32.44	5.63 ± 0.48
Larix gmelinii	6	3.82 ± 2.83	975.76 ± 384.57	0.58 ± 0.11	0.43 ± 0.07	1.48 ± 0.12	167.97 ± 40.13	3.68 ± 0.60
Picea abies	6	1.52 ± 0.52	1639.29 ± 220.22	0.42 ± 0.04	0.45 ± 0.05	2.01 ± 0.13	214.35 ± 19.09	4.44 ± 0.66
Tsuga canadensis	6	1.05 ± 0.66	689.36 ± 276.65	0.59 ± 0.07	0.60 ± 0.17	1.24 ± 0.42	125.24 ± 46.82	4.16 ± 0.63

Table 3) Type III Analysis of Variance Table for predicting root traits of 11 tree species in The Morton Arboretum (n =60) using a linear mixed model (dependent variables~myco x phylo x leaf + (1|species)). The species were nested in mycorrhizal types, phylogenetic groups and leaf habit. p-values (P<0.1) are highlighted in bold. The significance of fixed effects in the model was tested by Type III ANOVA with Wald chisquare tests. All measurements were log-transformed to ensure normality and improve homogeneity of variance.

Fixed Effect	Mass-specific Exudation rate			Root diameter (mm)			Specific root length (m g^{-1})			Specific root area (cm^2 g^{-1})		
	Chisq	Pr(>Chisq)	Adjusted p	Chisq	Pr(>Chisq)	Adjusted p	Chisq	Pr(>Chisq)	Adjusted p	Chisq	Pr(>Chisq)	Adjusted p
Mycorrhizal type (myco)	0.98	0.32	0.48	4.37	**0.04**	0.11	0.08	0.77	0.77	2.44	0.12	0.16
Phylogenetic group (phylo)	0.02	0.88	0.92	0.08	0.78	0.78	1.15	0.28	0.34	3.92	**0.048**	**0.10**
Leaf habit (leaf)	0.01	0.92	0.92	0.09	0.77	0.78	1.45	0.23	0.34	4.83	**0.028**	**0.08**
myco:phylo	9.27	**0.002**	**0.007**	2.58	0.11	0.22	2.22	0.14	0.34	0.66	0.42	0.42
myco:leaf	1.90	0.169	0.337	0.09	0.77	0.78	1.27	0.26	0.34	2.26	0.13	0.16
% of the total variance explained by random effect (Species plot)	> 0.1%			55.2% (p<0.001)			33.3% (p<0.001)			26% (p<0.01)		

Fixed Effect	Root tissue density (g cm^{-3})			Root N concentration (%)			Branching intensity (cm^{-1})		
	Chisq	Pr(>Chisq)	Adjusted p	Chisq	Pr(>Chisq)	Adjusted p	Chisq	Pr(>Chisq)	Adjusted p
Mycorrhizal type (myco)	9.42	**0.0021**	**0.006**	3.23	**0.072**	0.22	10.82	**0.0010**	**0.0057**
Phylogenetic group (phylo)	3.69	**0.055**	**0.082**	0.01	0.91	0.91	0.02	0.89	0.92
Leaf habit (leaf)	4.52	**0.034**	**0.067**	0.30	0.58	0.87	0.05	0.82	0.92
myco:phylo	0.42	0.52	0.52	0.13	0.72	0.87	0.59	0.442	0.88
myco:leaf	1.05	0.30	0.37	0.20	0.65	0.87	0.01	0.922	0.92
% of the total variance explained by random effect (Species plot)	45.5% (p<0.001)			55.4% (p<0.001)			74% (p<0.001)		

Note. Numerator Degrees of Freedom = 1; Denominator Degrees of Freedom =60. Mass-specific exudation rate (mgC*d^{-1}*g root^{-1})

Table 4) Mixed effects models for the effects of mycorrhizal type on exudation-to-trait relationships. The model considered root traits × mycorrhizal type + (1|species.plot) to predict exudation rates. 'species.plot' represents species-specific monodominant plots at The Morton Arboretum, Lisle, IL. The best model (exudation ~ SRA + Mycorrhizal type:Phylogeny) is selected via a stepwise reduction approach using mixed models. The other models are ordered from the lowest AIC values. The numbers in the table represent coefficients estimate (Std β) and standard error (Std SE) in brackets. Adjusted p-values were calculated using the Benjamini-Hochberg (BH) correction for multiple testing. Bold value indicates statistical significance of Adjusted p-values less than 0.05. Significance levels of Adjusted p-values: *p<0.05; **p<0.01; ***p<0.001. The comprehensive table for the effects of phylogeny on trait-exudation relationships is in Table S2.

Model to predict exudation rates - AIC values	Significant Fixed effect term - R^2m & R^2c	Std β (Std SE)	Adjusted p
SRA + Mycorrhizal type : Phylogeny	(Intercept)	-5.2936 (1.99)*	**0.013**
Best model selected with AIC =164.8	SRA	1.13 (0.39)**	**0.013**
	AM : Angiosperm	-1.05 (0.33)*	**0.013**
	EcM : Angiosperm	-1.13 (0.35)*	**0.013**
	AM : Gymnosperm	-1.23 (0.33)**	**0.013**
	- $R^2m = 0.34$; $R^2c= 0.36$		
RTD * Mycorrhizal type	RTD : EcM	-2.05 (0.64)**	**0.008**
- AIC=163.2	- $R^2m = 0.24$; $R^2c= 0.44$		
SRA * Mycorrhizal type	SRA	1.76 (0.64)*	**0.023**
- AIC=169.3	- $R^2m = 0.33$; $R^2c= 0.33$		
SRL * Mycorrhizal type	SRL	1.54 (0.45)**	**0.005**
- AIC=170.7	- $R^2m = 0.19$; $R^2c= 0.43$		
Diameter (mm) * Mycorrhizal type	ECM	2.91 (0.90)*	**0.016**
- AIC=173.0	Diameter : ECM	3.09 (1.09)*	**0.027**
	- $R^2m = 0.10$; $R^2c= 0.27$		

Note) Insignificant models (i.e., Root N * Mycorrhizal type; BI * Mycorrhizal type) and fixed effect terms are not reported in this table. See Table S2 for comprehensive results that include the effects from phylogeny on trait-exudation relationships. All measured variables were log-transformed to ensure normality of data. (Intercept) represents baseline exudation rate (AM reference). R^2m, Marginal R^2; R^2c, Conditional R^2, RTD. R^2m represents variance explained by fixed effects only, whereas R^2c indicates variance explained by both fixed and random effects. Root Tissue Density (g/cm^3); SRA, Specific Root Area (m^2/g); Specific Root Length (cm/g); Diameter, root mean diameter (mm); Root N, Root N Concentration (%); BI, Branching Intensity (cm^{-1}).

807
808