1

MinD proteins regulate CetZ1 localisation in *Haloferax volcanii*

Hannah J. Brown[1] and Iain G. Duggin[1,#]

[1]Australian Institute for Microbiology and Infection, University of Technology Sydney,
Ultimo, NWS, 2007, Australia.

[#]Correspondence: iain.duggin@uts.edu.au

Abstract

CetZ proteins are archaea-specific homologues of the cytoskeletal proteins FtsZ and
tubulin. In the pleomorphic archaeon *Haloferax volcanii*, CetZ1 contributes to the
development of rod shape and motility, and has been implicated in the proper
assembly and positioning of the archaellum and chemotaxis motility proteins. CetZ1
shows complex subcellular localization, including irregular midcell structures and
filaments along the long axis of developing rods and patches at the cell poles of the
motile rod cell type. The polar localizations of archaellum and chemotaxis proteins
are also influenced by MinD4, the only previously characterized archaeal member of
the MinD family of ATPases, which are better known for their roles in the positioning
of the division ring in bacteria. Using *minD* mutant strains and CetZ1 subcellular
localization studies, we show here that a second *minD* homolog, *minD2*, has a
strong influence on motility and the localization of CetZ1. Knockout of the *minD2*
gene altered the distribution of a fluorescent CetZ1-mTq2 fusion protein in a broad
midcell zone and along the edges of rod cells, and inhibited the localization of CetZ1-
mTq2 at the cell poles. MinD4 had a similar but weaker influence on motility and
CetZ1-mTq2 localization. The MinD2/4 mutant strains formed rod cell shapes like the
wildtype at an early log stage of growth. Our results are consistent with distinct roles
for CetZ1 in rod shape formation and at the poles of mature rods, that are positioned
through the action of the MinD proteins and contribute to the development of
swimming motility in multiple ways. They represent the first report of MinD proteins
controlling the positioning of tubulin superfamily proteins in archaea.

Introduction

Spatial and temporal regulation of protein subcellular localization is critical for the proper function of all cells. In bacteria, there are several different classes of proteins from the ParA/MinD superfamily [1], which coordinate positioning of flagella [2; 3], chemosensory arrays [4; 5], chromosomes [6], plasmids [7], cell division cytoskeletal proteins [8; 9], and various other molecules or structures [10; 11; 12; 13; 14]. A key characteristic of ParA/MinD superfamily proteins is their ATPase activity [15; 16] and dimerization [9; 17] involving amino acid residues within a deviant Walker A motif [1; 10]. In MinD proteins, a subset of proteins within the ParA/MinD superfamily, this ATP binding and hydrolysis drives intracellular Min system dynamics that can generate concentration gradients of Min proteins [7; 15; 16]. In bacteria, these gradients are critical for the correct positioning of FtsZ, a key cell division protein and tubulin superfamily homologue [18; 19; 20], at mid-cell to help ensure symmetric division of the mother cell into two daughter cells [9; 21; 22].

The Min system of *Escherichia coli* is the prototypical example of Min system dynamics and function in bacteria. MinD oscillates between poles of the rod-shaped cells, controlled by its regulator MinE, and carries MinC an FtsZ antagonist. Together, these proteins generate dynamic concentration gradients of MinCD which is on average highest at cell poles and lowest at mid-cell, preventing Z-ring formation anywhere other than at mid-cell [8]. MinE accelerates the ATPase activity of MinD [15] and drives its release from the membrane which it is bound to via a membrane targeting sequence [23; 24; 25]. Together these proteins constitute a reaction-diffusion system driven by the hydrolysis of ATP. Disruption of MinD ATPase activity blocks oscillation and function completely [26; 27].

ParA/MinD family homologues have been identified across the archaeal domain, but MinD homologues are particularly abundant in *Euryarchaea* [28]. In the model halophilic archaeon, *Haloferax volcanii*, there are four MinD paralogues. The functions of MinD1-3 (HVO_0225, HVO_0595, and HVO_1634, respectively) have not been described and no clear protein partners equivalent to bacterial MinE have been identified. MinD4 (HVO_0322) has been characterised and is known to oscillate between cell poles, as well as concentrating with a cap-like appearance at the poles [28]. Surprisingly, the oscillation was not dependent on the ATPase Walker A or Walker B motifs, but the cap-like structures were, suggesting that the oscillation is driven by an alternative energy source [28]. Furthermore, although *H. volcanii* possess FtsZ homologues, none of the MinD paralogues were found to have a role in the mid-cell positioning of FtsZ; simultaneous deletion of all four paralogues had no detected effect of FtsZ positioning. Instead, deletion of *minD4* caused a motility defect and reduced the polar positioning of key protein constituents of the archaellum and chemotaxis arrays, ArlD and CheW, respectively [28]. Furthermore, disruption of MinD4 predicted ATPase activity through mutations to the Walker A or Walker B motifs, or truncation of its extended C-terminal tail all result in motility

defects. Potential roles of the other MinD paralogues in motility of *H. volcanii* have
not been reported.

Although the archaellum and chemosensory arrays are considered the core motility
machinery in *H. volcanii*, there are many other contributing factors, and a range of
proteins with a variety of primary functions have been shown to influence motility [29;
30; 31; 32; 33; 34; 35; 36; 37; 38]. CetZ cytoskeletal proteins, a group of tubulin
superfamily proteins specific to archaea and abundant in Haloarchaea [37; 39; 40],
have been implicated in the control of cell shape and motility [37; 41]. The most well-
studied and highly conserved CetZ homologue, CetZ1, is necessary for rod-shape
development and swimming motility [37; 41]. Our recent results have suggested that
CetZ1 is an important factor for the proper assembly and positioning of motility
proteins ArlD and CheW [42]. It is not yet clear whether CetZ1 contributes to motility
solely via its essential role in rod shape development [37], or whether it has other
functions that promote motility. It also is not yet clear how MinD4 and CetZ1 might
coordinate their activities to regulate positioning and assembly of motility machinery.

Currently, no connection between Min proteins and FtsZ or other tubulin superfamily
homologues, such as the CetZ, has been reported in archaea. Here, we investigate
the potential for MinD homologues of *H. volcanii* to control positioning of CetZ1 and
identify MinD2 as a strong regulator of CetZ1 localisation in rod cells. This is
consistent with a model in which a hierarchy or network of proteins is involved the
promoting the development of motility structures at the cell poles.

Methods

Growth and culturing

Haloferax volcanii was routinely grown in Hv-Cab [43] medium supplemented with
uracil (50 μg/mL) to fulfil auxotrophic requirements if necessary, or L-Tryptophan (1
mM) to induce expression of genes under the control of the p.*tna* promotor. Liquid
cultures were grown at 42 °C with shaking (200 rpm).

Strain and plasmid construction

Strains, plasmids and oligonucleotides used in this study are listed in Tables S1, S2
and S3, respectively. *H. volcanii* H26 (Δ*pyrE2* ΔpHV2) was used as the wildtype
parent strain for construction of mutant strains. All strains investigated in experiments
contained a pHV2-based vector (pTA962 vector or derivatives) to complement the
pyrE2 auxotrophic marker and better match the original *H. volcanii* wild-type
genotype; previous work has shown that the base strains with Δ*pyrE2*, Δ*hdrB* or
ΔpHV2 show poorer expression of the cell morphology and motility phenotypes,
which are sensitive to nutrient and culture conditions [37; 43; 44].

To create chromosomal point mutations in *minD2*, generating strains ID810
(*minD2WA**) and ID811 (*minD2WB**)*,* H26 was transformed with vectors
pTA131_minD2WA or pTA131_minD2WB, respectively, using previously described
transformation methods [45] and the homologous recombination-based 'pop-in pop-
out' method [46] was used to replace the native *minD2* gene with Walker A (K16A,
AAG to GCC) or Walker B (D117A, GAT to GCC) mutant alleles. Primers
MinD2_USflank_extF (forward) and MinD2_DSflank_extR (reverse), designed to
bind externally to the regions of homologous recombination, were then used to
amplify the *minD2* locus. The resultant PCR products were purified, and then
sequenced by Sanger sequencing (Australian Genome Research Facility) to identify
transformants carrying the expected mutations. All strains using *minD2WA** (ID810)
as the parent strain were also verified by checking for complete conversion of alleles
and the absence of potential suppressor mutations (SNPs, indels) using Illumina
whole-genome sequencing with ~50X coverage.

Plasmids for the expression of *minD2*, *minD2WA**, and *minD2WB** were generated
by amplifying these open reading frames from *H. volcanii* H26 (ID621), ID810, and
ID811, respectively. Forward and reverse primers for this reaction incorporated NdeI
and BamHI restriction sites at 5' and 3' ends, respectively. ORFs were then ligated
between the same restriction sites in pTA962 [47], generating pHJB63-65 which
were sequence verified.

pHJB72 for the dual expression of *cetZ1-G-mTq2* and *minD2* was generated by
amplifying the *minD2* open reading frame from pHJB63, incorporating an XbaI site at
the 5' end. The amplicon was then ligated between NheI and NotI of pHVID135.

Plasmids and oligonucleotides used in this study are listed in Table S2 and S3,
respectively.

Soft-agar motility assays

138 Soft-agar motility assays were carried out as described in [42], using Hv-Cab with
139 0.25% (w/v) agar and 1 mM L-Tryptophan. Motility was quantified by averaging
140 perpendicular measurements of the motility halo diameter after three days of
141 incubation of the plates inside a plastic bag at 42 °C.

Light microscopy

To prepare samples for microscopy, single colonies were first used to inoculate 3 mL
Hv-Cab medium, each representing a single culture (biological) replicate. These
cultures were allowed to grow to OD_{600} ~0.6. Then, 3 µL of this culture was taken to
inoculate 10 mL Hv-Cab supplemented with 1 mM L-Tryptophan and grown for 18 h
(OD_{600} ~ 0.1). A 2 µL droplet of cells was spotted onto a 1.5% agarose pad
containing 18% BSW as described in [37]. Samples were then covered with a clean
#1.5 glass coverslip.

Imaging for phase-contrast and fluorescence microscopy was carried out using a V3
DeltaVision Elite (GE Healthcare) with an Olympus UPLSAPO 100X NA 1.4 objective
and pco.edge 5.5 sCMOS camera. A filter set for mTurquoise2/CFP (excitation=400-
453, emission=463-487) was used for imaging of CetZ1-mTq2 expressing strains.
3D Structured-illumination microscopy (3D-SIM) was conducted as described
previously [41] using a DeltaVision OMX SR microscope (GE Healthcare) and a
DAPI filter for excitation (405 nm) and an AF488 filter for emission (504-552 nm),
suitable for imaging of mTurquoise2. Raw images were reconstructed using
SoftWorx (Applied Precision, GE Healthcare), and 3D renders were generated
[48]using Bitplane IMARIS (v9.6.0).

Image analysis

Image analysis was conducted using the MicrobeJ (v5.13I) [49] plugin for FIJI
(v2.14.0/1.54f) [50]. Cell shape characteristics were analysed using phase-contrast
images and default settings. Fluorescent foci detection was carried out after
background subtraction with a ball size of 50 pixels, and the *point* detection method
was used for analysis of CetZ1-mTq2 localisation (Tolerance=200).

Fluorescence intensity profiles along the long axis of each cell were generated using
the *XStatProfile* function in MicrobeJ (the long axis was normalized between 0-1).
The mean and standard deviation of the raw fluorescence intensity over all cells
were then determined (Stat=mean±stdv, Y axis=MEDIAL.intensity.ch2, number of
bins=50). The intensity data were also normalized (0-100%) and the median
fluorescence intensity along the long axis determined for each culture replicate
(Stat=median, Y axis=MEDIAL.intensity.ch2, number of bins=50), and then the mean

and standard error of all culture replicates were plotted. These data were also
displayed as the difference in fluorescence intensity between *minD* deletion or
Walker mutant backgrounds and the wildtype background.

Heatmaps of foci localisation were generated using pooled data from all culture
replicates and the *XYCellDensity* function (Rendering=Density) with default settings.

Results

Deletion of minD2 or minD4 results in motility defects with normal rod cell
shape

We began by assessing the effect of *H. volcanii minD* gene deletions on motility,
focussing on MinD2 and MinD4, which are expressed significantly in standard culture
conditions [51; 52; 53]. MinD4 has previously been studied in this context [28]. The
minD deletion strains grew at rates like the wildtype in starter cultures, and motility
was assessed using soft-agar motility assays in which cells develop into a motile rod
form [37] and then swim out through the gel from a central inoculation point, forming
a halo.

Deletion of *minD2* resulted in a motility defect (approximately 25% of the halo
diameter compared to wildtype), which was significantly stronger than that caused by
deletion of *minD4* (approximately 50% of wildtype motility) (Fig. 1a, b). Deletion of
both *minD2* and *minD4* resulted in a stronger motility defect (approximately 16% of
wildtype motility). The deletion of all four *minD* homologues (Δ*minD1/2/3/4*) was
indistinguishable to that of the Δ*minD2/4.*

H. volcanii is pleomorphic and cultures can show a range of cell shapes including
irregular disk, or plate-shaped cells, and rods with varying widths and degrees of
elongation [37; 43]. Furthermore, a motile-rod cell type with regular cell widths
develops under certain conditions to promote swimming [37; 54]. To determine
whether the two MinD proteins are required for rod formation, which might contribute
to their effects on motility, we sampled wild-type and Δ*minD2/4* cells from the early-
log phase of liquid batch cultures (OD ~ 0.1), where motile rods are prevalent [43;
54]. By quantifying cell circularity as an indicator of the extent of cell elongation, we
observed no significant cell shape differences at this stage of growth in any of the
strains (Fig. 1c, d).

MinD2 and MinD4 influence the cellular positioning of CetZ1

MinD4 has been shown to have a role in the polar positioning of the archaellum and
chemotaxis arrays through the use of tagged marker proteins ArlD and CheW,
respectively [28]. We recently found that CetZ1 also influences the proper assembly
and positioning of ArlD and CheW [42]. It is unknown whether MinD4 and CetZ1
cooperate or work independently to promote motility. To begin investigating this, we
sought to determine whether the absence of MinD2 or MinD4 affects the localisation
of CetZ1.

To do this, we observed the localisation of a CetZ1-mTurquoise2 (mTq2) fusion
protein [55] in the wildtype and various *minD* deletion strain backgrounds.
Fluorescence microscopy of CetZ1-mTq2 in wildtype cells imaged in early-log growth
(OD_{600} ~ 0.1) (Fig. 2a), showed the expected patchy localisation with relatively high
fluorescence at cell poles and the mid-cell region [41; 42; 55]. To characterise these
patterns, we analysed the whole population of cells, in the wildtype and *minD* mutant

backgrounds, by generating heatmaps that represent the position and frequency of
fluorescent foci within a normalized cell shape (reflecting the averaged aspect ratio)
(Fig. 2b). We also generated plots of the total and normalized mean fluorescence
intensity of CetZ1-mTq2 along the long-axis of the cells (Fig. 2c-d). In the Δ*minD2*
background, CetZ1-mTq2 localisation remained patchy and concentrated in a broad
zone near midcell, which was noticeably broader compared to the wildtype, and the
high-intensity polar localisations of CetZ1-mTq2 were not observed (e.g., Fig. 2a, c,
d). These patterns were also reflected in the number of fluorescent foci detected
along the cell lengths (Fig. 2b); the wildtype background displayed clear peaks of
fluorescence around midcell and at the cell poles, but in the Δ*minD2* background the
midcell foci were more spread out along the cell length, and showed almost
complete loss of the polar clusters of foci.

To confirm whether the changes to CetZ1-mTq2 localization patterns were resultant
from the loss of *minD2* specifically*,* we observed CetZ1-mTq2 localisation when
minD2 deletion was complemented by expression of *minD2* from a plasmid
containing the open reading frames for *cetZ1-mTq2* and *minD2* in tandem, under the
control of the inducible *p.tna* promotor (Fig. S1). Dual expression of these genes in
the Δ*minD2* background using 1 mM L-Tryptophan did not clearly restore the CetZ1-
mTq2 localisation defect (Fig. S1a). However, when 2 mM L-Tryptophan was used,
the localisation of CetZ1-mTq2 was partially restored (Fig. S1c-g), confirming that
minD2 regulates the localization of CetZ1. The requirement for the additional
tryptophan to observe partial complementation is likely due to a sensitivity of the
phenotype to the expression level of *minD2*, which is likely to be influenced by its
second position in the tandem expression plasmid [55].

A weaker defect of CetZ1-mTq2 polar localization was also observed in the Δ*minD4*
background (compare the polar intensities between mutant and wildtype in Fig. 2d).
Additional control experiments demonstrated that the Δ*minD* strains producing
CetZ1-mTq2 had no significant differences in shape compared to the wildtype control
(OD ~ 0.1), and we also observed an equivalent motility defect with or without the
CetZ1-mTq2 fusion (Fig. S2a, b).The results in Figure 2 also revealed that the zone
of CetZ1-mTq2 localization around the midcell region was broadened in the absence
of either MinD2 or MinD4 (Fig. 2b, d). Interestingly, such broadening was not
apparent in strains in which both *minD2* and *minD4* were deleted, although the polar
localization of CetZ1-mTq2 was still strongly inhibited (Fig. 2b, d). This suggests that
the midcell zone of CetZ1-mTq2 localization is impacted when only one of MinD2 or
MinD4 is present and implies a genetic interaction or interference between *minD2*
and *minD4* that influences midcell CetZ1.

Structural characteristics of CetZ1-mTq2 localization in individual cells lacking
minD2* or *minD4

We further resolved structural characteristics of the CetZ1-mTq2 polar localizations
and the effects of the *minD* knock-outs, using three-dimensional structured-
illumination microscopy (3D-SIM). Multiple cells are shown as maximum intensity

projections in Fig. 3a-c to visualize the effect of the mutations on CetZ1 structures in
individual cells and facilitate comparisons to the aggregate data shown in Fig. 2b-d.

In wildtype early-log cells, CetZ1-mTq2 often appeared as a bright focus, doublet,
multi-lobed, or cap-like structure at the cell poles, suggestive of a ring or disc-like
structure at the tip of the cell pole (Fig. 3a, d). Patches and filaments were also
observed along the cell edges in the midcell zone. In the absence of *minD2*, the foci
of CetZ1-mTq2 appeared more randomly distributed around the cells, with no clear
preference for the cell poles or the midcell zone(Fig. 3b). In the absence of *minD4*,
this effect was weaker, and some of the cells still appeared to show the relatively
bright polar tip structures of CetZ1-mTq2 (Fig. 3c). The Δ*minD2* and Δ*minD4*
backgrounds also showed noticeably more of the CetZ1-mTq2 patches or foci
throughout the cell area compared to the wildtype (Fig. 3a-c). By inspecting the slice
series of images to resolve the 3D detail (Fig. 3e, Supplementary Video 4), CetZ1-
mTq2 foci appeared noticeably more prevalent at the top and bottom surfaces along
the length of the *minD* mutant cells, compared to the wildtype, and potentially in the
cytoplasm too. These observations are consistent with the broadening of the midcell
zone and reduced polar positioning of CetZ1-mTq2 in the *minD* mutants seen in the
aggregated epifluorescence data (Fig. 2b-d).

ATPase active-site residues of MinD2 are required for normal swimming
motility and polar localization of CetZ1

MinD proteins typically rely on ATP hydrolysis to drive their dynamic localization
patterns, such as oscillation between cell poles, generating concentration gradients
which form the basis for their spatial and temporal control of downstream proteins
[10; 27; 28]. When the Walker A and Walker B motifs of *H. volcanii* MinD4 were
previously mutated to disrupt ATP binding and hydrolysis, swimming motility was
reduced and polar localisation of MinD4 was disrupted, although not its oscillation
[28].

Given that MinD2 had the strongest effects on swimming motility and CetZ1
localization, we chose to further investigate the importance of the ATPase active-site
for MinD2 functionality in motility and CetZ1 localization. We generated
chromosomal point-mutations in the endogenous *minD2* gene in the ATPase active
site via the deviant Walker A (K16A) and Walker B (D117A) motifs, here forth referred
to as *minD2WA** and *minD2WB**, respectively. These mutations are equivalent to
those previously made in *minD4* [28].

Both *minD2WA** and *minD2WB** strains (containing the empty vector pTA962)
partially perturbed motility, resulting in ~78% and ~36% of the wildtype motility halo
diameter, respectively (Fig. 4a, c), compared to the complete deletion of *minD2*
(~25%, Fig. 1a, b). The motility defect of Δ*minD2* was complemented by expressing
minD2 from a plasmid (pTA962 backbone), confirming the important role of *minD2* in
motility (Fig. 4b, c).

When *minD2WA** was expressed from a plasmid in the Δ*minD2* background, this
strain had a more pronounced motility defect (~35%) compared to the chromosomal
mutant *minD2WA** (~78%). This might be related to different expression levels of
minD2WA* in the plasmid with p.*tna* compared to its native chromosomal site.
Expression of *minD2WB** from pTA962 in Δ*minD2* produced equivalent motility to the
complete *minD2* deletion, consistent with its much stronger effects on motility
observed with the chromosomal *minD2WB** mutant (Fig. 4b, c).

Phase-contrast microscopy and analysis of cell shape (circularity) confirmed that,
like Δ*minD2*, the *minD2WA** and *minD2WB** mutations did not significantly affect rod
shape (at OD ~0.1), whether expressed chromosomally (Fig. S3a, c) or from the
plasmid (Fig. S3b, d).

We then assessed the effect of the *minD2WA** and *minD2WB** chromosomal
mutants on CetZ1-mTq2 localisation. Control experiments showed that expression of
CetZ1-mTq2 did not impact cell circularity or motility of either the *minD2WA** or
minD2WB* strains (Fig. S3c, e). Fluorescence microscopy (Fig. 5a) showed that
both Walker mutations resulted in a substantial loss of CetZ1-mTq2 fluorescence
intensity (Fig. 5c, dc) and foci localisation at cell poles and showed the midcell zone
broadening (Fig. 5b) like in the Δ*minD2* background (Fig. 2).

Discussion

MinD proteins and related protein families are important for spatial and temporal positioning of a variety of molecules but have mainly been characterized in bacteria [10]. One of four MinD paralogues in the archaeal model organism *H. volcanii*, MinD4, has previously been described to affect motility via regulating the positioning of archaellum and chemotaxis arrays, via unknown mechanisms [28]. The tubulin-like cytoskeletal protein CetZ1 is another factor promoting the proper assembly and positioning of these proteins at the cell poles [42]. CetZ1 is needed for rod cell shape formation [37], which may contribute to motility in several ways, and shows complex subcellular localization along the cell body and then at the cell poles of motile rods, where it is hypothesised to form a scaffold-like structure, though its mechanism of action at these sites is unclear.

We found here that MinD2 is important for motility and for the subcellular positioning of CetZ1. The potential relationship between these two functions is yet to be elucidated. Deletion of *minD2* was found to have a more severe motility defect than deletion of *minD4*, and deletion of both genes resulted in an even greater motility defect, suggesting distinct roles or additive contributions of these two MinD paralogues in motility. MinD2 and to a lesser extent MinD4 also influenced the subcellular positioning of CetZ1 in the midcell region and at the poles, in a manner that was independent of cell shape. CetZ1 has been implicated in distinct roles in establishing rod-shape during motile cell differentiation, and then at the poles of the mature motile rods [37; 42]. Our results suggest that the effects of MinD2 on CetZ1 localization do not have an essential role in maintaining rod shape, because *ΔminD2* cells could still form normal shaped rods at an early-mid log stage of culture.

We found that mutations in the Walker A and B motifs of the MinD2 ATPase both strongly influenced CetZ1 positioning along the cell length and poles, like the *minD2* deletion. However, only the Walker B mutant (D117A) strongly perturbed motility, whereas the Walker A (K16A) mutant (in its chromosomal context) resulted in only a mild motility defect. This suggests that MinD2 has an impact on motility that is largely independent of the K16 residue and CetZ1 positioning. The K16 residue is expected to be involved in nucleotide dependent dimerization [27], so this function may not be critical to the overall motility function of MinD2, and instead plays a role in CetZ1 positioning for an as yet unresolved purpose. Notably, the equivalent MinD4 Walker A and B mutants reduced motility equivalently to *minD4* deletion [28]. The MinD2 and MinD4 proteins could therefore have multiple, differing functions that contribute to the proper development of motile *H. volcanii* cells.

The accompanying study by Patro et al. [56] found that MinD2 is required for normal motility and the proper polar localisation of the archaellum and chemotaxis proteins, ArlD and CheW, like previously seen with MinD4 [28]. They also reported a difference in cell shape during the development of rod cells in the *ΔminD2* compared to the wildtype. This likely represents an earlier phase of *H. volcanii* cultures, where

cells are actively shapeshifting into motile rods [43; 54], compared to the current
study where rods had already formed, and we found that their cell shapes in the
wildtype and Δ*minD2* strains were indistinguishable. Though we note that since *H.*
volcanii morphogenesis is highly sensitive to exact media composition and culture
conditions [43; 44], excessive comparisons of the exact magnitudes or timing of
events from separate studies should be avoided. Nevertheless, if the rate or timing of
rod formation is influenced by MinD2 [56], this may affect another developmental
stage of motility, such as the placement of Arl/Che proteins at the poles. However, it
is still unclear whether any effects on cell shape development directly contribute to
the *minD2*-dependent reduction of motility.

Considering the main findings from this and other recent studies [28; 56], we can
summarize several possibilities for the role of MinD proteins in motility and CetZ1
localization (Fig. 6). In this model, MinD2 and MinD4 help direct the archaellum (Arl)
and chemotaxis (Che) proteins to the cell poles directly or via positioning and
assembly of polar CetZ1. We recently found that CetZ1 plays a role in the proper
placement of the motility machinery [42]. However, the present finding that the
chromosomal *minD2WA** mutation blocked polar CetZ1 localization but only weakly
affected motility suggests that MinD proteins might separately control both the
position of CetZ1 and another aspect of motility, such as placement of the motility
machinery at the cell poles, independently of MinD2 regulated CetZ1 positioning.

Previous work showed that MinD4 forms dynamic cell cap-like structures in an
ATPase-dependent manner, and yet oscillates in the cells independently of ATPase
function (Fig. 6), suggesting that it interacts with another oscillator that drives the
movement [28]. The localization patterns of MinD2 are currently unknown due to the
absence of functional fluorescently tagged MinD2 [56]. However, our observations
that the complex patterns of CetZ1 localization are affected by MinD2 at multiple
sites and at a cellular scale would suggest that similar cellular level MinD2 pattern
formation might exist.

In summary, our findings have showed that multiple MinD homologues can have
differing functions and regulate the subcellular positioning of the archaeal tubulin-like
protein CetZ1 in the context of the development of swimming motility. They represent
the first demonstration of MinD proteins controlling the position of tubulin superfamily
proteins in archaea. The MinD-controlled positioning of CetZ1 in distinct subcellular
locations is consistent with multiple functions of CetZ1, evoking parallels with the
multiple functions of tubulins and other eukaryotic cytoskeletal proteins that are
extensively regulated in time and space.

Acknowledgements

The authors would like to acknowledge facilities support from L. Cole and A. Bottomley and the UTS Microbial Imaging Facility. The work was supported by the Australian Research Council (DP160101076 and FT160100010 to IGD).

Author Contributions

Conceptualisation – HJB & IGD; Data curation - HJB; Formal Analysis – HJB; Investigation – HJB; Methodology – HJB & IGD; Supervision – IGD; Original draft – HJB; Revision and reviewing the manuscript – HJB & IGD; Funding Acquisition – IGD.

 Figures

Figure 1. MinD2 and MinD4 have roles in motility without influencing shape. A) H26 wildtype
and *minD* deletion mutant backgrounds, carrying an empty pTA962 vector, were used to inoculate the
centre of a soft agar plate. **B)** The diameter of their motility halos was measured across six
independent liquid starter culture replicates (represented by individual points), each determined from
the mean of at least two replicate agar plates. Mean and standard error is shown. **C)** Phase-contrast
microscopy of cells grown in Hv-Cab liquid medium with 1 mM L-tryptophan, sampled at OD_{600} ~ 0.1.
Scale bars = 5 μm. **D)** Cell circularity distributions, quantified from phase-contrast images as a proxy
for cell elongation. Large data points represent mean circularity of one culture replicate, small data
points represent individual cells from all culture replicates for the indicated strain. Mean and standard
error of culture replicate means are shown (bars). The number of individual cells measured from
pooled culture replicates were as follows: WT + pTA962, n=996; Δ*minD2* + pTA962, n=247; Δ*minD4* +
pTA962, n=217; Δ*minD2/4* + pTA962, n=223; Δ*minD1/2/3/4* + pTA962, n=279. In **B** and **D**, one-way
ANOVA was used as a statistical test, and only significant and/or relevant comparisons are shown.
****$p<0.00005$, ns=not significant.

Figure 2. MinD2 controls polar positioning of CetZ1. H26 wildtype and minD deletion strains containing pHVID135 (for expression of CetZ1-mTq2) were grown in Hv-Cab with 1 mM L-Tryptophan and sampled at an approximate OD_{600} of 0.1. **A)** Epifluorescence images of CetZ1-mTq2 localisation in the indicated minD knockout strains (scale bar 5 µm). **B)** Heatmaps representing the position and number (heatmap density) of detected CetZ1-mTq2 fluorescence peak intensities (foci) along the long (vertical) and lateral (horizontal) axes of all cells combined. **C)** The median fluorescence intensity of CetZ1-mTq2 along the normalized long axis of each cell was used to generate a plot of the mean (bold line) and standard deviation (shading) of the raw fluorescence intensity data from all cells (n_c) (combined from at least two replicate cultures). Cultures of the wildtype strain were repeated in parallel alongside each mutant independently. **D)** The same intensity data were also normalized (0-100%) for each culture replicate and plotted as the mean (bold line) and standard error (shading) of all culture replicates (n_r) to enable comparison of protein distributions amongst the various strains.

**Figure 3. Subcellular structures of CetZ1-mTq2 in wildtype and *minD* mutant backgrounds, by
3D-SIM imaging. (A-C)** Maximum intensity projections of CetZ1-mTq2 in H26 wildtype, Δ*minD2*, and
Δ*minD4* backgrounds, viewed along the imaging z-axis (i.e., XY-views). Scale bar in panel C is 1 μm
and applies to panels A-C. **D)** XY-views of 3D rendered images, showing non-circular fluorescence
foci of CetZ1-mTq2 at the cell poles in H26 wildtype background. Red boxes indicate zoomed area
(bottom). **E)** Examples of 3D-SIM slice sections of CetZ1-mTq2 in the indicated strains at approximate
bottom, middle, and top z-positions of the cells (scale bar is 2 μm and applies to all images).

Figure 4. Chromosomal Walker mutants of *minD2* perturb motility. A) H26 wildtype, *minD2WA**, and *minD2WB** backgrounds containing pTA962, **B)** and Δ*minD2* containing pTA962 were compared to Δ*minD2* containing pHJB63-65 for expression of *minD2*, *minD2WA**, and *minD2WB**, respectively, and inoculated onto Hv-Cab soft agar (0.25 % w/v) supplemented with 1 mM L-Tryptophan. **C)** Halo diameters were quantified across four starter culture replicates with two technical replicates (motility agar plates) each (indicated by individual points). Mean and standard error are shown. One-way ANOVA was used as a statistical test, and only significant and relevant comparisons are shown. ****p<0.00005.

Figure 5. Chromosomal Walker mutants of *minD2* disrupt polar localisation of CetZ1. H26 wildtype and *minD2* walker mutant strains containing pHVID135 (for expression of CetZ1-mTq2) were grown in Hv-Cab with 1 mM L-Tryptophan and sampled at $OD_{600} \sim 0.1$. **A)** Cells were imaged using epifluorescence to observe CetZ1-mTq2 localisation. Scale bar 5 µm. **B)** Heat map representations of detected foci of CetZ1-mTq2 and their localisations. Heatmaps are coloured by density, referring to subcellular localisation along the relative length of lateral and longitudinal axes only and do not fluorescence intensity. **C)** normalised median fluorescence intensity of CetZ1-mTq2 along the long axis of the cell, represented as the mean (bold line) and standard deviation (shading) of biological replicates (as in Fig. 2d). Two biological replicates were carried out for the wildtype background, and four for each of *minD2WA** and *minD2WB**, and data for H26 wildtype is included on both graphs for reference. The number of cells measured from pooled biological replicates is as follows: WT + CetZ1-mTq2, n=2380; *minD2WA** + CetZ1-mTq2, n=2334; *minD2WB** + CetZ1-mTq2, n=1830.

Figure 6. Schematic model for assembly of motility proteins at cell poles. A) MinD4 oscillates
between cell poles and forms cap-like localisations at cell poles in an ATPase dependent manner [28].
B) MinD4 [28] and MinD2 direct protein constituents of the archaellum ("Arl") and chemosensory
arrays ("Che") such as ArlD and CheW, respectively, to cell poles [56]. CetZ1 is also directed to cell
poles, likely by both MinD paralogues, but more dominantly by MinD2. **C)** At cell poles, CetZ1 is
hypothesised to assemble as a sheet or disc-like structure, **D)** where it may contribute to assembly
and positioning of motility structures via an unknown mechanism [42].

References

[1] D.D. Leipe, Y.I. Wolf, E.V. Koonin, and L. Aravind, Classification and evolution of P-loop GTPases and related ATPases. Journal of molecular biology 317 (2002) 41-72.

[2] J.S. Schuhmacher, K.M. Thormann, and G. Bange, How bacteria maintain location and number of flagella? FEMS microbiology reviews 39 (2015) 812-822.

[3] J.S. Schuhmacher, F. Rossmann, F. Dempwolff, C. Knauer, F. Altegoer, W. Steinchen, A.K. Dörrich, A. Klingl, M. Stephan, and U. Linne, MinD-like ATPase FlhG effects location and number of bacterial flagella during C-ring assembly. Proceedings of the National Academy of Sciences 112 (2015) 3092-3097.

[4] S. Ringgaard, K. Schirner, B.M. Davis, and M.K. Waldor, A family of ParA-like ATPases promotes cell pole maturation by facilitating polar localization of chemotaxis proteins. Genes & Development 25 (2011) 1544-1555.

[5] M.A. Roberts, G.H. Wadhams, K.A. Hadfield, S. Tickner, and J.P. Armitage, ParA-like protein uses nonspecific chromosomal DNA binding to partition protein complexes. Proceedings of the National Academy of Sciences 109 (2012) 6698-6703.

[6] A.S. Jalal, and T.B. Le, Bacterial chromosome segregation by the ParABS system. Open biology 10 (2020) 200097.

[7] J.C. Baxter, and B.E. Funnell, Plasmid partition mechanisms. Plasmids: biology and impact in biotechnology and discovery (2015) 133-155.

[8] J. Lutkenhaus, Assembly dynamics of the bacterial MinCDE system and spatial regulation of the Z ring. Annu. Rev. Biochem. 76 (2007) 539-562.

[9] Z. Hu, and J. Lutkenhaus, Topological regulation of cell division in E. coli: spatiotemporal oscillation of MinD requires stimulation of its ATPase by MinE and phospholipid. Molecular cell 7 (2001) 1337-1343.

[10] J. Lutkenhaus, The ParA/MinD family puts things in their place. Trends in microbiology 20 (2012) 411-418.

[11] B.A. Perez‐Cheeks, P.J. Planet, I.N. Sarkar, S.A. Clock, Q. Xu, and D.H. Figurski, The product of tadZ, a new member of the parA/minD superfamily, localizes to a pole in Aggregatibacter actinomycetemcomitans. Molecular microbiology 83 (2012) 694-711.

[12] G. Laloux, and C. Jacobs-Wagner, How do bacteria localize proteins to the cell pole? Journal of cell science 127 (2014) 11-19.

[13] S.R. Thompson, G.H. Wadhams, and J.P. Armitage, The positioning of cytoplasmic protein clusters in bacteria. Proceedings of the National Academy of Sciences 103 (2006) 8209-8214.

[14] D.F. Savage, B. Afonso, A.H. Chen, and P.A. Silver, Spatially ordered dynamics of the bacterial carbon fixation machinery. Science 327 (2010) 1258-1261.

[15] M. Loose, E. Fischer-Friedrich, C. Herold, K. Kruse, and P. Schwille, Min protein patterns emerge from rapid rebinding and membrane interaction of MinE. Nature structural & molecular biology 18 (2011) 577-583.

[16] M. Howard, and K. Kruse, Cellular organization by self-organization: mechanisms and models for Min protein dynamics. The Journal of cell biology 168 (2005) 533.

[17] T.D. Dunham, W. Xu, B.E. Funnell, and M.A. Schumacher, Structural basis for ADP‑mediated transcriptional regulation by P1 and P7 ParAn. The EMBO journal 28 (2009) 1792-1802.

[18] E. Bi, K. Dai, S. Subbarao, B. Beall, and J. Lutkenhaus, FtsZ and cell division. Research in microbiology 142 (1991) 249-252.

[19] H.P. Erickson, FtsZ, a prokaryotic homolog of tubulin? Cell 80 (1995) 367-370.

[20] E. Nogales, K.H. Downing, L.A. Amos, and J. Löwe, Tubulin and FtsZ form a distinct family of GTPases. Nature Structural and Molecular Biology 5 (1998) 451.

[21] D.M. Raskin, and P.A. De Boer, Rapid pole-to-pole oscillation of a protein required for directing division to the middle of Escherichia coli. Proceedings of the National Academy of Sciences 96 (1999) 4971-4976.

[22] P.A. De Boer, R.E. Crossley, and L.I. Rothfield, A division inhibitor and a topological specificity factor coded for by the minicell locus determine proper placement of the division septum in E. coli. Cell 56 (1989) 641-649.

[23] L.L. Lackner, D.M. Raskin, and P.A. De Boer, ATP-dependent interactions between Escherichia coli Min proteins and the phospholipid membrane in vitro. Journal of bacteriology 185 (2003) 735-749.

[24] K.-T. Park, W. Wu, K.P. Battaile, S. Lovell, T. Holyoak, and J. Lutkenhaus, The Min oscillator uses MinD-dependent conformational changes in MinE to spatially regulate cytokinesis. Cell 146 (2011) 396-407.

[25] T.H. Szeto, S.L. Rowland, L.I. Rothfield, and G.F. King, Membrane localization of MinD is mediated by a C-terminal motif that is conserved across eubacteria, archaea, and chloroplasts. Proceedings of the National Academy of Sciences 99 (2002) 15693-15698.

[26] H. Zhou, R. Schulze, S. Cox, C. Saez, Z. Hu, and J. Lutkenhaus, Analysis of MinD mutations reveals residues required for MinE stimulation of the MinD ATPase and residues required for MinC interaction. Journal of bacteriology 187 (2005) 629-638.

[27] J. Lutkenhaus, and M. Sundaramoorthy, MinD and role of the deviant Walker A motif, dimerization and membrane binding in oscillation. Molecular microbiology 48 (2003) 295-303.

[28] P. Nußbaum, S. Ithurbide, J.C. Walsh, M. Patro, F. Delpech, M. Rodriguez-Franco, P.M. Curmi, I.G. Duggin, T.E. Quax, and S.-V. Albers, An oscillating MinD protein determines the cellular positioning of the motility machinery in archaea. Current Biology 30 (2020) 4956-4972. e4.

[29] M. Tripepi, S. Imam, and M. Pohlschroder, Haloferax volcanii flagella are required for motility but are not involved in PibD-dependent surface adhesion. Journal of bacteriology 192 (2010) 3093-3102.

[30] M. Tripepi, J. You, S. Temel, Ö. Önder, D. Brisson, and M. Pohlschröder, N-glycosylation of Haloferax volcanii flagellins requires known Agl proteins and is essential for biosynthesis of stable flagella. Journal of bacteriology 194 (2012) 4876-4887.

[31] M.F. Abdul Halim, F. Pfeiffer, J. Zou, A. Frisch, D. Haft, S. Wu, N. Tolić, H. Brewer, S.H. Payne, and L. Paša‑Tolić, H aloferax volcanii archaeosortase is required for motility, mating, and C‑terminal processing of the S‑layer glycoprotein. Molecular microbiology 88 (2013) 1164-1175.

[32] M.F. Abdul Halim, S. Schulze, A. DiLucido, F. Pfeiffer, A.W. Bisson Filho, and M. Pohlschroder, Lipid Anchoring of Archaeosortase Substrates and Midcell Growth in Haloarchaea. mBio 11 (2020) e00349-20.

[33] R.N. Esquivel, and M. Pohlschroder, A conserved type IV pilin signal peptide H – domain is critical for the post‐translational regulation of flagella‐dependent motility. Molecular microbiology 93 (2014) 494-504.

[34] M. Cerletti, M.J. Martínez, M.I. Giménez, D.E. Sastre, R.A. Paggi, and R.E. De Castro, The LonB protease controls membrane lipids composition and is essential for viability in the extremophilic haloarchaeon H aloferax volcanii. Environmental microbiology 16 (2014) 1779-1792.

[35] P. Chatterjee, M. Garcia, J. Cote, K. Yun, G. Legerme, R. Habib, M. Tripepi, C. Young, D.W. Kulp, and M. Dyall-Smith, Involvement of ArlI, ArlJ, and CirA in Archaeal Type-IV Pilin-Mediated Motility Regulation. bioRxiv (2024) 2024.03. 04.583388.

[36] R.K. Hackley, S. Hwang, J.T. Herb, P. Bhanap, K. Lam, A. Vreugdenhil, C.L. Darnell, M.M. Pastor, J.H. Martin, and J.A. Maupin‐Furlow, TbsP and TrmB jointly regulate gapII to influence cell development phenotypes in the archaeon Haloferax volcanii. Molecular Microbiology (2024).

[37] I.G. Duggin, C.H. Aylett, J.C. Walsh, K.A. Michie, Q. Wang, L. Turnbull, E.M. Dawson, E.J. Harry, C.B. Whitchurch, and L.A. Amos, CetZ tubulin-like proteins control archaeal cell shape. Nature 519 (2015) 362.

[38] H. Schiller, Y. Hong, J. Kouassi, T. Rados, J. Kwak, A. DiLucido, D. Safer, A. Marchfelder, F. Pfeiffer, and A. Bisson, Identification of structural and regulatory cell-shape determinants in Haloferax volcanii. Nature communications 15 (2024) 1414.

[39] C.H. Aylett, and I.G. Duggin, The tubulin superfamily in archaea, Prokaryotic Cytoskeletons, Springer, 2017, pp. 393-417.

[40] H.J. Brown, and I.G. Duggin, Diversity and Potential Multifunctionality of Archaeal CetZ Tubulin-like Cytoskeletal Proteins. Biomolecules 13 (2023) 134.

[41] R.T. de Silva, V. Shinde, H.J. Brown, Y. Liao, and I.G. Duggin, Dynamic self-association of archaeal tubulin-like protein CetZ1 drives Haloferax volcanii morphogenesis. bioRxiv (2024) 2024.04. 08.588506.

[42] H.J. Brown, M.I. Islam, J. Ruan, M.A. Baker, S. Ithurbide, and I.G. Duggin, CetZ1-dependent assembly and positioning of the motility machinery in haloarchaea. bioRxiv (2024) 2024.05. 02.592137.

[43] R.T. de Silva, M.F. Abdul-Halim, D.A. Pittrich, H.J. Brown, M. Pohlschroder, and I.G. Duggin, Improved growth and morphological plasticity of Haloferax volcanii. Microbiology (Reading) (2021).

[44] M. Patro, I.G. Duggin, S.V. Albers, and S. Ithurbide, "Influence of plasmids, selection markers and auxotrophic mutations on Haloferax volcanii cell shape plasticity". Front Microbiol 14 (2023) 1270665.

[45] S.W. Cline, W.L. Lam, R.L. Charlebois, L.C. Schalkwyk, and W.F. Doolittle, Transformation methods for halophilic archaebacteria. Canadian Journal of Microbiology 35 (1989) 148-152.

[46] T. Allers, H.-P. Ngo, M. Mevarech, and R.G. Lloyd, Development of additional selectable markers for the halophilic archaeon Haloferax volcanii based on the leuB and trpA genes. Applied and environmental microbiology 70 (2004) 943-953.

[47] T. Allers, S. Barak, S. Liddell, K. Wardell, and M. Mevarech, Improved strains and plasmid vectors for conditional overexpression of His-tagged proteins in Haloferax volcanii. Appl Environ Microbiol 76 (2010) 1759-69.

[48] B. Schmid, P. Tripal, T. Fraaß, C. Kersten, B. Ruder, A. Grüneboom, J. Huisken,
and R. Palmisano, 3Dscript: animating 3D/4D microscopy data using a
natural-language-based syntax. Nature methods 16 (2019) 278-280.
[49] A. Ducret, E.M. Quardokus, and Y.V. Brun, MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. Nature microbiology 1 (2016)
1-7.
[50] J. Schindelin, I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S.
Preibisch, C. Rueden, S. Saalfeld, and B. Schmid, Fiji: an open-source
platform for biological-image analysis. Nature methods 9 (2012) 676.
[51] J. Babski, K.A. Haas, D. Nather-Schindler, F. Pfeiffer, K.U. Forstner, M.
Hammelmann, R. Hilker, A. Becker, C.M. Sharma, A. Marchfelder, and J.
Soppa, Genome-wide identification of transcriptional start sites in the
haloarchaeon Haloferax volcanii based on differential RNA-Seq (dRNA-Seq).
BMC Genomics 17 (2016) 629.
[52] M. Cerletti, M.I. Gimenez, C. Troetschel, D.A. C, A. Poetsch, R.E. De Castro,
and R.A. Paggi, Proteomic Study of the Exponential-Stationary Growth Phase
Transition in the Haloarchaea Natrialba magadii and Haloferax volcanii.
Proteomics 18 (2018) e1800116.
[53] S. Laass, V.A. Monzon, J. Kliemt, M. Hammelmann, F. Pfeiffer, K.U. Forstner,
and J. Soppa, Characterization of the transcriptome of Haloferax volcanii,
grown under four different conditions, with mixed RNA-Seq. PLoS One 14
(2019) e0215986.
[54] Z. Li, Y. Kinosita, M. Rodriguez-Franco, P. Nußbaum, F. Braun, F. Delpech, T.E.
Quax, and S.-V. Albers, Positioning of the motility machinery in halophilic
archaea. MBio 10 (2019) e00377-19.
[55] S. Ithurbide, R.T. de Silva, H.J. Brown, V. Shinde, and I.G. Duggin, A vector
system for single and tandem expression of cloned genes and multi-colour
fluorescent tagging in Haloferax volcanii. Microbiology 170 (2024).
[56] M. Patro, S. Sivabalasarma, S. Gfrerer, M. Rodriguez-Franco, P. Nußbaum, S.
Ithurbide, and S.-V. Albers, MinD2 modulates cell shape and motility in the
archaeon *Haloferax volcanii*. bioRxiv (2024) 2024.08.01.606218.

654