Rapamycin increases murine lifespan but does not reduce mineral volume in the Matrix GLA Protein (MGP) knockout mouse model of medial arterial calcification.

Parya Behzadi[1], Rolando A. Cuevas[1], Alex Crane[1], Andrew A Wendling[1], Claire C. Chu[1], William J Moorhead III[1], Ryan Wong[1], Mark Brown[1], Joshua Tamakloe[1], Swathi Suresh[1], Payam Salehi[2], Iris Z. Jaffe[3], Allison L. Kuipers[4], Lyudmila Lukashova[5], Konstantinos Verdelis[5], and Cynthia St. Hilaire[1, 6]

[1] Department of Medicine, Division of Cardiology, and the Pittsburgh Heart, Lung, Blood and Vascular Medicine Institute, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

[2] CardioVascular Center, Vascular Surgery, Tufts Medical Center, 800 Washington Street, Boston, MA, 02111-1800, USA

[3] Molecular Cardiology Research Institute, Tufts Medical Center, 800 Washington Street, Boston, MA, 02111-1800, USA

[4] Department of Epidemiology, School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

[5] Departments of Endodontics and Oral Biology, School of Dental Medicine, University of Pittsburgh, Pittsburgh, PA, USA

[6] Department of Bioengineering, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

ABSTRACT: 235 words

MANUSCRIPT:

KEYWORDS: Medial Arterial Calcification, Rapamycin, Matrix GLA protein

ABBREVIATIONS

Adenosine monophosphate	AMP
Adenosine triphosphate	ATP
Alkaline phosphate	TNAP
Arterial calcification due to deficiency of CD73	ACDC
Ecto-5'-nucleotidase	NT5E
Induced pluripotent stem cells	iPSCs
Matrix GLA protein	MGP
Medial arterial calcification	MAC
mTORC complex 1	mTORC1
mTORC complex 2	mTORC2
Peripheral artery disease	PAD
Smooth muscle α-actin	SMA
Smooth muscle cells	SMCs
Myosin heavy chain 11	MYH11
Runt-related transcription factor 2	RUNX2

ABSTRACT

Peripheral artery disease (PAD) is the narrowing of the arteries that carry blood to the lower extremities. PAD has been traditionally associated with atherosclerosis. However, recent

studies have found that medial arterial calcification (MAC) is the primary cause of chronic limb
ischemia below the knee. MAC involves calcification of the elastin fibers surrounding smooth
muscle cells (SMCs) in arteries. Matrix GLA Protein (MGP) binds circulating calcium and inhibits
vascular calcification. $Mgp^{-/-}$ mice develop severe MAC and die within 8 weeks of birth due to
aortic rupture or heart failure. We previously discovered a rare genetic disease Arterial
Calcification due to Deficiency in CD73 (ACDC) in which patients present with extensive MAC in
their lower extremity arteries. Using a patient-specific induced pluripotent stem cell model we
found that rapamycin inhibited calcification. Here we investigated whether rapamycin could
reduce MAC in vivo using $Mgp^{-/-}$ mice as a model. $Mgp^{+/+}$ and $Mgp^{-/-}$ mice received 5mg/kg
rapamycin or vehicle. Calcification content was assessed via microCT, and vascular
morphology and extracellular matrix content assessed histologically. Immunostaining and
western blot analysis were used to examine SMC phenotypes and cellular functions. Rapamycin
prolonged $Mgp^{-/-}$ mice lifespan, decreased mineral density in the arteries, and increased smooth
muscle actin protein levels, however, calcification volume, vessel morphology, SMC
proliferation, and autophagy flux were all unchanged. These findings suggest that rapamycin's
effects in the $Mgp^{-/-}$ mouse are independent of the vascular phenotype.

68
INTRODUCTION
Peripheral artery disease (PAD) is the narrowing of blood vessels in the lower extremities due to
inward remodeling or thrombotic occlusion, both of which cause chronic limb ischemia and often
result in amputation of the lower leg (1). Traditionally atherosclerosis was the assumed cause of
PAD however recent studies now show that medial arterial calcification (MAC) promotes inward
remodeling, and stiffness related to medial dysplasia induces thrombosis (2-4). MAC is
characterized by the progressive buildup of calcium and phosphate within the arterial walls and
is typically not associated with lipid deposition, fibrous cap formation, or intimal hyperplasia (5).
These structural features distinguish MAC from atherosclerosis and indicate that MAC is a
distinct pathology driving adverse outcomes in PAD. Currently, there are no specific medical
therapies that target the pathogenesis of PAD or MAC. Finding therapies that can prevent, stop,
or even reverse MAC in PAD could greatly enhance the current standard of care.
81
Vascular calcification stems from the nucleation of calcium and phosphate into hydroxyapatite
crystals and the release of pro-mineralizing matrix vesicles from osteogenic-like arterial smooth
muscle cells (SMCs) that accumulate along the elastic lamina (6). Phosphate is a byproduct of
the breakdown of extracellular adenosine triphosphate (ATP) to adenosine, generating inorganic
phosphate at several points (7, 8). There are a number of genetic diseases that present with
MAC which harbor inactivating mutations in the genes related to extracellular ATP metabolism
(5). Patients with Arterial Calcification due to Deficiency of CD73 (ACDC, also known as CALJA,
OMIM # 211800) harbor inactivating mutations in the $NT5E$ gene, which encodes for the CD73
enzyme that metabolizes extracellular AMP to adenosine and inorganic phosphate. Key
signatures of this disease are calcification nodules in the small joint capsules and MAC in the
lower-extremity arteries that initiate along the elastic lamina (4, 9, 10). We previously found that
a lack of CD73-mediated adenosine enhanced the expression and activity of tissue-nonspecific
alkaline phosphatase (TNAP), a key enzyme that promotes calcification (11).
95
Matrix GLA protein (MGP) is a vitamin-K2-dependent protein with an unusual gamma-
carboxylation of five glutamate residues that enhance its affinity for calcium, preventing it from
mineralizing (12). MGP also inhibits calcification propagation by binding to hydroxyapatite
crystals and stimulating their uptake by local phagocytosing macrophages (13). MGP also
modulates osteogenic signaling by binding to BMP-2, preventing it from activating receptors and
upregulating RUNX2, the key regulator of osteoblast differentiation and maturation (14, 15).
Mice that lack MGP develop to term but die prematurely within approximately two months of age

due to extensive MAC, which leads to aortic rupture or heart failure (16). Aortas of MGP-
deficient mice exhibit increased collagen accumulation and elastin fiber fragmentation within the
medial layer of the arteries where calcification is also localized. This is phenotypically similar to
MAC observed in human patients with PAD MAC and ACDC patients (4).
107
The drug rapamycin inhibits the mammalian target of rapamycin (mTOR), a protein kinase
broadly expressed throughout the body, which controls many key processes such as energy
balance, autophagy, and proliferation (17-20). Rapamycin protects against the calcification of
vascular cells in in vitro and ex vivo models, as well as in in vivo models of chronic kidney
disease-induced MAC (21-23). We previously found rapamycin prevented calcification in in vitro
and in vivo disease models of ACDC patient-specific induced pluripotent stem cells (iPSCs)
(11).
115
116 While CD73-deficient humans develop extensive MAC in their lower extremity large vessels,
117 CD73-deficient mice do not phenocopy humans and do not exhibit calcification in their
118 vasculature.(24, 25) As the MAC observed in MGP-deficient mice mirrors that seen in ACDC
119 patients, we sought to investigate whether rapamycin prevented calcification in this genetic
120 mouse model of MAC.
121
MATERIALS AND METHODS

Availability of Materials
We abide by the NIH Grants Policy on Sharing of Unique Research Resources, including the
NIH Policy on Sharing of Model Organisms for Biomedical Research (2004), NIH Grants Policy
Statement (2003), and Sharing of Biomedical Research Resources: Principles and Guidelines
for Recipients of NIH Grants and Contracts (1999), and the Bayh-Dole Act and the Technology
Transfer Commercialization Act of 2000. Materials generated in our laboratory are made
available for non-commercial research per established University of Pittsburgh Office of
Research IRB and MTA protocols.
131
132 *Institutional Review Board Statement*
133 De-identified human tissues were obtained, with informed consent from subjects, who were
134 enrolled in studies approved by the University of Pittsburgh or Tufts Medical Center institutional
135 review boards, per the Declaration of Helsinki.
136
Animals and cell line generation
Animal use was approved by the Institutional Animal Care and Use Committee at the University
of Pittsburgh. MGP heterozygous mutant mice from a C57BL/6J background (strain# 023811
Jackson Laboratory, Bar Harbor, ME) were bred to produce +/+ and -/- littermate controls.
Smooth muscle lineage-specific *Raptor* ($Mgp^{-/-}; Raptor^{SMC-/-}$) and *Rictor* ($Mgp^{-/-}; Rictor^{SMC-/-}$)
knockout mice were produced by breeding $Mgp^{+/-}$ with Myh11-Cre-eGFP mice (strain# 007742
Jackson Laboratory, Bar Harbor, ME)(26), with Raptor-LoxP (strain# 013188 Jackson
Laboratory, Bar Harbor, ME)(27), or *Rictor*-LoxP (strain# 020649 Jackson Laboratory, Bar
Harbor, ME)(28). Genotypes were determined from tail snips that were incubated in DirectPCR
Lysis Reagent (Viagen Biotech) according to manufacturer instructions. The lysate was
amplified using OneTaq MasterMix (New England Biolabs) according to manufacturer
instructions with primers suggested by Jackson Laboratories. The PCR reaction was run on an
ethidium bromide agarose gel. SMC were obtained from three- to five-week-old male and
female $Mgp^{+/+}$ and $Mgp^{-/-}$ mice.

Cell culture

Mgp and *Mgp* aortic SMC lines from male and female murine aorta were cultured in
Dulbecco's modified Eagle's medium (DMEM; Gibco, Waltham, MA) supplemented with 20%
FBS (FBS; R&D Systems, Minneapolis, MN) and 100 U/mL penicillin-streptomycin (P/S; Gibco,
Waltham, MA). Growth media was changed every three days, and cells were split 1:2 when
confluent. Post-expansion, 25,000–50,000 cells/cm^2 were plated with serum reduced 10% FBS
in DMEM medium and grown to confluence. Before all experiments/treatments, cells were
serum-starved in DMEM with 0.5% FBS for 48 h. Incubation under reduced serum conditions
has been proven to be a beneficial technique for studying smooth muscle cell remodeling and
contractility in vitro (29, 30). Cells were treated with rapamycin (Novus Biologicals) at
concentrations of 200 nM or the same volume of DMSO as vehicle control for 12h. For
autophagy flux assessment, cells were exposed to either bafilomycin A1 100 nM (Sigma-
Aldrich) or rapamycin 200 nM for a duration of 24h following treatment intervention. All
chemicals were dissolved in DMSO and administered in equal volume as vehicle control.
165

Tissue Extraction
Mice were sacrificed by asphyxiation with carbon dioxide, followed by perforation of the
diaphragm. Tissue extraction began with dissection through the abdominal wall and then
perfusion of the heart with 50 mL of DPBS supplemented with 1:100 Amphotericin B (15290026,
Gibco) and 1:200 Gentamicin (15710064, Gibco). The aortas were maintained in this solution
while the adventitia was removed in a petri dish. Tissues were then fixed in 4% PFA in PBS for
2 hours and embedded in paraffin.
173

In Vivo Rapamycin Injections
Male and female *Mgp* mice received intraperitoneal injections of either vehicle or 5 mg/kg
rapamycin dissolved in 0.3 mg/ml peanut oil once a week beginning at 10 days of age. This
process continued until the natural death of all mice in each group. To assess the volume and
density of calcification, male and female *Mgp* mice received intraperitoneal injections of either
vehicle, 1x/week rapamycin or 3x/week rapamycin from 10 to 20 days of age.
180

Micro-Computed Tomography
A Scanco 30µCT microCT scanner (Scanco Medical, Bassersdorf, Switzerland) with a 45 kVp
beam energy (1,000 ms exposure) was used to assess the volume and density of calcification in
mouse aorta ex vivo, as previously described (11, 31). No filtering was used, allowing for
visibility of lower CT densities. Images were then analyzed for relative calcified mass within the
aorta (expressed as bone volume fraction) using the Scanco 3D Morphometry and Densitometry
Analysis software (Scanco Medical) after binarization of the images with a global threshold
within a user-defined aorta region of interest.
189

Western Blot Analysis
Cells were lysed in 1% CHAPS hydrate, 150 mmol/L sodium chloride, 25 mmol/L HEPES buffer
supplemented with 1× protease and phosphatase inhibitor (Sigma-Aldrich). Cells were scraped
into microcentrifuge tubes, vortexed for 5 minutes, freeze/thawed for 5 to 8 cycles, then
centrifuged at 12 000×g for 10 minutes at 4°C. Supernatant protein concentration was
determined using Pierce BCA protein assay kit (Thermo Fisher, Waltham, MA). Ten micrograms
of protein were used in the preparation of lysate with 1 × Pierce LDS sample buffer nonreducing
(Thermo Fisher, Waltham, MA) and 1 × NuPAGE sample reducing agent (Novex, Waltham,
MA). Lysates were denatured at 95°C for 15 min and then electrophoresed on 4%–20% TGX
stain-free polyacrylamide gel (Bio-Rad, Hercules, CA) in 1 × Tris/Glycine/SDS buffer (Bio-Rad,
Hercules, CA) at 120 V for 50 min. Protein was transferred onto a 0.2 µm nitrocellulose
membrane in prepared 1 × Towbin buffer with ethanol (EtOH) at 1 A and 25 V for 30 min using
the Trans-Blot Turbo Transfer System (Bio-Rad, Hercules, CA). Membranes were blocked in 1:1

Odyssey blocking buffer (Li-COR, Lincoln, NE) and PBS for 1 h at room temperature, followed by primary antibody incubation in 1:1 Odyssey blocking buffer and PBS plus 0.1% Tween 20 (PBS-T) at 4°C overnight. Membranes were washed in PBS-T three times for 5 min, then incubated in secondary antibody at room temperature for 1 h. Membranes belonging to the same experimental set were imaged simultaneously on an Odyssey CLx (LI-COR, Lincoln, NE), and band intensity quantification was performed with Image Studio (Version 5.2, LI-COR, Lincoln, NE) software. Individual bands were normalized to α-tubulin, and each treatment group's fold change was compared with each gel's vehicle control lanes. For sequential antibody incubations, membranes were stripped in 1 × NewBlot Nitro Western Blot Stripping Buffer (LI-COR, Lincoln, NE) for 10 min, followed by three washes in PBS.

Verhoeff Van Gieson, Von Kossa and Masson's Trichrome Staining
Mice aortic tissues were removed from mice through a modified bilateral thoracosternotomy under anesthesia conditions. Then aortic tissue was fixed with 10% paraformaldehyde and embedded in paraffin. The embedded specimens were transversely sectioned at 10 μm on a microtome cryostat (Microm HM 325). Slides with the adhered paraffin aortic sections were warmed to 65°C for 1 h and then deparaffinized through xylene, rehydrated with serial incubation in graded alcohol baths, and stained with Verhoeff–van Gieson for elastic fiber visualization (Polysciences, 25089-1) or Von Kossa for calcification using the Von Kossa Method of Calcium Kit (Polysciences, 24633-1) and Masson's for collagen fibers (Polysciences, 25088-1), according to manufacturer's instructions. Images were captured using PreciPoint scanner. The thickness of the artery was measured using ImageJ software (Java 8), where the medial thickness was determined by measuring the distance between the inner and outer elastic lamina at four sites per vessel. Measurements were normalized to the scale bar of each picture. Data is shown as a total pixel's length. The length of elastic fibers was measured by selecting three random areas within the vessel and using ImageJ to trace and measure each fiber. Measurements were normalized to the area unit square.

Immunofluorescence
Slides with the adhered paraffin aortic sections were warmed to 65°C for 1 h and then deparaffinized through xylene, rehydrated with serial incubation in graded alcohol baths. Slides were boiled in citric acid-based antigen retrieval solution (H-3300, Vector Labs) for 20 minutes. Slides were cooled in the unmasking solution for 1 hour, washed in 1x PBS and placed in blocking buffer (500 mL PBS 0.3g Fish Skin Gelatin) for 1 hour. Tissues were incubated overnight at 4°C with antibodies for MYH11 (Abnova, mab34251, 1:50) and ACTA2 (Abcam, ab5694, 1:50), RUNX2 (ABclonal, A2851, 1:100), LC3 (cell signaling,124714, 1:50), Ki67 (Abcam, ab15580, 1:250). All signals were optimized to IgG (Vector, 31235, 1:50). Cells were washed 3x for 5 minutes each with PBS, 0.1% TWEEN 20, then once with PBS, and then incubated with secondary antibody (A11006, A11012, Thermo Fisher) for 1 hour. Cells were washed 3x for 5 minutes each with PBS, 0.1% TWEEN 20, then once with PBS, then mounted with Fluoroshield Mounting Medium with DAPI (Ab104139, Abcam). Slides were imaged within 24 hours of mounting. Cells were imaged within 24 hours of mounting. In ImageJ, each image was split into blue, red, and green channels, and the pixel intensity of the red and blue channels was measured (32). The measured intensity of a given stain was normalized to DAPI intensity of that same image. We obtained the von Kossa staining images using PreciPoint scanner and the immunofluorescent image using a Nikon NI-E I microscope.

Statistics
Statistical analysis was performed with GraphPad Prism 9.2 (GraphPad Software, Inc) and data shown are mean±SD. Statistical significance between groups was assessed using the Kruskal Wallis test or by Two-way ANOVA according to the data set. Statistical analysis used, exact n

254   values, biological replicates, and P values are stated within each figure legend. A p-value equal
255   to or less than 0.05 will be considered statistically significant.
256
257   **RESULTS**
258
MAC in MGP-deficient mice phenocopies human MAC.
The main characteristic of MAC is calcification in the media along the elastic lamina. Von Kossa
staining for calcification illustrates that $Mgp^{-/-}$ mouse aortas have extensive mineralization in the
medial layer, which phenocopies the mineralization seen in MAC of the human tibial arteries
and is distinct from neointimal calcification found in the necrotic core of coronary artery with
atherosclerotic plaque (Figure 1A). MGP is a potent inhibitor of soft tissue calcification. $Mgp^{-/-}$
mice do not produce any MGP protein (Figure 1B) and, as a result, develop extensive MAC
pathology in large vessels.
267
Rapamycin reduces mineral density in $Mgp^{-/-}$ mice.
It has been shown that the $Nt5e$-knockout mice model shows altered renal function, diminished
control of the glomerular arteriolar tone, elevated atherogenesis, elevated thrombotic occlusion,
and elevated hypoxia-induced vascular leakage.(24) However, they do not develop ectopic
bone formation, do mimic the human vascular calcification phenotype, and do not have impaired
formation of bone and teeth, while CD73-deficient ACDC patients develop extensive ectopic
calcification (24, 25, 33). Jin et al. circumvented this issue using a patient-specific in vitro and in
vivo disease modeling system using iPSC technology. They demonstrated that patient-specific
ACDC-iPSCs developed extensive calcification in the in vivo teratoma model relative to control
patient iPSCs. Furthermore, they found that treating mice bearing ACDC-iPSC teratomas with
rapamycin reduced calcification (11).
279
As CD73-deficient mice do not sufficiently recapitulate the MAC observed in ACDC patients,
here we used the $Mgp^{-/-}$ mice as an in vivo model of MAC. Mice were administered 5 mg/kg
rapamycin injected three times per week starting at 10 days old until natural death (Figure 2A).
We discovered that rapamycin treatment extended the lifespan of $Mgp^{-/-}$ mice (T_{50} 61 days)
compared to the $Mgp^{-/-}$ mice treated with the DMSO vehicle (T_{50} 35 days), which start to die as
early as day 20 (Figure 2B). Rapamycin's effects on extending lifespan has been observed in
genetically heterogeneous non-diseased mice (34). Both male (n=11) and female (n=8) were
used however no sex differences were observed between $Mgp^{-/-}$ mice that had received either
vehicle or rapamycin treatment.
289
We also evaluated whether a low-frequency regime of rapamycin administration (one dose of 5
mg/kg injected weekly) versus a high-frequency regime of 5 mg/kg injected three times per
week affected the $Mgp^{-/-}$ vascular mineral density at day 20, before natural death (Figure 3A).
Whole aorta microCT analysis found no differences in the total calcification volume in either
treatment regime; however, mice receiving a high regime of rapamycin exhibited decreased
mineral density compared to the control group (Figure 3B). Von Kossa stain showed that the
aortas of $Mgp^{+/+}$ mice exhibit no mineralization, while both vehicle- and rapamycin-treated $Mgp^{-/-}$
mice showed extensive calcification with no discernible differences between vehicle or
rapamycin groups (Figure 3C). Medial wall thickness was not different between $Mgp^{+/+}$ and $Mgp^{-/-}$
that had received vehicle or rapamycin treatment (Figure 3C). RUNX2 is a key transcription
factor promoting vascular cell osteogenic differentiation (6). Immunofluorescent staining
revealed that while $Mgp^{-/-}$ mice showed elevated levels of RUNX2 compared to $Mgp^{+/+}$ mice,
rapamycin treatment did not reduce RUNX2 expression in $Mgp^{-/-}$ compared to $Mgp^{-/-}$ treated with
vehicle (Figure 3D). These findings suggest that while rapamycin doubles the lifespan of $Mgp^{-/-}$
mice, it has little to no effect in protecting the $Mgp^{-/-}$ mouse vasculature from calcification.

305
SMC phenotype remains unchanged after rapamycin treatment in vivo
The drug rapamycin has a broad spectrum of effects on SMCs. Rapamycin inhibits SMC migration and proliferation by inducing the cyclin-dependent kinase inhibitors p27^{kip} and p21^{cip} and promotes G1-S cell cycle arrest (35, 36). Rapamycin also induces SMC differentiation by inhibiting the mTOR-target S6K1 via AKT activation, promoting SMC contractile phenotype by enhancing SMA and MYH11 protein expression (37). Further, rapamycin was shown to protect against calcification of vascular cells in in vitro and ex vivo models (11, 21). While rapamycin did not reduce calcification volume in $Mgp^{-/-}$ vessels we hypothesized that it could perhaps prevent aortic rupture by maintaining SMC contractile phenotype.

Masson's trichrome staining further illustrates that $Mgp^{-/-}$ vessels are highly remodeled, with acellular areas exhibiting observable increased collagen staining (Figure 4A). Verhoeff-Van Gieson (VVG) staining revealed that the aortas of $Mgp^{-/-}$ mice exhibited less robust staining for elastin, and less tortuous fibers, in $Mgp^{-/-}$ compared to $Mgp^{+/+}$ However, rapamycin did not alter elastin fiber length in $Mgp^{-/-}$ mice (Figure 4B). Immunofluorescent staining was used to quantify the SMC contractility markers smooth muscle α-actin (SMA) and myosin heavy chain 11 (MYH11) (38, 39). In $Mgp^{+/+}$ we observed no differences in SMA levels between vehicle and rapamycin treatment, nor between $Mgp^{+/+}$ the $Mgp^{-/-}$ mice treated with vehicle. However, rapamycin treatment slightly, but significantly, increased SMA levels in $Mgp^{-/-}$ mice compared to vehicle alone. No differences in MYH11 levels were observed (Figure 4C). As the medial layer is remodeled in the $Mgp^{-/-}$ mice, we measured the mitotic marker Ki67 and found that at baseline, Ki67 levels were significantly elevated in $Mgp^{-/-}$ aorta while nearly undetectable in the $Mgp^{+/+}$. However, rapamycin treatment did not alter Ki67 levels in either of the two genotypes (Figure 4D). Together, these data suggest that rapamycin does not significantly alter the medial remodeling or SMC phenotype in $Mgp^{-/-}$ mice aorta.

Rapamycin lowered LC3 staining in $Mgp^{-/-}$ aorta but there were no observable changes in autophagy flux.
Autophagy is a ubiquitous process and contributes to bone development and the differentiation of osteoblasts and osteoclasts (40) In the vasculature, autophagy reduces vascular calcification by limiting the release of pro-calcific matrix vesicles (41). LC3 is a protein that functions at the initiation of autophagosome formation and serves as an autophagy marker. While there were no differences between the LC3 signal in $Mgp^{+/+}$ and $Mgp^{-/-}$ at baseline, we observed that rapamycin treatment significantly lowered LC3 puncta in $Mgp^{-/-}$ compared to $Mgp^{+/+}$ mice (Figure 4A).

Quantifying LC3 puncta hints at diminished autophagy but immunofluorescent staining of fixed tissues cannot assess autophagy flux. LC3-I is cleaved to LC3-II when it is incorporated into autophagosomes, but this process is dynamic and LC3-II can also be degraded during autophagy. The comparison of LC3-II to LC3-I in the presence of an autophagy inhibitor like bafilomycin enables autophagy flux observation (42). We isolated SMCs from $Mgp^{+/+}$ and $Mgp^{-/-}$ mice to evaluate autophagy flux. SMC were treated with vehicle (DMSO) or 200 nM rapamycin for 14h and then with vehicle or 100 uM or the late-phase autophagy inhibitor bafilomycin for 24h. Rapamycin alone did not increase LC3-II levels compared to vehicle controls in either the $Mgp^{+/+}$ and $Mgp^{-/-}$ SMC. Bafilomycin, an autophagy inhibitor, increased LC3-II levels in both $Mgp^{+/+}$ and $Mgp^{-/-}$ SMCs compared to their cognate vehicle controls, however the addition of rapamycin did not further alter LC3-II levels compared to bafilomycin (Figure 5B, left graph). Comparing LC3-II to LC3-I showed identical trends (Figure 5B, right graph), suggesting that global LC3 levels and flux are similar in $Mgp^{+/+}$ and $Mgp^{-/-}$ SMCs.

356
Rapamycin is not protecting the mineralized vasculature in *Mgp-/-* mice.
Figure 2 shows that rapamycin treatment doubles the lifespan of $Mgp^{-/-}$ mice compared to
vehicle alone. However, our subsequent inquiries revealed that rapamycin did not decrease the
volume of calcification, did not reverse the adverse remodeling of the medial layer, did not
enhance SMC contractile phenotype, or did not induce more autophagy. The mammalian target
of rapamycin, mTOR, can form two different protein complexes known as mTORC1, bound to
Raptor, and mTORC2, bound to Rictor. The former is acutely sensitive to rapamycin
administration, while the latter requires chronic rapamycin dosing for inhibition (19). To
investigate whether rapamycin's effect on the $Mgp^{-/-}$ mouse are acting on the vasculature, we
generated $Mgp^{-/-}$ mice with SMC-specific deletion of the mTOR complex proteins Raptor and
Rictor (26-28). $Mgp^{-/-};Myh11Cre-GFP;Raptor^{f/f}$ mice ($Mgp^{-/-};Raptor^{SMC-/-}$) were bred to examine
the effects of a nonfunctional mTORC1 complex, while $Mgp^{-/-};Myh11Cre-GFP;Rictor^{f/f}$ ($Mgp^{-/-}$
$;Rictor^{SMC-/-}$) mice knockout was bred to examine a nonfunctional mTORC2 complex. Fully
differentiated SMCs expressing *Myh11* will constitutively express Cre recombinase to inactive
Raptor and *Rictor* gene expression specifically in SMCs.

372
Similar to $Mgp^{-/-}$ mice treated with vehicle in Figure 2, Figure 6 shows that only 50% of the $Mgp^{-/-}$
$;Rictor^{SMC-/-}$ and $Mgp^{-/-};Rictor^{SMC+/+}$ are alive at 40 days, indicating that rapamycin acting via
mTORC2 does not recapitulate the effects of rapamycin on increasing $Mgp^{-/-}$ mice lifespan. The
50% survival rate of $Mgp^{-/-};Raptor^{SMC+/+}$ mice was similar to $Mgp^{-/-}$ treated with vehicle, $Mgp^{-/-}$
$;Rictor^{SMC-/-}$ and $Mgp^{-/-};Rictor^{SMC+/+}$, however the 50% survival rate of $MGP^{-/-};Raptor^{SMC-/-}$ mice
dropped to 22 days. As the Cre strain being used is a non-inducible system, this data in the
$MGP^{-/-};Raptor^{SMC-/-}$ mice suggest that mTORC1 is critical to the growth and development of the
vasculature. Together, these in vivo data suggest that the life-extending effects of rapamycin in
the $Mgp^{-/-}$ mouse are not acting to preserve or reverse the calcification or deleterious vascular
remodeling but are acting on another organ system in the animal.

383
DISCUSSION
In this study, we demonstrated that rapamycin treatment increased the lifespan in $Mgp^{-/-}$ mice,
but the effects of rapamycin do not reduce MAC nor rescue vessel integrity. In mice, rapamycin
has been shown to increase maximum lifespan and delay the onset of cancer as hyperactivity of
the mTOR pathway, frequently found in cancers, leads to the expression and activation of many
oncogenes like PI3K, Akt, and eIF4E. Hence, rapamycin can slow the proliferation of tumor cells
by causing cell arrest, promoting apoptosis, and blocking angiogenesis in tumor growths (43).
However, our data show that the increase in lifespan observed in $Mgp^{-/-}$ mice following
rapamycin treatment is not associated with improvements in SMC contractile protein expression
or enhanced autophagy flux.

394
While rapamycin is known to inhibit SMC proliferation and is used as such in drug-eluting stents,
there is limited research on the effects of rapamycin on vascular morphology in the presence of
calcification (36, 44). It was previously shown that treatment with rapamycin limits the
progression of abdominal and thoracic aortic aneurysms in mice and preserves elastic lamina
integrity (45, 46). In a hyperphosphatemic rat model, rapamycin treatment inhibited aortic
calcium deposition when ectopic calcification was induced by feeding the rats with a high-
phosphate, high-adenine diet. Rapamycin reduced the osteogenic markers MSX2 and OSX in
this model while increasing the *Opn* and *Acta2* gene expression via Klotho upregulation (47).
Our approached used a genetic model of MAC to investigate whether rapamycin could inhibit
calcification in vivo. MGP-deficient mice develop normally and at the time of birth are without
any overt abnormalities. At approximately one week old $Mgp^{-/-}$ mice start exhibiting calcification
of the vessels, suggesting that other gene products might compensate to prevent calcification

during development and immediately post-birth (16). These observations highlight the differences between genetically induced MAC and high phosphate feeding models. Diet-induced calcification must overcome the effect of endogenous MGP, which is a potent inhibitor of calcification, while the genetic removal of MGP results in calcification from the circulating calcium and phosphate and activation of BMP signaling (12, 14).

Rapamycin has also been shown to preserve the differentiated phenotype of vascular SMCs through IRS-1/P3K/AKT2 pathway (37). Our study did not observe rapamycin enhancing the SMC contractile phenotype in $Mgp^{-/-}$ mice. Sma expression typically occurs earlier than $Myh11$ expression during SMC development, even in undifferentiated mesenchymal cells, and its expression persists throughout the differentiation process (48). In contrast, $Myh11$ expression occurs later, during the maturation of SMCs. The first study related to SMC contractile gene expression in response to rapamycin was performed in vitro (37), while our study looked at SMC contractile markers in vivo. Further, the presence of calcification activates mechanosensing pathways that once activated may prevent full re-expression SMC contractile markers (49). However, further investigation would be needed to elucidate how these mechanisms are operating in the context of the $Mgp^{-/-}$ mouse.

MAC is characterized by mineral deposits in the medial vessel layer and frequently occurs in patients with PAD. Vascular calcification is an active biological process with many driving forces, including osteogenic dedifferentiation of SMCs, inflammatory signals, hyperphosphatemia, and extracellular matrix remodeling. In this study, rapamycin prolonged the lifespan of mice who developed serious MAC shortly after birth. Interestingly, even though rapamycin stimulates the cellular process of autophagy (50),-which can recycle components of the ECM, including mineral deposits (51)- calcification volume was not reduced in rapamycin treated $Mgp^{-/-}$ mice. However, microCT data showed that the mineral density was significantly reduced in $Mgp^{-/-}$ mice treated with rapamycin. This data perhaps suggests that changes in mineral structure, if explored at a further timepoint, may impact calcification volume. Alternatively, as MGP also functions to inhibit BMP2 signaling, rapamycin may not interfere with that process- such that changes in dosage or duration of rapamycin treatment would still not alter calcification content in the vessel wall. Another explanation is that the activated autophagy pathways cannot overcome the rapid deposition of hydroxyapatite that occurs in $Mgp^{-/-}$ mice. Frauscher et al. demonstrated that induction of autophagy by rapamycin treatment decreased calcification in the vasculature of uremic DBA/2 mice (52). This argument led us to examine autophagy flux in SMCs. However, we did not observe a difference in autophagy flux between $Mgp^{+/+}$ and $Mgp^{-/-}$. It is important to note that the $Mgp^{-/-}$ is on a C57BL6J background, perhaps explaining the differences observed in these in vivo models.

The mammalian target of rapamycin (mTOR) forms two distinct protein complexes known as mTORC1 and mTORC2. mTORC1, activated by amino acids and allosterically inhibited by rapamycin, requires a complex specific protein named raptor (53). After chronic administration, mTORC2 is only inhibited by rapamycin and requires a Rictor (54). In the $Mgp^{-/-};Raptor^{SMC-/-}$ mice, induction of Cre to knockout $Raptor$ was more lethal than the $Mgp^{-/-};Raptor^{SMC+/+}$ or Mgp-/- mice. Therefore, $Raptor$ is essential for the growth and maintenance of SMCs in the vasculature. This essential role in growth and development is perhaps unsurprising, considering that global $Raptor$ knockout mice die in utero (55). Moreover, there was no benefit to lifespan in the the $Mgp^{-/-}$:Rictor$^{SMC-/-}$ mice. The survival curves for the the $Mgp^{-/-}$:Rictor$^{SMC+/+}$ and $Mgp^{-/-};Rictor^{SMC-/-}$ mice, targeting mTORC2 signaling, do not diverge at any point and are statistically insignificant, and also mirror that seen in the $Mgp^{-/-}$ strain. Considering these data together, rapamycin likely extends the lifespan of $Mgp^{-/-}$ mice by targeting another organ system.

Arterial stiffness is strongly linked to heart failure (56), and these non-atherosclerotic vascular
pathologies are much more common in some populations, such as African-ancestry individuals
who are known to have a higher risk of cardiovascular disease events and mortality compared
to Caucasians(57-61). This excess cardiovascular disease risk is largely due to hypertensive
and peripheral vascular disease, versus atherosclerotic disease (57-59, 62-64). As such, it is
critical to define drivers of MAC such as MGP, which in some clinical studies in humans has
been associated with heart failure and MAC (65). Impairment in MGP activation, which would
phenocopy genetic deletion, is linked to a higher risk of heart failure and peripheral artery
disease due to excessive pulsatile afterloads on the left ventricle that reduce coronary artery
perfusion pressure during diastole (66, 67). As the mice that lack MGP are prone to aortic
rupture or heart failure, we speculate that rapamycin's effect on the mice's lifespan may be due
to rapamycin acting on the cardiac tissues.
470
Our study suggests that rapamycin's life-extending benefit in $Mgp^{-/-}$ mice may act through
another organ system, possibly the heart rather than the vessels. Future studies are needed to
examine the impact of rapamycin on cardiac function in $Mgp^{-/-}$ mice. Rapamycin has been
shown to increase cardiomyocyte autophagy (68), limit cardiomyocyte death, and attenuate
cardiomyocyte hypertrophy and cardiac remodeling by enhancing mTORC2 signaling while
simultaneously inhibiting mTORC1 signaling (69). Inhibition of mTOR signaling with rapamycin
improved cardiac function, such as left ventricular end-systolic dimensions, fractional
shortening, and ejection fraction in mice with decompensated cardiac hypertrophy compared to
the control group (70). Rapamycin has a protective effect on cardiac muscle contractility.
Cardiomyocytes isolated from the oldest mice treated with rapamycin show better contraction
(similar to that of young mice) than cardiomyocytes from old mice not treated with rapamycin
(71).
483
CONCLUSION
Rapamycin injection extended the lifespan of $Mgp^{-/-}$ mice that exhibit an advanced form of MAC
compared to $Mgp^{+/+}$ mice treated with vehicle, indicating a potential therapeutic benefit.
However, the mechanism of rapamycin-induced lifespan in relation to MAC must be clarified.
Rapamycin reduced mineral density but did not decrease calcification volume or show any
improvement in maintaining the SMC contractile phenotype in $Mgp^{-/-}$ mice. Rapamycin did not
alter proliferation in either $Mgp^{+/+}$ or $Mgp^{-/-}$ mice or induce autophagy. The rapamycin effect on
lifespan extension and mineral density reduction is independent of its inhibition of mTORC1 or
mTORC2 complexes in SMC, suggesting that it is not vascular dysfunction itself leading to
death. Thus, we predict that rapamycin is acting on the heart tissue to withstand failure due to
the stiffened, calcified vessels.
495

496 **ACKNOWLEDGEMENTS**
497 We want to thank the Center for Biologic Imaging at the University of Pittsburgh for their help
498 and support with microscopy imaging. Some figures were created with Biorender.com.
499
500 **SOURCES OF FUNDING**
501 This publication was supported by the NIH K22 HL117917-01A1, The Winters Foundation, The
502 McKamish Family Foundation (St. Hilaire), and the American Heart Association
503 24POST1186619 (Behzadi). MicorCT analysis and processing was performed on a system
504 supported by NIH S10-OD021533 (Verdelis).
505

506 **DISCLOSURES**
507 None.
508

REFERENCES

1. **St. Hilaire C**. Medial Arterial Calcification: A Significant and Independent Contributor of Peripheral Artery Disease. *Arteriosclerosis, Thrombosis, and Vascular Biology* 42: 253-260, 2022.

2. **O'Neill WC, Han KH, Schneider TM, and Hennigar RA**. Prevalence of nonatheromatous lesions in peripheral arterial disease. *Arterioscler Thromb Vasc Biol* 35: 439-447, 2015.

3. **Narula N, Dannenberg AJ, Olin JW, Bhatt DL, Johnson KW, Nadkarni G, Min J, Torii S, Poojary P, Anand SS, Bax JJ, Yusuf S, Virmani R, and Narula J**. Pathology of Peripheral Artery Disease in Patients With Critical Limb Ischemia. *J Am Coll Cardiol* 72: 2152-2163, 2018.

4. **Markello TC, Pak LK, St Hilaire C, Dorward H, Ziegler SG, Chen MY, Chaganti K, Nussbaum RL, Boehm M, and Gahl WA**. Vascular pathology of medial arterial calcifications in NT5E deficiency: implications for the role of adenosine in pseudoxanthoma elasticum. *Mol Genet Metab* 103: 44-50, 2011.

5. **Lanzer P, Hannan FM, Lanzer JD, Janzen J, Raggi P, Furniss D, Schuchardt M, Thakker R, Fok PW, Saez-Rodriguez J, Millan A, Sato Y, Ferraresi R, Virmani R, and St Hilaire C**. Medial Arterial Calcification: JACC State-of-the-Art Review. *J Am Coll Cardiol* 78: 1145-1165, 2021.

6. **Hortells L, Sur S, and St Hilaire C**. Cell Phenotype Transitions in Cardiovascular Calcification. *Front Cardiovasc Med* 5: 27, 2018.

7. **Villa-Bellosta R**. Role of the extracellular ATP/pyrophosphate metabolism cycle in vascular calcification. *Purinergic Signal* 19: 345-352, 2023.

8. **Rutsch F, Ruf N, Vaingankar S, Toliat MR, Suk A, Hohne W, Schauer G, Lehmann M, Roscioli T, Schnabel D, Epplen JT, Knisely A, Superti-Furga A, McGill J, Filippone M, Sinaiko AR, Vallance H, Hinrichs B, Smith W, Ferre M, Terkeltaub R, and Nurnberg P**. Mutations in ENPP1 are associated with 'idiopathic' infantile arterial calcification. *Nat Genet* 34: 379-381, 2003.

9. **St Hilaire C, Ziegler SG, Markello TC, Brusco A, Groden C, Gill F, Carlson-Donohoe H, Lederman RJ, Chen MY, Yang D, Siegenthaler MP, Arduino C, Mancini C, Freudenthal B, Stanescu HC, Zdebik AA, Chaganti RK, Nussbaum RL, Kleta R, Gahl WA, and Boehm M**. NT5E mutations and arterial calcifications. *N Engl J Med* 364: 432-442, 2011.

10. **Baral A, F FH, Lee SC, Matthew BP, Ferrante EA, Brofferio A, Cudrici CD, Rollison SF, Chen MY, Boehm M, and Wen H**. Images in Vascular Medicine: High-resolution CT imaging of arterial calcification in the hands and legs of patients with CD73 deficiency. *Vasc Med* 29: 342-344, 2024.

11. **Jin H, St Hilaire C, Huang Y, Yang D, Dmitrieva NI, Negro A, Schwartzbeck R, Liu Y, Yu Z, Walts A, Davaine JM, Lee DY, Donahue D, Hsu KS, Chen J, Cheng T, Gahl W, Chen G, and Boehm M**. Increased activity of TNAP compensates for reduced adenosine production and promotes ectopic calcification in the genetic disease ACDC. *Sci Signal* 9: ra121, 2016.

12. **Murshed M, Schinke T, McKee MD, and Karsenty G**. Extracellular matrix mineralization is regulated locally; different roles of two gla-containing proteins. *J Cell Biol* 165: 625-630, 2004.

13. **Hackeng TM, Rosing J, Spronk HM, and Vermeer C**. Total chemical synthesis of human matrix Gla protein. *Protein Sci* 10: 864-870, 2001.

14. **Zebboudj AF, Imura M, and Bostrom K**. Matrix GLA protein, a regulatory protein for bone morphogenetic protein-2. *J Biol Chem* 277: 4388-4394, 2002.

15. **Malhotra R, Burke MF, Martyn T, Shakartzi HR, Thayer TE, O'Rourke C, Li P, Derwall M, Spagnolli E, Kolodziej SA, Hoeft K, Mayeur C, Jiramongkolchai P, Kumar R, Buys ES, Yu PB, Bloch KD, and Bloch DB**. Inhibition of bone morphogenetic protein signal

transduction prevents the medial vascular calcification associated with matrix Gla protein deficiency. *PLoS One* 10: e0117098, 2015.

16. **Luo G, Ducy P, McKee MD, Pinero GJ, Loyer E, Behringer RR, and Karsenty G**. Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature* 386: 78-81, 1997.

17. **Haissaguerre M, Saucisse N, and Cota D**. Influence of mTOR in energy and metabolic homeostasis. *Mol Cell Endocrinol* 397: 67-77, 2014.

18. **Saxton RA, and Sabatini DM**. mTOR Signaling in Growth, Metabolism, and Disease. *Cell* 168: 960-976, 2017.

19. **Panwar V, Singh A, Bhatt M, Tonk RK, Azizov S, Raza AS, Sengupta S, Kumar D, and Garg M**. Multifaceted role of mTOR (mammalian target of rapamycin) signaling pathway in human health and disease. *Signal Transduct Target Ther* 8: 375, 2023.

20. **Arriola Apelo SI, and Lamming DW**. Rapamycin: An InhibiTOR of Aging Emerges From the Soil of Easter Island. *J Gerontol A Biol Sci Med Sci* 71: 841-849, 2016.

21. **Zhan JK, Wang YJ, Wang Y, Wang S, Tan P, Huang W, and Liu YS**. The mammalian target of rapamycin signalling pathway is involved in osteoblastic differentiation of vascular smooth muscle cells. *Can J Cardiol* 30: 568-575, 2014.

22. **He HQ, Law BYK, Zhang N, Qiu CL, Qu YQ, Wu AG, Han Y, Song Q, Zheng WL, Liu Y, He YZ, and Wong VKW**. Bavachin Protects Human Aortic Smooth Muscle Cells Against beta-Glycerophosphate-Mediated Vascular Calcification and Apoptosis via Activation of mTOR-Dependent Autophagy and Suppression of beta-Catenin Signaling. *Front Pharmacol* 10: 1427, 2019.

23. **Dai XY, Zhao MM, Cai Y, Guan QC, Zhao Y, Guan Y, Kong W, Zhu WG, Xu MJ, and Wang X**. Phosphate-induced autophagy counteracts vascular calcification by reducing matrix vesicle release. *Kidney Int* 83: 1042-1051, 2013.

24. **Joolharzadeh P, and St Hilaire C**. CD73 (Cluster of Differentiation 73) and the Differences Between Mice and Humans. *Arterioscler Thromb Vasc Biol* 39: 339-348, 2019.

25. **Ziegler SG, Ferreira CR, MacFarlane EG, Riddle RC, Tomlinson RE, Chew EY, Martin L, Ma CT, Sergienko E, Pinkerton AB, Millan JL, Gahl WA, and Dietz HC**. Ectopic calcification in pseudoxanthoma elasticum responds to inhibition of tissue-nonspecific alkaline phosphatase. *Sci Transl Med* 9: 2017.

26. **Xin HB, Deng KY, Rishniw M, Ji G, and Kotlikoff MI**. Smooth muscle expression of Cre recombinase and eGFP in transgenic mice. *Physiol Genomics* 10: 211-215, 2002.

27. **Sengupta S, Peterson TR, Laplante M, Oh S, and Sabatini DM**. mTORC1 controls fasting-induced ketogenesis and its modulation by ageing. *Nature* 468: 1100-1104, 2010.

28. **Magee JA, Ikenoue T, Nakada D, Lee JY, Guan KL, and Morrison SJ**. Temporal changes in PTEN and mTORC2 regulation of hematopoietic stem cell self-renewal and leukemia suppression. *Cell Stem Cell* 11: 415-428, 2012.

29. **Liu M, Espinosa-Diez C, Mahan S, Du M, Nguyen AT, Hahn S, Chakraborty R, Straub AC, Martin KA, Owens GK, and Gomez D**. H3K4 di-methylation governs smooth muscle lineage identity and promotes vascular homeostasis by restraining plasticity. *Dev Cell* 56: 2765-2782 e2710, 2021.

30. **Wanjare M, Kuo F, and Gerecht S**. Derivation and maturation of synthetic and contractile vascular smooth muscle cells from human pluripotent stem cells. *Cardiovasc Res* 97: 321-330, 2013.

31. **Zhang L, Yang C, Li J, Zhu Y, and Zhang X**. High extracellular magnesium inhibits mineralized matrix deposition and modulates intracellular calcium signaling in human bone marrow-derived mesenchymal stem cells. *Biochem Biophys Res Commun* 450: 1390-1395, 2014.

32. **Schneider CA, Rasband WS, and Eliceiri KW**. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9: 671-675, 2012.

33. **Li Q, Price TP, Sundberg JP, and Uitto J**. Juxta-articular joint-capsule mineralization in
CD73 deficient mice: similarities to patients with NT5E mutations. *Cell Cycle* 13: 2609-2615,
2014.
34. **Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, Nadon NL,**
Wilkinson JE, Frenkel K, Carter CS, Pahor M, Javors MA, Fernandez E, and Miller RA.
Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature* 460: 392-
395, 2009.
35. **Martin KA, Rzucidlo EM, Merenick BL, Fingar DC, Brown DJ, Wagner RJ, and**
Powell RJ. The mTOR/p70 S6K1 pathway regulates vascular smooth muscle cell
differentiation. *Am J Physiol Cell Physiol* 286: C507-517, 2004.
36. **Marx SO, Jayaraman T, Go LO, and Marks AR**. Rapamycin-FKBP inhibits cell cycle
regulators of proliferation in vascular smooth muscle cells. *Circ Res* 76: 412-417, 1995.
37. **Martin KA, Merenick BL, Ding M, Fetalvero KM, Rzucidlo EM, Kozul CD, Brown DJ,**
Chiu HY, Shyu M, Drapeau BL, Wagner RJ, and Powell RJ. Rapamycin promotes vascular
smooth muscle cell differentiation through insulin receptor substrate-1/phosphatidylinositol 3-
kinase/Akt2 feedback signaling. *J Biol Chem* 282: 36112-36120, 2007.
38. **Cheung C, Bernardo AS, Trotter MW, Pedersen RA, and Sinha S**. Generation of
human vascular smooth muscle subtypes provides insight into embryological origin-dependent
disease susceptibility. *Nat Biotechnol* 30: 165-173, 2012.
39. **Skalli O, Pelte MF, Peclet MC, Gabbiani G, Gugliotta P, Bussolati G, Ravazzola M,**
and Orci L. Alpha-smooth muscle actin, a differentiation marker of smooth muscle cells, is
present in microfilamentous bundles of pericytes. *J Histochem Cytochem* 37: 315-321, 1989.
40. **Qi M, Zhang L, Ma Y, Shuai Y, Li L, Luo K, Liu W, and Jin Y**. Autophagy Maintains
the Function of Bone Marrow Mesenchymal Stem Cells to Prevent Estrogen Deficiency-Induced
Osteoporosis. *Theranostics* 7: 4498-4516, 2017.
41. **Sun X, Zheng Y, Xie L, Zhou Y, Liu R, Ma Y, Zhao M, and Liu Y**. Autophagy reduces
aortic calcification in diabetic mice by reducing matrix vesicle body-mediated IL-1beta release.
Exp Cell Res 432: 113803, 2023.
42. **Mizushima N, and Yoshimori T**. How to Interpret LC3 Immunoblotting. *Autophagy* 3:
542-545, 2014.
43. **Ganesh SK, and Subathra Devi C**. Molecular and therapeutic insights of rapamycin: a
multi-faceted drug from Streptomyces hygroscopicus. *Mol Biol Rep* 50: 3815-3833, 2023.
44. **Sabate M**. Sirolimus Versus Paclitaxel: Second Round. *JACC Cardiovasc Interv* 15:
780-782, 2022.
45. **Rouer M, Xu BH, Xuan HJ, Tanaka H, Fujimura N, Glover KJ, Furusho Y, Gerritsen**
M, and Dalman RL. Rapamycin limits the growth of established experimental abdominal aortic
aneurysms. *Eur J Vasc Endovasc Surg* 47: 493-500, 2014.
46. **Zhou B, Li W, Zhao G, Yu B, Ma B, Liu Z, Xie N, Fu Y, Gong Z, Dai R, Zhang X, and**
Kong W. Rapamycin prevents thoracic aortic aneurysm and dissection in mice. *J Vasc Surg* 69:
921-932 e923, 2019.
47. **Zhao Y, Zhao MM, Cai Y, Zheng MF, Sun WL, Zhang SY, Kong W, Gu J, Wang X,**
and Xu MJ. Mammalian target of rapamycin signaling inhibition ameliorates vascular
calcification via Klotho upregulation. *Kidney Int* 88: 711-721, 2015.
48. **Li L, Miano JM, Cserjesi P, and Olson EN**. SM22α, a Marker of Adult Smooth Muscle,
Is Expressed in Multiple Myogenic Lineages During Embryogenesis. *Circulation Research* 78:
188-195, 1996.
49. **Krohn JB, Hutcheson JD, Martínez-Martínez E, Irvin WS, Bouten CVC, Bertazzo S,**
Bendeck MP, and Aikawa E. Discoidin Domain Receptor-1 Regulates Calcific Extracellular
Vesicle Release in Vascular Smooth Muscle Cell Fibrocalcific Response via Transforming
Growth Factor-β Signaling. *Arteriosclerosis Thrombosis and Vascular Biology* 36: 525-533,
2016.

50.   **Tanemura M, Ohmura Y, Deguchi T, Machida T, Tsukamoto R, Wada H, Kobayashi
S, Marubashi S, Eguchi H, Ito T, Nagano H, Mori M, and Doki Y**. Rapamycin causes
upregulation of autophagy and impairs islets function both in vitro and in vivo. *Am J Transplant*
12: 102-114, 2012.
51.   **Duan ZX, Tu C, Liu Q, Li SQ, Li YH, Xie P, and Li ZH**. Adiponectin receptor agonist
AdipoRon attenuates calcification of osteoarthritis chondrocytes by promoting autophagy. *J Cell*
Biochem 121: 3333-3344, 2020.
52.   **Frauscher B, Kirsch AH, Schabhuttl C, Schweighofer K, Ketszeri M, Pollheimer M,
Dragun D, Schroder K, Rosenkranz AR, Eller K, and Eller P**. Autophagy Protects From
Uremic Vascular Media Calcification. *Front Immunol* 9: 1866, 2018.
53.   **Szwed A, Kim E, and Jacinto E**. Regulation and metabolic functions of mTORC1 and
mTORC2. *Physiol Rev* 101: 1371-1426, 2021.
54.   **Sarbassov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, Markhard AL,
and Sabatini DM**. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB.
Mol Cell 22: 159-168, 2006.
55.   **Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, Brown M,
Fitzgerald KJ, and Sabatini DM**. Ablation in mice of the mTORC components raptor, rictor, or
mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKCalpha, but not
S6K1. *Dev Cell* 11: 859-871, 2006.
56.   **Schott A, Kluttig A, Mikolajczyk R, Greiser KH, Werdan K, Sedding D, and Nuding
S**. Association of arterial stiffness and heart failure with preserved ejection fraction in the elderly
population - results from the CARLA study. *J Hum Hypertens* 37: 463-471, 2023.
57.   **Writing Group M, Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De
Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N,
Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A,
McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W,
Sacco R, Sorlie P, Roger VL, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J,
American Heart Association Statistics C, and Stroke Statistics S**. Heart disease and stroke
statistics--2010 update: a report from the American Heart Association. *Circulation* 121: e46-
e215, 2010.
58.   **Allison MA, Ho E, Denenberg JO, Langer RD, Newman AB, Fabsitz RR, and Criqui
MH**. Ethnic-specific prevalence of peripheral arterial disease in the United States. *Am J Prev*
Med 32: 328-333, 2007.
59.   **Allison MA, Budoff MJ, Nasir K, Wong ND, Detrano R, Kronmal R, Takasu J, and
Criqui MH**. Ethnic-specific risks for atherosclerotic calcification of the thoracic and abdominal
aorta (from the Multi-Ethnic Study of Atherosclerosis). *Am J Cardiol* 104: 812-817, 2009.
60.   **Chaturvedi N, Bulpitt CJ, Leggetter S, Schiff R, Nihoyannopoulos P, Strain WD,
Shore AC, and Rajkumar C**. Ethnic differences in vascular stiffness and relations to
hypertensive target organ damage. *J Hypertens* 22: 1731-1737, 2004.
61.   **D'Agostino RB, Jr., Burke G, O'Leary D, Rewers M, Selby J, Savage PJ, Saad MF,
Bergman RN, Howard G, Wagenknecht L, and Haffner SM**. Ethnic differences in carotid wall
thickness. The Insulin Resistance Atherosclerosis Study. *Stroke* 27: 1744-1749, 1996.
62.   **Ali H, Zmuda JM, Cvejkus RK, Kershaw EE, Kuipers AL, Oczypok EA, Wheeler V,
Bunker CH, and Miljkovic I**. Wnt Pathway Inhibitor DKK1: A Potential Novel Biomarker for
Adiposity. *J Endocr Soc* 3: 488-495, 2019.
63.   **Bild DE, Detrano R, Peterson D, Guerci A, Liu K, Shahar E, Ouyang P, Jackson S,
and Saad MF**. Ethnic differences in coronary calcification: the Multi-Ethnic Study of
Atherosclerosis (MESA). *Circulation* 111: 1313-1320, 2005.
64.   **Budoff MJ, Nasir K, Mao S, Tseng PH, Chau A, Liu ST, Flores F, and Blumenthal
RS**. Ethnic differences of the presence and severity of coronary atherosclerosis. *Atherosclerosis*
187: 343-350, 2006.

65.   **Malhotra R, Nicholson CJ, Wang D, Bhambhani V, Paniagua S, Slocum C,**
Sigurslid HH, Lino Cardenas CL, Li R, Boerboom SL, Chen YC, Hwang SJ, Yao C,
Ichinose F, Bloch DB, Lindsay ME, Lewis GD, Aragam JR, Hoffmann U, Mitchell GF,
Hamburg NM, Vasan RS, Benjamin EJ, Larson MG, Zapol WM, Cheng S, Roh JD,
O'Donnell CJ, Nguyen C, Levy D, and Ho JE. Matrix Gla Protein Levels Are Associated With
Arterial Stiffness and Incident Heart Failure With Preserved Ejection Fraction. *Arterioscler*
Thromb Vasc Biol 42: e61-e73, 2022.
66.   **Dalmeijer GW, van der Schouw YT, Magdeleyns EJ, Vermeer C, Verschuren WM,**
Boer JM, and Beulens JW. Matrix Gla protein species and risk of cardiovascular events in type
2 diabetic patients. *Diabetes Care* 36: 3766-3771, 2013.
67.   **Chirinos JA**. Matrix Gla Protein, Large Artery Stiffness, and the Risk of Heart Failure
With Preserved Ejection Fraction. *Arterioscler Thromb Vasc Biol* 42: 223-226, 2022.
68.   **Buss SJ, Muenz S, Riffel JH, Malekar P, Hagenmueller M, Weiss CS, Bea F,**
Bekeredjian R, Schinke-Braun M, Izumo S, Katus HA, and Hardt SE. Beneficial effects of
Mammalian target of rapamycin inhibition on left ventricular remodeling after myocardial
infarction. *J Am Coll Cardiol* 54: 2435-2446, 2009.
69.   **Volkers M, Konstandin MH, Doroudgar S, Toko H, Quijada P, Din S, Joyo A,**
Ornelas L, Samse K, Thuerauf DJ, Gude N, Glembotski CC, and Sussman MA. Mechanistic
target of rapamycin complex 2 protects the heart from ischemic damage. *Circulation* 128: 2132-
2144, 2013.
70.   **McMullen JR, Sherwood MC, Tarnavski O, Zhang L, Dorfman AL, Shioi T, and**
Izumo S. Inhibition of mTOR signaling with rapamycin regresses established cardiac
hypertrophy induced by pressure overload. *Circulation* 109: 3050-3055, 2004.
71.   **Chakraborty AD, Kooiker K, Kobak KA, Cheng Y, Lee CF, Razumova M, Granzier**
HH, Regnier M, Rabinovitch PS, Moussavi-Harami F, and Chiao YA. Late-life Rapamycin
Treatment Enhances Cardiomyocyte Relaxation Kinetics and Reduces Myocardial Stiffness.
bioRxiv 2023.

740
741
742   **FIGURE LEGENDS**
743
Figure 1: Histological comparison of Matrix Gla Protein (*Mgp*) mice arteries, medial and
intimal calcification in human arteries. A. Histological sections of *Mgp*$^{-/-}$ mouse aorta, human
tibial plaque and human coronary artery with atherosclerotic plaque, stained for calcification
using Von Kossa Staining. *Mgp*$^{-/-}$ mouse aorta develop calcification (Calc) in the media layer of
artery similar to human tibial artery. **B.** Dot Blot Assay to identify MGP protein in Wild Type
(*Mgp*$^{+/+}$), Heterozygous (*Mgp*$^{+/-}$), and Knockout MGP (*Mgp*$^{-/-}$) mice. *Mgp*$^{-/-}$ showed no expression
of MGP (Scale Bars: human tissue 1 mm, mouse tissue 0.5 mm).
751
Figure 2: Impact of rapamycin treatment on survival in Matrix Gla Protein (*Mgp*)-deficient
mice A. Kaplan-Meier analysis of overall survival in *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice treated with vehicle
(Dimethyl sulfoxide; DMSO) or rapamycin (5mg/kg/week) for 3 months. Rapamycin treatment
significantly extends the survival of *Mgp*$^{-/-}$ mice (n=7) compared to those treated with the vehicle
(n=19). *Mgp*$^{+/+}$ mice treated with vehicle (n=5) or either rapamycin (n=8) showed the same
survival.
758
759   **Figure 3: Effect of rapamycin treatment on medial calcification in Matrix Gla Protein**
760   **(*Mgp*)-deficient mice a.** A cross-section of the aorta from *Mgp*$^{-/-}$ mice treated with vehicle (Veh)
761   or rapamycin (RAPA, [5 mg/kg] once or three times a week). Mineral density decreased with
762   three times a week rapamycin. **B.** Histological sections of *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mouse aortas

treated with vehicle or rapamycin three times a week, stained for calcification using Von Kossa staining. **C.** A cross-section of the aorta treated with vehicle or rapamycin three times a week were stained for calcification marker runt-related transcription factor 2 (RUNX2) by immunofluorescence. In $Mgp^{-/-}$ mice, RUNX2 expression was elevated with vehicle treatment but no significant difference with rapamycin treatment.

Figure 4: Histological structure of $Mgp^{+/+}$ and $Mgp^{-/-}$ mice aorta treated with vehicle or rapamycin. A. The cross sections of mice aorta were stained with Masson's Trichrome to identify collagen (stained blue). The staining revealed disruption in collagen fiber structure in $Mgp^{-/-}$ mice treated either with vehicle or rapamycin **B.** The cross sections of mice aorta were stained with Verhoeff-Van Gieson (VVG) to identify elastic fibers (stained black). The staining revealed disruption of elastic fiber structure and a reduction of elastic fibers in the $Mgp^{-/-}$ vehicle (Veh) group, and no difference observed with rapamycin (RAPA) treatment (Scale bar: 0.2 mm). **C.** The cross sections of mice aorta were stained with smooth muscle actin (SMA) and myosin heavy chain gene (MYH11) SMA. The rapamycin increased the SMA levels in $Mgp^{-/-}$ mice compared to the $Mgp^{-/-}$ vehicle. **D.** The cross section of mice aorta was stained for proliferation markers (ki67), which was higher in $Mgp^{-/-}$ compared to the $Mgp^{+/+}$.

Figure 5: The effect of rapamycin treatment on autophagy flux. A. Histological sections of $Mgp^{+/+}$ and $Mgp^{-/-}$ mouse aortas treated with vehicle (DMSO, Dimethyl sulfoxide) or rapamycin [5mg/kg] three times a week, stained for Microtubule-associated protein 1A/1B-light chain 3 (LC3) by immunofluorescence. LC3 decreased with rapamycin (RAPA) in $Mgp^{-/-}$ mice compared to the $Mgp^{+/+}$ mice. **B.** Smooth muscle cells from $Mgp^{+/+}$ and $Mgp^{-/-}$ mice were treated with vehicle (DMSO) or rapamycin (200 µM) for 14 hours, followed by the addition of bafilomycin (BafA, 100 µM), an autophagy inhibitor, for an additional 24 hours. Levels of LC3 (LC3-I and LC3-II) were quantified by western blot. Rapamycin treatment increases the levels of LC3-II/LC3I and LC3-II protein in both $Mgp^{+/+}$ and $Mgp^{-/-}$ mice smooth muscle cells.

Figure 6: Survival probability of Matrix Gla Protein (Mgp)-deficient mice with Rictor and Raptor knockouts in vascular smooth muscle cells (SMC). A. $Mgp^{-/-}$ using Cre recombinase that target the gene encoding Rictor in SMC ($SMC^{+/+}$ (n=7), $SMC^{-/-}$ (n=5)), a protein involved in mTORC2 signaling pathways. **B.** $Mgp^{-/-}$ using Cre recombinase that target the gene encoding Raptor in SMC ($SMC^{+/+}$ (n=6), $SMC^{-/-}$ (n=7)), a protein involved in mTORC1 signaling pathways. The presence or absence of Rictor, when combined with MGP deficiency, does not significantly affect the lifespan of the mice. However, Raptor plays a crucial role in the survival of MGP-deficient mice.

A

| Mouse
Mgp^{-/-} Aorta | Human
Tibial Artery | Human Coronary Artery |

B

| | *Mgp*^{+/+} | *Mgp*^{+/-} | *Mgp*^{-/-} |

MGP

Tubulin

A

B

A

B