Rapamycin Reduces Mineral Density and Promotes Beneficial Vascular Remodeling in a Murine Model of Severe Medial Arterial Calcification.

Parya Behzadi[1], Andrew A. Wendling[1], Rolando A. Cuevas[1], Alex Crane[1], Claire C. Chu[1], William J Moorhead III[1], Ryan Wong[1], Mark Brown[1], Joshua Tamakloe[1], Swathi Suresh[1], Payam Salehi[2], Iris Z. Jaffe[3], Allison L. Kuipers[4], Lyudmila Lukashova[5], Konstantinos Verdelis[5], and Cynthia St. Hilaire[1,6,7]

[1] Department of Medicine, Division of Cardiology, and the Pittsburgh Heart, Lung, Blood and Vascular Medicine Institute, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.

[2] CardioVascular Center, Vascular Surgery, Tufts Medical Center, 800 Washington Street, Boston, MA, 02111-1800, USA

[3] Molecular Cardiology Research Institute, Tufts Medical Center, 800 Washington Street, Boston, MA, 02111-1800, USA

[4] Departments of Medicine and Epidemiology & Biostatistics, College of Human Medicine, Michigan State University, Grand Rapids, MI, USA

[5] Departments of Endodontics and Oral Biology, School of Dental Medicine, University of Pittsburgh, Pittsburgh, PA, USA

[6] Department of Bioengineering, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

[7] Department of Cardiothoracic Surgery, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

CORRESPONDING AUTHOR CONTACT:
Cynthia St. Hilaire, PhD, FAHA
University of Pittsburgh
BSTWR 1744
200 Lothrop Street
Pittsburgh, PA 15261
sthilair@pitt.edu

ABSTRACT: 249 words

MANUSCRIPT: 6173 words

KEYWORDS: Medial Arterial Calcification, Rapamycin, Matrix GLA protein, Collagen, Elastin

ABBREVIATIONS

Adenosine monophosphate	AMP
Adenosine triphosphate	ATP
Alkaline phosphate	TNAP
Arterial calcification due to deficiency of CD73	ACDC
Ecto-5'-nucleotidase	NT5E
Induced pluripotent stem cells	iPSCs
Matrix GLA protein	Mgp
Medial arterial calcification	MAC
mTORC complex 1	mTORC1

52	mTORC complex 2
53	Peripheral artery disease
54	Smooth muscle α-actin
55	Smooth muscle cells
56	Myosin heavy chain 11
57	Runt-related transcription factor 2

mTORC complex 2   mTORC2
Peripheral artery disease   PAD
Smooth muscle α-actin   α-SMA
Smooth muscle cells   SMCs
Myosin heavy chain 11   Myh11
Runt-related transcription factor 2   Runx2

ABSTRACT
Peripheral artery disease (PAD) is the narrowing of the arteries that carry blood to the lower
extremities. PAD has been traditionally associated with atherosclerosis. However, recent
studies have found that thrombotic events triggered by medial arterial calcification (MAC) is the
primary cause of chronic limb ischemia below the knee. MAC is localized around the elastic
fibers surrounding smooth muscle cells (SMCs) in arteries. Matrix GLA protein (MGP) binds
circulating calcium and prevents hydroxyapatite mineral deposition, while also modulating pro-
osteogenic signaling by attenuating BMP-2-mediated activation of Mgp gene expression. $Mgp^{-/-}$
mice develop severe MAC and die around 8 weeks after birth due to aortic rupture or heart
failure. We previously discovered a rare genetic disease Arterial Calcification due to Deficiency
of CD73 (ACDC), in which patients present with extensive MAC in their lower extremity arteries.
Using a patient-specific induced pluripotent stem cell model, we found that rapamycin inhibited
calcification. Here we investigated whether rapamycin could reduce MAC in vivo using the $Mgp^{-/-}$
murine model. $Mgp^{+/+}$ and $Mgp^{-/-}$ mice received 5mg/kg rapamycin or vehicle. Calcification
content was assessed via microCT, and vascular morphology and extracellular matrix content
were assessed histologically. Immunostaining and western blot analysis were used to examine
SMC phenotype and extracellular matrix content. Rapamycin prolonged $Mgp^{-/-}$ mice lifespan,
decreased mineral density in the arteries, maintained SMC contractile phenotype, and improved
vessel structure, however, calcification volume was unchanged. $Mgp^{-/-}$ mice with SMC-specific
deletion of Raptor or Rictor did not recapitulate treatment with rapamycin. These findings
suggest rapamycin promotes beneficial vascular remodeling in vessels with MAC.

NEWS AND NOTEWORTHY
Peripheral artery disease (PAD) is associated with medial arterial calcification (MAC), which
involves calcification of arterial elastic fibers and smooth muscle cells (SMCs). Matrix GLA
protein (MGP) inhibits vascular calcification, and $Mgp^{-/-}$ mice develop severe MAC. Using this
model, we found rapamycin prolonged lifespan, reduced arterial mineral density, maintained
SMC contractile phenotype, and improved vessel structure, though calcification volume
remained unchanged. Findings highlight rapamycin's potential for vascular remodeling in MAC.

INTRODUCTION
Peripheral artery disease (PAD) is the narrowing of blood vessels in the lower extremities due to
inward remodeling or thrombotic occlusion, both of which cause chronic limb ischemia and often
result in amputation of the lower leg (1). Traditionally atherosclerosis has been the assumed
cause of PAD. However, recent studies now show that medial arterial calcification (MAC)
promotes inward remodeling, and stiffness related to medial dysplasia induces thrombosis (2-4).
MAC is characterized by the progressive buildup of calcium and phosphate along the elastic
fibers within the arterial walls and is typically not associated with lipid deposition or fibrous cap
formation (5). These structural features distinguish PAD from atherosclerosis and indicate that
MAC is a distinct pathology driving adverse outcomes in PAD. Currently, there are no specific
medical therapies that target the distinct pathogenesis of PAD or MAC. Finding therapies that
can prevent, stop, or even reverse MAC in PAD could greatly enhance the current standard of
care.
102

Vascular calcification stems from the nucleation of calcium and phosphate into hydroxyapatite
crystals and the release of pro-mineralizing matrix vesicles from osteogenic SMCs that
accumulate along the elastic lamina (6). Phosphate is a byproduct of the breakdown of
extracellular adenosine triphosphate (ATP) to adenosine, generating inorganic phosphate at
several points (7, 8). There are several genetic diseases that present with MAC, which harbor
inactivating mutations in the genes related to extracellular ATP metabolism (5). Patients with
Arterial Calcification due to Deficiency of CD73 (ACDC, OMIM # 211800) harbor inactivating
mutations in the *NT5E* gene, which encodes for the CD73 enzyme that primarily metabolizes
extracellular AMP to adenosine and inorganic phosphate. Key signatures of this disease are
calcification nodules in the small joint capsules and extensive MAC in the lower-extremity
arteries that initiate along the elastic fibers (4, 9, 10). We previously found that a lack of CD73-
mediated adenosine enhanced the expression and activity of tissue-nonspecific alkaline
phosphatase (TNAP), a key enzyme that promotes calcification (11, 12), as well as altered
elastin deposition and SMC contractility in response to TGF-β (13).

Matrix GLA protein (Mgp) is a vitamin-K2-dependent protein with an unusual gamma-
carboxylation of five glutamate residues that enhance its affinity for calcium while preventing it
from mineralizing (14). Mgp inhibits calcification propagation by binding hydroxyapatite crystals
and stimulating their uptake by local phagocytosing macrophages (15). Mgp also modulates
osteogenic signaling by binding to BMP-2, preventing it from activating receptors and
upregulating Runx2, the key regulator of osteoblast differentiation and maturation (16, 17). Mice
that lack *Mgp* develop to term but die prematurely within approximately two months of age due
to extensive MAC, which leads to aortic rupture or heart failure (18). Aortas of *Mgp*-deficient
mice exhibit increased collagen accumulation and elastic fiber fragmentation within the medial
layer of the arteries, where calcification is also localized. This phenocopies MAC observed in
human patients with PAD MAC and ACDC patients (4).

Rapamycin binds the FK506-binding protein (FKBP12), converting it to a potent allosteric
inhibitor of the mammalian target of rapamycin (mTOR) complex. mTOR is broadly expressed
throughout the body, which controls many key processes such as energy balance, autophagy,
and proliferation (19-22). Rapamycin has been found to protect against the calcification of
vascular cells in in vitro and ex vivo models, as well as in in vivo models of chronic kidney
disease-induced MAC (23-25). We previously found rapamycin prevented calcification in in vitro
and in in vivo disease models of ACDC patient-specific induced pluripotent stem cells (iPSCs)
(11).

While CD73-deficient humans develop extensive MAC and broken elastin fibers in their lower
extremity large vessels, CD73-deficient mice do not phenocopy humans and do not exhibit
calcification in their vasculature (26, 27). As the MAC observed in the aorta *Mgp*-deficient mice
mirrors that seen in the affected vessels of ACDC patients, we sought to investigate whether
rapamycin prevented calcification in this genetic mouse model of MAC.

MATERIALS AND METHODS

Availability of Materials
We abide by the NIH Grants Policy on Sharing of Unique Research Resources, including the
NIH Policy on Sharing of Model Organisms for Biomedical Research (2004), NIH Grants Policy
Statement (2003), and Sharing of Biomedical Research Resources: Principles and Guidelines
for Recipients of NIH Grants and Contracts (1999), and the Bayh-Dole Act and the Technology
Transfer Commercialization Act of 2000. Materials generated in our laboratory are made

available for non-commercial research per established University of Pittsburgh Office of
Research IRB and MTA protocols.
155
Human tissue collection
De-identified human tissues were obtained, with informed consent from subjects, who were
enrolled in studies approved by the institutional review board (IRB) of the University of
Pittsburgh or Tufts Medical Center, per the Declaration of Helsinki. Personnel involved with
specimen handling underwent all required institutional training. Human coronary artery and tibial
artery tissues were excised and washed with a sterile rinsing solution (sterile PBS
supplemented with 2.5 µg/mL of fungicide (Gibco, 15290026), 0.05 mg/mL of gentamicin
(Gibco, 15710064), and 5 µg/mL of plasmocin (InvivoGen, ant-mpt-1). Tissues were then fixed
in 4% paraformaldehyde in PBS for 2 hours, placed in PBS, embedded in paraffin blocks for
pathology, and cut to a thickness of 10 µm.
166
Animals and cell line generation
Animal use was approved by the Institutional Animal Care and Use Committee at the University
of Pittsburgh. Mice were under the veterinary care of the University of Pittsburgh Division of
Laboratory Animal Resources, which adheres to the NIH policy on the Animal Welfare Act and
all other applicable laws. Facilities are under the full-time supervision of veterinarians and are
AAALAC-accredited. Our protocols follow the AVMA Guidelines on Euthanasia. *Mgp*
heterozygous mutant mice from a C57BL/6J background (strain# 023811 Jackson Laboratory,
Bar Harbor, ME) were bred to produce wild-type (+/+) and knockout (-/-) littermate controls.
Smooth muscle lineage-specific *Raptor* (*Mgp$^{-/-}$; Raptor$^{SMC-/-}$*) and *Rictor* (*Mgp$^{-/-}$; Rictor$^{SMC-/-}$*)
knockout mice were produced by breeding *Mgp$^{+/-}$* with Myh11-Cre-eGFP mice (strain# 007742
Jackson Laboratory, Bar Harbor, ME)(28), with Raptor-LoxP (strain# 013188 Jackson
Laboratory, Bar Harbor, ME)(29), or *Rictor*-LoxP (strain# 020649 Jackson Laboratory, Bar
Harbor, ME)(30). Genotypes were determined from tail snips that were incubated in DirectPCR
Lysis Reagent (Viagen Biotech) according to manufacturer instructions. The lysate was
amplified using OneTaq MasterMix (New England Biolabs) according to manufacturer
instructions with primers suggested by Jackson Laboratories.
183

Cell culture
SMC lines were generated from three- to five-week-old male and female *Mgp$^{+/+}$* and *Mgp$^{-/-}$* mice
according to previously described protocols (13). *Mgp$^{+/+}$* and *Mgp$^{-/-}$* aortic SMC lines were
cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco, Waltham, MA) supplemented
with 20% FBS (FBS; R&D Systems, Minneapolis, MN) and 100 U/mL penicillin-streptomycin
(P/S; Gibco, Waltham, MA). Growth media was changed every three days, and cells were split
1:2 when confluent. Post-expansion, 25,000–50,000 cells/cm² were plated in DMEM
supplemented with 10% FBS and 100 U/mL penicillin-streptomycin, and grown to confluence.
Before all experiments/treatments, cells were serum-starved in DMEM with 0.5% FBS for 48 h.
Incubation under reduced serum conditions has been proven to be a beneficial technique for
studying smooth muscle cell remodeling and contractility in vitro (31, 32). Cells were treated
with rapamycin (Novus Biologicals) at concentrations of 200 nM or the same volume of DMSO
as vehicle control for 14h. For autophagy flux assessment, cells were exposed to either
bafilomycin A1 100 nM (Sigma-Aldrich) or rapamycin 200 nM for a duration of 24h following
treatment intervention. All chemicals were dissolved in DMSO and administered in equal volume
as vehicle control.
200
201   *Murine Tissue Extraction*

Mice were sacrificed by asphyxiation with carbon dioxide, followed by perforation of the diaphragm. Tissue extraction began with dissection through the abdominal wall and then perfusion of the heart with 50 mL of DPBS supplemented with 1:100 Amphotericin B (Gibco,15290026) and 1:200 Gentamicin (Gibco, 15710064). The aortas were maintained in this solution while the adventitia was removed in a petri dish. Tissues were then fixed in 4% paraformaldehyde in PBS for 2 hours and then placed in PBS. Tissues were brought to microCT and then embedded in paraffin and cut to a thickness of 10 μm.

In Vivo Rapamycin Injections
Male and female *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice received intraperitoneal injections (1x or 3x per week) of either vehicle or 5 mg/kg rapamycin dissolved in 0.3 mg/ml peanut oil beginning at 10 days of age. This process continued until the natural death or a collection timepoint.

Micro-Computed Tomography
Five-10mm in length segments of mouse arteries sealed in parafilm and stabilized inside a holder were imaged by a Scanco μCT 50 (Scanco Medical, Brüttisellen, Switzerland) microcomputed tomography system with a 3.4 μm voxel resolution, with a 55 KVp beam energy, and 145 μA (1500 ms exposure per view). The Scanco μCT software (HP, DECwindows Motif 1.6) was used for 3D volume reconstruction, and the Scanco morphometry and densitometry software was used for the processing of images. Measurements included the total volume and density of mineral deposits after segmentation from the background with a 200 mg/cc threshold.

Western Blot Analysis
Cells were lysed in 1% CHAPS hydrate, 150 mmol/L sodium chloride, and 25 mmol/L HEPES buffer supplemented with 1× protease and phosphatase inhibitor (Sigma-Aldrich). Cells were scraped into microcentrifuge tubes, vortexed for 5 minutes, frozen/thawed for 5 to 8 cycles, then centrifuged at 12 000×g for 10 minutes at 4°C. Supernatant protein concentration was determined using Pierce BCA protein assay kit (Thermo Fisher, Waltham, MA). Ten micrograms of protein were used to prepare lysate with 1 × Pierce LDS sample buffer nonreducing (Thermo Fisher, Waltham, MA) and 1 × NuPAGE sample reducing agent (Novex, Waltham, MA). Lysates were denatured at 95°C for 15 min and then electrophoresed on 4%–20% TGX stain-free polyacrylamide gel (Bio-Rad, Hercules, CA) in 1 × Tris/Glycine/SDS buffer (Bio-Rad, Hercules, CA) at 120 V for 50 min. Protein was transferred onto a 0.2 μm nitrocellulose membrane in prepared 1 × Towbin buffer with 100% ethanol (EtOH) at 1 A and 25 V for 30 min using the Trans-Blot Turbo Transfer System (Bio-Rad, Hercules, CA). Membranes were blocked in 1:1 Odyssey blocking buffer (Li-COR, Lincoln, NE) and PBS for 1 h at room temperature, followed by primary antibodies incubation of LC3 (Abcam, ab192890, RRID: AB_2827794) and α-tubulin (926-42213, LI-COR, RRID: AB_2637092) in 1:1 Intercept blocking buffer (LI-COR, 927-70001) plus 0.1% Tween 20 (PBS-T) at 4°C overnight. Membranes were washed in PBS-T three times for 5 min, then incubated in secondary antibodies, anti-rabbit(LI-COR, 926-32211, RRID: AB_621843) or anti-mouse antibody (LI-COR, 926-32210, RRID: AB_621842), at room temperature for 1 h. Membranes belonging to the same experimental set were imaged simultaneously on an Odyssey CLx (LI-COR, Lincoln, NE), and band intensity quantification was performed with Image Studio (Version 5.2, LI-COR, Lincoln, NE) software. Individual bands were normalized to α-tubulin, and each treatment group's fold change was compared with each gel's vehicle control lanes. For sequential antibody incubations, membranes were stripped in 1 × NewBlot Nitro Western Blot Stripping Buffer (LI-COR, Lincoln, NE) for 10 min, followed by three washes in PBS.

Verhoeff Van Gieson, Von Kossa, Masson's Trichrome and Picrosirius Red Staining

Mice aortic tissues were removed from mice through the abdominal wall under anesthesia conditions. Then aortic tissue was fixed with 4% paraformaldehyde and embedded in paraffin. The embedded specimens were transversely sectioned at 10 μm on a microtome cryostat (Microm HM 325). Slides with the adhered paraffin aortic sections were warmed to 65°C for 1 h and then deparaffinized through xylene, rehydrated with serial incubation in graded alcohol baths, and stained with Verhoeff–van Gieson for elastic fiber visualization (Polysciences, 25089-1), Von Kossa for calcification (Polysciences, 24633-1), Masson's for collagen fibers (Polysciences, 25088-1) and Picrosirius Red staining (Polysciences, 24901-500) for collagen quantification according to manufacturer's instructions. Verhoeff Van Gieson, Von Kossa, Masson's Trichrome images were captured using Fritz PreciPoint scanner with 20x Nikon objective. Collagen staining was assessed by measuring the intensity of collagen in the vessel wall normalized to scale bar. Picrosirius Red staining images captured collagen with polarized transmitted light microscopy (Nikon Eclipse NI-E) stand with a plan Apo 20x Nikon objective and a Teledyne Prime BSI Express camera with an exposure time of 100msec. The percentage of collagen in the medial layer was measured using Nikon Elements 6.02.03. The percentage of medial thickness of the medial layer was measured using ImageJ software (Java 8), where the medial thickness was determined by measuring the distance between the inner and outer elastic lamina at four sites per vessel. The percentage of calcification in the medial layer was measured using ImageJ software (Java 8) where the area of calcification was normalized to total medial area. The tortuosity of elastic fibers was measured by selecting three representative areas with the whole vessel. ImageJ was used to trace and measure the length of five elastic fibers and shown as a ratio over the vessel length. The number of breakages in the elastic lamina was assessed by counting the breaks throughout the entire aorta. All length measurements were normalized to the scale bar units.

Immunofluorescence

Slides with the adhered paraffin aortic sections were warmed to 65°C for 1 h and then deparaffinized through xylene, rehydrated with serial incubation in graded alcohol baths. Slides were boiled in citric acid-based antigen retrieval solution (H-3300, Vector Labs) for 20 minutes. Slides were cooled in the unmasking solution for 1 hour, washed in 1x PBS and placed in blocking buffer (500 mL PBS 0.3g Fish Skin Gelatin and 10% horse serum) for 1 hour. Tissues were incubated overnight at 4°C with antibodies for α-SMA (Abcam, ab5694, RRID: AB_2223021, 1:50), Ki67 (Abcam, ab15580, RRID: AB_443209, 1:250), LC3 (Abcam, ab192890, RRID: AB_2827794, 1:50), Mgp (Proteintech, 10734-I-AP, RRID: AB_2297660, 1:25), Myh11 (Invitrogen, MA5-4284, RRID: AB_2911986, 1:50), Runx2 (ABclonal, A2851, RRID: AB_2764676 1:100). Cells were washed 3x for 5 minutes each with PBS, 0.1% TWEEN 20, then once with PBS, and then incubated with secondary antibodies (Invitrogen, A-11012, A-11008) diluted in a blocking solution for 1 hour. Cells were washed 3x for 5 minutes each with PBS, 0.1% TWEEN 20, then once with PBS, then mounted with Fluoroshield Mounting Medium with DAPI (Ab104139, Abcam). All signals were optimized to a rabbit polyclonal IgG (Vector, I-1000, RRID: AB_2336355, 1:50) at the same concentration of the specific antibodies. Slides were imaged within 24 hours of mounting. Cells were imaged within 24 hours of mounting. Images were obtained using a Nikon NIE-I microscope at 20x objective. The measured intensity of a given stain was normalized to DAPI intensity of that same image. In ImageJ, each image was split into blue, red, and green channels, and the pixel intensity of the red and blue channels was measured (33). Representative images are selected based on quality, and if the assay includes quantification, the image that shows the average of the quantification across multiple images.

Blinding Procedures

Professionally trained technicians handled animal welfare, including weaning and mouse line expansion, diets, sacrifice, and surgery. For experiments, blinding procedures were used, and individuals conducting and analyzing experiments did not know the treatment groups until after the analysis was performed.

Sample Size and Statistics

Statistical analysis was performed with GraphPad Prism 9.2 (GraphPad Software, Inc) and data shown are mean ± standard deviation (SD). All experiments used at least n = 4 biological replicates and were run in technical duplicates, exact n per group specified in figure legends. Mantel-Cox test and Gehan-Breslow-wilcoxon test were used for Kaplan-Meier analysis. Statistical comparisons between the two groups were performed by either nonparametric Mann-Whitney U test or Kruskal-Wallis test when n ≤ 6 ≤ or parametric Two-way ANOVA when n > > 6. Statistical analysis used, exact n values, biological replicates, and P values are stated within each figure legend. A p-value equal to or less than 0.05 will be considered statistically significant.

RESULTS

MAC in MGP-deficient mice phenocopies human MAC.

The MAC is localized in the medial layer of arteries along the elastic fibers. Von Kossa staining for calcification illustrates that $Mgp^{-/-}$ mouse aortas have extensive mineralization in the medial layer that phenocopies the mineralization seen in MAC of the human tibial arteries, and is distinct from neointimal calcification found in the necrotic core of coronary artery with atherosclerotic plaque (Figure 1A). Mgp is a potent inhibitor of soft tissue calcification. $Mgp^{-/-}$ mice do not produce functional Mgp protein (Figure 1B, 1C) and, as a result, develop extensive MAC pathology in large vessels (18).

Rapamycin prolongs lifespan and reduces arterial mineral density in $Mgp^{-/-}$ mice.

Humans with homozygous inactivating mutations in *NT5E* develop extensive MAC in their lower extremity arteries (9). While the *Nt5e*-knockout mice model shows altered renal function, diminished control of the glomerular arteriolar tone, elevated atherogenesis, elevated thrombotic occlusion, and elevated hypoxia-induced vascular leakage (26), these mice do not mimic the human vascular calcification phenotype (26, 27, 34). Jin et al. circumvented this issue using a patient-specific in vitro and in vivo disease modeling system using iPSC technology. They demonstrated that patient-specific ACDC-iPSCs developed extensive calcification in the in vivo teratoma model relative to control patient iPSCs. Furthermore, they found that treating mice bearing ACDC-iPSC teratomas with rapamycin reduced calcification (11).

As CD73-deficient mice do not sufficiently recapitulate the MAC observed in ACDC patients, we used the $Mgp^{-/-}$ mice as an in vivo model of MAC to test the hypothesis that rapamycin could reduce vascular calcification in vivo. $Mgp^{+/+}$ and $Mgp^{-/-}$ mice were administered 5 mg/kg rapamycin or the same volume of vehicle (DMSO) three times per week starting at 10 days old until natural death (Figure 2A). We observed that rapamycin treatment extended the lifespan of $Mgp^{-/-}$ mice (T_{50} 61 days) compared to the $Mgp^{-/-}$ mice treated with the vehicle (T_{50} 35 days), which start to die as early as day 20 (Figure 2B). Both male (n=11) and female (n=8) mice were used; however, no sex differences were observed between $Mgp^{-/-}$ mice that had received either vehicle or rapamycin treatment.

We next investigated whether calcification content was altered in mice receiving rapamycin. Mice receiving vehicle, one (1x), or three doses (3x) of 5 mg/kg rapamycin per week, were sacrificed at day 25, and mineral content was assessed via microCT (Figure 3A). We selected

the 25-day time point because we observed that $Mgp^{-/-}$ mice start dying around that time (Figure
2B). Whole aorta microCT analysis found no differences in the total calcified deposits volume
under either treatment regime; however, mice receiving rapamycin 3x per week exhibited a
decreased mineral density compared to the control group (Figure 3B). Based on this data, we
proceeded with 3x/week for the remaining studies. Von Kossa stain showed that the aortas of
$Mgp^{+/+}$ mice exhibit no mineralization, while both vehicle- and rapamycin-treated $Mgp^{-/-}$ mice
showed extensive calcification compared to $Mgp^{+/+}$ mice. There were no discernible differences
in the calcified area or the medial thickness of vehicle or rapamycin groups of $Mgp^{-/-}$ (Figure
3C). RUNX2 is a key transcription factor promoting vascular cell osteogenic differentiation (6).
Immunofluorescent staining revealed that while $Mgp^{-/-}$ mice showed elevated levels of Runx2
compared to $Mgp^{+/+}$ mice, rapamycin treatment did not reduce Runx2 protein levels in $Mgp^{-/-}$
mice compared to $Mgp^{-/-}$ mice treated with vehicle (Figure 3D). These findings suggest that
while rapamycin doubles the lifespan of $Mgp^{-/-}$ mice, it does not lead to decreased calcification
volume, but does reduce the density of mineralized deposits in $Mgp^{-/-}$ mice.
367
Rapamycin alters SMC phenotype and ECM content in vivo
Rapamycin has a broad spectrum of effects on SMCs. Rapamycin inhibits SMC migration and
proliferation by inducing the cyclin-dependent kinase inhibitors p27^{kip} and p21^{cip} to promote G1-
S cell cycle arrest (35, 36). Rapamycin also induces SMC differentiation by inhibiting the mTOR-
target S6K1 via AKT activation, promoting a contractile phenotype by enhancing the protein
levels of SMC contractility markers smooth muscle α-actin (α-SMA) and myosin heavy chain 11
(Myh11) (37). Further, rapamycin was shown to protect against calcification of vascular cells in
in vitro and ex vivo models (11, 23). While rapamycin did not reduce calcification volume in $Mgp^{-/-}$
$^{/-}$ vessels we hypothesized that it could perhaps prevent aortic rupture by maintaining SMC
contractile phenotype or stabilize the ECM.
378
Masson's trichrome staining shows that $Mgp^{-/-}$ vehicle vessels are highly remodeled, with
acellular areas exhibiting observable increased collagen staining (Figure 4A). Picrosirius
staining enabled the quantification of collagen content and revealed that thicker mature collagen
in the red-orange colors and thinner immature collagen in green-yellow. In both the vehicle and
the rapamycin treatment groups, the $Mgp^{-/-}$ exhibited higher levels of both mature and immature
collagen compared to the cognate $Mgp^{+/+}$ mice. Within the $Mgp^{-/-}$ genotype, rapamycin treatment
reduced both types of collagen deposition compared to vehicle control (Figure 4B). This
suggests that rapamycin helps restore or stabilize collagen homeostasis within the ECM,
potentially preventing excessive fibrosis and maintaining tissue integrity (38), which may
improve resilience against rupture. Verhoeff-Van Gieson (VVG) staining revealed that the aortas
of $Mgp^{-/-}$ mice exhibited less robust staining for elastin and less tortuous fibers in $Mgp^{-/-}$
compared to $Mgp^{+/+}$. While rapamycin did not alter elastic fiber tortuosity in $Mgp^{-/-}$ mice
compared to vehicle controls, the number of elastin breakages was higher in $Mgp^{-/-}$ mice
compared to $Mgp^{+/+}$ in both groups and rapamycin treatment reduced the number of breakages
in $Mgp^{-/-}$ mice compared to vehicle (Figure 4C).
394
Immunofluorescent staining was used to quantify SMC contractile proteins α-SMA and Myh11.
In $Mgp^{+/+}$ we observed no differences in α-SMA levels between vehicle and rapamycin
treatment; however with vehicle treatment, $Mgp^{+/+}$ mice exhibit higher α-SMA than $Mgp^{-/-}$ mice.
Rapamycin treatment rescued α-SMA levels in $Mgp^{-/-}$ mice compared to vehicle alone, such that
no differences were observed between $Mgp^{-/-}$ mice treated with rapamycin and the $Mgp^{+/+}$
vehicle or $Mgp^{+/+}$ rapamycin groups. Similarly, $Mgp^{-/-}$ mice treated with the vehicle exhibit
reduced Myh11 levels compared to $Mgp^{+/+}$ mice treated with the vehicle (Figure 4D). These
results suggest rapamycin promotes the contractile phenotype of SMCs of $Mgp^{-/-}$ mice. As the

medial layer is remodeled in the *Mgp*$^{-/-}$ mice, we also measured the mitotic marker Ki67 and
found that at baseline, Ki67 levels were significantly elevated in *Mgp*$^{-/-}$ aorta while nearly
undetectable in *Mgp*$^{+/+}$. However, rapamycin treatment did not alter Ki67 levels in either of the
two genotypes (Figure 4E). Together, these data suggest that rapamycin promotes beneficial
medial layer remodeling and a more contractile SMC phenotype in *Mgp*$^{-/-}$ mice aorta.

408
Rapamycin lowered LC3 levels in *Mgp*$^{-/-}$ aorta, but there were no observable changes in
autophagy flux.
Autophagy is a ubiquitous process and contributes to bone development and the differentiation
of osteoblasts and osteoclasts (39). One study found that autophagy reduces vascular
calcification by limiting the release of pro-calcific matrix vesicles (40). Another found that in vitro
autophagy stimulation with valproic acid protects against phosphate-induced calcification, while
its inhibition increases extracellular vesicle release in SMCs (25). LC3 is a protein that functions
at the initiation of autophagosome formation and serves as an autophagy marker. LC3 signal
was significantly higher in *Mgp*$^{+/+}$ aorta at baseline compared to *Mgp*$^{-/-}$ mice. Within each
genotype, rapamycin treatment significantly lowered LC3 puncta in *Mgp*$^{+/+}$ compared to *Mgp*$^{-/-}$
mice (Figure 5A).

420
The quantification of LC3 puncta hints at diminished autophagy, but immunofluorescent staining
of fixed tissues cannot assess autophagy flux because it provides only a single snapshot of the
autophagic process, which does not reflect changes in the dynamic turnover of
autophagosomes or the lysosomal clearance of LC3-positive autophagic vesicles. LC3-I is
cleaved to LC3-II when it is incorporated into autophagosomes, but this process is dynamic, and
LC3-II can also be degraded during autophagy. The comparison of LC3-II to LC3-I in the
presence of an autophagy inhibitor like bafilomycin enables autophagy flux observation (41). We
isolated SMCs from *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice to evaluate autophagy flux. SMCs were treated with
vehicle (DMSO) or 200 nM rapamycin for 14h and then with vehicle or 100 uM or the late-phase
autophagy inhibitor bafilomycin for 24h. Immunoblot analysis revealed that rapamycin alone did
not increase LC3-II levels compared to vehicle controls in either the *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ SMCs.
Bafilomycin increased LC3-II levels in both *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ SMCs compared to their cognate
vehicle controls, however the addition of rapamycin did not further alter LC3-II levels compared
to bafilomycin (Figure 5B, left graph). Comparing LC3-II to LC3-I showed identical trends
(Figure 5B, right graph), suggesting that global LC3 levels and flux are similar in *Mgp*$^{+/+}$ and
Mgp$^{-/-}$ SMCs.

437
Genetic deletion of mTOR signaling in the SMCs does not replicate rapamycin treatment
***Mgp*$^{-/-}$ mice.**
Figure 2 shows that rapamycin treatment doubles the lifespan of *Mgp*$^{-/-}$ mice compared to
vehicle alone. However, our subsequent inquiries revealed that rapamycin did not decrease the
volume of calcification as was seen in in vivo iPSC teratoma models (11), while it did promote
less dense mineral formation, and beneficial SMC phenotype and EMC remodeling, albeit not to
levels seen in the *Mgp*$^{+/+}$ animals. mTOR can form two different protein complexes known as
mTORC1, bound to Raptor, and mTORC2, bound to Rictor. The former is acutely sensitive to
rapamycin administration, while the latter requires chronic rapamycin dosing for inhibition (21).
To investigate whether rapamycin's effects on the *Mgp*$^{-/-}$ mouse are acting through either of
these complexes, we generated *Mgp*$^{-/-}$ mice with SMC-specific deletion of the mTOR complex
proteins Raptor and Rictor (28-30). Fully differentiated SMCs expressing *Myh11* will
constitutively express Cre recombinase to inactive *Raptor* and *Rictor* gene expression
specifically in SMCs. *Mgp*$^{-/-}$;*Myh11Cre-GFP;Raptor*$^{f/f}$ mice (*Mgp*$^{-/-}$;*Raptor*$^{SMC-/-}$) were bred to
examine the effects of a nonfunctional mTORC1 complex, while *Mgp*$^{-/-}$;*Myh11Cre-GFP;Rictor*$^{f/f}$
(*Mgp*$^{-/-}$;*Rictor*$^{SMC-/-}$) mice knockout was bred to examine a nonfunctional mTORC2 complex.

454
Similar to $Mgp^{-/-}$ mice treated with vehicle in Figure 2, Figure 6 shows that only 50% of the $Mgp^{-/-}$
$;Rictor^{SMC-/-}$ and $Mgp^{-/-};Rictor^{SMC+/+}$ are alive at 40 days, indicating that rapamycin acting via
mTORC2 does not recapitulate the effects of rapamycin on increasing $Mgp^{-/-}$ mice lifespan. The
50% survival rate of $Mgp^{-/-};Raptor^{SMC+/+}$ mice was similar to $Mgp^{-/-}$ treated with vehicle, $Mgp^{-/-}$
$;Rictor^{SMC-/-}$ and $Mgp^{-/-};Rictor^{SMC+/+}$, however the 50% survival rate of $Mgp^{-/-};Raptor^{SMC-/-}$ mice
dropped to 22 days. As the Cre strain being used is a non-inducible constitutive expression
system, this data from $Mgp^{-/-};Raptor^{SMC-/-}$ mice suggest that mTORC1 is critical to the growth and
maturation of the vasculature post birth. Together, these in vivo data suggest that the life-
extending effects of rapamycin in the $Mgp^{-/-}$ mouse are not acting through the mTORC2
complex. Additional experiments, such as a tamoxifen inducible Cre system, would need to be
used to fully address the contribution of mTORC1 to mirror effects seen by rapamycin injection.
466
DISCUSSION
In this study, we demonstrated that rapamycin treatment increased the lifespan in $Mgp^{-/-}$ mice,
but rapamycin does not reduce MAC per se; while mineral density is lower with rapamycin
treatment the mineral volume remains unchanged. In mice, rapamycin has been shown to
increase maximum lifespan and delay the onset of cancer as hyperactivity of the mTOR
pathway leads to the expression and activation of many oncogenes like PI3K, Akt, and eIF4E.
Hence, rapamycin can slow the proliferation of tumor cells by causing cell arrest, promoting
apoptosis, and blocking angiogenesis in tumor growths (42). Our data show that the increase in
lifespan observed in $Mgp^{-/-}$ mice following rapamycin treatment is associated with improvements
in SMC contractile protein expression and a normalization of collagen content in the vessel wall,
but not with enhanced autophagy flux.
478
Collagen, the most abundant protein in the ECM, plays a crucial role in maintaining the
structural integrity of the blood vessels. Collagen dysregulation or degradation leads to
increased vessel distensibility and the risk of rupture (43, 44). Aging mice homozygous for a
deletion in the gene coding for collagen alpha 1(*Col1a1*) are at a significantly increased risk of
aortic dissection and rupture (45). Our results align with a study which showed elevated
collagen deposition in $Mgp^{-/-}$ mice aorta (18). In 25-day-old $Mgp^{-/-}$ mice, rapamycin treatment
reduced collagen content in the medial layer compared to vehicle-treated mice, bringing it to
levels observed in $Mgp^{+/+}$. Total adventitial collagen content in the mice aorta increases with age
(46), suggesting the differential distribution we observed was due to the young age of the mice
in our study. The decrease in collagen content following rapamycin treatment and the increase
in SMC contractile protein SMA may contribute to the prolonged lifespan of $Mgp^{-/-}$ mice by
improving both the structural stability and functional resilience of the vascular walls.
491
While rapamycin is known to inhibit SMC proliferation and is used in drug-eluting stents, there is
limited research on the effects of rapamycin on vascular morphology in the presence of
calcification (36, 47). It was previously shown that treatment with rapamycin limits the
progression of abdominal and thoracic aortic aneurysms in mice and preserves elastic lamina
integrity (48, 49). In a hyperphosphatemic (high-phosphate, high-adenine diet) rat model,
rapamycin treatment inhibited aortic calcium deposition and reduced the osteogenic markers
Msx2 and *Osx* while increasing the *Opn* and *Acta2* gene expression via Klotho upregulation
(50). Opposing this diet-induced model of calcification, we did not observe a rapamycin-induced
reduction of the osteogenic marker Runx2 in *Mgp*-deficient mice. Our approach used a genetic
model of MAC to investigate whether rapamycin could inhibit calcification in vivo. *Mgp*-deficient
mice develop normally to birth without any overt abnormalities. At approximately one week old
$Mgp^{-/-}$ mice start exhibiting calcification of the vessels, suggesting that other gene products
might compensate to prevent calcification during development and immediately post-birth (18).

These observations highlight the differences between genetically induced MAC and high
phosphate feeding models. Diet-induced calcification must overcome the effect of endogenous
Mgp, which is a potent inhibitor of calcification, while the genetic removal of *Mgp* results in
calcification from the circulating calcium and phosphate and activation of BMP signaling (14,
17).
510
Rapamycin has also been shown to preserve the differentiated phenotype of vascular SMCs
through IRS-1/P3K/AKT2 pathway (37). Our study did observe that rapamycin enhanced the
expression of a SMC contractile proteins in $Mgp^{-/-}$ mice. *Sma* expression typically occurs earlier
than *Myh11* expression during SMC development (51). In contrast, *Myh11* expression occurs
later, during the maturation of SMCs. One study examined SMC contractile gene expression in
response to rapamycin in vitro (37), while our study looked at the SMC contractile markers in
vivo. Further, the presence of calcification activates mechanosensing pathways that once
activated may prevent full re-expression of SMC contractile markers (52). However, further
investigation is needed to elucidate how these mechanisms are operating in the context of the
$Mgp^{-/-}$ mouse.
521
MAC is characterized by mineral deposits in the medial vessel layer and frequently occurs in
patients with PAD. Vascular calcification is an active biological process with many driving
forces, including osteogenic dedifferentiation of SMCs, inflammatory signals,
hyperphosphatemia, and extracellular matrix remodeling (53). In this study, rapamycin
prolonged the lifespan of mice who developed significant MAC shortly after birth. Interestingly,
even though rapamycin has been shown to stimulate the cellular process of autophagy (54)—
which can recycle components of the ECM, including mineral deposits (55)—calcification
volume was not reduced in rapamycin treated *Mgp-/-* mice. However, microCT data showed that
the mineral density was significantly reduced in $Mgp^{-/-}$ mice treated with rapamycin. The
decrease in mineral density without a change in calcification volume could suggest a shift from
compressed, organized macrocalcifications to more dispersed calcium-rich extracellular vesicles
and amorphous calcium phosphate deposits (7, 56). Amorphous calcium phosphate, primarily
composed of calcium and phosphate ions, represents the early stages of calcification (57). It
subsequently acts as a precursor for the nucleation and growth of hydroxyapatite crystals within
the ECM. Stabilization of hydroxyapatite crystals further contributes to the transition into a bone-
like phenotype. One reason that a lower mineral density did not affect total calcification volume
could be a difference in the pattern of mineral deposition. For example, the mineral may form in
a less compact, more porous manner, leading to a lower overall density despite occupying the
same spatial volume. Alternatively, the reduction in density could result from the breakdown or
remodeling of pre-existing mineral deposits. This suggests that changes in mineral structure, if
explored at a further timepoint may impact calcification volume. Alternatively, as Mgp also
functions to inhibit BMP2 signaling, rapamycin may not interfere with that process, such that
changes in dosage or duration of rapamycin treatment would still not alter calcification content in
the vessel wall. Another explanation is that the activated autophagy pathways cannot overcome
the rapid deposition of hydroxyapatite that occurs in $Mgp^{-/-}$ mice. Frauscher et al. demonstrated
that induction of autophagy by rapamycin treatment decreased calcification in the vasculature of
uremic DBA/2 mice (58). This argument, along with our prior data from iPSC-based disease
modeling (11), led us to examine autophagy flux in SMCs. However, we did not observe a
difference in autophagy flux between $Mgp^{+/+}$ and $Mgp^{-/-}$. It is important to note that the $Mgp^{-/-}$ is
on a C57BL6J background, perhaps explaining the differences observed in these in vivo
models.
553
554 Rapamycin is an inhibitor of mTOR, but mTOR pathways are complex. mTOR forms two distinct
555 protein complexes known as mTORC1 and mTORC2. mTORC1, activated by amino acids and

allosterically inhibited by rapamycin, requires the protein named Raptor (59). mTORC1 is acutely sensitive to rapamycin, while only after prolonged rapamycin treatment (>6 hours) does mTORC2 (requiring Rictor) acquire a sensitivity to rapamycin (22, 60). In the $Mgp^{-/-};Raptor^{SMC-/-}$ mice, induction of Cre to knockout *Raptor* was more lethal than the $Mgp^{-/-};Raptor^{SMC+/+}$ or $Mgp-/-$ mice. This suggests that *Raptor* is essential for the growth and maintenance of SMCs in the vasculature. This essential role in growth and development is perhaps unsurprising, considering that global *Raptor* knockout mice die in utero (61). Moreover, there was no benefit to lifespan in the $Mgp^{-/-}$:Rictor$^{SMC-/-}$ mice. The survival curves for the $Mgp^{-/-}$:Rictor$^{SMC+/+}$ and $Mgp^{-/-};Rictor^{SMC-/-}$ mice, targeting mTORC2 signaling, do not diverge at any point and are statistically insignificant and also mirror that seen in the $Mgp^{-/-}$ strain.

Arterial stiffness is strongly linked to heart failure (62), and these non-atherosclerotic vascular pathologies are much more common in some populations, such as African-ancestry individuals who are known to have a higher risk of cardiovascular disease events and mortality compared to Caucasians (63-67). This excess cardiovascular disease risk is largely due to hypertensive and peripheral vascular disease versus atherosclerotic disease (63-65, 68-70). As such, it is critical to define drivers of MAC such as MGP, which in some clinical studies in humans has been associated with heart failure and MAC (71). Impairment in MGP activation, which would phenocopy genetic deletion and lead to arterial stiffening and peripheral artery disease, is linked to a higher risk of heart failure due to chronically elevated afterload leading to left ventricle remodeling (72, 73). As the mice that lack *Mgp* are prone to aortic rupture or heart failure, we speculate that rapamycin's effect on the mice's lifespan may be due to rapamycin acting on the cardiac tissues.

Our study suggests that rapamycin's life-extending benefit in $Mgp^{-/-}$ mice may act through improvement in vascular function, which could prevent aortic rupture or through another organ system, possibly the heart. However, further studies are needed to examine the impact of rapamycin on cardiac function in $Mgp^{-/-}$ mice. Rapamycin has been shown to increase cardiomyocyte autophagy (74, 75), limit cardiomyocyte death, and attenuate cardiomyocyte hypertrophy and cardiac remodeling by enhancing mTORC2 signaling while simultaneously inhibiting mTORC1 signaling (76). Rapamycin had a protective effect on cardiomyocytes isolated from the old mice compared to old mice not treated with rapamycin, and improved ejection fraction, reduced cardiac hypertrophy and inflammation, and downregulated ANP-induced stress response in vivo (77, 78). In mice with decompensated cardiac hypertrophy, rapamycin improved cardiac function, such as left ventricular end-systolic dimensions, fractional shortening, and ejection fraction in mice compared to the control group (79). Three months of rapamycin treatment in aged mice increased lifespan, decreased heart rate and heart weight (80). However, a calcified phenotype was not noted in any of these animal studies.

CONCLUSION
This study provides novel insights into the effects of rapamycin on medial arterial calcification (MAC) using the *Mgp*-deficient murine model. While rapamycin did not reduce the volume of calcification in the vasculature, it significantly prolonged the lifespan of $Mgp^{-/-}$ mice, possibly through stabilizing vascular structure. This was evidenced by reduced collagen content and enhanced SMC contractile protein expression, which collectively suggest improved resilience against vascular rupture. Interestingly, while the volume of calcification was not changed, rapamycin treatment reduced mineral density. Further, our study underscores the importance of distinguishing between mTORC1- and mTORC2-mediated pathways. Rapamycin's life-extending effects appeared independent of autophagy flux or mTORC2 signaling, while the critical role of mTORC1 in vascular development and maintenance was highlighted, as $Mgp^{-/-}$

;Raptor^SMC-/- died earlier than *Mgp^-/-* mice. These findings illustrate rapamycin's role in modifying
calcification characteristics and enhancing vascular integrity without altering overall calcification
volume. Future investigations should focus on elucidating rapamycin's systemic effects, such as
its impact on cardiac function and/or the immune system, as well as understanding how it leads
to less-dense mineral deposits, to clarify its protective mechanisms in the context of MAC and
cardiovascular health.
612

613 **ACKNOWLEDGEMENTS**
614 We want to thank the histology core and the Center for Biologic Imaging at the University of
615 Pittsburgh for their help and support with microscopy imaging. Some figures were created with
616 Biorender.com.
617
618 **SOURCES OF FUNDING**
619 This publication was supported by the NIH grants K22HL117917, R01HL176595, The Winters
620 Foundation, The McKamish Family Foundation (St. Hilaire), and the American Heart Association
621 (Behzadi). MicroCT analysis and processing was performed on a system supported by NIH
622 S10-OD021533 (Verdelis).

623
624 **DISCLOSURES**
625 None.
626
REFERENCES

1.   **St Hilaire C**. Medial Arterial Calcification: A Significant and Independent Contributor of
Peripheral Artery Disease. *Arterioscler Thromb Vasc Biol* 42: 253-260, 2022.
2.   **O'Neill WC, Han KH, Schneider TM, and Hennigar RA**. Prevalence of
nonatheromatous lesions in peripheral arterial disease. *Arterioscler Thromb Vasc Biol* 35: 439-
447, 2015.
3.   **Narula N, Dannenberg AJ, Olin JW, Bhatt DL, Johnson KW, Nadkarni G, Min J,
Torii S, Poojary P, Anand SS, Bax JJ, Yusuf S, Virmani R, and Narula J**. Pathology of
Peripheral Artery Disease in Patients With Critical Limb Ischemia. *J Am Coll Cardiol* 72: 2152-
2163, 2018.
4.   **Markello TC, Pak LK, St Hilaire C, Dorward H, Ziegler SG, Chen MY, Chaganti K,
Nussbaum RL, Boehm M, and Gahl WA**. Vascular pathology of medial arterial calcifications in
NT5E deficiency: implications for the role of adenosine in pseudoxanthoma elasticum. *Mol
Genet Metab* 103: 44-50, 2011.
5.   **Lanzer P, Hannan FM, Lanzer JD, Janzen J, Raggi P, Furniss D, Schuchardt M,
Thakker R, Fok PW, Saez-Rodriguez J, Millan A, Sato Y, Ferraresi R, Virmani R, and St
Hilaire C**. Medial Arterial Calcification: JACC State-of-the-Art Review. *J Am Coll Cardiol* 78:
1145-1165, 2021.
6.   **Hortells L, Sur S, and St Hilaire C**. Cell Phenotype Transitions in Cardiovascular
Calcification. *Front Cardiovasc Med* 5: 27, 2018.
7.   **Villa-Bellosta R**. Role of the extracellular ATP/pyrophosphate metabolism cycle in
vascular calcification. *Purinergic Signal* 19: 345-352, 2023.
8.   **Rutsch F, Ruf N, Vaingankar S, Toliat MR, Suk A, Hohne W, Schauer G, Lehmann
M, Roscioli T, Schnabel D, Epplen JT, Knisely A, Superti-Furga A, McGill J, Filippone M,
Sinaiko AR, Vallance H, Hinrichs B, Smith W, Ferre M, Terkeltaub R, and Nurnberg P**.
Mutations in ENPP1 are associated with 'idiopathic' infantile arterial calcification. *Nat Genet* 34:
379-381, 2003.
9.   **St Hilaire C, Ziegler SG, Markello TC, Brusco A, Groden C, Gill F, Carlson-Donohoe
H, Lederman RJ, Chen MY, Yang D, Siegenthaler MP, Arduino C, Mancini C, Freudenthal**

B, Stanescu HC, Zdebik AA, Chaganti RK, Nussbaum RL, Kleta R, Gahl WA, and Boehm
M. NT5E mutations and arterial calcifications. *N Engl J Med* 364: 432-442, 2011.
10.   **Baral A, F FH, Lee SC, Matthew BP, Ferrante EA, Brofferio A, Cudrici CD, Rollison**
SF, Chen MY, Boehm M, and Wen H. Images in Vascular Medicine: High-resolution CT
imaging of arterial calcification in the hands and legs of patients with CD73 deficiency. *Vasc*
Med 29: 342-344, 2024.
11.   **Jin H, St Hilaire C, Huang Y, Yang D, Dmitrieva NI, Negro A, Schwartzbeck R, Liu**
Y, Yu Z, Walts A, Davaine JM, Lee DY, Donahue D, Hsu KS, Chen J, Cheng T, Gahl W,
Chen G, and Boehm M. Increased activity of TNAP compensates for reduced adenosine
production and promotes ectopic calcification in the genetic disease ACDC. *Sci Signal* 9: ra121,
2016.
12.   **Moorhead WJ, 3rd, Chu CC, Cuevas RA, Callahan Jt, Wong R, Regan C, Boufford**
CK, Sur S, Liu M, Gomez D, MacTaggart JN, Kamenskiy A, Boehm M, and St Hilaire C.
Dysregulation of FOXO1 (Forkhead Box O1 Protein) Drives Calcification in Arterial Calcification
due to Deficiency of CD73 and Is Present in Peripheral Artery Disease. *Arterioscler Thromb*
Vasc Biol 40: 1680-1694, 2020.
13.   **Cuevas RA, Wong R, Joolharzadeh P, Moorhead WJ, 3rd, Chu CC, Callahan Jt,**
Crane A, Boufford CK, Parise AM, Parwal A, Behzadi P, and St Hilaire C. Ecto-5'-
nucleotidase (Nt5e/CD73)-mediated adenosine signaling attenuates TGFbeta-2 induced elastin
and cellular contraction. *Am J Physiol Cell Physiol* 324: C327-C338, 2023.
14.   **Murshed M, Schinke T, McKee MD, and Karsenty G**. Extracellular matrix
mineralization is regulated locally; different roles of two gla-containing proteins. *J Cell Biol* 165:
625-630, 2004.
15.   **Hackeng TM, Rosing J, Spronk HM, and Vermeer C**. Total chemical synthesis of
human matrix Gla protein. *Protein Sci* 10: 864-870, 2001.
16.   **Malhotra R, Burke MF, Martyn T, Shakartzi HR, Thayer TE, O'Rourke C, Li P,**
Derwall M, Spagnolli E, Kolodziej SA, Hoeft K, Mayeur C, Jiramongkolchai P, Kumar R,
Buys ES, Yu PB, Bloch KD, and Bloch DB. Inhibition of bone morphogenetic protein signal
transduction prevents the medial vascular calcification associated with matrix Gla protein
deficiency. *PLoS One* 10: e0117098, 2015.
17.   **Bostrom K, Tsao D, Shen S, Wang Y, and Demer LL**. Matrix GLA protein modulates
differentiation induced by bone morphogenetic protein-2 in C3H10T1/2 cells. *J Biol Chem* 276:
14044-14052, 2001.
18.   **Luo G, Ducy P, McKee MD, Pinero GJ, Loyer E, Behringer RR, and Karsenty G**.
Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature*
386: 78-81, 1997.
19.   **Haissaguerre M, Saucisse N, and Cota D**. Influence of mTOR in energy and metabolic
homeostasis. *Mol Cell Endocrinol* 397: 67-77, 2014.
20.   **Saxton RA, and Sabatini DM**. mTOR Signaling in Growth, Metabolism, and Disease.
Cell 168: 960-976, 2017.
21.   **Panwar V, Singh A, Bhatt M, Tonk RK, Azizov S, Raza AS, Sengupta S, Kumar D,**
and Garg M. Multifaceted role of mTOR (mammalian target of rapamycin) signaling pathway in
human health and disease. *Signal Transduct Target Ther* 8: 375, 2023.
22.   **Arriola Apelo SI, and Lamming DW**. Rapamycin: An InhibiTOR of Aging Emerges
From the Soil of Easter Island. *J Gerontol A Biol Sci Med Sci* 71: 841-849, 2016.
23.   **Zhan JK, Wang YJ, Wang Y, Wang S, Tan P, Huang W, and Liu YS**. The mammalian
target of rapamycin signalling pathway is involved in osteoblastic differentiation of vascular
smooth muscle cells. *Can J Cardiol* 30: 568-575, 2014.
24.   **He HQ, Law BYK, Zhang N, Qiu CL, Qu YQ, Wu AG, Han Y, Song Q, Zheng WL, Liu**
Y, He YZ, and Wong VKW. Bavachin Protects Human Aortic Smooth Muscle Cells Against
beta-Glycerophosphate-Mediated Vascular Calcification and Apoptosis via Activation of mTOR-

Dependent Autophagy and Suppression of beta-Catenin Signaling. *Front Pharmacol* 10: 1427, 2019.

25.   **Dai XY, Zhao MM, Cai Y, Guan QC, Zhao Y, Guan Y, Kong W, Zhu WG, Xu MJ, and Wang X**. Phosphate-induced autophagy counteracts vascular calcification by reducing matrix vesicle release. *Kidney Int* 83: 1042-1051, 2013.

26.   **Joolharzadeh P, and St Hilaire C**. CD73 (Cluster of Differentiation 73) and the Differences Between Mice and Humans. *Arterioscler Thromb Vasc Biol* 39: 339-348, 2019.

27.   **Ziegler SG, Ferreira CR, MacFarlane EG, Riddle RC, Tomlinson RE, Chew EY, Martin L, Ma CT, Sergienko E, Pinkerton AB, Millan JL, Gahl WA, and Dietz HC**. Ectopic calcification in pseudoxanthoma elasticum responds to inhibition of tissue-nonspecific alkaline phosphatase. *Sci Transl Med* 9: 2017.

28.   **Xin HB, Deng KY, Rishniw M, Ji G, and Kotlikoff MI**. Smooth muscle expression of Cre recombinase and eGFP in transgenic mice. *Physiol Genomics* 10: 211-215, 2002.

29.   **Sengupta S, Peterson TR, Laplante M, Oh S, and Sabatini DM**. mTORC1 controls fasting-induced ketogenesis and its modulation by ageing. *Nature* 468: 1100-1104, 2010.

30.   **Magee JA, Ikenoue T, Nakada D, Lee JY, Guan KL, and Morrison SJ**. Temporal changes in PTEN and mTORC2 regulation of hematopoietic stem cell self-renewal and leukemia suppression. *Cell Stem Cell* 11: 415-428, 2012.

31.   **Liu M, Espinosa-Diez C, Mahan S, Du M, Nguyen AT, Hahn S, Chakraborty R, Straub AC, Martin KA, Owens GK, and Gomez D**. H3K4 di-methylation governs smooth muscle lineage identity and promotes vascular homeostasis by restraining plasticity. *Dev Cell* 56: 2765-2782 e2710, 2021.

32.   **Wanjare M, Kuo F, and Gerecht S**. Derivation and maturation of synthetic and contractile vascular smooth muscle cells from human pluripotent stem cells. *Cardiovasc Res* 97: 321-330, 2013.

33.   **Schneider CA, Rasband WS, and Eliceiri KW**. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9: 671-675, 2012.

34.   **Li Q, Price TP, Sundberg JP, and Uitto J**. Juxta-articular joint-capsule mineralization in CD73 deficient mice: similarities to patients with NT5E mutations. *Cell Cycle* 13: 2609-2615, 2014.

35.   **Martin KA, Rzucidlo EM, Merenick BL, Fingar DC, Brown DJ, Wagner RJ, and Powell RJ**. The mTOR/p70 S6K1 pathway regulates vascular smooth muscle cell differentiation. *Am J Physiol Cell Physiol* 286: C507-517, 2004.

36.   **Marx SO, Jayaraman T, Go LO, and Marks AR**. Rapamycin-FKBP inhibits cell cycle regulators of proliferation in vascular smooth muscle cells. *Circ Res* 76: 412-417, 1995.

37.   **Martin KA, Merenick BL, Ding M, Fetalvero KM, Rzucidlo EM, Kozul CD, Brown DJ, Chiu HY, Shyu M, Drapeau BL, Wagner RJ, and Powell RJ**. Rapamycin promotes vascular smooth muscle cell differentiation through insulin receptor substrate-1/phosphatidylinositol 3-kinase/Akt2 feedback signaling. *J Biol Chem* 282: 36112-36120, 2007.

38.   **Jiao Y, Li G, Li Q, Ali R, Qin L, Li W, Qyang Y, Greif DM, Geirsson A, Humphrey JD, and Tellides G**. mTOR (Mechanistic Target of Rapamycin) Inhibition Decreases Mechanosignaling, Collagen Accumulation, and Stiffening of the Thoracic Aorta in Elastin-Deficient Mice. *Arterioscler Thromb Vasc Biol* 37: 1657-1666, 2017.

39.   **Qi M, Zhang L, Ma Y, Shuai Y, Li L, Luo K, Liu W, and Jin Y**. Autophagy Maintains the Function of Bone Marrow Mesenchymal Stem Cells to Prevent Estrogen Deficiency-Induced Osteoporosis. *Theranostics* 7: 4498-4516, 2017.

40.   **Sun X, Zheng Y, Xie L, Zhou Y, Liu R, Ma Y, Zhao M, and Liu Y**. Autophagy reduces aortic calcification in diabetic mice by reducing matrix vesicle body-mediated IL-1beta release. *Exp Cell Res* 432: 113803, 2023.

41.   **Mizushima N, and Yoshimori T**. How to interpret LC3 immunoblotting. *Autophagy* 3: 542-545, 2007.

42. **Ganesh SK, and Subathra Devi C**. Molecular and therapeutic insights of rapamycin: a
multi-faceted drug from Streptomyces hygroscopicus. *Mol Biol Rep* 50: 3815-3833, 2023.
43. **Dobrin PB, and Mrkvicka R**. Failure of elastin or collagen as possible critical
connective tissue alterations underlying aneurysmal dilatation. *Cardiovasc Surg* 2: 484-488,
1994.
44. **Dobrin PB, Baker WH, and Gley WC**. Elastolytic and collagenolytic studies of arteries.
Implications for the mechanical properties of aneurysms. *Arch Surg* 119: 405-409, 1984.
45. **Rahkonen O, Su M, Hakovirta H, Koskivirta I, Hormuzdi SG, Vuorio E, Bornstein P,
and Penttinen R**. Mice with a deletion in the first intron of the Col1a1 gene develop age-
dependent aortic dissection and rupture. *Circ Res* 94: 83-90, 2004.
46. **Watson SR, Cooper KM, Liu P, Gharraee N, Du L, Han SM, Pena EA, Sutton MA,
Eberth JF, and Lessner SM**. Diet alters age-related remodeling of aortic collagen in mice
susceptible to atherosclerosis. *Am J Physiol Heart Circ Physiol* 320: H52-H65, 2021.
47. **Sabate M**. Sirolimus Versus Paclitaxel: Second Round. *JACC Cardiovasc Interv* 15:
780-782, 2022.
48. **Rouer M, Xu BH, Xuan HJ, Tanaka H, Fujimura N, Glover KJ, Furusho Y, Gerritsen
M, and Dalman RL**. Rapamycin limits the growth of established experimental abdominal aortic
aneurysms. *Eur J Vasc Endovasc Surg* 47: 493-500, 2014.
49. **Zhou B, Li W, Zhao G, Yu B, Ma B, Liu Z, Xie N, Fu Y, Gong Z, Dai R, Zhang X, and
Kong W**. Rapamycin prevents thoracic aortic aneurysm and dissection in mice. *J Vasc Surg* 69:
921-932 e923, 2019.
50. **Zhao Y, Zhao MM, Cai Y, Zheng MF, Sun WL, Zhang SY, Kong W, Gu J, Wang X,
and Xu MJ**. Mammalian target of rapamycin signaling inhibition ameliorates vascular
calcification via Klotho upregulation. *Kidney Int* 88: 711-721, 2015.
51. **Li L, Miano JM, Cserjesi P, and Olson EN**. SM22 alpha, a marker of adult smooth
muscle, is expressed in multiple myogenic lineages during embryogenesis. *Circ Res* 78: 188-
195, 1996.
52. **Krohn JB, Hutcheson JD, Martinez-Martinez E, Irvin WS, Bouten CV, Bertazzo S,
Bendeck MP, and Aikawa E**. Discoidin Domain Receptor-1 Regulates Calcific Extracellular
Vesicle Release in Vascular Smooth Muscle Cell Fibrocalcific Response via Transforming
Growth Factor-beta Signaling. *Arterioscler Thromb Vasc Biol* 36: 525-533, 2016.
53. **Behzadi P, and St Hilaire C**. Metabolites and metabolism in vascular calcification: links
between adenosine signaling and the methionine cycle. *Am J Physiol Heart Circ Physiol* 327:
H1361-H1375, 2024.
54. **Tanemura M, Ohmura Y, Deguchi T, Machida T, Tsukamoto R, Wada H, Kobayashi
S, Marubashi S, Eguchi H, Ito T, Nagano H, Mori M, and Doki Y**. Rapamycin causes
upregulation of autophagy and impairs islets function both in vitro and in vivo. *Am J Transplant*
12: 102-114, 2012.
55. **Duan ZX, Tu C, Liu Q, Li SQ, Li YH, Xie P, and Li ZH**. Adiponectin receptor agonist
AdipoRon attenuates calcification of osteoarthritis chondrocytes by promoting autophagy. *J Cell
Biochem* 121: 3333-3344, 2020.
56. **Jono S, McKee MD, Murry CE, Shioi A, Nishizawa Y, Mori K, Morii H, and Giachelli
CM**. Phosphate regulation of vascular smooth muscle cell calcification. *Circ Res* 87: E10-17,
2000.
57. **Millan A, Lanzer P, and Sorribas V**. The Thermodynamics of Medial Vascular
Calcification. *Front Cell Dev Biol* 9: 633465, 2021.
58. **Frauscher B, Kirsch AH, Schabhuttl C, Schweighofer K, Ketszeri M, Pollheimer M,
Dragun D, Schroder K, Rosenkranz AR, Eller K, and Eller P**. Autophagy Protects From
Uremic Vascular Media Calcification. *Front Immunol* 9: 1866, 2018.
59. **Szwed A, Kim E, and Jacinto E**. Regulation and metabolic functions of mTORC1 and
mTORC2. *Physiol Rev* 101: 1371-1426, 2021.

60.   **Goncharova EA**. mTOR and vascular remodeling in lung diseases: current challenges
and therapeutic prospects. *FASEB J* 27: 1796-1807, 2013.
61.   **Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, Brown M,**
Fitzgerald KJ, and Sabatini DM. Ablation in mice of the mTORC components raptor, rictor, or
mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKCalpha, but not
S6K1. *Dev Cell* 11: 859-871, 2006.
62.   **Schott A, Kluttig A, Mikolajczyk R, Greiser KH, Werdan K, Sedding D, and Nuding**
S. Association of arterial stiffness and heart failure with preserved ejection fraction in the elderly
population - results from the CARLA study. *J Hum Hypertens* 37: 463-471, 2023.
63.   **Writing Group M, Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De**
Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N,
Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A,
McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W,
Sacco R, Sorlie P, Roger VL, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J,
American Heart Association Statistics C, and Stroke Statistics S. Heart disease and stroke
statistics--2010 update: a report from the American Heart Association. *Circulation* 121: e46-
e215, 2010.
64.   **Allison MA, Ho E, Denenberg JO, Langer RD, Newman AB, Fabsitz RR, and Criqui**
MH. Ethnic-specific prevalence of peripheral arterial disease in the United States. *Am J Prev*
Med 32: 328-333, 2007.
65.   **Allison MA, Budoff MJ, Nasir K, Wong ND, Detrano R, Kronmal R, Takasu J, and**
Criqui MH. Ethnic-specific risks for atherosclerotic calcification of the thoracic and abdominal
aorta (from the Multi-Ethnic Study of Atherosclerosis). *Am J Cardiol* 104: 812-817, 2009.
66.   **Chaturvedi N, Bulpitt CJ, Leggetter S, Schiff R, Nihoyannopoulos P, Strain WD,**
Shore AC, and Rajkumar C. Ethnic differences in vascular stiffness and relations to
hypertensive target organ damage. *J Hypertens* 22: 1731-1737, 2004.
67.   **D'Agostino RB, Jr., Burke G, O'Leary D, Rewers M, Selby J, Savage PJ, Saad MF,**
Bergman RN, Howard G, Wagenknecht L, and Haffner SM. Ethnic differences in carotid wall
thickness. The Insulin Resistance Atherosclerosis Study. *Stroke* 27: 1744-1749, 1996.
68.   **Ali H, Zmuda JM, Cvejkus RK, Kershaw EE, Kuipers AL, Oczypok EA, Wheeler V,**
Bunker CH, and Miljkovic I. Wnt Pathway Inhibitor DKK1: A Potential Novel Biomarker for
Adiposity. *J Endocr Soc* 3: 488-495, 2019.
69.   **Bild DE, Detrano R, Peterson D, Guerci A, Liu K, Shahar E, Ouyang P, Jackson S,**
and Saad MF. Ethnic differences in coronary calcification: the Multi-Ethnic Study of
Atherosclerosis (MESA). *Circulation* 111: 1313-1320, 2005.
70.   **Budoff MJ, Nasir K, Mao S, Tseng PH, Chau A, Liu ST, Flores F, and Blumenthal**
RS. Ethnic differences of the presence and severity of coronary atherosclerosis. *Atherosclerosis*
187: 343-350, 2006.
71.   **Malhotra R, Nicholson CJ, Wang D, Bhambhani V, Paniagua S, Slocum C,**
Sigurslid HH, Lino Cardenas CL, Li R, Boerboom SL, Chen YC, Hwang SJ, Yao C,
Ichinose F, Bloch DB, Lindsay ME, Lewis GD, Aragam JR, Hoffmann U, Mitchell GF,
Hamburg NM, Vasan RS, Benjamin EJ, Larson MG, Zapol WM, Cheng S, Roh JD,
O'Donnell CJ, Nguyen C, Levy D, and Ho JE. Matrix Gla Protein Levels Are Associated With
Arterial Stiffness and Incident Heart Failure With Preserved Ejection Fraction. *Arterioscler*
Thromb Vasc Biol 42: e61-e73, 2022.
72.   **Dalmeijer GW, van der Schouw YT, Magdeleyns EJ, Vermeer C, Verschuren WM,**
Boer JM, and Beulens JW. Matrix Gla protein species and risk of cardiovascular events in type
2 diabetic patients. *Diabetes Care* 36: 3766-3771, 2013.
73.   **Chirinos JA**. Matrix Gla Protein, Large Artery Stiffness, and the Risk of Heart Failure
With Preserved Ejection Fraction. *Arterioscler Thromb Vasc Biol* 42: 223-226, 2022.

74.   **Buss SJ, Muenz S, Riffel JH, Malekar P, Hagenmueller M, Weiss CS, Bea F,**
Bekeredjian R, Schinke-Braun M, Izumo S, Katus HA, and Hardt SE. Beneficial effects of
Mammalian target of rapamycin inhibition on left ventricular remodeling after myocardial
infarction. *J Am Coll Cardiol* 54: 2435-2446, 2009.
75.   **Gu J, Hu W, Song ZP, Chen YG, Zhang DD, and Wang CQ**. Rapamycin Inhibits
Cardiac Hypertrophy by Promoting Autophagy via the MEK/ERK/Beclin-1 Pathway. *Front*
Physiol 7: 104, 2016.
76.   **Volkers M, Konstandin MH, Doroudgar S, Toko H, Quijada P, Din S, Joyo A,**
Ornelas L, Samse K, Thuerauf DJ, Gude N, Glembotski CC, and Sussman MA. Mechanistic
target of rapamycin complex 2 protects the heart from ischemic damage. *Circulation* 128: 2132-
2144, 2013.
77.   **Chakraborty AD, Kooiker K, Kobak KA, Cheng Y, Lee CF, Razumova M, Granzier**
HH, Regnier M, Rabinovitch PS, Moussavi-Harami F, and Chiao YA. Late-life Rapamycin
Treatment Enhances Cardiomyocyte Relaxation Kinetics and Reduces Myocardial Stiffness.
bioRxiv 2023.
78.   **Flynn JM, O'Leary MN, Zambataro CA, Academia EC, Presley MP, Garrett BJ,**
Zykovich A, Mooney SD, Strong R, Rosen CJ, Kapahi P, Nelson MD, Kennedy BK, and
Melov S. Late-life rapamycin treatment reverses age-related heart dysfunction. *Aging Cell* 12:
851-862, 2013.
79.   **McMullen JR, Sherwood MC, Tarnavski O, Zhang L, Dorfman AL, Shioi T, and**
Izumo S. Inhibition of mTOR signaling with rapamycin regresses established cardiac
hypertrophy induced by pressure overload. *Circulation* 109: 3050-3055, 2004.
80.   **Baghdadi M, Nespital T, Monzo C, Deelen J, Gronke S, and Partridge L**. Intermittent
rapamycin feeding recapitulates some effects of continuous treatment while maintaining lifespan
extension. *Mol Metab* 81: 101902, 2024.

884
885
886 **FIGURE LEGENDS**
887
Figure 1: Histological comparison of Matrix Gla Protein (Mgp) mice arteries, medial and
intimal calcification in human arteries. **A.** Histological sections of $Mgp^{-/-}$ mouse aorta, human
tibial plaque, and human coronary artery with atherosclerotic plaque, stained for calcification
(Cal) using Von Kossa Staining. Scale Bars: human tissue 1 mm, mouse tissue 0.5 mm. **B.** Dot
Blot Assay to identify Mgp protein in $Mgp^{+/+}$, $Mgp^{+/-}$, and $Mgp^{-/-}$ mice. Results are representative
of n=3 per group. **C.** Representative immunofluorescence images of Mgp protein in aortas from
$Mgp^{+/+}$, $Mgp^{+/-}$ and $Mgp^{-/-}$ mice. Scale Bars: 100 μm. n=6 $Mgp^{+/+}$ vehicle, n=6 $Mgp^{-/-}$ vehicle. P
values were calculated using Mann-Whitney U test. Data are shown as means ± SD.
896
Figure 2: Impact of rapamycin treatment on survival in Matrix Gla Protein (*Mgp*)-deficient
mice. A. Graphic illustration of vehicle (Dimethyl sulfoxide; DMSO) or rapamycin [5 mg/kg]
treatment protocol**. B.** Kaplan-Meier analysis of overall survival in $Mgp^{+/+}$ and $Mgp^{-/-}$ mice treated
with vehicle (DMSO) or rapamycin (5mg/kg/week) for 90 days until natural death. n=7 $Mgp^{-/-}$
rapamycin, n=19 $Mgp^{-/-}$ vehicle, n=8 $Mgp^{+/+}$ rapamycin, n=5 $Mgp^{+/+}$ vehicle. Mantel-Cox test and
Gehan-Breslow-wilcoxon test were used for curve comparison.
903
Figure 3: Effect of rapamycin treatment on medial calcification in *Mgp*-deficient mice. A.
Graphic illustration of vehicle (DMSO) or 5 mg/kg rapamycin treatment for 25 days in $Mgp^{+/+}$ and
$Mgp^{-/-}$ mice. **B.** Representative microCT measurement of axial and cross sections of $Mgp^{-/-}$ mice
aortas treated with vehicle or 5 mg/kg rapamycin, administered either once or three times a
week for 25 days. Scale bar 0.2 mm. n=4 $Mgp^{-/-}$ 1x rapamycin, n=10 $Mgp^{-/-}$ 3x rapamycin, n=12
$Mgp^{-/-}$ vehicle **C.** Representative Von Kossa staining images of $Mgp^{+/+}$ and $Mgp^{-/-}$ mice aortas

treated with vehicle or rapamycin three times a week for 25 days. Scale bar 0.2 mm. n=10 *Mgp*$^{-/-}$
rapamycin, n=10 *Mgp*$^{-/-}$ vehicle, n=10 *Mgp*$^{+/+}$ rapamycin, n=10 *Mgp*$^{+/+}$ vehicle. **D.** Representative
immunofluorescent images of Runx2 on the *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice aortas treated with vehicle
or rapamycin three times a week for 25 days. Scale bar 50 µm. n=10 *Mgp*$^{-/-}$ rapamycin, n=9
Mgp$^{-/-}$ vehicle, n=10 *Mgp*$^{+/+}$ rapamycin, n=10 *Mgp*$^{+/+}$ vehicle. Data are shown as means ± SD. P
values were calculated using Kruskal-Wallis test (Figure B) and Two-way ANOVA test (Figures
C, D).
917
Figure 4: Histological structure of *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice aorta treated with vehicle or
rapamycin. A. Representative Masson's Trichrome staining images of *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice
aortas treated with vehicle (Veh; DMSO) or rapamycin (RAPA; [5 mg/kg]) three times a week for
25 days. Collagen fiber structure in blue. Scale bar 0.2mm. **B.** Representative Picrosirius
staining images of collagen in *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice treated with vehicle or rapamycin three
times a week for 25 days. Scale bar: 50 µm. n=10 *Mgp*$^{-/-}$ rapamycin, n=10 *Mgp*$^{-/-}$ vehicle, n=10
Mgp$^{+/+}$ rapamycin, n=10 *Mgp*$^{+/+}$ vehicle. **C.** Representative Verhoeff-Van Gieson (VVG) staining
images of *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ treated with vehicle or rapamycin three times a week for 25 days.
Scale bar 0.2mm. n=7 *Mgp*$^{-/-}$ rapamycin, n=8 *Mgp*$^{-/-}$ vehicle, n=8 *Mgp*$^{+/+}$ rapamycin, n=9 *Mgp*$^{+/+}$
vehicle. **D.** Representative immunofluorescent images of smooth muscle actin (α-SMA) and
myosin heavy chain gene (Myh11) SMA on *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice aortas treated with vehicle or
rapamycin three times a week. Scale bar 50 µm. n=8 *Mgp*$^{-/-}$ rapamycin, n=9 *Mgp*$^{-/-}$ vehicle, n=10
Mgp$^{+/+}$ rapamycin, n=10 *Mgp*$^{+/+}$ vehicle. **E.** Representative immunofluorescent images of
proliferation markers (ki67) on *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice aortas treated with vehicle or rapamycin
three times a week for 25 days. Scale bar 50 µm. n=10 *Mgp*$^{-/-}$ rapamycin, n=10 *Mgp*$^{-/-}$ vehicle,
n=10 *Mgp*$^{+/+}$ rapamycin, n=10 *Mgp*$^{+/+}$ vehicle. Data are shown as means ± SD. P value were
calculated using Two-way ANOVA test.
935
936
Figure 5: The effect of rapamycin treatment autophagy flux. A. Representative
immunofluorescent images of LC3 on *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice aortas treated with vehicle (Veh;
DMSO) or rapamycin (RAPA; [5 mg/kg]) three times a week for 25 days. LC3 decreased with
rapamycin in *Mgp*$^{-/-}$ mice compared to the *Mgp*$^{+/+}$ mice. Scale bar 50 µm. n=9 *Mgp*$^{-/-}$ rapamycin,
n=10 *Mgp*$^{-/-}$ vehicle, n=10 *Mgp*$^{+/+}$ rapamycin, n=10 *Mgp*$^{+/+}$ vehicle. **B.** Smooth muscle cells from
Mgp$^{+/+}$ and *Mgp*$^{-/-}$ mice were treated with vehicle (DMSO) or rapamycin (200 µM) for 14 hours,
followed by the addition of bafilomycin (BafA, 100 µM), an autophagy inhibitor, for an additional
24 hours. Levels of LC3 (LC3-I and LC3-II) were quantified by western blot. n=6 *Mgp*$^{-/-}$
rapamycin, n=6 *Mgp*$^{-/-}$ vehicle, n=6 *Mgp*$^{+/+}$ rapamycin, n=6 *Mgp*$^{+/+}$ vehicle. Data are shown as
means ± SD. P values were calculated using Two-way ANOVA test (Figure A) and Kruskal-
Wallis test (Figure B).
948
Figure 6: Survival probability of Matrix Gla Protein (MGP)-deficient mice with Rictor and
Raptor knockouts in vascular smooth muscle cells (SMC). A. *Mgp*$^{-/-}$ using Cre recombinase
that targets the gene encoding Rictor in SMC, a protein involved in mTORC2 signaling
pathways. *Mgp*$^{-/-}$;*Rictor*$^{SMC-/-}$ (n=5) and *Mgp*$^{-/-}$;*Rictor*$^{SMC+/+}$ (n=7). **B.** *Mgp*$^{-/-}$ using Cre recombinase
that target the gene encoding Raptor in SMC a protein involved in mTORC1 signaling pathways.
Mgp$^{-/-}$;*Raptor*$^{SMC+/+}$ (n=6), *Mgp*$^{-/-}$;*Raptor*$^{SMC-/-}$ (n=7). Data are shown as means ± SD. Mantel-Cox
test and Gehan-Breslow-wilcoxon test were used for curve comparison.
956
957
958
959

A

Mouse *Mgp⁻ᐟ⁻* Aorta	Human Tibial Artery	Human Atherosclerotic Plaque

B

C

A

B

A

B

Mgp⁻ᐟ⁻ Veh

Mgp⁻ᐟ⁻ RAPA

Lifespan Average
32 days

Lifespan Average
57 days

**Elastic Fiber
Breakage**

Collagen

α-SMA

Calcification

ECM

Mgp

↓ **Elastic Fiber
Breakage**

↓ **Collagen**

↑ **α-SMA**

↔ **Calcification
Volume**

↓ **Mineral Density**

ECM

Mgp

★ Mineralization ⌇⌇ Elastin 〰〰 Collagen ◇ Smooth muscle cell with more a-SMA ◆ Smooth muscle cell with less a-SMA