1 **Rapamycin Reduces Arterial Mineral Density and Promotes Beneficial Vascular**
2 **Remodeling in a Murine Model of Severe Medial Arterial Calcification.**
3

4 Parya Behzadi[1], Rolando A. Cuevas[1], Alex Crane[1], Andrew A Wendling[1], Claire C. Chu[1], William
5 J Moorhead III[1], Ryan Wong[1], Mark Brown[1], Joshua Tamakloe[1], Swathi Suresh[1], Payam Salehi[2],
6 Iris Z. Jaffe[3], Allison L. Kuipers[4], Lyudmila Lukashova[5], Konstantinos Verdelis[5], and Cynthia St.
7 Hilaire[1,6,7]
8

9 [1] Department of Medicine, Division of Cardiology, and the Pittsburgh Heart, Lung, Blood and
10 Vascular Medicine Institute, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
11

12 [2] CardioVascular Center, Vascular Surgery, Tufts Medical Center, 800 Washington Street,
13 Boston, MA, 02111-1800, USA
14

15 [3] Molecular Cardiology Research Institute, Tufts Medical Center, 800 Washington Street,
16 Boston, MA, 02111-1800, USA
17

18 [4] Department of Epidemiology, School of Public Health, University of Pittsburgh, Pittsburgh,
19 Pennsylvania, USA
20

21 [5] Departments of Endodontics and Oral Biology, School of Dental Medicine, University of
22 Pittsburgh, Pittsburgh, PA, USA
23

24 [6] Department of Bioengineering, University of Pittsburgh, Pittsburgh, Pennsylvania, USA
25

26 [7] Department of Cardiothoracic Surgery, University of Pittsburgh, Pittsburgh, Pennsylvania, USA
27

28 **ABSTRACT**: 251 words
29

30 **MANUSCRIPT:**
31

32 **KEYWORDS:** Medial Arterial Calcification, Rapamycin, Matrix GLA protein, Collagen, Elastin
33

ABBREVIATIONS
Adenosine monophosphate   AMP
Adenosine triphosphate   ATP
Alkaline phosphate   TNAP
Arterial calcification due to deficiency of CD73   ACDC
Ecto-5'-nucleotidase   NT5E
Induced pluripotent stem cells   iPSCs
Matrix GLA protein   MGP
Medial arterial calcification   MAC
mTORC complex 1   mTORC1
mTORC complex 2   mTORC2
Peripheral artery disease   PAD
Smooth muscle α-actin   SMA
Smooth muscle cells   SMCs
Myosin heavy chain 11   MYH11
Runt-related transcription factor 2   RUNX2

50
ABSTRACT

Peripheral artery disease (PAD) is the narrowing of the arteries that carry blood to the lower extremities. PAD has been traditionally associated with atherosclerosis. However, recent studies have found that medial arterial calcification (MAC) is the primary cause of chronic limb ischemia below the knee. MAC involves calcification of the elastic fibers surrounding smooth muscle cells (SMCs) in arteries. Matrix GLA protein (MGP) inhibits vascular calcification by binding circulating calcium and preventing hydroxyapatite crystal deposition, while also modulating osteogenic signaling by blocking BMP-2 activation of RUNX2. $Mgp^{-/-}$ mice develop severe MAC and die around 8 weeks after birth due to aortic rupture or heart failure. We previously discovered a rare genetic disease Arterial Calcification due to Deficiency in CD73 (ACDC) in which patients present with extensive MAC in their lower extremity arteries. Using a patient-specific induced pluripotent stem cell model we found that rapamycin inhibited calcification. Here we investigated whether rapamycin could reduce MAC in vivo using the $Mgp^{-/-}$ murine model. $Mgp^{+/+}$ and $Mgp^{-/-}$ mice received 5mg/kg rapamycin or vehicle. Calcification content was assessed via microCT, and vascular morphology and extracellular matrix content assessed histologically. Immunostaining and western blot analysis were used to examine SMC phenotype and extracellular matrix content. Rapamycin prolonged $Mgp^{-/-}$ mice lifespan, decreased mineral density in the arteries, maintained SMC contractile phenotype, and improved vessel structure, however, calcification volume was unchanged. $Mgp^{-/-}$ mice with SMC-specific deletion of Raptor or Rictor, did not recapitulate treatment with rapamycin. These findings suggest rapamycin promotes beneficial vascular remodeling in vessels with MAC.

INTRODUCTION

Peripheral artery disease (PAD) is the narrowing of blood vessels in the lower extremities due to inward remodeling or thrombotic occlusion, both of which cause chronic limb ischemia and often result in amputation of the lower leg (1). Traditionally atherosclerosis was the assumed cause of PAD however recent studies now show that medial arterial calcification (MAC) promotes inward remodeling, and stiffness related to medial dysplasia induces thrombosis (2-4). MAC is characterized by the progressive buildup of calcium and phosphate within the arterial walls and is typically not associated with lipid deposition, fibrous cap formation, or intimal hyperplasia (5). These structural features distinguish MAC from atherosclerosis and indicate that MAC is a distinct pathology driving adverse outcomes in PAD. Currently, there are no specific medical therapies that target the pathogenesis of PAD or MAC. Finding therapies that can prevent, stop, or even reverse MAC in PAD could greatly enhance the current standard of care.

Vascular calcification stems from the nucleation of calcium and phosphate into hydroxyapatite crystals and the release of pro-mineralizing matrix vesicles from osteogenic-like arterial smooth muscle cells (SMCs) that accumulate along the elastic lamina (6). Phosphate is a byproduct of the breakdown of extracellular adenosine triphosphate (ATP) to adenosine, generating inorganic phosphate at several points (7, 8). There are a number of genetic diseases that present with MAC which harbor inactivating mutations in the genes related to extracellular ATP metabolism (5). Patients with Arterial Calcification due to Deficiency of CD73 (ACDC, also known as CALJA, OMIM # 211800) harbor inactivating mutations in the *NT5E* gene, which encodes for the CD73 enzyme that metabolizes extracellular AMP to adenosine and inorganic phosphate. Key signatures of this disease are calcification nodules in the small joint capsules and MAC in the lower-extremity arteries that initiate along the elastic lamina (4, 9, 10). We previously found that a lack of CD73-mediated adenosine enhanced the expression and activity of tissue-nonspecific alkaline phosphatase (TNAP), a key enzyme that promotes calcification (11).

99
Matrix GLA protein (MGP) is a vitamin-K2-dependent protein with an unusual gamma-
carboxylation of five glutamate residues that enhance its affinity for calcium, preventing it from
mineralizing (12). MGP also inhibits calcification propagation by binding to hydroxyapatite crystals
and stimulating their uptake by local phagocytosing macrophages (13). MGP also modulates
osteogenic signaling by binding to BMP-2, preventing it from activating receptors and upregulating
RUNX2, the key regulator of osteoblast differentiation and maturation (14, 15). Mice that lack
MGP develop to term but die prematurely within approximately two months of age due to
extensive MAC, which leads to aortic rupture or heart failure (16). Aortas of MGP-deficient mice
exhibit increased collagen accumulation and elastic fiber fragmentation within the medial layer of
the arteries where calcification is also localized. This is phenotypically similar to MAC observed
in human patients with PAD MAC and ACDC patients (4).
111
The drug rapamycin inhibits the mammalian target of rapamycin (mTOR), a protein kinase broadly
expressed throughout the body, which controls many key processes such as energy balance,
autophagy, and proliferation (17-20). Rapamycin protects against the calcification of vascular cells
in in vitro and ex vivo models, as well as in in vivo models of chronic kidney disease-induced MAC
(21-23). We previously found rapamycin prevented calcification in in vitro and in vivo disease
models of ACDC patient-specific induced pluripotent stem cells (iPSCs) (11).
118
119 While CD73-deficient humans develop extensive MAC in their lower extremity large vessels,
120 CD73-deficient mice do not phenocopy humans and do not exhibit calcification in their vasculature
121 (24, 25). As the MAC observed in MGP-deficient mice mirrors that seen in ACDC patients, we
122 sought to investigate whether rapamycin prevented calcification in this genetic mouse model of
123 MAC.
124
MATERIALS AND METHODS

Availability of Materials
We abide by the NIH Grants Policy on Sharing of Unique Research Resources, including the NIH
Policy on Sharing of Model Organisms for Biomedical Research (2004), NIH Grants Policy
Statement (2003), and Sharing of Biomedical Research Resources: Principles and Guidelines for
Recipients of NIH Grants and Contracts (1999), and the Bayh-Dole Act and the Technology
Transfer Commercialization Act of 2000. Materials generated in our laboratory are made available
for non-commercial research per established University of Pittsburgh Office of Research IRB and
MTA protocols.
134
135 *Institutional Review Board Statement*
136 De-identified human tissues were obtained, with informed consent from subjects, who were
137 enrolled in studies approved by the University of Pittsburgh or Tufts Medical Center institutional
138 review boards, per the Declaration of Helsinki.
139
Animals and cell line generation
Animal use was approved by the Institutional Animal Care and Use Committee at the University
of Pittsburgh. MGP heterozygous mutant mice from a C57BL/6J background (strain# 023811
Jackson Laboratory, Bar Harbor, ME) were bred to produce +/+ and -/- littermate controls. Smooth
muscle lineage-specific *Raptor* ($Mgp^{-/-}$; $Raptor^{SMC-/-}$) and *Rictor* ($Mgp^{-/-}$; $Rictor^{SMC-/-}$) knockout mice
were produced by breeding $Mgp^{+/-}$ with Myh11-Cre-eGFP mice (strain# 007742 Jackson
Laboratory, Bar Harbor, ME)(26), with Raptor-LoxP (strain# 013188 Jackson Laboratory, Bar

Harbor, ME)(27), or *Rictor*-LoxP (strain# 020649 Jackson Laboratory, Bar Harbor, ME)(28).
Genotypes were determined from tail snips that were incubated in DirectPCR Lysis Reagent
(Viagen Biotech) according to manufacturer instructions. The lysate was amplified using OneTaq
MasterMix (New England Biolabs) according to manufacturer instructions with primers suggested
by Jackson Laboratories. The PCR reaction was run on an ethidium bromide agarose gel. SMC
were obtained from three- to five-week-old male and female *Mgp*$^{+/+}$ and *Mgp*$^{-/-}$ mice.

Cell culture
Mgp$^{+/+}$ and *Mgp*$^{-/-}$ aortic SMC lines from male and female murine aorta were cultured in Dulbecco's
modified Eagle's medium (DMEM; Gibco, Waltham, MA) supplemented with 20% FBS (FBS; R&D
Systems, Minneapolis, MN) and 100 U/mL penicillin-streptomycin (P/S; Gibco, Waltham, MA).
Growth media was changed every three days, and cells were split 1:2 when confluent. Post-
expansion, 25,000–50,000 cells/cm² were plated in DMEM supplemented with 10% FBS and 100
U/mL penicillin-streptomycin, and grown to confluence. Before all experiments/treatments, cells
were serum-starved in DMEM with 0.5% FBS for 48 h. Incubation under reduced serum conditions
has been proven to be a beneficial technique for studying smooth muscle cell remodeling and
contractility in vitro (29, 30). Cells were treated with rapamycin (Novus Biologicals) at
concentrations of 200 nM or the same volume of DMSO as vehicle control for 14h. For autophagy
flux assessment, cells were exposed to either bafilomycin A1 100 nM (Sigma-Aldrich) or
rapamycin 200 nM for a duration of 24h following treatment intervention. All chemicals were
dissolved in DMSO and administered in equal volume as vehicle control.
167
Tissue Extraction
Mice were sacrificed by asphyxiation with carbon dioxide, followed by perforation of the
diaphragm. Tissue extraction began with dissection through the abdominal wall and then
perfusion of the heart with 50 mL of DPBS supplemented with 1:100 Amphotericin B (15290026,
Gibco) and 1:200 Gentamicin (15710064, Gibco). The aortas were maintained in this solution
while the adventitia was removed in a petri dish. Tissues were then fixed in 4% paraformaldehyde
in PBS for 2 hours and then placed in PBS. Tissues were brought to microCT and then embedded
in paraffin, or embedded in paraffin directly.
176
In Vivo Rapamycin Injections
Male and female *Mgp*$^{-/-}$ mice received intraperitoneal injections of either vehicle or 5 mg/kg
rapamycin dissolved in 0.3 mg/ml peanut oil once a week beginning at 10 days of age. This
process continued until the natural death of all mice in each group. To assess the volume and
density of calcification, male and female *Mgp*$^{-/-}$ mice received intraperitoneal injections of either
vehicle, 1x/week rapamycin or 3x/week rapamycin from 10 to 25 days of age.
183
Micro-Computed Tomography
A Scanco 30μCT microCT scanner (Scanco Medical, Bassersssdorf, Switzerland) with a 45 kVp
beam energy (1,000 ms exposure) was used to assess the volume and density of calcification in
mouse aorta ex vivo, as previously described (11, 31). No filtering was used, allowing for visibility
of lower CT densities. Images were then analyzed for relative calcified mass within the aorta
(expressed as bone volume fraction) using the Scanco 3D Morphometry and Densitometry
Analysis software (Scanco Medical) after binarization of the images with a global threshold within
a user-defined aorta region of interest.
192
Western Blot Analysis

Cells were lysed in 1% CHAPS hydrate, 150 mmol/L sodium chloride, 25 mmol/L HEPES buffer
supplemented with 1× protease and phosphatase inhibitor (Sigma-Aldrich). Cells were scraped
into microcentrifuge tubes, vortexed for 5 minutes, freeze/thawed for 5 to 8 cycles, then
centrifuged at 12 000×g for 10 minutes at 4°C. Supernatant protein concentration was determined
using Pierce BCA protein assay kit (Thermo Fisher, Waltham, MA). Ten micrograms of protein
were used in the preparation of lysate with 1 × Pierce LDS sample buffer nonreducing (Thermo
Fisher, Waltham, MA) and 1 × NuPAGE sample reducing agent (Novex, Waltham, MA). Lysates
were denatured at 95°C for 15 min and then electrophoresed on 4%–20% TGX stain-free
polyacrylamide gel (Bio-Rad, Hercules, CA) in 1 × Tris/Glycine/SDS buffer (Bio-Rad, Hercules,
CA) at 120 V for 50 min. Protein was transferred onto a 0.2 μm nitrocellulose membrane in
prepared 1 × Towbin buffer with ethanol (EtOH) at 1 A and 25 V for 30 min using the Trans-Blot
Turbo Transfer System (Bio-Rad, Hercules, CA). Membranes were blocked in 1:1 Odyssey
blocking buffer (Li-COR, Lincoln, NE) and PBS for 1 h at room temperature, followed by primary
antibody incubation in 1:1 Odyssey blocking buffer and PBS plus 0.1% Tween 20 (PBS-T) at 4°C
overnight. Membranes were washed in PBS-T three times for 5 min, then incubated in secondary
antibody at room temperature for 1 h. Membranes belonging to the same experimental set were
imaged simultaneously on an Odyssey CLx (LI-COR, Lincoln, NE), and band intensity
quantification was performed with Image Studio (Version 5.2, LI-COR, Lincoln, NE) software.
Individual bands were normalized to α-tubulin, and each treatment group's fold change was
compared with each gel's vehicle control lanes. For sequential antibody incubations, membranes
were stripped in 1 × NewBlot Nitro Western Blot Stripping Buffer (LI-COR, Lincoln, NE) for 10
min, followed by three washes in PBS.
216
Verhoeff Van Gieson, Von Kossa, Masson's Trichrome and Picrosirius Red Staining
Mice aortic tissues were removed from mice through the abdominal wall under anesthesia
conditions. Then aortic tissue was fixed with 4% paraformaldehyde and embedded in paraffin.
The embedded specimens were transversely sectioned at 10 μm on a microtome cryostat
(Microm HM 325). Slides with the adhered paraffin aortic sections were warmed to 65°C for 1 h
and then deparaffinized through xylene, rehydrated with serial incubation in graded alcohol baths,
and stained with Verhoeff–van Gieson for elastic fiber visualization (Polysciences, 25089-1), Von
Kossa for calcification (Polysciences, 24633-1), Masson's for collagen fibers (Polysciences,
25088-1) and Picrosirius Red staining for collagen quantification according to manufacturer's
instructions. Verhoeff Van Gieson, Von Kossa, Masson's Trichrome images were captured using
PreciPoint scanner. Collagen staining was assessed by measuring the intensity of collagen in the
vessel wall normalized to scale bar. Picrosirius Red staining images were captured collagen with
the Olympus VS200 Slide Scanner. The percentage of medial thickness of the medial layer was
measured using ImageJ software (Java 8), where the medial thickness was determined by
measuring the distance between the inner and outer elastic lamina at four sites per vessel. The
percentage of calcification in the medial layer was measured using ImageJ software (Java 8)
where the area of calcification was normalized to total medial area. The tortuosity of elastic fibers
was measured by selecting three representative areas with the whole vessel. ImageJ was used
to trace and measure the length of five elastic fibers and shown as a ration over the vessel length.
The number of breakages in the elastic lamina was assessed by counting the breaks throughout
the entire aorta. All length measurements were normalized to the scale bar units.
238
Immunofluorescence
Slides with the adhered paraffin aortic sections were warmed to 65°C for 1 h and then
deparaffinized through xylene, rehydrated with serial incubation in graded alcohol baths. Slides
were boiled in citric acid-based antigen retrieval solution (H-3300, Vector Labs) for 20 minutes.

Slides were cooled in the unmasking solution for 1 hour, washed in 1x PBS and placed in blocking
buffer (500 mL PBS 0.3g Fish Skin Gelatin) for 1 hour. Tissues were incubated overnight at 4°C
with antibodies for ACTA2 (Abcam, ab5694, 1:50), Ki67 (Abcam, ab15580, 1:250), LC3 (cell
signaling,124714, 1:50), MGP (10734-I-AP, 1:25), MYH11 (Invitrogen, MA5-4284, 1:50) , RUNX2
(ABclonal, A2851, 1:100). All signals were optimized to IgG (Vector, 31235, 1:50). Cells were
washed 3× for 5 minutes each with PBS, 0.1% TWEEN 20, then once with PBS, and then
incubated with secondary antibody (A11006, A11012, Thermo Fisher) for 1 hour. Cells were
washed 3× for 5 minutes each with PBS, 0.1% TWEEN 20, then once with PBS, then mounted
with Fluoroshield Mounting Medium with DAPI (Ab104139, Abcam). Slides were imaged within
24 hours of mounting. Cells were imaged within 24 hours of mounting. In ImageJ, each image
was split into blue, red, and green channels, and the pixel intensity of the red and blue channels
was measured (32). The measured intensity of a given stain was normalized to DAPI intensity of
that same image. We obtained the von Kossa staining images using PreciPoint scanner and the
immunofluorescent image using a Nikon NI-E I microscope.
257
Statistics
Statistical analysis was performed with GraphPad Prism 9.2 (GraphPad Software, Inc) and data
shown are mean±SD. Statistical significance between groups was assessed using the Kruskal
Wallis test or by Two-way ANOVA according to the data set. Statistical analysis used, exact n
values, biological replicates, and P values are stated within each figure legend. A p-value equal
to or less than 0.05 will be considered statistically significant.
264
265 **RESULTS**
266
MAC in MGP-deficient mice phenocopies human MAC.
The main characteristic of MAC is calcification in the media along the elastic lamina. Von Kossa
staining for calcification illustrates that *Mgp$^{-/-}$* mouse aortas have extensive mineralization in the
medial layer, which phenocopies the mineralization seen in MAC of the human tibial arteries and
is distinct from neointimal calcification found in the necrotic core of coronary artery with
atherosclerotic plaque (Figure 1A). MGP is a potent inhibitor of soft tissue calcification. *Mgp$^{-/-}$*
mice do not produce functional MGP protein (Figure 1B, 1C) and, as a result, develop extensive
MAC pathology in large vessels (16).
275
Rapamycin reduces mineral density in *Mgp$^{-/-}$* mice.
Humans with inactivating mutations in *NT5E* develop extensive MAC in their lower extremity
arteries (9). It has been shown that the *Nt5e*-knockout mice model shows altered renal function,
diminished control of the glomerular arteriolar tone, elevated atherogenesis, elevated thrombotic
occlusion, and elevated hypoxia-induced vascular leakage (24), these mice do not mimic the
human vascular calcification phenotype (24, 25, 33). Jin et al. circumvented this issue using a
patient-specific in vitro and in vivo disease modeling system using iPSC technology. They
demonstrated that patient-specific ACDC-iPSCs developed extensive calcification in the in vivo
teratoma model relative to control patient iPSCs. Furthermore, they found that treating mice
bearing ACDC-iPSC teratomas with rapamycin reduced calcification (11).
286
As CD73-deficient mice do not sufficiently recapitulate the MAC observed in ACDC patients, we
used the *Mgp$^{-/-}$* mice as an in vivo model of MAC to test the hypothesis that rapamycin could
reduce vascular calcification in vivo. *Mgp$^{+/+}$* and *Mgp$^{-/-}$* mice were administered 5 mg/kg rapamycin
of the same volume of vehicle (DMSO) three times per week starting at 10 days old until natural
death (Figure 2A). We discovered that rapamycin treatment extended the lifespan of *Mgp$^{-/-}$* mice

(T$_{50}$ 61 days) compared to the *Mgp*$^{-/-}$ mice treated with the vehicle (T$_{50}$ 35 days), which start to die
as early as day 20 (Figure 2B). Both male (n=11) and female (n=8) were used however no sex
differences were observed between *Mgp*$^{-/-}$ mice that had received either vehicle or rapamycin
treatment.
296
We next investigated whether calcification content was altered in mice receiving rapamycin. Mice
received vehicle, one dose of 5 mg/kg rapamycin per week, or 5 mg/kg rapamycin three times a
week. Mice were sacrificed at day 25 and mineral was assessed via microCT (Figure 3A). We
selected this time point because we observed that the mice began dying at 25 days (Figure 2B).
Whole aorta microCT analysis found no differences in the total calcification volume under either
treatment regime; however, mice receiving a rapamycin 3x per week exhibited decreased mineral
density compared to the control group (Figure 3B). Based on this data, we proceeded with
3x/week for the rest our studies. Von Kossa stain showed that the aortas of *Mgp*$^{+/+}$ mice exhibit
no mineralization, while both vehicle- and rapamycin-treated *Mgp*$^{-/-}$ mice showed extensive
calcification compared to *Mgp*$^{+/+}$ mice, there were no discernible differences in the calcified area
or the medial thickness vehicle or rapamycin groups of *Mgp*$^{-/-}$ (Figure 3C). RUNX2 is a key
transcription factor promoting vascular cell osteogenic differentiation (6). Immunofluorescent
staining revealed that while *Mgp*$^{-/-}$ mice showed elevated levels of RUNX2 compared to *Mgp*$^{+/+}$
mice, however rapamycin treatment did not reduce RUNX2 expression in *Mgp*$^{-/-}$ compared to *Mgp*$^{-}$
$^{/-}$ treated with vehicle (Figure 3D). These findings suggest that while rapamycin doubles the
lifespan of *Mgp*$^{-/-}$ mice, it does not lead to decreased calcification volume, but does promote les-
dense mineral formation in *Mgp*$^{-/-}$ mice.
314
Rapamycin alters SMC phenotype and ECM content in vivo
The drug rapamycin has a broad spectrum of effects on SMCs. Rapamycin inhibits SMC migration
and proliferation by inducing the cyclin-dependent kinase inhibitors p27^{kip} and p21^{cip} and
promotes G1-S cell cycle arrest (34, 35). Rapamycin also induces SMC differentiation by inhibiting
the mTOR-target S6K1 via AKT activation, promoting SMC contractile phenotype by enhancing
SMA and MYH11 protein expression (36). Further, rapamycin was shown to protect against
calcification of vascular cells in in vitro and ex vivo models (11, 21). While rapamycin did not
reduce calcification volume in *Mgp*$^{-/-}$ vessels we hypothesized that it could perhaps prevent aortic
rupture by maintaining SMC contractile phenotype or stabilize the ECM.
324
Masson's trichrome staining shows that *Mgp*$^{-/-}$ vehicle vessels are highly remodeled, with acellular
areas exhibiting observable increased collagen staining (Figure 4A). Picrosirius staining enabled
quantification of collagen content and shows that, while not at the levels seen in *Mgp*$^{+/+}$ vessels,
rapamycin increased the collagen content in the vessels of *Mgp*$^{-/-}$ mice (Figure 4B). This suggests
that rapamycin helps restore or stabilize the collagen within the ECM, which is likely offering a
protective effect by strengthening tissue structure to improve resilience against rupture. Verhoeff-
Van Gieson (VVG) staining revealed that the aortas of *Mgp*$^{-/-}$ mice exhibited less robust staining
for elastin, and less tortuous fibers, in *Mgp*$^{-/-}$ compared to *Mgp*$^{+/+}$. While rapamycin did not alter
elastic fiber tortuosity in *Mgp*$^{-/-}$ mice compared to vehicle controls, the number of elastin breakages
was higher in *Mgp*$^{-/-}$ mice compared to *Mgp*$^{+/+}$ in both groups, and rapamycin treatment reduced
the number of breakages in *Mgp*$^{-/-}$ mice compared to vehicle (Figure 4C).
336
Immunofluorescent staining was used to quantify the SMC contractility markers smooth muscle
α-actin (SMA) and myosin heavy chain 11 (MYH11) (37, 38). In *Mgp*$^{+/+}$ we observed no differences
in SMA levels between vehicle and rapamycin treatment, nor between *Mgp*$^{+/+}$ the *Mgp*$^{-/-}$ mice
treated with vehicle. However, rapamycin treatment slightly, but significantly, increased SMA

levels in *Mgp*[-/-] mice compared to vehicle alone. No differences in MYH11 levels were observed
(Figure 4D). These results suggest a slight increase in the contractile phenotype of SMCs of *Mgp*[-
/-] mice treated with, but not to levels see in *Mgp*[+/+] mice. As the medial layer is remodeled in the
Mgp[-/-] mice, we also measured the mitotic marker Ki67 and found that at baseline, Ki67 levels
were significantly elevated in *Mgp*[-/-] aorta while nearly undetectable in the *Mgp*[+/+]. However,
rapamycin treatment did not alter Ki67 levels in either of the two genotypes (Figure 4E). Together,
these data suggest that rapamycin does not significantly alter the medial remodeling or SMC
phenotype in *Mgp*[-/-] mice aorta.
349
Rapamycin lowered LC3 staining in *Mgp*[-/-] aorta but there were no observable changes in
autophagy flux.
Autophagy is a ubiquitous process and contributes to bone development and the differentiation
of osteoblasts and osteoclasts (39) One study found that autophagy reduces vascular calcification
by limiting the release of pro-calcific matrix vesicles (40). Another found that in vitro, autophagy
stimulation with valproic acid protects against phosphate-induced calcification, while its inhibition
increases extracellular vesicle release in SMCs (23). LC3 is a protein that functions at the initiation
of autophagosome formation and serves as an autophagy marker. While there were no
differences between the LC3 signal in *Mgp*[+/+] and *Mgp*[-/-] at baseline, we observed that rapamycin
treatment significantly lowered LC3 puncta in *Mgp*[-/-] compared to *Mgp*[+/+] mice (Figure 5A).
360
Quantifying LC3 puncta hints at diminished autophagy, but immunofluorescent staining of fixed
tissues cannot assess autophagy flux because it provides only a single snapshot of the
autophagic process, which does not reflect changes in the dynamic turnover of autophagosomes
or the lysosomal clearance of LC3-positive autophagic vesicles. LC3-I is cleaved to LC3-II when
it is incorporated into autophagosomes, but this process is dynamic, and LC3-II can also be
degraded during autophagy. The comparison of LC3-II to LC3-I in the presence of an autophagy
inhibitor like bafilomycin enables autophagy flux observation (41). We isolated SMCs from *Mgp*[+/+]
and *Mgp*[-/-] mice to evaluate autophagy flux. SMC were treated with vehicle (DMSO) or 200 nM
rapamycin for 14h and then with vehicle or 100 uM or the late-phase autophagy inhibitor
bafilomycin for 24h. Rapamycin alone did not increase LC3-II levels compared to vehicle controls
in either the *Mgp*[+/+] and *Mgp*[-/-] SMC. Bafilomycin, an autophagy inhibitor, increased LC3-II levels
in both *Mgp*[+/+] and *Mgp*[-/-] SMCs compared to their cognate vehicle controls, however the addition
of rapamycin did not further alter LC3-II levels compared to bafilomycin (Figure 5B, left graph).
Comparing LC3-II to LC3-I showed identical trends (Figure 5B, right graph), suggesting that global
LC3 levels and flux are similar in *Mgp*[+/+] and *Mgp*[-/-] SMCs.
376
Genetic deletion of mTOR signaling in the SMCs does not replicate rapamycin treatment
***Mgp*[-/-] mice.**
Figure 2 shows that rapamycin treatment doubles the lifespan of *Mgp*[-/-] mice compared to vehicle
alone. However, our subsequent inquiries revealed that rapamycin did not decrease the volume
of calcification, as was seen in in vivo iPSC teratoma models (11), while it did promote less dense
mineral formation, and beneficial SMC phenotype and EMC remodeling, albeit not to levels seen
in the *Mgp*[+/+] animals. The mammalian target of rapamycin, mTOR, can form two different protein
complexes known as mTORC1, bound to Raptor, and mTORC2, bound to Rictor. The former is
acutely sensitive to rapamycin administration, while the latter requires chronic rapamycin dosing
for inhibition (19). To investigate whether rapamycin's effects on the *Mgp*[-/-] mouse are acting
through either of these complexes, we generated *Mgp*[-/-] mice with SMC-specific deletion of the
mTOR complex proteins Raptor and Rictor (26-28). *Mgp*[-/-]*;Myh11Cre-GFP;Raptor*[f/f] mice (*Mgp*[-/-]
;Raptor[SMC-/-]) were bred to examine the effects of a nonfunctional mTORC1 complex, while *Mgp*[-

$^{-/-}$;*Myh11Cre-GFP*;*Rictor$^{f/f}$* (*Mgp$^{-/-}$*;*Rictor$^{SMC-/-}$*) mice knockout was bred to examine a nonfunctional
mTORC2 complex. Fully differentiated SMCs expressing *Myh11* will constitutively express Cre
recombinase to inactive *Raptor* and *Rictor* gene expression specifically in SMCs.
393
Similar to *Mgp$^{-/-}$* mice treated with vehicle in Figure 2, Figure 6 shows that only 50% of the *Mgp$^{-/-}$*
;Rictor$^{SMC-/-}$ and *Mgp$^{-/-}$*;*Rictor$^{SMC+/+}$* are alive at 40 days, indicating that rapamycin acting via
mTORC2 does not recapitulate the effects of rapamycin on increasing *Mgp$^{-/-}$* mice lifespan. The
50% survival rate of *Mgp$^{-/-}$*;*Raptor$^{SMC+/+}$* mice was similar to *Mgp$^{-/-}$* treated with vehicle, *Mgp$^{-/-}$*
;Rictor$^{SMC-/-}$ and *Mgp$^{-/-}$*;*Rictor$^{SMC+/+}$*, however the 50% survival rate of MGP$^{-/-}$;Raptor$^{SMC-/-}$ mice
dropped to 22 days. As the Cre strain being used is a non-inducible system, this data in the MGP-
$^{/-}$;Raptor$^{SMC-/-}$ mice suggest that mTORC1 is critical to the growth and development of the
vasculature. Together, these in vivo data suggest that the life-extending effects of rapamycin in
the *Mgp$^{-/-}$* mouse are not acting through the mTORC2 complex. Additional experiments, such as
a tamoxifen inducible Cre system, would need to be used to fully address the contribution of
mTORC1 to mirror effects seen by rapamycin injection.
405
DISCUSSION
In this study, we demonstrated that rapamycin treatment increased the lifespan in *Mgp$^{-/-}$* mice, but
the effects of rapamycin do not reduce MAC. In mice, rapamycin has been shown to increase
maximum lifespan and delay the onset of cancer as hyperactivity of the mTOR pathway,
frequently found in cancers, leads to the expression and activation of many oncogenes like PI3K,
Akt, and eIF4E. Hence, rapamycin can slow the proliferation of tumor cells by causing cell arrest,
promoting apoptosis, and blocking angiogenesis in tumor growths (42). Our data show that the
increase in lifespan observed in *Mgp$^{-/-}$* mice following rapamycin treatment is associated with
improvements in SMC contractile protein expression and collagen content in the vessel wall, but
not with enhanced autophagy flux.
416
Collagen, the most abundant protein in the ECM, plays a crucial role in maintaining the structural
integrity of the blood vessels. Collagen degradation leads to increased vessel distensibility and
the risk of rupture (43, 44). Aging mice homozygous for a deletion in the gene coding for collagen
alpha 1(Col1a1) are at a significantly increased risk of aortic dissection and rupture (45). The
increase in collagen content observed in *Mgp$^{-/-}$* mice treated with rapamycin compared to those
receiving vehicle treatment. The increase in collagen content following rapamycin treatment and
the increase in SMC contractile protein SMA may contribute to the prolonged lifespan of *Mgp$^{-/-}$*
mice by improving both the structural stability and functional resilience of the vascular walls.
425
While rapamycin is known to inhibit SMC proliferation and is used as such in drug-eluting stents,
there is limited research on the effects of rapamycin on vascular morphology in the presence of
calcification (35, 46). It was previously shown that treatment with rapamycin limits the progression
of abdominal and thoracic aortic aneurysms in mice and preserves elastic lamina integrity (47,
48). In a hyperphosphatemic rat model, rapamycin treatment inhibited aortic calcium deposition
when ectopic calcification was induced by feeding the rats with a high-phosphate, high-adenine
diet. Rapamycin reduced the osteogenic markers MSX2 and OSX in this model while increasing
the *Opn* and *Acta2* gene expression via Klotho upregulation (49). Our approached used a genetic
model of MAC to investigate whether rapamycin could inhibit calcification in vivo. MGP-deficient
mice develop normally and at the time of birth are without any overt abnormalities. At
approximately one week old *Mgp$^{-/-}$* mice start exhibiting calcification of the vessels, suggesting
that other gene products might compensate to prevent calcification during development and
immediately post-birth (16). These observations highlight the differences between genetically

induced MAC and high phosphate feeding models. Diet-induced calcification must overcome the
effect of endogenous MGP, which is a potent inhibitor of calcification, while the genetic removal
of MGP results in calcification from the circulating calcium and phosphate and activation of BMP
signaling (12, 15).
443
Rapamycin has also been shown to preserve the differentiated phenotype of vascular SMCs
through IRS-1/P3K/AKT2 pathway (36). Our study did observe rapamycin enhancing expression
of a SMC contractile protein in $Mgp^{-/-}$ mice. Sma expression typically occurs earlier than $Myh11$
expression during SMC development (50). In contrast, $Myh11$ expression occurs later, during the
maturation of SMCs. The first study examined SMC contractile gene expression in response to
rapamycin in vitro (36), while our study looked at the SMC contractile markers in vivo. Further,
the presence of calcification activates mechanosensing pathways that once activated may
prevent full re-expression SMC contractile markers (51). However, further investigation is needed
to elucidate how these mechanisms are operating in the context of the $Mgp^{-/-}$ mouse.
453
MAC is characterized by mineral deposits in the medial vessel layer and frequently occurs in
patients with PAD. Vascular calcification is an active biological process with many driving forces,
including osteogenic dedifferentiation of SMCs, inflammatory signals, hyperphosphatemia, and
extracellular matrix remodeling. In this study, rapamycin prolonged the lifespan of mice who
developed significant MAC shortly after birth. Interestingly, even though rapamycin stimulates the
cellular process of autophagy (52)—which can recycle components of the ECM, including mineral
deposits (53)—calcification volume was not reduced in rapamycin treated $Mgp^{-/-}$ mice. However,
microCT data showed that the mineral density was significantly reduced in $Mgp^{-/-}$ mice treated
with rapamycin. The decrease in mineral density without a change in calcification volume could
suggests a shift from dense, organized macrocalcifications to more dispersed calcium-rich
extracellular vesicles and amorphous calcium phosphate deposits (7, 54). Amorphous calcium
phosphate, primarily composed of calcium and phosphate ions, represents the early stages of
calcification (55). It subsequently acts as a precursor for the nucleation and growth of
hydroxyapatite crystals within the ECM. Stabilization of hydroxyapatite crystals further contribute
to the transition into a bone-like phenotype. One reason that a lower mineral density did not affect
total calcification volume could be a difference in the pattern of mineral deposition. For example,
the mineral may form in a less compact, more porous manner, leading to a lower overall density
despite occupying the same spatial volume. Alternatively, the reduction in density could result
from the breakdown, or remodeling of pre-existing mineral deposits. This suggests that changes
in mineral structure, if explored at a further timepoint, may impact calcification volume.
Alternatively, as MGP also functions to inhibit BMP2 signaling, rapamycin may not interfere with
that process, such that changes in dosage or duration of rapamycin treatment would still not alter
calcification content in the vessel wall. Another explanation is that the activated autophagy
pathways cannot overcome the rapid deposition of hydroxyapatite that occurs in $Mgp^{-/-}$ mice.
Frauscher et al. demonstrated that induction of autophagy by rapamycin treatment decreased
calcification in the vasculature of uremic DBA/2 mice (56). This argument led us to examine
autophagy flux in SMCs. However, we did not observe a difference in autophagy flux between
$Mgp^{+/+}$ and $Mgp^{-/-}$. It is important to note that the $Mgp^{-/-}$ is on a C57BL6J background, perhaps
explaining the differences observed in these in vivo models.
483
It is said that rapamycin is an inhibitor of mammalian target of rapamycin (mTOR), but there is
more complexity to that. mTOR forms two distinct protein complexes known as mTORC1 and
mTORC2. mTORC1, activated by amino acids and allosterically inhibited by rapamycin, requires
a complex specific protein named raptor (57). mTORC1 is acutely sensitive to rapamycin, while

only after prolonged rapamycin treatment (>6 hours) does mTORC2 acquires a sensitivity to
rapamycin (20, 58). In the $Mgp^{-/-};Raptor^{SMC-/-}$ mice, induction of Cre to knockout $Raptor$ was more
lethal than the $Mgp^{-/-};Raptor^{SMC+/+}$ or Mgp-/- mice. This suggests that $Raptor$ is essential for the
growth and maintenance of SMCs in the vasculature. This essential role in growth and
development is perhaps unsurprising, considering that global $Raptor$ knockout mice die in utero
(59). Moreover, there was no benefit to lifespan in the the $Mgp^{-/-}$:Rictor$^{SMC-/-}$ mice. The survival
curves for the the $Mgp^{-/-}$:Rictor$^{SMC+/+}$ and $Mgp^{-/-};Rictor^{SMC-/-}$ mice, targeting mTORC2 signaling, do
not diverge at any point and are statistically insignificant, and also mirror that seen in the $Mgp^{-/-}$
strain.
497
Arterial stiffness is strongly linked to heart failure (60), and these non-atherosclerotic vascular
pathologies are much more common in some populations, such as African-ancestry individuals
who are known to have a higher risk of cardiovascular disease events and mortality compared to
Caucasians(61-65). This excess cardiovascular disease risk is largely due to hypertensive and
peripheral vascular disease, versus atherosclerotic disease (61-63, 66-68). As such, it is critical
to define drivers of MAC such as MGP, which in some clinical studies in humans has been
associated with heart failure and MAC (69). Impairment in MGP activation, which would
phenocopy genetic deletion, is linked to a higher risk of heart failure and peripheral artery disease
due to excessive pulsatile afterloads on the left ventricle that reduce coronary artery perfusion
pressure during diastole (70, 71). As the mice that lack MGP are prone to aortic rupture or heart
failure, we speculate that rapamycin's effect on the mice's lifespan may be due to rapamycin
acting on the cardiac tissues.
510
Our study suggests that rapamycin's life-extending benefit in $Mgp^{-/-}$ mice may act through
improvement in vascular function, which could prevent aortic rupture, or through another organ
system, possibly the heart. However, further studies are needed to examine the impact of
rapamycin on cardiac function in $Mgp^{-/-}$ mice. Rapamycin has been shown to increase
cardiomyocyte autophagy (72, 73), limit cardiomyocyte death, and attenuate cardiomyocyte
hypertrophy and cardiac remodeling by enhancing mTORC2 signaling while simultaneously
inhibiting mTORC1 signaling (74).Rapamycin had a protective effect on cardiomyocytes isolated
from the old mice compared to old mice not treated with rapamycin, as well as improved ejection
fraction, reduced cardiac hypertrophy and inflammation, and downregulated ANP-induced stress
response in vivo (75, 76). In mice with decompensated cardiac hypertrophy, rapamycin improved
cardiac function, such as left ventricular end-systolic dimensions, fractional shortening, and
ejection fraction in mice with compared to the control group (77). Three months of rapamycin
treatment in aged mice increased lifespan, decrease heart rate and heart weight (78). However,
a calcified phenotype was not noted in any of these animal studies.
525
CONCLUSION
This study provides novel insights into the effects of rapamycin on medial arterial calcification
(MAC) using the MGP-deficient murine model. While rapamycin did not reduce the volume of
calcification in the vasculature, it significantly prolonged the lifespan of $Mgp^{-/-}$ mice, possibly
through stabilizing vascular structure. This was evidenced by increased collagen content and
enhanced SMC contractile protein expression, which collectively suggest improved resilience
against vascular rupture. Interestingly, while the volume of calcification was not changed,
rapamycin treatment reduced mineral density. Further, our study underscores the importance of
distinguishing between mTORC1- and mTORC2-mediated pathways. Rapamycin's life-extending
effects appeared independent of autophagy flux or mTORC2 signaling, while the critical role of
mTORC1 in vascular development and maintenance was highlighted, as $Mgp^{-/-};Raptor^{SMC-/-}$ died

earlier than *Mgp⁻/⁻* mice. These findings illustrate rapamycin's role in modifying calcification characteristics and enhancing vascular integrity without altering overall calcification volume. Future investigations should focus on elucidating rapamycin's systemic effects, such as its impact on cardiac function and/or the immune system, as well as understanding how it leads to less-dense mineral deposits, to clarify its protective mechanisms in the context of MAC and cardiovascular health.

ACKNOWLEDGEMENTS
We want to thank the histology core and the Center for Biologic Imaging at the University of Pittsburgh for their help and support with microscopy imaging. Some figures were created with Biorender.com.

SOURCES OF FUNDING
This publication was supported by the NIH K22 HL117917-01A1, The Winters Foundation, The McKamish Family Foundation (St. Hilaire), and the American Heart Association (Behzadi). MicorCT analysis and processing was performed on a system supported by NIH S10-OD021533 (Verdelis).

DISCLOSURES
None.

REFERENCES

1.   **St Hilaire C**. Medial Arterial Calcification: A Significant and Independent Contributor of Peripheral Artery Disease. *Arterioscler Thromb Vasc Biol* 42: 253-260, 2022.

2.   **O'Neill WC, Han KH, Schneider TM, and Hennigar RA**. Prevalence of nonatheromatous lesions in peripheral arterial disease. *Arterioscler Thromb Vasc Biol* 35: 439-447, 2015.

3.   **Narula N, Dannenberg AJ, Olin JW, Bhatt DL, Johnson KW, Nadkarni G, Min J, Torii S, Poojary P, Anand SS, Bax JJ, Yusuf S, Virmani R, and Narula J**. Pathology of Peripheral Artery Disease in Patients With Critical Limb Ischemia. *J Am Coll Cardiol* 72: 2152-2163, 2018.

4.   **Markello TC, Pak LK, St Hilaire C, Dorward H, Ziegler SG, Chen MY, Chaganti K, Nussbaum RL, Boehm M, and Gahl WA**. Vascular pathology of medial arterial calcifications in NT5E deficiency: implications for the role of adenosine in pseudoxanthoma elasticum. *Mol Genet Metab* 103: 44-50, 2011.

5.   **Lanzer P, Hannan FM, Lanzer JD, Janzen J, Raggi P, Furniss D, Schuchardt M, Thakker R, Fok PW, Saez-Rodriguez J, Millan A, Sato Y, Ferraresi R, Virmani R, and St Hilaire C**. Medial Arterial Calcification: JACC State-of-the-Art Review. *J Am Coll Cardiol* 78: 1145-1165, 2021.

6.   **Hortells L, Sur S, and St Hilaire C**. Cell Phenotype Transitions in Cardiovascular Calcification. *Front Cardiovasc Med* 5: 27, 2018.

7.   **Villa-Bellosta R**. Role of the extracellular ATP/pyrophosphate metabolism cycle in vascular calcification. *Purinergic Signal* 19: 345-352, 2023.

8.   **Rutsch F, Ruf N, Vaingankar S, Toliat MR, Suk A, Hohne W, Schauer G, Lehmann M, Roscioli T, Schnabel D, Epplen JT, Knisely A, Superti-Furga A, McGill J, Filippone M, Sinaiko AR, Vallance H, Hinrichs B, Smith W, Ferre M, Terkeltaub R, and Nurnberg P**. Mutations in ENPP1 are associated with 'idiopathic' infantile arterial calcification. *Nat Genet* 34: 379-381, 2003.

9.   **St Hilaire C, Ziegler SG, Markello TC, Brusco A, Groden C, Gill F, Carlson-Donohoe**
H, Lederman RJ, Chen MY, Yang D, Siegenthaler MP, Arduino C, Mancini C, Freudenthal
B, Stanescu HC, Zdebik AA, Chaganti RK, Nussbaum RL, Kleta R, Gahl WA, and Boehm
M. NT5E mutations and arterial calcifications. *N Engl J Med* 364: 432-442, 2011.
10.   **Baral A, F FH, Lee SC, Matthew BP, Ferrante EA, Brofferio A, Cudrici CD, Rollison**
SF, Chen MY, Boehm M, and Wen H. Images in Vascular Medicine: High-resolution CT
imaging of arterial calcification in the hands and legs of patients with CD73 deficiency. *Vasc*
Med 29: 342-344, 2024.
11.   **Jin H, St Hilaire C, Huang Y, Yang D, Dmitrieva NI, Negro A, Schwartzbeck R, Liu**
Y, Yu Z, Walts A, Davaine JM, Lee DY, Donahue D, Hsu KS, Chen J, Cheng T, Gahl W,
Chen G, and Boehm M. Increased activity of TNAP compensates for reduced adenosine
production and promotes ectopic calcification in the genetic disease ACDC. *Sci Signal* 9: ra121,
2016.
12.   **Murshed M, Schinke T, McKee MD, and Karsenty G**. Extracellular matrix
mineralization is regulated locally; different roles of two gla-containing proteins. *J Cell Biol* 165:
625-630, 2004.
13.   **Hackeng TM, Rosing J, Spronk HM, and Vermeer C**. Total chemical synthesis of
human matrix Gla protein. *Protein Sci* 10: 864-870, 2001.
14.   **Malhotra R, Burke MF, Martyn T, Shakartzi HR, Thayer TE, O'Rourke C, Li P,**
Derwall M, Spagnolli E, Kolodziej SA, Hoeft K, Mayeur C, Jiramongkolchai P, Kumar R,
Buys ES, Yu PB, Bloch KD, and Bloch DB. Inhibition of bone morphogenetic protein signal
transduction prevents the medial vascular calcification associated with matrix Gla protein
deficiency. *PLoS One* 10: e0117098, 2015.
15.   **Bostrom K, Tsao D, Shen S, Wang Y, and Demer LL**. Matrix GLA protein modulates
differentiation induced by bone morphogenetic protein-2 in C3H10T1/2 cells. *J Biol Chem* 276:
14044-14052, 2001.
16.   **Luo G, Ducy P, McKee MD, Pinero GJ, Loyer E, Behringer RR, and Karsenty G**.
Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature*
386: 78-81, 1997.
17.   **Haissaguerre M, Saucisse N, and Cota D**. Influence of mTOR in energy and metabolic
homeostasis. *Mol Cell Endocrinol* 397: 67-77, 2014.
18.   **Saxton RA, and Sabatini DM**. mTOR Signaling in Growth, Metabolism, and Disease.
Cell 168: 960-976, 2017.
19.   **Panwar V, Singh A, Bhatt M, Tonk RK, Azizov S, Raza AS, Sengupta S, Kumar D,**
and Garg M. Multifaceted role of mTOR (mammalian target of rapamycin) signaling pathway in
human health and disease. *Signal Transduct Target Ther* 8: 375, 2023.
20.   **Arriola Apelo SI, and Lamming DW**. Rapamycin: An InhibiTOR of Aging Emerges
From the Soil of Easter Island. *J Gerontol A Biol Sci Med Sci* 71: 841-849, 2016.
21.   **Zhan JK, Wang YJ, Wang Y, Wang S, Tan P, Huang W, and Liu YS**. The mammalian
target of rapamycin signalling pathway is involved in osteoblastic differentiation of vascular
smooth muscle cells. *Can J Cardiol* 30: 568-575, 2014.
22.   **He HQ, Law BYK, Zhang N, Qiu CL, Qu YQ, Wu AG, Han Y, Song Q, Zheng WL, Liu**
Y, He YZ, and Wong VKW. Bavachin Protects Human Aortic Smooth Muscle Cells Against
beta-Glycerophosphate-Mediated Vascular Calcification and Apoptosis via Activation of mTOR-
Dependent Autophagy and Suppression of beta-Catenin Signaling. *Front Pharmacol* 10: 1427,
2019.
23.   **Dai XY, Zhao MM, Cai Y, Guan QC, Zhao Y, Guan Y, Kong W, Zhu WG, Xu MJ, and**
Wang X. Phosphate-induced autophagy counteracts vascular calcification by reducing matrix
vesicle release. *Kidney Int* 83: 1042-1051, 2013.

24.   **Joolharzadeh P, and St Hilaire C**. CD73 (Cluster of Differentiation 73) and the
Differences Between Mice and Humans. *Arterioscler Thromb Vasc Biol* 39: 339-348, 2019.
25.   **Ziegler SG, Ferreira CR, MacFarlane EG, Riddle RC, Tomlinson RE, Chew EY,**
Martin L, Ma CT, Sergienko E, Pinkerton AB, Millan JL, Gahl WA, and Dietz HC. Ectopic
calcification in pseudoxanthoma elasticum responds to inhibition of tissue-nonspecific alkaline
phosphatase. *Sci Transl Med* 9: 2017.
26.   **Xin HB, Deng KY, Rishniw M, Ji G, and Kotlikoff MI**. Smooth muscle expression of
Cre recombinase and eGFP in transgenic mice. *Physiol Genomics* 10: 211-215, 2002.
27.   **Sengupta S, Peterson TR, Laplante M, Oh S, and Sabatini DM**. mTORC1 controls
fasting-induced ketogenesis and its modulation by ageing. *Nature* 468: 1100-1104, 2010.
28.   **Magee JA, Ikenoue T, Nakada D, Lee JY, Guan KL, and Morrison SJ**. Temporal
changes in PTEN and mTORC2 regulation of hematopoietic stem cell self-renewal and
leukemia suppression. *Cell Stem Cell* 11: 415-428, 2012.
29.   **Liu M, Espinosa-Diez C, Mahan S, Du M, Nguyen AT, Hahn S, Chakraborty R,**
Straub AC, Martin KA, Owens GK, and Gomez D. H3K4 di-methylation governs smooth
muscle lineage identity and promotes vascular homeostasis by restraining plasticity. *Dev Cell*
56: 2765-2782 e2710, 2021.
30.   **Wanjare M, Kuo F, and Gerecht S**. Derivation and maturation of synthetic and
contractile vascular smooth muscle cells from human pluripotent stem cells. *Cardiovasc Res* 97:
321-330, 2013.
31.   **Zhang L, Yang C, Li J, Zhu Y, and Zhang X**. High extracellular magnesium inhibits
mineralized matrix deposition and modulates intracellular calcium signaling in human bone
marrow-derived mesenchymal stem cells. *Biochem Biophys Res Commun* 450: 1390-1395,
2014.
32.   **Schneider CA, Rasband WS, and Eliceiri KW**. NIH Image to ImageJ: 25 years of
image analysis. *Nat Methods* 9: 671-675, 2012.
33.   **Li Q, Price TP, Sundberg JP, and Uitto J**. Juxta-articular joint-capsule mineralization in
CD73 deficient mice: similarities to patients with NT5E mutations. *Cell Cycle* 13: 2609-2615,
2014.
34.   **Martin KA, Rzucidlo EM, Merenick BL, Fingar DC, Brown DJ, Wagner RJ, and**
Powell RJ. The mTOR/p70 S6K1 pathway regulates vascular smooth muscle cell
differentiation. *Am J Physiol Cell Physiol* 286: C507-517, 2004.
35.   **Marx SO, Jayaraman T, Go LO, and Marks AR**. Rapamycin-FKBP inhibits cell cycle
regulators of proliferation in vascular smooth muscle cells. *Circ Res* 76: 412-417, 1995.
36.   **Martin KA, Merenick BL, Ding M, Fetalvero KM, Rzucidlo EM, Kozul CD, Brown DJ,**
Chiu HY, Shyu M, Drapeau BL, Wagner RJ, and Powell RJ. Rapamycin promotes vascular
smooth muscle cell differentiation through insulin receptor substrate-1/phosphatidylinositol 3-
kinase/Akt2 feedback signaling. *J Biol Chem* 282: 36112-36120, 2007.
37.   **Cheung C, Bernardo AS, Trotter MW, Pedersen RA, and Sinha S**. Generation of
human vascular smooth muscle subtypes provides insight into embryological origin-dependent
disease susceptibility. *Nat Biotechnol* 30: 165-173, 2012.
38.   **Skalli O, Pelte MF, Peclet MC, Gabbiani G, Gugliotta P, Bussolati G, Ravazzola M,**
and Orci L. Alpha-smooth muscle actin, a differentiation marker of smooth muscle cells, is
present in microfilamentous bundles of pericytes. *J Histochem Cytochem* 37: 315-321, 1989.
39.   **Qi M, Zhang L, Ma Y, Shuai Y, Li L, Luo K, Liu W, and Jin Y**. Autophagy Maintains
the Function of Bone Marrow Mesenchymal Stem Cells to Prevent Estrogen Deficiency-Induced
Osteoporosis. *Theranostics* 7: 4498-4516, 2017.

40. **Sun X, Zheng Y, Xie L, Zhou Y, Liu R, Ma Y, Zhao M, and Liu Y**. Autophagy reduces
aortic calcification in diabetic mice by reducing matrix vesicle body-mediated IL-1beta release.
Exp Cell Res 432: 113803, 2023.
41. **Mizushima N, and Yoshimori T**. How to interpret LC3 immunoblotting. *Autophagy* 3:
542-545, 2007.
42. **Ganesh SK, and Subathra Devi C**. Molecular and therapeutic insights of rapamycin: a
multi-faceted drug from Streptomyces hygroscopicus. *Mol Biol Rep* 50: 3815-3833, 2023.
43. **Dobrin PB, and Mrkvicka R**. Failure of elastin or collagen as possible critical
connective tissue alterations underlying aneurysmal dilatation. *Cardiovasc Surg* 2: 484-488,
1994.
44. **Dobrin PB, Baker WH, and Gley WC**. Elastolytic and collagenolytic studies of arteries.
Implications for the mechanical properties of aneurysms. *Arch Surg* 119: 405-409, 1984.
45. **Rahkonen O, Su M, Hakovirta H, Koskivirta I, Hormuzdi SG, Vuorio E, Bornstein P,
and Penttinen R**. Mice with a deletion in the first intron of the Col1a1 gene develop age-
dependent aortic dissection and rupture. *Circ Res* 94: 83-90, 2004.
46. **Sabate M**. Sirolimus Versus Paclitaxel: Second Round. *JACC Cardiovasc Interv* 15:
780-782, 2022.
47. **Rouer M, Xu BH, Xuan HJ, Tanaka H, Fujimura N, Glover KJ, Furusho Y, Gerritsen
M, and Dalman RL**. Rapamycin limits the growth of established experimental abdominal aortic
aneurysms. *Eur J Vasc Endovasc Surg* 47: 493-500, 2014.
48. **Zhou B, Li W, Zhao G, Yu B, Ma B, Liu Z, Xie N, Fu Y, Gong Z, Dai R, Zhang X, and
Kong W**. Rapamycin prevents thoracic aortic aneurysm and dissection in mice. *J Vasc Surg* 69:
921-932 e923, 2019.
49. **Zhao Y, Zhao MM, Cai Y, Zheng MF, Sun WL, Zhang SY, Kong W, Gu J, Wang X,
and Xu MJ**. Mammalian target of rapamycin signaling inhibition ameliorates vascular
calcification via Klotho upregulation. *Kidney Int* 88: 711-721, 2015.
50. **Li L, Miano JM, Cserjesi P, and Olson EN**. SM22 alpha, a marker of adult smooth
muscle, is expressed in multiple myogenic lineages during embryogenesis. *Circ Res* 78: 188-
195, 1996.
51. **Krohn JB, Hutcheson JD, Martinez-Martinez E, Irvin WS, Bouten CV, Bertazzo S,
Bendeck MP, and Aikawa E**. Discoidin Domain Receptor-1 Regulates Calcific Extracellular
Vesicle Release in Vascular Smooth Muscle Cell Fibrocalcific Response via Transforming
Growth Factor-beta Signaling. *Arterioscler Thromb Vasc Biol* 36: 525-533, 2016.
52. **Tanemura M, Ohmura Y, Deguchi T, Machida T, Tsukamoto R, Wada H, Kobayashi
S, Marubashi S, Eguchi H, Ito T, Nagano H, Mori M, and Doki Y**. Rapamycin causes
upregulation of autophagy and impairs islets function both in vitro and in vivo. *Am J Transplant*
12: 102-114, 2012.
53. **Duan ZX, Tu C, Liu Q, Li SQ, Li YH, Xie P, and Li ZH**. Adiponectin receptor agonist
AdipoRon attenuates calcification of osteoarthritis chondrocytes by promoting autophagy. *J Cell
Biochem* 121: 3333-3344, 2020.
54. **Jono S, McKee MD, Murry CE, Shioi A, Nishizawa Y, Mori K, Morii H, and Giachelli
CM**. Phosphate regulation of vascular smooth muscle cell calcification. *Circ Res* 87: E10-17,
2000.
55. **Millan A, Lanzer P, and Sorribas V**. The Thermodynamics of Medial Vascular
Calcification. *Front Cell Dev Biol* 9: 633465, 2021.
56. **Frauscher B, Kirsch AH, Schabhuttl C, Schweighofer K, Ketszeri M, Pollheimer M,
Dragun D, Schroder K, Rosenkranz AR, Eller K, and Eller P**. Autophagy Protects From
Uremic Vascular Media Calcification. *Front Immunol* 9: 1866, 2018.

57.   **Szwed A, Kim E, and Jacinto E**. Regulation and metabolic functions of mTORC1 and
mTORC2. *Physiol Rev* 101: 1371-1426, 2021.
58.   **Goncharova EA**. mTOR and vascular remodeling in lung diseases: current challenges
and therapeutic prospects. *FASEB J* 27: 1796-1807, 2013.
59.   **Guertin DA, Stevens DM, Thoreen CC, Burds AA, Kalaany NY, Moffat J, Brown M,
Fitzgerald KJ, and Sabatini DM**. Ablation in mice of the mTORC components raptor, rictor, or
mLST8 reveals that mTORC2 is required for signaling to Akt-FOXO and PKCalpha, but not
S6K1. *Dev Cell* 11: 859-871, 2006.
60.   **Schott A, Kluttig A, Mikolajczyk R, Greiser KH, Werdan K, Sedding D, and Nuding
S**. Association of arterial stiffness and heart failure with preserved ejection fraction in the elderly
population - results from the CARLA study. *J Hum Hypertens* 37: 463-471, 2023.
61.   **Writing Group M, Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De
Simone G, Ferguson TB, Ford E, Furie K, Gillespie C, Go A, Greenlund K, Haase N,
Hailpern S, Ho PM, Howard V, Kissela B, Kittner S, Lackland D, Lisabeth L, Marelli A,
McDermott MM, Meigs J, Mozaffarian D, Mussolino M, Nichol G, Roger VL, Rosamond W,
Sacco R, Sorlie P, Roger VL, Thom T, Wasserthiel-Smoller S, Wong ND, Wylie-Rosett J,
American Heart Association Statistics C, and Stroke Statistics S**. Heart disease and stroke
statistics--2010 update: a report from the American Heart Association. *Circulation* 121: e46-
e215, 2010.
62.   **Allison MA, Ho E, Denenberg JO, Langer RD, Newman AB, Fabsitz RR, and Criqui
MH**. Ethnic-specific prevalence of peripheral arterial disease in the United States. *Am J Prev
Med* 32: 328-333, 2007.
63.   **Allison MA, Budoff MJ, Nasir K, Wong ND, Detrano R, Kronmal R, Takasu J, and
Criqui MH**. Ethnic-specific risks for atherosclerotic calcification of the thoracic and abdominal
aorta (from the Multi-Ethnic Study of Atherosclerosis). *Am J Cardiol* 104: 812-817, 2009.
64.   **Chaturvedi N, Bulpitt CJ, Leggetter S, Schiff R, Nihoyannopoulos P, Strain WD,
Shore AC, and Rajkumar C**. Ethnic differences in vascular stiffness and relations to
hypertensive target organ damage. *J Hypertens* 22: 1731-1737, 2004.
65.   **D'Agostino RB, Jr., Burke G, O'Leary D, Rewers M, Selby J, Savage PJ, Saad MF,
Bergman RN, Howard G, Wagenknecht L, and Haffner SM**. Ethnic differences in carotid wall
thickness. The Insulin Resistance Atherosclerosis Study. *Stroke* 27: 1744-1749, 1996.
66.   **Ali H, Zmuda JM, Cvejkus RK, Kershaw EE, Kuipers AL, Oczypok EA, Wheeler V,
Bunker CH, and Miljkovic I**. Wnt Pathway Inhibitor DKK1: A Potential Novel Biomarker for
Adiposity. *J Endocr Soc* 3: 488-495, 2019.
67.   **Bild DE, Detrano R, Peterson D, Guerci A, Liu K, Shahar E, Ouyang P, Jackson S,
and Saad MF**. Ethnic differences in coronary calcification: the Multi-Ethnic Study of
Atherosclerosis (MESA). *Circulation* 111: 1313-1320, 2005.
68.   **Budoff MJ, Nasir K, Mao S, Tseng PH, Chau A, Liu ST, Flores F, and Blumenthal
RS**. Ethnic differences of the presence and severity of coronary atherosclerosis. *Atherosclerosis*
187: 343-350, 2006.
69.   **Malhotra R, Nicholson CJ, Wang D, Bhambhani V, Paniagua S, Slocum C,
Sigurslid HH, Lino Cardenas CL, Li R, Boerboom SL, Chen YC, Hwang SJ, Yao C,
Ichinose F, Bloch DB, Lindsay ME, Lewis GD, Aragam JR, Hoffmann U, Mitchell GF,
Hamburg NM, Vasan RS, Benjamin EJ, Larson MG, Zapol WM, Cheng S, Roh JD,
O'Donnell CJ, Nguyen C, Levy D, and Ho JE**. Matrix Gla Protein Levels Are Associated With
Arterial Stiffness and Incident Heart Failure With Preserved Ejection Fraction. *Arterioscler
Thromb Vasc Biol* 42: e61-e73, 2022.

70.   **Dalmeijer GW, van der Schouw YT, Magdeleyns EJ, Vermeer C, Verschuren WM,**
Boer JM, and Beulens JW. Matrix Gla protein species and risk of cardiovascular events in type
2 diabetic patients. *Diabetes Care* 36: 3766-3771, 2013.
71.   **Chirinos JA**. Matrix Gla Protein, Large Artery Stiffness, and the Risk of Heart Failure
With Preserved Ejection Fraction. *Arterioscler Thromb Vasc Biol* 42: 223-226, 2022.
72.   **Buss SJ, Muenz S, Riffel JH, Malekar P, Hagenmueller M, Weiss CS, Bea F,**
Bekeredjian R, Schinke-Braun M, Izumo S, Katus HA, and Hardt SE. Beneficial effects of
Mammalian target of rapamycin inhibition on left ventricular remodeling after myocardial
infarction. *J Am Coll Cardiol* 54: 2435-2446, 2009.
73.   **Gu J, Hu W, Song ZP, Chen YG, Zhang DD, and Wang CQ**. Rapamycin Inhibits
Cardiac Hypertrophy by Promoting Autophagy via the MEK/ERK/Beclin-1 Pathway. *Front*
Physiol 7: 104, 2016.
74.   **Volkers M, Konstandin MH, Doroudgar S, Toko H, Quijada P, Din S, Joyo A,**
Ornelas L, Samse K, Thuerauf DJ, Gude N, Glembotski CC, and Sussman MA. Mechanistic
target of rapamycin complex 2 protects the heart from ischemic damage. *Circulation* 128: 2132-
2144, 2013.
75.   **Chakraborty AD, Kooiker K, Kobak KA, Cheng Y, Lee CF, Razumova M, Granzier**
HH, Regnier M, Rabinovitch PS, Moussavi-Harami F, and Chiao YA. Late-life Rapamycin
Treatment Enhances Cardiomyocyte Relaxation Kinetics and Reduces Myocardial Stiffness.
bioRxiv 2023.
76.   **Flynn JM, O'Leary MN, Zambataro CA, Academia EC, Presley MP, Garrett BJ,**
Zykovich A, Mooney SD, Strong R, Rosen CJ, Kapahi P, Nelson MD, Kennedy BK, and
Melov S. Late-life rapamycin treatment reverses age-related heart dysfunction. *Aging Cell* 12:
851-862, 2013.
77.   **McMullen JR, Sherwood MC, Tarnavski O, Zhang L, Dorfman AL, Shioi T, and**
Izumo S. Inhibition of mTOR signaling with rapamycin regresses established cardiac
hypertrophy induced by pressure overload. *Circulation* 109: 3050-3055, 2004.
78.   **Baghdadi M, Nespital T, Monzo C, Deelen J, Gronke S, and Partridge L**. Intermittent
rapamycin feeding recapitulates some effects of continuous treatment while maintaining lifespan
extension. *Mol Metab* 81: 101902, 2024.

806
807
808 **FIGURE LEGENDS**
809
Figure 1: Histological comparison of Matrix Gla Protein (MGP) mice arteries, medial and
intimal calcification in human arteries. **A.** Histological sections of *Mgp$^{-/-}$* mouse aorta, human
tibial plaque and human coronary artery with atherosclerotic plaque, stained for calcification (Cal)
using Von Kossa Staining. Scale Bars: human tissue 1 mm, mouse tissue 0.5 mm. **B.** Dot Blot
Assay to identify MGP protein in *Mgp$^{+/+}$*, *Mgp$^{+/-}$*, and *Mgp$^{-/-}$* mice. Results representative of n=3
per group. **C.** Representative immunofluorescence images of MGP protein in aortas from *Mgp$^{+/+}$*,
Mgp$^{+/-}$ and *Mgp$^{-/-}$* mice. Scale Bars: 100 µm. n=6 *Mgp$^{+/+}$* vehicle, n=6 *Mgp$^{-/-}$* vehicle. P value
calculated using T-test. Data are shown as means ± SD.
818
Figure 2: Impact of rapamycin treatment on survival in Matrix Gla Protein (*Mgp*)-deficient
mice. A. Graphic illustration of vehicle (DMSO) or rapamycin [5 mg/kg] treatment protocol. **B.**
Kaplan-Meier analysis of overall survival in *Mgp$^{+/+}$* and *Mgp$^{-/-}$* mice treated with vehicle (Dimethyl
sulfoxide; DMSO) or rapamycin (5mg/kg/week) for 90 days until natural death. n=7 *Mgp$^{-/-}$*
rapamycin, n=19 *Mgp$^{-/-}$* vehicle, n=8 *Mgp$^{+/+}$* rapamycin, n=5 *Mgp$^{+/+}$* vehicle. Mantel-Cox test and
Gehan-Breslow-wilcoxon test were used for curve comparison.

Figure 3: Effect of rapamycin treatment on medial calcification in *Mgp*-deficient mice. A.
Graphic illustration of vehicle (DMSO) or 5 mg/kg rapamycin treatment for 25 days in *Mgp*^+/+^ and
Mgp^-/-^ mice. **B.** Representative microCT measurement of axial and cross sections of *Mgp*^-/-^ mice
aortas treated with vehicle or 5 mg/kg rapamycin, administered either once or three times a week
for 25 days. Scale bar 0.2 mm. n=4 *Mgp*^-/-^ 1x rapamycin, n=10 *Mgp*^-/-^ 3x rapamycin, n=12 *Mgp*^-
/-^ vehicle **C.** Representative Von Kossa staining images of *Mgp*^+/+^ and *Mgp*^-/-^ mice aortas treated
with vehicle or rapamycin three times a week for 25 days. Scale bar 0.2 mm. n=10 *Mgp*^-/-^
rapamycin, n=10 *Mgp*^-/-^ vehicle, n=10 *Mgp*^+/+^ rapamycin, n=10 *Mgp*^+/+^ vehicle. **D.** Representative
immunofluorescent images of Runx2 on *Mgp*^+/+^ and *Mgp*^-/-^ mice aortas treated with vehicle or
rapamycin three times a week for 25 days. Scale bar 50 µm. n=10 *Mgp*^-/-^ rapamycin, n=9 *Mgp*^-/-^
vehicle, n=10 *Mgp*^+/+^ rapamycin, n=10 *Mgp*^+/+^ vehicle. Data are shown as means ± SD. P value
were calculated using One-way ANOVA with multiple comparisons test (Figure B) and Two-way
ANOVA with multiple comparisons test (Figures C, D).

**Figure 4: Histological structure of *Mgp*^+/+^ and *Mgp*^-/-^ mice aorta treated with vehicle or
rapamycin. A.** Representative Masson's Trichrome staining images of *Mgp*^+/+^ and *Mgp*^-/-^ mice
aortas treated with vehicle (Veh; DMSO) or rapamycin (RAPA; [5 mg/kg]) three times a week for
25 days. Collagen fiber structure in blue. Scale bar 0.2mm. **B.** Representative Picrosirius staining
images of collagen in *Mgp*^+/+^ and *Mgp*^-/-^ mice treated with vehicle or rapamycin three times a week
for 25 days. Scale bar: 100 µm. n=10 *Mgp*^-/-^ rapamycin, n=10 *Mgp*^-/-^ vehicle, n=10 *Mgp*^+/+^
rapamycin, n=10 *Mgp*^+/+^ vehicle. **C.** Representative Verhoeff-Van Gieson (VVG) staining images
of *Mgp*^+/+^ and *Mgp*^-/-^ treated with vehicle or rapamycin three times a week for 25 days. Elastin
tortuosity and breakage quantified. Scale bar 0.2mm. n=7 *Mgp*^-/-^ rapamycin, n=8 *Mgp*^-/-^ vehicle,
n=8 *Mgp*^+/+^ rapamycin, n=9 *Mgp*^+/+^ vehicle. **D.** Representative immunofluorescent images of
smooth muscle actin (SMA) and myosin heavy chain gene (MYH11) SMA on *Mgp*^+/+^ and *Mgp*^-/-^
mice aortas treated with vehicle or rapamycin three times a week. Scale bar 50 µm. n=10 *Mgp*^-/-^
rapamycin, n=10 *Mgp*^-/-^ vehicle, n=10 *Mgp*^+/+^ rapamycin, n=10 *Mgp*^+/+^ vehicle. **E.** Representative
immunofluorescent images of proliferation markers (ki67) on *Mgp*^+/+^ and *Mgp*^-/-^ mice aortas treated
with vehicle or rapamycin three times a week for 25 days. Scale bar 50 µm. n=10 *Mgp*^-/-^
rapamycin, n=10 *Mgp*^-/-^ vehicle, n=10 *Mgp*^+/+^ rapamycin, n=10 *Mgp*^+/+^ vehicle. Data are shown as
means ± SD. P value were calculated using Two-way ANOVA with multiple comparisons test.

Figure 5: The effect of rapamycin treatment autophagy flux. A. Representative
immunofluorescent images of LC3 on *Mgp*^+/+^ and *Mgp*^-/-^ mice aortas treated with vehicle (Veh;
DMSO) or rapamycin (RAPA; [5 mg/kg]) three times a week for 25 days. LC3 decreased with
rapamycin in *Mgp*^-/-^ mice compared to the *Mgp*^+/+^ mice. Scale bar 50 µm. n=9 *Mgp*^-/-^ rapamycin,
n=10 *Mgp*^-/-^ vehicle, n=10 *Mgp*^+/+^ rapamycin, n=10 *Mgp*^+/+^ vehicle. **B.** Smooth muscle cells from
Mgp^+/+^ and *Mgp*^-/-^ mice were treated with vehicle (DMSO) or rapamycin (200 *µ*M) for 14 hours,
followed by the addition of bafilomycin (BafA, 100 *µ*M), an autophagy inhibitor, for an additional
24 hours. Levels of LC3 (LC3-I and LC3-II) were quantified by western blot. Rapamycin treatment
increases the levels of LC3-II/LC3I and LC3-II protein in both *Mgp*^+/+^ and *Mgp*^-/-^ mice smooth
muscle cells. n=6 *Mgp*^-/-^ rapamycin, n=6 *Mgp*^-/-^ vehicle, n=6 *Mgp*^+/+^ rapamycin, n=6 *Mgp*^+/+^ vehicle.
Data are shown as means ± SD. P value were calculated using One-way ANOVA with multiple
comparisons test (Figure A) and Two-way ANOVA with multiple comparisons test (Figure B).

**Figure 6: Survival probability of Matrix Gla Protein (MGP)-deficient mice with Rictor and
Raptor knockouts in vascular smooth muscle cells (SMC). A.** *Mgp*^-/-^ using Cre recombinase

that target the gene encoding Rictor in SMC, a protein involved in mTORC2 signaling pathways.
Mgp$^{-/-}$*;Rictor*$^{SMC-/-}$ (n=5) and *Mgp*$^{-/-}$*;Rictor*$^{SMC+/+}$ *(n=7)* . **B.** *Mgp*$^{-/-}$ using Cre recombinase that target
the gene encoding Raptor in SMC a protein involved in mTORC1 signaling pathways. *Mgp*$^{-/-}$
;Raptor$^{SMC+/+}$ (n=6), *Mgp*$^{-/-}$*;Raptor*$^{SMC-/-}$ (n=7), The presence or absence of Rictor, when combined
with MGP deficiency, does not significantly affect the lifespan of the mice. However, Raptor plays
a crucial role in the survival of MGP-deficient mice. Data are shown as means ± SD. Mantel-Cox
test and Gehan-Breslow-wilcoxon test were used for curve comparison.
881
882
883
884

A

Mouse *Mgp⁻ᐟ⁻* Aorta	Human Tibial Artery	Human Atherosclerotic Plaque

B

C

A

$Mgp^{+/+}$
$Mgp^{-/-}$

Vehicle or Rapamycin
[5 mg/kg] (3x/week)

d0   d20   Natural
                                   Death

B

Rapamycin
5mg/kg/week

Percent survival

Days old

$Mgp^{+/+}$ Vehicle
$Mgp^{+/+}$ Rapamycin
$Mgp^{-/-}$ Vehicle
$Mgp^{-/-}$ Rapamycin

**

A

LC3/DAPI

Mgp⁺/⁺ Veh   *Mgp*⁺/⁺ RAPA   *Mgp*⁻/⁻ Veh   *Mgp*⁻/⁻ RAPA

LC3 / DAPI

0.0502

○ *Mgp*⁺/⁺
○ *Mgp*⁻/⁻

B

Vehicle (DMSO)	Vehicle (DMSO)
Rapamycin	Rapamycin
Vehicle (DMSO)	Vehicle + BafA
Rapamycin	Rapamycin + BafA

14h   24h

Mgp⁺/⁺   *Mgp*⁻/⁻

Vehicle   LC3-I / LC3-II   — 15 KD
     Tubulin   — 57 KD

Rapamycin [200uM]   LC3-I / LC3-II   — 15 KD
     Tubulin   — 57 KD

BafA [100uM]   LC3-I / LC3-II   — 15 KD
     Tubulin   — 57 KD

Rapamycin [200uM] + BafA [100uM]   LC3-I / LC3-II   — 15 KD
     Tubulin   — 57 KD

LC3-II / Tubulin

0.0105
0.0466
0.0226
0.0194

LC3-II / LC3-I ratio

0.0217
0.0292
0.0419
0.0518

○ *Mgp*⁺/⁺
○ *Mgp*⁻/⁻

Collagen Breakage	↓ **Collagen Breakage**
Collagen	↑ **Collagen**
SMA	↑ **SMA**
Calcification Volume	↔ **Calcification Volume**
Mineral Density	↓ **Mineral Density**

Mgp⁻/⁻ Veh — Lifespan 13-43 days

Mgp⁻/⁻ RAPA — Lifespan 25-90 days

★ Mineralization   Elastin   Collagen   Smooth muscle cell with more SMA   Smooth muscle cell with less SMA

Matrix Gla protein (Mgp)-deficient mice develop extensive medial arterial calcification and die within 2 months of birth. We found that while treatment of these mice with rapamycin doubles their lifespan and improved the vascular structure. Vehicle (Veh), rapamycin (RAPA), Extracellular matrix (ECM), Smooth muscle alpha actin (SMA).