Systems-level reconstruction of kinase phosphosignaling networks

regulating endothelial barrier integrity using temporal data

Ling Wei[1,*], John D. Aitchison[1,2,3], Alexis Kaushansky[1,2,4], and Fred D. Mast[1,2,*]

[1]Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle, WA 98109, United States

[2]Department of Pediatrics, University of Washington, Seattle, WA 98105, United States

[3]Department of Biochemistry, University of Washington, Seattle, WA 98105, United States

[4]Department of Global Health, University of Washington, Seattle, WA 98105, United States

*Corresponding author. E-mail: ling.wei@seattlechildrens.org and fred.mast@seattlechildrens.org

Abstract

Phosphosignaling networks control cellular processes. We built kinase-mediated

regulatory networks elicited by thrombin stimulation of brain endothelial cells using two

computational strategies: Temporal Pathway Synthesizer (TPS), which uses

phosphoproteomics data as input, and Temporally REsolved KInase Network

Generation (TREKING), which uses kinase inhibitor screens. TPS and TREKING

predicted overlapping barrier-regulatory kinases connected with unique network

topology. Each strategy effectively describes regulatory signaling networks and is

broadly applicable across biological systems.

Keywords: endothelial barrier permeability; thrombin, phosphosignaling network;

Temporal Pathway Synthesizer (TPS); Temporally REsolved KInase Network

Generation (TREKING)

Protein kinases regulate numerous cellular processes through phosphorylation of

target proteins. These series of phosphosignaling events are triggered in response to

environmental changes and propagate within a network structure. The function of

kinases in cell signaling and regulation of cellular phenotypes has been intensively

explored in different contexts, including in mediating brain endothelial barrier integrity [1].

The integrity of the blood-brain barrier is essential for maintaining the homeostasis of the

central nervous system [2]. The permeability of brain endothelium is tightly controlled by

cell-cell junctions and focal adhesions, whose functions are heavily regulated by kinases

[1-3]. Therefore, understanding the network-level connectivity of these barrier-regulatory

kinases and their temporal dynamics is critical for developing targeted therapeutic

strategies for treating vascular leakage.

In recent years, computational tools and modeling approaches have been

advanced to facilitate reconstructing high-confident, systems-level protein signaling

networks based on proteomics data and prior knowledgebases (Table 1), but they often

fall short of providing detailed mechanistic understanding and/or fine temporal

information on cell signaling and their function in mediating cellular phenotypes. For

example, Babur *et al.* developed a pathway extraction method that maps high-

throughput proteomics data to the Pathway Commons database. They used a graphical

pattern search framework to identify the protein-protein interactions (PPIs) that explain

correlated changes in proteomics profiles [4]. Network inference methods like dynamic

Bayesian networks have been implemented on time course phosphorylation data to infer

protein signaling network structure specific to biological context without relying on user-

defined parameters [5-7]. Ordinary differential equation (ODE) models are kinetic models

that have been conventionally applied to analyze biochemical reaction networks, which

can capture the mechanistic details of biochemical processes and predict the temporal

evolution of variables that cannot be directly measured [8]. Unfortunately, building robust

medium- or large-scale ODE models is challenging due to a lack of detailed mechanistic

description of the biological system and insufficient experimental data to parameterize

the models [8,9]. Unbiased network reconstruction has benefitted from the development of

logic modeling, which employs either discrete logic models or continuous ODEs

transformed from discrete logic models, followed by optimization to fit experimental data,

to generate predictive models that describe context-specific signaling within the generic

protein signaling networks [10,11].

Besides building signaling networks from phosphoproteomics data, which are

typically collected at relatively sparse time points, network generation pipelines can use

functional screens with fine temporal resolution as input along with known PPIs [12,13]

(Table 1). For example, Bello *et al.* extended the kinase regression (KiR) approach that

was initially designed to identify key kinase regulators of a particular cellular phenotype

from quantitative drug screens. Their KiRNet method maps KiR hits onto existing PPI

networks using network propagation to create kinase-centered, functional network

models associated with the phenotype of interest [12]. We recently developed a novel

approach called TREKING that integrates temporal drug screen data with a known

kinase-substrate phosphorylation database to infer functional kinase signaling networks

with temporal resolution [13].

Köksal *et al.* have also developed a network inference algorithm specifically to

model time course data [14]. The Temporal Pathway Synthesizer (TPS) uses time-

resolved phosphoproteomics data as input [14]. TPS assembles signaling pathways by

first extracting a subnetwork that connects the phosphorylated proteins to the source

node(s) from a background PPI network. It then uses the temporal phosphorylation data

75 to systematically examine all directed, signed edges in the subnetwork to determine how

76 signals propagate from source node(s). Finally, TPS consolidates all valid paths into a

77 unified network that represents the pathway structures facilitating phosphosignal

78 propagation.

79 **Table 1. List of computational approaches for building and analyzing protein**

80 **phosphosignaling networks.**

Methodology	Description	Input experimental data	Output model	References
CausalPath	Extract causal links in pathways by analyzing changes in high-throughput proteomics profiles on the background of prior knowledge captured in biochemical reaction knowledgebases	High-throughput proteomics data	A set of causal links between measurable molecular features that can explain correlated changes in a given set of proteomics and other molecular profiles	Babur *et al.* [4]
Dynamic Bayesian inference	Infer network structure and probabilistic relationships among a set of variables over time without user-set tuning parameters	Time-course phosphoprotein data	A graphic model consisting of kinase nodes and casual relationships between them with probability distribution function assigned	Hill *et al.* [5] Hill *et al.* [6] Merrell and Gitter [7]
Information theory	Quantitatively predict strength of underlying complex relationships between proteins from noisy data based on the formalism of information theory	Single-cell protein phosphorylation data	A network denoted by the amount of mutual information transduced by the signaling pathways in response to external signals	Cheong *et al.* [15] Krishnaswamy *et al.* [16]
Differential	Describe biochemical	Kinetic	A set of	Frohlich,

equations	processes in the system using a set of coupled differential equations and predict temporal and spatial evolution of the latent variables	parameters of biochemical reactions	interacting molecular species trackable (amount and cellular localization) over time	Loos, and Hasenauer [9]
Logic modeling	Model predictive signaling pathways based on networks of logic gates that specify how input signals lead to changes of cellular state	Functional biochemical data and phosphoproteomics data	A data-optimized Boolean model (PPIs are in logical representations) that can predict cell responses to specific biological stimuli	Saez-Rodriguez et al. [10] Vaga et al. [11]
KiRNet	Generate kinase-centered, functional network models by mapping kinase hits from functional kinase inhibitor screens onto a PPI network using network propagation approach	Phenotypic data from small-scale chemical screens	A focused subnetwork centered around the predicted key kinases that represents the proteins and relationships most critical for the phenotype of interest	Bello et al. [12]
TREKING	Inform functionality of kinases and kinase signaling networks with temporal resolution based on temporal phenotypic readouts and existing biochemical data	Phenotypic data from small-scale chemical screens	A set of local kinase phosphosignaling networks describing the connections between functional kinases associated with temporal cellular phenotypes	Wei et al. [13]
TPS	Assemble paths through which phosphosignals propagate from source node(s) based on the temporal activity of proteins and known PPIs	Temporal phosphoproteomics data	A summary network that compiles the valid pathway models that depict how phosphosignals may propagate from source	Köksal et al. [14]

			node(s)	

Phosphoproteomics-based kinase signaling networks offer a distinct perspective

from those derived from functional assays. While phosphoproteomics help elucidate how

phosphosignals propagate within phosphosignaling networks, functional assays can link

kinase activity directly to cellular phenotypes. Comparing these two network

reconstruction approaches enhances our understanding and interpretation of

phosphosignaling events crucial to the phenotype of interest.

Here, we reconstructed a phosphosignaling network regulating thrombin-induced

endothelial barrier disruption and recovery using the TPS algorithm (Fig. 1). We then

compared this network with a TREKING network we previously assembled using kinase

inhibitor screens in the same cellular background. The input time series data for TPS

were collected from western blot experiments as described in [13]. Briefly, human brain

microvascular endothelial cells (HBMECs) were treated with thrombin and the

phosphorylation status on the activation sites of 25 kinases, and 3 non-kinase proteins

known to regulate barrier function, were collected at nine time points across a 6-hour

time window after thrombin treatment. Data on 17 kinases have been published

previously [13], and data generated in this study on the remaining 8 kinases and 3 non-

kinase proteins are available in Supplementary Data S1. The input network that

connects phosphorated proteins to the source node, proteinase-activated receptor 1

(F2R/PAR1), was generated using Omics Integrator [17,18] (Supplementary Data S2). The

input files for TPS were prepared according to [14] and are described briefly in Methods.

The TPS-generated network resulted in 966 directed edges, with only five

signed, directed edges due to the sparsity of the input time series data (Fig. 2a,

Supplementary Data S3). The network recapitulates interactions between extracellular

signal-regulated kinases (ERKs; MAPK1/ERK2 and MAPK3/ERK1) and β-arrestins

(ARRB1 and ARRB2), key components of canonical thrombin signaling through PARs

(Reactome pathway R-HSA-456926: thrombin signaling through proteinase activated

receptors (PARs), Supplementary Data S5). The model predicts that both ABL proto-

oncogene 1 non-receptor tyrosine kinase (ABL1/c-Abl) and mitogen-activated protein

kinase 8 (MAPK8/JNK1) activate paxillin (PXN), consistent with previous findings [19,20].

However, the model's prediction that protein tyrosine kinase 2 (PTK2/FAK) inhibits PXN

contradicts established literature [21] (Fig. 2a). To make comparisons with the TREKING

network, which currently includes only kinase-to-kinase connections, we extracted

kinase-kinase interactions from the TPS network, yielding a subnetwork of 153 edges

(Fig. 2b (bottom left), Supplementary Data S4). Among the 126 kinases predicted by

TREKING, 56 (44.4%) kinases were also inferred by the TPS model, demonstrating the

ability of these two distinct methodologies to infer the set of kinases activated and play a

functional role in barrier regulation. Multiple members of the canonical MAPK cascades

were predicted by both TPS and TREKING, including ERKs (MAPK1/ERK2 and

MAPK3/ERK1), Jun N-terminal kinases (JNKs; MAPK8/JNK1 and MAPK9/JNK2) and

MAPK14/p38α, whose roles in temporal barrier function have been characterized [13].

Additionally, one of the switch kinases, MAPK activated protein kinase 2

(MAPKAPK2/MK2), identified by TREKING as having both barrier-disruptive and barrier-

restorative roles, was also predicted by TPS to mediate barrier properties, though TPS

does not distinguish between barrier-disruptive or barrier-protective functions.

Among the 153 directed kinase-to-kinase connections in the TPS network, 50

(32.7%) are also present in the TREKING network (Fig. 2b, bottom). Both TPS and

TREKING predicted regulation of MAPKAPK2/MK2 by MAPK1/ERK2 and MAPK14/p38α

in thrombin-treated endothelial cells (Fig. 2c). TREKING further predicted that the

MAPK1/ERK2-MAPKAPK2/MK2 interaction is crucial for barrier disruption, while

MAPK14/p38α-MAPKAPK2/MK2 interaction is important for late barrier recovery. Unlike

TREKING, TPS does not include phenotypic data in its network reconstruction. Instead,

TPS inferred 103 kinase-to-kinase connections not captured by TREKING, including

several signaling hubs such as outgoing signals from protein kinase cAMP-activated

catalytic subunit alpha (PRKACA), SRC and protein kinase C alpha (PRKCA), and

incoming signals to glycogen synthase kinase 3 beta (GSK3B) (Fig. 2c).

137     TPS's ability to infer interactions between kinase regulators and their non-kinase

substrates makes it a powerful tool for linking upstream kinase signaling to cellular

phenotypes, as proteins directly associated with phenotypes are often better

characterized and usually not kinases. Despite relying on a sparse input dataset, TPS

predicted that cadherin 5 (CDH5), an endothelial adherens junction protein, is regulated

by SRC. It also predicted that heat shock protein family B member 1 (HSPB1/HSP27),

which directly regulates actin organization in endothelial cells [22], is regulated by MAPK

kinases MAPK14/p38α and MAPKAPK2/MK2, PRKACA, and protein kinase CGMP-

dependent 1 (PRKG1) (Supplementary Data S3); these predictions are supported by

experimental validations [22-24]. Additionally, TPS predicted that PXN, a cytoskeletal

protein involved in actin-membrane attachment at focal adhesions, is regulated by Src

family kinases FYN and SRC, p21 (RAC1) activated kinase 1 (PAK1), and several

MAPK kinases, including MAPK1/ERK2, MAPK3/ERK1, MAPK8/JNK1 and

MAPK14/p38α (Supplementary Data S3). These regulatory interactions on PXN and

their roles in barrier regulation are consistent with previous studies [25-27].

152     Along with providing a summary network, the TPS pipeline also provides activity

windows for each node. However, due to the sparse sampling of the input time series

data, TPS could not consistently identify precise activity states (activation, inhibition, or

inactivity) for proteins at specific time points, except for those with time course data. This

underscores a limitation of TPS in reliably inferring global signaling cascades when

working with small input datasets.

We systematically evaluated the accuracy of the predicted barrier-regulatory

networks in response to thrombin by comparing the model's predictions with

independent experimental data not used to build the networks (Fig. 3a). For the TPS

network, we calculated the cumulative inhibition of each kinase inhibitor on the network

(see Methods) as a surrogate for overall perturbation of the barrier-regulatory kinase

activity, using the kinase-compound biochemical data [28]. A cutoff for cumulative

inhibition – 25% of the kinases in the TPS network are inhibited by 50% – was set,

above which the kinase inhibitor is considered to substantially perturb the network and

therefore predicted to alter the barrier permeability. Among the 29 kinase inhibitors in the

phenotypic screen [29], the model predicts 20 kinase inhibitors to alter the barrier

permeability, 18 (90%) of which agree with the measurements (Fig. 3b). We also

generated random networks by randomizing both input networks and protein

phosphorylation data and ran them through the TPS algorithm (see Methods). The TPS

network outperforms random networks in predicting kinase-targeting compounds that

modulate barrier permeability (Fig. 3b).

To evaluate the accuracy of the TREKING model for predicting kinase activity,

we compared the model predictions of barrier-regulatory kinases with the western blot

data used for building the TPS network (Fig. 3a). We first used the TREKING model to

generate time-resolved predictions of the activation states of kinases in response to

thrombin. To assess the barrier function of a kinase at a particular time point, a 10-

minute time window centered at that time point was applied. If any of the signaling

pathways with which the kinase is associated are active within the time window, the

model predicts the kinase to be functionally active within that time window; otherwise,

the model predicts the kinase to be functionally inactive within the time window.

TREKING predicts 129 instances of active barrier function across all time points; 97

(75.2%) differential phosphorylation events were observed by western blot (Fig. 3c). We

also generated random networks by randomly sampling temporal profiles for kinases,

assigning kinases to random groups, and building phosphosignaling networks based on

randomly generated kinase-substrate phosphorylation interactions (see Methods).

Overall, the TREKING network outperforms random models in predicting barrier-

regulatory kinases with temporal resolution (Fig. 3c).

In this study, we compared two computational approaches, TPS and TREKING,

in reconstructing kinase signaling networks describing phosphosignal propagation within

brain endothelial cells after thrombin stimulation. Overall, TPS and TREKING provide

different insights into kinase regulatory networks associated with cellular phenotypes.

TPS focuses on the activity of functional proteins, while TREKING focuses on the

functionality of kinases altering phenotypes. Since the phosphorylation state of a kinase

does not directly imply its function, which is also impacted by its subcellular localization

and presence of substrates, discrepancies between the networks reconstructed by these

two approaches are expected. Both methodologies are easily generalizable to various

biological systems and, in the context of infectious diseases, can deconvolve kinase-

mediated signaling mechanisms in host cells or parasites. In the future, integration of

TPS and TREKING will facilitate developing more informative network models.

Integrating TPS and TREKING methodologies would allow 1) revealing the functionality

of kinases and their inter-connections that alter cellular phenotypes along with their

temporal enzymatic activity during changes of cellular phenotypes, and 2) linking

functional non-kinase proteins that directly modulate cellular phenotypes to their

upstream, functional kinase regulatory pathways that control the activity and function of

the proteins.

Methods

Cell lines

Primary human brain microvascular endothelial cells (HBMECs; Cell Systems Cat# ACBRI 376) were cultured on rat tail collagen type I (5 µg/cm^2; Corning Cat# 354236) in HBMEC culture media (Lonza Cat# CC-3202) at 37°C and 5% CO_2. HBMECs were obtained at passage 3 and used until passage 9.

Lysate preparation

HBMECs were seeded in 6-well plates (Corning Cat# 353046) at 55,000 cells/well and grown for 4 days with media change every other day. On the day of lysate collection, cells were equilibrated in serum-free culture media for 1 hour and then treated with thrombin at a final concentration of 5 nM for 5, 15, 30, 60, 120, 180, 240, 360 minutes. After the indicated incubation periods, cells were washed twice with ice-cold phosphate buffered saline (PBS) and lysed in sodium dodecyl sulfate (SDS) lysis buffer (50 mM Tris-HCl, 2% SDS, 5% glycerol, 5 mM ethylenediaminetetraacetic acid, 1 mM sodium fluoride, 10 mM β-glycerophosphate, 1 mM phenylmethylsulfonyl fluoride, 1 mM sodium orthovanadate, 1 mM dithiothreitol, supplemented with a cocktail of protease inhibitors (Roche Cat# 4693159001) and phosphatase inhibitors (Sigma-Aldrich Cat# P5726)). Cell lysates were clarified in filter plates (Pall Cat# 8075) at 2,671$\times g$ for 30 minutes, after which they were stored at -80°C until use. Cell lysates from three biological replicates were collected.

Western blot

All the gel electrophoresis was performed using Bolt™ 4-12% Bis-Tris mini protein gels. Proteins were transferred to PVDF membranes using iBlot 2 (Thermo Fisher Scientific

Cat# IB21001) or iBlot 3 (Thermo Fisher Scientific Cat# A56727) dry blotting system.

Primary antibodies were used at concentrations recommended by the manufacturer (see

Supplementary Data S1 for antibody information). Antibody to GAPDH (Cell Signaling

Technology Cat# 97166) was used as loading control at 1:2000 dilution. Blots were

imaged using Bio-Rad ChemiDoc imaging system and signals were quantified using

ImageJ2 (https://imagej.nih.gov/ij/, version 2.3.0). Background correction was done for

each band by subtracting background signals nearby the band. The signals from

proteins of interest were first normalized to the signals from GAPDH, and then the

signals at each time point were normalized to the signals at time zero for the fold change

of phosphorylation from basal level. Phosphorylation on activation sites of 25 kinases

and 3 non-kinase proteins were evaluated at nine time points (0, 5, 15, 30, 60, 120, 180,

240 and 360 minutes after thrombin treatment). Western blot was performed on cell

lysates collected from three biological replicates.

Reconstruction of TPS network

The input time series data for TPS were collected from western blot experiments. A

portion of the data were previously published [13]. The input network, an undirected

subnetwork of the PPI network that connects the phosphorylated proteins to the source

node(s), was generated using the Omics Integrator implementation of the Prize-

Collecting Steiner Forest algorithm [17,18]. Time series phosphorylation data, protein prizes

and interactome network were required for generating the input network. Protein prizes

were obtained using significance scores for each time point relative to the first time point

and to the previous time point for each profile. The interactome network was compiled by

combining human PPIs from iRefIndex [30] (https://irefindex.vib.be//) and kinase-substrate

phosphorylation interactions from PhosphoSitePlus [31] (https://www.phosphosite.org).

Omics Integrator was run with dummy edge weight of 10, edge reliability of 10, degree

penalty of 0, and 100 randomize prize runs. Significance scores for TPS run were

computed with paired Student's t-tests comparing the phosphorylation intensity at each

time point with the first time point ("firstscores") or comparing the phosphorylation

intensity at the current time point with the preceding time point ("prevscores"). The

threshold for significance scores was set to 0.05, above which measurements are

considered non-significant. Proteinase-activated receptor 1 (F2R/PAR1) was designated

as the sole source node for TPS run, as it is the predominant thrombin receptor and the

only member of PARs mediating thrombin-induced phosphoregulation in endothelial

cells [32,33]. An optional partial model was provided for TPS run, which is a directed

kinase-substrate phosphorylation interaction network from PhosphoSitePlus [31]

(https://www.phosphosite.org).

Generation of random models

The random TPS summary networks were generated following the TPS pipeline. During

each iteration, a random subnetwork and a random partial model were generated by

randomly shuffling the nodes in the original networks. Random time series

phosphorylation data, random significance scores with respect to the first time point

("firstscores") or the previous time point ("prevscores") of the profile were sampled by

randomly shuffling the values in the original datasets. A threshold for significance scores

was set to 0.05. F2R/PAR1 was designated as the sole source node for generating

random summary networks. In total, 100 random summary networks were generated

and used to compare with the TPS network.

The random TREKING models were generated following the TREKING pipeline.

During each iteration, a random background phosphosignaling network with the same

network topology as the background phosphosignaling network used in TREKING was

generated using the configuration model function in NetworkX package. All the kinases

were shuffled and randomly assigned to each of the nodes in the random network. At

each time point, kinases of barrier-weakening or barrier-strengthening functionality were

randomly selected from the 300 kinases used for KiR, with the same probabilities as in

the TREKING model at the corresponding time point. The random sampling of kinases

was performed at all time points corresponding to the TREKING model. Each of the

kinases selected at any time point during the time course was randomly assigned to one

of the 36 clusters, corresponding to the total number of neurons generated by SOMs.

For each cluster, the phosphosignaling network was reconstructed by searching for the

shortest paths between any pair of kinases within the cluster, using the random

phosphosignaling network generated for this iteration as the background network. In

total, 100 random models were generated and used to compare with the TREKING

predictions.

Visualization

Figs. 1-3 were made in Inkscape (https://inkscape.org, version 1.2.1), a free and open-

source vector graphics editor licensed under the GPL. In Fig. 1, the hierarchically-

clustered heatmap was generated using seaborn (https://seaborn.pydata.org, version

0.12.0), a Python data visualization library based on Matplotlib; protein-protein

interaction network and phosphosignaling networks were visualized using Cytoscape

(https://cytoscape.org, version 3.9.1), with layouts generated from yFiles layout

algorithms (https://www.yworks.com/products/yfiles-layout-algorithms-for-cytoscape,

version 1.1.4); the logo for Omics Integrator was obtained from the website of its original

developers (https://fraenkel-nsf.csbi.mit.edu/omicsintegrator/). In Fig. 2, panel (a) was

created in BioRender (https://app.biorender.com); phosphosignaling networks in panel

(b) were visualized using NetworkX (https://github.com/networkx/networkx, version

2.8.6), a Python package for analyzing complex network structures; phosphosignaling

network in panel (c) was visualized using Cytoscape (https://cytoscape.org, version

3.9.1), with its layout generated from yFiles hierarchic layout algorithm

(https://www.yworks.com/products/yfiles-layout-algorithms-for-cytoscape, version 1.1.4).

In Fig. 3, panel (a) was created in BioRender (https://app.biorender.com); panels (b)-(c)

were generated using Matplotlib (https://matplotlib.org, version 3.5.3) in Python.

Quantification and statistical analysis

The kinase-compound biochemical data [28] were used to calculate the cumulative

inhibition of kinase inhibitors on the TPS network. For each kinase inhibitor in the

screen, its cumulative inhibition was calculated by summing the percentage of inhibition

(100 – residual activity) across all the kinases in TPS network (if they are also in the

biochemical data).

Student's t-test was used to evaluate the difference of barrier permeability

between kinase inhibitor- and DMSO-treated conditions. The permeability data are from

Dankwa *et al.* [29] For each kinase inhibitor treatment, both p-value and percent change of

normalized area under the curve (AUC) across a 6-hour time window with respect to

DMSO-treated condition were calculated to determine if the two sets of data differ from

each other. A p-value below 0.05 or all biological replicates reporting a percent change

of normalized AUC greater than 10% above/below the DMSO control indicates that the

barrier permeability is different between kinase inhibitor- and DMSO-treated conditions.

Student's t-test was also used to evaluate the difference of kinase activity between non-

treated and thrombin-treated conditions. At each time point, both p-value and fold

change of phosphorylation with respect to non-treated cells were reported to determine if

the two sets of data differ. A p-value below 0.05 or all biological replicates reporting a

fold change increasing/decreasing by at least 20% compared to non-treated cells

indicates that the kinase activity is different between non-treated and thrombin-treated

conditions. SciPy statistical package (version 1.9.1) was used to perform the t-tests.

Data availability

A portion of the data used in this study have been published previously. The remaining data generated in this study is available in the supplementary information.

Acknowledgments

The authors would like to thank Joseph D. Smith and Selasi Dankwa for helpful discussions. This work was funded by National Institutes of Health (NIH) grants R01 AI148802, R61 HL154250 and R33 HL154250 (to A.K.), and R01 DK041737 and R01 GM109824 (to J.D.A.). F.D.M. is the recipient of a Career Development Award from Seattle Children's Research Institute. The funders played no role in study design, data collection, analysis and interpretation of data, or the writing of this manuscript.

Author contributions

Ling Wei: Conceptualization, Investigation, Writing – Original Draft, Writing – Review & Editing. **John D. Aitchison:** Funding acquisition, Supervision, Writing – Review & Editing. **Alexis Kaushansky:** Conceptualization, Funding acquisition, Supervision, Writing – Review & Editing. **Fred D. Mast:** Conceptualization, Investigation, Supervision, Writing – Review & Editing.

Competing Interests

All authors declare no financial or non-financial competing interests.

References

References

1   Kuppers, V., Vockel, M., Nottebaum, A. F. & Vestweber, D. Phosphatases and kinases as
regulators of the endothelial barrier function. *Cell Tissue Res* **355**, 577-586 (2014).
https://doi.org:10.1007/s00441-014-1812-1

2   Daneman, R. & Prat, A. The blood-brain barrier. *Cold Spring Harb Perspect Biol* **7**,
a020412 (2015). https://doi.org:10.1101/cshperspect.a020412

3   Mehta, D. & Malik, A. B. Signaling mechanisms regulating endothelial permeability.
Physiol Rev **86**, 279-367 (2006). https://doi.org:10.1152/physrev.00012.2005

4   Babur, O. *et al.* Causal interactions from proteomic profiles: Molecular data meet
pathway knowledge. *Patterns (N Y)* **2**, 100257 (2021).
https://doi.org:10.1016/j.patter.2021.100257

5   Hill, S. M. *et al.* Bayesian inference of signaling network topology in a cancer cell line.
Bioinformatics **28**, 2804-2810 (2012). https://doi.org:10.1093/bioinformatics/bts514

6   Hill, S. M. *et al.* Context Specificity in Causal Signaling Networks Revealed by
Phosphoprotein Profiling. *Cell Syst* **4**, 73-83 e10 (2017).
https://doi.org:10.1016/j.cels.2016.11.013

7   Merrell, D. & Gitter, A. Inferring signaling pathways with probabilistic programming.
Bioinformatics **36**, i822-i830 (2020). https://doi.org:10.1093/bioinformatics/btaa861

8   Klipp, E., Herwig, R., Kowald, A., Wierling, C. & Lehrach, H. *Systems biology in practice:*
concepts, implementation and application. (John Wiley & Sons, 2005).

9   Frohlich, F., Loos, C. & Hasenauer, J. Scalable Inference of Ordinary Differential Equation
Models of Biochemical Processes. *Methods Mol Biol* **1883**, 385-422 (2019).
https://doi.org:10.1007/978-1-4939-8882-2_16

10   Saez-Rodriguez, J. *et al.* Discrete logic modelling as a means to link protein signalling
networks with functional analysis of mammalian signal transduction. *Mol Syst Biol* **5**,
331 (2009). https://doi.org:10.1038/msb.2009.87

11   Vaga, S. *et al.* Phosphoproteomic analyses reveal novel cross-modulation mechanisms
between two signaling pathways in yeast. *Mol Syst Biol* **10**, 767 (2014).
https://doi.org:10.15252/msb.20145112

12   Bello, T. *et al.* KiRNet: Kinase-centered network propagation of pharmacological screen
results. *Cell Rep Methods* **1** (2021). https://doi.org:10.1016/j.crmeth.2021.100007

13   Wei, L., Dankwa, S., Vijayan, K., Smith, J. D. & Kaushansky, A. Interrogating endothelial
barrier regulation by temporally resolved kinase network generation. *Life Sci Alliance* **7**
(2024). https://doi.org:10.26508/lsa.202302522

14   Koksal, A. S. *et al.* Synthesizing Signaling Pathways from Temporal Phosphoproteomic
Data. *Cell Rep* **24**, 3607-3618 (2018). https://doi.org:10.1016/j.celrep.2018.08.085

15   Cheong, R., Rhee, A., Wang, C. J., Nemenman, I. & Levchenko, A. Information
transduction capacity of noisy biochemical signaling networks. *Science* **334**, 354-358
(2011). https://doi.org:10.1126/science.1204553

16   Krishnaswamy, S. *et al.* Systems biology. Conditional density-based analysis of T cell
signaling in single-cell data. *Science* **346**, 1250689 (2014).
https://doi.org:10.1126/science.1250689

17   Tuncbag, N. *et al.* Network-Based Interpretation of Diverse High-Throughput Datasets
through the Omics Integrator Software Package. *PLoS Comput Biol* **12**, e1004879 (2016).
https://doi.org:10.1371/journal.pcbi.1004879

18   Kedaigle, A. J. & Fraenkel, E. Discovering Altered Regulation and Signaling Through
Network-based Integration of Transcriptomic, Epigenomic, and Proteomic Tumor Data.
Methods Mol Biol **1711**, 13-26 (2018). https://doi.org:10.1007/978-1-4939-7493-1_2

19   Fu, P. *et al.* Role played by paxillin and paxillin tyrosine phosphorylation in hepatocyte
growth factor/sphingosine-1-phosphate-mediated reactive oxygen species generation,
lamellipodia formation, and endothelial barrier function. *Pulm Circ* **5**, 619-630 (2015).
https://doi.org:10.1086/683693

20   Huang, C., Rajfur, Z., Borchers, C., Schaller, M. D. & Jacobson, K. JNK phosphorylates
paxillin and regulates cell migration. *Nature* **424**, 219-223 (2003).
https://doi.org:10.1038/nature01745

21   Sato, T. *et al.* A FAK/HDAC5 signaling axis controls osteocyte mechanotransduction. *Nat*
Commun **11**, 3282 (2020). https://doi.org:10.1038/s41467-020-17099-3

22   Rada, C. C. *et al.* Heat shock protein 27 activity is linked to endothelial barrier recovery
after proinflammatory GPCR-induced disruption. *Sci Signal* **14**, eabc1044 (2021).
https://doi.org:10.1126/scisignal.abc1044

23   Klomp, J. E. *et al.* Time-Variant SRC Kinase Activation Determines Endothelial
Permeability Response. *Cell Chem Biol* **26**, 1081-1094 e1086 (2019).
https://doi.org:10.1016/j.chembiol.2019.04.007

24   Chen, W. *et al.* Atrial natriuretic peptide-mediated inhibition of microcirculatory
endothelial Ca2+ and permeability response to histamine involves cGMP-dependent
protein kinase I and TRPC6 channels. *Arterioscler Thromb Vasc Biol* **33**, 2121-2129
(2013). https://doi.org:10.1161/ATVBAHA.113.001974

25   Shikata, Y., Birukov, K. G., Birukova, A. A., Verin, A. & Garcia, J. G. Involvement of site-
specific FAK phosphorylation in sphingosine-1 phosphate- and thrombin-induced focal
adhesion remodeling: role of Src and GIT. *FASEB J* **17**, 2240-2249 (2003).
https://doi.org:10.1096/fj.03-0198com

26   Zhang, P. *et al.* Polychlorinated biphenyl quinone induces endothelial barrier
dysregulation by setting the cross talk between VE-cadherin, focal adhesion, and MAPK
signaling. *Am J Physiol Heart Circ Physiol* **308**, H1205-1214 (2015).
https://doi.org:10.1152/ajpheart.00005.2015

27   Birukova, A. A., Alekseeva, E., Cokic, I., Turner, C. E. & Birukov, K. G. Cross talk between
paxillin and Rac is critical for mediation of barrier-protective effects by oxidized

phospholipids. *Am J Physiol Lung Cell Mol Physiol* **295**, L593-602 (2008).
https://doi.org:10.1152/ajplung.90257.2008

28   Anastassiadis, T., Deacon, S. W., Devarajan, K., Ma, H. & Peterson, J. R. Comprehensive
assay of kinase catalytic activity reveals features of kinase inhibitor selectivity. *Nat*
Biotechnol **29**, 1039-1045 (2011). https://doi.org:10.1038/nbt.2017

29   Dankwa, S. *et al.* Exploiting polypharmacology to dissect host kinases and kinase
inhibitors that modulate endothelial barrier integrity. *Cell Chem Biol* (2021).
https://doi.org:10.1016/j.chembiol.2021.06.004

30   Turinsky, A. L. *et al.* Navigating the Global Protein-Protein Interaction Landscape Using
iRefWeb. *Methods Mol Biol* **2199**, 191-207 (2021). https://doi.org:10.1007/978-1-0716-
0892-0_12

31   Hornbeck, P. V. *et al.* PhosphoSitePlus, 2014: mutations, PTMs and recalibrations.
Nucleic Acids Res **43**, D512-520 (2015). https://doi.org:10.1093/nar/gku1267

32   van den Eshof, B. L. *et al.* Paradigm of Biased PAR1 (Protease-Activated Receptor-1)
Activation and Inhibition in Endothelial Cells Dissected by Phosphoproteomics.
Arterioscler Thromb Vasc Biol **37**, 1891-1902 (2017).
https://doi.org:10.1161/ATVBAHA.117.309926

33   Vu, T. K., Hung, D. T., Wheaton, V. I. & Coughlin, S. R. Molecular cloning of a functional
thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* **64**,
1057-1068 (1991). https://doi.org:10.1016/0092-8674(91)90261-v

Fig. 1. Workflow for reconstructing a kinase-mediated phosphosignaling network controlling brain endothelial barrier integrity after thrombin stimulation using TPS. HBMECs were treated with thrombin for 0, 5, 15, 30, 60, 120, 180, 240 or 360 minutes. Phosphorylation levels (at activation sites) of 28 proteins (25 kinases and 3 non-kinase proteins) were evaluated by western blot and quantified by fold change from the basal level (media only) after normalization to GAPDH. The interactome network was compiled by combining human PPIs from iRefIndex and kinase-substrate phosphorylation interactions from PhosphoSitePlus. An undirected subnetwork was generated using the Omics Integrator implementation of the Prize-Collecting Steiner Forest algorithm, incorporating the time series phosphorylation data and the interactome network. The subnetwork, kinase-substrate phosphorylation interactions, and time series phosphorylation data were input into the TPS algorithm, with PAR1/F2R as the source node. TPS outputs included 1) the kinase-mediated signaling network (directed and signed) that controls protein phosphorylation in HBMECs after thrombin treatment, and 2) activity of each protein in the subnetwork at the corresponding time points.

Fig. 2. TPS infers common and unique kinases and signaling cascades compared to TREKING models. a Signed edges in the summary network generated by TPS. **b** Comparison of kinase phosphosignaling networks built by TPS and TREKING. The background network in gray includes all kinase-kinase phosphorylation interactions from the kinase-substrate phosphorylation database PhosphoSitePlus. Turquoise nodes are the kinases with time series phosphorylation data used to inform the TPS network; turquoise edges are the kinase-kinase phosphorylation interactions inferred by TPS algorithm. Blue and red nodes are the kinases predicted by KiR to have only barrier-weakening and barrier-strengthening functionality, respectively; purple nodes are the switch kinases having both barrier-weakening and barrier-strengthening functionalities but at different time frames. Blue and red edges are the interactions predicted by TREKING that are associated with only barrier-weakening and barrier-strengthening activity, respectively; purple edges are the interactions associated with both barrier-weakening and barrier-strengthening activities. **c** The combined kinase signaling network inferred by TPS and TREKING methodologies. Kinases and kinase-kinase interactions inferred by TPS or TREKING only are in turquoise or moccasin, respectively; kinases and kinase-kinase interactions inferred by both TPS and TREKING are in sky blue.

Fig. 3. Evaluate the accuracy of model predictions. a Schematic showing the types of

data used for reconstructing and validating phosphosignaling networks built by TPS and

TREKING approaches. **b** Validation on the TPS network using kinase-compound

biochemical data [28] and kinase inhibitor screen data [29]. Left: light sky blue indicates that

TPS correctly predicted the impact of kinase inhibitors on endothelial barrier

permeability, gray indicates incorrect predictions, and white indicates that TPS did not

predict substantial inhibition of the network by the kinase inhibitors and no comparison

was made with the phenotypic data. Middle: number of target kinases (residual activity <

50%) in the TPS network. Right: overall prediction accuracy of TPS compared to random

models on predicting impact of kinase inhibitors on barrier permeability (light sky blue:

correct predictions; gray: incorrect predictions). If the model predicts a kinase inhibitor to

alter barrier permeability and the barrier permeability in the presence of the kinase

inhibitor is different from DMSO-treated control, the model makes correct prediction on

the kinase inhibitor; otherwise, the model makes incorrect prediction on the kinase

inhibitor. **c** Validation on the TREKING network using western blot data. Left: pink

indicates that TREKING correctly predicted the kinase functionality in barrier regulation,

gray indicates incorrect predictions, and white indicates that TREKING did not predict

the barrier function, and no comparison was made with the western blot data. Middle:

kinase connectivity (in/out-degrees of freedom) in the TREKING network. Right: overall

prediction accuracy of TREKING compared to random models on predicting the barrier

activity of kinases (pink: correct predictions; gray: incorrect predictions). If the model

predicts a kinase to regulate barrier permeability and its activity is different from basal

level at the corresponding time point, the model makes correct prediction on the kinase

barrier activity; otherwise, the model makes incorrect prediction on the kinase barrier

activity.

Interactome network
- Protein-protein interactions: iRefIndex
- Kinase-substrate interactions: PhosphoSitePlus

Time series phosphorylation data

Fold change of phosphorylation

10.0
7.5
5.0
2.5
1.0

HSPB1
MAPK1
MAPK3
ABL1
MAP3K14
MAP2K1
MAP2K2
CHUK
CHEK1
AKT1
BRAF
PTK2
RAF1
RPS6KB1
PRKCD
CDH5
MAP3K20
PDPK1
PXN
SRC
MAPK8
MAPK9
MKNK1
MAPKAPK2
MAPK14
MAPK13
MAPK11
MAPK12

0 5 15 30 60 120 180 240 360

Time (min)

Partial model: signed, directed network of known kinase-substrate interactions

PhosphoSitePlus

Omics Integrator
Prize-Collecting Steiner Forest algorithm

Subnetwork connecting phosphorylated proteins to the source nodes

Temporal Pathway Synthesizer ← **Source:** PAR1/F2R

Summary network

5 min   15 min   360 min

...