Biased estimates of phylogenetic branch lengths resulting from the discretised Gamma model of site rate heterogeneity

Luca Ferretti[1,2], Tanya Golubchik[3], Francesco Di Lauro[1,2],
Mahan Ghafari[1,2,4], Julian Villabona-Arenas[5], Katherine E. Atkins[5,6],
Christophe Fraser[1,2], Matthew Hall[1,2,7,*]

[1] Pandemic Sciences Institute, Nuffield Department for Medicine, University of Oxford, Old Road Campus, Oxford, UK

[2] Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, Nuffield Department of Medicine, Old Road Campus, University of Oxford, UK

[3] Sydney Infectious Diseases Institute, School of Medical Sciences, Faculty of Medicine and Health, University of Sydney, Australia

[4] Department of Biology, University of Oxford, Oxford, UK

[5] Centre for Mathematical Modelling of Infectious Diseases and Department of Infectious Disease Epidemiology and Dynamics, Faculty of Epidemiology and Population Health, London School of Hygiene and Tropical Medicine, London, UK.

[6] Centre for Global Health, Usher Institute, University of Edinburgh, UK

[7] Department of Clinical Research, Faculty of Infectious and Tropical Disease, London School of Hygiene and Tropical Medicine, London, UK.

* Corresponding author: matthew.hall@lshtm.ac.uk

April 24, 2025

Abstract

A standard method in phylogenetic reconstruction for representing variation in substitution rates between sites in the genome is the discrete Gamma model (DGM). Relative rates are assumed to be distributed according to a discretised Gamma distribution, where the probabilities that a site is included in each class are equal. Here, we identify a serious bias in the branch lengths of reconstructed phylogenies when the DGM is used, whereby lengths are usually, and often substantially, overestimated, and the magnitude of this effect varies (usually increasing) with the number of sequences in the alignment. We show that the alternative "FreeRate" model, which assumes no parametric distribution and allows the class probabilities to vary, is not subject to the issue. We further establish that the reason for the behaviour is the equal class probabilities, not the discretisation itself. We also explore the mathematical reasons for the phenomenon, showing that the effect is determined by the bias in the mean relative rate of evolution amongst the observed sites and that branch lengths will be overestimated where there is a long-tail of fast-evolving sites in the true rate distribution, the usual situation in real datasets. We recommend that the DGM be retired from general use. While FreeRate is an immediately available replacement, it is highly parameterised and known to be difficult to fit, and thus there is scope for innovation in rate heterogeneity models.

Introduction

It was recognised early in the development of molecular phylogenetics that it is unrealistic to assume that mutation rates are invariant between genomic sites (Fitch and Margoliash 1967; Uzzell and Corbin 1971). One key reason for the existence of variation is the degeneracy of the genetic code, where a random mutation is least likely to be synonymous at the first codon position and most likely at the third, making mutations at the latter more likely to be neutral (Fitch 1986; Wolfe et al. 1989). In more general terms, the functional and structural role of

sites in the genome vary very widely, and thus the evolutionary pressures on each, and hence the probability of observing a mutation there, cannot help but differ considerably (Yang 1996).

By the 1990s it was clear that methods were needed to account for between-site rate variation, and it became standard to use a gamma distribution to model it. An overall mutation rate μ is modified at each site in the genome by a draw from a gamma distribution with shape α and scale $1/\alpha$. (The mean of the distribution is thus 1 and the mean rate over the entire genome remains μ.) This was first employed in distance matrix calculations (Nei and Gojobori 1986), before Yang (1993) integrated it into maximum-likelihood algorithms. A year later, due to the computational limitations of research in the early 1990s, Yang (1994) introduced a pair of approximations. The first of these, the "discrete gamma model" (DGM), approximates the gamma distribution by dividing its support into a finite number (four was suggested) of contiguous regions of equal probability. The mean (or median) value of the gamma probability density function (PDF) on each region is calculated, and each site in the genome takes each of those values with equal probability. In subsequent years, this approximation became the absolutely standard approach to rate heterogeneity in almost all major software packages for maximum likelihood and Bayesian phylogenetic reconstruction, including PhyML (Guindon et al. 2010), RAxML (Stamatakis 2014), IQ-TREE (Minh et al. 2020), MrBayes (Ronquist et al. 2012), and both versions of BEAST (Suchard et al. 2018; Bouckaert et al. 2019).

It might well be asked why, if one was to assume a discrete probability distribution for between-site rate variation, it needed to be assumed that either it took a gamma form or that every rate had an equal probability, and both assumptions can be relaxed. Yang (1995) outlined a general non-parametric approach, in which there are k rates $\rho_1, \ldots, \rho_k$ and the probability that a site evolves according to each ρ_i is w_i. Because $\sum_i w_i = 1$ and the mean relative rate across all sites is always assumed to be 1 (i.e. $\sum_i w_i \rho_i = 1$) this model has $2(k-1)$ free parameters. In contrast, the DGM has only one, the shape of the gamma distribution itself, regardless of how many categories are chosen. This major increase in the number of parameters causes computational problems that continue to this day (Morel et al. 2021).

The non-parametric version could be viewed as a type of mixture model, and the use of mixture models for between-site rate variation has been a subject of continued development (Mayrose, Friedman, et al. 2005; Huelsenbeck and Suchard 2007; Evans and Sullivan 2012; Wu

et al. 2013). The version of Yang (1995) was belatedly christened "FreeRate" by Soubrier et al. (2012) and it is implemented under that name in the current versions of PhyML (Guindon et al. 2010), RAxML-NG (Kozlov et al. 2019), and IQ-TREE (Minh et al. 2020).

Previous critiques of the DGM have tended to demonstrate that a mixture model is superior if the underlying rate distribution is not well-described by a gamma distribution (Mayrose, Friedman, et al. 2005; Soubrier et al. 2012). Here we demonstrate a more fundamental problem that occurs even if true rates are actually drawn from a continuous gamma distribution, and which has profound implications for phylogenetic reconstruction. We demonstrate that branch lengths estimated using a DGM are biased, sometimes considerably so, and usually in the direction of overestimation. We further show that this effect increases with the number of sequences in the alignment. This phenomenon does not occur when FreeRate is used instead, and are a result of the equal size of the categories used in the DGM (rather than the discretisation itself or the choice of a gamma distribution). We demonstrate the problem using real and simulated data, and then provide an analytic demonstration of why it happens. We recommend, as a result, that the DGM be retired from use and that, if FreeRate is too highly-parameterised for general use, work is needed to produce a robust alternative.

Materials and Methods

Site rate heterogeneity models

Models of rate heterogeneity assume that each site D_i in the genome $\mathbf{D}$ is assigned a variable rate of evolution, $\rho_i \in \mathbb{R}^+$. For the category of model we are dealing with, the ρ_i are assumed to be independent draws from a probability distribution H. This may be discrete or continuous. The rate at each D_i is then taken to be an independent draw from the distribution whose probability mass or density function is f_H.

This distribution is of a set of *relative* rates, and describes the heterogeneity among sites independently of the absolute rate of evolution. Thus its mean is taken to be 1, i.e.

$$\sum_{\rho \in \mathbf{R}} \rho \, f_H(\rho) = 1 \tag{1}$$

for a discrete H taking values in a finite set $\mathbf{R}$ of positive reals, or

$$\int_0^\infty \rho f_H(\rho)d\rho = 1 \tag{2}$$

for a continuous H. When H is a gamma distribution, the requirement of a mean of 1 allows the distribution to be defined only by its shape k; the scale must then be $\frac{1}{k}$.

A model of nucleotide substitution with a rate heterogeneity component has two random outcomes: the assignment of rates to sites, and the allelic states along the tree at each site. This latter, at the tips, produces the alignment, the data source in phylogenetic inference. The former are not directly observed and are usually integrated over in reconstruction algorithms (Yang 1994).

Suppose H is discrete, the parameters of the rate heterogeneity model are P, and those of the nucleotide substitution model are M. The likelihood of a phylogeny $\mathbf{T}$ given a multiple sequence alignment $\mathbf{D}$ with n_{sites} positions is:

$$\mathrm{L}(\mathbf{T}, P, M|\mathbf{D}) = \prod_{i=1}^{n_{\text{sites}}} \sum_{\rho \in \mathbf{R}} \mathbb{P}(D_i|\mathbf{T}, \rho, M)\mathbb{P}(\rho|P) \tag{3}$$

where D_i denotes the content of $\mathbf{D}$ at site i. Maximisation of this likelihood leads to maximum likelihood estimates for all parameters, denoted here by $\hat{\mathbf{T}}$, $\hat{P}$ and $\hat{M}$. Henceforth we refer to the terms $\mathbb{P}(\rho|P)$ as *weights*; they must sum to 1 over $\rho \in \mathbf{R}$.

The classic DGM belongs to a larger class of "equal-probability models", where sites are assigned to one of n_{cats} rate categories where n_{cats} is finite. The important property here is that these categories are equiprobable, i.e. there is a fixed probability $1/n_{\text{cats}}$ that a site belong to any one of them. Equivalently, in expression (3), the weights are invariant.

Each category is assigned a specific relative rate. In the case of the DGM, the rates are chosen as either the mean or the median (Yang 1994) of the rates in the corresponding n_{cats}-ile of a Gamma distribution. In the case of the mean, to which we give the full treatment here (the behaviour observed in this paper is unaffected by choosing the median instead) this is given by:

$$\tilde{\rho}^{(i)} = n_{\text{cats}} \int_{q_\Gamma\left((i-1)/n_{\text{cats}}|k,\frac{1}{k}\right)}^{q_\Gamma\left(i/n_{\text{cats}}|k,\frac{1}{k}\right)} x\, f_\Gamma\left(x|k, \frac{1}{k}\right) dx, \quad i \in \{1, \ldots, n_{\text{cats}}\} \tag{4}$$

where f_Γ and q_Γ are respectively the PDF and quantile function of a Gamma distribution with shape k and scale $\frac{1}{k}$. Note that the mean of the rates among categories is $\sum_i \tilde{\rho}^{(i)}/n_{\mathrm{cats}} = 1$.

HIV phylogenetic reconstruction

For real data, we took an alignment of 1000 full-genome subtype C HIV sequences from the PopART Phylogenetics study (Hall et al. 2024; Hayes et al. 2019), available via the PANGEA-HIV consortium (Abeler-Dörner et al. 2019). Sequences can be downloaded at `http://github.com/PANGEA-HIV/PANGEA-Sequences`. Positions where more than 10% of sequences had an N or gap character were removed, and the whole alignment consists of 7854 base pairs. This alignment was progressively downsampled by subtracting 25 random sequences from it, to give a total of 39 alignments of between 50 and 1000 sequences; the 50 in the former are shared between all of them. Phylogenies for this alignment were reconstructed using IQ-TREE 2.2.0 (Minh et al. 2020) specifying GTR+F+G4 (4-category DGM) and GTR+F+R4 (4-category FreeRate) as substitution models.

As a senstivity analysis we also performed this exercise using 6-category, 8-category and 10-category DGMs (GTR+F+G6, GTR+F+G8 and GTR+F+G10) as well as the 4-category DGM with invariant sites (GTR+F+I+G4), using the same set of alignments as above, and also repeating it with three other alignments of subtype C PANGEA sequences.

Simulations

A simulated HIV transmission tree was generated using PopART-IBM (Pickles et al. 2021). The result was a completely sampled epidemic of 1771 cases. This was transformed to a timed phylogeny using VirusTreeSimulator (Ratmann et al. 2017), and that, in turn, was transformed to a tree with branch lengths in units of substitutions per site by simply assuming a strict molecular clock with a substitution rate of 0.001 substitutions per site per year (i.e. by multiplying each branch length by 0.001). This was then used to generate sequences with AliSim (Ly-Trong et al. 2022), with 10,000 bases, the GTR substitution model with equal base frequencies. and as a site heterogeneity model, one of the following two options:

1. A continuous gamma distribution with a shape parameter of 0.5 and a scale parameter of 2.

2. A 4-category discretised version of the same gamma distribution, with each category taking up 1/4 of the probability space and its value on that quarter being the mean of the PDF in that interval.

Option 1 is *true* gamma-distributed rate heterogeneity, while in option 2 sequences are generated using the DGM itself. The alignments generated were downsampled to 500 sequences and then further to 100. (The contents of the two alignments of 500 sequences correspond to the same 500 tips of the input tree, as do the alignments of 100 sequences.) Phylogenetic reconstruction, again using GTR+F+R4 and GTR+F+G4, was performed as above.

Results

Branch lengths in HIV phylogenies increase with sample size with the DGM, but not FreeRate

In figure 1, we plot, for the downsampled alignments up to size 250, patristic distances between the tips representing the 50 shared sequences, in the reconstruction using just those sequences (x-axis) and in that using larger alignments (y-axis). A clear effect whereby the addition of more sequences to the alignment has a scaling effect on all branch lengths, when the phylogeny is reconstructed using DGM, can be observed. This is absent for FreeRate.

Figure 2 gives the gradients of regression lines for the comparisons, up to 1000 tips. A rapid increase in the gradient when moving from from 50 to 125 tips indicates corresponding to a swift increase in the amount by which branch lengths are overestimated. This effect does appear to eventually saturate after 125 tips, after which the increase in gradient becomes more gradual.

Perhaps contrary to intuition, increasing the number of rate categories in the DGM actually exacerbates the problem. In Fig. **??**, we plot gradients for models with a varying number of categories for four different 1000-sequence alignments; replicate 1 is that of figure 2. A 6-category DGM results in considerably more branch length expansion than a 4-category one. 8- and 10-category versions show more expansion still, although the progression is not straightforward and the 8-category version is often worse than the 10-category. On the other hand, adding an invariant sites category does mitigate the issue. This effect is, however, slight and the problem

Figure 1: Patristic distances between the tips representing 50 subtype C HIV sequences in phylogenetic reconstructions. The x-axis gives these when the alignment consists of just those 50 sequences, while the y-axis are the same from a larger alignment, with the total number of sequences increasing in the facets from left to right. Top row: reconstruction uses 4-category DGM. Bottom row: reconstruction uses 4-category FreeRate.

Figure 2: The gradient of a linear regression line plotting patristic distances in phylogenies generated using the DGM on a 50-sequence alignment and in phylogenies from larger alignments. The x-axis is the size of the larger alignment. The error bars give the 95% confidence interval of the regression coefficient. In the 50-tip case the tree on both axes is the same one.

remains very much in evidence. Also of note is the considerable inconsistency in the behaviour of the expansion between replicates for larger number of categories; while the 4-category DGM converges to a gradient of around 1.5 in all four replicates (and whether or not an invariant sites category is present), this kind of consistent behaviour is not in evidence for more complex models. In some cases (most clearly replicate 2 for the 6-category DGM), the largest alignments show less expansion than some of the smaller ones.

Biased branch length estimates as the result of equiprobable rate categories

In figure 4, we plot patristic distances between the tips in the 100-sequence reconstructions of the simulated datasets to those between the same 100 tips in the 500-sequence reconstructions. The only situation in which branch lengths increase with sample size is where the generating model is the continuous gamma distribution and the reconstruction model is the DGM.

While figure 4, top left, suggests that DGM estimates of branch lengths are unreliable for a continuous input distribution, it is still a comparison of outputs and does not indicate the presence or direction of any bias. In figure 5 we plot patristic distances in the true tree

Figure 3: The gradient of a linear regression line plotting patristic distances in phylogenies generated using the DGM on a 50-sequence alignment and in phylogenies from larger alignments. The x-axis is the size of the larger alignment. The error bars give the 95% confidence interval of the regression coefficient. In the 50-tip case the tree on both axes is the same one. Points are coloured by the version of the DGM used for reconstruction. The four facets correspond to different initial 1000-sequence alignments which were progressively downsampled, all consisting of subtype C HIV sequences from PANGEA-HIV.

Figure 4: Patristic distances between the tips representing 100 simulated HIV sequences in phyloenetic reconstructions. The x-axis gives these when the alignment consists of just those 100 sequences, while the y-axis are the same from a larger alignment of 500. Vertical facets represent the rate heterogenity model of the simulation (continuous gamma distribution or DGM) and horizontal the reconstruction model in IQ-TREE (DGM or FreeRate).

Figure 5: Comparison of patristic distances between 100 tips in the input phylogeny to a simulated alignment (x-axis) and estimates from phylogenetic reconstructions (y-axis). 400 other sequences were added to the alignments used for those reconstructions. Estimates from DGM reconstructions are in orange and FreeRate in green.

against those obtained from FreeRate and DGM reconstructions, when the generating model is the continuous version. Branch lengths as reconstructed by FreeRate are unbiased; those reconstructed using the DGM are systematically overestimated.

To summarise these results, FreeRate provides accurate estimates regardless of the generating model. The DGM gives good estimates of branch lengths when the generating model is itself discretised, but an obvious bias occurs when attempting to fit it to sequences generated using the true, continuous gamma model.

This simulation notably lacks any process mimicking misalignment, recombination, sequencing error, selection, or any other complex processes that might drive the phenomenon of branch length bias. As none of these are present, they cannot explain the results. The bias is due simply to the generation of a model using a continuous distribution and reconstruction using the DGM.

As FreeRate is correctly estimating branch lengths, it would seem natural that it is also correctly estimating the form of the input distribution even if its nature as a gamma distribution is not specified, and this, indeed, is the case. The fitted FreeRate distribution consists of the weights for the (four) rate classes and the estimated rate in each. In figure 6 we plot those estimated rates on their quantiles of the unit interval, as calculated in the analysis of the 500-sequence alignment generated using the continuous model, against the actual mean of the generating gamma distribution on the intervals defined by those weights. The match is

Figure 6: The actual quantile plot function of a gamma distribution with shape parameter 0.5 and scale parameter 2 (black) compared with the fitted FreeRate distribution from a simulated 500-sequence alignment (green) and the result of calculating the mean value of the true distribution on the weight boundaries estimated using FreeRate (red).

almost exact, and it is plain that FreeRate can successfully estimate a version of the input rate distribution that is discretised in a way that does not bias branch length estimates. No standard maximum likelihood phylogenetics package currently allows the user to specify, as a rate heterogeneity model, a discretised gamma distribution whose weights are variable and estimated by the algorithm. However, it would naturally follow that if FreeRate is able to mimic the form of one without being subject to branch length bias in the output, such a "gamma + variable weights" option in a package would also be able to. The problem thus cannot be the discretisation, and must instead lie with the equal weights.

We conclude that reconstruction of phylogenies using the DGM from alignments whose true rate heterogeneity model is a continuous gamma distribution are systematically biased, and that the reason is the constraint that the probabilities of a site belonging to each category are equal.

Branch expansion is a consequence of model mis-specification and a tail of rapidly evolving sites

In the appendix to this paper, we delve further into the mathematics of the phenomenon. We summarise those results here. Under "Likelihood expression for star-like phylogenies" we

Figure 7: Ratios of the maximum likelihood estimate of the (constant) phylogenetic branch length in a star-like phylogeny to its true length, for different sizes of alignment (x-axis) and lengths of the true branch (substitutions per site, facets). Calculations are the result of simulating 100 alignments of 5,000 sites each with a gamma shape parameter of 0.5. The central line connects the mean maximum likeihood estimate across those 100 alignments, with the error bars giving the 95% symmetric credible interval.

give an expression for the likelihood of branch lengths in an ultrametric starlike tree (a tree with a single internal node and whose branch lengths are all equal), after some simplifying assumptions. Maximising this expression allows us to estimate the magnitude of branch length overestimation for various scenarios of true branch length and alignment size (figure 7). We see that a longer branch length is initially associated with a faster increase in maximum likelihood branch length as the nynber of tree tips increases, but that there is also an asymptotic limit to the bias which is obtained sooner, and at a lower value.

That the DGM is a misspecified model of the true distribution of rates would seem clear for real data, as it was always intended as a simplified method (Yang 1994). However, the bias exists even if the generating rate distribution is a continuous gamma distribution, and, as shown above ("Branch lengths in HIV phylogenies increase with sample size with the DGM, but not FreeRate") it clearly also occurs in real datasets.

In the section entitled "The impact of the equal-weights discretisation in the two-category and n-category case" we give an expression for the bias in branch lengths when a FreeRate distribution with equal weights is assumed but the true model of evolution has unequal weights. We show that branch lengths are overestimated when there is a negative correlation between

the weight of each category w and its relative rate ρ in the true model of evolution. More precisely, if the true branch length is λ and the estimate $\hat{\lambda}$, and $\mathbf{P}$ and $\mathbf{W}$ are the vectors of rates and weights respectively, the relative bias is given by:

$$\frac{\lambda - \lambda^{\text{true,local}}}{\lambda^{\text{true,local}}} = -\text{Cov}\left[\mathbf{W}/\overline{\mathbf{W}}, \mathbf{P}\right] \tag{5}$$

If rates and weights are positively correlated, i.e. larger rates are most common, this expression is negative and branch lengths are underestimated. However, this will be rare in the real world as most inferred rate distributions have a mode of less than 1 (Jia et al. 2014; Silvestro et al. 2024). It would be much more common to see a negative correlation between rate and weight, leading to overestimation of branch lengths.

Discussion

This is, to our knowledge, the first time this bias has been described in the literature, despite the almost thirty-year history of the DGM. Brown et al. (2010) attributed a similar phenomenon, observed in the Bayesian realm only, to either poor Markov Chain Monte Carlo mixing or excessive posterior weight to trees with long branches. On the maximum likelihood side, Nguyen et al. (2018) found that that problematic heuristic optimisation routines in the major packages were responsible for misestimation of the shape parameter of the DGM and tree length, as well as the proportion of invariant sites. Mannino et al. (2020) identified a different problem with the equal-probabilities specification, in that it places a hard upper bound on the variance of the rate distribution, but did not explore branch lengths. Schwartz and Mueller (2010) explored bias in branch lengths in relatively small trees using both Bayesian and maximum likelihood methods. They observed a tendency for underestimation in both cases, along with some overestimation of longer branches for maximum likelihood. Their focus was in comparing estimates for individual branches to their true lengths, rather than comparing reconstruction results as we do here, and while they did employ the DGM, the datasets used may have been too small (in terms of numbers of taxa) for the phenomenon we observe here to have become obvious. The phenomenon we identify here is broader than all of these, with major implications for the field.

The equal weights specification is a highly problematic choice, which causes systematic bias in the estimation of phylogenetic branch lengths, usually in the direction of overestimating them. This effect increases with sample size but can still be significant for small alignments if branch lengths are long enough. It should be emphasised that nothing about this phenomenon is biological. It is not specific to any organism, although its magnitude will depend on quantities such as the branch lengths in a typical phylogeny. No analysis that has previously used the DGM should be regarded as exempt from a re-exploration in light of this finding. It is not even specific to biological organisms at all; studies using phylogenetic reconstruction in the realm of linguistics have also used the DGM in the past (e.g. Lee and Hasegawa (2011)). The effect is more pronounced when mutation rates are fast (Fig. 7), and as such it is perhaps not surprising that we first observed it when analysing the genomes of RNA viruses. However, given the huge body of phylogenetic research previously conducted using the DGM and the simplifying assumptions we needed to make in exploring the mathematics of the issue, we consider it advisable for researchers studying even much more slowly-evolving organisms to investigate if their work would be affected.

An obvious line of further enquiry, which we do not explore here but is the subject of intensive ongoing work, is the effect of this bias on time tree inference (Sagulenko et al. 2018; Suchard et al. 2018; Bouckaert et al. 2019). In a time tree, a molecular clock model is employed so that branch lengths are in units of calendar time and tips appear at points on the timeline corresponding to the dates upon which the corresponding samples were taken. Longer branches in the underlying divergence tree will not necessarily correspond to longer branches in a time tree, because a model can instead increase the rate of evolution dictated by the molecular clock. We can compensate for the allocation of a greater amount of evolution to a particular phylogenetic branch by, in turn, increasing our estimate for how quickly that evolution happened. However, time tree models are often complex, and the extent to which the effects of this bias affect branch lengths and the extent to which they affect clock rates requires urgent investigation across a whole range of organisms.

Increasing the number of categories in the DGM does not resolve the problem and in fact makes it worse (Fig. 3). On the other hand, adding an invariant sites category appears to be slightly mitigating, at least for our HIV alignments. This is perhaps because the size of that

category is estimated and as such the model does, in a limited way, enable variable weights. Nevertheless, the reduction is modest and we would not recommend it as a long-term solution. For six or more categories, the behaviour of the expansion can be strikingly different when applied to different alignments. Identifying the reasons for this is beyond the scope of this work, but we would argue that possible limitations of the heuristic maximum-likelihood routine would need to be excluded as a cause before a mathematical explanation was sought.

FreeRate is the most immediate option for an alternative to the DGM that is not subject to the problem of branch length bias. It is implemented and documented in state of the art maximum likelihood phylogenetics packages (Stamatakis 2014; Minh et al. 2020) although much less common in Bayesian packages. It is, however, a model with many more free parameters than the DGM and it is known to be difficult to fit (Morel et al. 2021). We stress that our objective in this paper is to inform the research community of a particular issue with the DGM, not to advocate for FreeRate as a gold standard alternative. A potential alternative from the existing literature, proposed by Felsenstein (2001) is to use generalised Gauss-Laguerre quadrature to approximate the integral of the likelihood over the rate categories. We would also suggest that time is right for new innovations in site rate heterogeneity models, after so many years during which the DGM has been used by default. This process may be underway, as examples taking a machine learning approach are now starting to emerge (Silvestro et al. 2024).

In summary, we would argue the problem that we identify here regarding the DGM is one that should preclude its future use. For our HIV examples we show that branch lengths are highly influenced by the size of the alignment, a situation that cannot be compatible with robust inference. FreeRate provides one option, but has its own disadvantages. The impact of this problem on phylogenetic dating is less clear and work is needed to establish the extent to which estimates may be changed in the light of this finding.

Funding Acknowledgment

The computational aspects of this research were supported by the Wellcome Trust Core Award Grant Number 203141/Z/16/Z and the NIHR Oxford BRC. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

This study was supported by the Bill & Melinda Gates Foundation through the PANGEA

consortium (OPP1175094 and OPP1084362) and by the Wellcome Trust through the ARTIC Network Collaboration Award (206298/Z/17/Z).

CJVA and KEA were funded by an ERC Starting Grant (award number 757688) awarded to KEA.

TG is supported by an Investigator Grant (GNT2025445) from the National Health and Medical Research Council, Australia (NHMRC).

Acknowledgment

We thank Sergei Kosakovsky Pond and Andrew Rambaut for helpful suggestions.

References

Abeler-Dörner L., Grabowski M.K., Rambaut A., Pillay D., Fraser C., and PANGEA consortium. 2019. PANGEA-HIV 2: Phylogenetics And Networks for Generalised Epidemics in Africa. *Curr. Opin. HIV AIDS* 14:173–180.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchêne S., Fourment M., Gavryushkina A., Heled J., Jones G., Kühnert D., De Maio N., Matschiner M., Mendes F.K., Müller N.F., Ogilvie H.A., Plessis L. du, Popinga A., Rambaut A., Rasmussen D., Siveroni I., Suchard M.A., Wu C.-H., Xie D., Zhang C., Stadler T., and Drummond A.J. 2019. BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLOS Comput. Biol.* 15:e1006650.

Brown J.M., Hedtke S.M., Lemmon A.R., and Lemmon E.M. 2010. When trees grow too long: investigating the causes of highly inaccurate bayesian branch-length estimates. *Syst. Biol.* 59:145–161.

Darriba D., Posada D., Kozlov A.M., Stamatakis A., Morel B., and Flouri T. 2020. ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Mol. Biol. Evol.* 37:291–294.

Evans J. and Sullivan J. 2012. Generalized mixture models for molecular phylogenetic estimation. *Syst. Biol.* 61:12–21.

Felsenstein J. 1973. Maximum likelihood and minimum-steps methods for estimating evolu-
tionary trees from data on discrete characters. *Syst. Biol.* 22:240–249.

Felsenstein J. 2001. Taking variation of evolutionary rates between sites into account in inferring
phylogenies. *J. Mol. Evol.* 53:447–455.

Fitch W.M. 1986. The estimate of total nucleotide substitutions from pairwise differences is
biased. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 312:317–324.

Fitch W.M. and Margoliash E. 1967. A method for estimating the number of invariant amino
acid coding positions in a gene using cytochrome c as a model case. *Biochem. Genet.* 1:65–
71.

Guindon S., Dufayard J.-F., Lefort V., Anisimova M., Hordijk W., and Gascuel O. 2010.
New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. *Syst. Biol.* 59:307–321.

Hall M., Golubchik T., Bonsall D., Abeler-Dörner L., Limbada M., Kosloff B., Schaap A.,
Cesare M. de, MacIntyre-Cockett G., Otecko N., Probert W., Ratmann O., Bulas Cruz A.,
Piwowar-Manning E., Burns D.N., Cohen M.S., Donnell D.J., Eshleman S.H., Simwinga M.,
Fidler S., Hayes R., Ayles H., Fraser C., HPTN 071 (PopART) Phylogenetics protocol team,
and PANGEA consortium. 2024. Demographics of sources of HIV-1 transmission in Zambia:
a molecular epidemiology analysis in the HPTN 071 PopART study. *Lancet Microbe* 5:e62–
e71.

Hayes R.J., Donnell D., Floyd S., Mandla N., Bwalya J., Sabapathy K., Yang B., Phiri M.,
Schaap A., Eshleman S.H., Piwowar-Manning E., Kosloff B., James A., Skalland T., Wilson
E., Emel L., Macleod D., Dunbar R., Simwinga M., Makola N., Bond V., Hoddinott G.,
Moore A., Griffith S., Deshmane Sista N., Vermund S.H., El-Sadr W., Burns D.N., Harg-
reaves J.R., Hauck K., Fraser C., Shanaube K., Bock P., Beyers N., Ayles H., Fidler S., and
HPTN 071 (PopART) Study Team. 2019. Effect of universal testing and treatment on HIV
incidence - HPTN 071 (PopART). *N. Engl. J. Med.* 381:207–218.

Huelsenbeck J.P. and Suchard M.A. 2007. A nonparametric method for accommodating and
testing across-site rate variation. *Syst. Biol.* 56:975–987.

Jia F., Lo N., and Ho S.Y.W. 2014. The impact of modelling rate heterogeneity among sites on phylogenetic estimates of intraspecific evolutionary rates and timescales. *PLOS One* 9:e95722.

Kozlov A.M., Darriba D., Flouri T., Morel B., and Stamatakis A. 2019. RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35:4453–4455.

Lee S. and Hasegawa T. 2011. Bayesian phylogenetic analysis supports an agricultural origin of Japonic languages. *Proc. R. Soc. London. B Biol. Sci.* 278:3662–3669.

Mannino F., Wisotsky S., Kosakovsky Pond S.L., and Muse S.V. 2020. Equiprobable discrete models of site-specific substitution rates underestimate the extent of rate variability. *PLOS One* 15:e0229493.

Mayrose I., Friedman N., and Pupko T. 2005. A gamma mixture model better accounts for among site rate heterogeneity. *Bioinformatics* 21 Suppl 2:ii151–8.

Mayrose I., Graur D., Ben-Tal N., and Pupko T. 2004. Comparison of site-specific rate-inference methods for protein sequences: empirical Bayesian methods are superior. *Mol. Biol. Evol.* 21:1781–1791.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., Haeseler A. von, and Lanfear R. 2020. IQ-TREE 2: New models and efficient methods for phylogenetic inference in the genomic era. *Mol. Biol. Evol.* 37:1530–1534.

Morel B., Barbera P., Czech L., Bettisworth B., Hübner L., Lutteropp S., Serdari D., Kostaki E.-G., Mamais I., Kozlov A.M., Pavlidis P., Paraskevis D., and Stamatakis A. 2021. Phylogenetic analysis of SARS-CoV-2 data is difficult. *Mol. Biol. Evol.* 38:1777–1791.

Nei M. and Gojobori T. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* 3:418–426.

Nguyen L.-T., Haeseler A. von, and Minh B.Q. 2018. Complex models of sequence evolution require accurate estimators as exemplified with the invariable site plus gamma model. *Syst. Biol.* 67:552–558.

Pickles M., Cori A., Probert W.J.M., Sauter R., Hinch R., Fidler S., Ayles H., Bock P., Donnell D., Wilson E., Piwowar-Manning E., Floyd S., Hayes R.J., Fraser C., and HPTN 071 (PopART) Study Team. 2021. PopART-IBM, a highly efficient stochastic individual-

based simulation model of generalised HIV epidemics developed in the context of the HPTN 071 (PopART) trial. *PLOS Comput. Biol.* 17:e1009301.

Ratmann O., Hodcroft E.B., Pickles M., Cori A., Hall M., Lycett S., Colijn C., Dearlove B., Didelot X., Frost S., Hossain A.S.M.M., Joy J.B., Kendall M., Kühnert D., Leventhal G.E., Liang R., Plazzotta G., Poon A.F.Y., Rasmussen D.A., Stadler T., Volz E., Weis C., Leigh Brown A.J., Fraser C., and PANGEA-HIV Consortium. 2017. Phylogenetic Tools for Generalized HIV-1 Epidemics: Findings from the PANGEA-HIV Methods Comparison. *Mol. Biol. Evol.* 34:185–203.

Ronquist F., Teslenko M., Mark P.v.d., Ayres D.L., Darling A., Höhna S., Larget B., Liu L., Suchard M.A., and Huelsenbeck J.P. 2012. MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61:539–542.

Sagulenko P., Puller V., and Neher R.A. 2018. TreeTime: Maximum-likelihood phylodynamic analysis. *Virus Evol.* 4:vex042.

Schwartz R.S. and Mueller R.L. 2010. Branch length estimation and divergence dating: estimates of error in Bayesian and maximum likelihood frameworks. *BMC Evol. Biol.* 10:5.

Silvestro D., Latrille T., and Salamin N. 2024. Toward a semi-supervised learning approach to phylogenetic estimation. *Syst. Biol.* 73:789–806.

Soubrier J., Steel M., Lee M.S.Y., Der Sarkissian C., Guindon S., Ho S.Y.W., and Cooper A. 2012. The influence of rate heterogeneity among sites on the time dependence of molecular rates. *Mol. Biol. Evol.* 29:3345–3358.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313.

Suchard M.A., Lemey P., Baele G., Ayres D.L., Drummond A.J., and Rambaut A. 2018. Bayesian phylogenetic and phylodynamic data integration using BEAST 1.10. *Virus Evol.* 4:vey016.

Ly-Trong N., Naser-Khdour S., Lanfear R., and Minh B.Q. 2022. AliSim: a fast and versatile phylogenetic sequence simulator for the genomic era. *Mol. Biol. Evol.* 39:msac092.

Uzzell T. and Corbin K.W. 1971. Fitting discrete probability distributions to evolutionary events. *Science* 172:1089–1096.

Wolfe K.H., Sharp P.M., and Li W.H. 1989. Mutation rates differ among regions of the mammalian genome. *Nature* 337:283–285.

Wu C.-H., Suchard M.A., and Drummond A.J. 2013. Bayesian selection of nucleotide substitution models and their site assignments. *Mol. Biol. Evol.* 30:669–688.

Yang Z. 1993. Maximum-likelihood estimation of phylogeny from DNA sequences when substitution rates differ over sites. *Mol. Biol. Evol.* 10:1396–1401.

Yang Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *J. Mol. Evol.* 39:306–314.

Yang Z. 1995. A space-time process model for the evolution of DNA sequences. *Genetics* 139:993–1005.

Yang Z. 1996. Among-site rate variation and its impact on phylogenetic analyses. *Trends Ecol. Evol.* 11:367–372.

Appendix

Back to basics

Statistical phylogenetic inference, whether maximum likelihood or Bayesian, has as its core the likelihood function (Felsenstein 1973):

$$\mathrm{L}(\mathbf{T}, P, M, |\mathbf{D}) \equiv \mathbb{P}(\mathbf{D}|\mathbf{T}, P, M)$$

where $\mathbf{T}$ is the tree, $\mathbf{D}$ the sequence alignment (that is, the data), P the parameters of the rate heterogeneity model and M the parameters of the nucleotide substitution model. Implicit in this formulation is that $\mathbf{T}$ exists independently of $\mathbf{D}$; there is an underlying "true" tree, with branch lengths in substitutions per site, even if no set of sites is specified. (This is perhaps most easily recognised in the scenario of generating simulated sequences, as packages generally take a fixed tree as input and then provide an alignment of any desired length.)

A component of M is an overall rate of evolution μ, such that the expected number of mutations at a single position during an "evolutionary time period" of length t is μt. As the units of "evolutionary time" are arbitrary, one can actually assume $\mu = 1$, as is common in phylogenetics packages.

480 **D** must have a finite length; suppose this is n_{sites}. P is a vector of positive reals $(\rho_1, \ldots, \rho_{n_{\text{sites}}})$
481 providing a scale factor for μ at each site D_i of **D**. The likelihood is then calculated as:

$$482 \quad \mathrm{L}(\mathbf{T}, M, \rho | \mathbf{D}) = \prod_{i=1}^{l} \mathbb{P}(D_i | \mathbf{T}, M, \rho_i)$$
483

If the expected number of mutations in t is to remain t then we need $\sum_{i=1}^{l} \rho_i / n_{\text{sites}} = 1$. To
write down an exact analytical expression for this likelihood requires a large number of choices
regarding how the mutational process is assumed to behave. For example, one must choose
how many of the relative rates of different types of base substitution are taken to be the same
(Darriba et al. 2020), and how the ρ_i are distributed.

Maximum likelihood phylegenetics provides an estimate $\hat{\mathbf{T}}$ for $\mathbf{T}$, as well as $\hat{P}$ for P and $\hat{M}$
for M.

Simplifying assumptions

Let us simplify **T** considerably by assuming the following:

1. **T** is star-like; all tips are direct children of the root, which is a polytomy of order n_{tips}.

2. All branches of **T** have the same length λ. The total length of the branches of **T** is $n_{\text{tips}}\lambda$.

3. The alignment **D** consists of n_{sites} sites which take only two allelic states (0 and 1). The
substitution model, before rate heterogeneity is considered, is simply transition from 0
to 1 or vice versa at an instantaneous rate of 1, and nucleotide frequencies are equal.
Rate heterogeneity parameters ρ_i are drawn from an unspecified probability distribution
H with support on the positive reals and a mean of 1.

4. The root state of the phylogeny is known to be 0 at every site.

5. The topology of **T** and the form and parameters of the substitution model and rate
heterogeneity model are known and need not be estimated.

As the expected number of mutations at site i along a branch of length λ is $\lambda\rho_i$, the expected
number of mutations across all sites is $\sum_i \lambda\rho_i$ and the expected number of substitutions per
site is $\sum_i \lambda\rho_i / n_{\text{sites}} = \lambda$.

Likelihood expression for star-like phylogenies

We want to obtain an estimate $\hat{\lambda}$ for λ using the observed alignment $\mathbf{D}$. Let $K_i \leq n_{\text{tips}}$ be the number of sequences with an observed mutation (i.e. a 1) at the ith site. As the overall mutation rate μ is 1, the mutation rate at site i is just ρ_i. The Q matrix of the continuous time Markov chain at that site is $\begin{pmatrix} -\rho_i & \rho_i \\ \rho_i & -\rho_i \end{pmatrix}$, which has eigenvectors $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ for its eigenvalues 0 and $-2\rho_i$. The matrix of eigenvectors is $\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ with inverse $\begin{pmatrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{pmatrix}$. Then

$$
\begin{pmatrix} \mathbb{P}_{00}(\lambda|\rho_i) \\ \mathbb{P}_{01}(\lambda|\rho_i) \end{pmatrix} = \frac{1}{2}e^{0\lambda}\begin{pmatrix} 1 \\ 1 \end{pmatrix} + \frac{1}{2}e^{-2\rho_i\lambda}\begin{pmatrix} 1 \\ -1 \end{pmatrix}
$$

$$
= \begin{pmatrix} \frac{1}{2}(1 + e^{-2\rho_i\lambda}) \\ \frac{1}{2}(1 - e^{-2\rho_i\lambda}) \end{pmatrix}
$$

is the vector of probabilities for going from the state 0 to 0 or to 1 along a branch of length λ.

Suppose we are to estimate λ by maximum likelihood, but using a discretised rate heterogeneity distribution $\widetilde{H}$ with equal weights. $\widetilde{H}$ has n_{cats} rate categories with a set of relative rates $(\tilde{\rho}^{(1)}, \ldots, \tilde{\rho}^{(n_{\text{cats}})})$ such that $\frac{1}{n_{\text{cats}}}\sum_j \tilde{\rho}^{(j)} = 1$. The expected number of mutations at site i along a branch of length κ, if i belongs to the category with rate $\tilde{\rho}^{(j)}$, is $\tilde{\rho}^{(j)}\kappa$. Thus if $\mathbb{P}(D_i|\tilde{\rho}^{(j)}, \kappa)$ is the probability of observing the data at site i if all branch lengths are κ,

$$
\mathbb{P}(D_i|\tilde{\rho}^{(j)}, \kappa) = \left(\frac{1 - e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{K_i}\left(\frac{1 + e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{n_{\text{tips}} - K_i}
$$

If $\tilde{\rho}^{(j)}$ is unknown then it needs to be summed over:

$$
\mathbb{P}(D_i|\kappa) = \frac{1}{n_{\text{cats}}}\sum_{j=1}^{n_{\text{cats}}}\left(\frac{1 - e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{K_i}\left(\frac{1 + e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{n_{\text{tips}} - K_i}
$$

Ferreti et al.

and then the probability of the entire alignment is:

$$\mathbb{P}(\mathbf{D}|\kappa) = \prod_{i=1}^{n_{\text{sites}}} \left[\frac{1}{n_{\text{cats}}} \sum_{j=1}^{n_{\text{cats}}} \left(\frac{1 - e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right)^{K_i} \left(\frac{1 + e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right)^{n_{\text{tips}} - K_i} \right]$$

Treated as a likelihood (i.e. a function of κ), and taking logs, we get an estimated branch length $\hat{\lambda}$:

$$\hat{\lambda} = \underset{\kappa}{\text{argmax}} \sum_{i=1}^{n_{\text{sites}}} \log \left[\frac{1}{n_{\text{cats}}} \sum_{j=1}^{n_{\text{cats}}} \left(\frac{1 - e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right)^{K_i} \left(\frac{1 + e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right)^{n_{\text{tips}} - K_i} \right] \tag{6}$$

Numerical results

We can calculate $\hat{\lambda}$ using (6), for a known n_{sites} and set of $\tilde{\rho}^{(j)}$, using standard numerical optimisation. We simulate evolution along n_{tips} branches of length λ according to the true rates ρ and calculate K_i by counting the number of 1s at the end. Figure 7 plots maximum-likelihood estimates of $\hat{\lambda}$ inferred using the 4-category Gamma model with alpha=0.5, against the actual branch length λ, for an alignment of 5,000 sites and a true gamma shape parameter of 0.5.

For small trees and slow rates, no bias in branch lengths is in evidence. The branch expansion effect is plain for longer branches, and the rate at which it occurs is correlated with λ. However, this increase eventually converges to an asymptote which is also reached faster for longer branch lengths, with the asymptotic value actually declining as the branch length increases. The latter effect may well be due to mutational saturation, as the increasing prevalence of unobserved back-mutations will tend to decrease the branch lengths in the reconstruction.

Biased per-site rate estimates lead to biased branch lengths

Several well-known software packages (Guindon et al. 2010; Kozlov et al. 2019; Minh et al. 2020) have been developed to estimate the maximum likelihood values of $\mathbf{T}$, M and ρ for a given alignment $\mathbf{D}$. These estimates can then be used to estimate the expected substitution rate $\text{E}(\rho_i|\mathbf{D})$ at every site D_i in $\mathbf{D}$. For example, IQ-TREE includes an empirical Bayes routine (Mayrose, Graur, et al. 2004) for this purpose.

Let us temporarily move back to a general phylogenetic scenario, without the simplifying

assumptions above. Let Λ be the total length of all branches of the true tree $\mathbf{T}$, and $\hat{\Lambda}$ the total

length of all branches of a maximum likelihood estimate $\hat{\mathbf{T}}$. We show using simulated data that

the ratio $\hat{\Lambda}/\Lambda$ is approximately equal to the mean expected substitution rate $\sum_i \mathrm{E}(\rho_i|\mathbf{D})/n_{\text{sites}}$.

We repeated the simulation exercise described under "Biased branch length estimates as the

result of equiprobable rate categories" above on a much larger scale. We used the same input

tree, but ran AliSim 50 times to generate 50 distinct simulated alignments using the continuous

Gamma site rate variation model. For each of these, we picked 500 random sequences from

each and progressively downsampled these to 100 in decrements of 100. We then performed

that downsampling procedure 4 more times each. This gave us a total of 1250 alignments, 250

of each size. We ran IQ-TREE using both the DGM and FreeRate on all of them.

In figure 8, left, the x-axis represents the ratio of the sum of all branch lengths of the

reconstructed DGM tree to that of the true tree. The y-axis is the reciprocal of $\sum_i \mathrm{E}(\rho_i|\mathbf{D})/n_{\text{sites}}$

as estimated using the empirical Bayes routine. Grey lines join results from the same sequencing

and downsampling replicate. There is a very clear correlation, albeit with some noise around

the line of equality. However, it should be noted that the maximum likelihood estimate of a

phylogeny derived from a simulated alignment would not be expected to exactly match the

tree used to generate that alignment. We can, however, get an estimate for the ML phylogeny

of the simulated alignment by using FreeRate (Fig 8, right). Here the points are very close

indeed to the line of equality, albeit with increasing deviation from it as tree sizes get smaller.

We are unsure about the reasons for the latter phenomenon, which may simply be to do with

limitations of the FreeRate model or the heuristic maximum likelihood approximation, and

leave it for future work.

As branch lengths are a measure of the amount of elapsed mutation, it is entirely to be

expected that if per-site mutation rates are systematically misestimated then lengths would

also be. The true ρ_is have a mean of 1 by definition, but their estimated values may not. In

the next section we show this formally in our simplified scenario of a star-like tree.

Likelihood in the large n_{tips} limit

581 To recap, ρ_i is the genuine relative substitution rate at site i, drawn from the distribution H.

582 The mean ρ_i over $\mathbf{D}$ is 1 and thus the ρ_is sum to n_{sites}. The probability of seeing a mutation at

Figure 8: Left: For 1250 simulated alignments (points), we plot, on the x-axis, the ratio of the total tree length of a phylogeny reconstructed using the DGM to the total length of the true tree, versus, on the y-axis, the reciprocal of mean reconstructed site rate across the genome. The thick black line is equality; thin black lines connect alignments from the same simulation and downsampling replicate. Right: The same, except the x-axis is now the ratio of the total branch length from a DGM reconstruction to the total branch length from a FreeRate reconstruction.

i on any branch is $\frac{1-2e^{-\rho_i\lambda}}{2}$. In contrast, $\tilde{\rho}^{(j)}$ is one of the n_{cats} discrete rates in the DGM which

each D_i's rate is equal to with probability $1/n_{\text{cats}}$. Define $f_i = K_i/n_{\text{tips}}$. Now, at a single site

D_i:

$$\log(\mathbb{P}(D_i|\kappa)) = \log\left[\frac{1}{n_{\text{cats}}}\sum_{j=1}^{n_{\text{cats}}}\left(\frac{1-e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{K_i}\left(\frac{1+e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{n_{\text{tips}}-K_i}\right]$$

$$= \log\left[\frac{1}{n_{\text{cats}}}\sum_{j=1}^{n_{\text{cats}}}\left(\left(\frac{1-e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{f_i}\left(\frac{1+e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{1-f_i}\right)^{n_{\text{tips}}}\right]$$

In the limit of large n_{tips}, the sum here is dominated by its largest term, and we can write

$$\log(\mathbb{P}(D_i|\kappa)) \approx \log\left[\frac{1}{n_{\text{cats}}}\max_j\left(\left(\frac{1-e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{f_i}\left(\frac{1+e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{1-f_i}\right)^{n_{\text{tips}}}\right]$$

$$= \log\left[\max_j\left(\left(\frac{1-e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{f_i}\left(\frac{1+e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{1-f_i}\right)^{n_{\text{tips}}}\right] - \log(n_{\text{cats}})$$

$$= \max_j\left[\log\left(\left(\frac{1-e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{f_i}\left(\frac{1+e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)^{1-f_i}\right)^{n_{\text{tips}}}\right] - \log(n_{\text{cats}})$$

$$= n_{\text{tips}}\max_j\left[f_i\log\left(\frac{1-e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right) + (1-f_i)\log\left(\frac{1+e^{-2\tilde{\rho}^{(j)}\kappa}}{2}\right)\right] - \log(n_{\text{cats}})$$

and the log-likelihood (6) can be written (up to additive constants) as

$$\log(\mathrm{L}(\kappa)) \approx n_{\text{tips}} \sum_{i=1}^{n_{\text{sites}}} \max_{j} \left[f_i \log \left(\frac{1 - e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right) + (1 - f_i) \log \left(\frac{1 + e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right) \right]$$

The value of f_i is fixed by the data rather than dependent on the value of κ, and for large n_{tips}, it converges towards its expected value $\frac{1 - e^{-2\rho_i\lambda}}{2}$. Therefore, the log likelihood can be well approximated by

$$\log(\mathrm{L}(\kappa)) \approx n_{\text{tips}} \sum_{i=1}^{n_{\text{sites}}} \max_{j} \left[\frac{1 - e^{-2\rho_i\lambda}}{2} \log \left(\frac{1 - e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right) + \frac{1 + e^{-2\rho_i\lambda}}{2} \log \left(\frac{1 + e^{-2\tilde{\rho}^{(j)}\kappa}}{2} \right) \right]$$

Likelihood in the "infinite sites model"

If λ and κ are both small, we can approximate $1 - e^{-2\rho_i\lambda}$ as $2\rho_i\lambda$ and $1 - e^{-2\tilde{\rho}^{(j)}\kappa}$ as $2\tilde{\rho}^{(j)}\kappa$, leading to the approximation

$$\log(\mathrm{L}(\kappa)) \approx n_{\text{tips}} \sum_{i=1}^{n_{\text{sites}}} \max_{j} \left[\rho_i\lambda \log \left(\tilde{\rho}^{(j)}\kappa \right) + (1 - \rho_i\lambda) \log \left(1 - \tilde{\rho}^{(j)}\kappa \right) \right]$$

$$\approx n_{\text{tips}} \sum_{i=1}^{n_{\text{sites}}} \max_{j} \left[\rho_i\lambda \log \left(\tilde{\rho}^{(j)}\kappa \right) - \tilde{\rho}^{(j)}\kappa \right]$$

where in the last expression we ignore quadratic and higher terms in λ and κ. In order to move the maximisation over j out of the summation, separate it into a variable j_i for each position i:

$$\log(\mathrm{L}(\kappa)) \approx n_{\text{tips}} \max_{j_1, \ldots, j_{n_{\text{sites}}}} \sum_{i=1}^{n_{\text{sites}}} \left[\rho_i\lambda \log \left(\tilde{\rho}^{(j_i)}\kappa \right) - \tilde{\rho}^{(j_i)}\kappa \right] \tag{7}$$

Note that if $g(\kappa) = \lambda\rho \cdot \log(\tilde{\rho}\kappa) - \tilde{\rho}\kappa$ then $g'(\kappa) = \frac{\lambda\rho}{\kappa} - \tilde{\rho}$ and g is maximised when $\tilde{\rho} = \frac{\lambda\rho}{\kappa}$, and thus if $\lambda = \kappa$ and the $\tilde{\rho}^{(j_i)}$s can be freely assigned to any value (i.e. $n_{\text{cats}} = n_{\text{sites}}$; a FreeRate model with one category per site), the maximum likelihood is obtained when each $\tilde{\rho}^{(j_i)} = \rho_i$.

We now want to maximise (7) over κ to find $\hat{\lambda}$.

$$\max_{\kappa} \log(\mathrm{L}(\kappa)) \approx \max_{\kappa} \left[n_{\text{tips}} \max_{j_1, \ldots, j_{n_{\text{sites}}}} \sum_{i=1}^{n_{\text{sites}}} \left(\rho_i\lambda \log \left(\tilde{\rho}^{(j_i)}\kappa \right) - \tilde{\rho}^{(j_i)}\kappa \right) \right]$$

$$= n_{\text{tips}} \max_{j_1, \ldots, j_{n_{\text{sites}}}} \max_{\kappa} \left[\sum_{i=1}^{n_{\text{sites}}} \left(\rho_i\lambda \log \left(\tilde{\rho}^{(j_i)}\kappa \right) - \tilde{\rho}^{(j_i)}\kappa \right) \right] \tag{8}$$

The first derivative of the internal expression with respect to κ is

$$\lambda \sum_{i=1}^{n_{\text{sites}}} \frac{\rho_i}{\kappa} - \sum_{i=1}^{n_{\text{sites}}} \tilde{\rho}^{(j_i)}$$

which gives a maximum at:

$$\kappa = \lambda \frac{\sum_{i=1}^{n_{\text{sites}}} \rho_i}{\sum_{i=1}^{n_{\text{sites}}} \tilde{\rho}^{(j_i)}} \tag{9}$$

$$= \lambda \frac{n_{\text{sites}}}{\sum_{i=1}^{n_{\text{sites}}} \tilde{\rho}^{(j_i)}}$$

$$= \frac{\lambda}{\bar{\rho}}$$

where $\bar{\rho} = \sum_i \tilde{\rho}^{j_i} / n_{\text{sites}}$. Inserting this into (8) gives:

$$\max_{\kappa} \log(\mathrm{L}(\kappa)) \approx n_{\text{tips}} \max_{j_1,\ldots,j_{n_{\text{sites}}}} \left[\sum_{i=1}^{n_{\text{sites}}} \left(\rho_i \lambda \log\left(\tilde{\rho}^{(j_i)} \frac{\lambda}{\bar{\rho}} \right) - \tilde{\rho}^{(j_i)} \frac{\lambda}{\bar{\rho}} \right) \right] \tag{10}$$

$$= n_{\text{tips}} \max_{j_1,\ldots,j_{n_{\text{sites}}}} \left[\lambda \sum_{i=1}^{n_{\text{sites}}} \rho_i \log\left(\tilde{\rho}^{(j_i)} \frac{\lambda}{\bar{\rho}} \right) - \sum_{i=1}^{n_{\text{sites}}} \tilde{\rho}^{(j_i)} \frac{\lambda}{\bar{\rho}} \right]$$

$$= n_{\text{tips}} \max_{j_1,\ldots,j_{n_{\text{sites}}}} \left[\lambda \left(\sum_{i=1}^{n_{\text{sites}}} \rho_i \log\left(\frac{\tilde{\rho}^{(j_i)}}{\bar{\rho}} \right) + \sum_{i=1}^{n_{\text{sites}}} \rho_i \log \lambda - n_{\text{sites}} \right) \right]$$

$$= n_{\text{tips}} \max_{j_1,\ldots,j_{n_{\text{sites}}}} \left[\lambda \left(\sum_{i=1}^{n_{\text{sites}}} \rho_i \log\left(\frac{\tilde{\rho}^{(j_i)}}{\bar{\rho}} \right) + n_{\text{sites}}(\log \lambda - 1) \right) \right] \tag{11}$$

Every site i has a value of j_i, $\hat{j}_i$, that maximises $\sum_{i=1}^{n_{\text{sites}}} \rho_i \log(\tilde{\rho}^{(j_i)}/\bar{\rho})$. Maximising all of them simultaneously gives the maximum likelihood estimate of λ as

$$\hat{\lambda} = \lambda \frac{n_{\text{sites}}}{\sum_i \tilde{\rho}^{(\hat{j}_i)}} = \frac{\lambda}{\bar{\rho}_{ml}} \tag{12}$$

where $\bar{\rho}_{ml} = \sum_i \tilde{\rho}^{(\hat{j}_i)}/n_{\text{sites}}$. This formally demonstrates the validity of the findings of the previous section for our simplified tree structure.

The impact of the equal-weights discretisation in the two-category and n-category case

If (7) is the approximate log likelihood for a given κ integrating over site rate assignments, then

$$\log(\mathrm{L}(\kappa, \{j_i\})) \approx n_{\text{tips}} \sum_{i=1}^{n_{\text{sites}}} \left[\rho_i \lambda \log \left(\tilde{\rho}^{(j_i)} \kappa \right) - \tilde{\rho}^{(j_i)} \kappa \right] \tag{13}$$

is the approximate log likelihood for a given κ and known rate assignments.

Consider the case where the true site-specific rates are defined by a FreeRate distribution with two kinds of sites, with "slow" and "fast" rates ρ^s and ρ^f, $\rho^s < \rho^f$. The weights are w^s and w^f such that $w^s + w^f = 1$ and $w^s \rho^s + w^f \rho^f = 1$.

In the last section we showed that if each site is assigned to its correct underlying rate and $\lambda = \kappa$ then the likelihood is maximised when each D_i is assigned its correct rate. It should also be clear that if each $\rho_i = \tilde{\rho}^{j_i}$ then $\lambda = \kappa$ in (9). Thus the maximum likelihood values of λ and the $\tilde{\rho}^{j_i}$s are the ones that generated the data. The log value of this likelihood is approximately:

$$\max \log(\mathrm{L}) \approx n_{\text{tips}} n_{\text{sites}} \lambda \left(w^s \rho^s \left(\log \left(\rho^s \lambda \right) - 1 \right) + w^f \rho^f \left(\log \left(\rho^f \lambda \right) - 1 \right) \right)$$

$$= n_{\text{tips}} n_{\text{sites}} \lambda \left(w^s \rho^s \log \left(\rho^s \lambda \right) + w^f \rho^f \log \left(\rho^f \lambda \right) - 1 \right) \tag{14}$$

Now consider a model with rates $\tilde{\rho}^s$ and $\tilde{\rho}^f$ such that $\tilde{\rho}^s = 2\rho^s/(\rho^s + \rho^f)$, $\tilde{\rho}^f = 2\rho^f/(\rho^s + \rho^f)$ and equal weights of $1/2$. (Equal weights are actually required for the mean rate across sites to be 1 here.) If we assign ρ^s as the rate to the $n_{\text{sites}} w^s$ "slow" sites and ρ^f to the $n_{\text{sites}} w^f$ "fast" ones, the mean rate across sites is:

$$\bar{\rho} = w^s \tilde{\rho}^s + w^f \tilde{\rho}^s$$

$$= \frac{2(w^s \rho^s + w^f \rho^f)}{\rho^s + \rho^f}$$

$$= \frac{2}{\rho^s + \rho^f}$$

663 The likelihood for this rate assignment and a branch length of $\lambda/\bar{\rho}$ is:

664
$$\log(\mathrm{L}) = n_{\text{tips}} n_{\text{sites}} \lambda \left(w^s \left(\rho^s \log \left(\frac{\tilde{\rho}^s \lambda}{\bar{\rho}} \right) - \frac{\tilde{\rho}^s}{\bar{\rho}} \right) + w^f \left(\rho^s \log \left(\frac{\tilde{\rho}^f \lambda}{\bar{\rho}} \right) - \frac{\tilde{\rho}^f}{\bar{\rho}} \right) \right)$$

665
$$= n_{\text{tips}} n_{\text{sites}} \lambda \left(w^s \left(\rho^s \log \left(\rho^s \lambda \right) - \rho^s \right) + w^f \left(\rho^f \log \left(\rho^f \lambda \right) - \rho^f \right) \right)$$

666
667
$$= n_{\text{tips}} n_{\text{sites}} \lambda \left(w^s \rho^s \log \left(\rho^s \lambda \right) + w^f \rho^f \log \left(\rho^f \lambda \right) - 1 \right)$$

This matches (14). Thus the likelihood of the equal weights model with rates $\tilde{\rho}^s$ and $\tilde{\rho}^f$ is equal

to the maximum likelihood of the FreeRate model. As the equal weights model is nested in the

FreeRate model, this must also be the *maximum* likelihood of the former.

So the likelihood of the equal weights model is maximised by assigning individual rates to

sites in a way that departs from the equal weights assumption. But when this is done, the

mean rate $\bar{\rho} = w^s \tilde{\rho}^s + w^f \tilde{\rho}^f$ is not 1 and thus the maximum likelihood estimate $\hat{\lambda}$ of the branch

length is not λ.

The relative bias in the branch length estimate is given by $\frac{\hat{\lambda}-\lambda}{\lambda}$ and by using the identities

$w^s = 1 - w^f$ and $\rho^f = (1 - w^s \rho^s)/w^f$ this can be written as:

677
$$\frac{\hat{\lambda} - \lambda}{\lambda} = \frac{1}{\bar{\rho}} - 1$$

678
679
$$= \frac{1 - \rho^s}{2} \left(\frac{1}{w^f} - 2 \right) \tag{15}$$

This result shows that a) the magnitude of the bias is larger for a larger separation between

fast and slow rates (bias increases as $\rho^s \to 0$) and for a smaller fraction of rapidly evolving sites

(bias increases as $w^f \to 0$). As ρ^s is always less than 1, the sign of the bias is driven by w^f: if

this is greater than 0.5 then branches will contract; otherwise they expand. The former case is

rather unlikely in real-world situations, where rate distributions generally have a mode of less

than 1 (Jia et al. 2014; Silvestro et al. 2024).

We can generalise this to the situation of a tree generated using a n_{cats}-category FreeRate

distribution and reconstructed with an equal-weight model also with n_{cats} categories. Denote

the rates and weights of the FreeRate distribution by n_{cats}-dimensional vectors $\mathbf{P} = \{\rho^{(c)}\}$

and $\mathbf{W} = \{w^{(c)}\}$ satisfying $\sum_c w^{(c)} = 1$ and $\sum_c \rho^{(c)} w^{(c)} = 1$. If we assign site-specific rates as

$\tilde{\rho}^{(c)} = \rho^{(c)} n_{\text{cats}} / \sum_c \rho^{(c)}$ and assign the sites to categories correspondingly, these rates correspond

to the maximum of the unconstrained likelihood (9), and thus must maximise the likelihood

of the reconstruction model. The value of $\bar{\rho}$ for this realisation can be easily computed as

$n_{\text{cats}}/\sum_c \rho^{(c)}$ and therefore $\hat{\lambda} = \lambda \sum_c \rho^{(c)}/n_{\text{cats}}$. Then the relative bias can be written as:

694
695
$$\frac{\hat{\lambda} - \lambda}{\lambda} = \frac{\sum_c \rho^{(c)}}{n_{\text{cats}}} - 1 = -\frac{1}{n_{\text{cats}}} \sum_c \left(w^{(c)} n_{\text{cats}} - 1\right) \left(\rho^{(c)} - \frac{\sum_{c'} \rho^{(c')}}{n_{\text{cats}}}\right) = -\text{Cov}\left(\mathbf{W}/\overline{\mathbf{W}}, \mathbf{P}\right)$$

696 Thus the bias is inversely proportional to the covariance between rates and relative weights

697 among categories. When the distribution includes a tail of rare but fast-evolving sites, there is

698 a negative correlation between weight and rate and therefore a branch expansion effect.

Fitting Gamma-distributed rates with $k = 1$ by a discretised Gamma with $n_{\text{cats}} = 2$

Here we allow the underlying rates to be drawn from the simplest possible continuous Gamma

distribution, that with shape and scale 1 (i.e. an exponential distribution with rate 1). We will

to fit a two-category discrete model with equal weights. As in the previous section, we take

two rates $\tilde{\rho}^{(s)}$ and $\tilde{\rho}^{(f)}$, with $\tilde{\rho}^{(s)} < \tilde{\rho}^{(f)}$, such that $\tilde{\rho}^{(s)} + \tilde{\rho}^{(f)} = 2$. We can write $\rho^{(s)} = 1 - \epsilon$ and

$\rho^{(f)} = 1 + \epsilon$ for $0 \leq \epsilon \leq 1$. Both weights are $1/2$. In contrast to the previous sections, where

category rates are known, we do not assume we know ϵ and must estimate it.

The expression which is to be maximised with respect to both the rates and their site

assignments is derived from (11):

709
$$\log(\text{L}(\tilde{\rho}^{(s)}, \tilde{\rho}^{(f)}, \{j_i\})) \approx n_{\text{tips}} \max_{j_1,\dots,j_{n_{\text{sites}}}} \left[\lambda \left(\sum_{i=1}^{n_{\text{sites}}} \rho_i \log\left(\frac{\tilde{\rho}^{(j_i)}}{\bar{\rho}}\right) + n_{\text{sites}}(\log\lambda - 1)\right)\right]$$

710
711
$$= n_{\text{tips}} \left[\max_{j_1,\dots,j_{n_{\text{sites}}}} \left[\lambda \sum_{i=1}^{n_{\text{sites}}} \rho_i \log\left(\frac{\tilde{\rho}^{(j_i)}}{\bar{\rho}}\right)\right] + n_{\text{sites}}(\log\lambda - 1)\right] \tag{16}$$

where each $j_i \in \{s, f\}$. We first need to show that, for some α between 0 and 1, the likelihood

is maximised when the slowest αn_{sites} sites are allocated to $\tilde{\rho}^{(s)}$ and the fastest $(1 - \alpha)n_{\text{sites}}$ to

$\tilde{\rho}^{(f)}$. To see this, suppose you know the value of α that maximises the likelihood L and calculate

L under this allocation. Now switch the assignments of two sites a and b, where a was assigned

to $\tilde{\rho}^{(s)}$ and b to $\tilde{\rho}^{(f)}$ in the assignment maximimising L, and recalculate the likelihood as L′.

Using (10) as the expression for the likelihoods, L and L′ differ only in as much as L contains

the term:

$$M = \lambda \left(\rho_a \log \left(\frac{\tilde{\rho}^{(s)} \lambda}{\bar{\rho}} \right) - \frac{\tilde{\rho}^{(s)}}{\bar{\rho}} + \rho_b \log \left(\frac{\tilde{\rho}^{(f)} \lambda}{\bar{\rho}} \right) - \frac{\tilde{\rho}^{(f)}}{\bar{\rho}} \right)$$

721 while L' contains:

$$M' = \lambda \left(\rho_a \log \left(\frac{\tilde{\rho}^{(f)} \lambda}{\bar{\rho}} \right) - \frac{\tilde{\rho}^{(f)}}{\bar{\rho}} + \rho_b \log \left(\frac{\tilde{\rho}^{(s)} \lambda}{\bar{\rho}} \right) - \frac{\tilde{\rho}^{(s)}}{\bar{\rho}} \right)$$

724 where $\bar{\rho} = \alpha n_{\text{sites}} \tilde{\rho}^{(s)} + (1 - \alpha) n_{\text{sites}} \tilde{\rho}^{(f)}$. It remains solely to show that

$$\rho_a \log \left(\tilde{\rho}^{(s)} \right) + \rho_b \log \left(\tilde{\rho}^{(f)} \right) > \rho_a \log \left(\tilde{\rho}^{(f)} \right) + \rho_b \log \left(\tilde{\rho}^{(s)} \right)$$

727 but straightforwardly:

$$\rho_a \log \left(\tilde{\rho}^{(s)} \right) + \rho_b \log \left(\tilde{\rho}^{(f)} \right) - \rho_a \log \left(\tilde{\rho}^{(f)} \right) - \rho_b \log \left(\tilde{\rho}^{(s)} \right) = \rho_a \log \left(\frac{\tilde{\rho}^{(s)}}{\tilde{\rho}^{(f)}} \right) + \rho_b \log \left(\frac{\tilde{\rho}^{(f)}}{\tilde{\rho}^{(s)}} \right)$$

$$= \rho_a \log \left(\frac{\tilde{\rho}^{(s)}}{\tilde{\rho}^{(f)}} \right) - \rho_b \log \left(\frac{\tilde{\rho}^{(s)}}{\tilde{\rho}^{(f)}} \right)$$

$$= (\rho_a - \rho_b) \cdot \log \left(\frac{\tilde{\rho}^{(s)}}{\tilde{\rho}^{(f)}} \right)$$

$$> 0$$

733 (the last because both $\rho_a < \rho_b$ and $\tilde{\rho}^{(s)} < \tilde{\rho}^{(f)}$). Thus the maximum likelihood allocation of

734 sites to rates must work in this way and it remains only to estimate α and ϵ.

735   Rearrange the index of the sum in (16) such that $\rho_j > \rho_i$ if $j > i$. To get an analytic

736 expression for the log likelihood, consider that

$$\lim_{n_{\text{sites}} \to \infty} \sum_{i=1}^{n_{\text{sites}}} \rho_i \log \left(\frac{\tilde{\rho}^{(j_i)}}{2\bar{\rho}} \right) = \int_0^1 \nu(q) \log \left(\frac{1 - (-1)^{i(q)} \epsilon}{1 + \epsilon(1 - 2\alpha)} \right) dq$$

739 where $\nu(q) = -\log(1 - q)$ is the quantile function of the true rate distribution and $i(q) = 1$ if

740 $q \leq \alpha$ and 0 otherwise. If we take n_{sites} to be large enough that the integral can be taken as an

741 approximation of the sum, then the term $\frac{1}{2} n_{\text{tips}} n_{\text{sites}} (\log \lambda - 1)$ in (16) is invariant by site and

it remains only to maximise the integral over α and ϵ:

$$\log(\mathrm{L}(\alpha, \epsilon)) \propto -\int_0^\alpha \log(1-q)\log\left(\frac{1-\epsilon}{1+\epsilon(1-2\alpha)}\right)dq - \int_\alpha^1 \log(1-q)\log\left(\frac{1+\epsilon}{1+\epsilon(1-2\alpha)}\right)dq$$

$$= -\log\left(\frac{1-\epsilon}{1+\epsilon(1-2\alpha)}\right)\int_0^\alpha \log(1-q)dq - \log\left(\frac{1+\epsilon}{1+\epsilon(1-2\alpha)}\right)\int_\alpha^1 \log(1-q)dq$$

$$= -\log\left(\frac{1-\epsilon}{1+\epsilon(1-2\alpha)}\right)[(q-1)\log(1-q)+1-q]_0^\alpha$$
$$- \log\left(\frac{1+\epsilon}{1+\epsilon(1-2\alpha)}\right)[(q-1)\log(1-q)+1-q]_\alpha^1$$

$$= -\log\left(\frac{1-\epsilon}{1+\epsilon(1-2\alpha)}\right)((\alpha-1)\log(1-\alpha)-\alpha)$$
$$- \log\left(\frac{1+\epsilon}{1+\epsilon(1-2\alpha)}\right)((1-\alpha)\log(1-\alpha)+\alpha-1)$$

To maximise this, we calculate its partial derivatives with respect to ϵ and α, to obtain:

$$\frac{\partial \log(\mathrm{L}(\alpha,\epsilon))}{\partial \epsilon} = -\frac{1}{1-\epsilon}[\alpha+(1-\alpha)\log(1-\alpha)]$$
$$+ \frac{1}{1+\epsilon}[1-\alpha-(1-\alpha)\log(1-\alpha)] - \frac{1-2\alpha}{1+\epsilon(1-2\alpha)}$$

$$\frac{\partial^2 \log(\mathrm{L}(\alpha,\epsilon))}{\partial\alpha\partial\epsilon} = \log(1-\alpha)\log\left(\frac{1+\epsilon}{1-\epsilon}\right) + \frac{2\epsilon}{1+\epsilon(1-2\alpha)}$$

If solved numerically this gives $\epsilon \approx 0.65$ and $\alpha \approx 0.64$, from which $\bar{\rho} \approx 0.81$. If we look at the expression for of $\hat{\lambda}$ from equation (12), the maximisation of the likelihood results in a branch expansion of roughly 20%.