Microplastics as a novel facilitator for antimicrobial resistance: Effects of concentration,
composition, and size on *Escherichia coli* multidrug resistance

Neila Gross[1*], Johnathan Muhvich[2], Carly Ching[2], Bridget Gomez[2], Evan Horvath[2], Yanina
Nahum[3], and Muhammad H. Zaman[2, 3]

Affiliations

[1]Department of Materials Science and Engineering, Boston University, Boston, MA 02215

[2]Department of Biomedical Engineering, Boston University, Boston, MA 02215

[3]Center on Forced Displacement, Boston University, 111 Cummington Mall, Boston, MA, 02215

*Corresponding author: ngross1@bu.edu

Word count: 5554

26
27

Abstract

Microplastics (MPs) have emerged as a significant environmental pollutant with profound implications for public health, particularly as substrates to facilitate bacterial antimicrobial resistance (AMR). Recently, studies have shown that MPs may accommodate microbial communities, chemical contaminants and genetic material containing AMR genes. This study investigated the effects of MP concentration, composition, and size on the development of multidrug resistance in *Escherichia coli*. Specifically, we exposed *E. coli* to varying concentrations of different MP types, including polyethylene (PE), polystyrene (PS), and polypropylene (PP), across a range of sizes (3-10 µm, 10-50 µm, and 500 µm). Results indicated a direct correlation between MP presence and elevated multidrug-resistant (MDR) in *E. coli*. Notably, MPs exhibited a higher propensity for facilitating resistance than control substrates such as glass, likely due to their hydrophobicity, greater adsorption capacities, and surface chemistries. Furthermore, we observed that co-culture with MPs resulted in biofilm formation. Notably, we found that the bacteria from passaged MPs formed stronger biofilms once the MPs were removed, associated with changes in motility. Thus, we find that MPs also select for cells that are better at forming biofilms, which can lead to recalcitrant infections in the environment and healthcare setting. Our study highlights the immediate need for comprehensive environmental management strategies to mitigate the risk posed by MPs.

Importance

Antimicrobial resistance is one of the world's most pressing global health crises, with an estimated 10 million deaths per year forecasted by 2050. With the pipeline of antibiotics running dry, it is imperative that mitigation strategies understand the mechanisms that drive the genesis of antimicrobial resistance. One emerging dimension of antimicrobial resistance is the environment. This study highlights the relationship between a widespread environmental pollutant, (MPs), and the rise of drug-resistant bacteria. While it is known that MPs facilitate resistance through several modes (biofilm formation, plastic adsorption rates, etc.), this study fills the knowledge gap on how different types of MPs are contributing to antimicrobial resistance.

Introduction

Global plastic use and mismanaged disposal are significant environmental and public health concerns. Plastic use has increased twenty-fold since 1964, and prevailing estimates suggest global unmanaged trash will reach 155-265 megatons per year in 2060.[1] Not surprisingly, the detection of (MPs) has significantly increased. MPs are canonically insoluble synthetic particles or polymer matrices with regular or irregular shapes and sizes ranging from the micrometer to millimeter range.[2] Primary sources of MPs include polyethylene, polypropylene, and polystyrene particles in cosmetic and medicinal products.[3] Several countries have banned the use of primary MPs within certain industry segments; however, MPs are still generated through the degradation of existing plastics, creating contamination beyond legislation around new product formulations.[3] These degradation products, characterized as secondary MPs, originate from physical, chemical,

and biological processes, resulting in plastic debris fragmentation. Different sources of MPs and
their varying surface chemistry also cause them to occur in varying shapes and sizes, such as
pellets, fibers, and fragments in environmental samples.[3] Depending on the conditions plastics
are exposed to and the resulting change in surface properties, MPs arise as a unique substrate that
has proved to be very difficult to control.[25, 26]

MPs have infiltrated various ecosystems on the planet, ranging from submarine canyons[4] to the
summit of Mount Everest.[5] Additionally, wastewater has become a significant reservoir for MPs
and other anthropological wastes. Despite global awareness, MPs can persist through wastewater
treatment plants, disseminating them into surrounding environments. Their persistence can be
attributed to their small size, buoyancy, and hydrophobic properties, allowing them to adhere to
organic matter and avoid sedimentation.[6] Consequently, treated wastewater effluents serve as a
main source of MP pollution in aquatic environments where they accumulate in sediments and
surface waters and interact with the organisms around them.[6] The repercussions of MPs in
wastewater are manifold, especially considering their impacts on human health.

Contemporaneously, increased rates of antimicrobial resistance (AMR)—the ability of microbes
to protect themselves against antimicrobials—have been observed in bacterial populations across
the globe. AMR can be influenced by a multifaceted network of factors, including the overuse
and misuse of antibiotics, poor sanitation and hygiene, and environmental contamination with
antibiotic residues in wastewater.[27] Recent studies show that MPs might also play an important
role in the development of AMR.[28, 29, 30] This is mainly due to their ability to accommodate
microbial communities and chemical contaminants and genetic material containing antibiotic-
resistant genes (ARGs) through biofilm formation.[19] In these communities of bacteria that grow
together colonizing MP surfaces, ARGs can be transferred to pathogenic bacteria through
horizontal gene transfer.[19] While it has been established that bacteria on the surface of MPs host
ARGs, there is limited knowledge on AMR development as a function of MP properties
(composition, size, concentration, etc.) as well as their interactions with various antibiotics.[8,11]

Understanding the interplay between AMR and MPs is critical, especially in places with high
infection rates and significant plastic waste, such as low-resource settings. The difficulty in
treating infectious diseases in these areas, combined with inadequate wastewater treatment—
which may result in higher concentrations of MPs—may contribute to the observed increase in
AMR cases among vulnerable populations. Therefore, understanding the fundamental
interactions between MPs and AMR development is imperative. To date, few studies have
looked at the effect of MP size, structure, and other features on the development of drug
resistance in the presence of antibiotics. Those that have observed ARGs and increased tolerance
to antibiotics.[38,39] In this study, we investigate the impact of different MP characteristics,
including MP concentration, surface composition (between plastics and other materials), size,
and surface area, on the *in vitro* development of resistance to ampicillin, ciprofloxacin,
doxycycline, and streptomycin in *E. coli*. These antibiotics are broad-spectrum, represent

different antibiotic classes, and are readily found in wastewater systems.[12] More specifically, we
probed if MP concentration (no. plastics/µL), size, surface area, and composition (between
plastics types) had any impact on (1) bacterial growth alone, (2) antibiotic-specific AMR and (3)
multidrug resistance (MDR) between the four antibiotics. We then examined potential
mechanisms behind difference in AMR development, specifically studying biofilm formation.
Our results identify that MPs play an important and significant role in the development of AMR.
Overall, our findings may provide context to wastewater surveillance data and provide insights
into waste management and associated disease burdens.

Methods

<u>Strains and culture conditions</u>

E. coli MG1655 (ATCC 700926) was used in all experiments. All liquid cultures were grown in
Lysogeny Broth (Miller, LB) medium under shaking conditions at 180 rpm at 37 °C. First, MPs
(Table S.1) were added to the media for 48 hours to allow biofilm surface attachment and full
maturity.[40] Following the initial MP exposure, any additional amendments were added, which
included subinhibitory levels of ampicillin (Sigma Aldrich), ciprofloxacin (MP Biomedicals,
86393-32-0), doxycycline (Sigma-Aldrich), streptomycin (Sigma-Aldrich), as indicated below.

<u>Growth Curve Data</u>

Wild-type *E. coli* MG1655 was cultured at 37 in 4mL of the LB with varying MP
concentrations (100, 500, 1000, 2000, 4000, 5000, 10000, and 15000 plastics/µL) in glass culture
tubes and sampled at 0, 1, and 24 hours. These samples were plated in triplicate on LB agar to
determine the CFU/mL at the time point of interest. Cultures were plated before and after being
vortexed, to compensate for cells attached to the MP surface, and CFUs were counted 24 hours
after incubation (Fig. 1)

Antibiotic and MP susceptibility testing

To determine the initial Minimum Inhibitory Concentration (MIC) of bacteria that is used as the
Day 0 for the MIC fold change comparison, cells were grown in drug-free media until substantial
biofilm growth was detected (~48 hours) using confocal microscopy (procedure below).
Microparticles (plastics and glass) were vortexed for one minute and spun down to release the
biofilm on the surface (Fig. S.1). The supernatant of the vortexed solution was used in a broth
microdilution MIC in a 96-well plate using LB media to determine the MIC, effectively testing
the MIC of attached cells and effluent cells (Fig. S.2).[34] The microbroth dilutions begin in the
first column with 100µL of an antibiotic and are diluted two-fold into the next column and so on.
10 µL of the samples from the culture are then placed in each well and grown in a static
incubator for 24 hours. Following this, the optical density is checked in a spectrophotometer at

600 nm and recorded against a negative control. This was done every 24 hours as shown in
Figure S.2.

MP-antibiotic combination assays

After the initial 48 hour growth period and MIC testing was completed as described above,
saturated liquid cultures (cultures that had reach carrying capacity) were passaged once a day
into fresh amended media for the antibiotics only or the WT (no antibiotics or MPs)–4 mL of LB
broth, 40% of the initial MIC (3.2 µg/mL, 0.0074 µg/mL, 1.6 µg/mL, and 6.4 µg/mL for
ampicillin, ciprofloxacin, doxycycline, and streptomycin, respectively) of a singular antibiotic,
and their respective growth conditions--via 1:100 dilution until day ten (40µL of the previous
day's culture passaged into 4 mL of the new media). For the MP samples, after each 24-hour
exposure period, samples were vortexed to release a portion of the biofilm cells (Fig. S1) to be
passaged and tested in MIC plates (Fig. S2B). Not all cells are released from the surface of the
biofilm but enough that you can see biofilm in the surrounding media. Then, the MPs were spun
down (if they were 10 µm) or allowed enough time to settle (between five and ten minutes) to
make sure none were taken out in the following step, in which the supernatant was taken out of
the culture tube (40µL saved to passage into the new media and the rest used for the MIC
assays), and new media (along with 40µL of the supernatant) was allotted into the *same* culture
tube containing the *same* MPs (with less biofilm than pre vortexing) throughout the entire
experiment (Fig. S2B). All groups were tested for antibiotic susceptibility every two days, as
described above. As a control, cells in drug-free media with the same variable of MPs
(concentration, size, composition), cells with no adhesion sites (no MPs or glass spheres) but
with drug (40% MIC), as well cells in drug-free and particle free media were also tested (Wild
Type (WT)). A schematic of the experiment is displayed in Fig. S.2.

Confocal Laser Scanning Microscopy (CLSM) image collection and analysis

Visualizing the cells on the surface of the particles was done using confocal microscopy. Using
the cultures described in the susceptibility testing section, 500 µm diameter polystyrene spheres
were randomly selected out of each culture and dyed using the LIVE/DEAD *Bac*Light Bacterial
Viability Kit Protocol, per manufacturer instructions. Briefly, SYTO 9 was used to analyze
viable cells, as it can permeate all bacterial cell membranes. In contrast, propidium iodide was
used to count dead cells, as it can only enter cells with disrupted membranes. Each plastic was
put into a centrifuge tube with 1 mL of sterile water, and three µL of staining solution (1.5 µL of
each dye) was injected into the tube. Samples were incubated for 20 minutes at room temperature
and protected from light.

Stained samples were imaged with an inverted CLSM (Olympus DSU spinning disk confocal)
using a 10X oil immersion objective. Samples individually stained with propidium iodide and
SYTO 9 were first analyzed separately to ensure clear signals without overlap. The particles

were removed from the dye and suspended in a costar clear round bottom 96 well plate to
preserve the MP shape while imaging. After finding the sample and adjusting brightness
parameters, z-stacks with an optimized step size were taken for each sample to obtain a 3D
visualization of biofilm viability, starting at the base of the spherical particle where growth
started and finishing at the top of the sphere where the growth ended.

Z-stacks were analyzed using FIJI.[14] 3-D renderings were made by taking the .oir file raw data
from the microscope and merging the red and green channels. Following the merge, the channels
were elucidated by merging the stacks via image -> stacks -> z project with a "Max Intensity"
projection type.

Biofilm formation and Crystal Violet Staining
E. coli was inoculated in a 3-to-5-ml culture and grown to stationary phase for 24 hours. The
following day, sterile, nontreated 24-well plates (Celltreat ® #229524, Celltreat Scientific
Products, Pepperel, MA, USA) were inoculated in 400-µl LB medium per well. The plates were
covered and incubated statically at 37° C for 48 hours. After the inoculation period, cells were
stained per Merritt et al.'s protocol.[35] Well optical densities (OD) were then measured in a
spectrophotometer at a wavelength of 500 to 600 nm.
191
Motility Assay
Following the protocol by (Patridge and Harshey, 2020)*,* 0.3% Agar plates were made and
poured to the same thickness (20mL), and let dry for 1 hr.[36] Using a P20 tip, 5ul of culture was
pipetted, inserted into the agar about halfway through, and dispensed. Plates were incubated at
30 C and the diameter of the radial movement (indicated by the visible growth of cells) was then
measured using a ruler in centimeters (cm) after 20 hrs.

<u>Statistical Analysis</u>

The significance of MIC fold changes was determined using an ordinary one-way ANOVA.
Before performing the ANOVA, the residuals were tested for normality using the Shapiro-Wilk
test, confirming a Gaussian distribution. Each variable was then compared to the mean of a
control, which varied depending on the study. Multiple comparisons were corrected with the
Dunnett test, with P values adjusted accordingly. Finally, the residuals were tested for
homogeneity of variances and potential clustering through the Brown-Forsythe and Bartlett's
tests. The one-way ANOVA was chosen because it allows for comparing mean MIC fold
changes across groups under the assumption of normally distributed residuals, which was
confirmed here with the Shapiro-Wilk test. Dunnett's test was applied for post hoc comparisons,
as it is well-suited for comparing each group to a control with minimized type I error. Finally,
the Brown-Forsythe and Bartlett's tests were used to assess variance homogeneity, ensuring the
robustness of the ANOVA results.

Results

MPs impact growth at high concentrations

Eight different concentrations of MPs were grown with *E. coli* for 18 hours to determine whether MPs influence bacterial growth. The MPs used for testing concentration dependence were 10-µm diameter polystyrene spheres at concentrations of 100 MP/µL, 500 MP/µL, 1,000 MP/µL, 2,000 MP/µL, 4,000 MP/µL, 5,000 MP/µL, 10,000 MP/µL, and 15,000 MP/µL. Colony-forming units per mL (CFU/mL) were tested at one hour and 24 hrs after growth in the different concentrations. After one hour of growth, there were no significant differences in growth between the concentrations; however, at 24 hours, two MP solutions (10,000 and 15,000 MP/µL) showed a significantly lower number of CFU/mL compared to the WT (Fig. 1A, 1B). Notably, counts were higher in vortexed samples at 24 hrs, indicating the presence of surface-attached cells on the MPs, which is corroborated by confocal imaging (Fig. S1).

---**Insert Fig 1**--

Exposure to polystyrene MPs increases MDR.

We next sought to determine if cells exposed to MPs and subinhibitory levels of a given antibiotic, another common environmental contaminant, had altered patterns of resistance to other classes of antibiotics, rendering the bacteria multidrug-resistant. The four antibiotics we tested were ampicillin (β-lactam antibiotic), ciprofloxacin (fluoroquinolone), doxycycline (tetracycline), and streptomycin (aminoglycoside), which are all commonly found within the environment.[12,15] First, we measured changes in MIC in cells grown with MPs (500 µm diameter polystyrene spheres due to their ease of passaging) alone, to determine the role of MP on AMR in the absence of antibiotics (Fig. 2). In these susceptibility assays, specifically, we are testing the MIC of effluent containing attached cells released from the MPs upon vortexing (Fig. S1).

--**Insert Fig 2** --

Strikingly, the presence of MPs alone led to increased resistance to all the antibiotics listed above compared to media containing no MPs. While the MIC of bacteria grown in only LB stayed relatively stable, the samples co-cultured with MPs displayed a significantly higher resistance at a faster rate (Fig. 2). Most notably, the cells from the passaged MPs samples had an almost 100-fold higher MIC by day ten compared to day zero for ciprofloxacin, doxycycline and streptomycin (Table S.1). Next, we measured the MIC of cells grown with a single subinhibitory antibiotic with or without MPs (500 µm diameter polystyrene spheres).

--**Insert Fig 3** --

Our results suggest that the addition of MPs led to an increase in AMR for nearly all antibiotics. In the first four rows of Fig. 3, we show the fold change of antibiotics relative to WT for bacteria

grown in either antibiotics alone or antibiotics and MPs. In each case where bacteria were grown
and tested in the same antibiotic, the addition of MPs to antibiotics in the media led to an MIC
increase of at least 5 times more compared to cells grown in the antibiotics alone. Interestingly,
bacteria grown in ciprofloxacin with MPs generally had higher levels of multidrug resistance (up
to 171 times that of the control grown in the antibiotic) (Fig. 3, Table S.1). The other antibiotics
with MPs displayed resistance to ciprofloxacin of up to 75-fold higher than the antibiotic control
(Table S.1). Additionally, bacteria grown in streptomycin with MPs developed extremely high
levels of resistance to ciprofloxacin, doxycycline, and streptomycin (Table S.1).

After ten days of exposure to subinhibitory antibiotics, exposure to subinhibitory antibiotics was
halted, and bacteria were grown in antibiotic-free media for five days. The MIC was tested daily
to determine if resistance was stable. Of the sixteen conditions, 81.25% (13 out of 16) of the
bacteria grown in MPs and subinhibitory antibiotics retained the day ten resistance to their
respective antibiotic or even gained resistance (Fig. S.3). Conversely, 43.75% (7 out of 16) of the
bacteria grown in just MPs retained or grew in resistance to their respective antibiotics.
Surprisingly, only 18.75% (3 out of 16) of the bacteria grown in the subinhibitory antibiotics
alone retained resistance. Two of the three conditions that expressed stable resistance was grown
in doxycycline (Fig. S.3).

Exposure to different plastic characteristics (concentration and size) does not affect
resistance.

Next, we sought to investigate whether different MP characteristics (namely size and
concentration) affected antibiotic resistance. For simplicity, we used ciprofloxacin in these
experiments as the antibiotic showed the highest resistance changes in the abovementioned
studies. Thus, the effect of MP concentration (number of plastics/μL) on the development of
ciprofloxacin resistance was investigated under subinhibitory ciprofloxacin conditions. To
empirically understand whether the concentration of MPs affected ciprofloxacin resistance
development and magnitude of resistance, we exposed the bacteria to subinhibitory levels of
ciprofloxacin (7.5e-6 mg/mL, 40% of the initial MIC detailed above) and different
concentrations of polystyrene MPs. Resistance was determined in MIC fold changes relative to
the control (WT *E. coli* grown without MPs or subinhibitory levels of antibiotics) which were
tested every 48 hours.

The concentrations of MPs tested with the 500-μm diameter polystyrene spheres were 40 MP/mL
and 100 MP/mL. Conversely, the concentrations tested with the 10-μm polystyrene diameter
beads were 1,000 MP/μL, 500 MP/μL, 100 MP/μL, and 10 MP/μL. There was a significant
difference in ciprofloxacin MIC fold changes between each concentration of MP and the MP-
free counterpart (bacteria with subinhibitory levels of ciprofloxacin but no MPs) as well as the
WT (no MPs or antibiotics introduced during growth) (Fig. 4A, C), however, there was no
significant difference in changes in ciprofloxacin MIC between the different concentrations of

MPs (Fig. 4B, D); however, the absolute MIC values increased up to 3.713 ± 0.66 µg/mL of
ciprofloxacin—over three times the Clinical & Laboratory Standards Institute (CLSI) defined 1
µg/mL clinical breakpoint—for the larger diameter beads and 2.508 ± 4 µg/mL of ciprofloxacin
for the smaller beads (Table S.3 and S.4).[34]

--**Insert Fig 4** ---

MIC variation based on MPs' size and surface area was investigated following the results above.
The sizes of polystyrene MPs investigated were 5 µm (100 MP/µL), 10 µm (100 MP/µL), and
500 µm (40 MP/µL) diameter spheres. All three size variations were exposed to subinhibitory
ciprofloxacin along with a control that had no MPs added. Results showed that the difference
between the MPs themselves had no significant statistical difference. While the results showed
no significant difference between the different surface areas of the beads, the cells grown with
MPs and antibiotics again consistently had higher absolute MIC values (above the clinical
breakpoint) compared to the antibiotic-only comparators (Fig. 5A, 5B) and the WT.

--**Insert Fig 5** ---

Plastic compositions affects antibiotic resistance.

Next, we sought to determine if plastic composition affected the development and magnitude of
ciprofloxacin resistance (Fig. 6). This experimental design tested and compared the three most
common plastic types–polystyrene, polyethylene, and polypropylene.[24] We found that all three
conditions with varying MP compositions had a significantly higher MIC after ten days of
exposure than the MP-free control exposed to only subinhibitory levels of ciprofloxacin. The
polystyrene samples reached a significantly different absolute MIC value (1.65 ± 0.572 µg/mL)
than polyethylene (0.743 ± 0.36 µg/mL), polypropylene (0.528 ± 0.162 µg/mL) and the control
of only ciprofloxacin (0.20625 ± 0.04 µg/mL) (Fig. 6, Table S5). Furthermore, the polystyrene
MPs facilitated an absolute MIC higher than the ciprofloxacin-*E. coli* clinical breakpoint (1
µg/mL). Interestingly, the rates of AMR growth in all three plastic samples were similar in terms
of fold change (Fig. 6B).

--**Insert Fig 6** ---

Increased resistance and biofilm on polystyrene compared to glass

We also studied the difference in resistance development between polystyrene MP and glass
spheres (500-µm diameter spheres at the same concentration of 40 MP/µL) to determine if plastic
substrates had a specific effect or if any small particles would lead to an increase in AMR (Fig.
7). Similar to previous experiments, the microparticles were exposed to subinhibitory levels of

ciprofloxacin and compared to a control, which only had the subinhibitory ciprofloxacin and no glass or plastic particles. The polystyrene plastic spheres facilitated a statistically higher absolute MIC value by the end of the ten-day study from its glass counterpart and antibiotic control (Fig. 7A). Like Figure 6, the fold changes of the attachment surfaces were not statistically different (Fig. 7B). This indicates that the rate of AMR growth is similar in terms of fold change.

---**Insert Fig 7** --

To elucidate the mechanism behind the high levels of resistance found in the samples containing the MPs (particularly polystyrene, Fig. 2 and 3) and the discrepancies behind the resistance profiles of the particles of different compositions, we wanted to visualize the surface of the particles. We did this qualitatively by using confocal microscopy. The glass and polystyrene spheres were dyed on day 10 of subinhibitory ciprofloxacin exposure and compared to each other (Fig. 8) using a live dead stain. Green pixels indicate viable growth (e.g., live cells). In contrast, red indicates nonviable growth (dead cells). The glass (Fig. 8A) depicts two spheres, each having a significantly decreased amount of growth on the surface of the sphere, and most of the growth exhibited was colonized by dead cells. Conversely, the polystyrene condition had mostly live cells, and bacteria colonized the whole surface.

---**Insert Fig 8** --

Exposure to MPs select for increased biofilm

Following the visualization of robust biofilms on the surfaces of the polystyrene MPs, we sought to understand the role of biofilm further. Specifically, we investigated whether cells passaged with MPs (Fig. 2 and 3) formed more robust biofilms in the absence of MPs (i.e. does the presence of MPs select for better biofilm formers). Crystal violet staining can detect biofilm formation as it binds to bacterial cells and the extracellular matrix, making biofilms both visible and quantifiable. The bacteria grown with and without MPs along with various antibiotics and controls for 10 days (Fig. 2 and 3) were purified of MPs and trace antibiotics and grown in a 24-well polystyrene dish for 48 hours, and biofilm (surface attachment) was quantified using 0.1% crystal violet solution (Fig. 9).

---Insert Figure 9---

All samples grown with MPs had significantly more biofilm growth than those grown without MPs, indicated by their higher OD readings and increase in stained surface-attached cells (Fig. 9, Fig. S.4). Bacterial motility is a broad mechanism involved in biofilm formation; specifically, impaired motility is associated with increased biofilm due to several interconnected factors, mainly related to regulatory shifts, changes in surface properties, and environmental sensing mechanisms.[37] Thus, swimming motility was also assayed using the bacteria grown with and without MPs along with various antibiotics and controls for 10 days (Fig.2 and 3) on 0.3% soft

agar. Cells not grown with MPs had significantly larger diameters than those grown with MPs,
indicating that the bacteria exposed to MPs had impaired motility (Fig. 10, Fig. S.5).

--Insert Figure 10--

Discussion

MPs are significant environmental pollutants with critical implications for public health,
particularly in the context of increasing AMR. Understanding how MPs affect the development
of AMR is a crucial step in better understanding how environmental factors shape AMR in
bacteria. This will only become more important as both MPs and AMR become more prevalent.
Our study showed that MPs significantly impact the AMR rate of development and its
magnitude. More specifically, MPs facilitated post-breakpoint multidrug resistance to four
distinct families of antibiotics (Fig. 2A, 2B). Breakpoints are an integral part of modern
microbiology practice and define susceptibility and resistance to antibacterials in the clinical
setting (e.g., related to human health).[16]

Polystyrene, in particular, had the most significant impact on resistance development, which was
surprising given its relative hydrophilicity compared to polyethylene and polypropylene (Fig. 6).
Previous studies have shown that bacteria have an affinity for plastic due to their high
hydrophobicity and oxidation, which can lead to easy adhesion.[31, 32] However, it is essential to
note that studies have shown that *E. coli* prefers hydrophilic surfaces over hydrophobic ones.[17]
Of the three plastics used in this study, polystyrene is the most hydrophilic, while polypropylene
is the most hydrophobic, so these findings are in line with *E. coli's* preference to hydrophilic
surfaces. With this in mind, we expected glass beads of similar diameter to have a higher
adhesion rate and, therefore, a higher resistance to the tested antimicrobials. This, however, was
not the case, as polystyrene had a higher absolute MIC value and greater MIC fold change over
the glass condition. This indicates that plastics may be a unique substrate for bacteria to develop
and maintain resistance to.

While the complete mechanism is not yet known for AMR on MPs, the current prevailing theory
indicates that biofilm formation upon the plastics allows for higher resistance rates.[29] As
depicted in the CLSM images in Fig. 8, we observed a dense biofilm that spans the entire surface
of the MPs, compared to the glass substrate, which has uneven clusters of cells on the surface,
with a large portion of them nonviable post-subinhibitory antibiotic exposure. This would help
explain the differences in MIC shown in Fig. 7. We investigated the effects the cells had post-
MP exposure to see if they had a propensity for forming biofilms over their counterparts grown
in the same media without MPs. Qualitatively, confocal imaging found that not only did the
bacteria grown with MPs develop biofilms, but they grew biofilms with more biomass on the
plastics than glass surfaces (Fig. 8). Notably, cells grown in subinhibitory levels of doxycycline
and MPs had the highest biofilm growth. Doxycycline was the only bacteriostatic antibiotic—an
antibiotic that attacks cell reproduction rather than the cell itself—used in this study.

Doxycycline is known to target the 30S ribosomal subunit and inhibit protein synthesis, which
can trigger stress pathways that upregulate biofilm-associated genes and extra polymeric
substance production.[41] We next examined bacterial motility to investigate further the bacteria's
ability to grow biofilms. In these experiments, we found that samples grown with MPs had
impaired motility, which can influence biofilm formation (Fig. 10).[37] Overall, these data suggest
that the presence of MP not only presents a surface for biofilm formation but also selects for
cells that are better at biofilm formation. We hypothesize that, especially in the presence of low
levels of antibiotics, there is a combinatorial effect between selection for antibiotic resistance
target genes and biofilm formation that leads to high levels of resistance and recalcitrance.
Further research is needed to investigate the cellular-level interactions and material properties of
plastics that contribute to such high resistance rates.

It is well known that biofilms play a crucial role in the spread of AMR. Bacteria within biofilms
produce persister cells that are metabolically inert, which is one mechanism for evading
antibiotics.[22] These cells can survive even in high concentrations of antibiotics.[22] Furthermore,
current research suggests that biofilms act as refugia of MDR plasmids by retaining them, even
in the absence of antibiotics.[23.] This would support our results with respect to the MP properties
we investigated. First, MP concentration was not factored into different resistance rates. Instead,
size and composition had more of an impact on the rate of resistance development and
magnitude of resistance. A larger-sized MP would have a larger biofilm and, therefore, a greater
capacity to develop resistance. The plastic composition can also play a role in both bacterial
attachment and biofilm growth. MPs are known to serve as electron donors for bacterial biofilms
to feed on, inducing a faster or easier attachment of the bacteria to the surface and promoting
bacterial growth and colonization.[31, 35] This may explain the higher concentration of biofilm on
polystyrene MP compared to glass (Fig. 7), which in turn would explain polystyrene's larger
resistance load.

The proposed mechanisms of biofilms and selective pressures are assumed to have accumulated
overtime and compounded off of each other, creating high resistance levels as the time trial went
on. We believe that the potential ramifications of high-level multidrug-resistant bacteria
facilitated by the addition of MPs are significant. Moreover, we found bacteria exhibiting this
behavior within ten days of subinhibitory antibiotics and MP exposure. The rate of AMR
development and the surpassing of clinical breakpoints in both single and multidrug tests
highlight a need to monitor MPs and antibiotic levels in the environment. This is especially true
in areas with inadequate waste disposal and substandard public health infrastructure, such as
low-and middle-income countries (LMICs) and vulnerable populations.[18] Future studies in this
area should focus on the disparities of wastewater treatment in LMICs and high-income
countries and how the different environmental factors shape AMR development. Additionally,
wastewater and environmental monitoring should also include the presence of MPs, as they have
the potential to exacerbate AMR outbreaks. Our work can inform the ongoing development of
AMR surveillance strategies, helping to predict and prevent future outbreaks.

Acknowledgments
Thank you to Matt Lim for guidance at the confocal microscope and Boston University's shared CORE facility for the usage of their instruments.

Funding
This study was made possible by NSF GRFP grant number 2023354788 and supported by funds from the College of Engineering and Center on Forced Displacement at Boston University. The funders had no role in study design, data collection and interpretation, or the discussion to submit the work for publication.

CRediT Statement:
NG: Conceptualization, Methodology, Investigation, Writing-Original Draft, Writing- Review and Editing
JM: Investigation, Writing-Review and Editing
CC: Conceptualization, Writing-Review and Editing
BG: Investigation, Writing-Review and Editing
EH: Writing-Review and Editing
YN: Writing- Review and Editing
MHZ: Writing-Review and Editing

Competing Interests
Authors have no competing interests to declare.

Data Sharing
All data used for this study has been included in the manuscript or supplementary material.

References

1. Lebreton, L. & Andrady, A. Future scenarios of global plastic waste generation and disposal. *Palgrave Communications* **5**, 1–11 (2019).
2. Microplastics: Finding a consensus on the definition. *Mar. Pollut. Bull.* **138**, 145–147 (2019).
3. Microplastics in freshwater and terrestrial environments: Evaluating the current understanding to identify the knowledge gaps and future research priorities. *Sci. Total Environ.* **586**, 127–141 (2017).
4. Pham, C. K. *et al.* Marine Litter Distribution and Density in European Seas, from the Shelves to Deep Basins. *PLoS One* **9**, e95839 (2014).
5. Reaching New Heights in Plastic Pollution—Preliminary Findings of Microplastics on Mount Everest. *One Earth* **3**, 621–630 (2020).
6. Microplastics in the environment: A critical review of current understanding and identification of future research needs. *Environ. Pollut.* **254**, 113011 (2019).
7. Impact of wastewater from different sources on the prevalence of antimicrobial-resistant Escherichia coli in sewage treatment plants in South India. *Ecotoxicol. Environ. Saf.* **115**, 203–208 (2015).
8. Sulis, G., Sayood, S. & Gandra, S. Antimicrobial resistance in low- and middle-income

countries: current status and future directions. *Expert Rev. Anti. Infect. Ther.* (2022) doi:10.1080/14787210.2021.1951705.

9. Fouz, N. *et al.* The Contribution of Wastewater to the Transmission of Antimicrobial Resistance in the Environment: Implications of Mass Gathering Settings. *Tropical Medicine and Infectious Disease* **5**, 33 (2020).

10. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* **399**, 629–655 (2022).

11. Wastewater plastisphere enhances antibiotic resistant elements, bacterial pathogens, and toxicological impacts in the environment. *Sci. Total Environ.* **841**, 156805 (2022).

12. Sutradhar, I. *et al.* Computational Model To Quantify the Growth of Antibiotic-Resistant Bacteria in Wastewater. *mSystems* **6**, e0036021 (2021).

13. Brewster, J. D. A simple micro-growth assay for enumerating bacteria. *J. Microbiol. Methods* **53**, 77–86 (2003).

14. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

15. Sutradhar, I. *et al.* Effects of antibiotic interaction on antimicrobial resistance development in wastewater. *Sci. Rep.* **13**, 7801 (2023).

16. Park, G. E. *et al.* Clinical Impact of Revised Ciprofloxacin Breakpoint in Patients with Urinary Tract Infections by Enterobacteriaceae. *Antibiotics* **10**, (2021).

17. Hamadi, F. *et al.* The relation between Escherichia coli surface functional groups' composition and their physicochemical properties. *Braz. J. Microbiol.* **39**, 10–15 (2008).

18. Ferronato, N. & Torretta, V. Waste Mismanagement in Developing Countries: A Review of Global Issues. *Int. J. Environ. Res. Public Health* **16**, (2019).

19. Tuvo B, Scarpaci M, Bracaloni S, Esposito E, Costa AL, Ioppolo M, Casini B. Microplastics and Antibiotic Resistance: The Magnitude of the Problem and the Emerging Role of Hospital Wastewater. Int J Environ Res Public Health. 2023 May 18;20(10):5868. doi: 10.3390/ijerph20105868. PMID: 37239594; PMCID: PMC10218389.

20. Brewster JD. A simple micro-growth assay for enumerating bacteria. J Microbiol Methods. 2003 Apr;53(1):77-86. doi: 10.1016/s0167-7012(02)00226-9. PMID: 12609726.

21. CLSI. [CLSI vs. FDA Breakpoints, 1st ed. CLSI guideline. Clinical and Laboratory Standards Institute; [2023].

22. Abebe GM. The Role of Bacterial Biofilm in Antibiotic Resistance and Food Contamination. Int J Microbiol. 2020 Aug 25;2020:1705814. doi: 10.1155/2020/1705814. PMID: 32908520; PMCID: PMC7468660.

23. Metzger, G.A., Ridenhour, B.J., France, M. *et al.* Biofilms preserve the transmissibility of a multi-drug resistance plasmid. *npj Biofilms Microbiomes* **8**, 95 (2022). https://doi.org/10.1038/s41522-022-00357-1

24. Joint Group of Experts on the Scientific Aspects of Marine Environmental Protection (GESAMP). 2015. Sources, fate and effects of microplastics in the marine environment: a global assessment. Kershaw, P. J., ed. (IMO/FAO/UNESCO-IOC/UNIDO/WMO/IAEA/UN/UNEP/UNDP). Rep. Stud. GESAMP No. 90, 96 pp.

25. Jenna R. Jambeck *et al.*, Plastic waste inputs from land into the ocean. *Science* **347**, 768-771
(2015). DOI:10.1126/science.1260352

26 De Souza Machado AA, Kloas W, Zarfl C, Hempel S, Rillig MC. Microplastics as an
emerging threat to terrestrial ecosystems. *Glob Change Biol*. 2018; 24: 1405–
1416. https://doi.org/10.1111/gcb.14020

27 Jose Luis Martinez, Environmental pollution by antibiotics and by antibiotic resistance
determinants, *Environmental Pollution*, Volume 157, Issue 11, 2009,
https://doi.org/10.1016/j.envpol.2009.05.051.

28. Huixiang Wang, Kaiwen Xu, Jing Wang, Chong Feng, Yihan Chen, Jianghong Shi, Yan
Ding, Chengxun Deng, Xiaowei Liu, Microplastic biofilm: An important microniche that may
accelerate the spread of antibiotic resistance genes via natural transformation, *Journal of
Hazardous Materials*, 2023, https://doi.org/10.1016/j.jhazmat.2023.132085.

29. María Belén Sathicq, Raffaella Sabatino, Gianluca Corno, Andrea Di Cesare. Are
Microplastic Particles a Hotspot for the Spread and the Persistence of Antibiotic Resistance in
Aquatic Systems? 2021. *Environmental Pollution* 279 (June): 116896.

30 Yuanze Sun, Na Cao, Chongxue Duan, Qian Wang, Changfeng Ding, Jie Wang. "Selection of
Antibiotic Resistance Genes on Biodegradable and Non-Biodegradable Microplastics." 2021.
Journal of Hazardous Materials 409 (May): 124979.

31. Debroy, A., George, N. and Mukherjee, G. (2022), Role of biofilms in the degradation of
microplastics in aquatic environments. *J Chem Technol Biotechnology*, 97: 3271-3282.
https://doi.org/10.1016/j.jhazmat.2023.132085

33. Lihua Niu, Yuanyuan Li, Yi Li, Qing Hu, Chao Wang, Jiaxin Hu, Wenlong Zhang, Longfei
Wang, Chi Zhang, Huanjun Zhang. New insights into the vertical distribution and microbial
degradation of microplastics in urban river sediments, *Water Research*, 2021,
https://doi.org/10.1016/j.watres.2020.116449.

34. Chantell, Cristina, Humphries, Romney M., Lewis, James S. II, "Selection of Antibiotic
Resistance Genes on Biodegradable and Non-Biodegradable Microplastics" 2021,
https://clsi.org/media/3011/mr02ed1.pdf

35. Merritt JH, Kadouri DE, O'Toole GA. Growing and analyzing static biofilms. Curr Protoc
Microbiol. 2005 Jul;Chapter 1:Unit 1B.1. doi: 10.1002/9780471729259.mc01b01s00. PMID:
18770545; PMCID: PMC4568995.

36. Partridge, J. D., Harshey, R. M. Investigating Flagella-Driven Motility in *Escherichia coli* by
Applying Three Established Techniques in a Series. *J. Vis. Exp.* (159), e61364,
doi:10.3791/61364 (2020)

37. Benyoussef W, Deforet M, Monmeyran A, Henry N. Flagellar Motility During E. coli
Biofilm Formation Provides a Competitive Disadvantage Which Recedes in the Presence of Co-
Colonizers. Front Cell Infect Microbiol. 2022 Jul 8;12:896898. doi: 10.3389/fcimb.2022.896898.
PMID: 35880077; PMCID: PMC9307998.

38. Joannard, B., Sanchez-Cid, C. Bacterial dynamics of the plastisphere microbiome exposed to
sub-lethal antibiotic pollution. *Microbiome* **12**, 97 (2024). https://doi.org/10.1186/s40168-024-
01803-2

39. Zadjelovic V, Wright RJ, Borsetto C, Quartey J, Cairns TN, Langille MGI, Wellington EMH,
Christie-Oleza JA. Microbial hitchhikers harbouring antimicrobial-resistance genes in the
riverine plastisphere. Microbiome. 2023 Nov 1;11(1):225. doi: 10.1186/s40168-023-01662-3.
PMID: 37908022; PMCID: PMC10619285.

40. Hung, Chia, Yizhou Zhou, Jerome S. Pinkner, Karen W. Dodson, Jan R. Crowley, John
Heuser, Matthew R. Chapman, Maria Hadjifrangiskou, Jeffrey P. Henderson, and Scott J.
Hultgren. 2013. "Escherichia Coli Biofilms Have an Organized and Complex Extracellular
Matrix Structure." mBio, September. https://doi.org/10.1128/mbio.00645-13.

41. Chukwudi, C.U., Good, L. Doxycycline inhibits pre-rRNA processing and mature rRNA
formation in *E. coli*. *J Antibiot* **72**, 225–236 (2019). https://doi.org/10.1038/s41429-019-0149-0

Figures:

Figure 1. *CFU/mL readings after 1 hour of growth (A) and 24 hours of growth (B) in WT and*
MP concentrations (100, 500, 1000, 2000, 4000, 5000, 10000 and 15000 MP/μL; 10μm
diameter polystyrene spheres). Samples that were vortexed are presented in black and non-
vortexed samples are depicted in red.

Figure 2. *Normalized 10-day time (x-axis) series fold change (y-axis) of 500-μm diameter polystyrene MPs at 40 MP/mL, grown up in LB and then tested for the MIC in all four antibiotics (left to right: ampicillin, ciprofloxacin, doxycycline and streptomycin).*

Figure 3. *Normalized 10-day time (x-axis) series fold change (y-axis) of 500-μm diameter polystyrene MPs at 40 MP/mL, grown up in one of four antibiotics left to right: ampicillin,*

ciprofloxacin, doxycycline, and streptomycin and then tested for the MIC in top to bottom:

ampicillin, ciprofloxacin, doxycycline and streptomycin

Figure 4. *Absolute ciprofloxacin MIC values for 500-μm diameter polystyrene MPs at two*

different MP concentrations with the dashed line indicating the ciprofloxacin clinical breakpoint

concentration at Day 10 of exposure(A), time series fold change of 500-μm diameter polystyrene

MPs at different MP concentrations and the subinhibitory antibiotic control (ciprofloxacin)

relative to the WT (B), Absolute ciprofloxacin MIC values for 10-μm diameter polystyrene MPs

at four different MP concentrations with the dashed line indicating the ciprofloxacin clinical

breakpoint concentration at Day 10 of exposure (C)), time series fold change of 10-μm diameter

polystyrene MPs at different MP concentrations and the subinhibitory antibiotic control
(ciprofloxacin) relative to the WT (D)

Figure 5. *Absolute ciprofloxacin MIC values for various sized polystyrene MPs at 100 MP/μL*
with the dashed line indicating the ciprofloxacin clinical breakpoint concentration at Day 10 of
exposure (A), time series fold change of various sized polystyrene MPs at 100 MP/μL, and the
subinhibitory antibiotic control relative to the WT (B)

597
Figure 6. *Absolute ciprofloxacin MIC values for 10-μm diameter polystyrene, polyethylene, and*
polypropylene MPs at 100 MP/μL with the dashed line indicating the ciprofloxacin clinical
breakpoint concentration at Day 10 of exposure (A), time series fold change of 10-μm diameter
polystyrene, polyethylene, polypropylene MPs at 100 MP/μL, and the subinhibitory antibiotic
control relative to the WT (B)

Figure 7. *Absolute ciprofloxacin MIC values for 500-μm diameter polystyrene and glass spheres*
at 40 MP/mL at Day 10 of exposure (A), with the dashed line indicating the ciprofloxacin
clinical breakpoint concentration. Time series fold change of the 500-μm (B) glass, polystyrene,
and the subinhibitory antibiotic control relative to the WT.

Figure 8. *CLSM image of 500-μm diameter glass (left) and polystyrene (right) spheres at day 10*
of subinhibitory ciprofloxacin exposure. Green pixels indicate live cells, while red indicate dead
cells.

Figure 9. *OD readings of crystal violet staining for E. coli grown with or without MPs in media*
containing subinhibitory levels of four antibiotics left to right: ampicillin, ciprofloxacin,
doxycycline, or streptomycin or no antibiotics (LB).

Figure 10. *Motility in centimeters (cm) of E. coli grown with or without MPs in media*
containing subinhibitory levels of four antibiotics left to right: ampicillin, ciprofloxacin,
doxycycline, or streptomycin or no antibiotics (LB).

625