Dysregulation of cholesterol and bile acid homeostasis in the gut-liver axis following *Cryptosporidium* intestinal infection

Shuhong Wang[1], Ai-Yu Gong[1], Edward Barker[1], David L. Williams[1], Christopher Forsyth[2], Liqing Yu[3], and Xian-Ming Chen[1,4]

[1]Department of Microbial Pathogens and Immunity, Rush University Medical Center, Chicago, Illinois, USA; [2]Department of Anatomy and Cell Biology, Rush University Medical Center, Chicago, Illinois, USA; [3]Division of Endocrinology, Diabetes and Nutrition, Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland 21201, USA.

[4]Correspondence: xian_m_chen@rush.edu

Abstract

Cryptosporidium spp., an apicomplexan protozoan, is one of the most common pathogens causing moderate-to-severe diarrhea in children under 2 years of age and is also an important opportunistic pathogen for patients with AIDS. There are currently no effective vaccines or therapies available. Infection in children is associated with malnutrition, growth defects, and even impaired cognitive development, but underlying mechanisms remain unclear. We report here that *C. parvum* infection in neonatal mice impairs bile acid reabsorption in the ileum, disrupts lipid metabolism in the liver, and alters bile acid homeostasis in the enterohepatic circulation. The reduction of bile acid pool further impairs lipid absorption in the small intestine. Additionally, replenishing bile prevents the decline in lipid absorption in infected neonatal mice. Strikingly, bile gavage significantly reduces the infection burden and ameliorates the dysregulated homeostasis of cell proliferation and migration in intestinal epithelium following infection. These findings may guide novel therapeutic approaches for cryptosporidiosis.

Introduction

Cryptosporidium is a significant opportunistic intestinal pathogen for immunocompromised individuals and one of the leading causes of moderate-to-severe diarrhea in children under 2 years of age[1]. Infection with this parasite in young kids is associated with mortality, growth stunting, and developmental deficits[2]. Recent estimates from the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) indicate *Cryptosporidium* spp. is responsible for approximately 7.6 million cases and between 48,000 and 202,000 deaths annually among young children in low-resource settings[3,4]. Cryptosporidiosis in humans is primarily caused by *C. parvum* and *C. hominis* species[2]. Nitazoxanide, the only approved drug for the treatment of cryptosporidiosis, is ineffective in immunocompromised patients and malnourished children[2,5–7]. Pathologically, *Cryptosporidium* infections are primarily confined to the distal ileum and it is speculated that the combined loss of microvillus borders and villus height reduces the absorptive surface area of the intestine, leading to diarrhea and malnutrition in children[8]. However, the current understanding of the underlying molecular mechanisms of malnutrition remains unclear, hampering the development of effective therapeutics. The intestinal mucosa consists of a monolayer of rapidly self-renewing epithelial cells. Intestinal stem cells in the crypts generate new functional epithelial cells, which mature and differentiate as they move up the villus before being shed at the tip[9,10]. *Cryptosporidium* infection is confined to the villi (mainly enterocytes) of the intestinal epithelium, where the organism forms an intracellular yet extra-cytoplasmic parasitophorous vacuole at the brush border following cell entry[11]. Increased cell death during infection triggers crypt hyperplasia and branching due to accelerated cell division to compensate for the loss of cells at the villi[8,12], however, its underlying molecular mechanisms remain unknown.

As a major source of nutrition for infants, milk provides a significant amount of fatty acids, phospholipids, and cholesterol. Bile acids are amphipathic molecules synthesized from cholesterol in the liver, stored in the gallbladder, and released into the duodenum after a meal[13]. Bile acids facilitate the emulsification of insoluble lipids into tiny mixed micelles, promoting lipid digestion and absorption in the intestine[14]. More than 95% of the bile acids are reabsorbed in the terminal ileum through the apical sodium-dependent bile salt transporter (ASBT; SLC10A2) and recirculated to the liver via the portal blood in a process known as enterohepatic circulation[13]. Bile acids have been identified as endogenous ligands with high affinity for the

Farnesoid X receptor (FXR), a ligand-activated transcription factor in the nuclear receptor (NR)-
family [15–17]. Thus, the Bile acid-FXR network plays a crucial role in maintaining whole-body bile
acid homeostasis by regulating the expression of target genes involved in both hepatic bile acid
synthesis and enterohepatic circulation[18]. Using a murine neonatal infection model, we report
that *Cryptosporidium* infection significantly disrupts bile acid and lipid homeostasis in both the
intestine and liver. Functionally, bile intervention greatly reduces the infection burden, enhances
intestinal lipid absorption, and ameliorates dysregulated intestinal homeostasis. These findings
suggest that bile intervention may have therapeutic potential for intestinal cryptosporidiosis.
86
Results
87
C. parvum infection disrupts cholesterol homeostasis in the small intestine
88
Cryptosporidium lacks the capability to synthesize cholesterol and needs to acquire it both from
the gut lumen and the host cell to meet its metabolite[19,20]. Therefore, we investigated whether *C.*
parvum infection affects intestinal cholesterol absorption and transport. To explore this, we
performed RNA-seq analysis to examine genome-wide changes in gene expression in the ileal
epithelium of infected neonatal mice. Consistent with results from previous studies[21], RNA-Seq
analysis of the ileal epithelium from infected neonatal mice [5-days old, 72 h post-infection (p.i.)]
reveals significant changes in host gene expression (Supplementary Table 1). Gene set
enrichment analysis (GSEA) indicates marked alterations in pathways related to inflammatory
responses (Extended Data Fig. 2b), defense response (Extended Data Fig. 2c), and the regulation
of sterol homeostasis in the infected ileum epithelium (Extended Data Fig. 2d).
99
Cellular cholesterol homeostasis is finely regulated through the dynamic balance between
biosynthesis, uptake, export, and esterification[22]. The regulation of cellular cholesterol transport
involves low-density lipoprotein receptor (LDLR)-mediated absorption and chylomicron
secretion, as well as transmembrane influx/efflux through transporters such as Niemann-Pick C1-
Like 1 (NPC1L1), the heterodimeric ATP-binding cassette transporters G5 and G8 (ABCG5/G8),
and ATP-binding cassette transporter A1 (ABCA1) (Fig. 1a). This process is governed by
intricate feedback regulation mechanisms, including the liver X receptor (LXR) signaling[23].
Further GSEA analysis indicates that cholesterol efflux, import, and esterase activity were
significantly downregulated in the infected ileal epithelium (Fig. 1b and Extended Data Fig.
2a,d), while most genes involved in cholesterol biosynthesis were upregulated (Fig. 1b).
Specifically, reduced expression of several cholesterol efflux genes, including ABCG5, ABCG8
and ABCA1, was confirmed by reverse transcription–quantitative PCR (RT–qPCR) (Fig. 1d).
Immunoblot analyses confirmed reduced protein levels of ABCG5 and ABCG8 from the ileal
epithelium of infected mice (72 h and 96 h p.i.) compared to control animals (Fig. 1f). Two
LXRs, LXRα (NR1H3) and LXRβ (NR1H2), play key roles in the transcriptional regulation of
genes involved in cholesterol homeostasis[23]. Interestingly, the protein level of LXRβ was
decreased, while LXRα level remained unchanged in the ileum of infected neonates compared to
the control groups (72 h and 96 h p.i.) (Fig. 1f), suggesting repression of the LXR pathway
following infection. In contrast, most cholesterol biosynthesis genes were upregulated after *C.*
parvum infection (Fig. 1b,c), as validated by RT-qPCR (Fig. 1e). The rate-limiting enzyme in
cholesterol biosynthesis is 3-hydroxy-3-methylglutaryl-CoA reductase (HMGCR)[24]. Immunoblot
analyses of cytosolic protein extracts confirmed the higher levels of HMGCR protein in the
ileum of infected neonates compared to the control groups (Fig. 1f). Furthermore, we found a

significant reduction of free cholesterol in the ileum of infected neonatal mice compared to
control mice (Fig. 1h).
125
Strikingly, the proximal small intestine, which is primarily responsible for nutrient absorption
and is not infected with *C. parvum* (Extended Data Fig. 1a,b), also exhibited similar changes in
the expression levels of cholesterol homeostasis genes (Fig. 1g and Extended Data Fig. 2e).
Notably, RT-qPCR showed an increased expression of NPC1L1 in the proximal small intestine
but decreased expression in the ileum (Fig. 1d and Extended Data Fig. 2e). Furthermore, a
significant reduction in total cholesterol was detected in the proximal small intestinal epithelium
of infected neonates compared to the control groups (Fig. 3g and Extended Data Fig. 6a). Most
of the cholesterol absorbed by the intestine is secreted as chylomicrons into the circulation[25].
Expression of genes involved in the chylomicron assembly was significantly reduced in both the
proximal small intestine and ileum of infected animals (Extended Data Fig. 2f-h). Consistent
with results from previous studies[8,21], the villi length in the infected ileum was significantly
shorter (Extended Data Fig. 2j), whereas no such change was observed in the non-directly
infected region of the intestine (Extended Data Fig. 2j). Additionally, the small intestine length
in infected neonates was shorter compared to control animals (Extended Data Fig. 2i), suggesting
a reduction in the absorptive surface area in the intestine. Collectively, these findings indicate
that *C. parvum* infection disrupts cholesterol homeostasis in the small intestine.
142
***C. parvum* infection alters hepatic lipid metabolism**
The liver is the primary organ for cholesterol metabolism, and most lipids in the intestine are
absorbed by enterocytes in the small intestine, transported through the lymphatic system into the
circulation, then to the liver, and eventually to peripheral tissues[26,27]. Based on this, we
hypothesized that intestinal infection by *C. parvum* could impact cholesterol and bile acid
metabolism in the liver. To test this, we infected neonatal mice with *C. parvum* oocysts for 3
days, collected liver tissues, and performed RNA-seq (Fig. 2a). Consistent with our hypothesis,
C. parvum infection caused significant alterations in gene expression profiles in the liver
(Extended Data Fig. 3a and Supplementary Table 2). GSEA indicated that the major pathways
associated with the altered gene expression profiles are related to immune response regulation
(Extended Data Fig. 3a). Gene set variation analysis (GSVA) revealed an overall downregulation
of genes involved in lipid metabolism pathways in the liver of infected mice, including these
related to cholesterol and bile acid metabolism (Supplementary Table 2). RT-qPCR further
validated the reduced levels of most genes involved in lipid metabolism (Extended Data Fig. 3b-
f). Immunoblot analyses of APOB100 (Apolipoprotein B100), HMGCR, and HMGCS2
(hydroxymethylglutaryl CoA synthase 2) confirmed reduced protein levels in the infected
animals (Extended Data Fig. 3g). The protein levels of hepatic FXR and LXRβ were
significantly decreased, whereas LXRα remained unchanged after infection (Fig. 2d). Consistent
with this, the mRNA expression of most FXR and LXR target genes was significantly reduced in
the liver of infected mice (Fig. 2b,c). Immunoblot of selected canonical target genes of LXRs
and FXRs, including *ABCG5*, *ABCG8*, *CYP7A1* and *NR0B2* (*Shp*), confirmed decreased protein
levels (Fig. 2d). Accordingly, lipidomics analysis revealed altered lipid profiles in the liver of
infected mice (Fig. 2a and Extended Data Fig. 4a,b). There was no significant reduction in total
lipids in the liver after 4 days infection compared to the control group (Extended Data Fig. 4d,e).
However, reduced amounts of bile acids and cholesterol esters, but not total cholesterol, were
detected in the liver of infected mice (Fig. 2e and Extended Data Fig. 4f-h). As an important

component of milk, fat provides about 50% of the energy for infants[28]. The main fat component
is triacylglycerol (TG), which makes up more than 98% of milk fat[29]. We observed significant
reductions in most TGs, diacylglycerols (DG), and free fatty acids (FFA) in the liver after *C.*
parvum intestinal infection (Fig. 2e and Extended Data Fig. 4i-n). Kyoto Encyclopedia of Genes
and Genomes (KEGG) pathway enrichment analysis of the lipidomics results revealed
downregulation of lipid metabolic pathways in the liver of infected mice (Extended Data Fig. 4c),
consistent with the GSVA results from the RNA-seq (Supplementary Table 2). To further
investigate the alterations in bile acids, we performed a comprehensive, quantitative bile acid
targeted assay (Fig. 2a), which revealed broad metabolic downregulation in the liver of infected
mice (Fig. 2f-h). Taken together, these data suggest that *C. parvum* intestinal infection disturbs
cholesterol and bile acid homeostasis in the liver.
180
***C. parvum* infection decreases bile acid levels in the small intestine and reduces the**
absorption of dietary lipids
Given the significant dysregulation of cholesterol and bile acid homeostasis in the liver
following *C. parvum* intestinal infection, particularly the decreased expression of FXR-targeted
genes, we next investigated lipid metabolism with a focus on the Bile acid-FXR signaling
pathway in the infected intestine. FXR is essential for maintaining whole-body bile acid
homeostasis by regulating the expression of target genes involved in both hepatic bile acid
synthesis and enterohepatic circulation[18]. The enterohepatic circulation of bile acids and the FXR
pathway involve the synthesis of bile acids from cholesterol in the liver and the finely controlled
expression of genes for bile acids reabsorption in the intestinal epithelium, as shown in Fig. 3a.
GSEA indicates that many genes involved in the bile acid metabolic process in the ileum were
significantly reduced in infected mice (Fig. 3b). The reduction of FXR and FXR-targeted genes
Shp and *Fgf15* suggests a robust suppression of FXR signaling in the small intestine following
infection (Fig. 3b-d and Extended Data Fig. 5a). Bile acids generate negative feedback on ileal
ASBT transcription through FXR-mediated induction of *Shp*[30]. Notably, despite a substantial
increase in its mRNA expression (Fig. 3b,c), ASBT was markedly reduced at the protein level
(Fig. 3d), which could substantially impact bile reabsorption in the ileum.
198
To address whether dysregulation of liver and intestinal bile acid metabolism affects the bile
levels in the small intestine following infection, we infected neonatal mice with *C. parvum*
oocysts for 4 days. We then collected small intestine tissues, including the luminal contents, and
performed a comprehensive, quantitative bile acid targeted assay. A marked decrease of most
intestinal bile acids was observed in the infected mice (Fig. 3h and Extended Data Fig. 6d). In
addition, the ileum appears paler, possibly due to a reduction in intestinal bile acids following the
infection (Extended Data Fig. 5d). RNA-seq analysis of ileal tissues revealed a global decrease in
the expression of genes involved in lipid and vitamin absorption in the infected mice (Extended
Data Fig. 5g,h). Imaging of neutral lipids by oil red O (ORO) staining[31] showed a reduction in
neutral lipids in the proximal small intestine and ileum following infection (Fig. 3e). Consistent
with this, lipidomics analysis demonstrated a significant decrease in total lipids (Fig. 3f),
particularly bile acid, cholesterol, phosphatidylcholines (PC), hexosylceramides (HexCer-NS)
and Ceramide-NS (Cer-NS) in the proximal small intestinal epithelium of infected neonates (Fig.
3g and Extended Data Fig. 6a). However, lipidomics analysis showed no significant alteration in
total TG level (Fig. 3g and Extended Data Fig. 6a). Additionally, the proximal small intestinal
epithelium and ileum from infected neonates exhibited reduced expression of several genes

involved in fatty acid oxidation, TG catabolism (Extended Data Fig. 5b,c,e,f), and chylomicron assembly (Extended Data Fig. 2f-h). Furthermore, lower levels of L-palmitoylcarnitine were detected in the proximal small intestinal epithelium of *C. parvum* infected neonates (Extended Data Fig. 6c), indicating impaired fatty acid oxidation. These alterations may contribute to the accumulation of TG in the proximal small intestinal epithelium following infection. Linoleic acid is an essential lipid that cannot be synthesized in mammalian cells and is exclusively acquired from the diet[32]. The KEGG pathway enrichment analysis of lipidomics revealed a reduction in linoleic acid metabolism (Extended Data Fig. 6b), further supporting the reduced absorption of dietary lipids in the intestine after *C. parvum* infection.

Bile intervention shows therapeutic potential for intestinal cryptosporidiosis
To assess the impact of reduced bile on the pathogenesis of *C. parvum* infection, we administered whole bile collected from the gallbladders of adult mice to neonates (5 days old) at 24 h after oral gavage of *C. parvum* oocysts, followed by daily administration of bile for an additional 3 days (Fig. 4a). Bile feeding activated intestinal transcription of FXR target genes (Fig. 4c) and surprisingly, greatly reduced the infection burden compared to the animals receiving PBS (Fig. 4b). Our data revealed significant variation in infection burden across different regions, with the highest levels typically observed in the distal ileum (Fig. 4d). Notable reductions in the infected area of the distal ileum and overall reduced infection burden were observed in the intestinal tissues of animals receiving bile, compared to the PBS-treated control group (Fig. 4d). Additionally, replenishing bile restored the ileal expression of genes involved in cholesterol homeostasis in the infected mice (Extended Data Fig. 7a,b). Furthermore, human intestinal epithelial HCT-8 cells were first exposed to *C. parvum* infection for 4 h (a time frame for parasite attachment and invasion)[33], after which free-parasites were removed after washing. Cell cultures were then exposed to various concentrations of bile or PBS for an additional 20 h, followed by measurement of infection burden. A substantial reduction in infection burden was observed in cells treated with higher concentrations of bile (0.5% and 1%) compared to lower bile concentrations (0.2%, 0.25% and 0.33%) (Fig. 4e,f). Interestingly, an increased infection burden was noted in cultures treated with lower concentrations of bile compared to the PBS group (Fig. 4e,f). Primary 2D intestinal epithelial monolayers derived from mice for *C. parvum ex vivo* infection further confirmed this phenomenon (Extended Data Fig. 7e). Data from previous reports show that bile salt sodium taurocholate enhances excystation and the entry of *Cryptosporidium* into host cells[34]. Consistent with this, we observed an overall increase in infection burden when HCT-8 cells were exposed to *C. parvum* oocysts in the presence of various bile concentrations (from 0.07%-2% bile acids as tested) during the 4 h parasite attachment and invasion time frame (Extended Data Fig. 7d). HCT-8 cells exposed to different concentrations of bile, with or without *C. parvum* infection, showed no significant reduction in cell viability (Extended Data Fig. 8a-d). To further assess the therapeutic potential of bile for cryptosporidiosis. We orally inoculated neonatal mice with excessive dose of *C. parvum* (2×10^6 oocysts per animal). At 24 h after *C. parvum* infection, mice were fed different doses of whole bile collected from the gallbladders of adult mice (0.5, 1.5, and 3 μl of bile per animal, diluted with PBS), followed by daily for another 3 days (Fig. 5a). Resultant data further support a role for bile in reducing infection and increasing lipid uptake following *C. parvum* infection (Fig. 5b-f and Extended Data Fig. 7c). Specifically, feeding 1.5 and 3 μl of bile per day for 4 days activated intestinal transcription of FXR target genes (Fig. 5e) and greatly decreased the infection burden compared to the animals receiving PBS (Fig. 5d,f and Extended Data Fig. 7c),

while feeding 0.5 µl of bile per day did not cause significant changes (Fig. 5b,c). Notably, in the presence of excessive dose of *C. parvum* inoculation, treatment with 3 µl of bile per day for 4 days greatly reduced the infected area from distal ileum in mice, which was negatively correlated with the area and extent of intestinal lipid absorption (Fig. 5f and Extended Data Fig. 7c).

Bile acid-FXR signaling pathway is involved in the regulation of intestinal epithelium turnover in response to *C. parvum* infection

To monitor the cell proliferation and epithelial turnover in the intestine following infection, we examined the incorporation of ethynyl-2'-deoxyuridine (EdU) and tracked the migration of EdU-labeled cells at 4 and 48 h after EdU pulse labeling (Fig. 6a). To explore the potential effects of bile, we gavaged infected and uninfected mice with the whole bile collected from the gallbladders of adult mice (Fig. 6a). As shown in Fig. 6e and Extended Data Fig. 9a, there was a significant increase in the number of EdU-positive proliferating cells in the crypt regions of infected mice at 4 h. These proliferative cells also migrated much higher along the villus compared to their control counterparts at 48 h p.i. (Fig. 6g,j). Interestingly, quantification showed an appreciable decrease in EdU-positive proliferating and migrating cells in infected mice treated with bile compared to those treated with PBS (Fig. 6e,g,j and Extended Data Fig. 9a). Notably, we found that the dysregulation of cell proliferation and migration induced by infection, as well as the effects of bile supplement, were not limited to the ileum but extended to the entire small intestine and even the colon (Fig. 6c-j and Extended Data Fig. 9a).

There was a significant increase in the transcription levels of FXR target genes in infected mice treated with bile compared to those treated with PBS (Fig. 4c and Fig. 5e). We then examined whether activation of FXR pathway could attenuate crypt hyperproliferation and cell migration *in vivo*. We administrated mice with Fexaramine (FexD) (Fig. 7a), which has previously been shown to activate FXR pathways *in vivo* [35]. Quantification showed a significant decrease in EdU-positive proliferating and migrating cells (jejunum and ileum) in uninfected mice treated with FexD compared to those treated with vehicle control (Fig. 7b-g). This was consistent with previous literature that the activation of FXR pathway inversely correlates with epithelial proliferation, colorectal cancer progression and malignancy [36–38]. Interestingly, EdU-labeling analysis demonstrated an even more pronounced decrease in EdU-positive proliferating and migrating cells between infected mice treated with FexD and vehicles, compared to those uninfected mice treated with FexD and vehicles (Fig. 7b-g). Notably, FXR activation significantly increased the infection burden (Fig. 7h,i), further emphasizing the critical role of FXR signaling in maintaining crypt-villus homeostasis during *Cryptosporidium* infection. Importantly, this effect appears to be independent of the infection burden. RT-PCR indicated that bile treatment for 6 h can activate intestinal transcription of FXR target genes in uninfected mice (Fig. 6b). However, in contrast to *C. parvum* infected mice, no significant differences were observed in uninfected mice treated with bile and PBS (Fig. 6c-j and Extended Data Fig. 9a). The negative feedback of bile acid-FXR axis might restrain the constant activation of FXR pathway in uninfected mice [18]. Overall, these data suggest that the Bile acid-FXR signaling pathway is involved in regulating cell proliferation and migration of the intestinal epithelium following *C. parvum* infection.

Discussion

The preferred site of *Cryptosporidium* infection is the ileum, where sporozoites penetrate individual epithelial cells (mainly enterocytes). After cell entry, they form a unique apical localization (intracellular but extra-cytoplasmic), preventing the infection from further spreading to deeper epithelial tissues or entering the circulation and infecting other organs[39,40]. The significant alterations in gene expression profiles and metabolic processes observed in the liver, and proximal small intestine - regions where no parasites were detected - suggest that the pathological changes extend beyond the directly infected ileum. From a pathological perspective, the metabolic dysregulation observed beyond the ileum may play an important role in the growth stunting and malnutrition seen in young children following cryptosporidial infection.

The correlation between lipid and bile acid metabolic dysregulation observed in the liver and intestine from infected mice provides further evidence of the role of the gut-liver axis in the pathogenesis of various enteritis. Given the preferred infection site of *Cryptosporidium* in the ileum, where over 95% of the bile acids are reabsorbed[13], it maybe pathologically significant that the expression of ASBT is downregulated at the protein level in the infected ileum. This downregulation could reduce bile acid reabsorption, contributing to the decreased bile acid levels in the enterohepatic circulation. Depletion of intestinal cholesterol may also contribute to the decreased bile acid pool, as ASBT function depends on the cholesterol content of lipid raft domains in the plasma membrane[41]. Moreover, the decreased absorptive surface area in the intestine, due to the loss of microvilli and shorten of villus height and intestinal length following *Cryptosporidium* infection, could decrease the brush border apical surface. Surprisingly, the reduction of the bile acid pool and the downregulation of the FXR pathway in the enterohepatic circulation following *C. parvum* infection did not lead to an increase in hepatic bile synthesis as compensation. Instead, there was an overall decrease in bile acid metabolism in the liver. Hepatic bile synthesis generally accounts for only a small fraction of the bile acid pool, as over 95% bile acids are conserved through intestinal reabsorption in the enterohepatic circulation[42].

While most bile acids are transported back into the liver via enterohepatic circulation, a small fraction of this pool (roughly 5%) escapes reabsorption in the ileum and is further converted into secondary bile acids in the colon by the gut microbiota[13]. Previous studies have demonstrated controversial effects of several bile acids on *Cryptosporidium* infection. Three secondary bile acids (deoxycholate, deoxycholic acid, and lithocholic acid) have been reported to impair *Cryptosporidium* growth *in vitro*[43]. The bile salt sodium taurocholate enhances microneme secretion and increases the entry of *Cryptosporidium* into host cells[34]. Our studies indicate that total bile collected from the gallbladder of adult mice can enhance the entry of *Cryptosporidium* into cultured host epithelial cells. However, adequate supplementation with collected bile significantly reduces the infection burden both *in vivo* and *in vitro*. Additionally, treatment with bile collected from the gallbladder of adult mice or the FXR agonist FexD ameliorates dysregulated cell proliferation and migration in the intestine, including both the directly infected ileum region and the non-infected regions. Several previous studies have suggested a potential association between cryptosporidiosis and the presence of digestive neoplasia[44]. It remains to be determined whether there is an association between Bile acid-FXR pathway and the development of digestive neoplasia following *Cryptosporidium* infection.

Although obvious disturbances in cholesterol and bile homeostasis in the gut-liver axis were observed following *Cryptosporidium* infection, the underlying mechanisms driving these

metabolic disturbances remain unclear. In addition to reduced bile acid pool and subsequent lipid
malabsorption, the general inflammatory responses triggered by infection as previously
reported[45] may play a role. If so, it would be interesting to explore whether these findings are
applicable to infection caused by other gastrointestinal pathogens. In the intestinal epithelium,
these disturbances may also reflect the immature state of the epithelial cell population during
infection. Previous studies have shown that infection with this parasite accelerates ileal epithelial
development, resulting in an epithelium dominated by immature cells[46]. The outgrowth of
immature epithelial cells, which are actively progressing through the cell cycle and not yet fully
differentiated, may contribute to lipid malabsorption. Biliary infection by *Cryptosporidium*, as
reported in AIDS patients and other weakened immune systems[11], may have additional effects on
cholesterol and bile homeostasis. Additionally, bile acids have been shown to drive the
maturation of the newborn's gut microbiota[47]. Gut microbiome development is closely linked to
the overall development of the child, with changes in its composition and maturity potentially
affecting immune functions, intestinal development, neurodevelopment, nutritional energy
harvest, growth, and weight gain[48]. The composition and diversity of the gut microbiome during
early development play a crucial role in the incidence and severity of *Cryptosporidium*
infection[49–51]. Whether bile treatment influences the maturation of the gut microbiota and, in turn,
improves long-term deficits in growth, weight gain, and cognitive development following
Cryptosporidium infection warrants further exploration.
371
Methods

Ethics statement
374 All research studies involving the use of animals were reviewed and approved by the
375 Institutional Animal Care and Use Committee (IACUC) of Rush University at Chicago, Illinois,
376 and were carried out in strict accordance with the recommendations in the Guide for the Care and
377 Use of Laboratory Animals.
378
C. parvum and Cell lines
380 *C. parvum* oocysts of the Iowa strain were obtained from a commercial source (Bunch Grass
381 Farm, Deary, ID). HCT-8 cells were purchased from ATCC (Manassas, Virginia). The culture
382 media were supplied with 10% fetal bovine serum (FBS) (Ambion, Austin, Texas) and
383 antibiotics (100 IU/ml of penicillin and 100 µg/ml of streptomycin).
384
Histology of intestinal tissue
Neonatal mice were killed and small intestine was dissected. Given the stand 'swiss-roll' was not
feasible with the unfixed small intestines of neonatal mice due to their fragility. We collected
fresh tissue, gently rolled it into a spiral shape, and embedded it in OCT compound for
subsequent analysis. Slides were evaluated in a blinded fashion for measurements of parasite
burden, and Oil red staining (see Fig. 5f and Extended Data Fig. 7c). A section four centimeters
from the stomach of neonatal mice as the proximal small intestine (jejunum), and the distal small
intestine four centimeters from the cecum as the ileum for subsequent analysis (RT-qPCR,
Immunoblotting, Immunohistochemistry, and Oil red staining).
394
Bile collection
396 Gall bladders were removed from healthy adult C57BL/6 mice after 24 hours of fasting,
397 punctured with forceps to collect bile, and stored at -80°C until processing.

398
In vitro C. parvum infection experiments
399
Models of intestinal cryptosporidiosis using cultured cell lines were employed as previously
described[52]. For *C. parvum* infection burden experiments, infection was done in serum-free
culture medium for 4 h with a 1:4 ratio of *C. parvum* oocysts to HCT-8 cells. The culture
medium was changed to remove free parasites. The medium was supplemented with mouse bile
at concentrations of 0.2%, 0.25%, 0.33%, 0.5% and 1% (collected by gall bladder puncture from
healthy adult C57BL/6 mice) and cultured for an additional 20 h (a total 24 h p.i.), followed by
infection burden measurement. Infection burden was quantified by measuring levels of *C.*
parvum 18S gene (*cp18S*)[53]. Immunofluorescent staining of *C. parvum* was carried out using
rabbit antiserum against *C. parvum* membrane proteins as previously described[54,55]. For *C.*
parvum invasion experiments, HCT-8 cells were incubated with *C. parvum* oocysts (1:4 ratio of
oocysts to cells) in the presence of various concentrations of bile (0.07%, 0.2%, 1%, and 2%) or
PBS for 4 h, followed by RT-qPCR of *cp18S*.
412
In vivo C. parvum infection experiments
413
A well-established infection model of cryptosporidiosis in neonatal mice was used for *in vivo*
experiments[56,57]. Neonatal mice (5 days old) were orally inoculated with *C. parvum* (10^6 oocysts
per animal). Mice receiving PBS by oral gavage were used as control. At 72 and 96 h after *C.*
parvum oocysts or PBS administration, the animals were sacrificed, and intestine and liver
tissues were collected for biochemical analyses. For *C. parvum* infection burden experiments,
neonatal mice (5 days old) were orally inoculated with *C. parvum* (at a lower dose of 10^5 oocysts
per animal). At 24 h after *C. parvum* infection, the animals were fed with PBS-diluted bile (0.5,
1.5, 3, and 3,5 µl bile per animal, diluted in PBS to a total volume of 7 µl) and PBS (7µl per
animal), followed by daily for 4 days. The animals were sacrificed, and intestine tissues were
collected for biochemical analyses.
424
RT-qPCR
425
Total RNA was isolated using TRIzol™ Reagent (Thermo Fisher Scientific, catalogue no.
15596018), and genomic DNA was removed using the DNA-free™ DNA Removal Kit (Thermo
Fisher Scientific, catalogue no. AM1906). The reverse-transcription reaction was performed on
2 µg total RNA with M-MLV Reverse Transcriptase (Thermo Fisher Scientific, catalogue no.
28025013). Quantitative PCR assays were conducted using Bio-Rad CFX Manager v3.1 using
the iTaq™ Universal SYBR® Green Supermix (Bio-rad, Hercules, CA, USA, catalogue no.
1725124). RNA expression levels were calculated using the ΔΔCt method.
433
RNA-Seq and bioinformatic analysis
434
Total RNAs isolated with Rneasy Mini Kit (QIAGEN, catalogue no. 74104). Three biological
replicates (all mice from the same litter) were performed for each condition. The samples were
analyzed by BGI Americas Corporation (Cambridge, MA) using the DNBSEQ™ sequencing
technology platform, as previously reported[21]. Differential expression analysis and statistical
significance were assessed using DESeq in R with a standard negative binomial fit for the
aligned counts data and are described relative to the indicated control (Table S1 and S2). Gene
set enrichment analysis (GSEA) was performed using the R package "clusterProfiler"[58] with
gene sets from the MsigDB database (http://www.gsea-msigdb.org/gsea/index.jsp)[59]. Gene set
variation analysis (GSVA) was conducted with a significance threshold of $p < 0.05$ to explore

correlated pathways, using gene sets from the MsigDB database[60]. Bioinformatic analysis was
carried out and results were employed as we previously reported[21,61].
446
Immunoblotting
Membrane and cytosolic protein extracts were isolated from frozen mouse liver or small intestine
(30 mg) using the Mem-PER Plus Membrane Protein Extraction Kit (Thermo Fisher Scientific,
Catalog no. 89842). Membrane and cytosolic fractions were separated by SDS-PAGE and
transferred to a nitrocellulose membrane. Images were captured using the ChemiDoc MP
Imaging System (Bio-Rad, Catalog no. 12003154). The following primary and secondary
antibodies were used: anti-β Tubulin (Santa Cruz, catalogue no. sc-5274, 1:1000), anti-GAPDH
(Santa Cruz, catalogue no. sc-32233, 1:1000), anti-ABCG5 (Thermo Fisher Scientific, catalogue
no. BS-5013R, 1:1000), anti-ABCG8 (Novus Biologicals, catalogue no. 1B10A5, 1:1000), anti-
SLC10A2 (Abcam, catalogue no. ab203205, 1:1000), anti-FGF15 (Santa Cruz, catalogue no. sc-
514647, 1:1000), anti-NR1H4 (Proteintech, catalogue no. 25055-1-AP, 1:3000), anti-NR1H3
(Proteintech, catalogue no. 14351-1-AP, 1:5000), anti-NR1H2 (Proteintech, catalogue no.
60345-1-Ig, 1:1000), anti-Na+/K+-ATPase (Abclonal, catalogue no. A12405, 1:2000), anti-
HMGCR (Abclonal, catalogue no. A1633, 1:2000), anti-CYP7A1 (Abclonal, catalogue no.
A10615, 1:1000), anti-NR0B2 (Abclonal, catalogue no. A1836, 1:500), anti-HMGCS2
(Abclonal, catalogue no. A19232, 1:2000), anti-rabbit B700 (Bio-Rad, catalogue no. 12004161,
1:2500), anti-mouse B700 (Bio-Rad, catalogue no. 12004158, 1:2500), anti-rabbit B520 (Bio-
Rad, catalogue no. 12005869, 1:2500), anti-mouse B520 (Bio-Rad, catalogue no. 12005866,
1:2500), anti-GAPDH hFAB™ Rhodamine (Bio-Rad, catalogue no. 12004167, 1:2500), anti-
Tubulin hFAB™ Rhodamine (Bio-Rad, catalogue no. 12004165, 1:2500).
467
Immunohistochemistry
For tissue staining, frozen sections (8 μm) were fixed in 4% PFA in PBS for 15 min, then
washed twice with PBS. The sections were permeabilized by adding 0.5% Triton X-100 (Bio-rad,
catalogue no. 1610407) in PBS for 20 min, followed by removal of the permeabilization buffer
and two PBS washes. Sections were then blocked for 30 min in 10% normal donkey serum, 0.1%
Triton X-100, 0.01% sodium azide in PBS before incubation with the primary antibody for 2 h at
room temperature or overnight at 4 °C. Slides were washed three times with PBS and incubated
in block solution containing secondary antibody for 30 min at room temperature. After antibody
incubation, sections were washed three times with PBS (5 min per wash) and mounted with
Antifade Mounting Medium with DAPI (Vector, catalogue no. H-1200). Images were captured
using a Keyence BZ-X810 microscope. For cell staining, HCT-8 cells were fixed in 4% PFA in
PBS for 15 min and washed twice with PBS. Fixed cells were permeabilized with 0.1% Triton
X-100 in PBS for 5 min and washed three times with PBS. Cells were blocked for 30 min in 10%
normal donkey serum, 0.1% Triton X-100, 0.01% sodium azide in PBS, followed by incubation
with primary antibody for 1 h at room temperature. Cells were then washed three times with PBS
and incubated in block solution containing secondary antibody for 30 min at room temperature.
Following antibody incubation, cells were washed three times with PBS (5 min per wash) and
mounted with Antifade Mounting Medium with DAPI (Vector, catalogue no. H-1200). Images
were taken using a Keyence BZ-X810 microscope. Immunofluorescent staining of *C. parvum*
was carried out using rabbit antiserum against *C. parvum* membrane proteins as previously
described[54,55]. Other antibodies used for immunofluorescent staining include anti-Ki67 antibody
(Cell Signaling Technology, catalogue no. 9129, 1:200), anti-ABCA1 antibody (Thermo Fisher
Scientific, catalogue no. PA1-16789, 1:200), goat-anti-bovine Alexa fluor 488 (Jackson Immuno

Research, catalogue no. 101-545- 003, 1:250), goat-anti-bovine Alexa fluor 594 (Jackson
Immuno Research, catalogue no. 101-585- 003, 1:250), donkey-anti-rabbit Alexa fluor 488
(Jackson Immuno Research, catalogue no. 711-545-152, 1:500), donkey-anti-rabbit Rhodamine
Red (Jackson Immuno Research, catalogue no. 711-295-152, 1:500).
495
496   **Oil red staining**
497   Lipid vesicle storage in tissue was assessed with Oil Red O staining, following the
498   manufacturer's protocol (Abcam, catalogue no. ab150678). Images were captured using a
499   Keyence BZ-X810 microscope.
500
501   **Cell viability**
502   Cell viability was assessed using a Neutral Red Assay Kit, following the manufacturer's
503   instructions (Abcam, catalogue no. ab234039). The absorbance of each well at 540 nm was
504   measured using a microplate reader (SpectraMax iD3, Molecular Devices, USA) to evaluate the
505   cell viability.
506
507   **Bile Acid Targeted Metabolomics Assay**
508   Metabolites were extracted from the livers and small intestines of control and *C. parvum*-infected
509   neonatal mice, with three biological replicates for each condition. The samples were analyzed by
510   Metware Biotechnology Inc (Metware). Metabolomics profiling was employed as previously
511   described[62].
512
513   **Quantitative Lipidomics**
514   Metabolites were extracted from the livers and proximal intestinal epithelium of control and *C.*
515   *parvum* infected neonatal mice, with four biological replicates for each condition. The samples
516   were analyzed by Metware Biotechnology Inc (Metware). Metabolomics profiling was employed
517   as previously described[63].
518
Cholesterol assay
Cholesterol Quantification Assay Kit (Sigma-Aldrich, CS0005) was used to measure the
cholesterol concentration of samples. We sampled the distal small intestine four centimeters
from the cecum of neonatal mice and collected epithelial cells. Briefly, 46 μL Assay Buffer, 2 μL
Probe, 2 μL Enzyme Mix, and 50 μL sample were mixed and incubated at 37 °C for 30 min in
each well. A calibration curve was established for every measurement with standard samples
with 0–5 μg cholesterol. All samples were diluted to the range of the calibration curve with the
Assay Buffer. Absorbance at 570 nm was measured and compared to the standards on the same
plate to determine total cholesterol.
528
***In vivo* EdU assay**
Neonatal mice (5 days old) were orally inoculated with *C. parvum* (10^5 oocysts per animal).
Mice receiving PBS by oral gavage were used as control. For bile treatment, 24 h after *C.*
parvum oocysts or PBS administration, control mice were treated with PBS, *C. parvum*-infected
mice were fed PBS-diluted bile (7μl per animal,1:1 ratio of bile to PBS) or PBS only (7μl per
animal), followed by daily treatments for 3 days. Mice were i.p. injected with EdU and sacrificed
4 h post injection. In another group, mice were i.p. injected with EdU, treated with PBS-diluted
bile or PBS for an additional 2 days, and sacrificed 48 h post injection. For FXR agonists
fexaramine D (FexD) treatment, 24 h after *C. parvum* oocysts or PBS administration, control

mice were i.p. injected with vehicle, and *C. parvum*-infected mice were i.p. injected with vehicle
with or without 100 mg/kg of FexD, followed by daily treatments for 3 days. Mice were i.p.
injected with EdU and sacrificed 4 h post injection, or were i.p. injected with EdU, treated with
vehicle with or without FexD for an additional 2 days and sacrificed 48 h post injection. Intestine
tissues were frozen in OCT and cryosectioned (10 μm). EdU staining was performed using the
Click-iT EdU imaging kit (Thermofisher cat# C10337) according to the manufacturer's protocol.
To evaluate the proliferation and migration of EdU-labeled cells, we calculated the average
values across all villi/crypts within the same group as previously reported[64].
546
Bioinformatic analysis of metabolomics
Multivariate analysis of Variable Importance in Projection (VIP) from OPLS-DA modeling
(MetaboAnalystR, v.1.0.1) was used to preliminarily select differential metabolites from
different samples. For two-group analysis, differential metabolites were determined by VIP
(VIP > 1) and P-value (P-value < 0.05, Student's t test). The data were log-transformed ($\log_2$)
and mean-centered prior to OPLS-DA. To avoid overfitting, a permutation test (200
permutations) was performed. For KEGG annotation and enrichment analysis, identified lipids
were annotated using the KEGG Compound database (http://www.kegg.jp/kegg/compound/).
Annotated lipids were then mapped to the KEGG Pathway database
(http://www.kegg.jp/kegg/pathway.html). Pathways with significantly regulated lipids were
further analyzed using the MSEA (lipid sets enrichment analysis), with significance determined
by the hypergeometric test's p-values.
559
Statistical analysis
For all quantitative analyses, a minimum of three biological replicates were included. Statistical
tests were selected based on the assumption that sample data are from a population following a
probability distribution with a fixed set of parameters. Student's t-tests were used to assess the
statistical significance of differences between two groups. One-way AVOVA was used for
multiple comparison tests. The following values were considered statistically significant:
*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$. Calculations were performed using the
GraphPad Prism 10 software package. Data are presented as mean $\pm$ the standard error of the
mean.
569
Data availability
The RNA-seq data generated in this study have been deposited into NCBI's Gene Expression
Omnibus (GEO) and are accessible through GEO series accession number GSE234579 and
GSE272799. The metabolomics analysis data are presented in Table S3-S6. Uncropped images
of immunoblots presented in the figures are included in Supplementary Figure 1. Source data are
available with this paper.
576
Figure legends
Fig.1 | *C. parvum* infection disrupts cholesterol homeostasis in the small intestine.
a, Schematic diagram summarizing the molecular mechanism of cholesterol transport.
b, Changes in cholesterol homeostasis-related gene expression upon *C. parvum* infection (72 h
p.i.). The enrichment of cholesterol biosynthesis and efflux gene sets is shown at the bottom. The
upregulated and downregulated genes (log2FC > 0.5, adj. p < 0.05) are indicated by the light
red– and light blue–shaded rectangles underlaid on plot, respectively.

c, Schematic of the cholesterol biosynthesis pathway, where genes are represented in blue.

d, e, Relative amounts of mRNAs analyzed by RT–qPCR. Gene expression was presented as fold
change to uninfected control.

f, g, Immunoblotting analysis of ileum and proximal intestinal epithelial cells (IECs) in neonates
with and without *C. parvum* infection (72 h and 96 h p.i.). GAPDH and TUBULIN were used as
loading control of cytosolic protein.

h, Quantification of free cholesterol in ileum of control and infected mice (96 h p.i.).

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test
for (d, e). Data are expressed as mean ± SEM, and representative of at least two independent
experiments, n = 3–5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel
source data, see Supplementary Figure 1.

Fig.2 | *C. parvum* infection alters hepatic lipid metabolism.

a, Experiment schematics: Neonatal mice (5 days old) were orally inoculated with *C. parvum*
oocysts (10^6 oocysts per animal), and liver tissues were collected for deep sequencing (72 h p.i.)
and metabolomics (96 h p.i.).

b, Clustered heatmap of normalized gene counts of LXR and FXR target genes expression in the
liver of *C. parvum* infected neonatal mice versus control mice (72 h p.i.).

c, RT-qPCR analysis confirming hepatic mRNA expression of LXR and FXR target genes in
neonates with and without *C. parvum* infection at 72 h p.i.

d, Immunoblotting analysis of liver samples in control and *C. parvum* infected neonatal mice (72
h p.i.). GAPDH and TUBULIN were used as loading control of cytosolic protein.

e, Scatter plot showing the change of hepatic lipids between control or *C. parvum* infected mice
in different subclasses (96 h p.i.). Lipid with VIP ≥ 1 represents significant difference.

f, Heatmap of differential hepatic bile acids expression in control versus *C. parvum* infected mice
(96 h p.i.).

g, The total quantification of all bile acids in liver between control and *C. parvum* infected
neonatal mice from (f).

h, Structures of cholesterol and bile acids. Bile acids are synthesized from cholesterol and can be
amidated by taurine (or glycine).

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test
for (c, g). Data are expressed as mean ± SEM, and representative of at least two independent
experiments, n = 3–5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel
source data, see Supplementary Figure 1.

Fig.3 | *C. parvum* infection decreases bile acid levels in the small intestine and reduces the
absorption of dietary lipids.

a, Schematic overview of the enterohepatic circulation of bile acids and the FXR pathway.

b, Differential gene expression analysis upon *C. parvum* infected neonatal mice (72 h p.i.). The
enrichment of bile acid metabolic gene set is shown in blue and the distribution of Shp and
ASBT is shown in orange.

c, RT-qPCR analysis confirming ileum mRNA expression of FXR, Fgf15, Shp and ASBT from
infected neonates (72 h p.i.).

d, Immunoblotting analysis of ileum samples in control and *C. parvum* infected neonatal mice
(72 h and 96 h p.i.). GAPDH and TUBULIN were used as loading control of cytosolic protein,
Na+/K+ ATPase was used as loading control of membrane protein.

e, Oil Red O stains of the proximal small intestine and ileum section in neonates with and
without *C. parvum* infection (96 h p.i.).
f, The total quantification of all lipids in proximal IECs between control and *C. parvum* infected
neonatal mice.
g, Scatter plot showing the change of proximal IECs lipids between control or *C. parvum*
infected mice in different subclasses. Neonatal mice (5 days old) were orally inoculated with *C.*
parvum oocysts (10^6 oocysts per animal) and proximal IECs were collected (96 h p.i.) for
lipidomics analysis. Lipid with VIP $\geq$ 1 represents significant difference.
h, Heatmap of differential intestinal bile acids expression in control versus *C. parvum* infected
mice. Neonatal mice (5 days old) were orally inoculated with *C. parvum* oocysts (10^6 oocysts per
animal) and small intestine was collected (96 h p.i.) for bile acid targeted assay.
Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test
for (c, f). Data are expressed as mean ± SEM, and representative of at least two independent
experiments, n = 3–4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel
source data, see Supplementary Figure 1.

Fig.4 | Bile intervention promotes intestinal anti-*C. parvum* defense.
a, Experiment schematics: Neonatal mice (5 days old) were orally administered 1×10^5 *C. parvum*
oocysts; PBS and bile were given at 24 h p.i. followed by daily for 4 days, respectively;
Intestinal tissues were collected, and the mRNA levels of infection burden (b) and FXR target
genes (c) were measured by RT-qPCR.
d, Immunofluorescent staining of *C. parvum* from (a). A higher magnification of the boxed
region to visualize the infection burden of *C. parvum* from different portions of ileum is shown
(d). Blue: DAPI, red: Ki67, green: *C. parvum*. Bars, 100 μm.
e, Effects of bile on *C. parvum* infection of cultured HCT-8 cells. Cells were incubated with
oocysts for 4 h first, subsequently exposed to different concentrations of bile or PBS for another
20 h, followed by immunofluorescence staining of *C. parvum*. Blue: DAPI (nuclei), red: *C.*
parvum. Bars, 100 μm.
f, Effects of bile on *C. parvum* infection of cultured HCT-8 cells. Cells were incubated with
oocysts for 4 h first, subsequently exposed to different concentration of bile or PBS for another
20 h, followed by RT-qPCR of *cp18S* (*p<0.05 vs. PBS, ^ p<0.05 vs. 0.2% bile, #p<0.05 vs.
0.25% bile, + p<0.05 vs. 0.33% bile).
Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test
for (b, c), one-way analysis of variance (ANOVA) with Tukey tests was used for correction of
multiple comparisons in f. Data are expressed as mean ± SEM, and representative of at least two
independent experiments, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Fig.5 | Effects of different doses of bile treatment on *C. parvum* infection *in vivo*.
a, Experiment schematics: Neonatal mice (5 days old) were orally administered 2×10^6 *C. parvum*
oocysts; PBS and bile were given at 24 h p.i. followed by daily for 4 days, respectively;
Intestinal tissues were collected, and the mRNA levels of infection burden (b, d) and FXR target
genes (c, e) were measured by RT-qPCR.
f, Immunofluorescent and Oil Red O staining of the distal small intestine sections from (a). A
higher magnification of the boxed region (orange) to visualize the background staining; A higher
magnification of the boxed region (red) to visualize the *C. parvum* staining; Red arrow indicates
C. parvum staining. Blue: DAPI, red: Ki67, green: *C. parvum*. Statistical analysis: unpaired t test

for (b-e). Data are expressed as mean ± SEM, and representative of at least two independent
experiments, n = 4-5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.
678
Fig.6 | Bile acid treatment attenuates the dysregulated homeostasis in the intestinal
epithelium following *C. parvum* infection.
a, Experiment schematics: Neonatal mice (5 days old) were orally administered $1×10^5$ *C. parvum*
oocysts; PBS or bile was given at 24 h p.i. followed by daily for 3 days; Mice were i.p. injected
with EdU and sacrificed at 4 h post injection; Mice were i.p. injected with EdU, treated with PBS
or bile for another 2 days and sacrificed at 48 h post injection.
b, RT-qPCR analysis confirming mRNA expression of FXR, Shp, and Fgf15 in the small
intestine of uninfected neonates after bile treatment.
c-f, Quantification of EdU staining of proliferating cells in crypts at 4 h post injection (>50
crypts per mouse, 3 mice/group).
g, Representative images of Ki67, EdU staining of proliferating cells at 48 h post injection. Blue:
DAPI (nuclei), red: Ki67, green: EdU. Bars, 100 µm.
h-j, Quantification of cell migration length along the crypt-villus axis at 48 h post injection.
Quantification of cell migration length by measuring the distance from the crypt base to the
highest labelled cell in the villus (>30 villus per mouse, 3 mice/group).
Statistical analysis: unpaired t test for (b); One-way analysis of variance (ANOVA) with Tukey
tests was used for correction of multiple comparisons in c-f, h-j, and representative of at least
two independent experiments, n = 3 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P <
0.0001.
698
Fig.7 | FXR activation attenuates the dysregulated homeostasis in the intestinal epithelium
following *C. parvum* infection.
a, Experiment schematics: Neonatal mice (5 days old) were orally administered $1×10^5$ *C. parvum*
oocysts; FexD or vehicle were given at 24 h p.i., followed by daily for 3 days; Mice were i.p.
injected with EdU and sacrificed at 4 h post injection; Mice were i.p. injected with EdU, treated
with FexD or vehicle for another 2 days and sacrificed at 48 h post injection.
b, Representative images of Ki67, EdU staining of proliferating cells in crypts at 4 h post
injection. Blue: DAPI (nuclei), red: Ki67, green: EdU. Bars, 50 µm.
c-e, Quantification of cell migration length along the crypt-villus axis at 48 h post injection.
Quantification of cell migration length by measuring the distance from the crypt base to the
highest labelled cell in the villus (>20 villus per mouse, 3 mice/group).
f, Quantification of EdU staining of proliferating cells in crypts at 4 h (b) post injection (>50
crypts per mouse, 3 mice/group).
g, Representative images of Ki67, EdU staining of proliferating cells at 48 h post injection. Blue:
DAPI (nuclei), red: Ki67, green: EdU. Bars, 100 µm.
h, i, Neonatal mice (5 days old) were orally administered $1×10^5$ *C. parvum* oocysts; FexD or
vehicle was given at 24 h p.i., followed by daily for 3 days; Ileum tissues were collected, and
the mRNA levels of FXR target genes (h) and infection burden (i) were measured by RT-
qPCR.
Statistical analysis: unpaired t test for (h, i); One-way analysis of variance (ANOVA) with Tukey
tests was used for correction of multiple comparisons in c-f, and representative of at least two
independent experiments, n = 3 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.
721

Supplemental Figures Legends

Extended Data Fig.1 | *C. parvum* **intestinal infection burden in neonates with and without** *C. parvum* **infection.**

a, Representative images of duodenum, jejunum and ileum sections from *C. parvum* infected and control mice immunostained for *C. parvum* and Ki67 (96 h p.i.). Blue: DAPI (nuclei), red: Ki67, green: *C. parvum*. Bars, 100 μm.

b, Relative amounts of mRNAs analysed by RT–qPCR. Neonatal mice (5 days old) were inoculated with *C. parvum* oocysts (10^6 oocysts per animal) for 96 h. Infection burden was evaluated by RT-qPCR of *cp18S* gene in the duodenum, jejunum and ileum, respectively. The cycle threshold (Ct) values were used to determine the infection burden.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for b. Data is expressed as mean ± SEM, and representative of at least two independent experiments, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.2 | *C. parvum* **infection disrupts cholesterol homeostasis in the small intestine.**

a, Differential gene expression analysis upon *C. parvum* infection (72 h p.i.). The enrichment of sterol esterase activity gene set is shown at the bottom.

b-d, GSEA showing the enriched pathways of inflammatory reactions (b), defense response (c), sterol homeostasis and sterol transport (d) (|NES| > 1, p < 0.05, FDR < 0.25).

e, g, h, Relative amounts of mRNAs analyzed by RT–qPCR. Gene expression was presented as fold change to uninfected control.

f, Immunoblotting analysis of proximal and ileal IECs samples. GAPDH and TUBULIN were used as loading control of cytosolic protein.

i, Small intestine length of control and *C. parvum* infected mice (120 h p.i.).

j, Sections from the proximal/Jejunum and distal small intestine of neonatal mice 4 days after *C. parvum* infection were immunostaining for Ki67 and *C. parvum*. Blue: DAPI (nuclei), red: Ki67, green: *C. parvum*.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (e, g-i). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 3–5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel source data, see Supplementary Figure 1.

Extended Data Fig.3 | *C. parvum* **infection alters hepatic genes expression.**

a, Differential hepatic gene expression analysis upon *C. parvum* infection (72 h p.i.). The enrichment of positive regulation of immune response gene set is shown at the bottom.

b-f, Relative mRNA expression of lipid metabolic genes associated with the indicated processes were measured by RT-qPCR.

g, Immunoblotting analysis of liver samples in neonates with and without *C. parvum* infection (72 h p.i.). GAPDH and TUBULIN were used as loading control of cytosolic protein.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (b-f). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 4–5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel source data, see Supplementary Figure 1.

Extended Data Fig.4 | Impaired lipid homeostasis in the livers of *C. parvum* **infected mice.**

a, Hierarchical clustering showing the similarity between control and *C. parvum* infected neonatal mice (96 h p.i.).

b, Volcano plot illustrating the hepatic lipids expression between control and *C. parvum* infected neonatal mice (96 h p.i.). Lipid with VIP ≥ 1 represents significant difference.

c, KEGG pathway enrichment analysis and differential abundance (DA) scores of differential metabolites in the liver of control and *C. parvum* infected neonatal mice (96 h p.i.). DA score = (up regulated lipids in a pathway-down regulated lipids in a pathway) / (Total number of lipids annotation in a pathway).

d, Oil Red O stains of the liver sections in neonates with and without *C. parvum* infection (96 h p.i.).

e, The total quantification of all hepatic lipids between control and *C. parvum* infected neonatal mice (96 h p.i.).

f-k, Graphs depicting quantification of the differences in hepatic BA (f), CE (g), Cholesterol (h), FFA (i), DG (j) and TG (k) between control or *C. parvum* infected mice (96 h p.i.).

l-n, Graphs showing the fold change of hepatic FFA (l), DG (m) and TG (n) with different chain lengths between control or *C. parvum* infected mice (96 h p.i.). Each point represents a lipid, the size of the point represents the P-value.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (e-k). Data are expressed as mean ± SEM, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.5 | *C. parvum* infection alters intestinal lipid metabolic genes expression.

a-c, Relative amounts of mRNAs analyzed by RT–qPCR. Gene expression was presented as fold change to uninfected control.

d, Representative images of ileum (72 h and 96 h p.i.).

e, Immunoblot analysis of proximal and ileal IECs from control and *C. parvum* infected neonatal mice (72 h and 96 h p.i.). GAPDH was used as loading control of cytosolic protein.

f-h, GSEA showing the enriched pathways of fatty acid metabolic process (f), intestinal lipid absorption process (g) and vitamin metabolic process (h) (|NES| > 1, p < 0.05, FDR < 0.25).

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (a-c). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 3–4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel source data, see Supplementary Figure 1.

Extended Data Fig.6 | Impaired lipid homeostasis in the proximal IECs of *C. parvum* infected mice.

a, Graphs depicting quantification of the differences in proximal IECs lipid abundance of each subclass between control or *C. parvum* infected mice (96 h p.i.).

b, KEGG pathway enrichment analysis and DA scores of differential metabolites in proximal IECs of control and *C. parvum* infected neonatal mice (96 h p.i.).

c, Graphs showing the quantification of L-palmitoylcarnitine in proximal IECs between control or *C. parvum* infected mice (96 h p.i.).

d, Violin plot showing the result of top 18 differentially bile acids with the largest log2FC value. The box in the middle represents the interquartile range, and the middle box represents the 95% confidence interval. The black horizontal line is the median, and the outer shape represents the distribution density of the data.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (a,c). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 3–4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.7 | Effects of bile on *C. parvum* infection.
a, Effects of bile on *C. parvum* intestinal infection in neonates. Neonatal mice (5 days old) were orally administered *C. parvum* oocysts; PBS or bile were given at 24 h p.i. followed by daily for 4 days, respectively; Intestinal tissues were collected, and the mRNA levels of cholesterol biosynthesis genes (a) and cholesterol efflux genes (b) were measured by RT-qPCR. Gene expression was presented as fold change to uninfected control.
c, Immunofluorescent and Oil Red O staining of the distal small intestine sections from (a). A higher magnification of the boxed region (red) to visualize the *C. parvum* staining; Red arrow indicates *C. parvum* staining. Blue: DAPI, red: Ki67, green: *C. parvum*.
d, Effects of bile on *C. parvum* invasion of cultured HCT-8 cells. Cells were incubated with oocysts for 4 h in the presence of different concentrations of bile or PBS, followed by RT-qPCR of *cp18S*.
e, Effects of bile on *C. parvum* infection of 2D monolayers. Cells were incubated with oocysts for 4 h first, subsequently exposed to different concentration of bile or PBS for another 20 h, followed by RT-qPCR of *cp18S*.
Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (a, b, d, e). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.8 | Effects of bile on HCT-8 cells viability with or without *C. parvum* infection.
a, Effects of bile on HCT-8 cells viability in the presence of *C. parvum* infection. Cells were incubated with oocysts for 4 h first, subsequently exposed to different concentrations of bile or PBS for another 20 h, followed by cell viability assay.
b, Cell survival of HCT-8 cells treated with different concentration of bile or PBS for 24 h.
(c, d) Cell survival of HCT-8 cells exposed to different concentration of bile or PBS with (c) or without (d) oocysts for 4 h. One-way analysis of variance (ANOVA) with Dunnett's multiple comparison test was used for correction of multiple comparisons in a-d, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.9 | Bile intervention attenuates the dysregulated cell proliferation in crypts following *C. parvum* infection.
a, Neonatal mice (5 days old) were orally administered 1×10^5 *C. parvum* oocysts; PBS or bile was given at 24 h p.i. followed by daily for 3 days; Mice were i.p. injected with EdU and sacrificed at 4 h post injection. Representative images of Ki67, EdU staining of proliferating cells in crypts at 4 h post injection. Blue: DAPI (nuclei), red: Ki67, green: EdU. Bars, 50 μm.

References

1. Kotloff, K. L. *et al.* Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet Lond. Engl.* **382**, 209–222 (2013).

2. Checkley, W. *et al.* A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for cryptosporidium. *Lancet Infect. Dis.* **15**, 85–94 (2015).

3. Sow, S. O. *et al.* The Burden of Cryptosporidium Diarrheal Disease among Children < 24 Months of Age in Moderate/High Mortality Regions of Sub-Saharan Africa and South Asia, Utilizing Data from the Global Enteric Multicenter Study (GEMS). *PLoS Negl. Trop. Dis.* **10**, e0004729 (2016).

4. Khalil, I. A. *et al.* Morbidity, mortality, and long-term consequences associated with diarrhoea from Cryptosporidium infection in children younger than 5 years: a meta-analyses study. *Lancet Glob. Health* **6**, e758–e768 (2018).

5. Costa, L. B. *et al.* Cryptosporidium-malnutrition interactions: mucosal disruption, cytokines, and TLR signaling in a weaned murine model. *J. Parasitol.* **97**, 1113–1120 (2011).

6. Kirkpatrick, B. D. *et al.* Cryptosporidiosis stimulates an inflammatory intestinal response in malnourished Haitian children. *J. Infect. Dis.* **186**, 94–101 (2002).

7. Mondal, D., Haque, R., Sack, R. B., Kirkpatrick, B. D. & Petri, W. A. Attribution of malnutrition to cause-specific diarrheal illness: evidence from a prospective study of preschool children in Mirpur, Dhaka, Bangladesh. *Am. J. Trop. Med. Hyg.* **80**, 824–826 (2009).

8. Tzipori, S. & Ward, H. Cryptosporidiosis: biology, pathogenesis and disease. *Microbes Infect.* **4**, 1047–1058 (2002).

9. Watson, A. J. M., Duckworth, C. A., Guan, Y. & Montrose, M. H. Mechanisms of epithelial cell shedding in the Mammalian intestine and maintenance of barrier function. *Ann. N. Y. Acad. Sci.* **1165**, 135–142 (2009).

10. Delgado, M. E., Grabinger, T. & Brunner, T. Cell death at the intestinal epithelial front line. *FEBS J.* **283**, 2701–2719 (2016).

11. Chen, X.-M., Keithly, J. S., Paya, C. V. & LaRusso, N. F. Cryptosporidiosis. *N. Engl. J. Med.* **346**, 1723–1731 (2002).

12. Pardy, R. D., Wallbank, B. A., Striepen, B. & Hunter, C. A. Immunity to Cryptosporidium: insights into principles of enteric responses to infection. *Nat. Rev. Immunol.* **24**, 142–155 (2024).

13. Wahlström, A., Sayin, S. I., Marschall, H.-U. & Bäckhed, F. Intestinal Crosstalk between Bile Acids and Microbiota and Its Impact on Host Metabolism. *Cell Metab.* **24**, 41–50 (2016).

14. Chiang, J. Y. L. Bile acids: regulation of synthesis. *J. Lipid Res.* **50**, 1955–1966 (2009).

15. Makishima, M. *et al.* Identification of a nuclear receptor for bile acids. *Science* **284**, 1362–1365 (1999).

16. Parks, D. J. *et al.* Bile acids: natural ligands for an orphan nuclear receptor. *Science* **284**, 1365–1368 (1999).

17. Forman, B. M. *et al.* Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* **81**, 687–693 (1995).

18. Matsubara, T., Li, F. & Gonzalez, F. J. FXR signaling in the enterohepatic system. *Mol. Cell. Endocrinol.* **368**, 17–29 (2013).

19. Marzook, N. B. *et al.* The essential host genome for Cryptosporidium survival exposes metabolic dependencies that can be leveraged for treatment. *Cell* S0092-8674(25)00751–2 (2025) doi:10.1016/j.cell.2025.07.001.

20. Ehrenman, K., Wanyiri, J. W., Bhat, N., Ward, H. D. & Coppens, I. Cryptosporidium parvum scavenges LDL-derived cholesterol and micellar cholesterol internalized into enterocytes. *Cell. Microbiol.* **15**, 1182–1197 (2013).

21. Deng, S. *et al.* Cryptosporidium uses CSpV1 to activate host type I interferon and attenuate antiparasitic defenses. *Nat. Commun.* **14**, 1456 (2023).

22. Luo, J., Yang, H. & Song, B.-L. Mechanisms and regulation of cholesterol homeostasis. *Nat. Rev. Mol. Cell Biol.* **21**, 225–245 (2020).

23. Wang, B. & Tontonoz, P. Liver X receptors in lipid signalling and membrane homeostasis. *Nat. Rev. Endocrinol.* **14**, 452–463 (2018).

24. Xu, D. *et al.* Cholesterol sulfate alleviates ulcerative colitis by promoting cholesterol biosynthesis in colonic epithelial cells. *Nat. Commun.* **13**, 4428 (2022).

25.   Tso, P. & Balint, J. A. Formation and transport of chylomicrons by enterocytes to the lymphatics. *Am. J.*
Physiol. **250**, G715-726 (1986).

26.   Ko, C.-W., Qu, J., Black, D. D. & Tso, P. Regulation of intestinal lipid metabolism: current concepts and
relevance to disease. *Nat. Rev. Gastroenterol. Hepatol.* **17**, 169–183 (2020).

27.   Li, H., Yu, X.-H., Ou, X., Ouyang, X.-P. & Tang, C.-K. Hepatic cholesterol transport and its role in non-
alcoholic fatty liver disease and atherosclerosis. *Prog. Lipid Res.* **83**, 101109 (2021).

28.   Jensen, R. G. Lipids in human milk. *Lipids* **34**, 1243–1271 (1999).

29.   Liu, Z., Rochfort, S. & Cocks, B. Milk lipidomics: What we know and what we don't. *Prog. Lipid Res.* **71**, 70–
85 (2018).

30.   Chen, F. *et al.* Liver receptor homologue-1 mediates species- and cell line-specific bile acid-dependent
negative feedback regulation of the apical sodium-dependent bile acid transporter. *J. Biol. Chem.* **278**, 19909–
19916 (2003).

31.   Moschandrea, C. *et al.* Mitochondrial dysfunction abrogates dietary lipid processing in enterocytes.
Nature **625**, 385–392 (2024).

32.   Spector, A. A. & Kim, H.-Y. Discovery of essential fatty acids. *J. Lipid Res.* **56**, 11–21 (2015).

33.   Chen, X. M. & LaRusso, N. F. Mechanisms of attachment and internalization of Cryptosporidium parvum to
biliary and intestinal epithelial cells. *Gastroenterology* **118**, 368–379 (2000).

34.   Feng, H., Nie, W., Sheoran, A., Zhang, Q. & Tzipori, S. Bile acids enhance invasiveness of Cryptosporidium
spp. into cultured cells. *Infect. Immun.* **74**, 3342–3346 (2006).

35.   Fang, S. *et al.* Intestinal FXR agonism promotes adipose tissue browning and reduces obesity and insulin
resistance. *Nat. Med.* **21**, 159–165 (2015).

36.   Fu, T. *et al.* FXR Regulates Intestinal Cancer Stem Cell Proliferation. *Cell* **176**, 1098-1112.e18 (2019).

37.   Maran, R. R. M. *et al.* Farnesoid X receptor deficiency in mice leads to increased intestinal epithelial cell
proliferation and tumor development. *J. Pharmacol. Exp. Ther.* **328**, 469–477 (2009).

38.   Modica, S., Murzilli, S., Salvatore, L., Schmidt, D. R. & Moschetta, A. Nuclear bile acid receptor FXR
protects against intestinal tumorigenesis. *Cancer Res.* **68**, 9589–9594 (2008).

39.   Ming, Z., Zhou, R. & Chen, X.-M. Regulation of host epithelial responses to Cryptosporidium infection by
microRNAs. *Parasite Immunol.* **39**, (2017).

40.   Thomson, S. *et al.* Bovine cryptosporidiosis: impact, host-parasite interaction and control strategies. *Vet.*
Res. **48**, 42 (2017).

41.   Annaba, F. *et al.* Modulation of ileal bile acid transporter (ASBT) activity by depletion of plasma
membrane cholesterol: association with lipid rafts. *Am. J. Physiol. Gastrointest. Liver Physiol.* **294**, G489-497 (2008).

42.   Cohen, D. E. Balancing cholesterol synthesis and absorption in the gastrointestinal tract. *J. Clin. Lipidol.* **2**,
S1-3 (2008).

43.   Funkhouser-Jones, L. J. *et al.* Microbiota-produced indole metabolites disrupt mitochondrial function and
inhibit Cryptosporidium parvum growth. *Cell Rep.* **42**, 112680 (2023).

44.   Kalantari, N. *et al.* Association between Cryptosporidium infection and cancer: A systematic review and
meta-analysis. *Parasitol. Int.* **74**, 101979 (2020).

45.   Russler-Germain, E. V. *et al.* Commensal Cryptosporidium colonization elicits a cDC1-dependent Th1
response that promotes intestinal homeostasis and limits other infections. *Immunity* **54**, 2547-2564.e7 (2021).

46.   Wallbank, B. A., Pardy, R. D., Brodsky, I. E., Hunter, C. A. & Striepen, B. Cryptosporidium impacts epithelial
turnover and is resistant to induced death of the host cell. *mBio* **15**, e0172024 (2024).

47.   van Best, N. *et al.* Bile acids drive the newborn's gut microbiota maturation. *Nat. Commun.* **11**, 3692
(2020).

48.   Fontaine, F., Turjeman, S., Callens, K. & Koren, O. The intersection of undernutrition, microbiome, and
child development in the first years of life. *Nat. Commun.* **14**, 3554 (2023).

49.   VanDussen, K. L. *et al.* Neonatal Mouse Gut Metabolites Influence Cryptosporidium parvum Infection in
Intestinal Epithelial Cells. *mBio* **11**, e02582-20 (2020).

50.   Charania, R., Wade, B. E., McNair, N. N. & Mead, J. R. Changes in the Microbiome of Cryptosporidium-
Infected Mice Correlate to Differences in Susceptibility and Infection Levels. *Microorganisms* **8**, 879 (2020).

51.   Carey, M. A. *et al.* Megasphaera in the Stool Microbiota Is Negatively Associated With Diarrheal
Cryptosporidiosis. *Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.* **73**, e1242–e1251 (2021).

52.   Wang, Y. *et al.* Delivery of parasite Cdg7_Flc_0990 RNA transcript into intestinal epithelial cells during
Cryptosporidium parvum infection suppresses host cell gene transcription through epigenetic mechanisms. *Cell.*
Microbiol. **19**, (2017).
53.   Ferguson, S. H. *et al.* Interferon-λ3 Promotes Epithelial Defense and Barrier Function Against
Cryptosporidium parvum Infection. *Cell. Mol. Gastroenterol. Hepatol.* **8**, 1–20 (2019).
54.   Zhu, G. *et al.* Expression and functional characterization of a giant Type I fatty acid synthase (CpFAS1)
gene from Cryptosporidium parvum. *Mol. Biochem. Parasitol.* **134**, 127–135 (2004).
55.   Chen, X.-M. *et al.* Cryptosporidium parvum invasion of biliary epithelia requires host cell tyrosine
phosphorylation of cortactin via c-Src. *Gastroenterology* **125**, 216–228 (2003).
56.   Lacroix, S., Mancassola, R., Naciri, M. & Laurent, F. Cryptosporidium parvum-specific mucosal immune
response in C57BL/6 neonatal and gamma interferon-deficient mice: role of tumor necrosis factor alpha in
protection. *Infect. Immun.* **69**, 1635–1642 (2001).
57.   Kapel, N. *et al.* Kinetics of mucosal ileal gamma-interferon response during cryptosporidiosis in
immunocompetent neonatal mice. *Parasitol. Res.* **82**, 664–667 (1996).
58.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R package for comparing biological themes
among gene clusters. *Omics J. Integr. Biol.* **16**, 284–287 (2012).
59.   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst.* **1**,
417–425 (2015).
60.   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for microarray and RNA-seq
data. *BMC Bioinformatics* **14**, 7 (2013).
61.   Klomp, J. A. *et al.* Defining the KRAS- and ERK-dependent transcriptome in KRAS-mutant cancers. *Science*
384, eadk0775 (2024).
62.   Yin, Q. *et al.* Enhancing milk quality and modulating rectal microbiota of dairy goats in starch-rich diet: the
role of bile acid supplementation. *J. Anim. Sci. Biotechnol.* **15**, 7 (2024).
63.   Wang, J. *et al.* Identification and evaluation of a lipid-lowering small compound in preclinical models and
in a Phase I trial. *Cell Metab.* **34**, 667-680.e6 (2022).
64.   Wang, B. *et al.* Phospholipid Remodeling and Cholesterol Availability Regulate Intestinal Stemness and
Tumorigenesis. *Cell Stem Cell* **22**, 206-220.e4 (2018).
990

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Extended Data Fig.1 | *C. parvum* intestinal infection burden in neonates with and without *C. parvum* infection.
a, Representative images of duodenum, jejunum and ileum sections from *C. parvum* infected and control mice immunostained for *C. parvum* and Ki67 (96 h p.i.). Blue: DAPI (nuclei), red: Ki67, green: *C. parvum*. Bars, 100 μm.
b, Relative amounts of mRNAs analysed by RT–qPCR. Neonatal mice (5 days old) were inoculated with *C. parvum* oocysts (10^6 oocysts per animal) for 96 h. Infection burden was evaluated by RT-qPCR of *cp18S* gene in the duodenum, jejunum and ileum, respectively. The cycle threshold (Ct) values were used to determine the infection burden.
Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for b. Data is expressed as mean ± SEM, and representative of at least two independent experiments, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.2 | *C. parvum* infection disrupts cholesterol homeostasis in the small intestine.

a, Differential gene expression analysis upon *C. parvum* infection (72 h p.i.). The enrichment of sterol esterase activity gene set is shown at the bottom.

b-d, GSEA showing the enriched pathways of inflammatory reactions (b), defense response (c), sterol homeostasis and sterol transport (d) (|NES| > 1, p < 0.05, FDR < 0.25).

e, g, h, Relative amounts of mRNAs analyzed by RT–qPCR. Gene expression was presented as fold change to uninfected control.

f, Immunoblotting analysis of proximal and ileal IECs samples. GAPDH and TUBULIN were used as loading control of cytosolic protein.

i, Small intestine length of control and *C. parvum* infected mice (120 h p.i.).

j, Sections from the proximal/Jejunum and distal small intestine of neonatal mice 4 days after *C. parvum* infection were immunostaining for Ki67 and *C. parvum*. Blue: DAPI (nuclei), red: Ki67, green: *C. parvum*.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (e, g-i). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 3–5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel source data, see Supplementary Figure 1

Extended Data Fig.3 | *C. parvum* infection alters hepatic genes expression.
a, Differential hepatic gene expression analysis upon *C. parvum* infection (72 h p.i.). The enrichment of positive regulation of immune response gene set is shown at the bottom.
b-f, Relative mRNA expression of lipid metabolic genes associated with the indicated processes were measured by RT-qPCR.
g, Immunoblotting analysis of liver samples in neonates with and without *C. parvum* infection (72 h p.i.). GAPDH and TUBULIN were used as loading control of cytosolic protein.
Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (b-f). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 4–5 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel source data, see Supplementary Figure 1.

Extended Data Fig.4 | Impaired lipid homeostasis in the livers of *C. parvum* infected mice.

a, Hierarchical clustering showing the similarity between control and *C. parvum* infected neonatal mice (96 h p.i.).

b, Volcano plot illustrating the hepatic lipids expression between control and *C. parvum* infected neonatal mice (96 h p.i.). Lipid with VIP ≥ 1 represents significant difference.

c, KEGG pathway enrichment analysis and differential abundance (DA) scores of differential metabolites in the liver of control and *C. parvum* infected neonatal mice (96 h p.i.). DA score = (up regulated lipids in a pathway-down regulated lipids in a pathway) / (Total number of lipids annotation in a pathway).

d, Oil Red O stains of the liver sections in neonates with and without *C. parvum* infection (96 h p.i.).

e, The total quantification of all hepatic lipids between control and *C. parvum* infected neonatal mice (96 h p.i.).

f-k, Graphs depicting quantification of the differences in hepatic BA (f), CE (g), Cholesterol (h), FFA (i), DG (j) and TG (k) between control or *C. parvum* infected mice (96 h p.i.).

l-n, Graphs showing the fold change of hepatic FFA (l), DG (m) and TG (n) with different chain lengths between control or *C. parvum* infected mice (96 h p.i.). Each point represents a lipid, the size of the point represents the P-value.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (e-k). Data are expressed as mean ± SEM, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.5 | *C. parvum* infection alters intestinal lipid metabolic genes expression.

a-c, Relative amounts of mRNAs analyzed by RT–qPCR. Gene expression was presented as fold change to uninfected control.

d, Representative images of ileum (72 h and 96 h p.i.).

e, Immunoblot analysis of proximal and ileal IECs from control and *C. parvum* infected neonatal mice (72 h and 96 h p.i.). GAPDH was used as loading control of cytosolic protein.

f-h, GSEA showing the enriched pathways of fatty acid metabolic process (f), intestinal lipid absorption process (g) and vitamin metabolic process (h) (|NES| > 1, p < 0.05, FDR < 0.25).

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (a-c). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 3–4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. For gel source data, see Supplementary Figure 1.

Extended Data Fig.6 | Impaired lipid homeostasis in the proximal IECs of *C. parvum* infected mice.
a, Graphs depicting quantification of the differences in proximal IECs lipid abundance of each subclass between control or *C. parvum* infected mice (96 h p.i.).
b, KEGG pathway enrichment analysis and DA scores of differential metabolites in proximal IECs of control and *C. parvum* infected neonatal mice (96 h p.i.).
c, Graphs showing the quantification of L-palmitoylcarnitine in proximal IECs between control or *C. parvum* infected mice (96 h p.i.).
d, Violin plot showing the result of top 18 differentially bile acids with the largest log2FC value. The box in the middle represents the interquartile range, and the middle box represents the 95% confidence interval. The black horizontal line is the median, and the outer shape represents the distribution density of the data.
Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (a,c). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 3–4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.7 | Effects of bile on *C. parvum* infection.

a, Effects of bile on *C. parvum* intestinal infection in neonates. Neonatal mice (5 days old) were orally administered *C. parvum* oocysts; PBS or bile were given at 24 h p.i. followed by daily for 4 days, respectively; Intestinal tissues were collected, and the mRNA levels of cholesterol biosynthesis genes (a) and cholesterol efflux genes (b) were measured by RT-qPCR. Gene expression was presented as fold change to uninfected control.

c, Immunofluorescent and Oil Red O staining of the distal small intestine sections from (a). A higher magnification of the boxed region (red) to visualize the *C. parvum* staining; Red arrow indicates *C. parvum* staining. Blue: DAPI, red: Ki67, green: *C. parvum*.

d, Effects of bile on *C. parvum* invasion of cultured HCT-8 cells. Cells were incubated with oocysts for 4 h in the presence of different concentrations of bile or PBS, followed by RT-qPCR of *cp18S*.

e, Effects of bile on *C. parvum* infection of 2D monolayers. Cells were incubated with oocysts for 4 h first, subsequently exposed to different concentration of bile or PBS for another 20 h, followed by RT-qPCR of *cp18S*.

Each symbol represents one mouse, all from the same litter. Statistical analysis: unpaired t test for (a, b, d, e). Data are expressed as mean ± SEM, and representative of at least two independent experiments, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.8 | Effects of bile on HCT-8 cells viability with or without *C. parvum* infection.
a, Effects of bile on HCT-8 cells viability in the presence of *C. parvum* infection. Cells were incubated with oocysts for 4 h first, subsequently exposed to different concentrations of bile or PBS for another 20 h, followed by cell viability assay.
b, Cell survival of HCT-8 cells treated with different concentration of bile or PBS for 24 h.
(c, d) Cell survival of HCT-8 cells exposed to different concentration of bile or PBS with (c) or without (d) oocysts for 4 h. One-way analysis of variance (ANOVA) with Dunnett's multiple comparison test was used for correction of multiple comparisons in a-d, n = 4 per group. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Fig.9 | Bile intervention attenuates the dysregulated cell proliferation in crypts following *C. parvum* infection.
a, Neonatal mice (5 days old) were orally administered 1×10^5 *C. parvum* oocysts; PBS or bile was given at 24 h p.i. followed by daily for 3 days; Mice were i.p. injected with EdU and sacrificed at 4 h post injection. Representative images of Ki67, EdU staining of proliferating cells in crypts at 4 h post injection. Blue: DAPI (nuclei), red: Ki67, green: EdU. Bars, 50 μm.