Qing-Luo-Yin eases angiogenesis in adjuvant-induced arthritis rats by activating

PPARγ

Meng-Ke Song [a,1], Qin Yin [b,c,1], Meng-Fan Gu [a,b], Wen-Gang Chen [c], Opeyemi Joshua

Olatunji [d], Yan Li [a,e,*] Jian Zuo [a,e,f,*]

[a] Xin'an Medicine Research Center, the First Affiliated Hospital of Wannan Medical

College (Yijishan Hospital), Wuhu, 241000, China;

[b] Research Center of Integration of Traditional Chinese and Western Medicine,

Wannan Medical College, Wuhu, 241000, China;

[c] Department of Pharmacy, the Second Affiliated Hospital of Wannan Medical

College, Wuhu, 241000, China;

[d] African Genome Center, Mohammed VI Polytechnic University, Ben Guerir, 43150,

Morocco;

[e] Center for Xin'an Medicine and Modernization of Traditional Chinese Medicine,

Institution of Health and Medicine, Hefei Comprehensive National Science Center,

Hefei 230000, China;

[F] Anhui Province Key Laboratory of Non-coding RNA Basic and Clinical

Transformation, Wannan Medical College, Wuhu, 241000, China.

[1] These authors contributed equally to this work.

Correspondence to:

[*] Prof. Yan Li

No 2, West Zheshan Road, Wuhu 241000, Anhui, China; liyan.0301@163.com

Phone +86-13093626158; fax +86-553-5739999

[*] Dr. Jian Zuo

No 2, West Zheshan Road, Wuhu 241000, Anhui, China; zuojian8178@163.com

Phone +86-15056477665; fax +86-553-5739999

ABSTRACT

Objective: Qing-Luo-Yin (QLY) is an anti-rheumatic herbal formula with potentials

activating PPARγ. This study investigated if its anti-angiogenesis effects are related to

immune modulation.

Method: Adjuvant-induced arthritis (AIA) rats were orally treated by QLY or

rosiglitazone (a PPARγ agonist) for 30 days. Their immune and metabolism statues

were investigated afterward. Isolated monocytes and lymphocytes were co-cultured

reciprocally, and treated by different serums. Healthy rats received blood transfusion

from QLY-treated or AIA model rats. Two days ahead of sacrifice, a matrigel plug

was planted. The plug and some blood immune indicators were examined. AIA rat

serum-incubated THP-1 and Jurkat cells were treated by sinomenine, berberine and

palmatine. The medium and T0070907 (a PPARγ inhibitor) were used to stimulate

HUVEC cells.

Results: QLY showed similar therapeutic effects on AIA to rosiglitazone, alleviating

joint injuries, synovial angiogenesis, and metabolic disorders. Although QLY

impaired inflammatory phenotype of AIA monocytes in vivo, the effect was hardly

achieved or sustained in vitro. T cells from QLY-treated AIA rats showed the

weakened inflammatory phenotype, and were unable to induce monocytes

inflammatory polarization. AIA rat lymphocytes induced angiogenesis in the matrigel

plug in healthy recipients. In lymphocytes enrichment site, QLY reduced the secretion

of IL-17A, IFNγ, and many angiogenesis-related cytokines. QLY-related components

affected Jurkat but not THP-1 cells. Jurkat T cells induced angiogenesis of HUVEC

cells when cultured by AIA rat serum. Inhibitory effects of the compounds on it were

abolished by T0070907.

Conclusion: PPARγ activation in T cells is a foundation for the anti-angiogenesis

property of QLY.

Keywords:

Rheumatoid arthritis; Traditional Chinese Medicine; immune, monocyte; vascular

endothelial cell; angiogenesis

Abbreviations:

WAT, white adipose tissues; AIA, adjuvant-induced arthritis; QLY, Qing-Luo-Yin;

RA, rheumatoid arthritis; TCM, Traditional Chinese Medicine; IFA, Incomplete

Freud's adjuvant; BCG, Bacillus Calmette-Guérin; RSG, rosiglitazone; RF,

rheumatoid factor; HIF-1, hypoxia inducible factor 1; ARG-1, arginase 1; IL-1β,

interleukin 1 beta; TGF-β1, transforming growth factor beta1; IFNγ, interferon

gamma; VEGF, vasoactive endothelial growth factor; ICAM-1, intercellular cell

adhesion molecule 1; CXCL1, C-X-C motif ligand 1; MCP-1, monocyte chemotactic

protein 1; IGF-1, insulin-like growth factor 1; TG, triglyceride; NEFA,

nonestesterified fatty acid; GSH, reduced glutathione; MDA, malonaldehyde; iNOS,

inducible nitric oxide synthase; CAT, catalase; NOX, NADH oxidase; AKP, alkaline

phosphatase; AST, aspartate aminotransferase; FBS, fetal bovine serum; PCR,

polymerase chain reaction; WB, western-blot; FCM, flow cytometry; DMARD,

disease modifying antirheumatic drug.

1. Introduction

The successful treatments of rheumatoid arthritis (RA) are still challenging due to its elusive etiological mechanisms. Many pathological changes contribute to its onset and progression. According to the current understanding, synovitis is the key driving force underlying RA-related structure degradation and functional loss of joints [1-2]. In addition to synovial invasion, angiogenesis fuels this abnormality directly [3]. Because articular manifestations are still the priority of anti-rheumatic therapies now, synoviocytes and vascular endothelial cells should be the main therapeutic targets. Despite many studies have validated plausibility of this strategy [4-5], there is no such targeting drug available now. A fundamental reason is that RA pathogenesis is driven by immune hyper-activation rather than specific local changes [2]. Various outcomes from effective RA therapies can be explained by the improved immune milieu, and immune-regulating approach would bring more benefits than the other strategies.

Thanks to the accumulating knowledge about the immune characteristics of RA, some biological agents have been successfully developed and introduced into clinical practices [6]. Compared to the conventional disease modifying anti-rheumatic drugs (DMARDs), they exhibit advantages in treating various RA-related pathological changes [6]. They mainly target immune cells, the dominant roles in RA pathology, as well as the signals controlling their differentiation. But these drugs are unaffordable in developing countries. Hence, there is still a great demand for cheap and effective therapies.

In addition to DMARDs, herbal medicines are widely used worldwide. They might be even more effective than the synthesized compounds, benefited from their functions on alleviating symptom combinations by numerous chemical combinations and therapeutic targets, which have been explored by novel technologies such as network pharmacology [7-9]. As a successful representative, the Traditional Chinese Medicine (TCM) formula Qing-Luo-Yin (QLY) has been used for over 40 years. Its efficacy has been thoroughly validated by both clinical and pharmacological

researches [10-13]. Of note, it exhibited eyes catching effects on synovial

angiogenesis [14].

Great endeavor is devoted to clarify the anti-rheumatic mechanisms of QLY, and

we confirm that SIRT1/PPARγ is one of its molecular targets [12-13]. Interestingly,

SIRT1 and PPARγ function as a pair of rivals, from both the immune and metabolism

perspectives. Furthermore, they negatively regulate each other's expression [15-16].

QLY shows potentials in inhibiting SIRT1 and activating PPARγ [13]. Since SIRT1 is

generally recognized as a negative regulator of inflammation, its effect on PPARγ

would bring more benefits.

PPARγ functional deficiency is common in rheumatic subjects, and implicated in

many RA-related pathological changes [16]. This abnormality doesn't only occur in

joints, and in fact PPARγ down-regulation is a key factor driving the imbalanced

differentiation of immune cells [16-17]. These factors make PPARγ a promising

anti-rheumatic target. Many compounds with PPARγ activating potentials have been

confirmed with good therapeutic effects on RA, and rosiglitazone (RSG) is the most

well-known representative [18]. Thanks to the pleiotropic roles of PPARγ, the related

treatments can cause series of outcomes in rheumatic subjects, including angiogenesis

inhibition [18-19]. Considering the crucial impacts of immune cells on RA pathology,

it is apparent that the anti-angiogenesis effects of these agents are immune dependent.

Inspired by the above clues, it is reasonable to assume that QLY anti-rheumatic

treatments may ease immune disruption-related angiogenesis by activating PPARγ. In

this study, we further ascertained the involvement of PPARγ activation during QLY

treatment on adjuvant-induced arthritis (AIA) rats by comparing its effects to RSG, a

PPARγ agonist. Importantly, we identified T cells as its preferential targeting immune

cells, and revealed that via this approach, potential PPARγ agonists in QLY improved

the overall immune environment, and consequently inhibited synovial angiogenesis.

2. Materials and Methods

2.1.Chemicals and reagents

Incomplete Freund's Adjuvant (IFA) and Bacillus Calmette-Guérin (BCG) were purchased from Chondex (Redmond, WA) and Rebio Scientific (Shanghai, China), respectively. Rheumatoid factor (RF), hypoxia inducible factor-1(HIF-1), and arginase-1 (ARG-1) ELISA kits were supplied by Jianglai Biotechnology (Shanghai, China). Interleukin 1 beta (IL-1β), IL-6, IL-10, IL-17A, transforming growth factor beta1 (TGF-β1), interferon gamma (IFNγ), vasoactive endothelial growth factor (VEGF), intercellular cell adhesion molecule 1(ICAM-1), C-X-C motif ligand 1 (CXCL1), monocyte chemotactic protein 1 (MCP-1), and insulin-like growth factor 1 (IGF-1) ELISA kits were supplied by Multi-Science (Hangzhou, Zhejiang, China). Glucose, lactic acid, pyruvic acid, triglyceride (TG), nonestesterified fatty acid (NEFA), reduced glutathione (GSH), malonaldehyde (MDA), inducible nitric oxide synthase (iNOS), catalase (CAT), NADH oxidase (NOX), alkaline phosphatase (AKP), and aspartate aminotransferase (AST) quantification kits were bought from Jiancheng Bioengineering Institute (Nanjing, Jiangsu, China). ReverAid first-strand cDNA synthesis kits and universal qPCR master mix were purchased from Thermo Fisher Scientific (Rockford, IL, USA) and New England Biolabs (Ipswich, MA, USA), respectively. Gene-specific primers used were synthesized by General Biology (Chuzhou, Anhui, China). Primary β-ACTIN, GAPDH, TLR4, STAT3, P-STAT3, JNK, p-JNK, CPT1A, CD36, PPARγ, and ATGL antibodies were obtained from ABclonal Technology (Wuhan, Hubei, China) or Affinity Biosciences (Changzhou, Jiangsu, China). HRP-conjugated IgG antibodies were purchased from Boster Biological Technology (Wuhan, Hubei, China). Red donkey anti-rabbit and orange donkey anti-mouse secondary antibodies were bought from Abbkine (Wuhan, Hubei, China). PE-tagged CD172a, APC-tagged CD43, PE-tagged CD16, FITC-tagged CD3, PE-tagged CD4 antibodies were procured from BioLegend (SanDiego, CA, USA) or Invitrogen (Carlsbad, CA, USA). RPMI-1640 medium, polyformaldehyde, crystal violet staining solution were procured from Biosharp (Beijing, China). Fetal bovine serum (FBS) was the product of Wisent (Montreal, Quebec, Canada). Pure

compounds sinomenine, berberine, and palmatine were purchased from Bencao

Yikang (Nanjing, Jiangsu, China). T0070907 (a PPARγ selective inhibitor) was

purchased from Aladdin (Shanghai, China). Rat peripheral blood lymphocyte

isolation kit and rat/human peripheral blood monocyte isolation kit were procured

from Solarbio (Beijing, China). Fluorescent dye CFDA-SE and matrigel were bought

from from MedChemExpress (Monmouth Junction, NJ, USA) and Thermo Fisher

Scientific (Rockford, IL, USA), respectively.

2.2.AIA induction and treatments

The *in vivo* experiments were based on Wistar rats, which were supplied by Tianqin

Biotechnology (Changsha, Hunan, China). They were kept in a qualified SPF lab with

the standard conditions, and fed by the commercial rodent chow and boiled tap water.

All invasive operations were conducted under anesthesia by the help of pentobarbital

sodium (30 mg/kg). The greatest efforts were devoted to minimize the experimental

animals' suffering.

AIA induction was conducted based on a previously detailed protocol using male

rats (8 weeks-old) [11-13]. IFA and BCG were mixed to prepare complete Freund's

adjuvant (CFA), which was used to provoke immune reactions in all the rats except

those assigned into normal group. Since CFA injection, the immunized rats were

divided randomly into 3 groups, and received various treatments for 30 days. Each

group contained 5 rats, and the grouping is as below: healthy controls (normal, treated

by 0.5% CMC-Na), AIA controls (AIA, treated by 0.5% CMC-Na), QLY-treated AIA

rats (QLY, treated by QLY extract dispersed in 0.5% CMC-Na, equivalent to 10 g/kg

of crude drugs), and RSG-treated AIA rats (RSG, treated by RSG resolved in 0.5%

CMC-Na, 2 mg/kg). All the treatments were administrated daily by gavage. During

the treatment period, arthritis scores and body weights were periodically recorded.

The qualification criterion of arthritis scores was based on a previous report [20].

2.3.Therapeutic outcomes evaluation

One hour after the last administration on day 30, all the rats were anesthetized. The maximum amounts of blood were collected through abdominal aorta until the animals were euthanized. Blood plasma and serum were prepared by centrifugation. Levels of representative cytokines, metabolites and diagnostic indicators of RA in blood were detected by commercial quantification kits. Rat monocytes and lymphocytes were prepared using anti-coagulation blood by the aid of isolation kits. Blood serum and immune cells were used in the *in vitro* experiments. Main organs were dissected and weighted. Joint injuries were assessed by histological examinations with the reported protocols [21-24]. The livers were homogenized. Levels of metabolites and cytokines in the homogenates were detected by the corresponding kits. Expression of certain immune and metabolism-related proteins in the livers and spleens was studied by western-blot (WB) method with routine procedures [23-25].

2.4.Assessment of QLY-caused impacts on monocytes and lymphocytes

Distribution of monocyte subsets was investigated by flow cytometry (FCM) method [11, 26]. The whole blood collected above were spiked into 6-fold volumes of red cells lysis buffer, incubated, and centrifuged. The precipitates were re-suspended, and incubated with fluorescein-tagged antibodies (APC-CD43 and PE-CD172a) in dark. The stained cells were further washed by PBS, filtered (200 mesh), and fed to a BD FACSVerse instrument. Some of the isolated monocytes were cultured in complete RPMI-1640 medium for 24 h. Levels of certain cytokines in the medium were determined by ELISA kits. The remaining medium was used to incubate healthy rat lymphocytes, whose status was evaluated by polymerase chain reaction (PCR) with routine procedures [27-28]. The primed used were synthesized by Universal Biology (Chuzhou, Anhui, China), and their sequences are shown in **Supplementary S1**. Afterwards, we cultured the lymphocytes isolated from QLY-treated or AIA model rats for 24 h, and detected representative cytokines in the medium. The remaining medium was used to culture healthy rat monocytes, and their secretion capacity was evaluated by ELISA tests.

Next, we treated normal monocytes with different blood serums. To prepare

QLY-containing serum, 5 healthy rats were treated by QLY for 3 consecutive days at

the dose mentioned above. Five untreated rats from the same batch were killed at the

same day to harvest blood serum. After normal rat monocytes were cultured in these

serums for 24 h, levels of IL-1β and IL-6 in the medium were determined by ELISA

kits. The rinsed cells were treated by Triton X-100, bovine serum albumin, primary

antibodies (IL-1β and PPARγ), and fluorescein-tagged secondary antibodies in turns.

Finally, cell nucleus was stained by DAPI [29]. In a replicate experiment, RA patients'

monocytes were used instead. In addition to ELISA analysis, $CD16^+$ cell subset was

analyzed by FCM method in this experiment.

2.5.Blood transfusion-based angiogenesis experiments

Six pairs of sex-matured male and female rats were kept separately for breeding. The

pups were raised for 8 weeks. By then, 10 male offspring were induced with AIA, and

half of them received QLY treatment. When inflammation became obvious in AIA

rats (day 14), 1 ml of peripheral blood was sampled from each donator through fossa

orbitalis vein every 3 days, and injected into caudal vein of their female littermates.

To investigate distribution of the implanted immune cells, whole blood from an

AIA rat was used to prepare peripheral blood mononuclear cells based on a gradient

centrifugation method, which were then dyed with CFDA-SE. Since cell injection, the

rat was observed by an Optima Max-XP instrument at pre-determined intervals. The

excitation and emission wavelengths were set at 500 and 520 nm, respectively.

On the day of the 8[th] blood injection, 0.4 ml of matrigel supplemented with

VEGF (10 ng/ml) was injected subcutaneously on abdomen [30]. The matrigel was

thawed under 4° overnight beforehand. Three days later, the recipients were sacrificed.

The formed plugs were retrieved, photographed, and subjected to H&E histological

examination. Blood was collected through abdominal aorta for FCM analysis. Hinder

paws were homogenized. Levels of cytokines and biochemical indicators in rats' knee

homogenates and remaining blood samples were determined by corresponding kits.

2.6.Evaluating immune-mediated anti-angiogenesis effects of compounds

To precipitate proteins, 100 μl of QLY-containing serum was spiked into 400 μl of

methanol [31-32]. After a centrifugation (12000 rpm for 10 min), the supernatant was

injected into a 1260 Agilent Infinity II LC system for quantification analysis. The

separation was achieved on a Waters C18 column (4.6 × 250 mm, 8 μm). A gradient

elution program was adopted, and sodium dihydrogen phosphate aqueous solution (10

mM, pH = 8.0) and acetonitrile served as mobile phases A and B respectively: 0-5

min, 20% A; 5-10 min, 50% A; 10-15 min, 20% A. The flow rate, injection volume

and detection wavelength were set at 1 ml/min, 20 μl and 270 nm, respectively. The

quantification curves for sinomenine, berberine and palmatine were developed by

analyzing a serial of diluted solutions, which contained all the reference compounds.

THP-1 and Jurkat cells were cultured and passaged under ordinary conditions

[33]. They were seeded in 6-wells plates. After 12 h adaptive growth, the supernatant

was replaced by fresh RPMI-1640 medium supplemented with 10% normal healthy or

AIA rat serums. Some of the AIA serum-stimulated cells were treated by a mixture

(sinomenine, berberine and palmatine) at various concentrations. Twenty-four hours

later, some cytokines in medium were detected, and the mediums were collected.

HUVEC cells were seeded in a 6-wells plate at an appropriate density. After the

adaptive culture, they were incubated in the medium from immune cells culture. In

this experiment, only one optimized concentration of the compounds mixture was

adopted when treating immune cells, and one of the treatment wells was added by

T0070907 (18 μM). HUVEC cells monolayer was scratched by a pipette tip. Twelve

hours later, the wound was observed again, and VEGF levels in the medium were

detected [30]. Next, we seeded HUVEC cells in a matrigel-coated 6-wells plate. After

attachment, they were treated by the same medium and chemical as above for 24 h.

Capillary tubules were observed and photographed [30]. In a replicate experiment,

HUVEC cells were seeded in matrigel-coated transwells (8 μm). The lower chambers

were filled with the medium mentioned above. After 24 h, the cells infiltrated through

pores were stained with crystal violet and photographed.

2.7.Statistical analysis

All the data were presented as mean ± standard deviation. Statistical differences

among groups were analyzed based on T-test or one-way analysis of variance coupled

with Tukey post hoc test using GraphPad Prism 8.0 (GraphPad Software, Cary, NC,

USA).

3. Results

3.1.QLY and RSG alleviated AIA-related angiogenesis in rats

AIA-related articular manifestations reach peak about day 14. Since then, therapeutic

effects of the two treatments became obvious, causing profound decrease in arthritis

scores. But they didn't prevent AIA-caused decrease in body weight gain (**Figure 1A**).

By the end of the *in vivo* experiment, paw swelling was still evident in AIA rats, and

this situation was greatly eased in all the treated rats (**Figure 1B**). Among the main

organs, liver and spleen showed the most significant weight changes. Hepatomegaly

and splenomegaly were obvious in AIA rats, which were greatly reduced by QLY and

RSG treatments (**Figure 1C**). This serves as a sign of the improved immune status,

which was solidly confirmed by ELISA analysis.

Compared to normal controls, all the tested RA-related indicators in blood were

elevated in AIA rats. RF levels were increased by approximately 4-fold, indicating the

successful duplication of RA models. The increase in levels of CXCL1, IL-1β, IL-6

and MCP-1 shows inflammatory activation of immune system. Interestingly, levels of

IL-10 and TGF-β1, two anti-inflammatory cytokines, were also up-regulated. VEGF

increase in AIA rats demonstrates an environment favorable for angiogenesis. All the

abnormal changes were improved by the treatments. Although QLY performed better

than RSG in controlling CXCL1, IL-1β and IL-6 levels, they were similarly effective

in reducing VEGF (**Figure 1D**). Benefited from that, the two treatments both greatly

reduced angiogenesis in the joints of AIA rats (H&E staining sections). As the result,

joint degradation was prevented (SOFG staining sections) (**Figure 1E**). These facts

above suggest that the therapeutic actions of QLY were similar to RSG, and they can

both improve immune environment and ease abnormal angiogenesis in AIA rats.

3.2.QLY and RSG relieved AIA-related metabolic abnormalities by activating PPARγ

Because PPARγ controls metabolism, we observed metabolic changes in the treated

rats to ascertain the effects of the treatments on this signaling. We observed lactic acid

and pyruvic acid increase as well as glucose decrease in AIA rats, demonstrating the

accelerated glycolysis, a sign of inflammation [30]. Meanwhile, TG was increased a

bit. QLY and RSG generally narrowed these AIA-caused metabolic changes (**Figure**

2A). The altered metabolism conditions would impact oxidative balance *in vivo*. The

increase of MDA/CAT ratio exhibits escalated oxidative stress, which was restored by

QLY and RSG more or less. Confusingly, GSH levels were increased in AIA rats too,

and barely affected by the two treatments (**Figure 2B**). IGF-1 is a novel RA-related

indictor, controlling glucose and mineral metabolism [34]. It levels were decreased

under AIA conditions. But neither of the treatments improved this situation (**Figure**

2C).

Because metabolism is largely controlled by the liver, we next focused changes

there. ALP and AST levels were increased a lot, which serve the solid evidence for

hepatic injury (**Figure 2D**). It was attributed to AIA-related systemic inflammation,

and levels of IL-1β and IL-6 were increased significantly in AIA rats' liver (**Figure**

2E). After QLY and RSG treatments, inflammatory liver injuries were thoroughly

cured (**Figure 2D-E**). These changes must affect metabolism functions of the liver.

Compared to the normal controls, TG and NEFA levels were decreased a lot in the

liver of AIA rats. Although the treatments didn't affect TG, they effectively restored

NEFA levels (**Figure 2F**). To clarify the reason accounting for the metabolic changes

there, we performed WB analysis (**Figure 2G**). All the uncropped raw band images

were included in **Supplementary S2**. We observed that expression of ATGL, CD36,

and CPT1A was unchanged, up-regulated and down-regulated in AIA rats' liver,

respectively (**Figure 2H**). It can be inferred that liver took in more NEFA, but utilized

less under AIA conditions. Meanwhile, fat mobilization remained stable. Hence, the

reduced TG and NEFA must be caused by the impaired synthesis, which reflects

substantial deficiency of PPARγ functions. Indeed, compared to the normal controls,

PPARγ expression in AIA rats' liver was reduced. Similar to RSG, QLY restored its

levels, and promoted CPT1A expression, a downstream target of PPARγ (**Figure 2H**).

This experiment also provided some useful clues to explain the reason for AIA-related

hepatic inflammation. TLR4 as well as its downstream JNK were barely affected by

AIA conditions, while STAT3 phosphorylation was greatly promoted (**Figure 2G**).

Hence, Th17 cells but not monocytes/macrophages may initiate inflammation there

[35]. RSG and QLY greatly reduced p-STAT3 expression, but showed weak effects

on TLR4/JNK pathway. It suggests that PPARγ regulation may exert more impacts on

T cells than monocytes/macrophages.

3.3.T cells were preferential targets of QLY treatment in AIA rats

In addition to T cells, monocytes/macrophages also play a crucial role in RA-related

inflammatory manifestations, and are important anti-rheumatic targets [36-37]. We

performed FCM analysis to investigate the impacts of QLY and RSG treatments on

monocytes (**Figure 3A**). It was revealed that non-classic $CD43^+CD172a^+$ monocyte

subset was increased by 3-fold in AIA rats, but classic $CD43^+CD172a^-$ monocytes

were reduced. Both the treatments restored these abnormal changes (**Figure 3B**). Also,

they showed potentials in down-regulating NOX and iNOS levels in the blood of AIA

rats, two indicators of inflammatory monocytes/macrophages (**Figure 3C**) [37]. To

confirm whether immune phenotype of these cells was fundamentally changed, we

isolated them from blood and cultured *in vitro*. AIA rat monocytes were distinctly

different from the normal controls. They released more IL-1β, IL-6, TGF-β1 and

IL-10. However, no improvement concerning this was observed in the monocytes

isolated from QLY and RSG-treated AIA rats (**Figure 3D**). This fact questioned if

these PPARγ activating treatments can substantially affect monocytes/macrophages.

Next, we co-cultured normal rat lymphocytes with the monocytes from either

QLY-treated or model AIA rats, and performed PCR analysis using the lymphocytes.

This experiment confirmed that QLY didn't fundamentally change immune nature of

AIA monocytes, because the monocytes from QLY-treated AIA rats showed similar

impacts on expression of different Th cell subsets-related genes (*IL-6*, *IFNγ*, *IL-10*,

IL-17A, *FOXP3*, and *ROR-γT*) to those from AIA models (**Figure 4A**). In accordance

to this conclusion, we found QLY-containing rat serum showed no difference with the

normal serum on IL-1β and IL-10 secretion of AIA monocytes *in vitro* (**Figure 4B**).

Immunofluorescence assay obtained a similar result. QLY-containing serum neither

reduced IL-1β secretion nor promoted PPARγ expression (**Figure 4C**). We noticed

that the expression of IL-1β and PPARγ was not always relevant. Some cells could

highly express IL-1β and PPARγ at the same time. It further confirms that monocytes

are not preferential targets of PPARγ agonists. To confirm this, we repeated the above

cell culture by using RA patients' monocytes. Once again, QLY-containing rat serum

showed no effect on the immune indicators IL-1β, IL-10, IL-6 and ARG-1 (**Figure**

4D). FCM analysis shows than it even increased the distribution of CD16$^+$ subset in

the cultured human monocytes, the traditionally defined inflammatory cells (**Figure**

4E-F) [38]. Regardless of this mystery, we can conclude that QLY treatment exert no

anti-inflammatory effects on monocytes directly at treatment doses.

Hence, the observed changes of monocytes in treated AIA rats must be mediated

by T cells, the dominate role in RA pathology [2]. As anticipated, the lymphocytes

isolated from QLY-treated AIA rats released much less IL-17A, IL-6 and IFN-γ than

the model controls, indicating impaired Th1/17 differentiation. IL-10 levels remained

unchanged, showing unaffected Treg differentiation (**Figure 4G**). Furthermore, it was

revealed that compared to the normal monocytes co-cultured by AIA rat lymphocytes,

those co-cultured by the lymphocytes from QLY-treated AIA rats produced less IL-1β,

IL-10, and IL-6. The opposite outcome occurred on ARG-1 levels (**Figure 4H**). These

evidences confirm that QLY inhibited inflammatory polarization of monocytes by

inhibiting Th1/17 differentiation. Spleen is the immune organ where lymphocytes

reside and mature. We performed a WB assay using these samples (**Figure 4I**), and

observed that PPARγ expression in AIA rats' spleen was reduced dramatically, which

was restored by QLY and RSG to certain extents (**Figure 4J**). It validates the claim

that QLY reshapes rheumatic subjects' T cells phenotypes by activating PPARγ, and

improves overall immune milieu by affecting other immune cells differentiation.

3.4.T cells-mediated anti-angiogenesis effects of QLY in AIA rats

To test the central role of T cells in QLY anti-rheumatic treatment, we transplanted

the peripheral immune cells from QLY-treated or AIA model rats in healthy recipients

and compared the consequences. During the intermittent whole blood transfusion, the

impacts from short-lived granulocytes would be negligible. Because of much larger

population size, lymphocytes will bring more obvious impacts than monocytes on the

recipients. The immune cells from AIA rats' blood were first widely distributed in the

body through circulation channels, and viscera, spine and joints were their enrichment

sites. Gradually, the cells in chest were eliminated. The remaining cells were enriched

in knee joints and spine (**Figure 5A**). It exhibits that T cells were the main long-lived

immune cells from AIA rats' blood functional in the recipients, became they are

autoimmune reactive to cartilage [2]. Similar to the outcomes above, QLY-treated

AIA rats' blood transfusion caused a decline in $CD43^+CD172a^+$ cell counts in the

recipients, albeit the change wasn't significant (**Figure 5B-C**). We analyzed a panel

of cytokines to better demonstrate immune cells transplant-caused consequences.

Disappointingly, blood transfusion didn't cause notable systematic changes. In fact,

the counts of $CD3^+CD4^+$ T cells in the recipients of the two groups were also similar

(**Supplementary S3**). Among all the investigated indicators, only IL-1β and ARG-1

were found changed in the recipients receiving QLY-treated AIA rats' blood (**Figure

5D**). The decrease in IL-1β and increase in ARG-1 exhibit the rebalanced monocytes

polarization [37]. But iNOS levels reduction in the recipients' blood wasn't obvious

(**Figure 5E**). There was no difference about MDA, GSH and CAT levels between the

two groups, further showing their similar immune and metabolism conditions (**Figure 5F**). These results basically show that the ability of QLY-treated AIA rats' lymphocytes in inducing inflammatory polarization of monocytes was impaired a bit.

Because the donators' immune cells were enriched in joints, we believed that changes there would be more obvious. Hence, we repeated the above analyses using knee joint homogenates, which basically verified the hypothesis. Level differences of IL-1β and ARG-1 in joints between the groups were similar to those found in blood. TGF-β1 decrease didn't only indicate altered immune conditions, but also promised the eased angiogenesis in the recipients receiving QLY-treated AIA rats' blood [3-4]. In line with this, some other angiogenesis-related cytokines including VEGF, HIF-1α and ICAM-1 were also reduced. IL-17A and IFNγ decrease in these animals confirm that Th1/17 differentiation was hampered by QLY therapy (**Figure 6A**). Due to T cells enrichment in joints, functional difference of monocytes/macrophages between the two groups became obvious there. QLY group showed much lower levels of iNOS expression than AIA group (**Figure 6B**). Despite of these immune changes, oxidative stress was unaffected (**Figure 6C**). The above ELISA tests confirm that angiogenesis is a T cells-mediated event. To vividly demonstrate the effects brought by the immune cells transplant, we planted a matrigel plug in the healthy recipients. Obviously, AIA rats' immune cells caused angiogenesis, indicated by dark red color of the plug, and their potentials were greatly weakened after QLY therapy (**Figure 6D**). Examination on H&E stained sections revealed dense new vessels in the plug planted in healthy recipients transfused with AIA models' blood. A large amount of immune cells were infiltrated through these channels. These situations were improved in QLY group (**Figure 6E**). The above results show that it is plausible that QLY reshaped T cells, and then exerted anti-angiogenesis effects in AIA rats.

3.5.QLY-derived PPARγ agonists inhibited angiogenesis by targeting T cells

Main bioactive components in QLY are alkaloid derivatives, which contribute greatly to its anti-rheumatic effects [7-10]. In QLY-containing serum, we detected 3 alkaloids,

sinomenine, berberine and palmatine (**Figure 7A**). Our unpublished study revealed

that all of them are potential PPARγ agonists. Using the quantification curves (**Figure**

7B), we calculated their concentrations in the blood. The mean levels of sinomenine,

berberine and palmatine were 2.08, 0.51 and 1.12 μg/ml, respectively (**Figure 7C**).

The values were adopted as the medium treatment dose of the compounds mixture in

following experiments. The low and high doses were divided and multiplied by 2,

respectively. The mixture failed to reduce IL-1β and TGF-β1 secretion in AIA rat

serum-primed THP-1 monocytes at all the three doses (**Figure 7D**). Differently, AIA

serum-induced IFNγ and IL-17A production in Jurkat T cells was reduced a lot by

them, generally in a dose-dependent manner (**Figure 7E**). Based on these results, we

then treated rat lymphocytes at the high dose, and added a PPARγ antagonist in some

treatment wells. The resulting medium was used to culture HUVEC cells. The culture

medium from AIA rat serum-primed lymphocytes greatly augmented VEGF secretion

in normal HUVEC cells. QLY-related compounds impaired this capacity of these rat

lymphocytes, while T0070907 completely abrogated this effect (**Figure 7F**). VEGF

level differences eventually affected angiogenesis potentials of HUVEC cells. Twelve

hours after growth in the lymphocytes culture mediums, wound healing phenomenon

was obvious in AIA group, but the compounds treatment on lymphocytes suppressed

HUVEC cells migration (**Figure 7G**). Subsequently, we confirmed that the potentials

of HUVEC cells infiltrating through transwell and forming tubes were reinforced

when cultured by the medium from AIA rat serum-primed lymphocytes, which were

inhibited by the compounds too (**Figure 7H-J**). T0070907 showed the antagonistic

effects to the compounds mixture in all the above experiments. The facts confirm that

T cells-mediated anti-angiogenesis effects of QLY rely on PPARγ activation.

4. Discussion

Because of the distinctly different theories and therapeutic notions from contemporary

medicine system, TCM is generally unfamiliar to the western world. Some of TCM

viewpoints seem to be groundless, and lack of supporting evidence. We have to admit

that being an empirical medicine, TCM always emphasize clinical efficacy rather than

theory evolution. To overcome its instinct shortage, more efforts should be devoted to

narrow this gap with evolving research strategy and new knowledge [39].

In RA field, humankind has entered a new era. It is clear now that immune cells

are the dominant players in RA pathology, although some other cells also participate

in development of RA-related manifestations. Accordingly, immune cells become the

preferential targets of RA therapies. It does not only lay a foundation for developing

new biological anti-rheumatic agents, but also successfully explains the therapeutic

mechanisms of some conventional DMARDs. Effective therapies always achieve

comprehensive improvements, but their direct targets were largely unknown. It will

impose negative impacts on rational and precise use of the relevant drugs. By taking

methotrexate as an example, it can ease many aspects of RA, including inflammation,

angiogenesis, joints degradation, and even metabolic complications [40-41]. We can't

attribute all of the outcomes to its medicinal properties, and it is inappropriate to use

this drug to treat metabolic diseases. Actually, adenosine receptors activation-caused

immune changes serve as the foundation for the other effects. It is highly possible that

certain natural drugs also achieve comprehensive anti-rheumatic effects by regulating

immune status. Many researches support this theory [42-43].

QLY is a TCM formula showing multiple functions in treating RA [44]. The

relevant researches started from investigating its effects on synovial angiogenesis [14].

But in fact, it can also treat systemic inflammation, joint injuries, and metabolic

alterations. Inspired by the aforementioned clues, we believe that immune-regulating

property is the key for QLY to achieve the overall anti-rheumatic outcomes. An

important clue is that all the herbal components in it (Radix Sophorae flavescentis,

Caulis Sinomenium acutum, Cortex Phellodendri chinensis, and Rhizoma Dioscoreae

hypoglaucae) are discovered with the relevant functions [45-48].

It is a common sense that T cells drive the development and progression of RA.

Meanwhile, monocytes/macrophages are the main source of inflammatory mediators

and deeply implicated in RA-related inflammation. Therefore, the two representatives

of adaptive and innate immune components are usually emphasized in RA researches.

Our works confirmed that QLY can inhibit them from acquiring the inflammatory

phenotypes [11-12]. As immune environment is shaped based on sophisticated

interplay of different cells, these findings brought another puzzle: which kind of

immune cells is the main targets of QLY. The factors mediating development and

differentiation of T cells and monocytes/macrophages are largely different. It is not

very possible that QLY is equally effective in regulating them.

One of our previous studies showed that QLY therapy hampered inflammatory

differentiation of T cells and monocytes/macrophages simultaneously in rheumatic

rodents [11]. Importantly, this work found that T cells were much more sensitive to

QLY treatments than monocytes/macrophages at treatment doses, demonstrating the

central role of T cells in QLY-induced immune changes for the first time. QLY didn't

only affect Th1/17 differentiation, but also impaired generation and maturation of T

cells. The current study confirms this conclusion and provides more convincing

evidence. Although QLY globally decreased inflammatory cytokines in AIA rats

(**Figure 1D**), it selectively targeted T cells rather than innate immunocytes in the liver,

indicated by the varied effects on STAT3 and TLR4/JNK signals (**Figure 2G**). At the

same time, QLY-induced monocytes/macrophages changes cannot be sustained *in*

vitro (**Figure 3D**). It hints that these outcomes relied on internal environment changes.

In another word, impacts of QLY on them were indirect and mediated by some other

factors. Indeed, the monocytes from AIA rats and RA patients were both unaffected

by QLY-containing serum (**Figure 4**). QLY-related compounds cannot impact them

too at therapeutic concentrations (**Figure 7D**). Both the *in vitro* (**Figure 4**) and *in vivo*

(**Figure 5**) evidences show that T cells responded well to QLY therapy, and mediated

the global immune changes. In fact as early as 2018, we had noticed that QLY would

preferentially target inflammatory T cells, as it inhibited pentose phosphate pathway

[10]. Since T cells initiate RA-related autoimmune reactions and most of pathological

changes, this property explains many anti-rheumatic functions of QLY, including its

effects on synovial angiogenisis. Meanwhile, this previous work revealed PPARγ as a

key molecular target of QLY.

PPARγ is a well-known governor of lipid metabolism. Therefore, QLY-induced

PPARγ activation must affect adipose tissues and liver. Previously, we showed that

QLY can attenuate monocytes/macrophages-mediated inflammation in AIA rats by

activating PPARγ in adipose tissues [12]. Here, we further confirm its activation

effects on PPARγ by examining the changes in liver. As expected, QLY didn't only

restore the altered lipids levels there, but also directly promoted expression of PPARγ

and its downstream CPT1A (**Figure 2G**). A recent study suggests that PPARγ

activation during QLY treatments would be resulted from the inhibition on SIRT1

[10]. But it is difficult to explain QLY-induced PPARγ up-regulation in some cases

when SIRT1 signal was normal [12]. Possibly, there exist PPARγ agonists in QLY.

Our unpublished work confirmed this, and revealed that many related alkaloids in

QLY are PPARγ agonist candidates. As their representatives, sinomenine, berberine

and palmatine can potently inhibit inflammatory secretion of T cells, and

compromised their ability in inducing angiogenesis (**Figure 7**). This finding suggests

that PPARγ activation in T cells is especially beneficial in the context of RA therapies,

despite of the fact that monocytes/macrophages are usually defined as

PPARγ-relevant immune cells, since the transcriptional factors driving their

inflammatory polarization like AP-1 and NF-κB are negatively regulated by PPARγ

[16-17]. Although many studies confirm that PPARγ activation would inhibit

inflammatory T cells differentiation, the detailed mechanism is still not thoroughly

elucidated [49-50]. Hence, more works should be done in this aspect, which is crucial

for exploiting the potentials of PPARγ-targeting anti-rheumatic therapies.

5. Conclusion

Similar to RSG, QLY also effectively reduced AIA severity in rats, reducing systemic

inflammation, joint injuries as well as synovial angiogenesis by activating PPARγ. T

cells were more sensitive to than monocytes QLY treatment. Through cell interactions, QLY improved overall immune environment by inhibiting Th1/17 differentiation. The inflammatory T cells were unable to induce obvious angiogenesis both *in vivo* and *in vitro*. The collective evidences show that QLY inhibited immune-mediated abnormal angiogenesis in AIA rats by activating PPARγ in T cells.

CRediT authorship contribution statement

Jian Zuo and **Yan Li** conceived the idea; **Meng-Ke Song** and **Qin Yin** performed majority of the experiments; **Meng-Fan Gu** and **Wen-Gang Chen** assisted all the experiments; **Jian Zuo** wrote the manuscript; **Opeyemi Joshua Olatunji** proof read the manuscript. All the authors gave final approval of the version to be published.

Funding

This work was supported by National Natural Science Foundation of China (82274465), Plans for Major Provincial Science & Technology Projects (202303a07020001), Excellent Research and Innovation Team of Anhui Provincial Colleges (2023AH010075), Anhui Provincial research projects of Traditional Chinese Medicine (2020ccyb03), and Scientific Research Fund of Wannan Medical College (WK2023XS54).

Data Availability

All data generated or analyzed during this study are included in this article and its supplementary information file.

Declaration of Competing Interest

The authors declare no conflict of interest.

References

1. Fukae J, Tanimura K, Atsumi T, Koike T. Sonographic synovial vascularity of synovitis in rheumatoid arthritis. Rheumatology 2014; 53(4): 586-91.

2. McInnes IB, Schett G. The pathogenesis of rheumatoid arthritis. N Engl J Med 2011; 365(23):2205-19.

3. Elshabrawy HA, Chen Z, Volin MV, Ravella S, Virupannavar S, Shahrara S. The

pathogenic role of angiogenesis in rheumatoid arthritis. Angiogenesis 2015; 18(4): 433-48.

4. Wang Y, Wu H, Deng R. Angiogenesis as a potential treatment strategy for rheumatoid arthritis. Eur J Pharmacol 2021; 910: 174500.

5. Tsaltskan V, Firestein GS. Targeting fibroblast-like synoviocytes in rheumatoid arthritis. Curr Opin Pharmacol 2022; 67: 102304.

6. Smolen JS, Landewé RBM, Bergstra SA, Kerschbaumer A, Sepriano A, Aletaha D, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2022 update. Ann Rheum Dis. 2023; 82(1): 3-18.

7. Li S, Wang R, Zhang Y, Zhang X, Layon AJ, Li Y, Chen M. Symptom combinations associated with outcome and therapeutic effects in a cohort of cases with SARS. Am J Chin Med. 2006;34(6):937-47.

8. □□□,□□. □□□□□□□□□□□□□□□□□□□□ [J]. □□□□□□□□□□□, 2015, 29 (06): 883-892.

9. Wang ZY, Wang X, Zhang DY, Hu YJ, Li S. [Traditional Chinese medicine network pharmacology: development in new era under guidance of network pharmacology evaluation method guidance]. Zhongguo Zhong Yao Za Zhi. 2022 Jan;47(1):7-17.

10. Zuo J, Wang X, Liu Y, Ye J, Liu Q, Li Y, Li S. Integrating network pharmacology and metabolomics study on anti-rheumatic mechanisms and antagonistic effects against methotrexate-induced toxicity of Qing-Luo-Yin. Front Pharmacol 2018; 9: 1472.

11. Wang DD, Wu XY, Dong JY, Cheng XP, Gu SF, Olatunji OJ, Li Y, Zuo J. Qing-Luo-Yin alleviated experimental arthritis in rats by disrupting immune feedback between inflammatory T cells and monocytes: key evidences from its effects on immune cell phenotypes. J Inflamm Res 2021; 14: 7467-86.

12. Wang R, Li DF, Hu YF, Liao Q, Jiang TT, Olatunji OJ, Yang K, Zuo J.

Qing-Luo-Yin alleviated monocytes/macrophages-mediated inflammation in rats

with adjuvant-induced arthritis by disrupting their interaction with (pre)-adipocytes

through PPAR-γ signaling. Drug Des Devel Ther 2021; 15: 3105-3118.

13. Ye P, Wang QH, Liu CS, Li GH, Olatunji OJ, Lin JT, Zuo J. SIRT1 inhibitors

within Qing-Luo-Yin alleviated white adipose tissues-mediated inflammation in

antigen-induced arthritis mice. Phytomedicine 2024; 122: 155132.

14. Li S, Lu AP, Wang YY, Li YD. Suppressive effects of a Chinese herbal medicine

Qing-Luo-Yin extract on the angiogenesis of collagen-induced arthritis in rats. Am

J Chin Med 2003; 31(5): 713-20.

15. Wu YJ, Fang WJ, Pan S, Zhang SS, Li DF, Wang ZF, Chen WG, Yin Q, Zuo J.

Regulation of Sirt1 on energy metabolism and immune response in rheumatoid

arthritis. Int Immunopharmacol 2021; 101(Pt A): 108175.

16. Liu Y, Wang J, Luo S, Zhan Y, Lu Q. The roles of PPARγ and its agonists in

autoimmune diseases: A comprehensive review. J Autoimmun 2020; 113: 102510.

17. Nobs SP, Kopf M. PPAR-γ in innate and adaptive lung immunity. J Leukoc Biol

2018; 104(4): 737-41.

18. Kwon EJ, Park EJ, Choi S, Kim SR, Cho M, Kim J. PPARγ agonist rosiglitazone

inhibits migration and invasion by downregulating Cyr61 in rheumatoid arthritis

fibroblast-like synoviocytes. Int J Rheum Dis 2017; 20(10): 1499-509.

19. Wagner N, Wagner KD. PPARs and angiogenesis-implications in pathology. Int J

Mol Sci 2020; 21(16): 5723.

20. Zhou Y, Xiang R, Qin G, Ji B, Yang S, Wang G, Han J. Xanthones from Securidaca

inappendiculata Hassk. attenuate collagen-induced arthritis in rats by inhibiting the

nicotinamide phosphoribosyltransferase/glycolysis pathway and macrophage

polarization. Int Immunopharmacol 2022; 111: 109137.

21. Liu JQ, Zhao XT, Qin FY, Zhou JW, Ding F, Zhou G, et al. Isoliquiritigenin

mitigates oxidative damage after subarachnoid hemorrhage in vivo and in vitro by

regulating Nrf2-dependent signaling pathway via targeting of SIRT1.

Phytomedicine 2022; 105: 154262.

22. Zhang S, Zhu P, Yuan J, Cheng K, Xu Q, Chen W, Pan Z, Zheng Y. Non-alcoholic

fatty liver disease combined with rheumatoid arthritis exacerbates liver fibrosis by

stimulating co-localization of PTRF and TLR4 in rats. Front Pharmacol 2023; 14:

1149665.

23. Sun Y, Bai YP, Wang DG, Xing YJ, Zhang T, Wang W, et al. Protective effects of

metformin on pancreatic β-cell ferroptosis in type 2 diabetes in vivo. Biomed

Pharmacother 2023; 168: 115835.

24. Hua Z, Hui LI, Haihua W. Potential protective effects of the water-soluble Chinese

propolis on experimental ulcerative colitis. J Tradit Chin Med 2023; 43(5): 925-33.

25. Wang X, Shen C, Wang X, Tang J, Wu Z, Huang Y, et al. Schisandrin protects

against ulcerative colitis by inhibiting the SGK1/NLRP3 signaling pathway and

reshaping gut microbiota in mice. Chin Med 2023; 18(1): 112.

26. Wu YJ, Zhang SS, Yin Q, Lei M, Wang QH, Chen WG, et al. α-Mangostin

inhibited M1 polarization of macrophages/monocytes in antigen-induced arthritis

mice by up-regulating silent information regulator 1 and peroxisome

proliferators-activated receptor γ simultaneously. Drug Des Devel Ther 2023; 17:

563-77.

27. Yin M, Ding X, Yin S, Wang L, Zhang K, Chen Y, et al. Exosomes from hepatitis B

virus-infected hepatocytes activate hepatic stellate cells and aggravate liver fibrosis

through the miR-506-3p/Nur77 pathway. J Biochem Mol Toxicol 2023; 37(10):

e23432.

28. Xu S, He L, Ding K, Zhang L, Xu X, Wang S, Qian X. Tanshinone IIA ameliorates

streptozotocin-induced diabetic nephropathy, partly by attenuating PERK

pathway-induced fibrosis. Drug Des Devel Ther 2020; 14:5773-82.

29. Zhou HH, Zhang YM, Zhang SP, Xu QX, Tian YQ, Li P, et al. Suppression of

PTRF alleviates post-infectious irritable bowel syndrome via downregulation of the

TLR4 pathway in rats. Front Pharmacol 2021; 12: 724410.

30. Jiang TT, Ji CF, Cheng XP, Gu SF, Wang R, Li Y, Zuo J, Han J. α-Mangostin
alleviated HIF-1α-mediated angiogenesis in rats with adjuvant-induced arthritis by
suppressing aerobic glycolysis. Front Pharmacol 2021; 12: 785586.

31. Liu Y, Sun H, Li C, Pu Z, Wu Z, Xu M, et al. Comparative HPLC-MS/MS-based
pharmacokinetic studies of multiple diterpenoid alkaloids following the
administration of Zhenwu Tang and Radix Aconiti Lateralis Praeparata extracts to
rats. Xenobiotica 2021; 51(3): 345-54.

32. Li H, Wang Y, Fan R, Lv H, Sun H, Xie H, et al. The effects of ferulic acid on the
pharmacokinetics of warfarin in rats after biliary drainage. Drug Des Devel Ther
2016; 10: 2173-80.

33. Cheng L, Wang H, Wang Z, Huang H, Zhuo D, Lin J. Leflunomide inhibits
proliferation and induces apoptosis via suppressing autophagy and PI3K/Akt
signaling pathway in human bladder cancer cells. Drug Des Devel Ther 2020; 14:
1897-908.

34. Lee H, Suh YS, Lee SI, Cheon YH, Kim M, Noh HS, Kim HO. Serum IGF-1 in
patients with rheumatoid arthritis: correlation with disease activity. BMC Res
Notes 2022; 15(1): 128.

35. Chen JY, Tian XY, Wei SS, Xu W, Pan RR, Chen LL, et al. Magnolol as STAT3
inhibitor for treating multiple sclerosis by restricting Th17 cells. Phytomedicine
2023; 117: 154917.

36. Yin Q, Wu YJ, Pan S, Wang DD, Tao MQ, Pei WY, Zuo J. Activation of
cholinergic anti-inflammatory pathway in peripheral immune cells involved in
therapeutic actions of α-mangostin on collagen-induced arthritis in rats. Drug Des
Devel Ther 2020; 14: 1983-93.

37. Lei M, Tao MQ, Wu YJ, Xu L, Yang Z, Li Y, et al. Metabolic enzyme
triosephosphate isomerase 1 and nicotinamide phosphoribosyltransferase, two
independent inflammatory indicators in rheumatoid arthritis: evidences from
collagen-induced arthritis and clinical samples. Front Immunol 2022; 12: 795626.

38. Kapellos TS, Bonaguro L, Gemünd I, Reusch N, Saglam A, Hinkley ER, Schultze

JL. Human monocyte subsets and phenotypes in major chronic inflammatory

diseases. Front Immunol 2019; 10: 2035.

39. Xu H, Li S , Liu J, Cheng J, Kang L, Li W, et al. Bioactive compounds from Huashi

Baidu decoction possess both antiviral and anti-inflammatory effects against

COVID-19. Proceedings of the National Academy of Sciences 2023;120(18):

e2301775120 .

40. Friedman B, Cronstein B. Methotrexate mechanism in treatment of rheumatoid

arthritis. Joint Bone Spine 2019; 86(3): 301-7.

41. Zhao Z, Hua Z, Luo X, Li Y, Yu L, Li M, et al. Application and pharmacological

mechanism of methotrexate in rheumatoid arthritis. Biomed Pharmacother 2022;

150: 113074.

42. Chen WG, Zhang SS, Pan S, Wang ZF, Xu JY, Sheng XH, et al. α-Mangostin treats

early-stage adjuvant-induced arthritis of rat by regulating the CAP-SIRT1 pathway

in macrophages. Drug Des Devel Ther 2022; 16: 509-20.

43. Wang Z, Cheng G, Wu M. Columbianetin improve rheumatoid arthritis and relative

mechanisms. Lat Am J Pharm 2021; 40(8): 1910-7.

44. Zhou W, Yang K, Zeng J, Lai X, Wang X, Ji C, Li Y, Zhang P, Li S. FordNet:

Recommending traditional Chinese medicine formula via deep neural network

integrating phenotype and molecule. Pharmacol Res. 2021 Nov;173:105752.

45. Zhou BH, Yang JY, Ding HY, Chen QP, Tian EJ, Wang HW. Anticoccidial effect

of toltrazuril and Radix Sophorae Flavescentis combination: Reduced inflammation

and promoted mucosal immunity. Vet Parasitol 2021; 296: 109477.

46. Zhao XX, Peng C, Zhang H, Qin LP. Sinomenium acutum: a review of chemistry,

pharmacology, pharmacokinetics, and clinical use. Pharm Biol 2012; 50(8):

1053-61.

47. Yin MC, Chang CH, Su CH, Yu B, Hsu YM. Pteris multifida, Cortex phellodendri,

and probiotics attenuated inflammatory status and immunity in mice with a

Salmonella enterica serovar Typhimurium infection. Biosci Biotechnol Biochem

2018; 82(5): 836-47.

48. Guo C, Ding G, Huang W, Wang Z, Meng Z, Xiao W. Total saponin of Dioscoreae

hypoglaucae rhizoma ameliorates streptozotocin-induced diabetic nephropathy.

Drug Des Devel Ther 2016; 10: 799-810.

49. Miao Y, Wu X, Xue X, Ma X, Yang L, Zeng X, et al. Morin, the PPARγ agonist,

inhibits Th17 differentiation by limiting fatty acid synthesis in collagen-induced

arthritis. Cell Biol Toxicol 2023; 39(4): 1433-52.

50. Wen S, He L, Zhong Z, Zhao R, Weng S, Mi H, Liu F. Stigmasterol restores the

balance of Treg/Th17 cells by activating the butyrate-PPARγ axis in colitis. Front

Immunol 2021; 12: 741934.

Figure 1. QLY reduced AIA severity in rats. **A**, periodic changes of arthritis scores and body weights; **B**, morphological changes of hind paws; **C**, relative weights of liver and spleen; **D**, levels of RA diagnostic indicators in blood; **E**, joint injuries in hind knee joints (arrows in the above H&E staining sections indicate angiogenesis; arrows in the below SOFG staining sections indicate cartilage erosion). Statistical significance: $^{*}p < 0.05$ and $^{**}p < 0.01$ compared with AIA model rats.

Figure 2. QLY improved metabolic disorders in AIA rats by activating PPARγ. **A**,

levels of representative metabolites in blood; **B**, levels of oxidative indicators in blood;

C, levels of IGF-1 in blood; **D**, levels of liver injuries indicators in blood; **E**, levels of

inflammatory cytokines in liver; **F**, levels of representative lipid metabolites in liver;

G, WB assay performed using liver samples; **H**, the quantification results of assay G.

Statistical significance: *p < 0.05 and **p < 0.01 compared with AIA model rats; $^{#}$p <

0.05 and $^{##}$p < 0.01 compared with normal healthy rats.

Figure 3. QLY-induced monocyte phenotype changes were unsustainable *in vitro*. **A**,

FCM analysis of different monocyte subsets in the rats' blood; **B**, the quantification

results of assay A; **C**, levels of inflammatory monocyte indicators in blood; **D**, levels

of cytokines released by the *in vitro* cultured rat monocytes. Statistical significance:

*p < 0.05 and **p < 0.01 compared with AIA model rats.

Figure 4. QLY preferentially targeted inflammatory T cells when treating AIA rats. **A**, expression of various Th subsets-related genes in the normal lymphocytes grew in the monocyte culture medium from the experiment above; **B**, levels of cytokines released by AIA rats' monocytes cultured by normal or QLY-containing serum; **C**, expression and co-localization of IL-1β and PPARγ in these monocytes; **D**, levels of cytokines released by RA patients' monocytes cultured by normal or QLY-containing serum; **E**, FCM analysis of these cells; **F**, the quantification results of assay E; **G**, levels of cytokines released by the *in vitro* cultured lymphocytes from different rats; **H**, levels of cytokines released by normal monocytes grew in the lymphocyte culture medium above; **I**, WB assay performed using spleen samples; **J**, the quantification results of assay I. Statistical significance: $^{*}p < 0.05$ and $^{**}p < 0.01$ compared with AIA group.

Figure 5. Transfusion using different rats' blood brought little impacts on the internal environment of healthy recipients. **A**, the dynamic distribution of AIA rats' peripheral immune cells in the recipient after transplant; **B**, FCM analysis of monocyte subsets in the recipients' blood; **C**, the quantification results of assay B; **D**, a comprehensive evaluation of the internal environment changes by determining a panel of cytokines in blood; **E**, levels of iNOS in the recipients' blood; **F**, levels of oxidative indicators in the recipients' blood. Statistical significance: $^{*}p < 0.05$ compared with AIA group.

Figure 6. QLY impaired the ability of AIA rats' lymphocytes inducing angiogenesis *in vivo*. **A**, levels of immune/angiogenesis-related cytokines in the recipients' joints; **B**, levels of iNOS in the joints; **C**, levels of oxidative indicators in the joints; **D**, matrigel plugs retrieved from the recipients transfused by different rats' blood; **E**, histological examination of H&E stained plug sections. Statistical significance: [*]p < 0.05 and [**]p < 0.01 compared with AIA group.

Figure 7. QLY-related compounds inhibited T cells-meditated angiogenesis *in vitro*.
A, HPLC detection of the compounds in QLY-containing serum; **B**, the quantification
curves developed using the reference compounds; **C**, the compounds concentrations
detected in QLY-containing serum; **D**, levels of cytokines released by THP-1 cells
stimulated by AIA serum and the compound mixture; **E**, levels of cytokines released
by Jurkat cells stimulated by AIA serum and the compound mixture; **F**, levels of
VEGF released by HUVEC cells grew in different Jurkat cells culture mediums; **G**-**H**,
the impacts of different Jurkat cells culture mediums on the wound healing, transwell
infiltration, and tube-forming capacities of HUVEC cells. Statistical significance: *p <
0.05 and **p < 0.01 compared with AIA group; $^{\#\#}$p < 0.01 compared with compounds
treatment group.