Title: Social and environmental predictors of gut microbiome age in wild baboons

Authors

Mauna R. Dasari[1,2,3,*], Kimberly E. Roche[4], David Jansen[1], Jordan Anderson[5], Susan C.

Alberts[5,6,7], Jenny Tung[5,6,7,8,9,10], Jack A. Gilbert[11], Ran Blekhman[12], Sayan

Mukherjee[13,14,15], Elizabeth A. Archie[1,*]

Affiliations

[1] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA

[2] Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA

[3] California Academy of Sciences, San Francisco, CA, USA

[4] Program in Computational Biology and Bioinformatics, Duke University, Durham, NC, USA

[5] Department of Evolutionary Anthropology, Duke University, Durham, NC, USA

[6] Department of Biology, Duke University, Durham, NC, USA

[7] Duke University Population Research Institute, Duke University, Durham, NC, USA

[8] Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary Anthropology,

04103 Leipzig, Germany

[9] Canadian Institute for Advanced Research, Toronto, Ontario, Canada

[10] Faculty of Life Sciences, Institute of Biology, Leipzig University, Leipzig, Germany

[11] Department of Pediatrics and the Scripps Institution of Oceanography, University of California, San

Diego, San Diego, CA, USA

[12] Section of Genetic Medicine, Department of Medicine, University of Chicago, Chicago, IL, USA.

[13] Departments of Statistical Science, Mathematics, Computer Science, and Bioinformatics & Biostatistics,

Duke University, Durham, NC, USA

[14] Center for Scalable Data Analytics and Artificial Intelligence, University of Leipzig, Leipzig Germany

[15] Max Planck Institute for Mathematics in the Natural Sciences, Leipzig, Germany

*Corresponding authors. mauna.dasari@gmail.com (M.R.D.), earchie@nd.edu (E.A.A.)

Abstract

Mammalian gut microbiomes are highly dynamic communities that shape and are shaped

by host aging, including age-related changes to host immunity, metabolism, and behavior. As such,

gut microbial composition may provide valuable information on host biological age. Here we test

this idea by creating a microbiome-based age predictor using 13,563 gut microbial profiles from

479 wild baboons collected over 14 years. The resulting "microbiome clock" predicts host

chronological age. Deviations from the clock's predictions are linked to some demographic and

socio-environmental factors that predict baboon health and survival: animals who appear old-for-

age tend to be male, sampled in the dry season (for females), and have high social status (both

sexes). However, an individual's "microbiome age" does not predict the attainment of

developmental milestones or lifespan. Hence, in our host population, gut microbiome age largely

reflects current, as opposed to past, social and environmental conditions, and does not predict the

pace of host development or host mortality risk. We add to a growing understanding of how age is

reflected in different host phenotypes and what forces modify biological age in primates.

MAIN TEXT

INTRODUCTION

For most vertebrate species, physical declines with age are inevitable. These changes define the concept of biological aging and contribute to increased disease burden in older individuals (*1*, *2*). While vertebrates have species-typical patterns of biological aging, those patterns can also differ between individuals within species. Hence, an animal's age in years—i.e., its chronological age— is not an exact reflection of age-related decline in physical functioning (*3–6*). Measuring individual differences in biological age is an important first step to understand how socio-environmental conditions influence aging processes and to identify strategies to improve health in old age.

In mammals, one valuable marker of biological aging may lie in the composition and dynamics of the gut microbiome (*7–10*). Age-related changes in gut microbiomes are well documented in humans and other animals, and the gut microbiome has the potential to reflect a wide variety of aging processes for individual hosts (*11–24*). Mammalian gut microbiomes interact with the immune, endocrine, nervous, and digestive systems, all of which change with age (*25–28*). Gut microbiomes are also sensitive to host environments and behaviors that change with age, including host diet, living conditions, and social integration (*23*, *29–32*). Finally, gut microbiomes may play a causal role in age-related changes in host development and longevity and may, therefore, be directly involved in individual differences in biological age (*16*, *17*, *22*, *33*, *34*). For example, children who experience famine exhibit developmentally immature gut microbiomes that, when transplanted into mice, delay growth and alter bone morphology (*35–38*). Experiments in flies, mice, and killifish find that the gut microbiome can also influence longevity (*16*, *17*, *22*, *39*).

One strategy for testing if gut microbiomes reflect host biological age is to apply supervised machine learning to microbiome compositional data to develop a model for predicting host chronological age, and then to test whether deviations from the resulting "microbiome clock" age predictions are explained by socio-environmental drivers of biological age and/or predict host development or mortality. A parallel approach is commonly applied to patterns of DNA methylation, and epigenetic clock age estimates have been shown to predict disease and mortality risk more accurately than chronological age alone (*40–45*). To date, at least five microbiome age-predicting clocks have been built for humans, which predict sample-specific age with median error of 6 to 11 years (*46–49*). However, to our knowledge, no clocks have tested whether microbiome age is sensitive to potential socio-environmental drivers of biological age or predicts host development or mortality.

Here we create a microbiome-based age-predicting clock using 13,476 16S rRNA gene sequencing-based gut microbiome compositional profiles from 479 known-age, wild baboons (*Papio sp.*) sampled over a 14-year period (**Figures 1A and 1B**). These microbiome profiles represent a subset of a data set previously described in Grieneisen et al. (*50*), Björk et al. (*51*) and Roche et al. (*52*), filtered to include only the baboon hosts whose ages were known precisely (within a few days' error). Important to human aging, baboons share many developmental similarities with humans, including an extended juvenile period, followed by sexual maturation and non-seasonal breeding across adulthood (**Figures 1C and 1D**; [*51–54*]).

The baboons in our data set were members of the well-studied Amboseli baboon population in Kenya, which has continuous, individual-based data on early-life environments, maturational milestones, social relationships, and mortality (*57–62*). Relevant to measuring biological age, prior research in Amboseli has identified several demographic, environmental, and social conditions that predict physical condition, the timing of development, or survival, including sex, season, social status (i.e., dominance rank), and early-life adversity (*53–55*, *63–69*). Consistent with the possibility that these associations arise from causal effects of harsh or stressful conditions on

biological aging (*70–72*), and with the idea that microbiomes serve as a marker of biological age (*7–10*), we tested whether socio-environmental conditions predicted microbiome age estimates. Our predictions about the direction of these effects varied depending on the socio-environmental condition and the developmental stage of the animal (i.e., juvenile or adult).

In terms of sex, adult male baboons exhibit higher mortality than adult females. Hence, we predicted that adult male baboons would exhibit gut microbiomes that are old-for-age, compared to adult females (by contrast, we expected no sex effects on microbiome age in juvenile baboons).

In terms of season, the Amboseli ecosystem is a semi-arid savannah with a 5-month long dry season during which little rain falls, often leading to nutritional hardship (*73*). We predicted that samples from the dry season might appear to be old-for-age, compared to samples from the wet season due to nutritional stress in this difficult season.

In terms of social status, baboons experience linear, sex-specific hierarchies. Female ranks are nepotistic, with little social mobility, and low-rank is linked to low priority of access to food (*55, 68, 74–76*). In contrast, adult male rank is determined by strength and fighting ability and is dynamic across adulthood (*77*). High-ranking males experience high energetic costs of mating effort, have altered immune responses, and exhibit old-for-age epigenetic age estimates compared to low-ranking males (*45, 64, 65*). We expected that individuals who pay the largest energetic costs—low-ranking adult females and high ranking adult males (*45*)—would appear old-for-age.

In terms of early-life adversity, prior research in Amboseli has identified six conditions whose cumulative, and sometimes individual, effects predict adult female mortality, including maternal loss prior to age 4 years, drought in the first year of life, birth into an especially large social group, the presence of a close-in-age competing younger sibling, and having a low-ranking or socially isolated mother (*64, 65, 67*). For adult female baboons, experiencing multiple sources of adversity in early life is the strongest socio-environmental predictor of mortality; hence, we expected that these individuals would have old-for-age clock estimates in adulthood (*58, 59, 64, 66, 67, 78*). However, we also expected that some sources of early-life adversity might be linked to young-for-age gut microbiomes in juvenile baboons. For instance, famine is linked to gut microbial immaturity (*35–38*), and maternal social isolation might delay gut microbiome development due to less frequent microbial exposures from conspecifics. In contrast to the expectation that harsh early-life conditions are linked to old-for-age microbiomes in adult baboons, a viable alternative is that gut microbiome age might be better predicted by an individual's current environmental or social conditions (e.g., season or social status), rather than past events. Indeed, gut microbiomes are highly dynamic and can change rapidly in response to host diet or other aspects of host physiology, behavior, or environments (*79–81*). Such results would support recency models for biological aging (*82, 83*) and would be consistent with findings from a recent epigenetic clock study in Amboseli (*45*).

We began our analyses by identifying gut microbiome features that change reliably with host age. We then constructed a microbiome clock by comparing the performance of several supervised machine learning algorithms to predict host age from gut microbial composition in each sample from each host. We evaluated the clock's performance for male and female baboons and tested whether deviations from clock performance were predicted by the baboons' social and environmental conditions (guided by the predictions outlined above). Lastly, we tested whether baboons with young-for-age gut microbiomes have correspondingly late developmental timelines or longer lifespans. In general, our results support the idea that a baboon's current socio-environmental conditions, especially their current social rank and the season of sampling, have stronger effects on microbiome age than early-life events—many of which occurred many years prior to sampling. As such, the dynamism of the gut microbiome may often overwhelm and erase

early-life effects on gut microbiome age. Our work highlights the diversity of ways that social and
environmental conditions shape microbiome aging in natural systems.

Figure 1. Microbiome sampling time series and developmental milestones for the Amboseli
baboons. Plot (A) shows microbiome samples from female baboons, and plot (B) shows samples
from male baboons. Each point represents a microbiome sample from an individual subject (y-axis)
collected at a given host age in years (x-axis; n=8,245 samples from 234 females shown in yellow;
n=5,231 samples from 197 males shown in blue). Light and dark point colors indicate whether the
baboon was sexually mature at the time of sampling, with lighter colors reflecting samples collected
prior to menarche for females (n=2,016 samples) and prior to testicular enlargement for males
(n=2399 samples). Due to natal dispersal in males, we have fewer samples after the median age of
first dispersal in males (n=1,705 samples, 12.6% of dataset) than from females after the same age
(n=4408 samples, 32.6% of dataset). The timelines below the plots indicate the median age in years
at which (C) female baboons attain the developmental milestones analyzed in this paper—adult

rank, menarche and first live birth—and (D) males attain adult rank, testicular enlargement, and
disperse from their natal groups (*55*, *56*). The age at which 75% of animals in the population have
died is shown to indicate different life expectancies for females versus males (*63*). Baboon
illustrations courtesy of Emily (Lee) Nonnamaker.

RESULTS

Many microbiome features change with age

Before creating the microbiome clock, we characterized microbiome features that change
reliably with the age of individual hosts. Our subjects were 479 known-age baboons (264 females
and 215 males) whose microbiome taxonomic compositions were characterized using 13,476 fecal-
derived 16S rRNA gene sequencing profiles over a 14-year period (**Figures 1A and 1B;** baboon
age ranged from 7 months to 26.5 years; 8,245 samples from females; 5,231 samples from males;
range=3-135 samples per baboon; mean=35 samples per female and 26 samples per male).

We tested age associations for 1,440 microbiome features, including: (i) five metrics of
alpha diversity; (ii) the top 10 principal components of Bray-Curtis dissimilarity (which collectively
explained 57% of the variation in microbiome community composition); and (iii) centered log-ratio
transformed abundances (*84*) of each microbial phylum (n=30), family (n=290), genus (n=747),
and amplicon sequence variance (ASV) detected in >25% of samples (n=358; **Table S1**; see
methods for details on centered log-ratio transformations). We tested these different taxonomic
levels in order to learn whether the degree to which coarse and fine-grained designations categories
were associated with host age. For each of these 1,440 features, we tested its association with host
age by running linear mixed effects models that included linear and quadratic effects of host age
and four other fixed effects: sequencing depth, the season of sample collection (wet or dry), the
average maximum temperature for the month prior to sample collection, and the total rainfall in the
month prior to sample collection (*50*, *51*, *62*). Baboon identity, social group membership,
hydrological year of sampling, and sequencing plate (as a batch effect) were modeled as random
effects.

We found that many aspects of microbiome community composition changed with host age
(**Figure 2; Figure S1**). All alpha diversity metrics, except richness, the only unweighted metric,
exhibited U-shaped relationships with age, with high values in early life and old age, and low values
in young adulthood. While we should interpret this pattern with caution due to the small sample
size beyond age 20 (n=18 females), this U-shaped pattern differs somewhat from patterns in humans
and chimpanzees: most human populations exhibit an asymptotic increase in alpha diversity with
age (*85*, *86*) while in chimpanzees alpha diversity is highest in early life (*23*) (FDR < 0.05; **Figures
2C and 2E**; **Table S1**). Further, seven of the ten principal components (PCs) of microbiome
composition exhibited linear, and in some cases quadratic, relationships with age, with PC1, PC2,
PC4 and PC6 exhibiting the strongest age-associations (FDR < 0.05; **Figures 2C and 2F**; **Table
S1**).

51.6% of the 1,440 features exhibited significant linear or quadratic relationships with host
age (**Figure S1 and S2**; **Table S1**; FDR < 0.05). 60% of phyla (18 of 30) decreased proportionally
with age, while only three phyla—Kiritimatiellaeota, Firmicutes, and Chlamydiae—increased
proportionally with age (FDR < 0.05; **Table S1**). Similarly, 66% (66 of 100) of age-associated
families and 55% (115 of 209) of age-associated genera exhibited declining proportions with age
(**Table S1**). Consistent with the idea that age-related taxa differ across host populations and host
taxa, none of the taxa that changed in this baboon population were commonly age-associated in
humans (*85*). The taxa most consistently linked to human aging include *Akkermansia,*
Faecalibacterium, *Bacteroidaceae*, and *Lachnospiraceae* (*34*, *85*) while in our sample of baboons,

the strongest age-related changes were seen in the families *Campylobacteraceae, Clostridiaceae*, *Elusimicrobiaceae Enterobacteriaceae*, *Peptostreptococcaceae*, and an uncharacterized family within Gastranaerophilales (**Figure 2D and 2F**; **Table S1**). The genera that had the strongest relationships with age included *Camplylobacter*, *Catenibacterium, Elusimicrobium*, *Prevotella*, *Romboutsia*, and *Ruminococcaceae UCG-011* (**Figure 2D and 2F**; **Table S1**).

Figure 2: Microbiome features change with age. (A) and (B) show the percent mean relative abundance of microbial phyla across life for females and males respectively. Panel (C) shows the estimates of the linear associations between mean-centered age for metrics of microbiome alpha diversity and principal components of microbiome compositional variation that exhibited

significant associations with age (FDR < 0.05). Positive values are more abundant in older hosts. Panel (D) shows the estimate of the linear association between mean-centered age and the top 50 microbiome features that exhibited significant associations with age. Positive values are more abundant in older hosts. Panel (E) shows the average value of the microbiome features from (C) as a function of age, across all subjects. Note that sample sizes for patterns beyond age 20 years rely on 256 samples from just 18 females; hence, we interpret the pattern in these oldest animals with caution. Panel (F) shows the average prevalence of the higher taxonomic designations from (D) as a function of age, across all subjects. In (C-F) points are colored by the category of the feature (see legend). UC is an abbreviation for uncharacterized. Features that had a significant quadratic age term are indicated by * (see also **Figure S2; Table S1**).

Microbiome clock calibration and composition

We next turned our attention to building a microbiome clock using all 9,575 microbiome compositional and taxonomic features present in at least 3 samples (**Table S2**; the 1,440 features discussed above only included taxa present in >25% of samples). We included all 9,575 microbiome features in our age predictions, as opposed to just those that were statistically significantly associated with age because removing these non-significant features could exclude features that contribute to age prediction via interactions with other taxa. In developing the clock, we compared the performance of three supervised machine learning methods to predict the chronological age of individual hosts at the time each microbiome sample was collected. The three machine learning methods were elastic net regression, Random Forest regression, and Gaussian process regression (see Supplementary Methods and Results). Because gut microbiomes are highly personalized in ours and other host populations (*51*), at least one sample from each host was always present in the training sets for these models (see Methods).

We found that the most accurate age predictions were produced by a Gaussian process regression model with a kernel customized to account for heteroscedasticity (**Figure 3**; **Figure S3**). This model predicted host chronological age (age$_c$), with an adjusted R^2 of 48.8% and a median error of 1.96 years across all individuals and samples (**Figure 3A**, **Table 1**). As has been observed in some previous age clocks (e.g., *39*, *44*, *46*), microbial age estimates (age$_m$) were compressed relative to the x=y line, leading the model to systematically over-predict the ages of young individuals and under-predict the ages of old individuals (**Figure 3A**).

When we subset our age$_m$ estimates by sex, we found that the microbiome clock was slightly more accurate for males than for females (**Figure 3B**, **Table 1**). The adjusted R^2 for the correlation between age$_c$ and age$_m$ for males was 50.0%, with a median prediction error of 1.71 years as compared to an adjusted R^2 of 48.9% and median error of 2.15 years for female baboons (**Table 1**). Male baboons also exhibit significantly older gut microbial age than females (**Figure 3B**, chronological age by sex interaction: β=0.18, p<0.001, **Table S3**). Across the lifespan, males show a 1.4-fold higher rate of change in age$_m$ as a function of age$_c$ compared to females (relationship between age$_c$ and age$_m$ in males: β=0.63, p< 0.001; relationship between age$_c$ and age$_m$ in females: β=0.45, p < 0.001; **Table S4**). Similar to patterns from a recent epigenetic age-predicting clock developed for this population (*45*), this effect was only present after sexual maturity: when we subset the age$_m$ estimates to microbiome samples collected prior to the median age at sexual maturity (5.4 years for testicular enlargement in males and 4.5 years for menarche in females; *54*), we found no significant interaction between sex and age (age$_c$ by host sex interaction prior to median age of maturity: β= -0.09, p=0.203; age$_c$ by host sex interaction after median age of maturity: β=0.15, p<0.001; **Table S3**). After maturity, we recapitulate the 1.4-fold higher rate of change in males compared to females observed in the full data set (relationship between age$_c$ and

age$_m$ in males only: β=0.53, p<0.001; relationship between age$_c$ and age$_m$ in females only: β=0.38,
p<0.001; **Table S4**).

Overall, age$_m$ estimates performed reasonably well compared to other known predictors of
age in the Amboseli baboons (**Table S5;** [*44*]). Age$_m$ performed similarly to the non-invasive
physiology and behavior clock (NPB clock) recently developed for this population (*87*); R^2=51%;
median error=2.33 years). Age$_m$ was a stronger predictor of host chronological age than body mass
index (BMI; except juvenile male BMI), blood cell composition from flow cytometry, and
differential white blood cell counts from blood smears (**Table S5**; *45, 54, 87–89*). However, age$_m$
was a less accurate predictor of chronological age than both dentine exposure (males, females
respectively: adjusted R^2=73%, 85%; median error=1.11 years, 1.12 years; **Table S5**) and an
epigenetic clock based on DNA methylation data (males, females respectively: adjusted R^2=74%,
60%; median error=0.85 years, 1.62 years; **Table S5**; *44*).

Table 1. Comparison of Gaussian process regression model performance between sexes. Model
accuracy was determined based on the correlation between known chronological age (age$_c$) and
predicted age (age$_m$), the variance explained in age$_c$ by age$_m$ (R^2), and the median absolute
difference between age$_c$ and age$_m$ (*40*).

Subset	Sample Size	R^2	Pearson's R	Median Error (years)
All Subjects	13,476	48.8%	0.698	1.962
Females Only	8,245	48.9%	0.699	2.150
Males Only	5,231	50.0%	0.707	1.706

Figure 3. Microbiome clock age predictions in wild baboons. Panels (A) and (B) show predicted
microbiome age in years (age$_m$) from a Gaussian process regression model, relative to each
baboon's true chronological age in years (age$_c$) at the time of sample collection. Each point

represents a microbiome sample. Panel (A) shows linear fit for all subjects in the model; (B) shows
separate linear fits for each sex (**Table S4**). Dashed lines show the 1-to-1 relationship between age_c
and age_m. Panel (A) also shows the measurement of sample-specific microbiome Δage compared
to chronological age. Whether an estimate is old- or young-for-chronological age is calculated for
each microbiome sample as the difference between age_m and age_c. Because of model compression
relative to the 1-to-1 line, we correct for host chronological age by including chronological age in
any model. An example of an old-for-age sample is shown as a red point, with dashed lines showing
the value of age_c for a given sample with its corresponding age_m.

Social and environmental conditions predict variation in microbiome age

To test whether deviations in microbiome age for a given chronological age are correlated
with socio-environmental predictors of health and mortality risk, we calculated whether
microbiome age estimates from individual samples were older or younger than their hosts' known
chronological ages (Δage in years; **Figure 3C**). We then tested whether several social and
environmental variables predicted individual variation in microbiome Δage in years (**Table S7;**
note that whether microbiome ages are old- or young-for-age is correlated with host age, hence our
models always included host chronological age as a covariate). Overall, we expected that adult
baboons who experienced harsh conditions in early-life adversity (the strongest socio-
environmental predictor of adult mortality in Amboseli) would tend to look old-for-age based on
the microbiome clock (*58, 59, 64, 66, 67, 78*). Alternatively, microbiome deviations from
chronological age might be best predicted by an individual's current social status or season, rather
than past events. These results would support recency models of biological aging (*82, 83*) and would
be consistent with findings from a recent epigenetic clock study in Amboseli (*45*).

We found that individual baboons varied considerably in gut microbiome Δage. For
instance, in mixed effects models, individual identity explained ~25% to ~50% of the variance in
Δage for females and males respectively over the course of their lives (**Table S6**). Further, we found
that season, dominance rank, and some aspects of early-life adversity (large group size, drought,
and maternal social isolation) were linked to small deviations from chronological age. In support
of our expectation that microbiome samples collected in the dry season are old-for-chronological
age, we found that age estimates based on microbiome samples collected from female baboons in
the dry season were ~2 months older than the host's true chronological age (β=-0.180, p=0.021,
Table 2; Table S6C; Figure S6A). However, season did not significantly predict the difference
between microbiome age and known age in male baboons.

In terms of social status, we expected to observe that low-ranking females and high-ranking
males would be old-for-chronological age (*45*). In support, we found that estimates from high-
ranking males were old-for-age compared to estimates from low-ranking males, but these effects
were relatively weak and noisy (rank effect: β=0.033, p<0.001; **Figure 4A; Table 2; Tables S6B**
and **D**). Specifically, controlling for chronological age, alpha male gut microbiomes (ordinal
rank=1) appeared to be approximately 4 months older than microbiomes sampled from males with
an ordinal rank of 10 (**Table 2**). However, contrary to our expectations, high-ranking female
baboons also had old-for-age estimated when compared to low-ranking females (rank effect:
β=1.745, p<0.001; **Figure 4B; Table 2; Tables S6A** and **C**). Specifically, controlling for
chronological age, the microbiome of an alpha female (proportional rank=1) appeared to be
approximately 1.75 years older than the lowest-ranking females in the population (proportional
rank=0; **Table 2**).

Some forms of early-life adversity also predicted variation in microbiome Δage, but only in
males, and in inconsistent directions. For instance, males born into the highest quartile of observed

group sizes had old-for-age estimates. Males experiencing this source of early-life adversity had
gut microbiomes that were predicted to be ~5.4 months older than males not experiencing this
source of adversity (β=0.471, p=0.033, **Table 2; Table S6D; Figure S6C**). However, early-life
drought and maternal social isolation were linked to young-for-age gut microbiomes in males
(drought effect: β=-0.451, p=0.021; maternal social isolation effect: β=-0.395, p=0.006, **Table 2;**
Table S6D; Figure S6D). Probably as a result of these conflicting effects, we found no effect of
cumulative early adversity on microbiome Δage in males (**Table S6B**).

Table 2. Social and environmental factors predicting variation in microbiome Δage in female and
male baboons. Models below only show variables that minimize the Akaike information criterion
(AIC) for each model; see **Table S6** for full models. Coefficients for social dominance rank are
transformed so higher values reflect higher rank/social status (see footnotes).

Fixed Effect	β	p-value	Interpretation
Predictors of microbiome Δage in females (n=6,743 samples from 192 females)			
Chronological age	-0.551	<0.001	Included to control for the correlation between chronological age and microbiome Δage (Fig. 3)
Season	-0.180	0.021	Dry season samples are microbially old-for-age
Proportional rank*	1.745	<0.001	Low-ranking females are microbially young-for-age
Predictors of microbiome Δage in males (n=4,355 samples from 168 males)			
Chronological age	-0.404	<0.001	Included to control for the correlation between chronological age and microbiome Δage (Fig. 3)
Ordinal rank**	0.033	<0.001	Low-ranking males are microbially young-for-age
Born in a drought	-0.451	0.021	Males born during a drought are microbially young-for-age
Born into a large group	0.471	0.033	Males born into large groups are microbially old-for-age
Socially isolated mother	-0.395	0.006	Males with a socially isolated mother are microbially young-for-age

337 *proportional rank ranges from 0 to 1, with higher values reflecting higher social status
338 **ordinal rank is an integer ranking, with lower values reflecting higher social status; we have inverted the sign
339 of the coefficient so higher numbers reflect higher rank to facilitate comparison to females

Figure 4. Social dominance rank predicts gut microbiome Δage in male and female baboons (corrected for confounders). Panels (A) and (B) show the relationship between host proportional dominance rank and corrected gut microbiome Δage in (A) males (blue points) and (B) females (yellow points). Each point represents an individual gut microbiome sample. Corrected microbiome Δage is calculated as the residuals of age$_m$ correcting for host chronological age, season, monthly temperature, monthly rainfall, and social group and hydrological year at the time of collection.

Microbiome age does not predict the timing of development or survival

Finally, we tested whether variation in microbiome Δage predicted the timing of individual maturational milestones or survival using Cox proportional hazards models (**Table S8**). Maturational milestones for females were the age at which they attained their first adult dominance rank, reached menarche, or gave birth to their first live offspring (**Figure 1C**). Male maturational milestones were the age at which they attained testicular enlargement, dispersed from their natal social group, or attained their first adult dominance rank (**Figure 1D**). We also tested if microbiome Δage predicted juvenile survival (in females and males) or adult survival (females only). We did not test adult survival in males because male dispersal makes it difficult to know age at death for most males (*90*).

Contrary to our expectations, microbiome Δage did not predict the timing of any baboon developmental milestone or measure of survival **(Tables S9, S10)**. However, these patterns should be treated with caution, as reflected by the large number of censored animals, large hazard ratios, and small sample sizes for some tests.

DISCUSSION

We report three main findings. First, similar to humans and other animals (*11–24*), baboon gut microbiomes show age-related changes in taxonomic composition that produce a dependable microbiome-based age predictor—a microbiome clock. This clock explains nearly half the variance in true host chronological age, and variation in its age predictions recapitulate well-known patterns of faster male senescence in humans and other primates (*91*). Second, parallel to a recent epigenetic clock in the Amboseli baboons (*45*), deviations from microbiome age predictions are explained by

socio-environmental conditions experienced by individual hosts, especially recent conditions, although the effect sizes are small and are not always directionally consistent. Notably, recent social competition as reflected in social dominance rank predicts both microbiome and epigenetic age . Third, microbiome age did not predict the timing of individual development or survival (but caution is warranted because sample sizes were small for some tests). Hence, in our host population, gut microbial age reflects current social and environmental conditions, but not necessarily the pace of development or mortality risk.

To date, at least five other microbiome clocks have been built—all in human subjects—that predict host chronological age (*46–49*). Compared to these clocks, our clock in baboons has comparable or better predictive power, with a median error of 1.96 years, compared to 6 to 11 years in human age-predicting clocks ((*46–49*); baboon lifespans are approximately one third of a human lifespan). Age prediction from gut microbial composition may be more successful in baboons than humans for at least three reasons. First, signs of age in the baboon gut microbiome may be more consistent across hosts, perhaps because of the relative homogeneity in host environments and lifestyles in baboons compared to humans (*51*). Second, the baboon clock relies on dense longitudinal sampling for each host, and because the training data set included at least one sample from each host, the microbiome clock may be better able to address personalized microbiome compositions and dynamics than clocks that rely on cross-sectional data (e.g. *46–48*). However, because our training data set was not naïve to information from the host being predicted, this approach could leak information between the training and test set. Third, our Gaussian process regression approach allowed us to account for non-linear and interactive relationships between microbes with age, leveraging a wider variety of age-related signatures in the microbiome than other machine learning approaches (e.g., elastic net regression or random forest regression).

Despite the relative accuracy of the baboon microbiome clock compared to similar clocks in humans, our clock has several limitations. First, the clock's ability to predict individual age is lower than for age clocks based on patterns of DNA methylation—both for humans and baboons (*40–43, 45*). One reason for this difference may be that gut microbiomes can be influenced by several non-age-related factors, including social group membership, seasonal changes in resource use, and fluctuations in microbial communities in the environment. Second, gut microbiomes are highly personalized: each host species, population, and even host individuals within populations have distinctive, characteristic microbiomes, which likely limits the utility of our clock beyond our study population (*51, 60, 62, 92*). To make a more generalizable clock, an important next step would be to train the clock on data from many more host populations and incorporate features of the gut microbiome that are broader and more universal across host species and populations. Third, the relationships between potential socio-environmental drivers of biological aging and the resulting biological age predictions were inconsistent. For instance, some sources of early-life adversity were linked to old-for-age gut microbiomes (e.g., males born into large social groups), while others were linked to young-for-age microbiomes (e.g., males who experienced maternal social isolation or early-life drought), or were unrelated to gut microbiome age (e.g., males who experienced maternal loss; any source of early-life adversity in females).

The most consistent socio-ecological driver of baboon microbiome age was individual dominance rank. Microbiome samples from high-ranking males and females both appeared old-for-age. These results are interesting considering a growing body of evidence that finds rank-related differences in immunity and metabolism, including costs of high rank, especially for males (*45, 64, 65, 93–95*). For instance, in Amboseli, high social status in males is linked to old-for-age epigenetic age estimates (*45*), differences in immune regulation (*65, 93*), and, for alpha males, elevated glucocorticoid levels (*64*). These patterns, together with our evidence that high-ranking males tend to look 'old-for-age', are consistent with the idea that high-ranking males pursue a "live fast die young" life history strategy (*45*).

However, we also found evidence for old-for-age microbiome age estimates in samples from high-ranking females who do not seem to be "living fast" in the same sense as high-ranking males (indeed alpha females have *lower* glucocorticoid hormones than other females (*96*)). This outcome points towards a shared driver of high social status in shaping gut microbiome age in both males and females. While it is difficult to identify a plausible shared driver, one benefit shared by both high-ranking males and females is priority of access to food. This access may result in fewer foraging disruptions and a higher quality, more stable diet. At the same time, prior research in Amboseli suggests that as animals age, their diets become more canalized and less variable (*50*). Hence aging and priority of access to food might both be associated with dietary stability and old-for-age microbiomes. However, this explanation is speculative and more work is needed to understand the relationship between rank and microbiome age.

While some social and environmental conditions associated with baboon development or mortality predict microbiome age, our microbiome clock predictions do not themselves predict baboon development or mortality. This finding supports the idea that microbiome age is sensitive to transient social and environmental conditions; However, these patterns do not have long-term consequences for development and mortality. One reason for this may be that the biological drivers of development and mortality are too diverse to be well reflected in gut microbial communities. For instance, animals in Amboseli die for many reasons, including interactions with predators and humans, conflict with conspecifics (*97*), and disease, and the biological predictors of these events in the gut microbiome are likely weaker or more diverse than the biological signals that predict developmental milestones (i.e., sex steroids, growth hormones, metabolic status, and physical condition).

Three important future directions of our work will be to test: (i) whether microbiome age is correlated with other hallmarks of biological age in this population; (ii) whether it is possible to build a microbiome-based predictor of individual lifespan (i.e., *remaining* lifespan as opposed to years already lived); and (iii) the relationships between microbiome compositional features and individual survival. While several metrics of biological age have been developed for this population (*45*, *54*, *87–89*), testing the correlations between them is complex because these metrics were measured for different sets of subjects at different ages and with different sampling designs (e.g., longitudinal versus cross-sectional measures). We also hope to measure epigenetic age in fecal samples, leveraging methods developed in Hanski et al. (*98*).

In sum, our findings support the hypothesis that the gut microbiome is a biomarker of some aspects of host age. By leveraging microbial, social, environmental, and life history data on host individuals followed from birth to death, we bolster the validity of microbiome clock studies in humans and find some socio-environmental predictors of microbiome age. Future work may benefit from searching for more universal aspects of the microbiome that may predict host aging across populations and even host species.

MATERIALS AND METHODS

Study population and subjects

Our study subjects were 479 wild baboons (215 males and 264 females) living in the Amboseli ecosystem in Kenya between April 2000 to September 2013. The Amboseli baboon population is primarily composed of yellow baboons (*Papio cynocephalus*) with some admixture from nearby anubis baboon (*Papio anubis*) populations (*99–101*). Prior research in our population finds no link between host hybrid ancestry and microbiome composition (*92*). Since 1971, the Amboseli Baboon Research Project (ABRP) has been collecting continuous observations of the baboons' demography, behavior, and environment (*73*). The baboons are individually identified by expert observers who visit and collect data on each social group 3 to 4 times per week (the subjects lived in up to 12 different social groups over the study period). During each monitoring visit, the

469 observers conduct group censuses and record all demographic events, including births, maturation
470 events, and deaths, allowing us to calculate age at maturity and lifespan with precision. This
471 research was approved by the IACUC at Duke University, University of Notre Dame, and Princeton
472 University and the Ethics Council of the Max Planck Society and adhered to all the laws and
473 guidelines of Kenya.

Sample collection, DNA extraction, and 16S data generation

The 13,476 gut microbiome compositional profiles in this analysis represent a subset of
17,277 profiles, which were previously described in (*50*, *51*). The 13,476 samples in our analyses
include those from baboons whose birthdates, and hence individual ages, were known with just a
few days' error. Each baboon had on average 33 samples collected across 6 years of their life
(**Figure 1A and 1B**; range=3-135 samples per baboon; median days between samples=44 days).

Samples were collected from the ground within 15 minutes of defecation. For each sample,
approximately 20 g of feces was collected into a paper cup, homogenized by stirring with a wooden
tongue depressor, and a 5 g aliquot of the homogenized sample was transferred to a tube containing
95% ethanol. While a small amount of soil was typically present on the outside of the fecal sample,
mammalian feces contains 1000 times the number of microbial cells in a typical soil sample (*102*,
103), which overwhelms the signal of soil bacteria in our analyses (*92*). Samples were transported
from the field in Amboseli to a lab in Nairobi, freeze-dried, and then sifted to remove plant matter
prior to long term storage at -80°C.

DNA from 0.05 g of fecal powder was manually extracted using the MoBio (Catalog
No. 12955-12) and QIAGEN (Catalog No. 12955-4) PowerSoil HTP kits for 96-well plates using
a modified version of the MoBio PowerSoil-HTP kit. Specifically, we followed the manufacturers'
instructions but increased the amount of PowerBead solution to 950 μL/well and incubated the
plates at 60°C for 10 minutes after the addition of PowerBead solution and lysis buffer C1. We
included one extraction blank per batch, which had significantly lower DNA concentrations than
sample wells (t-test; t=-50, $p < 2.2 \times 10^{-16}$; *50*). We also included technical replicates, which were
the same fecal sample sequenced across multiple extraction and library preparation batches.
Technical replicates from different batches clustered with each other rather than with their batch,
indicating that true biological differences between samples are larger than batch effects.

Following DNA extraction, a ~390 bp segment of the V4 region of the 16S rRNA gene was
amplified and libraries prepared following standard protocols from the Earth Microbiome Project
(*104*). Libraries were sequenced on the Illumina HiSeq 2500 using the Rapid Run mode (2 lanes
per run). Sequences were single indexed on the forward primer and 12 bp Golay barcoded. The
resulting sequencing reads were processed following a DADA2 pipeline (*105*), with the following
additional quality filters: we removed samples with low DNA extraction concentrations (< 4x the
plate's blank DNA extraction concentration), samples with <1,000 reads, and amplicon sequence
variants that appeared in one sample (see (*50*) for details). ASVs were assigned to microbial taxa
using the IdTaxa(…) function in the DECIPHER package, against the Silva reference database
SILVA_SSU_r132_March2018.RData (*106*, *107*). The final set of samples had 1,017 to 427,454
reads (median =51,839 reads), with 8,492 total ASVs.

Identifying microbiome features that contribute to age predictions and that change with age

To identify microbiome features that change with host age, we ran linear mixed models on
1,440 microbiome features (**Table S1**). Models were run using the R package *lme4*, with p-value
estimates from *lmerTest* (*108*, *109*). These features included: (i) five metrics of alpha diversity; (ii)

the top 10 principal components of microbiome compositional variation; (iii) centered log ratio
(CLR) transformed abundances (i.e., read counts) of each microbial phyla (n=30), family (n=290),
genus (n=747), and amplicon sequence variance (ASV) detected in >25% of samples (n=358). CLR
transformations are a recommended approach for addressing the compositional nature of 16S rRNA
amplicon read count data (84). Alpha diversity metrics were calculated using the R package *vegan*
and principal components of microbiome compositional variation were calculated using the R
package *labdsv* (*110*, *111*).

For each feature, we modeled its relationship to host chronological age using both linear
and quadratic terms. To make our quadratic terms more interpretable, we centered our age estimates
on zero by subtracting the average age in the dataset from each age value. Specifically, when a
quadratic term is negative, the curve is concave, whereas when the term is positive, the curve is
convex. We also included season (wet or dry) and z-scored read count, rainfall, and temperature as
fixed effects, and individual identity, social group at time of collection, hydrological year, and the
DNA extraction/PCR plate identity were modeled as random effects. We did not model individual
social network position because prior analyses of this data set find no evidence that close social
partners have more similar gut microbiomes, probably because we lack samples from close social
partners sampled close in time (*50*, *51*). All community features (i.e., alpha diversity and principal
components) and all taxa present in >25% of samples were modeled using a Gaussian error
distribution. We extracted the coefficient, standard error, and p-value for the age term, then
corrected for multiple testing using the false discovery rate approach of Benjamini and Hochberg
(*112*).

Building the gut microbiome clock

We created a microbiome clock by fitting a Gaussian process (GP) regression model (with
a kernel customized to account for heteroskedasticity) to predict each baboon's chronological age
at the time of sample collection using 9,575 microbiome compositional and taxonomic features
present in at least 3 samples (**Table S2**; i.e., we did not restrict the features in the clock to the 1,440
most abundant features used in the age-association analyses described above). The GP regression
model with heteroskedasticity correction was the best performing of four supervised machine
learning approaches we considered, including elastic net, random forest, and Gaussian process
regression with and without the heteroskedasticity kernel (**Figure S3**; **Table S11**; See Supplemental
Methods for a comparison of other algorithms). Pearson's correlations between age predictions
across the four methods ranged from 0.69 between the random forests and the GP regression model
without the heteroskedasticity kernel to 0.96 between the two GP regression models (with and
without the heteroskedasticity kernel; **Table S11**).

Gaussian process regressions were conducted in Python 3 using scikit-learn (*113*, *114*). As
a nonparametric, Bayesian approach that infers a probability distribution over all the potential
functions that fit the data, the Gaussian process regression does not assume a linear relationship
between chronological and predicted age (*115*). For the prior distribution in the Gaussian process
regression, we used a radial basis function as our kernel and set the scale parameter to the mean
Euclidean distance of the dataset, as calculated in the R package vegan (*110*). Because initial,
exploratory models exhibited heteroskedasticity (**Figure S7**), we multiplied the variance in the
training data by the radial basis function, which distributed the higher variance in later life more
evenly across lifespan.

To calculate a microbial age estimate for every sample, and to estimate generalization error,
we used nested five-fold cross validation. In each of the five model runs, we used 80% of the data
to train the model, and the remaining 20% of the dataset as the test data. Because host identity has
a strong effect on microbiome composition in our population (*51*), we distributed samples from

each host across the five test/training data sets by randomly assigning each sample a test set without replacement. Hence, the training data set was not naïve to information from the given host being predicted as some samples for that host were included in the training set. For each model run, four of the test datasets were treated altogether as training data and the 5th set was the validation test set. We then took the estimates from all five model runs and estimated global model accuracy on the aggregated estimates.

We assessed the accuracy of our microbiome clock by regressing each sample's chronological age (age_c) against the model's predicted microbial age (age_m) and determining the R^2 value and Pearson's correlation between age_c and age_m. We also calculated the median error of the model fit as the median absolute difference between age_c and age_m across all samples (*40*).

Calculating microbiome Δage estimates

To characterize patterns of microbiome age from our microbiome clock, we calculated sample-specific microbiome Δage in years as the difference between a sample's microbial age estimate, age_m from the microbiome clock, and the host's chronological age in years at the time of sample collection, age_c. Higher microbiome Δages indicate old-for-age microbiomes, as $age_m >$ age_c, and lower values (which are often negative) indicate a young-for-age microbiome, where age_c $> age_m$ (see **Figure 3**). Because the microbiome clock systematically over predicted the ages of young animals and under predicted the ages of old animals), we also calculated a "corrected microbiome Δage" as the residuals of age_m correcting for host chronological age, season, monthly temperature, monthly rainfall, and social group and hydrological year at the time of collection. This measure is used for visualizations of the predictors of microbiome age and for testing whether average microbiome Δage predicts developmental milestones or survival.

Testing sources of variation in microbiome Δage

Many social and environmental factors have been shown to predict fertility and survival in the Amboseli baboons (*54, 59, 67, 68, 75*). To test if some of the most important known factors also predict patterns of microbiome age, we used linear mixed models to test predictors of microbiome age in individual samples separately for males and females.

In these models, the response variable was the sample-specific measure of Δage (age_m - age_c). All models included the following fixed effects: individual chronological age at the time of sample collection, to correct for model compression; the average maximum temperature during the 30 days before the sample was collected, total rainfall during the 30 days before the sample was collected, and the season (wet or dry) during sample collection. Every model also included, as fixed effects, measures of early-life adversity the individual experienced prior to 4 years of age (**Table S7**). These were modeled as either six, individual, binary variables, reflecting the presence or absence of each source of adversity in the first four years of life, or as a cumulative sum of the number of sources of adversity the individual experienced, also in the first four years of life (*67*). Social rank at the time of sampling was also modeled as a fixed effect. For males we used ordinal rank, and for females we used proportional rank (*116*). To make model interpretation more intuitive (high rank corresponds to higher values), we multiplied the coefficients for ordinal rank and maternal rank by -1. Random effects included individual identity, the social group the individual lived in at the time of collection, and hydrological year. In models of microbiome age in females, the number of adult females in the group at the time of sample collection was included as female-specific measure of resource competition.

Testing whether microbiome Δage predicts baboon maturation and survival

We used Cox proportional hazards models to test whether microbiome Δage predicted the age at which females and males attained maturational milestones and the age at death for juveniles and adult females (**Table S8**). We only measured adult survival in females because males disperse between social groups, often repeatedly across adulthood, making it is difficult to know if male disappearances are due to dispersal or death (*90*). For females, the maturational milestones of interest were the age at adult rank attainment (median age 2.24 in Amboseli), age at menarche (median age 4.51 in Amboseli), and the age at first live birth (median age 5.97 in Amboseli). For males, these milestones were the age of testicular enlargement (median age 5.38 in Amboseli), the age of dispersal from natal group (median age 7.47 in Amboseli), and the age of first adult rank attainment (i.e., when a male first outranks another adult male in his group's dominance hierarchy; median age 7.38 in Amboseli) (*55, 56*). See full descriptions of each milestone in **Table S8**. To be included in these analyses, animals must have reached the milestone after the onset of sampling (April 2000) and had at least three samples available in the timeframe of interest. We verified that none of our models violated the proportional hazards assumption of a Cox regression.

The variables we modeled differed based on the event of interest. However, all models included as fixed effects corrected Δage as the residuals of gut microbiome Δage averaged over the timeframe. All models of developmental milestones also included variables tested in Charpentier et al. 2008 (*55, 56*): (iii) maternal presence at the time of the milestone, (iv) the number of maternal sisters in the social group, averaged over the timeframe, (v) rainfall averaged over the timeframe, and (vi) whether the subject's mother was low ranked (was in the lowest quartile for female ordinal rank). For female-specific milestones, we also included (vii) the average number of adult females in the group averaged over the timeframe, and for male-specific milestones we included the number of excess cycling females in the group averaged over the timeframe, or the difference between the number of cycling females and the number of mature males within a subject's social group. Last, we included (viii) the subject's hybrid score, which is an estimation of the proportion of an individual's genetic ancestry attributable to anubis or yellow baboon ancestry (*101*).

All juvenile survival models included as fixed effects the residuals of microbiome Δage averaged over the timeframe and measures the cumulative number of sources of early-life adversity each individual experienced (*67*). Additionally, we ran three versions of the juvenile survival analysis: two subset to each sex, and one version that included both sexes. In the model including both sexes, we included sex as a predictor.

Adult female survival models included the same variables as for juvenile survival, but additionally included average lifetime dyadic social connectedness to adult females, average lifetime dyadic social connectedness to adult males, and average lifetime proportional rank. Full descriptions of all predictors are available in **Table S12**.

Acknowledgments: We thank Jeanne Altmann for her essential role in stewarding the Amboseli Baboon Project, and in collecting many of the fecal samples used in this manuscript. In Kenya, we thank the Kenya Wildlife Service, the Wildlife Training and Research Institute, the National Council for Science, Technology, and Innovation, and the National Environment Management Authority for permission to conduct research and collect biological samples. We also thank the University of Nairobi, Institute of Primary Research, National Museums of Kenya, the Amboseli-Longido pastoralist communities, the Enduimet Wildlife Management Area, Ker & Downey Safaris, Air Kenya, and Safarilink for their cooperation and assistance in the field. We thank Karl Pinc for managing and designing the database. We thank Raphael Mututua, Serah Sayialel,

Kinyua Warutere, and Long'ida Siodi for collecting the field data and fecal samples. We also thank Tawni Voyles, Anne Dumaine, Yingying Zhang, Meghana Rao, Tauras Vilgalys, Amanda Lea, Noah Snyder-Mackler, Paul Durst, Jay Zussman, Garrett Chavez, and Reena Debray for contributing to fecal sample processing. Complete acknowledgments for the ABRP can be found online at https://amboselibaboons.nd.edu/acknowledgements/.

Funding: This work was supported by the National Institutes of Health and National Science Foundation, especially the National Institute on Aging for R01 AG071684 (EAA), R21 AG055777 (EAA, RB), NIH R01 AG053330 (EAA), NIH R35 GM128716 (RB), NSF DBI 2109624 (MRD), and NSF DEB 1840223 (EAA, JAG). We also thank the Duke University Population Research Institute P2C-HD065563 (pilot to JT), the University of Notre Dame's Eck Institute for Global Health (EAA), and the Notre Dame Environmental Change Initiative (EAA). We also thank Duke University, Princeton University, the University of Notre Dame, the Chicago Zoological Society, the Max Planck Institute for Demographic Research, the L.S.B. Leakey Foundation and the National Geographic Society for support at various times over the years.

Author contributions:
Conceptualization: MRD, JT, EAA
Methodology: MRD, KER, DJ, JA, SM, JT, EAA
Supervision: SCA, LBB, JAG, RB, SM, JT, EAA
Writing—original draft: MRD, EAA
Writing—review & editing: MRD, KER, DJ, JA, SCA, LBB, JAG, RB, SM, JT, EAA

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: All data for these analyses are available on Dryad at https://doi.org/10.5061/dryad.b2rbnzspv. The 16S rRNA gene sequencing data are deposited on EBI-ENA (project ERP119849) and Qiita (study 12949). Code is available at the following GitHub repository: https://github.com/maunadasari/Dasari_etal-GutMicrobiomeAge

References

1. M. Komanduri, S. Gondalia, A. Scholey, C. Stough, The microbiome and cognitive aging: a review of mechanisms. *Psychopharmacology*, doi: 10.1007/s00213-019-05231-1 (2019).

2. C. López-Otín, M. A. Blasco, L. Partridge, M. Serrano, G. Kroemer, The Hallmarks of Aging. *Cell* **153**, 1194–1217 (2013).

3. E. Nakamura, K. Miyao, A method for identifying biomarkers of aging and constructing an index of biological age in humans. *J. Gerontol. A Biol. Sci. Med. Sci.* **62**, 1096–1105 (2007).

4. D. Gems, L. Partridge, Genetics of Longevity in Model Organisms: Debates and Paradigm Shifts. *Annual Review of Physiology* **75**, 621–644 (2013).

5. D. W. Belsky, A. Caspi, R. Houts, H. J. Cohen, D. L. Corcoran, A. Danese, H. Harrington, S. Israel, M. E. Levine, J. D. Schaefer, K. Sugden, B. Williams, A. I. Yashin, R. Poulton, T. E. Moffitt, Quantification of biological aging in young adults. *Proc. Natl. Acad. Sci. U.S.A.* **112**, E4104-4110 (2015).

6. A. D. Hayward, J. Moorad, C. E. Regan, C. Berenos, J. G. Pilkington, J. M. Pemberton, D. H. Nussey, Asynchrony of senescence among phenotypic traits in a wild mammal population. *Experimental Gerontology* **71**, 56–68 (2015).

7.   M. J. Claesson, I. B. Jeffery, S. Conde, S. E. Power, E. M. O'Connor, S. Cusack, H. M. B. Harris, M. Coakley,
B. Lakshminarayanan, O. O'Sullivan, G. F. Fitzgerald, J. Deane, M. O'Connor, N. Harnedy, K. O'Connor, D.
O'Mahony, D. van Sinderen, M. Wallace, L. Brennan, C. Stanton, J. R. Marchesi, A. P. Fitzgerald, F.
Shanahan, C. Hill, R. P. Ross, P. W. O'Toole, Gut microbiota composition correlates with diet and health in
the elderly. *Nature* **488**, 178–184 (2012).

8.   C. Heintz, W. Mair, You Are What You Host: Microbiome Modulation of the Aging Process. *Cell* **156**, 408–
411 (2014).

9.   P. W. O'Toole, I. B. Jeffery, Gut microbiota and aging. *Science* **350**, 1214–1215 (2015).

10.   B. Sadoughi, D. Schneider, R. Daniel, O. Schülke, J. Ostner, Aging gut microbiota of wild macaques are
equally diverse, less stable, but progressively personalized. *Microbiome* **10**, 95 (2022).

11.   F. Bäckhed, R. E. Ley, J. L. Sonnenburg, D. A. Peterson, J. I. Gordon, Host-bacterial mutualism in the human
intestine. *Science* **307**, 1915–1920 (2005).

12.   S. Mueller, K. Saunier, C. Hanisch, E. Norin, L. Alm, T. Midtvedt, A. Cresci, S. Silvi, C. Orpianesi, M. C.
Verdenelli, T. Clavel, C. Koebnick, H.-J. F. Zunft, J. Doré, M. Blaut, Differences in Fecal Microbiota in
Different European Study Populations in Relation to Age, Gender, and Country: a Cross-Sectional Study.
Applied and Environmental Microbiology **72**, 1027–1033 (2006).

13.   J. E. Koenig, A. Spor, N. Scalfone, A. D. Fricker, J. Stombaugh, R. Knight, L. T. Angenent, R. E. Ley,
Succession of microbial consortia in the developing infant gut microbiome. *Proceedings of the National
Academy of Sciences* **108**, 4578–4585 (2011).

14.   T. Yatsunenko, F. E. Rey, M. J. Manary, I. Trehan, M. G. Dominguez-Bello, M. Contreras, M. Magris, G.
Hidalgo, R. N. Baldassano, A. P. Anokhin, A. C. Heath, B. Warner, J. Reeder, J. Kuczynski, J. G. Caporaso, C.
A. Lozupone, C. Lauber, J. C. Clemente, D. Knights, R. Knight, J. I. Gordon, Human gut microbiome viewed
across age and geography. *Nature* **486**, 222–227 (2012).

15.   A. Bergström, T. H. Skov, M. I. Bahl, H. M. Roager, L. B. Christensen, K. T. Ejlerskov, C. Mølgaard, K. F.
Michaelsen, T. R. Licht, Establishment of Intestinal Microbiota during Early Life: a Longitudinal, Explorative
Study of a Large Cohort of Danish Infants. *Applied and Environmental Microbiology* **80**, 2889–2900 (2014).

16.   M. G. I. Langille, C. J. Meehan, J. E. Koenig, A. S. Dhanani, R. A. Rose, S. E. Howlett, R. G. Beiko, Microbial
shifts in the aging mouse gut. *Microbiome* **2**, 1–12 (2014).

17.   R. I. Clark, A. Salazar, R. Yamada, S. Fitz-Gibbon, M. Morselli, J. Alcaraz, A. Rana, M. Rera, M. Pellegrini,
W. W. Ja, D. W. Walker, Distinct Shifts in Microbiota Composition during Drosophila Aging Impair Intestinal
Function and Drive Mortality. *Cell Reports* **12**, 1656–1667 (2015).

18.   X. Cong, W. Xu, S. Janton, W. A. Henderson, A. Matson, J. M. McGrath, K. Maas, J. Graf, Gut Microbiome
Developmental Patterns in Early Life of Preterm Infants: Impacts of Feeding and Gender. *PLoS One* **11**,
e0152751 (2016).

19.   M. Yassour, T. Vatanen, H. Siljander, A.-M. Hämäläinen, T. Härkönen, S. J. Ryhänen, E. A. Franzosa, H.
Vlamakis, C. Huttenhower, D. Gevers, E. S. Lander, M. Knip, R. J. Xavier, Natural history of the infant gut
microbiome and impact of antibiotic treatment on bacterial strain diversity and stability. *Science Translational
Medicine* **8**, 343ra81-343ra81 (2016).

20.   E. Biagi, C. Franceschi, S. Rampelli, M. Severgnini, R. Ostan, S. Turroni, C. Consolandi, S. Quercia, M.
Scurti, D. Monti, M. Capri, P. Brigidi, M. Candela, Gut Microbiota and Extreme Longevity. *Current Biology*
26, 1480–1485 (2016).

21.   T. Odamaki, K. Kato, H. Sugahara, N. Hashikura, S. Takahashi, J. Xiao, F. Abe, R. Osawa, Age-related
changes in gut microbiota composition from newborn to centenarian: a cross-sectional study. *BMC
Microbiology* **16**, 1–12 (2016).

22.   P. Smith, D. Willemsen, M. Popkes, F. Metge, E. Gandiwa, M. Reichard, D. R. Valenzano, Regulation of life
span by the gut microbiota in the short-lived African turquoise killifish. *eLife* **6**, e27014 (2017).

23.   A. T. Reese, S. R. Phillips, L. A. Owens, E. M. Venable, K. E. Langergraber, Z. P. Machanda, J. C. Mitani, M.
N. Muller, D. P. Watts, R. W. Wrangham, T. L. Goldberg, M. Emery Thompson, R. N. Carmody, Age
Patterning in Wild Chimpanzee Gut Microbiota Diversity Reveals Differences from Humans in Early Life.
Current Biology, doi: 10.1016/j.cub.2020.10.075 (2020).

24.   A. Baniel, L. Petrullo, A. Mercer, L. Reitsema, S. Sams, J. Beehner, T. Bergman, N. Snyder-Mackler, A. Lu,
Maternal effects on early-life gut microbiome maturation in a wild nonhuman primat. [Preprint] (2021).
https://doi.org/10.1101/2021.11.06.467515.

25.   F. Bäckhed, C. M. Fraser, Y. Ringel, M. E. Sanders, R. B. Sartor, P. M. Sherman, J. Versalovic, V. Young, B.
B. Finlay, Defining a Healthy Human Gut Microbiome: Current Concepts, Future Directions, and Clinical
Applications. *Cell Host & Microbe* **12**, 611–622 (2012).

26.   K. R. Foster, J. Schluter, K. Z. Coyte, S. Rakoff-Nahoum, The evolution of the host microbiome as an
ecosystem on a leash. *Nature* **548**, 43–51 (2017).

27.   J. B. Clayton, A. Gomez, K. Amato, D. Knights, D. A. Travis, R. Blekhman, R. Knight, S. Leigh, R. Stumpf,
T. Wolf, K. E. Glander, F. Cabana, T. J. Johnson, The gut microbiome of nonhuman primates: Lessons in
ecology and evolution. *American Journal of Primatology* **80**, e22867 (2018).

28.   A. M. Martin, E. W. Sun, D. J. Keating, Mechanisms controlling hormone secretion in human gut and its
relevance to metabolism. *Journal of Endocrinology* **244**, R1–R15 (2020).

29.   S. Bengmark, Ecological control of the gastrointestinal tract. The role of probiotic flora. *Gut* **42**, 2–7 (1998).

30.   M. J. Claesson, S. Cusack, O. O'Sullivan, R. Greene-Diniz, H. de Weerd, E. Flannery, J. R. Marchesi, D.
Falush, T. Dinan, G. Fitzgerald, C. Stanton, D. van Sinderen, M. O'Connor, N. Harnedy, K. O'Connor, C.
Henry, D. O'Mahony, A. P. Fitzgerald, F. Shanahan, C. Twomey, C. Hill, R. P. Ross, P. W. O'Toole,
Composition, variability, and temporal stability of the intestinal microbiota of the elderly. *Proceedings of the
National Academy of Sciences* **108**, 4586–4591 (2011).

31.   G. K. Gerber, The dynamic microbiome. *FEBS Letters* **588**, 4131–4139 (2014).

32.   C. Palmer, E. M. Bik, D. B. DiGiulio, D. A. Relman, P. O. Brown, Development of the Human Infant Intestinal
Microbiota. *PLOS Biology* **5**, e177 (2007).

33.   A. Salosensaari, V. Laitinen, A. S. Havulinna, G. Meric, S. Cheng, M. Perola, L. Valsta, G. Alfthan, M.
Inouye, J. D. Watrous, T. Long, R. A. Salido, K. Sanders, C. Brennan, G. C. Humphrey, J. G. Sanders, M. Jain,
P. Jousilahti, V. Salomaa, R. Knight, L. Lahti, T. Niiranen, Taxonomic signatures of cause-specific mortality
risk in human gut microbiome. *Nat Commun* **12**, 2671 (2021).

34.   T. Wilmanski, C. Diener, N. Rappaport, S. Patwardhan, J. Wiedrick, J. Lapidus, J. C. Earls, A. Zimmer, G.
Glusman, M. Robinson, J. T. Yurkovich, D. M. Kado, J. A. Cauley, J. Zmuda, N. E. Lane, A. T. Magis, J. C.
Lovejoy, L. Hood, S. M. Gibbons, E. S. Orwoll, N. D. Price, Gut microbiome pattern reflects healthy ageing
and predicts survival in humans. *Nature Metabolism*, 1–13 (2021).

35.   M. I. Smith, T. Yatsunenko, M. J. Manary, I. Trehan, R. Mkakosya, J. Cheng, A. L. Kau, S. S. Rich, P.
Concannon, J. C. Mychaleckyj, J. Liu, E. Houpt, J. V. Li, E. Holmes, J. Nicholson, D. Knights, L. K. Ursell, R.
Knight, J. I. Gordon, Gut Microbiomes of Malawian Twin Pairs Discordant for Kwashiorkor. *Science* **339**,
548–554 (2013).

36.   S. Subramanian, S. Huq, T. Yatsunenko, R. Haque, M. Mahfuz, M. A. Alam, A. Benezra, J. DeStefano, M. F.
Meier, B. D. Muegge, M. J. Barratt, L. G. VanArendonk, Q. Zhang, M. A. Province, W. A. Petri Jr, T. Ahmed,
J. I. Gordon, Persistent gut microbiota immaturity in malnourished Bangladeshi children. *Nature* **510**, 417–421
(2014).

37. L. V. Blanton, M. R. Charbonneau, T. Salih, M. J. Barratt, S. Venkatesh, O. Ilkaveya, S. Subramanian, M. J.
Manary, I. Trehan, J. M. Jorgensen, Y. Fan, B. Henrissat, S. A. Leyn, D. A. Rodionov, A. L. Osterman, K. M.
Maleta, C. B. Newgard, P. Ashorn, K. G. Dewey, J. I. Gordon, Gut bacteria that prevent growth impairments
transmitted by microbiota from malnourished children. *Science* **351** (2016).

38. J. L. Gehrig, S. Venkatesh, H.-W. Chang, M. C. Hibberd, V. L. Kung, J. Cheng, R. Y. Chen, S. Subramanian,
C. A. Cowardin, M. F. Meier, D. O'Donnell, M. Talcott, L. D. Spears, C. F. Semenkovich, B. Henrissat, R. J.
Giannone, R. L. Hettich, O. Ilkayeva, M. Muehlbauer, C. B. Newgard, C. Sawyer, R. D. Head, D. A.
Rodionov, A. A. Arzamasov, S. A. Leyn, A. L. Osterman, M. I. Hossain, M. Islam, N. Choudhury, S. A.
Sarker, S. Huq, I. Mahmud, I. Mostafa, M. Mahfuz, M. J. Barratt, T. Ahmed, J. I. Gordon, Effects of
microbiota-directed foods in gnotobiotic animals and undernourished children. *Science* **365** (2019).

39. X. Tian, A. Seluanov, V. Gorbunova, Molecular mechanisms determining lifespan in short- and long-lived
species. *Trends Endocrinol Metab* **28**, 722–734 (2017).

40. S. Horvath, DNA methylation age of human tissues and cell types. *Genome Biol.* **14**, R115 (2013).

41. R. E. Marioni, S. Shah, A. F. McRae, B. H. Chen, E. Colicino, S. E. Harris, J. Gibson, A. K. Henders, P.
Redmond, S. R. Cox, A. Pattie, J. Corley, L. Murphy, N. G. Martin, G. W. Montgomery, A. P. Feinberg, M. D.
Fallin, M. L. Multhaup, A. E. Jaffe, R. Joehanes, J. Schwartz, A. C. Just, K. L. Lunetta, J. M. Murabito, J. M.
Starr, S. Horvath, A. A. Baccarelli, D. Levy, P. M. Visscher, N. R. Wray, I. J. Deary, DNA methylation age of
blood predicts all-cause mortality in later life. *Genome Biology* **16**, 25 (2015).

42. B. H. Chen, R. E. Marioni, E. Colicino, M. J. Peters, C. K. Ward-Caviness, P.-C. Tsai, N. S. Roetker, A. C.
Just, E. W. Demerath, W. Guan, J. Bressler, M. Fornage, S. Studenski, A. R. Vandiver, A. Z. Moore, T.
Tanaka, D. P. Kiel, L. Liang, P. Vokonas, J. Schwartz, K. L. Lunetta, J. M. Murabito, S. Bandinelli, D. G.
Hernandez, D. Melzer, M. Nalls, L. C. Pilling, T. R. Price, A. B. Singleton, C. Gieger, R. Holle, A.
Kretschmer, F. Kronenberg, S. Kunze, J. Linseisen, C. Meisinger, W. Rathmann, M. Waldenberger, P. M.
Visscher, S. Shah, N. R. Wray, A. F. McRae, O. H. Franco, A. Hofman, A. G. Uitterlinden, D. Absher, T.
Assimes, M. E. Levine, A. T. Lu, P. S. Tsao, L. Hou, J. E. Manson, C. L. Carty, A. Z. LaCroix, A. P. Reiner, T.
D. Spector, A. P. Feinberg, D. Levy, A. Baccarelli, J. van Meurs, J. T. Bell, A. Peters, I. J. Deary, J. S. Pankow,
L. Ferrucci, S. Horvath, DNA methylation-based measures of biological age: meta-analysis predicting time to
death. *Aging (Albany NY)* **8**, 1844–1859 (2016).

43. A. M. Binder, C. Corvalan, V. Mericq, A. Pereira, J. L. Santos, S. Horvath, J. Shepherd, K. B. Michels, Faster
ticking rate of the epigenetic clock is associated with faster pubertal development in girls. *Epigenetics* **13**, 85–
94 (2018).

44. K. Declerck, W. Vanden Berghe, Back to the future: Epigenetic clock plasticity towards healthy aging.
Mechanisms of Ageing and Development **174**, 18–29 (2018).

45. J. A. Anderson, R. A. Johnston, A. J. Lea, F. A. Campos, T. N. Voyles, M. Y. Akinyi, S. C. Alberts, E. A.
Archie, J. Tung, High social status males experience accelerated epigenetic aging in wild baboons. *eLife* **10**,
e66128 (2021).

46. J. de la Cuesta-Zuluaga, S. T. Kelley, Y. Chen, J. S. Escobar, N. T. Mueller, R. E. Ley, D. McDonald, S.
Huang, A. D. Swafford, R. Knight, V. G. Thackray, Age- and Sex-Dependent Patterns of Gut Microbial
Diversity in Human Adults. *mSystems* **4** (2019).

47. F. Galkin, P. Mamoshina, A. Aliper, E. Putin, V. Moskalev, V. N. Gladyshev, A. Zhavoronkov, Human Gut
Microbiome Aging Clock Based on Taxonomic Profiling and Deep Learning. *iScience* **23**, 101199 (2020).

48. S. Huang, N. Haiminen, A.-P. Carrieri, R. Hu, L. Jiang, L. Parida, B. Russell, C. Allaband, A. Zarrinpar, Y.
Vázquez-Baeza, P. Belda-Ferre, H. Zhou, H.-C. Kim, A. D. Swafford, R. Knight, Z. Z. Xu, Human Skin, Oral,
and Gut Microbiomes Predict Chronological Age. *mSystems* **5** (2020).

49. Y. Chen, H. Wang, W. Lu, T. Wu, W. Yuan, J. Zhu, Y. K. Lee, J. Zhao, H. Zhang, W. Chen, Human gut
microbiome aging clocks based on taxonomic and functional signatures through multi-view learning. *Gut*
Microbes **14**, 2025016 (2022).

50. L. E. Grieneisen, M. Dasari, T. J. Gould, J. R. Björk, J.-C. Grenier, V. Yotova, D. Jansen, N. Gottel, J. B.
Gordon, N. H. Learn, L. R. Gesquiere, T. L. Wango, R. S. Mututua, J. K. Warutere, L. Siodi, J. A. Gilbert, L.
B. Barreiro, S. C. Alberts, J. Tung, E. A. Archie, R. Blekhman, Gut microbiome heritability is nearly universal
but environmentally contingent. *Science* **373**, 181–186 (2021).

51. J. R. Björk, M. R. Dasari, K. Roche, L. Grieneisen, T. J. Gould, J.-C. Grenier, V. Yotova, N. Gottel, D. Jansen,
L. R. Gesquiere, J. B. Gordon, N. H. Learn, T. L. Wango, R. S. Mututua, J. Kinyua Warutere, L. Siodi, S.
Mukherjee, L. B. Barreiro, S. C. Alberts, J. A. Gilbert, J. Tung, R. Blekhman, E. A. Archie, Synchrony and
idiosyncrasy in the gut microbiome of wild baboons. *Nat Ecol Evol*, 1–10 (2022).

52. K. E. Roche, J. R. Bjork, M. R. Dasari, L. Grieneisen, D. A. Jansen, T. J. Gould, L. R. Gesquiere, L. B.
Barreiro, S. C. Alberts, R. Blekhman, J. A. Gilbert, J. Tung, S. Mukherjee, E. A. Archie, Universal gut
microbial relationships in the gut microbiome of wild baboons. *eLife* **12**, e83152 (2023).

53. S. C. Alberts, J. Altmann, Preparation and activation: determinants of age at reproductive maturity in male
baboons. *Behavioral Ecology and Sociobiology* **36**, 397–406 (1995).

54. J. Altmann, L. Gesquiere, J. Galbany, P. O. Onyango, S. C. Alberts, Life history context of reproductive aging
in a wild primate model. *Annals of the New York Academy of Sciences* **1204**, 127–138 (2010).

55. M. J. E. Charpentier, J. Tung, J. Altmann, S. C. Alberts, Age at maturity in wild baboons: genetic,
environmental and demographic influences. *Molecular Ecology* **17**, 2026–2040 (2008).

56. P. O. Onyango, L. R. Gesquiere, J. Altmann, S. C. Alberts, Puberty and dispersal in a wild primate population.
Hormones and Behavior **64**, 240–249 (2013).

57. S. C. Alberts, E. A. Archie, L. R. Gesquiere, J. Altmann, J. W. Vaupel, K. Christensen, "The Male-Female
Health-Survival Paradox: A Comparative Perspective on Sex Differences in Aging and Mortality" in *Advances
in Biodemography: Cross-Species Comparisons of Social Environments and Social Behaviors, and Their
Effects on Health and Longevity* (The National Academies Press, Washington D.C., 2014), pp. 339–363.

58. E. A. Archie, J. Altmann, S. C. Alberts, Costs of reproduction in a long-lived female primate: injury risk and
wound healing. *Behavioral Ecology and Sociobiology* **68**, 1183–1193 (2014).

59. E. A. Archie, J. Tung, M. Clark, J. Altmann, S. C. Alberts, Social affiliation matters: both same-sex and
opposite-sex relationships predict survival in wild female baboons. *Proc. R. Soc. B* **281**, 20141261 (2014).

60. L. E. Grieneisen, J. Livermore, S. Alberts, J. Tung, E. A. Archie, Group Living and Male Dispersal Predict the
Core Gut Microbiome in Wild Baboons. *Integr Comp Biol* **57**, 770–785 (2017).

61. T. Ren, L. E. Grieneisen, S. C. Alberts, E. A. Archie, M. Wu, Development, diet and dynamism: longitudinal
and cross-sectional predictors of gut microbial communities in wild baboons. *Environmental Microbiology*,
1312–1325 (2015).

62. J. Tung, L. B. Barreiro, M. B. Burns, J. C. Grenier, J. Lynch, L. E. Grieneisen, J. Altmann, S. C. Alberts, R.
Blekhman, E. A. Archie, Social networks predict gut microbiome composition in wild baboons. *Elife* **4** (2015).

63. A. M. Bronikowski, J. Altmann, D. K. Brockman, M. Cords, L. M. Fedigan, A. Pusey, T. Stoinski, W. F.
Morris, K. B. Strier, S. C. Alberts, Aging in the Natural World: Comparative Data Reveal Similar Mortality
Patterns Across Primates. *Science* **331**, 1325–1328 (2011).

64. L. R. Gesquiere, N. H. Learn, M. C. Simao, P. O. Onyango, S. C. Alberts, J. Altmann, Life at the top: rank and
stress in wild male baboons. *Science* **333**, 357–360 (2011).

65. E. A. Archie, J. Altmann, S. C. Alberts, Social status predicts wound healing in wild baboons. *Proceedings of
the National Academy of Sciences* **109**, 9017–9022 (2012).

66. A. J. Lea, J. Altmann, S. C. Alberts, J. Tung, Developmental Constraints in a Wild Primate. *The American
Naturalist* **185**, 809–821 (2015).

67. J. Tung, E. A. Archie, J. Altmann, S. C. Alberts, Cumulative early life adversity predicts longevity in wild
baboons. *Nature Communications* **7**, 11181 (2016).

68. L. R. Gesquiere, J. Altmann, E. A. Archie, S. C. Alberts, Interbirth intervals in wild baboons: Environmental
predictors and hormonal correlates. *American Journal of Physical Anthropology* **166**, 107–126 (2018).

69. M. N. Zipple, E. A. Archie, J. Tung, J. Altmann, S. C. Alberts, Intergenerational effects of early adversity on
survival in wild baboons. *eLife* **8**, e47433 (2019).

70. T. Jovanovic, L. A. Vance, D. Cross, A. K. Knight, V. Kilaru, V. Michopoulos, T. Klengel, A. K. Smith,
Exposure to Violence Accelerates Epigenetic Aging in Children. *Sci Rep* **7**, 8962 (2017).

71. A. S. Zannas, J. Arloth, T. Carrillo-Roa, S. Iurato, S. Röh, K. J. Ressler, C. B. Nemeroff, A. K. Smith, B.
Bradley, C. Heim, A. Menke, J. F. Lange, T. Brückl, M. Ising, N. R. Wray, A. Erhardt, E. B. Binder, D. Mehta,
Lifetime stress accelerates epigenetic aging in an urban, African American cohort: relevance of glucocorticoid
signaling. *Genome Biology* **16**, 266 (2015).

72. L. Raffington, D. W. Belsky, M. Malanchini, E. M. Tucker-Drob, K. P. Harden, Analysis of socioeconomic
disadvantage and pace of aging measured in saliva DNA methylation of children and adolescents. bioRxiv
[Preprint] (2020). https://doi.org/10.1101/2020.06.04.134502.

73. S. C. Alberts, J. Altmann, "The Amboseli Baboon Research Project: Themes of continuity and change" in
Long-Term Field Studies of Primates, P. Kappeler, W. DP, Eds. (Springer Verlag, 2012), pp. 261–288.

74. D. J. Melnick, M. C. Pearl, "Cercopithecines in multimale groups: genetic diversity and population structure"
in *Primate Societies* (University of Chicago Press, Chicago, 1987).

75. J. Altmann, S. C. Alberts, Growth rates in a wild primate population: ecological influences and maternal
effects. *Behavioral Ecology and Sociobiology* **57**, 490–501 (2005).

76. J. B. Silk, S. C. Alberts, J. Altmann, Social Bonds of Female Baboons Enhance Infant Survival. *Science* **302**,
1231–1234 (2003).

77. S. C. Alberts, H. E. Watts, J. Altmann, Queuing and queue-jumping: long-term patterns of reproductive skew
in male savannah baboons, Papio cynocephalus. *Animal Behaviour* **65**, 821–840 (2003).

78. A. J. Lea, M. Y. Akinyi, R. Nyakundi, P. Mareri, F. Nyundo, T. Kariuki, S. C. Alberts, E. A. Archie, J. Tung,
Dominance rank-associated gene expression is widespread, sex-specific, and a precursor to high social status in
wild male baboons. *Proc Natl Acad Sci U S A* **115**, E12163–E12171 (2018).

79. A. L. Hicks, K. J. Lee, M. Couto-Rodriguez, J. Patel, R. Sinha, C. Guo, S. H. Olson, A. Seimon, T. A. Seimon,
A. U. Ondzie, W. B. Karesh, P. Reed, K. N. Cameron, W. I. Lipkin, B. L. Williams, Gut microbiomes of wild
great apes fluctuate seasonally in response to diet. *Nature Communications* **9**, 1786 (2018).

80. O. Kolodny, M. Weinberg, L. Reshef, L. Harten, A. Hefetz, U. Gophna, M. W. Feldman, Y. Yovel,
Coordinated change at the colony level in fruit bat fur microbiomes through time. *Nat Ecol Evol* **3**, 116–124
(2019).

81. A. Risely, K. Wilhelm, T. Clutton-Brock, M. B. Manser, S. Sommer, Diurnal oscillations in gut bacterial load
and composition eclipse seasonal and lifetime dynamics in wild meerkats. *Nat Commun* **12**, 6017 (2021).

82. D. Kuh, Y. Ben-Shlomo, J. Lynch, J. Hallqvist, C. Power, Life course epidemiology. *J Epidemiol Community
Health* **57**, 778–783 (2003).

83. L. Shanahan, W. E. Copeland, E. J. Costello, A. Angold, Child-, adolescent- and young adult-onset
depressions: differential risk factors in development? *Psychol. Med.* **41**, 2265–2274 (2011).

84. G. B. Gloor, J. M. Macklaim, V. Pawlowsky-Glahn, J. J. Egozcue, Microbiome Datasets Are Compositional:
And This Is Not Optional. *Front. Microbiol.* **8** (2017).

85.   V. D. Badal, E. D. Vaccariello, E. R. Murray, K. E. Yu, R. Knight, D. V. Jeste, T. T. Nguyen, The Gut
Microbiome, Aging, and Longevity: A Systematic Review. *Nutrients* **12**, 3759 (2020).

86.   J. G. Caporaso, C. L. Lauber, E. K. Costello, D. Berg-Lyons, A. Gonzalez, J. Stombaugh, D. Knights, P. Gajer,
J. Ravel, N. Fierer, J. I. Gordon, R. Knight, Moving pictures of the human microbiome. *Genome Biol* **12**
(2011).

87.   C. J. Weibel, M. R. Dasari, D. A. Jansen, L. R. Gesquiere, R. S. Mututua, J. K. Warutere, L. I. Siodi, S. C.
Alberts, J. Tung, E. A. Archie, Using non-invasive behavioral and physiological data to measure biological age
in wild baboons. *GeroScience* **46**, 4059–4074 (2024).

88.   L. Jayashankar, K. M. Brasky, J. A. Ward, R. Attanasio, Lymphocyte Modulation in a Baboon Model of
Immunosenescence. *Clinical and Vaccine Immunology* **10**, 870–875 (2003).

89.   J. Galbany, J. Altmann, A. Pérez-Pérez, S. C. Alberts, Age and individual foraging behavior predict tooth wear
in Amboseli baboons. *American Journal of Physical Anthropology* **144**, 51–59 (2011).

90.   F. A. Campos, F. Villavicencio, E. A. Archie, F. Colchero, S. C. Alberts, Social bonds, social status and
survival in wild baboons: a tale of two sexes. *Philosophical Transactions of the Royal Society B: Biological
Sciences* **375**, 20190621 (2020).

91.   J.-F. Lemaître, V. Ronget, M. Tidière, D. Allainé, V. Berger, A. Cohas, F. Colchero, D. A. Conde, M. Garratt,
A. Liker, G. A. B. Marais, A. Scheuerlein, T. Székely, J.-M. Gaillard, Sex differences in adult lifespan and
aging rates of mortality across wild mammals. *PNAS* **117**, 8546–8553 (2020).

92.   L. E. Grieneisen, M. J. E. Charpentier, S. C. Alberts, R. Blekhman, G. Bradburd, J. Tung, E. A. Archie, Genes,
geology and germs: gut microbiota across a primate hybrid zone are explained by site soil properties, not host
species. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20190431 (2019).

93.   J. A. Anderson, A. J. Lea, T. N. Voyles, M. Y. Akinyi, R. Nyakundi, L. Ochola, M. Omondi, F. Nyundo, Y.
Zhang, F. A. Campos, S. C. Alberts, E. A. Archie, J. Tung, Distinct gene regulatory signatures of dominance
rank and social bond strength in wild baboons. *Philosophical Transactions of the Royal Society B: Biological
Sciences* **377**, 20200441 (2022).

94.   B. Habig, E. A. Archie, Social status, immune response and parasitism in males: a meta-analysis. *Philosophical
Transactions of the Royal Society B: Biological Sciences* **370**, 20140109 (2015).

95.   N. Snyder-Mackler, J. Sanz, J. N. Kohn, J. F. Brinkworth, S. Morrow, A. O. Shaver, J.-C. Grenier, R. Pique-
Regi, Z. P. Johnson, M. E. Wilson, L. B. Barreiro, J. Tung, Social status alters immune regulation and response
to infection in macaques. *Science* **354**, 1041–1045 (2016).

96.   E. J. Levy, L. R. Gesquiere, E. McLean, M. Franz, J. K. Warutere, S. N. Sayialel, R. S. Mututua, T. L. Wango,
V. K. Oudu, J. Altmann, E. A. Archie, S. C. Alberts, Higher dominance rank is associated with lower
glucocorticoids in wild female baboons: A rank metric comparison. *Hormones and Behavior* **125**, 104826
(2020).

97.   E. N. Paietta, C. J. Weibel, D. A. Jansen, R. S. Mututua, J. K. Warutere, I. Long'ida Siodi, L. R. Gesquiere, V.
Obanda, S. C. Alberts, E. A. Archie, Troubled waters: Water availability drives human-baboon encounters in a
protected, semi-arid landscape. *Biological Conservation* **274**, 109740 (2022).

98.   E. Hanski, S. Joseph, A. Raulo, K. M. Wanelik, Á. O'Toole, S. C. L. Knowles, T. J. Little, Epigenetic age
estimation of wild mice using faecal samples. *Molecular Ecology* **33**, e17330 (2024).

99.   A. Samuels, J. Altmann, Immigration of a Papio anubis male into a group of Papio cynocephalus baboons and
evidence for an anubis-cynocephalus hybrid zone in Amboseli, Kenya. *International Journal of Primatology* **7**,
131–138 (1986).

100.   J. Tung, M. J. E. Charpentier, D. A. Garfield, J. Altmann, S. C. Alberts, Genetic evidence reveals temporal
change in hybridization patterns in a wild baboon population. *Molecular Ecology* **17**, 1998–2011 (2008).

101. T. P. Vilgalys, A. S. Fogel, J. A. Anderson, R. S. Mututua, J. K. Warutere, I. L. Siodi, S. Y. Kim, T. N. Voyles,
J. A. Robinson, J. D. Wall, E. A. Archie, S. C. Alberts, J. Tung, Selection against admixture and gene
regulatory divergence in a long-term primate field study. *Science* **377**, 635–641 (2022).

102. R. Sender, S. Fuchs, R. Milo, Revised Estimates for the Number of Human and Bacteria Cells in the Body.
PLoS Biology **14**, e1002533 (2016).

103. X. Raynaud, N. Nunan, Spatial Ecology of Bacteria at the Microscale in Soil. *PLOS ONE* **9**, e87217 (2014).

104. J. A. Gilbert, J. K. Jansson, R. Knight, The Earth Microbiome project: successes and aspirations. *BMC Biology*
12, 69 (2014).

105. B. J. Callahan, P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, S. P. Holmes, DADA2: High-
resolution sample inference from Illumina amplicon data. *Nat Meth* **13**, 581–583 (2016).

106. E. S. Wright, L. S. Yilmaz, D. R. Noguera, DECIPHER, a search-based approach to chimera identification for
16S rRNA sequences. *Appl Environ Microbiol* **78**, 717–725 (2012).

107. C. Quast, E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, F. O. Glöckner, The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res* **41**,
D590–D596 (2013).

108. D. Bates, M. Mächler, B. Bolker, S. Walker, Fitting Linear Mixed-Effects Models Using lme4. *Journal of*
Statistical Software **67**, 1–48 (2015).

109. A. Kuznetsova, P. B. Brockhoff, R. H. B. Christensen, lmerTest Package: Tests in Linear Mixed Effects
Models. *Journal of Statistical Software* **82**, 1–26 (2017).

110. P. Dixon, VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science* **14**, 927–
930 (2003).

111. D. W. Roberts, labdsv: Ordination and Multivariate Analysis for Ecology, (2019); https://CRAN.R-
project.org/package=labdsv.

112. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful Approach to
Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)* **57**, 289–300 (1995).

113. G. Van Rossum, F. L. Drake, Python 3 Reference Manual, CreateSpace (2009);
https://docs.python.org/3/reference/.

114. F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R.
Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, E. Duchesnay, Scikit-
learn: Machine Learning in Python. *Journal of Machine Learning Research* **12**, 2825–2830 (2011).

115. C. E. Rasmussen, C. K. I. Williams, *Gaussian Processes for Machine Learning (Adaptive Computation and*
Machine Learning) (The MIT Press, 2005).

116. E. J. Levy, M. N. Zipple, E. McLean, F. A. Campos, M. Dasari, A. S. Fogel, M. Franz, L. R. Gesquiere, J. B.
Gordon, L. Grieneisen, B. Habig, D. J. Jansen, N. H. Learn, C. J. Weibel, J. Altmann, S. C. Alberts, E. A.
Archie, A comparison of dominance rank metrics reveals multiple competitive landscapes in an animal society.
Proc Biol Sci **287** (2020).

1001