Endocytic dynamin deficiency impairs the host colonization of phytopathogenic fungi

Xin Chen[1,2], Xiaomin Chen[3], Yunfei Long[1,2], Zhenyu Fang[1,2], Yakubu Saddeeq Abubakar[1,5], Huawei

Zheng[4], Zonghua Wang[1,4*], Wenhui Zheng[1,2*]

[1] State Key Laboratory for Ecological Pest Control of Fujian and Taiwan Crops, College of Plant

Protection, Fujian Agriculture and Forestry University, Fuzhou, 350000 China;

[2] Key Laboratory of Bio-pesticide and Chemical Biology, Ministry of Education, College of Plant

Protection, Fujian Agriculture and Forestry University, Fuzhou, 350000 China;

[3] Fujian University Key Laboratory for Plant Microbe Interaction, College of Life Sciences, Fujian

Agriculture and Forestry University, Fuzhou 350002, China;

[4] Ministerial and Provincial Joint Innovation Centre for Safety Production of Cross-Strait Crops,

College of Geography and Oceanography, Minjiang University, Fuzhou 350108, China;

[5] Department of Biochemistry, Faculty of Life Science, Ahmadu Bello University, Zaria 810281,

Nigeria.

* Authors for correspondence:

Wenhui Zheng (wenhuiz@fafu.edu.cn), Zonghua Wang (wangzh@fafu.edu.cn).

Abstract

During host-pathogen interactions, pathogens employ endolysosomal trafficking to fine-tune host and environmental cues, facilitating disease progression. Inhibiting the functions of the endolysosomal network could be promising for the control of plant fungal diseases. Herein, we showed that inhibition of the endocytic dynamin FgVps1 by the phenothiazines chlorpromazine (CPZ) and prochlorperazine (PCZ) leads to endolysosomal dysfunction. Further analyses revealed that the phenothiazines directly bind the GTPase domain of FgVps1 to inhibit its activity. Moreover, our results showed that FgVps1 is recruited to the endosome by the Rab GTPases FgRab51 and FgRab7 to facilitate the release of retromer- and sorting nexin-coated vesicles, thereby ensuring the proper recycling of some cargos and promoting fungal development and pathogenicity. Direct treatment of the phytopathogens *F. graminearum*, *Magnaporthe oryzae* and *Fusarium oxysporum f. sp. cubense* with CPZ and PCZ also impairs their growth, development and virulence. These results unveil the underlying mechanisms of endocytic dynamin actions and provide a potential strategy for broad-spectrum control of phytopathogenic fungi.

Key words

endocytic trafficking, dynamin, Rab GTPase, sorting machinery, phenothiazines, phytopathogenic fungi

Introduction

Each year, crop losses caused by emerging fungal pathogens pose a seriously risk to global food security [1, 2]. The progression of these diseases depends on the complex interplay between environmental factors, hosts and pathogens [3, 4]. Understanding the multicellular strategies that pathogens use to maintain homeostasis and adapt to environmental conditions during interactions with hosts will enhance our knowledge of disease development and lead to more effective control strategies.

The maintenance of cellular homeostasis when exposed to biotic and abiotic stresses is spatiotemporally controlled by the endolysosomal network in eukaryotes [5]. Cell surface cargo proteins entry the endolysosomal network following endocytosis and reach the early endosomes

where cargo sorting is initiated. At the early endosome, the fates of the cargos are decided: to be

recycled or degraded [5, 6]. For recycling, cargo proteins can be directly delivered to the cell surface

or transit through the endocytic recycle compartment to be delivered to the cell surface. In addition,

some selected cargo proteins can be retrieved into the trans-Golgi by the retromer complex for

subsequent delivery to the cell surface through the secretory pathway. For degradation, cargo

proteins at the late endosome are sorted into the multivesicular body (MVB) which further fuses

with the vacuole to empty its contents for degradation. This is fine-tuned by Rab GTPases, PIPs and

ESCRT machinery [6, 7, 8].

Highlighting its importance, the endolysosomal network is associated with the pathogenesis of plant

pathogenic fungi. Dysfunction of the endolysosomal network impairs the vegetative growth and

infection-related development of various fungal pathogens [9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24,

25, 26, 27, 28, 29, 30]. Additionally, the endolysosomal network is required for effective modulation of

autophagy process, which helps in adaptation to metabolic changes and environmental cues during

pathogen-host interactions[21, 26, 27, 28, 29]. To suppress host immune responses, fungal pathogens use

the endolysosomal network to facilitate secretion of effector proteins during host infection [14].

Furthermore, the endolysosomal network is highly linked with mycotoxin production, the toxic

secondary metabolites that seriously threaten food security and animal health but are useful for host

colonization [10, 11, 12, 13, 23, 24, 25, 31]. Given the accumulating evidence showing the important roles of

the endolysosomal network in the pathogenesis of fungal phytopathogens, we hypothesized that

disrupting the endolysosomal network might have potential applications in fungal disease control.

Endocytosis, the beginning of the endolysosomal network, governs the internalization of cargo

proteins and molecules and has shown its potential in clinical treatment of various diseases [32, 33, 34,

35]. Among the various endocytosis pathways identified, clathrin-mediated endocytosis (CME) is the

major and well-characterized endocytic pathway [33, 36, 37]. Briefly, adaptor protein-2 binds a cargo

protein from one end and binds clathrin at the other end to initiate the assembly of clathrin-coated

pits (CCP). Other adaptors are then recruited to the CCP to ensure its stabilization and maturation.

After maturation, dynamins drive the scission of the clathrin-coated vesicle to allow for

internalization and subsequent fusion of the vesicle with early endosome [33, 36, 37]. In addition to

mediating the release of the clathrin-coated vesicle, dynamins also participate in membrane

remodeling events during vesicle trafficking [38, 39]. However, whether antifungal agents exert their

functions by targeting endocytic dynamins remains largely unknown.

In the present study, we evaluated the mechanistic relationship between phenothiazines (a class of

endocytosis inhibitors) and the activity of the dynamin Vps1 [34, 40]. Treatment with phenothiazines

significantly impairs the development and virulence of phytopathogenic fungi. We also found that

phenothiazines target FgVps1 and inhibit its GTPase activity via binding to its G domain region,

consequently disrupting endocytosis process in *F. graminearum*. Furthermore, FgVps1 is recruited

to the endosome by the Rab GTPases FgRab51 and FgRab7 to facilitate endocytic trafficking by

mediating the scission of retromer- and sorting nexins-coated endosomal tubules. Moreover, FgSnc1

is identified as the cargo protein in FgVps1-associated endocytic trafficking, and its intracellular

trafficking requires the GTPase activity of FgVps1. These results provide insight into the complex

interplay of the endolysosomal network in pathogenic fungi and highlight the potentials of

phenothiazines for the control of phytopathogenic fungal diseases.

Results

Phenothiazines impair the development and pathogenicity of fungal pathogens

Previously, we highlighted the important roles of endolysosomal network in the development and

pathogenicity of *F. graminearum*, the causal agent of wheat head blight [9, 10, 11, 12, 23, 24, 25, 41]. Since

the endolysosmal network begins with endocytosis of surface cargos, inhibition of endocytosis

could impair the homeostasis of the network. Phenothiazines have previously been shown to inhibit

endocytosis in mammals [5, 34, 40, 42], but their effects on phytopathogens remain unclear. To clarify

this, we first treated *F. graminearum*, *Magnaporthe oryzae* (the causal agent of rice blast) and

Fusarium oxysporum f. sp. cubense tropical race 4 (FocTR4, the causal agent of Fusarium wilt of

banana) with the phenothiazine derivatives chlorpromazine (CPZ) and prochlorperazine (PCZ) (Fig.

S1A). As shown in Fig. 1A, the growth of the indicated fungal pathogens were significant reduced

under treatments with CPZ and PCZ. Next, we examined the potential effects of these

phenothiazines on the fungal pathogenicity. Notably, when treated with CPZ and PCZ, *F.*

graminearum, FocTR4 and *M. oryzae* all displayed serious defects in their invasive growths and

virulence on wheat coleoptiles, banana and rice leaves, respectively (Fig. 1B, C and D). We further

treated the host plants with these phenothiazines to investigate whether the drugs exert any impacts

on the plant development. As shown in Fig. S1B, treatments with these inhibitors have no obvious

effects on the development of the host plants. Overall, we conclude that phenothiazine treatment

perturbs phytopathogenic fungal development and host invasion.

Phenothiazines target FgVps1 to compromise *F. graminearum* development and pathogenicity

A previous study indicated that phenothiazines target dynamin to inhibit endocytosis [40]. We

therefore used the endocytic marker dye FM4-64 to stain *F. graminearum* and monitor the impact

of the substances on endocytosis efficiency. We found that internalization of FM4-64 was

significantly reduced under phenothiazines treatment, suggesting the conserved impact of

phenothiazines on endocytosis in *F. graminearum* (Fig. 2A, Fig. S2A). We also observed that the

pathogenicity of *F. graminearum* on wheat heads was also reduced following treatments with the

endocytic inhibitors (Fig. S2B). We further conducted a phylogenetic analysis to identify the

potential dynamin target proteins in *F. graminearum*. As shown in Fig. 2B, FgVps1(FGSG_07172)

is the only member of the dynamin family of proteins that is involved in endocytic vesicle scission.

To test the role of FgVps1 in endocytosis, we generated Δ*Fgvps1* mutant and confirmed the gene

deletion by Southern blot assay (Fig. S2C). Consistent with phenothiazine treatment, loss of

FgVPS1 impaired the internalization of FM4-64 into the vacuole, indicating the importance of

FgVps1 in endocytosis efficiency in *F. graminearum* (Fig. 2C). To further explore the biological

functions of FgVps1, we generated a complemented strain Δ*Fgvps1*-com by expressing a vector

harboring GFP-*FgVPS1* into the Δ*Fgvps1* mutant (Fig. S2D). Phenotypic analyses revealed that

FgVps1 is important for hyphal and conidial morphogenesis, sexual reproduction and virulence of

the wheat head blight fungus (Fig. S2 E to G). These results demonstrate that, similar to

phenothiazines treatment, FgVps1 loss-of-function perturbs *F. graminearum* development and

pathogenicity.

To investigate the relationship between the phenothiazines treatment and FgVps1 GTPase activity,

we purified a GST-FgVps1 recombinant protein and determined its GTPase activity in the

presence/absence of phenothiazines. It was observed that the GTPase activity of FgVps1 was

significantly reduced under the phenothiazines treatment, compared to the untreated reaction system

(Fig. 2D). Next, we used molecular docking approach (AutoDock [43]) to unveil the possibility of

interaction between FgVps1 and the phenothiazines CPZ and PCZ. Interestingly, we found that

CPZ and PCZ share the same binding pocket on FgVps1, which involves the amnio acids ranging

from D239 to I247 (Fig. 2E). Structurally, FgVps1 possesses three domains: G (GTPase) domain,

DM (dynamin central region) domain and GED (dynamin GTPase effector domain) (Fig. S3A).

Through domain mutation, we found that all these domains contribute to the full functions and

subcellular distribution of FgVps1 (Fig. S3B and S3C). The phenothiazines binding pocket fall

within the G domain which is involved in guanine nucleotides binding and hydrolysis [39]. To validate

whether the binding pocket contributes to FgVps1 GTPase activity, we deleted the residues that

constitute the binding pocket (D239 to I247) and obtained FgVps1$^{\Delta POC}$ mutant expressing the

truncated protein. Next, we purified GST-FgVps1$^{\Delta POC}$ recombinant protein and analyzed its GTPase

activity using a GTPase activity assay kit. As shown in Fig. 2F, the binding pocket mutation results

in significant reduction in FgVps1 GTPase activity. In addition, the deletion of the binding pocket

also hampers the efficiency of endocytosis in *F. gramienarum* (Fig. 2G). Similar to the G domain

mutation, loss of the phenothiazines binding pocket impairs the full function and subcellular

localization of FgVps1, with the FgVps1 punctate localization detected in the cytosol (Fig. S3 and

Fig. 2H and 2I). Taken together, these results suggest that phenothiazines target the dynamin

FgVps1 to inhibit its GTPase activity, thereby compromising endocytosis and pathogenicity of *F.*

graminearum.

FgVps1 regulates the scission of retromer and sorting nexins coated vesicles

Endocytic cargos enter the endolysosomal pathway to the endosomes where their fates are

determined [5]. Since our earlier results identified FgVps1 as the target of phenothiazines (Fig. 2), we

decided to further dissect the mechanism of FgVps1-dependent endocytosis in relation to the fungal

development and pathogenicity. Therefore, we conducted GFP pull-down tandem LC-MS/MS mass

spectrometry to identify potential FgVps1-associated proteins in *F. graminearum*. As shown in Fig.

3A and Table S3, the various proteins involved in endolysososmal sorting were captured and

identified, including Rab GTPases, the retromer complex and sorting nexins. In *F. graminearum*,

the retromer complex and sorting nexins (FgSnx4-FgSnx41) both contribute to endosomal

trafficking and are required for the fungal development and pathogenicity [11, 12]; but their association

with the dynamin FgVps1 remain has not been established. Through time-lapse imaging and co-

immunoprecipitation (Co-IP) assays, we found that FgVps1 is closely associated with the retromer

and sorting nexins *in vivo* (Fig. S4A and S4B). Previously, the Vps1 was reported to mediate the

scission of endosomal tubules in yeast [44, 45]. This prompted us to hypothesize that FgVps1

contributes to the scission of retromer- and sorting nexins-associated tubules in *F. graminearum*.

To test this, we transformed FgSnx4-GFP and FgVps35-GFP (the core retromer subunit) vectors

into the wild type and Δ*Fgvps1* mutant. In the wild type, the fusion proteins displayed punctate

localization predominantly along the plasma membrane (Fig. S4C). In the Δ*Fgvps1* mutant,

however, the fluorescence signals were trapped in some tubules in the cytosol, suggesting the

involvement of FgVps1 in the scission of retromer- and sorting nexins-associated tubules.

In yeast, the retromer and Snx4-Snx41 dimer mediate distinct recycling pathways [44, 46, 47]. Also, our

previously studies showed that the retromer and Snx4-Snx41 dimer both contribute to the recycling

of the v-SNARE Snc1 in *M. oryzae* and *F. graminearum* [11, 14, 48]. Further analysis of our FgVps1,

FgVps35 and FgSnx4 pull-down data revealed that FgVps35 and FgVps1 have many potential

interacting partners in common (Fig. S4D). Thus, we hypothesized that FgVps35 and FgSnx4 may

interact to jointly execute their biological functions. To verify this hypothesis, yeast two-hybrid and

Co-IP assays were carried to check for possible interaction between FgVps35 and FgSnx4. As

shown in Fig. S4E and S4F, FgVps35 directly interacts with FgSnx4 both *in vivo* and *in vitro*.

Consistently, co-expression of FgVps35-GFP and FgSnx4-mCherry the in wild type and Δ*Fgvps1*

mutant showed close association of FgVps35 and FgSnx4, and both fusion proteins are trapped in

the same tubules in the Δ*Fgvps1* mutant (Fig. 3B). To further validate the association of FgVps35

and FgSnx4, in silico interaction model of FgVps35 and FgSnx4 was analyzed using AlphaFold

Multimer. As shown in Fig. 3C, N383 residue of FgVps35 was observed to interact with E373

residue of FgSnx4 via hydrogen bonding. To validate this prediction, we disrupted this hydrogen

bond by introducing N383A and E373A point mutations in FgVps35 and FgSnx4, respectively, and

then analyzed the mutated proteins for possible interaction by yeast two-hybrid assay. We found

that the interaction of FgVps35 with FgSnx4 was obviously weakened by N383A mutation but

completely abolished by E373A mutation (Fig. 3D). Moreover, pathogenicity tests on wheat heads

223 revealed that the FgSnx4 E373A mutation causes reduced virulence compared to the wild type (Fig.

224 3E). These results suggest that FgSnx4 E373 residue is critical for the interaction of FgSnx4 with

225 FgVps35.

227 **FgVps1 is recruited to the endosome by FgRab51 and FgRab7 to engage in endolysosomal**

228 **sorting**

229 In addition to the retromer and sorting nexins identified, the Rab GTPases FgRab51 and FgRab7

230 were also repeated captured in our FgVps1 pulldown. By co-expressing mCherry-FgVps1 with

231 GFP-FgRab51 and GFP-FgRab7, we found that FgVps1 partially co-localized with the early

232 endosomal marker FgRab51 but completely co-localized with FgRab7-marked late endosomes and

233 vacuoles. Moreover, Co-IP assays indicated that FgVps1 positively interacts with FgRab51 and

234 FgRab7 *in vivo* (Fig. 3F and 3G). In yeast, Mvp1 was shown to recruit Vps1 to the endosomes,

235 while in *F. graminearum*, Δ*Fgmvp1* mutant showed phenotypic differences with Δ*Fgvps1* [46, 49].

236 Also, Rab GTPases are required for the recruitment of retromer and sorting nexins to the endosomes

237 [11, 14, 41, 50]. Therefore, we proposed that FgRab51 and FgRab7 could mediate the recruitment of

238 FgVps1 to the endosomes in *F. graminearum*. We checked the localization of GFP-FgVps1 in the

239 wild type, Δ*Fgrab51* and Δ*Fgrab7* mutants. As shown in Fig. 3H, loss of the Rab GTPases resulted

240 in mis-localization of GFP-FgVps1 into the cytosol. We therefore conclude that FgVps1 is recruited

241 to the endosome by FgRab51 and FgRab7 to facilitate the scission of endosomal tubules for

242 retrograde trafficking in *F. graminearum* (Fig. 3I).

244 **Inhibiting the GTPase activity of FgVps1 by phenothiazine treatment causes mis-sorting of**

245 **Snc1**

246 Given that recruitment of FgVps1 to endosomes is required for effective scission of FgVps35 and

247 FgSnx4 endosomal tubules, and that FgVps35 and FgSnx4 are both required for the recycling of

248 FgSnc1 (Fig. 3, Fig. S4) [11, 48], we reasoned that FgSnc1 recycling could also be required for the

249 functioning of FgRab51, FgRab7 and FgVps1. First, we transformed GFP-FgSnc1 into the wild type

250 and Δ*Fgrab51* and Δ*Fgrab7* mutants and checked for possible alteration in FgSnc1 localization.

251 We found that the plasma membrane localization of FgSnc1 was significantly compromised in the

252 Rab GTPase mutants (Fig. S5A), indicating that the recycling of FgSnc1 also depends on the Rab

GTPases in *F. graminearum*. Next, we investigated the role of FgVps1 in FgSnc1 recycling. Similar

to FgVps35 and FgSnx4, FgVps1 can also interact with FgSnc1, and loss of FgVps1 has a severe

impact on the trafficking of FgSnc1 back to the plasma membrane (Fig. S5B and S5C).

Our earlier results showed that phenothiazines inhibit FgVps1 GTPase activity, and FgVps1-

dependent endosomal tubule scission is required for FgSnc1 sorting (Fig. 2, Fig. S5). It is therefore

possible that FgSnc1 recycling would be disrupted under phenothiazine treatments in *F.*

graminearum. First, we co-transformed GFP-FgSnc1 with mCherry-FgVps1 and mCherry-

FgVps1$^{\Delta POC}$ into $\Delta Fgvps1$ mutant to monitor the dynamics of FgSnc1, FgVps1 and the GTPase-

deficient form of FgVps1. We noticed that the mutation of the phenothiazine binding pocket altered

the localization of FgSnc1 (Fig. 4A and Fig. S6A). We further transformed GFP-FgSnc1 into the

wild type, $\Delta Fgvps1$ and Fgvps1$^{\Delta POC}$ mutants and analyzed its localization in these strains. We found

that the proportion of FgSnc1 detected on the plasma membrane was significantly low in the

$\Delta Fgvps1$ and Fgvps1$^{\Delta POC}$ mutants (Fig. 4B, Fig. S6B). Similarly, degradation level of FgSnc1 also

increased in the $\Delta Fgvps1$ and Fgvps1$^{\Delta POC}$ mutants following analysis by western blot (Fig. 4C).

These results indicate that the GTPase activity is required for the localization of FgSnc1 on plasma

membrane. We also found that the reduced GTPase activity of FgVps1 negates the scission of

retromer- and sorting nexin-coated endolysosomal tubules (Fig. S6C). Given that phenothiazines

inhibit FgVps1 GTPase activity, and considering the conserved role of SNARE Snc1 in the

development and pathogenicity of fungal pathogens (Fig. 1 and Fig. 4D), it is possible that Snc1

might be a downstream target of phenothiazines in phytopathogens. We tested this by treating the

strains expressing GFP-labeled Snc1 with CPZ and PCZ to evaluate the effects of the phenothiazines

on Snc1 proteins distribution. As shown in Fig. 4E, treatment with phenothiazines changed the

distribution of Snc1 in the different fungal pathogens. Overall, these results indicate that

phenothiazines inhibit the GTPase activity FgVps1 and causes mis-sorting of Snc1, which

consequently compromised phytopathogenic fungal development and host infection.

Discussion

During pathogen-host interactions, pathogens exploit the endolysosomal network to fine-tune

intracellular homeostasis, thereby facilitating host colonization [9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23,

24, 25, 26, 27, 28, 29, 30]. However, the complex interplay within the endolysosomal network and its

potential in fungal disease control remain largely unknown. Here, we identified the endocytic

dynamin Vps1 as a key regulator of endolysosomal network in phytopathogens. FgVps1 participates

in endocytic trafficking process through association with Rab GTPases, retromer and sorting nexins.

Inhibition of FgVps1 by phenothiazine treatments disrupts endocytosis and vesicle trafficking

efficiency, which impairs the development and pathogenesis of phytopathogens. (Fig. 5). These

results offer novel insights into the mechanism of how endolysosomal network functions, and

provide a potentially effective pathogenic fungal control strategy by disrupting endocytic dynamin

through repurposing of phenothiazines.

Phenothiazines, the clinically used antipsychotic drugs, also display antimicrobial effects against

human pathogenic bacterial and fungi [51, 52, 53]. Although the mechanisms of phenothiazines activity

have been described [52], the underlying mechanism of its antifungal effect remains unclear. Here, we

showed that phenothiazines displayed broad-spectrum inhibition against phytopathogens. (Fig. 1).

Our results indicated that phenothiazines can interact with the dynamin FgVps1 (through binding to

amino acids from D239 to I247 residues of the FgVps1 G domain) to inhibit its GTPase activity,

resulting in endocytosis inhibition in *F. graminearum* (Fig. 2). In addition, FgVps1-dependent

endocytic trafficking of FgSnc1 was also disrupted under phenothiazine treatments, and this effect

is conserved among plant pathogenic fungi (Fig. 4). These results suggest that the anti-fungal effect

of phenothiazines can be associated with endolyssosomal network dysfunction. Additionally, we

noticed that at a high concentration (100 μg/mL), phenothiazines cause more severe phenotypic

defects in the fungi than *FgVPS1* deletion (Fig. 1, Fig. 2 and Fig. S2). Previous studies indicated

that phenothiazines cause side effects related to impairment of calcium-calmodulin system,

membrane stabilization and efflux pumps [54, 55]. It is possible that these effects also contribute to the

antifungal property of phenothiazines. Since phenothiazines are synthetic drugs with diverse

biological activities [51], further understanding their mechanism of action may provide an avenue for

drug development and repurposing.

The integrity and functionality of endolysosomal network are vital for successful microbial infection

[56, 57, 58]. Additionally, pathogens can manipulate the host's endolysosomal network to enhance their

colonization efficiency [57, 59, 60, 61]. Given the importance of endocytosis in endolysosomal network,

its inhibition could be a novel approach for the control of fungal pathogens. To date, various

inhibitors of endocytosis have been characterized [35, 40]. Among them, phenothiazines were shown

to target dynamin GTPase activity in mammals [40]. Here, we provide evidence that the

phenothiazines CPZ and PCZ exert the conserved role of inhibiting dynamin GTPase activity, which

also broaden our understanding of how the dynamin FgVps1 function in the endolysosomal system

in fungal pathogens (Fig. 2, Fig. 3 and Fig. S4). Interestingly, although phenothiazines displayed

significant impacts on fungal pathogens, short-term treatment with the drugs does not affect the host

plant's growth and development (Fig. S1). However, one of the limitations of the study is our

inability to analyze the effects of long-term treatment on the hosts and field experiment.

Dynamins are multi-domain GTPases that are involved in multiple membrane remodeling

(including scission and fusion of endocytic vesicles and organelles) and cell division events [38, 39, 62].

Unlike the various dynamin proteins identified in mammals and plants [38, 39], fungi contain three

dynamin proteins namely Dnm1, Vps1 and Sey1; Dnm1-dependent mitochondrial fission and Sey1-

mediated ER fusion have been shown to be important for development, fungicide resistance and

pathogenicity in phytopathogens [63, 64, 65]. Although studies have implicated Vps1 in endocytosis,

vesicle scission and peroxisome fission and recycling events [44, 45, 46, 66, 67, 68], the mechanism of its

functions in fungal phytopathogens remains largely unknown. Our results showed that, in addition

to its role in endocytosis, FgVps1 can be recruited to endosomes by Rab GTPases to mediate

endosomal tubule scission, which further facilitates the intracellular trafficking of FgSnc1 (Fig.3,

Fig. S5 and Fig. 5). Disruption of FgVps1, either by gene deletion or phenothiazine treatments,

significantly impaired the development and pathogenicity of *F. graminearum* (Fig. 2, Fig. S2).

Interestingly, unlike the precise organization in yeast and mammals, the endosomal network in

fungal pathogens appears to be 'coarse' but efficient. Through the tight interconnection among

endolysosomal components, Rab GTPases (Rab5 and Rab7), dynamin (FgVps1) and sorting

machineries (retromer and sorting nexins), the endocytic trafficking required for the host

colonization process in phytopathogenic fungi can be fine-tuned (Fig. 3 and Fig. S4) [11, 41, 46, 49, 69].

Pathogens might have developed this for effective host colonization during long-term co-evolution

with their hosts. As endocytic dynamin is conserved among fungi (Fig. 2) [39], inhibition of endocytic

dynamin may act as an effective fungal control strategy.

Materials and methods

Fungal strains and culture conditions

For *F. graminearum*, the PH-1 strain was used as the wild type. For *M. oryzae and F. oxysporum*,

70-15 strain and FocTR4 strains were used as the wild type, respectively. A list of all the strains

used in this study is provided in Table S1.

In brief, the culture condition of *F. graminearum* were performed based on previously described [10].

F. graminearum strains were cultured on CM, CMC and TBI media for growth, conidial production

and DON induction, respectively. For vegetative growth and conidial production assays of *M.*

oryzae, the wild-type strain 70-15 was cultured on complete medium as previously described [70]. For

F. oxysporum growth and conidial production, wild type strains FocTR4 was culture on CM or PDB

media as previously described [71].

Generation of the *FgVPS1* deletion mutants and complementation

For the generation of Δ*Fgvps1* mutants, split-marker approach was utilized as described previously

[70]. Briefly, approximately 1kb upstream and downstream sequences of *FgVPS1* gene were amplified

using specific primers in Table S2. The resulting fragments were then fused with hygromycin

resistance gene cassette through splicing by overlap extension (SOE) PCR and then transformed

into the wild type PH-1 protoplasts. The resulting transformants were further screened by PCR and

southern blot assay. For Δ*Fgvps1* complementation, the GFP-Fgvps1 vector was transformed into

the Δ*Fgvps1* protoplasts and then the positive transformants were screened using confocal

microscopy. Additionally, the selected transformants were further verified through phenotypic

assays.

Plant infection assays

For *F. graminearum* infection on wheat coleoptiles, conidia suspensions (10^6 spores/mL) of the

indicated strains were inoculated on wheat coleoptiles for 7 days as previous described [72]. For

pathogenicity on wheat heads, mycelial plugs of the related strains were inoculated into flowering

wheat heads for 14 days as previously described [10]. For the monitor of invasive growth, conidia

suspensions were injected into wheat coleoptiles and incubated at 28 °C in dark with proper

humidity for 3 days and then observed under confocal microscopy.

For *M. oryzae* pathogenicity on barley, 10 μL conidial suspension (5×10^4 conidia/mL) of the

indicated strains were inoculated on barley leaves for 5 days. For rice spray assay, 10 mL conidial

suspensions (5×10^4 conidia/mL) were sprayed on rice 3-week-old CO39 rice seedlings and then

incubated at 28 °C with high humidity, lesions were recorded at 7 dpi (day post inoculation). For

the monitor of invasive growth, the mixture of conidial suspension and phenothiazines were injected

into rice leave sheath for 36 h.

For *F. oxysporum* host infection, hyphal blocks of the wild type were inoculated on wound banana

leaves with phenothiazines for 3 days, DMSO was used as control. Visualization of invasive growth

was performed as previously described [71].

Confocal microscopy

For monitoring endocytosis, conidia of the fungal strain were incubated with 10 μg/mL FM4-64 dye

(Cat. T3166, Invitrogen, Carlsbad, California) for 1 min, 10 min, and 25 min, respectively. For

staining the plasma membrane, hyphae of the indicated strains were stained with FM4-64 for 1 min

and then observed under confocal microscopy. Confocal examination was performed using a Nikon

CSU-W1 spinning disk confocal microscope (Nikon, Japan). The excitation and emission

wavelengths used were 488 nm/500-550 nm for GFP and 561 nm/570-620 nm for mCherry as

previously described [14].

Immunoprecipitation-mass spectrometry

For the immunoprecipitation-mass spectrometry assays, wild type strain PH-1 and strains

expressing the GFP and GFP-FgVps1 were cultured in liquid CM for two days. Mycelia were

harvested, ground into a fine powder, and then incubated in lysis buffer containing 2 mM PMSF

and protease inhibitor cocktail (Cat. GRF101, Epizyme Biotech, Shanghai, China) for 30 min.

Samples were centrifuged at 12,000 rpm for 20 minutes, and then the total cell lysates were further

incubated with 30 μL GFP-Trap A beads (Cat. no. gta-20; ChromoTek Inc., Planegg Martinsried,

Germany) for 4 hours and then heated at 100℃ with protein loading buffer (Cat. LT103, Epizyme

Biotech, Shanghai, China) for 10 minutes. Finally, the samples were separated by SDS-PAGE and

mass spectrometry analysis was conducted to identify the related proteins bound to the GFP-FgVps1.

Yeast two hybrid and co-immunoprecipitation (Co-IP) assays

For yeast two hybrid assays, the CDS sequence of the related genes were amplified using the specific

primers listed in table S2, the amplified sequences were fused with pGADT7 or pGBKT7 vectors

and further verified by sequencing (Sangon biotech, Shanghai, China). The resulting vectors were

transformed into yeast strain AH109 according to Matchmaker™ GAL4 Two-Hybrid System 3

protocol (Clontech, Mountain View, CA, USA).

For Co-IP assay, the strains expressing the fusion proteins were inoculated in liquid CM at 28℃,

110 rpm for 36 h. The sampled were collected and ground into fine powder in liquid nitrogen, and

then lysed in lysis buffer. The total cell lysates were then incubated with GFP-Trap A beads at 4 ℃

for 4 h. The bound proteins were subsequently eluted by heating at 100℃ for 10 min with protein

loading buffer and further analyzed by western blot with anti-GFP (Cat. M20004M, Abmart,

shanghai, China) and anti-Myc (Cat. ab1326, Abcam, Cambridge, United Kingdom) antibodies.

Phylogeny analysis, protein structure modeling and molecular docking

For phylogeny analysis, the proteins sequences of dynamin family proteins were download from the

NCBI (https://www.ncbi.nlm.nih.gov/). The related phylogeny tree was constructed using the

Neighbor-joining method through MEGA-X and further modified using iTOL [73]. The protein

structure of FgVps1, the FgVps35-FgSnx4 multimer were predicted using AlphaFold2 [74] and further

visualized using PyMOL. The structures of the prochlorperazine and chlorpromazine were

downloaded from the NCBI PubChem and the molecular docking of FgVps1with the

prochlorperazine and chlorpromazine were further performed using AutoDock and visualized by

PyMOL.

Protein purification and GTPase activity assay

For the purification of GST-FgVps1, GST-FgVps1$^{\Delta POC}$ recombinant proteins, the related CDS

sequences of FgVps1, FgVps1$^{\Delta POC}$ were amplified using the specific primers listed in Table S2 and

then fused with the pGET-4T-2 vector. The resulting vectors were expressed in *Escherichia coli*

BL21 cells and further sonicated in the lysis buffer (150 mM NaCl, 50 mM Tris HCL, pH 8.0, 10%

glycerol) containing 2 mM PMSF and protease inhibitor cocktail (Cat. GRF101, Epizyme Biotech,

Shanghai, China). After sonication, the suspensions were centrifuged at 4 ℃, 12000rpm for 20 min

and incubated with GST beads (Cat. A10018, Abmart, Shanghai, China) for 2 h. The beads were

then washed with the lysis buffer and eluted by incubating with the elution buffer (150 mM NaCl,

50 mM Tris HCL, pH 8.0, 150 mM glutathione) at 4 ℃ for 2 h.

For the effects of the phenothiazines on FgVps1 GTPase activity, the purified GST-FgVps1 was

incubated with the phenothiazines for 30 min and then the GTPase activity was measured using the

GTPase assay kit (Cat. ab272520, Abcam, Cambridge, United Kingdom) and Tristar² S LB 942

multimode reader (Berthold, Germany). For the effects of the pocket mutation on FgVps1 GTPase

activity, the purified GST-FgVps1 and GST-FgVps1$^{\Delta POC}$ were quantified using the Bradford Protein

Assay Kit (Cat. P0006C, Beyotime, Shanghai, China) and then the GTPase activity were measured

using the GTPase assay kit. GST was used as the negative control. Data were collected from three

biological repeats.

Author contributions

Conceptualization, Zonghua Wang and Wenhui Zheng; methodology, Xin Chen, Xiaomin Chen and

Wenhui Zheng; investigation, Xin Chen, Xiaomin Chen, Yunfei Long and Zhenyu Fang;

visualization, Xin Chen, Zonghua Wang and Wenhui Zheng; Writing - Original Draft, Xin Chen;

Writing - Review & Editing, Xin Chen, Xiaomin Chen, Yakubu Saddeeq Abubakar, Huawei Zheng,

Zonghua Wang and Wenhui Zheng; resources and funding acquisition, Wenhui Zheng.

Acknowledgements

This research was supported by the National Natural Science Foundation of China (32122071,

32272481), and the Natural Science Foundation of Fujian Province (2021J06015). The authors

declared no financial or other potential conflict of interest.

References

1. Fones HN, Bebber DP, Chaloner TM, Kay WT, Steinberg G, Gurr SJ. Threats to global food security from emerging fungal and oomycete crop pathogens. *Nat Food* **1**, 332-342 (2020).

2. Dean R, *et al.* The Top 10 fungal pathogens in molecular plant pathology. *Mol Plant Pathol* **13**, 414-430 (2012).

3. van der Does HC, Rep M. Adaptation to the Host Environment by Plant-Pathogenic Fungi. *Annu Rev Phytopathol* **55**, 427-450 (2017).

4. Wang Y, Pruitt RN, Nurnberger T, Wang Y. Evasion of plant immunity by microbial pathogens. *Nat Rev Microbiol* **20**, 449-464 (2022).

5. Cullen PJ, Steinberg F. To degrade or not to degrade: mechanisms and significance of endocytic recycling. *Nat Rev Mol Cell Biol* **19**, 679-696 (2018).

6. Borchers AC, Langemeyer L, Ungermann C. Who's in control? Principles of Rab GTPase activation in endolysosomal membrane trafficking and beyond. *J Cell Biol* **220**, (2021).

7. Migliano SM, Wenzel EM, Stenmark H. Biophysical and molecular mechanisms of ESCRT functions, and their implications for disease. *Curr Opin Cell Biol* **75**, 102062 (2022).

8. Posor Y, Jang W, Haucke V. Phosphoinositides as membrane organizers. *Nat Rev Mol Cell Biol* **23**, 797-816 (2022).

9. Zheng H, *et al.* Rab GTPases are essential for membrane trafficking-dependent growth and pathogenicity in Fusarium graminearum. *Environmental Microbiology* **17**, 4580-4599 (2015).

10. Yang C, *et al.* FgVps9, a Rab5 GEF, Is Critical for DON Biosynthesis and Pathogenicity in Fusarium graminearum. *Front Microbiol* **11**, 1714 (2020).

11. Zheng W, *et al.* The endosomal recycling of FgSnc1 by FgSnx41-FgSnx4 heterodimer is essential for polarized growth and pathogenicity in Fusarium graminearum. *New Phytol* **219**, 654-671 (2018).

12. Zheng W, *et al.* Retrograde trafficking from the endosome to the trans-Golgi network mediated by the retromer is required for fungal development and pathogenicity in Fusarium graminearum. *New Phytol* **210**, 1327-1343 (2016).

13. Wang S, *et al.* The regulatory role of the Aspergillus flavus core retromer complex in aflatoxin metabolism. *J Biol Chem* **298**, 102120 (2022).

506   14.   Chen X, *et al.* Rab7/Retromer-based endolysosomal trafficking is essential for proper host
507   invasion in rice blast. *New Phytologist* **239**, 1384-1403 (2023).
508

509   15.   Zhang X, *et al.* The small GTPase Ypt7 of Penicillium expansum is required for growth, patulin
510   biosynthesis and virulence. *Food Microbiol* **119**, 104434 (2024).
511

512   16.   Wu C, *et al.* The retromer CSC subcomplex is recruited by MoYpt7 and sequentially sorted by
513   MoVps17 for effective conidiation and pathogenicity of the rice blast fungus. *Mol Plant Pathol*
514   **22**, 284-298 (2021).
515

516   17.   Huang LY, Wu M, Yu XY, Li L, Lin FC, Liu XH. Physical interactions and mutational analysis
517   of MoYpt7 in Magnaporthe oryzae. *J Zhejiang Univ Sci B* **19**, 79-84 (2018).
518

519   18.   Liu J, Lei M, Zhou Y, Chen F. A Comprehensive Analysis of the Small GTPases Ypt7 Involved
520   in the Regulation of Fungal Development and Secondary Metabolism in Monascus ruber M7.
521   *Front Microbiol* **10**, 452 (2019).
522

523   19.   Li XY, Zhang SP, He L. Retromer subunit, CfVps35 is required for growth development and
524   pathogenicity of Colletotrichum fructicola. *BMC Genom Data* **23**, 68 (2022).
525

526   20.   Li S, Li X, Li H. The Retromer Subunit CfVps29 Is Involved in the Growth, Development, and
527   Pathogenicity of Colletotrichum fructicola. *J Fungi (Basel)* **8**, (2022).
528

529   21.   Sun L, *et al.* A Subunit of ESCRT-III, MoIst1, Is Involved in Fungal Development,
530   Pathogenicity, and Autophagy in Magnaporthe oryzae. *Front Plant Sci* **13**, 845139 (2022).
531

532   22.   Cheng J, Yin Z, Zhang Z, Liang Y. Functional analysis of MoSnf7 in Magnaporthe oryzae.
533   *Fungal Genet Biol* **121**, 29-45 (2018).
534

535   23.   Xie Q, *et al.* Component Interaction of ESCRT Complexes Is Essential for Endocytosis-
536   Dependent Growth, Reproduction, DON Production and Full Virulence in Fusarium
537   graminearum. *Front Microbiol* **10**, 180 (2019).
538

539   24.   Xie Q, *et al.* ESCRT-III accessory proteins regulate fungal development and plant infection in
540   Fusarium graminearum. *Curr Genet* **65**, 1041-1055 (2019).
541

542   25.   Xie Q, *et al.* Endosomal sorting complexes required for transport-0 is essential for fungal
543   development and pathogenicity in Fusarium graminearum. *Environ Microbiol* **18**, 3742-3757
544   (2016).
545

546   26.   Sun LX, *et al.* Endosomal sorting complexes required for transport-0 (ESCRT-0) are essential
547   for fungal development, pathogenicity, autophagy and ER-phagy in Magnaporthe oryzae.
548   *Environ Microbiol* **24**, 1076-1092 (2022).
549

27. Li B, Mao HY, Zhang ZY, Chen XJ, Ouyang SQ. FolVps9, a Guanine Nucleotide Exchange Factor for FolVps21, Is Essential for Fungal Development and Pathogenicity in Fusarium oxysporum f. sp. lycopersici. *Front Microbiol* **10**, 2658 (2019).

28. Zheng W, *et al.* Retromer Is Essential for Autophagy-Dependent Plant Infection by the Rice Blast Fungus. *PLoS Genet* **11**, e1005704 (2015).

29. Liu XH, *et al.* The small GTPase MoYpt7 is required for membrane fusion in autophagy and pathogenicity of Magnaporthe oryzae. *Environ Microbiol* **17**, 4495-4510 (2015).

30. Zheng H, *et al.* Sorting nexin (MoVps17) is required for fungal development and plant infection by regulating endosome dynamics in the rice blast fungus. *Environ Microbiol* **19**, 4301-4317 (2017).

31. Sumarah MW. The Deoxynivalenol Challenge. *J Agric Food Chem* **70**, 9619-9624 (2022).

32. Doherty GJ, McMahon HT. Mechanisms of endocytosis. *Annu Rev Biochem* **78**, 857-902 (2009).

33. Smith SM, Smith CJ. Capturing the mechanics of clathrin-mediated endocytosis. *Curr Opin Struct Biol* **75**, 102427 (2022).

34. Chew HY, *et al.* Endocytosis Inhibition in Humans to Improve Responses to ADCC-Mediating Antibodies. *Cell* **180**, 895-914 e827 (2020).

35. Szewczyk-Roszczenko OK, *et al.* The Chemical Inhibitors of Endocytosis: From Mechanisms to Potential Clinical Applications. *Cells* **12**, (2023).

36. Kaksonen M, Roux A. Mechanisms of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* **19**, 313-326 (2018).

37. Mettlen M, Chen PH, Srinivasan S, Danuser G, Schmid SL. Regulation of Clathrin-Mediated Endocytosis. *Annu Rev Biochem* **87**, 871-896 (2018).

38. Ferguson SM, De Camilli P. Dynamin, a membrane-remodelling GTPase. *Nat Rev Mol Cell Biol* **13**, 75-88 (2012).

39. Ramachandran R, Schmid SL. The dynamin superfamily. *Curr Biol* **28**, R411-R416 (2018).

40. Daniel JA, *et al.* Phenothiazine-derived antipsychotic drugs inhibit dynamin and clathrin-mediated endocytosis. *Traffic* **16**, 635-654 (2015).

41. Abubakar YS, *et al.* FgRab5 and FgRab7 are essential for endosomes biogenesis and non-redundantly recruit the retromer complex to the endosomes in Fusarium graminearum. *Stress Biology* **1**, (2021).

42.   Simanjuntak Y, Liang JJ, Lee YL, Lin YL. Repurposing of prochlorperazine for use against
dengue virus infection. *J Infect Dis* **211**, 394-404 (2015).

43.   Morris GM*, et al.* AutoDock4 and AutoDockTools4: Automated docking with selective receptor
flexibility. *J Comput Chem* **30**, 2785-2791 (2009).

44.   Ma M, Burd CG, Chi RJ. Distinct complexes of yeast Snx4 family SNX-BARs mediate
retrograde trafficking of Snc1 and Atg27. *Traffic* **18**, 134-144 (2017).

45.   Chi RJ, Liu J, West M, Wang J, Odorizzi G, Burd CG. Fission of SNX-BAR-coated endosomal
retrograde transport carriers is promoted by the dynamin-related protein Vps1. *J Cell Biol* **204**,
793-806 (2014).

46.   Suzuki SW, Oishi A, Nikulin N, Jorgensen JR, Baile MG, Emr SD. A PX-BAR protein
Mvp1/SNX8 and a dynamin-like GTPase Vps1 drive endosomal recycling. *Elife* **10**, (2021).

47.   Hettema EH. Retromer and the sorting nexins Snx4/41/42 mediate distinct retrieval pathways
from yeast endosomes. *The EMBO Journal* **22**, 548-557 (2003).

48.   Long Y*, et al.* Golgi-associated retrograde protein (GARP) complex recruits retromer to trans-
Golgi network for FgKex2 and FgSnc1 recycling, necessary for development and pathogenicity
of Fusarium graminearum. *bioRxiv*, (2023).

49.   Lou Y*, et al.* Genome-Wide Characterization of PX Domain-Containing Proteins Involved in
Membrane Trafficking-Dependent Growth and Pathogenicity of Fusarium graminearum. *mBio*
12, e0232421 (2021).

50.   Rojas R*, et al.* Regulation of retromer recruitment to endosomes by sequential action of Rab5
and Rab7. *J Cell Biol* **183**, 513-526 (2008).

51.   Jaszczyszyn A*, et al.* Chemical structure of phenothiazines and their biological activity.
Pharmacol Rep **64**, 16-23 (2012).

52.   Varga B, Csonka A, Csonka A, Molnar J, Amaral L, Spengler G. Possible Biological and
Clinical Applications of Phenothiazines. *Anticancer Res* **37**, 5983-5993 (2017).

53.   Grimsey EM, Piddock LJV. Do phenothiazines possess antimicrobial and efflux inhibitory
properties? *FEMS Microbiol Rev* **43**, 577-590 (2019).

54.   Sharma S, Kaur H, Khuller GK. Cell cycle effects of the phenothiazines: trifluoperazine and
chlorpromazine in Candida albicans. *FEMS Microbiol Lett* **199**, 185-190 (2001).

55.   Amaral L*, et al.* Phenothiazines, bacterial efflux pumps and targeting the macrophage for

638   enhanced killing of intracellular XDRTB. *In Vivo* **24**, 409-424 (2010).

56.   Shoji JY, Kikuma T, Kitamoto K. Vesicle trafficking, organelle functions, and unconventional

secretion in fungal physiology and pathogenicity. *Curr Opin Microbiol* **20**, 1-9 (2014).

57.   Elwell C, Engel J. Emerging Role of Retromer in Modulating Pathogen Growth. *Trends*

Microbiol **26**, 769-780 (2018).

58.   Abubakar YS, Sadiq IZ, Aarti A, Wang Z, Zheng W. Interplay of transport vesicles during plant-

fungal pathogen interaction. *Stress Biol* **3**, 35 (2023).

59.   Cossart P, Helenius A. Endocytosis of viruses and bacteria. *Cold Spring Harb Perspect Biol* **6**,

(2014).

60.   Simonetti B, Daly JL, Cullen PJ. Out of the ESCPE room: Emerging roles of endosomal SNX-

BARs in receptor transport and host-pathogen interaction. *Traffic* **24**, 234-250 (2023).

61.   Yuen ELH, Shepherd S, Bozkurt TO. Traffic Control: Subversion of Plant Membrane

Trafficking by Pathogens. *Annu Rev Phytopathol* **61**, 325-350 (2023).

62.   Antonny B*, et al.* Membrane fission by dynamin: what we know and what we need to know.

EMBO J **35**, 2270-2284 (2016).

63.   Zhong K*, et al.* MoDnm1 Dynamin Mediating Peroxisomal and Mitochondrial Fission in

Complex with MoFis1 and MoMdv1 Is Important for Development of Functional Appressorium

in Magnaporthe oryzae. *PLoS Pathog* **12**, e1005823 (2016).

64.   Kang J-b*, et al.* Mitochondrial dynamics caused by QoIs and SDHIs fungicides depended on

FgDnm1 in Fusarium graminearum. *Journal of Integrative Agriculture* **22**, 481-494 (2023).

65.   Chong X*, et al.* The Dynamin-Like GTPase FgSey1 Plays a Critical Role in Fungal

Development and Virulence in Fusarium graminearum. *Appl Environ Microbiol* **86**, (2020).

66.   Arlt H, Reggiori F, Ungermann C. Retromer and the dynamin Vps1 cooperate in the retrieval of

transmembrane proteins from vacuoles. *J Cell Sci* **128**, 645-655 (2015).

67.   Ekal L, Alqahtani AMS, Hettema EH. The dynamin-related protein Vps1 and the peroxisomal

membrane protein Pex27 function together during peroxisome fission. *J Cell Sci* **136**, (2023).

68.   Palmer SE*, et al.* A dynamin-actin interaction is required for vesicle scission during endocytosis

in yeast. *Curr Biol* **25**, 868-878 (2015).

69.   Wang J, Fedoseienko A, Chen B, Burstein E, Jia D, Billadeau DD. Endosomal receptor

trafficking: Retromer and beyond. *Traffic* **19**, 578-590 (2018).

70.   Chen X, *et al.* Trehalose Phosphate Synthase Complex-Mediated Regulation of Trehalose 6-
684   Phosphate Homeostasis Is Critical for Development and Pathogenesis in Magnaporthe oryzae.
685   *mSystems* **6**, (2021).
686

687   71.   Fang Z, *et al.* Two distinct SNARE complexes mediate vesicle fusion with the plasma
688   membrane to ensure effective development and pathogenesis of Fusarium oxysporum f. sp.
689   cubense. *Molecular Plant Pathology* **25**, (2024).
690

691   72.   Jia LJ, Wang WQ, Tang WH. Wheat Coleoptile Inoculation by Fusarium graminearum for
692   Large-scale Phenotypic Analysis. *Bio Protoc* **7**, e2439 (2017).
693

694   73.   Letunic I, Bork P. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree
695   display and annotation tool. *Nucleic Acids Res*, (2024).
696

697   74.   Mirdita M, Schutze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making
698   protein folding accessible to all. *Nat Methods* **19**, 679-682 (2022).
699

    Figure legend

Fig.1 Treatment of phenothiazines impairs the pathogenicity of phytopathogens.

A. Phenothiazines treatment has significant impacts on the fungal growth of the indicated

phytopathogens. For evaluating the effects of phenothiazines on fungal growth, mycelial blocks of

F. graminearum, *F. oxysporum*, and *M. oryzae* were inoculated on complete medium with or without

phenothiazines for 2 days, 3 days, and 7 days, respectively. B. Treatment of CPZ and PCZ impaired

the host invasion of the *F. graminearum*. For wheat infection, 50 μL conidia suspensions (10^6

spores/mL) of the *F. graminearum* PH-1 strain were inoculated on wheat coleoptiles with or without

CPZ or PCZ for 7 days. For the observation of invasive growth, conidia suspensions of *F.*

graminearum PH-1 strain expressing GFP were injected into wheat coleoptiles with CPZ or PCZ;

DMSO treatment was used as a control. No obvious invasive hyphae were detected at 3 dpi under

the treatment of phenothiazines. C. Effects of CPZ and PCZ treatment on *F. oxysporum* invasion.

For pathogenicity assays, hyphal blocks were inoculated on wounded banana leaves for 3 days with

CPZ or PCZ treatment, respectively. For the observation of invasive growth, conidial suspensions

of *F. oxysporum* expressing GFP were co-incubated with banana roots for 3 days and then the cross

sections of the banana roots were observed under confocal microscopy. white asterisks indicate the

invasion hyphae of the *F. oxysporum*. The invasive hyphae were hardly detectable in phenothiazine-

treated banana roots. D. Effects of PCZ and CPZ treatment on *M. oryzae* host invasion. For the

pathogenicity of *M. oryzae* on barley leaves, 10 μL conidia suspensions were inoculated on barley

leaves with PCZ or CPZ treatment for 5 days. For the rice spray assay, a 10 mL mixture of conidia

suspension (at a final concentration of $5*10^4$) with phenothiazines was sprayed on rice leaves,

photos were taken at 7 dpi (days post inoculation). DMSO treatment was used as a control. For the

monitor of invasive growth, the mixture of conidia suspension with phenothiazines was injected

into rice leaf sheaths for 36 h. No invasive hyphae were detected when co-incubated with the

phenothiazines CPZ and PCZ at 36 hpi (hours post inoculation).

Fig.2 Phenothiazines inhibit FgVps1 GTPase activity to impair the endocytosis of *F.*

graminearum.

A. Effects of phenothiazines (PCZ and CPZ) treatment on the endocytosis of *F. graminearum*. B.

Phylogeny analysis of the dynamin family proteins. The phylogenetic tree was constructed using

MEGA-X and then modified with iTOL. C. FM4-64 staining to observed the effects of *FgVPS1* on

endocytosis. D. Purification of the GST and GST-FgVps1 recombinant protein and subsequent

GTPase activity assays. To evaluate the effects of phenothiazines on FgVps1 GTPase activity, 5 μL

of the purified GST-FgVps1 was co-incubated with 5 μL of phenothiazines (at a final concentration

of 100 μg/mL) for 30 min and then used for GTPase activity assay. E. Molecular docking of the

prochlorperazine and chlorpromazine with FgVps1. The molecular docking of FgVps1with the

phenothiazines was analyzed using AutoDock and further visualized by PyMOL. FgVps1 is

depicted in surface representation (gray), and PCZ and CPZ are depicted in sticks representation. In

the right panel, the binding pocket is depicted in lines (salmon), PCZ is represented in blue, and

CPZ is represented in light blue. F. Purification of the GST, GST-FgVps1and GST-FgVps1$^{\Delta POC}$

recombinant proteins and subsquent GTPase activity assays. The concentration of the purified

recombinant proteins was quantified, and then 10 μL of each protein was used for further GTPase

activity assay. G. Effects of phenothiazine drugs binding pocket on the endocytosis of *F.*

graminearum. H. Confocal examination for the localization of GFP-FgVps1 and GFP-FgVps1$^{\Delta POC}$.

I. Pathogenicity assay of the binding pocket mutation on wheat heads.

Fig. 3 Rab GTPases-dependent FgVps1 recruitment facilitates FgVps1-mediated

endolysosomal tubule fission in *F. graminearum*.

A. GFP-FgVps1 pull down tandem LC-MS/MS assays for the identification of putative FgVps1

interacting proteins. B. Confocal examination for the co-localization of FgVps35-GFP with FgSnx4-

mCherry in wild type and Δ*Fgvps1* mutant. White boxes indicate the selected area used for time-

lapse imaging. C. Putative interacting model of FgVps35 with sorting nexin FgSnx4. FgVps35 is

represented in light blue, FgSnx4 is represented in wheat, and the interface is represented in pale

green. Black box indicates the enlarged area. Yellow dash represents the hydrogen bond formed

between FgSnx4 E373 and FgVps35 N383. D. Yeast two hybrid assay to verify the effects of point

mutations on the interaction between FgVps35 and FgSnx4. E. Pathogenicity assays of the point

mutation strains on wheat heads. F. Confocal examination and Co-IP assays to verify the interaction

between FgVps1 and the Rab GTPase FgRab51.White boxes indicate the area selected for time-

lapse imaging. G. Confocal examination and Co-IP assays to verify the interaction between FgVps1

and the Rab GTPase FgRab7.White boxes indicate the area selected for time-lapse imaging. H.

Effects of the Rab GTPase on the subcellular distribution of FgVps1. I. Diagram represents the roles of dynamin FgVps1 in the endolysosomal network of *F. graminearum*.

Fig. 4 Inhibition of GTPase activity by phenothiazine drugs impair the homeostasis of SNARE proteins Snc1 in different phytopathogens.

A. Time-lapse imaging for the dynamics of GFP-FgSnc1 with mCherry-FgVps1 and mCherry-FgVps1$^{\Delta POC}$. B. Confocal examination of the effects of the FgVps1$^{\Delta POC}$ mutation on the distribution of FgSnc1. C. Western blot assay for the protein levels of GFP and GFP-FgSnc1 in the indicated strains. D. Pathogenicity assays of the related *SNC1* mutants in different species. For the pathogenicity of the *F. graminearum*, hyphal blocks of the wild type and Δ*Fgsnc1* mutant were inoculated on wheat heads and then photographed at 14 dpi. For the pathogenicity of Δ*Mosnc1* mutant on rice, conidial suspensions of indicated strains were sprayed on three-weeks rice cultivar CO-39 for 7 days. For the pathogenicity assay of the *F. oxysporum,* hyphal blocks of the wild type and Δ*Focsnc1* were inoculated on wounded banana leaves for 3 days. For the observation of invasive growth, conidial suspensions of the wild type and Δ*Focsnc1* mutant expressing GFP were co-incubated with banana roots for 3 days and then the cross sections of the banana roots were observed under confocal microscopy. E. Confocal examination for the effects of phenothiazine drugs treatment on the distribution of Snc1 in phytopathogens. White boxes indicate the enlarged area in the right panel. The plasma membrane and septa were marked by FM4-64 dye.

Fig. 5 Proposed model of dynamin FgVps1 in endocytic trafficking and pathogenicity of *F. graminearum*.

For endolysosomal trafficking, the endocytic dynamin FgVps1 engages in the release of the endocytosed vesicles to allow the cell surface cargos to enter the endolysosomal system. Moreover, FgVps1 is recruited to the endosome by Rab GTPases (FgRab51 and FgRab7) to mediate the fission of retromer and sorting nexins endolysosomal tubules, facilitating the recycling of related cargos (e.g., FgSnc1). Dynamin GTPase inhibition, caused by dynamin FgVps1 disruption or phenothiazines treatment, significantly inhibits the endocytosis and endocytic trafficking process, resulting in the dysfunction of the endolysosomal system and leading to defects in host colonization.

A

FM4-64 detected in vacuole at 25 min

Wild type

Wild type + Prochlorperazine **

Wild type + Chlorpromazine **

B

bootstrap
0.54 0.66 0.77 0.89 1

● Endocytic vesicles fission ● Mitochondrial fission ● ER fusion

Sey1 Fusarium graminearum XP 011316549

Vps1 Fusarium graminearum XP 011326892

Dnm1 Fusarium graminearum XP 011317106

C

Wild type

FM4-64 1 min   Enlarged
FM4-64 25 min
10 μm

ΔFgvps1

FM4-64 1 min   Enlarged
FM4-64 25 min
10 μm

FM4-64 detected in vacuole at 25 min

Wild type

ΔFgvps1 **

D

Marker GST GST-FgVps1

GTPase activity (U/L)

FgVps1
FgVps1 + Prochlorperazine **
FgVps1 + Chlorpromazine **

E

FgVps1 with Prochlorperazine

Enlarged
3.0
2.9 Asp 246
-6.46 Kcal/mol Asp 243

FgVps1 with Chlorpromazine

Enlarged
2.9 Asp 243
-5.33 Kcal/mol

D239
M238
G241
T242
D243
2.9
2.9 V244
I245
I247
D246 3.0

Chlorpromazine
Prochlorperazine

F

Marker GST GST-FgVps1 GST-FgVps1^ΔPOC

GTPase activity (U/L)

FgVps1
FgVps1^ΔPOC **

G

FM4-64 detected in vacuole at 25 min

Wild type
ΔFgvps1 **
FgVps1^ΔPOC **

H

GFP-FgVps1
Merge GFP

GFP-FgVps1^ΔPOC
Merge GFP
20 μm

I

Wild type ΔFgvps1 FgVps1^ΔPOC

Disease index

Wild type
ΔFgvps1 **
FgVps1^ΔPOC **

+ *Fusarium graminearum*

+ Phenothiazines
+ *Fusarium graminearum*

Wheat

PM

Vps1 mediated
fission of endocytic vesicle

Vps1 mediated
endolysosomal tubule fission

recruitment

fission

Rab GTPases

FgVps1

FgSnc1

FgVps35

FgSnx4

Golgi

Vps1 GTPase activity

Golgi

Vacuole

Fusarium graminearum