Title: Synapse weakening-induced caspase-3 activity confers specificity to microglia-mediated synapse elimination

Authors: Zhou Yu[1,2†*], Andrian Gutu[1†], Namsoo Kim[1], Erin K. O'Shea[1*]

Affiliations:

[1]Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA.

[2]Current address: Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA.

[†]These authors contributed equally.

[*]Correspondence: zyu3@bwh.harvard.edu, osheae@hhmi.org.

Abstract: During brain development, synapses are initially formed in excess and are later eliminated in an activity-dependent manner, with weak synapses being preferentially removed. Previous studies identified glia as mediators of synapse removal, but it is unclear how glia specifically target weak synapses. Here we show that, in the developing mouse visual pathway, inhibiting synaptic transmission induces postsynaptic activation of caspase-3. Caspase-3 is essential for synapse elimination driven by both spontaneous and experience-dependent neural activity. Synapse weakening-induced caspase-3 activation determines the specificity of synapse elimination mediated by microglia but not astrocytes. Furthermore, in a mouse model of Alzheimer's disease, caspase-3 deficiency protects against synapse loss induced by amyloid-β deposition. Our results reveal caspase-3 activation as a key step in activity-dependent synapse elimination during development and synapse loss in neurodegeneration.

Main Text:

Synapse elimination is the process in which excess synapses formed early during

brain development are subsequently eliminated to form mature circuits (*1-3*). A key

feature of synapse elimination is that the outcome critically depends on both spontaneous

and experience-driven neural activity, and less active synapses are preferentially

removed (*2, 4, 5*). Previous studies have established that microglia and astrocytes, two

major glial cell types of the brain, engage in synapse engulfment by using phagocytic

receptors to detect 'eat-me' signals displayed on synapses (*6-14*). Although both

microglia- and astrocyte-mediated synapse elimination are known to be activity-

dependent (*7, 10*), it is unclear how the intrinsic strength of a synapse is converted into

external molecular cues that can be recognized by glial cells to guide the selective

removal of weaker synapses. A recent study proposed a molecular mechanism for

activity-dependent synapse elimination in which the JAK2-STAT1 pathway is activated in

inactive callosal projection neurons and regulates the removal of synapses and axons

(*15*). However, the JAK-STAT pathway canonically functions through transcriptional

regulation (*16*), which affects entire cells. If not all synapses made by a neuron are

destined for removal (*3*), activation of the JAK2-STAT1 pathway alone may not provide

sufficient elimination specificity. Therefore, other unidentified, locally-effective

mechanisms likely exist that confer elimination specificity at the synapse level.

Caspase-3 is a protease crucial for the execution of apoptosis (*17*). In hippocampal

neurons, transient activation of caspase-3 is required for long-term depression (LTD), a

form of synaptic plasticity that induces long-lasting decreases in synaptic strength (*18*).

Caspase-3 activation is also known to trigger the display of 'eat-me' signals on apoptotic

cells, facilitating their clearance by phagocytes through the engagement of phagocytic

receptors (*19, 20*). These findings led us to hypothesize that caspase-3 may link synapse

weakening with glial recognition and subsequent synapse removal. Although caspases

were previously implicated in dendrite pruning in metamorphosis, degeneration-like axon

elimination, and the regulation of hippocampal spine density (*21-24*), the role of caspase-

3 in synapse elimination in response to neural activity was not clear. In this work, we

identify caspase-3 as a pivotal molecule that is activated in postnatal development in

dendritic compartments upon synaptic weakening and is necessary to direct microglia to

selectively eliminate weak synapses. Furthermore, we discovered that caspase-3

deficiency protects against amyloid-β-induced synapse loss in a mouse model of

Alzheimer's disease, highlighting its significance not only in development synapse

elimination but also in adult neurodegenerative diseases.

Synapse inactivation induces postsynaptic caspase-3 activity

We chose the mouse retinogeniculate visual pathway as a model system to study

synapse elimination (*25*). In mice, retinal ganglion cells (RGCs) in each eye send out

axons to form retinogeniculate synapses with relay neurons in the dorsal lateral

geniculate nuclei (dLGN) of the thalamus on both sides of the brain (Fig. S1) (*25*). The

majority of RGC axons from each eye innervate the contralateral (on the side opposite to

the originating eye) dLGN while a smaller fraction innervate the ipsilateral (on the same

side as the originating eye) dLGN (Fig. S1) (*25*). Regions in each dLGN innervated by the

two eyes initially overlap but later segregate into eye-specific territories (Fig. S1). This

segregation process is a hallmark of retinogeniculate pathway development at the

morphological level and depends on synapse elimination and spontaneous RGC activity

(*25-27*).

To investigate the role of caspase-3 in activity-dependent synapse elimination, we

first needed to establish a method that could manipulate the strength of a selective subset

of retinogeniculate synapses. For this purpose, we used adeno-associated-virus (AAV) to

deliver a construct expressing tetanus toxin light chain (TeTxLC) under the control of the

neuron-specific human synapsin promoter (AAV-hSyn-TeTxLC) (Fig. 1A) (*28*). *In utero*

intraocular injection of AAV-hSyn-TeTxLC at embryonic day 15 (E15) leads to TeTxLC

expression in RGCs by the time of birth, blocking neurotransmitter release at

retinogeniculate synapses by cleaving synaptobrevin (*28, 29*). As anterograde tracers

and controls, we used AAV to express a fluorescent protein (mTurquiose2, eGFP, or

tdTomato, depending on the experiment) either alone or together with TeTxLC (Fig. 1A).

To validate our method, we injected AAV-hSyn-TeTxLC into the right eye of wildtype E15

embryos and analyzed eye-specific territories at postnatal day 8 (P8) (see Fig. S2 and

text associated with Fig. 3 for details of the analysis), when the segregation process is

largely complete (*25*). Consistent with previous results (*26, 27*), inactivating right eye-

associated retinogeniculate synapses led to contraction of right eye-specific territories

and expansion of left eye-specific territories relative to control animals (Fig. S3A and S3B).

In addition, segregation of eye-specific territories in TeTxLC-injected mice was defective

(Fig. S3C and S3D), confirming that our method potently inactivated synapses and

perturbed activity-dependent refinement of the retinogeniculate pathway.

Is caspase-3 activated in response to synapse weakening? To address this

question, we injected AAV-hSyn-TeTxLC into the right eyes of E15 embryos and looked

for caspase-3 activation in dLGNs of TeTxLC-injected mice at the age of P5 (a time when

synapse elimination is most active) (Fig. 1A) (*7, 25*). TeTxLC expression causes a

widespread inactivation of retinogeniculate synapses and enhances potential synapse

inactivation-induced caspase-3 activity. We measured caspase-3 activity by detecting the

cleaved and active form of caspase-3 through immunohistochemistry (*17*). If synapse

inactivation leads to caspase-3 activation, we would expect higher levels of caspase-3

activity in dLGNs with more inactivated synapses (Fig. 1A). In agreement with this

hypothesis, normalized caspase-3 activity in the left dLGN of right eye TeTxLC-injected

animals (Fig. 1B and 1E) was significantly higher compared to that in the right dLGN of

right eye TeTxLC-injected animals (Fig. 1C and 1E) and compared to that in the left dLGN

of Turq-injected control animals (Fig. 1D and 1E). We did not detect differences in

caspase-3 activity between the right dLGN of right eye TeTxLC-injected animals and the

dLGN of control animals (Fig. 1C, 1D and 1E), presumably due to low levels of synaptic

inactivation in the former and/or insufficient signal-to-background ratio. Reassuringly,

caspase-3 activities in left and right dLGN of control animals were comparable (Fig. 1E),

suggesting that surgical manipulation and viral transduction did not contribute to caspase-

3 activation. These findings support the model that inactivation of retinogeniculate

synapses induces caspase-3 activation.

Does inactivation of retinogeniculate synapses activate caspase-3 pre-synaptically

in RGCs or post-synaptically in dLGN relay neurons? Caspase-3 activity in dLGNs of

TeTxLC-injected mice was present either as discrete, punctate signals (Fig. 1F) or as

staining of the entire cell (Fig. 1B and Fig. S4Bi-iii). Both types of signals were more

abundant in dLGNs with strong synapse inactivation than in control dLGNs (Fig. 1B, 1D

and 1F). Intriguingly, punctate caspase-3 activity was not found within TeTxLC-expressing

RGC axons (Fig. 1G). Instead, punctate caspase-3 activity closely juxtaposed inactivated

axon terminals (Fig. 1G) and co-localized with a dendritic marker, MAP2 (Fig. S4A). In

the case where caspase-3 was activated in entire cells, the cells were positive for nuclear

NeuN (Fig. S4C) and possessed relatively large round somas and multipolar dendritic

arbors (Fig. S4Bi-iii), all of which are characteristics of dLGN relay neurons (*30, 31*).

These observations suggest that caspase-3 activation triggered by synapse inactivation

occurs predominantly in the dendritic compartments of dLGN relay neurons that are

postsynaptic to the inactivated synapses. It is important to note that the whole-cell

caspase-3 activation observed in TeTxLC-inactivated dLGNs is likely a consequence of

strong synapse inactivation and is not a normal developmental phenomenon (compare

Fig. 1B and 1D). Consistent with this idea, we observed a transitional state of caspase-3

activation where multiple dendrites of a relay neuron were positive for activate caspase-

3 without the cell body becoming apoptotic (Fig. S4Biv), suggesting that, in TeTxLC-

inactivated dLGNs, strong synapse inactivation causes excessive accumulation of

caspase-3 activity in dendrites and eventually leads to caspase-3 activation in entire relay

neurons, possibly through caspase-3 mediated feedback amplifications of the apoptosis

signaling cascade (*32*).

Synapse inactivation-induced caspase-3 activation requires synaptic competition

Is synapse inactivation-induced caspase-3 activity a result of synapse inactivation

per se, or is it a result of competition between inactive and active synapses? Previous

studies have either hyperactivated or inactivated RGCs in one of the two eyes while

leaving the other eye unperturbed (*26, 27*). In the scenario involving hyperactivation, both

the less active (unperturbed) and the more active (hyperactivated) eye had higher activity

than their counterparts in the scenario involving inactivation. However, in both

experiments, the dLGN territory of the less active eye contracted while that of the more

active eye expanded (*26, 27*), suggesting that relative, rather than absolute, synapse

strength determines the outcome of synapse elimination in the retinogeniculate pathway.

Consistent with this idea, a recent study demonstrated that elimination of inactive

synapses in callosal projections occurred only when other active synapses are present

(*28, 33*).

To test if synapse inactivation-induced caspase-3 activation requires synaptic

competition, we devised two conditions where either one (single inactivation) or both eyes

(dual inactivation) were injected with AAV-hSyn-TeTxLC (Fig. 2A). In single inactivation,

retinogeniculate synapses originating from different eyes have different strengths and

compete within each dLGN (Fig. 2A), whereas in dual inactivation, all retinogeniculate

synapses are inactivated and no competition occurs (Fig. 2A). We first confirmed that, in

accordance with previous findings (*26, 34*), inactivating RGCs in both eyes with TeTxLC

disrupted segregation of eye-specific regions (Fig. S5A-D). We then measured caspase-

3 activity in the single- or dual-inactivated dLGNs and in the control dLGN, where

retinogeniculate synapses were unperturbed (Fig. 2A). Intriguingly, while single

inactivation induced significantly higher levels of caspase-3 activity in the dLGN

compared to controls (Fig. 2B and 2C), active caspase-3 signal in the dual-inactivated

dLGN was not significantly different from the signal in the control dLGN (Fig. 2B and 2C)

and was significantly lower than that in the single-inactivated dLGN (Fig. 2B and 2C).

Although not reaching statistical significance, caspase-3 activity on average was still

higher in the dLGN in dual inactivation than in the control dLGN (Fig. 2C), possibly due

to incomplete inactivation of retinogeniculate synapses, or competition between

retinogeniculate synapses and synapses from other pathways, or additional mechanisms

that activate caspase-3 independently of synapse competition. These results suggest that

competition between active and inactive retinogeniculate synapses is required to induce

caspase-3 activity in dendritic compartments of dLGN relay neurons.

Caspase-3 is required for segregation of eye-specific territories

Is synapse inactivation-induced caspase-3 activity important for synapse

elimination in the retinogeniculate pathway? To address this question, we investigated

whether segregation of eye-specific dLGN territories is defective if caspase-3 activation

does not occur. As we were unable to find a Cre-expressing mouse line that satisfactorily

depletes caspase-3 in dLGN relay neurons, we used full-body caspase-3 deficient

(*Casp3$^{-/-}$*) mice (*35, 36*). These mice were backcrossed to congenicity on the C57Bl/6J

background and were viable, fertile, and did not exhibit overt abnormalities (*36*).

To measure segregation of eye-specific territories, we injected the eyes of

Casp3$^{+/+}$ and *Casp3$^{-/-}$* littermate mice at P9 with fluorophore-conjugated cholera toxin

subunit B (CTB), using a different fluorophore for each eye. CTB molecules bind to

ganglioside molecules on RGC surfaces and anterogradely label RGC axon terminals

(*37*). At P10, when CTB labeling and eye-specific segregation were complete (*25, 38*),

we collected the brains and imaged eye-specific territories in each dLGN. By visual

inspection, eye-specific territories in *Casp3$^{+/+}$* dLGNs overlapped minimally (Fig. 3A). By

contrast, *Casp3*^{-/-} dLGNs showed clear defects in the separation of eye-specific territories

(Fig. 3B) (*39*). To quantify the overlap between eye-specific territories in an unbiased

manner, we applied a set of increasing thresholds to signals from both eyes (Fig. S2) (*38*).

We calculated the percentage overlap of eye-specific territories as the ratio between

dLGN area with above-threshold signals from both eyes and the total dLGN area (Fig. S2

and Fig. 3C). Percentage overlap in *Casp3*^{-/-} dLGNs was consistently higher than that in

Casp3^{+/+} dLGNs across all thresholds (Fig. 3C and D). Importantly, the fold difference and

statistical significance of the difference in percentage overlap between the two groups of

animals increased as the cutoff threshold increased (Fig. 3D), suggesting the defects

observed in *Casp3*^{-/-} animals cannot be attributed to artifacts generated from background

noise.

As RGCs are known to undergo extensive apoptosis during the first week of

postnatal development (*40, 41*), we wondered if caspase-3 deficiency might block RGC

turnover and result in more RGC inputs in the dLGN, thereby confounding the overlap

analysis. To address this concern, we quantified the density of RGCs in whole-mount

retinae using a RGC marker, RBPMS (Fig. S6A) (*42*). Reassuringly, RGC densities in

retinae of *Casp3*^{+/+} and *Casp3*^{-/-} mice were not different (Fig. S6B), indicating that RGC

apoptosis still occurred at normal rates in the absence of caspase-3, presumably through

redundant mechanisms. Collectively, our findings suggest that caspase-3 is required for

normal segregation of eye-specific territories in dLGN.

Caspase-3 is required for retinogeniculate circuit refinement

Segregation of eye-specific territories in the mouse dLGN occurs prior to eye

opening (which happens at ~P13) and relies predominantly on spontaneous RGC activity

(*25*). After eye opening, sensory-dependent RGC activity drives a second phase of

refinement at the circuit level (*25*). At the time of eye opening, each dLGN relay neuron

receives numerous weak RGC inputs. With visual experience, a minority of these inputs

are strengthened while other weak inputs are eliminated, so that by P30 only a few

(typically ≤3) strong RGC inputs innervate each dLGN relay neuron (*25*).

To test if caspase-3 is required for visual experience-driven circuit refinement, we

prepared acute brain slices from P30 *Casp3*$^{+/+}$ and *Casp3*$^{-/-}$ littermate mice and

investigated the electrophysiological properties of retinogeniculate synapses (Fig. S7A)

(*43, 44*). We stimulated the optic tract while recording both AMPA receptor (AMPAR)- and

NMDA receptor (NMDAR)-mediated excitatory postsynaptic currents (EPSCs) in dLGN

relay neurons (Fig. S7A and Fig. 4A-B). Because each RGC axon has a distinct excitation

threshold, as we gradually increase the stimulation intensity on the optic tract, we recruit

individual RGC axons in a serial manner (Fig. S7A). Excitation of each RGC axon triggers

a step increase in the EPSC amplitude (Fig. S7A), allowing us to count the number of

steps in both AMPAR- and NMDAR-mediated response curves and infer the number of

RGC inputs on each relay neuron (Fig. 4C and S8A). In agreement with previous studies,

retinogeniculate circuits in *Casp3*$^{+/+}$ animals were refined, with dLGN relay neurons most

frequently receiving 2 strong RGC inputs, and the great majority of neurons receiving no

more than 3 inputs (Fig. 4A, 4C and S8A). In contrast, RGC input counts of dLGN relay

neurons in *Casp3*$^{-/-}$ animals followed a long-tailed distribution that was significant right-

shifted from that of *Casp3*$^{+/+}$ animals (Fig. 4C and S8A). The majority of *Casp3*$^{-/-}$ relay

neurons received more than 3 inputs (Fig. 4B-C and S8A), and many inputs elicited only

small increments in EPSC amplitude (Fig. 4B). To investigate whether the defects

observed in *Casp3*[-/-] animals could be due to general synaptic dysfunction, we compared

AMPAR- and NMDAR-mediated EPSCs in *Casp3*[+/+] and *Casp3*[-/-] animals and observed

no difference in maximum EPSC amplitude (Fig. S8B and S8C) or the ratio of the two

types of EPSCs (Fig. S8D). Additionally, we calculated fiber fraction (FF) as the ratio

between the first non-zero response (elicited by a single axon fiber) and the maximum

response in each dLGN relay neuron (Fig. S8E). FF increases as RGC inputs are

strengthened during development (*44*). We observed comparable FF values in *Casp3*[+/+]

and *Casp3*[-/-] animals (Fig. S8E), which suggests that defective circuit refinement in

Casp3[-/-] animals was not due to lack of RGC input strengthening but instead an inability

to eliminate weak inputs.

An increased number of RGC axons innervating dLGN relay neurons in *Casp3*[-/-]

mice should correspond to greater numbers of presynaptic release sites. To infer the

number of release sites, we stimulated the optic tract with a pair of stimuli in rapid

succession and measured the paired-pulse response ratio (PPR) in dLGN relay neurons

(Fig. S7B and Fig. 4D-E) (*43, 45*). In retinogeniculate synapses, the first stimulus

significantly depletes the readily releasable pool (RRP) of neurotransmitters and reduces

the response to the second stimulus, resulting in a small PPR (calculated as second

response over first response) (Fig. S7B) (*43, 45*). If there is an increase in the number of

release sites in *Casp3*[-/-] mice, we expect the RRP to expand, leading to a larger second

response and PPR (Fig. S7C) (*45*). Indeed, we found that the PPR was significantly

greater in *Casp3*[-/-] dLGN relay neurons than in controls (Fig. 4D-E), suggesting that the

inability to eliminate weak inputs resulted in an increase in the number of release sites

from RGC inputs in *Casp3*[-/-] mice. Taken together, our results demonstrate that caspase-

3 is required for experience-driven retinogeniculate circuit refinement. In the absence of

caspase-3, weak RGC inputs cannot be eliminated, and dLGN relay neurons remain

multiply innervated.

Microglia-mediated synapse elimination depends on caspase-3

Our results reveal activity-dependent caspase-3 activation at synapses as a key

molecular event that directs synapse elimination in the retinogeniculate pathway but do

not identify the cell types involved. Previous studies have demonstrated that both

microglia and astrocytes participate in synapse elimination by engulfing synaptic material

(*7, 10*). It is possible that caspase-3 activation at synapses might induce the display

apoptosis-related cell surface markers that can be recognized by microglia and astrocytes

for synapse elimination.

If microglia-mediated synapse elimination depends on caspase-3 activation, then

microglia in *Casp3*[-/-] mice should engulf less synaptic material. Similar to a previous study

(*7*), we utilized the *Cx3cr1-Gfp* transgenic mouse line to visualize microglia *in vivo.* In this

line, a *Gfp* transgene is expressed under the control of the native *Cx3cr1* promoter to

enable specific labeling of microglia in the brain (Fig. 5A) (*46*). As a proxy for labeling

retinogeniculate synapses, we labeled presynaptic RGC axon terminals by injecting

Casp3[+/-]; *Cx3cr1-Gfp*[+/-] and *Casp3*[-/-]; *Cx3cr1-Gfp*[+/-] littermate pups (*Casp3*[+/-] instead of

Casp3[+/+] mice were used to enable more efficient breeding) at the age of P4 with

fluorophore-conjugated CTB in the two eyes and harvested the brains at P5 (Fig. 5A) (*7*).

We used automatic segmentation to isolate volumes corresponding to microglia and RGC

axon terminals (Fig. 5B) from regions in the dLGN where RGC inputs from both eyes were

present (Fig. 5A). The presence of small soma and thin processes of microglia in *Casp3*⁻

$^{/-}$; *Cx3cr1-Gfp*$^{+/-}$ mice indicates that caspase-3 deficiency did not result in microglial

activation (Fig. 5B), which is further validated by immunohistochemical analysis of

microglia in caspase-3 deficient mice (Fig. S9A-D). To quantify synapse engulfment, we

calculated the volume of labeled RGC terminals that were fully enclosed in each microglia

(Fig. 5C). As expected, we observed that microglia in *Casp3*$^{-/-}$; *Cx3cr1-Gfp*$^{+/-}$ mice

engulfed significantly less axonal material compared to those of *Casp3*$^{+/-}$; *Cx3cr1-Gfp*$^{+/-}$

littermates (Fig. 5B-C). The same trend was observed when RGC axon terminals

originating from the two eyes were analyzed separately (Fig. S10A-B), suggesting that

defects in microglia-mediated synapse engulfment in *Casp3*$^{-/-}$ mice affect inputs from both

eyes. As the engulfment capacity of microglia is limited by their size, we normalized the

volume of engulfed synaptic material in each microglia to the microglial volume (Fig. 5D).

Microglia in both groups of mice are comparable in size (Fig. S10C), and the defect in

synaptic engulfment observed in *Casp3*$^{-/-}$ mice persisted after normalization to microglial

volume (Fig. 5D). Our results demonstrate that microglia-mediated synapse elimination

depends on caspase-3.

Astrocytes utilize phagocytic receptors MerTK and Megf10 to eliminate synapses

(*10*). Both receptors have been implicated in the clearance of apoptotic cells (*47, 48*). We

asked whether caspase-3 activation is also required for astrocyte-mediated synapse

elimination. To visualize astrocytes, we used the *Aldh1l1-Gfp* transgenic mouse line,

where a *Gfp* transgene under the control of the promoter of *Aldh1l1* is specifically

expressed in astrocytes (Fig. S11A) (*49*). We again labeled RGC axon terminals in

Casp3$^{+/-}$; *Aldh1l1-Gfp$^{+/-}$* and *Casp3$^{-/-}$*; *Aldh1l1-Gfp$^{+/-}$* littermate mice with intraocular

injection of CTB conjugates (Fig. S11A). Astrocytes were present in the P5 dLGN at a

much higher density than microglia and possess very fine processes (Fig. S11A). To

ensure segmentation fidelity, we only analyzed astrocyte cell bodies and their proximal

processes (Fig. S11B). Unexpectedly, we found that the volume of axonal material

engulfed by astrocytes was comparable in the two groups of mice (Fig. S11C). On

average, astrocytes in caspase-3 deficient mice did engulf less axonal material (Fig.

S11C), but the effect size appears too small to be biologically relevant, and the statistical

significance was small considering that the sample size was large (Fig. S11C). The same

trend was observed when we normalized the volume of engulfed synaptic material to

astrocyte volume (Fig. S11D-E). It is possible that a more pronounced difference would

be observed if *Casp3$^{+/+}$* mice were used for comparison. Nevertheless, the differential

response to caspase-3 deficiency between microglia- and astrocyte-mediated synapse

elimination raises the possibility that the two cell types may target distinct sets of

synapses for elimination (see discussion).

Caspase-3 activation determines synapse elimination specificity

Our data demonstrate that caspase-3 is activated at weak/inactive synapses, and

that microglia eliminate less synaptic material when caspase-3 activation is blocked.

Combining the two observations, we hypothesize that activity-dependent caspase-3

activation at synapses is a key molecular event that directs microglia to specifically

eliminate weaker synapses. This model predicts that in caspase-3 deficient mice,

microglia should lose the ability to preferentially target weaker synapses.

To test this model, we inactivated retinogeniculate synapses originating from right

eyes in $Casp3^{+/+}$ and $Casp3^{-/-}$ mice with AAV-hSyn-TeTxLC injections (Fig. 6A). A group

of control animals was included for each genotype that received AAV-hSyn-Turq injections.

We then labeled RGC axon terminals from both eyes with CTB conjugates in all animals

and analyzed microglia-mediated synapse engulfment (Fig. 6B-G). According to our

model, in TeTxLC-injected $Casp3^{+/+}$ mice, microglia should preferentially engulf inactive

retinogeniculate synapses originating from the right eye (Fig. 6A), whereas in TeTxLC-

injected $Casp3^{-/-}$ mice, synapses from the right and the left eye should be engulfed at

similar levels regardless of their strength (Fig. 6A). To quantitatively detect preferential

synapse engulfment by microglia, a comparison between TeTxLC-injected and control-

injected animals is necessary (Fig. 6B, C, and F). We first calculated the ratio between

the volume of right eye-originated RGC axon terminals and left eye-originated terminals

that were engulfed by each microglia. Then, by comparing this right-to-left engulfment

ratio computed in TeTxLC-injected and control-injected animals, preferential engulfment

of inactivated synapses from right eyes could be detected as higher engulfment ratios in

TeTxLC-injected animals (Fig. 6F). It is worth noting that microglia-related confounding

factors, such as possible alterations in microglia physiology due to caspase-3 deficiency

or surgical manipulations, cancel out when calculating engulfment ratios, allowing

information pertaining only to substrate preference to be isolated.

Consistent with previous studies (7), in $Casp3^{+/+}$ mice inactivation of

retinogeniculate synapses from the right eye led to enhanced microglia-mediated

engulfment of right eye-originated RGC axon terminals but not left eye-originated

terminals (Fig. S12A-B), resulting in a significant increase in right-to-left engulfment ratios

in microglia of TeTxLC-injected animals compared to control-injected animals (Fig. 6B, C,

and F). In contrast, engulfment ratios were similar between TeTxLC-injected and control-

injected $Casp3^{-/-}$ mice (Fig. 6D, E and G and Fig. S12C-D), suggesting that microglia

could no longer distinguish between strong and weak synapses in the absence of

caspase-3 activation. Summing the evidence obtained thus far, we propose the activity-

dependent caspase-3 activation is a crucial signaling event that integrates upstream

information about synaptic activity, takes place at weak/inactive synapses, and recruits

downstream microglia to eliminate synapses in an activity-dependent manner.

Caspase-3 deficiency protects against amyloid-β-induced synapse loss

Lastly, we wanted to test if caspase-3 regulates synapse loss in neurodegenerative

diseases through mechanisms analogous to its role in activity-dependent synapse

elimination. To this end, we focused on Alzheimer's disease (AD). AD is the most common

cause of dementia (*50*), and synapse loss is the strongest correlate of cognitive decline

in AD patients (*51*). Accumulation and deposition of oligomeric or fibril amyloid-

β (Aβ) peptide has been proposed to be the cause of neurodegeneration in AD (*50*).

Intriguingly, oligomeric Aβ impairs long-term potentiation (LTP) (*52*), and this suppression

of LTP requires caspase-3 activation (*53*). Additionally, caspase-3 activation has been

linked to early synaptic dysfunction in an AD mouse model (*54*).

To test if synapse loss in AD requires caspase-3, we utilized the APP/PS1 mouse

line, in which amyloid precursor protein (APP) and presenilin 1 (PS1) carrying mutations

associated with early-onset familial AD are overexpressed in CNS neurons (*55*). We

observed that female *App/Ps1+/-* mice developed amyloid deposits in the hippocampus

and cortex by 5 months (Fig. S13Aii), and the amyloid burden worsened by 6 months (Fig.

S13Aiii). In male *App/Ps1+/-* mice, amyloid deposits only became apparent at 6 months

and occurred at a lower level compared to age-matched females (Fig. S13Aiv-v). We

focused on analyzing *App/Ps1+/-* female mice in subsequent experiments to obtain robust

phenotypes. To quantify synapse loss, we labeled presynaptic and postsynaptic

compartments in the dentate gyrus of 6 month-old *App/Ps1-/-* and *App/Ps1+/-* littermates

by detecting synaptic vesicle protein 2 (SV2) and Homer1, respectively, and selected

multiple fields of interest per animal for imaging analysis while avoiding regions adjacent

to amyloid deposits (Fig. S14). We fitted ellipsoids to SV2 and Homer1 signals to identify

pre- and post-synaptic puncta and defined synapses as Homer1-SV2 punctum pairs that

were less than 300 nm apart (Fig. 7A). We observed that synapse density in *App/Ps1+/-*

mice was significantly reduced compared to that in *App/Ps1-/-* littermate controls (Fig. 7A-

B), and that this reduction could be attributed to lower densities of both pre- and post-

synaptic puncta in *App/Ps1+/-* mice (Fig. S15A-B). We then crossed APP/PS1 mice to

caspase-3 deficient mice and quantified synapse density in 6 month-old *Casp3-/-*;

App/Ps1-/- and *Casp3-/-*; *App/Ps1+/-* littermate mice (Fig. S14). We observed that, in the

caspase-3 deficient background, overexpression of mutant APP and PS1 no longer

resulted in a reduction in the density of synapses (Fig. 7C-D) or pre-/post-synaptic puncta

(Fig. S15C-D), suggesting that caspase-3 activity is required for Aβ-induced synapse loss

in the APP/PS1 mouse model. Intriguingly, levels of amyloid deposition were comparable

between *App/Ps1+/-* and *Casp3-/-*; *App/Ps1+/-* mice (Fig. S13Avi and B), and reactive

microglia were observed near amyloid deposits regardless of caspase-3 deficiency (Fig.

S16), indicating that protection conferred by caspase-3 deficiency against Aβ-induced

synapse loss might not require inhibition of amyloid accumulation or microgliosis. We also

investigated whether Aβ accumulation induced elevated levels of caspase-3 activity. We

analyzed 4, 5, and 6 month-old *App/Ps1*$^{-/-}$ and *App/Ps1*$^{+/-}$ mice but only observed robust

upregulation of caspase-3 activity in the dentate gyrus of a subset of 4 month-old

App/Ps1$^{+/-}$ mice (Fig. S17A-C). Upregulated caspase-3 activity in these mice remained in

a punctate pattern without inducing neuronal death (Fig. S17D-F). It is possible that Aβ

transiently induces caspase-3 activation in *App/Ps1*$^{+/-}$ mice, but stochasticity in the onset

and duration of such induction prevents reliable detection with our sample size and

temporal resolution. Overall, our results demonstrate caspase-3 as an important regulator

of Aβ-induced synapse loss and a potential therapeutic target for AD treatment.

Discussion:

A model for caspase-3 dependent synapse elimination

Our findings support a model where competition between strong/active synapses

and weak/inactive synapses in the retinogeniculate pathway triggers caspase-3 activation

in the postsynaptic compartments of weaker synapses. This activation signals microglia

to selectively eliminate these weaker synaptic connections, thereby facilitating the

establishment of mature circuits in the visual pathway.

Mechanisms of activity-dependent caspase-3 activation

In the experiment where we demonstrated activity-dependent caspase-3 activation,

we inactivated all retinogeniculate synapses from one eye to induce sufficiently high

caspase-3 activity that can be discerned from background. It is important to note that this

degree of inactivation is not typical for normal development. Consequently, caspase-3

activation in entire dLGN relay neurons observed in this experimental setting should not

be interpreted as a physiological outcome of activity-dependent synapse elimination. In

fact, apoptotic neurons were infrequently observed in the dLGN of control animals (Fig.

1D and 2B), suggesting that during normal development, activity-dependent caspase-3

activation is predominantly localized and generally does not lead to neuronal death.

Previous studies have proposed ubiquitin-dependent proteasomal degradation as a

mechanism that can restrain caspase-3 activity to local sites in dendrites (*24, 56, 57*).

Intriguingly, synaptic inactivation in our experiment was achieved through TeTxLC

expression in presynaptic RGCs, but caspase-3 activation was observed in postsynaptic

relay neurons. Conversely, while caspase-3 is activated in the postsynaptic compartment,

we were able to detect defects in synapse elimination using assays that measure

presynaptic inputs. These observations suggest that activity-dependent caspase-3

activation requires synaptic transmission, and that caspase-3 dependent synapse

elimination is not strictly axonal or dendritic but involves both pre- and post-synaptic

compartments. An important limitation of this study is that experiments in full-body *Casp3*[-]

[/-] mice cannot query the roles of caspase-3 in presynaptic and postsynaptic compartments

separately. Future experiments that perturb caspase-3 activation and other signaling

events specifically in the pre- or post-synaptic compartments are needed to reveal more

detailed mechanisms of activity-dependent synapse elimination.

The observation that activity-dependent caspase-3 activation requires synaptic

competition resonates with previous research demonstrating that relative synaptic

efficacy biases the outcome of synapse elimination during development of neuromuscular

junctions and the callosal pathway (*15, 33, 58, 59*). However, the mechanisms linking

synaptic competition with caspase-3 activation remains elusive. During LTD, caspase-3

is activated through mitochondrial release of cytochrome-c (*18*). Whether synaptic

competition could induce caspase-3 activation via mechanisms such as calcium signaling,

which regulates both neurotransmission and cytochrome-c release (*60*), is an outstanding

question.

Roles of microglia and astrocytes in synapse elimination

The differential impact of caspase-3 deficiency on microglia- and astrocyte-

mediated synapses elimination in the visual pathway suggest that the two glial cell types

have nonidentical substrate specificities when engulfing synaptic material. Indeed,

considering that astrocytes are present at nearly ten-fold greater density than microglia

in the developing dLGN (Fig. 5A and S11A) and have comparable engulfing capacities

(Fig. 5D and S11D), contribution from microglia in synapse elimination would be negligible

if the two cell types target the same substrates, which contradicts previous findings (*7,*

14). A recent study discovered that during clearance of apoptotic neurons in adult brains,

microglia engulf the soma and proximal dendrites while astrocytes target distal dendrites,

and this difference in substrate preference is at least partially attributed to differential

kinetics in binding and engulfing substrates by the two cell types (*61*). Investigating

whether a similar division-of-labor exists during development could provide critical

insights into the mechanisms of activity-dependent, glia-mediated synapse elimination.

It is often assumed in the literature that microglia directly engulf intact synapses,

but existing data also support an alternative model where microglia indirectly contribute

to synapse elimination by engulfing synaptic debris that is shed through neuron-

autonomous mechanisms (*62*). The observation that synaptic activity instructs microglial

engulfment through caspase-3 activation raises the possibility that weakened synapses

partially disintegrate through apoptosis-like mechanisms prior to being engulfed by

microglia. Further studies are needed to elucidate the intermediate steps between

caspase-3 activation and microglial engulfment of synapses.

Molecular link between caspase-3 activation and microglial engulfment

How does caspase-3 activation at synapses recruit microglia for engulfment?

Caspase-3 activation is known to trigger the display of apoptosis markers that can be

recognized by phagocytes (*19*). One of the best understood markers is

phosphatidylserine (PtdSer), a phospholipid that translocates from the cytoplasmic leaflet

of the plasma membrane to the exoplasmic leaflet during apoptosis (*19*). Importantly,

caspase-3 directly regulates the enzymes that translocate PtdSer through enzymatic

cleavage (*63, 64*). Two recent studies demonstrated that PtdSer is displayed on

retinogeniculate synapses during postnatal development, and microglia use the exposed

PtdSer as a cue to target and engulf synapses (*8, 9*). Whether PtdSer exposure is

upregulated in response to synapse weakening and whether caspase-3 activity is

required for such exposure are important questions that require further investigation.

The complement protein C1q has been observed to bind to PtdSer on apoptotic

cells and facilitate phagocytic clearance (*19*). Additionally, a previous study found

caspase-3 activity in C1q-bound synaptosomes (*65*), suggesting that C1q may bridge

caspase-3 activation and synapse elimination. However, we investigated C1q expression

and localization in the dLGN of P5 TeTxLC-injected mice but did not observe upregulation

of C1q abundance or colocalization between C1q and caspase-3 activity (data not shown).

Caspase-3 as a potential target for AD treatment

Our findings suggest that caspase-3 deficiency confers protection against Aβ-

induced synapse loss in a mouse model of AD. Notably, the absence of caspase-3 does

not seem to affect Aβ deposition or microglia reactivity, indicating that caspase-3 depletion

may mitigate synapse loss by directly countering Aβ-induced synaptic toxicity. Given the

prominence of Aβ and inflammation as targets for AD drug development (*66*), inhibiting

caspase-3 activity offers a potential alternative therapeutic strategy that could

complement and enhance current treatment approaches. Therefore, it is important to test

whether pharmacological inhibition of caspase-3 can effectively slow synapse loss and

cognitive decline in both mouse models of AD and human patients.

Acknowledgement: We thank Crystall Lopez, Benjamin Gantz, and Brooke Groff for help

with animal husbandry and breeding. We thank Sarah Lindo, Colin Morrow, Claire Boyer,

and Rae Demars for help with surgery. We thank Monique Copeland, Amy Hu, Morgan

Clarke, and Benjamin Foster for help with histology. We are also grateful to Sarah

Kivimaki and the Viral Tools team of Janelia Core Services for help with AAV preparation.

We thank Dr. Cagla Eroglu, Dr. Richard Axel, and Dr. David Clapham for comments on

the manuscript. This work is supported by the Howard Hughes Medical Institute.

References:

1.   M. Kano, K. Hashimoto, Synapse elimination in the central nervous system. *Curr Opin*
Neurobiol **19**, 154-161 (2009).
2.   T. E. Faust, G. Gunner, D. P. Schafer, Mechanisms governing activity-dependent synaptic
pruning in the developing mammalian CNS. *Nat Rev Neurosci* **22**, 657-673 (2021).
3.   M. M. Riccomagno, A. L. Kolodkin, Sculpting neural circuits by axon and dendrite pruning.
Annu Rev Cell Dev Biol **31**, 779-805 (2015).
4.   L. C. Katz, C. J. Shatz, Synaptic activity and the construction of cortical circuits. *Science* **274**,
1133-1138 (1996).
5.   J. Y. Hua, S. J. Smith, Neural activity and the dynamics of central nervous system
development. *Nat Neurosci* **7**, 327-332 (2004).
6.   B. Stevens *et al.*, The classical complement cascade mediates CNS synapse elimination.
Cell **131**, 1164-1178 (2007).
7.   D. P. Schafer *et al.*, Microglia sculpt postnatal neural circuits in an activity and
complement-dependent manner. *Neuron* **74**, 691-705 (2012).
8.   T. Li *et al.*, A splicing isoform of GPR56 mediates microglial synaptic refinement via
phosphatidylserine binding. *Embo J* **39**, e104136 (2020).
9.   N. Scott-Hewitt *et al.*, Local externalization of phosphatidylserine mediates developmental
synaptic pruning by microglia. *Embo J* **39**, e105380 (2020).
10.   W. S. Chung *et al.*, Astrocytes mediate synapse elimination through MEGF10 and MERTK
pathways. *Nature* **504**, 394-400 (2013).
11.   D. K. Wilton, L. Dissing-Olesen, B. Stevens, Neuron-Glia Signaling in Synapse Elimination.
Annu Rev Neurosci **42**, 107-127 (2019).
12.   W. S. Chung, N. J. Allen, C. Eroglu, Astrocytes Control Synapse Formation, Function, and
Elimination. *Cold Spring Harb Perspect Biol* **7**, a020370 (2015).
13.   Q. Li, B. A. Barres, Microglia and macrophages in brain homeostasis and disease. *Nat Rev*
Immunol **18**, 225-242 (2018).
14.   R. C. Paolicelli *et al.*, Synaptic pruning by microglia is necessary for normal brain
development. *Science* **333**, 1456-1458 (2011).
15.   M. Yasuda, S. Nagappan-Chettiar, E. M. Johnson-Venkatesh, H. Umemori, An activity-
dependent determinant of synapse elimination in the mammalian brain. *Neuron* **109**,
1333-1349 e1336 (2021).
16.   R. L. Philips *et al.*, The JAK-STAT pathway at 30: Much learned, much more to do. *Cell* **185**,
3857-3876 (2022).
17.   D. R. McIlwain, T. Berger, T. W. Mak, Caspase functions in cell death and disease. *Cold*
Spring Harb Perspect Biol **5**, a008656 (2013).

18.   Z. Li *et al.*, Caspase-3 Activation via Mitochondria Is Required for Long-Term Depression
and AMPA Receptor Internalization. *Cell* **141**, 859-871 (2010).

19.   S. Y. Park, I. S. Kim, Engulfment signals and the phagocytic machinery for apoptotic cell
clearance. *Exp Mol Med* **49**, (2017).

20.   L. A. Trouw, A. M. Blom, P. Gasque, Role of complement and complement regulators in the
removal of apoptotic cells. *Mol Immunol* **45**, 1199-1207 (2008).

21.   D. W. Williams, S. Kondo, A. Krzyzanowska, Y. Hiromi, J. W. Truman, Local caspase activity
directs engulfment of dendrites during pruning. *Nat Neurosci* **9**, 1234-1236 (2006).

22.   C. T. Kuo, S. Zhu, S. Younger, L. Y. Jan, Y. N. Jan, Identification of E2/E3 ubiquitinating
enzymes and caspase activity regulating Drosophila sensory neuron dendrite pruning.
Neuron **51**, 283-290 (2006).

23.   D. J. Simon *et al.*, A caspase cascade regulating developmental axon degeneration. *J
Neurosci* **32**, 17540-17553 (2012).

24.   A. Erturk, Y. Wang, M. Sheng, Local pruning of dendrites and spines by caspase-3-
dependent and proteasome-limited mechanisms. *J Neurosci* **34**, 1672-1688 (2014).

25.   L. Liang, C. Chen, Organization, Function, and Development of the Mouse
Retinogeniculate Synapse. *Annu Rev Vis Sci* **6**, 261-285 (2020).

26.   A. A. Penn, P. A. Riquelme, M. B. Feller, C. J. Shatz, Competition in retinogeniculate
patterning driven by spontaneous activity. *Science* **279**, 2108-2112 (1998).

27.   D. Stellwagen, C. J. Shatz, An instructive role for retinal waves in the development of
retinogeniculate connectivity. *Neuron* **33**, 357-367 (2002).

28.   M. Yasuda, S. Nagappan-Chettiar, H. Umemori, In utero intraocular AAV injection for early
gene expression in the developing rodent retina. *STAR Protoc* **2**, 100742 (2021).

29.   G. Schiavo *et al.*, Tetanus and botulinum-B neurotoxins block neurotransmitter release by
proteolytic cleavage of synaptobrevin. *Nature* **359**, 832-835 (1992).

30.   T. E. Krahe, R. N. El-Danaf, E. K. Dilger, S. C. Henderson, W. Guido, Morphologically distinct
classes of relay cells exhibit regional preferences in the dorsal lateral geniculate nucleus
of the mouse. *J Neurosci* **31**, 17437-17448 (2011).

31.   J. G. Parnavelas, E. J. Mounty, R. Bradford, A. R. Lieberman, The postnatal development of
neurons in the dorsal lateral geniculate nucleus of the rat: a Golgi study. *J Comp Neurol*
171, 481-499 (1977).

32.   S. McComb *et al.*, Efficient apoptosis requires feedback amplification of upstream
apoptotic signals by effector caspase-3 or -7. *Sci Adv* **5**, eaau9433 (2019).

33.   S. Nagappan-Chettiar, M. Yasuda, E. M. Johnson-Venkatesh, H. Umemori, The molecular
signals that regulate activity-dependent synapse refinement in the brain. *Curr Opin
Neurobiol* **79**, 102692 (2023).

34.   F. M. Rossi *et al.*, Requirement of the nicotinic acetylcholine receptor beta 2 subunit for
the anatomical and functional development of the visual system. *Proc Natl Acad Sci U S A*
98, 6453-6458 (2001).

35.   K. Kuida *et al.*, Decreased apoptosis in the brain and premature lethality in CPP32-
deficient mice. *Nature* **384**, 368-372 (1996).

36.   C. Houde *et al.*, Caspase-7 expanded function and intrinsic expression level underlies
strain-specific brain phenotype of caspase-3-null mice. *J Neurosci* **24**, 9977-9984 (2004).

37.   L. De Haan, T. R. Hirst, Cholera toxin: a paradigm for multi-functional engagement of
585   cellular mechanisms (Review). *Mol Membr Biol* **21**, 77-92 (2004).
586   38.   C. L. Torborg, M. B. Feller, Unbiased analysis of bulk axonal segregation patterns. *J Neurosci*
587   *Meth* **135**, 17-26 (2004).
588   39.   N. Otsu, A Threshold Selection Method from Gray-Level Histograms. *IEEE Transactions on*
589   *Systems, Man, and Cybernetics* **9**, 62-66 (1979).
590   40.   R. W. Young, Cell death during differentiation of the retina in the mouse. *J Comp Neurol*
591   **229**, 362-373 (1984).
592   41.   B. Dreher, R. A. Potts, M. R. Bennett, Evidence that the early postnatal reduction in the
593   number of rat retinal ganglion cells is due to a wave of ganglion cell death. *Neurosci Lett*
594   **36**, 255-260 (1983).
595   42.   A. R. Rodriguez, L. P. de Sevilla Muller, N. C. Brecha, The RNA binding protein RBPMS is a
596   selective marker of ganglion cells in the mammalian retina. *J Comp Neurol* **522**, 1411-1443
597   (2014).
598   43.   C. Chen, W. G. Regehr, Developmental remodeling of the retinogeniculate synapse.
599   *Neuron* **28**, 955-966 (2000).
600   44.   B. M. Hooks, C. Chen, Distinct roles for spontaneous and visual activity in remodeling of
601   the retinogeniculate synapse. *Neuron* **52**, 281-291 (2006).
602   45.   W. G. Regehr, Short-term presynaptic plasticity. *Cold Spring Harb Perspect Biol* **4**, a005702
603   (2012).
604   46.   S. Jung *et al.*, Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and
605   green fluorescent protein reporter gene insertion. *Mol Cell Biol* **20**, 4106-4114 (2000).
606   47.   T. Iram *et al.*, Megf10 Is a Receptor for C1Q That Mediates Clearance of Apoptotic Cells by
607   Astrocytes. *J Neurosci* **36**, 5185-5192 (2016).
608   48.   G. Lemke, Biology of the TAM receptors. *Cold Spring Harb Perspect Biol* **5**, a009076 (2013).
609   49.   S. Gong *et al.*, A gene expression atlas of the central nervous system based on bacterial
610   artificial chromosomes. *Nature* **425**, 917-925 (2003).
611   50.   D. S. Knopman *et al.*, Alzheimer disease. *Nat Rev Dis Primers* **7**, 33 (2021).
612   51.   S. Meftah, J. Gan, Alzheimer's disease as a synaptopathy: Evidence for dysfunction of
613   synapses during disease progression. *Front Synaptic Neurosci* **15**, 1129036 (2023).
614   52.   G. M. Shankar *et al.*, Amyloid-beta protein dimers isolated directly from Alzheimer's brains
615   impair synaptic plasticity and memory. *Nat Med* **14**, 837-842 (2008).
616   53.   J. Jo *et al.*, Abeta(1-42) inhibition of LTP is mediated by a signaling pathway involving
617   caspase-3, Akt1 and GSK-3beta. *Nat Neurosci* **14**, 545-547 (2011).
618   54.   M. D'Amelio *et al.*, Caspase-3 triggers early synaptic dysfunction in a mouse model of
619   Alzheimer's disease. *Nat Neurosci* **14**, 69-76 (2011).
620   55.   J. L. Jankowsky *et al.*, Mutant presenilins specifically elevate the levels of the 42 residue
621   beta-amyloid peptide in vivo: evidence for augmentation of a 42-specific gamma secretase.
622   *Hum Mol Genet* **13**, 159-170 (2004).
623   56.   Y. Suzuki, Y. Nakabayashi, R. Takahashi, Ubiquitin-protein ligase activity of X-linked
624   inhibitor of apoptosis protein promotes proteasomal degradation of caspase-3 and
625   enhances its anti-apoptotic effect in Fas-induced cell death. *Proc Natl Acad Sci U S A* **98**,
626   8662-8667 (2001).

57. Y. E. Choi *et al.*, The E3 ubiquitin ligase cIAP1 binds and ubiquitinates caspase-3 and -7 via
unique mechanisms at distinct steps in their processing. *J Biol Chem* **284**, 12772-12782
(2009).
58. M. Buffelli *et al.*, Genetic evidence that relative synaptic efficacy biases the outcome of
synaptic competition. *Nature* **424**, 430-434 (2003).
59. Y. I. Lee, Developmental neuromuscular synapse elimination: Activity-dependence and
potential downstream effector mechanisms. *Neurosci Lett* **718**, 134724 (2020).
60. C. Garrido *et al.*, Mechanisms of cytochrome c release from mitochondria. *Cell Death*
Differ **13**, 1423-1433 (2006).
61. E. C. Damisah *et al.*, Astrocytes and microglia play orchestrated roles and respect
phagocytic territories during neuronal corpse removal in vivo. *Sci Adv* **6**, eaba3239 (2020).
62. U. Eyo, A. V. Molofsky, Defining microglial-synapse interactions. *Science* **381**, 1155-1156
(2023).
63. J. Suzuki, D. P. Denning, E. Imanishi, H. R. Horvitz, S. Nagata, Xk-Related Protein 8 and CED-
8 Promote Phosphatidylserine Exposure in Apoptotic Cells. *Science* **341**, 403-406 (2013).
64. K. Segawa *et al.*, Caspase-mediated cleavage of phospholipid flippase for apoptotic
phosphatidylserine exposure. *Science* **344**, 1164-1168 (2014).
65. B. A. Gyorffy *et al.*, Local apoptotic-like mechanisms underlie complement-mediated
synaptic pruning. *Proc Natl Acad Sci U S A* **115**, 6303-6308 (2018).
66. P. van Bokhoven *et al.*, The Alzheimer's disease drug development landscape. *Alzheimers*
Res Ther **13**, 186 (2021).

659 **Figures:**

660
Figure 1. Inactivation of retinogeniculate synapses induces caspase-3 activity. (**A**) Schematics
of experimental setup. AAVs expressing tetanus toxin light chain (TeTxLC) and/or mTurquiose2 (Turq)
were injected into the right eye of E15 mice (left). By P5, retinogeniculate synapses in dLGN were
inactivated to varying extents depending on injection and side (right). (**B-D**) Confocal images of Turq
(left panels) and activated caspase-3 (right panels) in left dLGN (**B**) and right dLGN (**C**) of a TeTxLC-
injected P5 animal and in left dLGN of a control P5 animal (**D**). Images from the same fluorescent
channel were adjusted to the same contrast. Dotted lines delineate dLGN boundaries. The compass
in **B** marks tissue orientation. Scale-bars: 100 μm. D, dorsal; V, ventral; M, medial; L, lateral. (**E**)
Quantification of caspase-3 activity in indicated dLGNs. Activated caspase-3 signals in each dLGN
(highlighted areas in **B-D**) were summed and normalized to dLGN area. Each point represents the
result from one dLGN. Data from two dLGNs of the same animal were paired for analysis (grey lines).
n=10 for TeTxLC-injected animals and n=9 for control animals. Mean and standard deviation (S.D.)
are shown. P-values were calculated from two-tailed t-tests (paired when applicable). (**F**) Example
images showing punctate caspase-3 activities in ventral-medial regions of indicated dLGNs. Images
were adjusted to the same contrast. Scale-bar: 20 μm. (**G**) High-resolution images of dLGN showing
TeTxLC-expressing RGC axons (yellow) and activated caspase-3 (magenta). Two regions of interest
(dotted squares) are magnified to illustrate that caspase-3 activity was found juxtaposing TeTxLC-
expressing axon terminals but not within them. Scale-bar: 5 μm.

Figure 2. Synapse inactivation-induced caspase-3 activation requires synaptic competition. (**A**) Schematics illustrating different experimental conditions. Presynaptic (green) and postsynaptic (cyan) compartments and synaptic strength (yellow arrowheads) of retinogeniculate synapses are shown. No synapses are inactivated in the control (left); expressing TeTxLC only in right eye inputs results in synapse inactivation and competition (middle); expressing TeTxLC in both eyes results in synapse inactivation but no competition (right). (**B**) Confocal images of dLGNs in the three conditions showing activated caspase-3 signal. The dLGN shown for the single inactivation condition was from the left side. Dotted lines mark dLGN boundaries. Scale-bars: 100 μm. (**C**) Quantification of caspase-3 activity in the dLGN in the indicated conditions. Activated caspase-3 signal in each dLGN were summed and normalized to the dLGN area. For the single inactivation condition, values were from the left dLGN only. For the other two conditions, values from both dLGNs were averaged. n=8 animals for control, n=11 animals for single inactivation, and n=10 animals for dual inactivation. Mean and S.D. are shown. P-values were calculated from Tukey's multiple comparison tests.

717
Figure 3. Caspase-3 is required for segregation of eye-specific territories. (**A**-**B**) Representative
confocal images of retinogeniculate inputs in the dLGN of P10 wild-type (**A**) and *Casp3*[-/-] (**B**) mice.
Contralateral inputs are labeled with AlexaFlour488 (AF488) conjugated cholera toxin subunit B (CTB)
and ipsilateral inputs with AF594-CTB. Original images were thresholded into 0-or-1 images using the
Otsu method (*34*), and the overlap between thresholded contralateral and ipsilateral inputs is shown.
(**C**) Percentage overlap between eye-specific territories in wildtype and *Casp3*[-/-] mice under a series
of increasing signal cutoff thresholds. Note that the percentage overlap is plotted on a log scale. Each
circle represents one animal. Mean and S.D. are shown. n=9 for wildtype mice and n=6 for *Casp3*[-/-]
mice. (**D**) Mean percentage overlap values in wildtype and *Casp3*[-/-] mice and p-values of two-tailed t-
tests between the two genotypes are listed for each cutoff threshold.

741
Figure 4. Caspase 3 is required for retinogeniculate circuit refinement. (**A**-**B**) Example recordings
of dLGN relay neuron responses in P30 wildtype (**A**) and *Casp3*[-/-] (**B**) mice. Excitatory postsynaptic
currents (EPSCs) were evoked by increasing stimulation currents in the optic tract. Both AMPAR-
mediated inward current at -70 mV membrane potential and AMPAR and NMDAR-mediated outward
current at +40 mV membrane potential are shown. Peak response amplitudes at each stimulation
intensity are plotted to the right of recording traces. Scale bars represent 0.5 nA and 10 ms. (**C**)
Distribution of RGC input numbers on individual dLGN relay neurons in wildtype and *Casp3*[-/-] mice.
The number of RGC inputs was inferred by manually counting the number of steps in AMPAR-
mediated EPSC response curves (lower right in **A** and **B**) while blind to the genotypes. P-value was
calculated from two-tailed t-test. n=37 cells for wildtype mice and n=32 cells for *Casp3*[-/-] mice. (**D**)
Example recordings from paired pulse measurements at -70 mV membrane potential in wildtype and
Casp3[-/-] mice. Traces from experiments with 50, 150, 250, 500, and 1000 ms inter-stimulus intervals
are overlayed. Stimulus artifacts were removed from the traces for clarity. Scale-bars represent 100
ms (horizontal) and 0.3 nA (vertical). (**E**) Paired-pulse ratio (calculated as peak amplitude of the second
response over that of the first response) in wildtype and *Casp3*[-/-] mice at various inter-stimulus intervals.
Mean and standard error of the mean (SEM) are shown. P-values were calculated from Bonferroni's
multiple comparison test. p=0.0067 for 50 ms interval, p=0.0369 for 150 ms interval, and p=0.0097 for
250 ms interval. n=16 cells for wildtype mice and n=13 for *Casp3*[-/-] mice.

Figure 5. Microglia-mediated synapse elimination depends on caspase-3. (**A**) Representative 3D-reconstructed images of a P5 *Casp3*$^{+/-}$; *Cx3cr1-Gfp*$^{+/-}$ mouse dLGN with microglia displayed in green, contralateral RGC axon terminals in red, and ipsilateral RGC terminals in blue. In the merged image, the region from which microglia are selected for analysis is indicated with the dashed line. The scale-bar represents 100 μm. (**B**) Representative surface rendering of microglia (green) from P5 dLGNs of *Casp3*$^{+/-}$; *Cx3cr1-Gfp*$^{+/-}$ and *Casp3*$^{-/-}$; *Cx3cr1-Gfp*$^{+/-}$ mice. Intracellular contralateral (red) and ipsilateral (blue) RGC axon terminals are shown. Microglia from caspase-3 deficient mice engulf visibly less synaptic material. Scale-bars represent 10 μm. (**C**) Total volume of engulfed synaptic material in individual microglia from *Casp3*$^{+/-}$; *Cx3cr1-Gfp*$^{+/-}$ and *Casp3*$^{-/-}$; *Cx3cr1-Gfp*$^{+/-}$ mice. (**D**) Total volume of engulfed synaptic material in each microglia (from **C**) is normalized to the volume of that microglia. In **C**-**D**, each point represents one microglia. Mean and S.D. are shown. p-values were calculated from unpaired two-tailed t-tests. n=61 microglia from 8 *Casp3*$^{+/-}$; *Cx3cr1-Gfp*$^{+/-}$ mice and n=54 microglia from 5 *Casp3*$^{-/-}$; *Cx3cr1-Gfp*$^{+/-}$ mice.

Figure 6. Caspase-3 activation determines specificity of microglia-mediated synapse elimination. (**A**) Schematics illustrating the experimental rationale. In wildtype mice (upper panel), inactivating retinogeniculate synapses from the right eye activates caspase-3 (magenta pie) and recruits microglia to engulf more right eye-originated synapses (red dots) than left eye-originated synapses (green dots). If caspase-3 activation is blocked (lower panel), we expect preferential engulfment of inactive synapses to be attenuated. (**B**-**E**) Surface rendering of representative microglia from P5 left dLGN of *Casp3*[+/+] (**B**-**C**) or *Casp3*[-/-] (**D**-**E**) mice injected with AAV expressing mTurquoise2 (**B** and **D**) or TeTxLC (**C** and **E**) in the right eye at E15. RGC axon terminals from the left eye are shown in yellow and terminals from the right eye in red. Scale bars represent 15 μm. (**F**-**G**) Ratio between volumes of right-eye and left-eye-originated synaptic material engulfed by microglia from *Casp3*[+/+] (**F**) or *Casp3*[-/-] (**G**) mice injected with AAV expressing mTurquoise2 (blue) or TeTxLC (red). Each dot represents one microglia. Engulfment ratios are displayed on a log scale. 0, 25, 50, 75, and 100 percentiles are shown. p-values were calculated from unpaired two-tailed Mann-Whitney tests. n=52 microglia from 7 Turq-injected *Casp3*[+/+] mice, n=50 microglia from 8 TeTxLC-injected *Casp3*[+/+] mice, n=64 microglia from 5 Turq-injected *Casp3*[-/-] mice, and n=51 microglia from 6 TeTxLC-injected *Casp3*[-/-] mice.

833
Figure 7. Caspase-3 deficiency protects against Aβ induced synapse loss. (**A** and **C**)
Representative 3D-reconstructed images (3 μm in z) showing presynaptic (SV2, in green) and
postsynaptic (Homer1, in red) signals in dentate gyrus of female 6 month-old APP/PS1 mice on
caspase-3 wildtype (**A**) and deficient (**C**) backgrounds. Ellipsoids were fitted to the original images
(upper panels) to isolate pre- (green) and post-synaptic (red) puncta (lower panels). Homer1 ellipsoids
found with 300 nm of a SV2 ellipsoid are highlighted in white in the fitted images. Original images are
adjusted to the same contrast. For the fitted images, only ellipsoids from the upper half of the z-stack
are shown. Scale-bar represents 4 μm. (**B** and **D**) Quantification of synapse density in APP/PS1 mice
on caspase-3 wildtype (**B**) and deficient (**D**) backgrounds. Mean and S.D. are shown. p-values were
calculated from unpaired two-tailed t-tests. n=5 for *App/Ps1*[-/-] mice, n=6 for *App/Ps1*[+/-] mice, n=6 for
Casp3[-/-]; *App/Ps1*[-/-] mice, and n=4 for *Casp3*[-/-]; *App/Ps1*[+/-] mice.
845

Figure S1. Segregation of eye-specific territories in the mouse retinogeniculate pathway. In the mouse retinogeniculate pathway, retinal ganglion cells (RGCs) in the retina of each eye innervate relay neurons in both the contralateral (opposite side as the originating RGC) dorsal lateral geniculate nucleus (dLGN) and the ipsilateral (same side from the originating RGC) dLGN to form retinogeniculate synapses (upper right). Within each dLGN, the majority of retinogeniculate synapses receive inputs from the contralateral eye, while the minority receive inputs from the ipsilateral eye (upper right). At the age of P3, regions in each dLGN receiving inputs from each of the two eyes overlap significantly (left). Through a process that requires synapse elimination and spontaneous RGC activity, these regions are refined into non-overlapping eye-specific territories by the age of P8 (upper right). If the refinement process is defective because neural activity or synapse elimination is disrupted, eye-specific territories fail to completely segregate, and regions innervated by the two eyes remain overlapping (lower right).

Figure S2. Quantifying eye-specific segregation with multi-threshold overlap analysis. For each dLGN, RGC inputs from the two eyes were imaged using separate fluorescence channels. A small area in the thalamus outside of each dLGN was chosen, and average signal intensity in each channel within that area was calculated and used as background (upper panel). For each channel, background was subtracted, and signals were normalized to between 0 and 1. To calculate overlap between eye-specific territories, a threshold, x, was chosen between 0 and 1 and applied to both channels (middle panel). The overlap between eye-specific territories were defined as the set of pixels with above-threshold signals in both channels. Percentage overlap was then calculated as the ratio between the area of the dLGN where eye-specific territories overlapped and the total area of the dLGN (lower panel). To avoid introducing biases by artificially selecting one threshold, we repeated the analysis with a set of increasingly stringent thresholds (e.g., from 0.1 to 0.25 in Fig. 3C and 3D).

A Tomato (L)　Turq (R)

B Tomato (L)　TeTxLC+Turq (R)

C

D

Threshold (%)	Control mean	TeTxLC mean	P-value
5.0	0.2172	0.2839	0.135
7.5	0.1045	0.1809	0.00601
10.0	0.05357	0.1187	0.00089
12.5	0.02736	0.07447	0.00014
15.0	0.01427	0.04548	0.000022
17.5	0.006909	0.02745	0.000005
20.0	0.003429	0.01568	0.000003

Figure S3. Blocking spontaneous RGC activity in one eye with TeTxLC disrupts eye-specific segregation in the retinogeniculate pathway. (**A** and **B**) Confocal images of P8 left dLGN (left panels) and right dLGN (right panels) of an animal receiving control injections (**A**) and an animal receiving TeTxLC injection in the right eye (**B**). Eye-specific regions were labeled with tdTomato (red, left eye input) and mTurquiose2 (cyan, right eye input). Right eye territory (cyan) contracted and left eye territory (red) expanded in TeTxLC-injected animals (**B**) compared to controls (**A**). Scale-bars: 100 μm. (**C**) Quantification of overlap between eye-specific territories in dLGN of P8 control and TeTxLC-injected animals. Analysis was done at multiple thresholds to avoid biases introduced with threshold selection. Note that percentage overlap is displayed on a log scale. For details on the analysis, see Fig. S2 and text associated with Fig. 3. n=7 for control animals and 5 for TeTxLC-injected animals. Mean and S.D. are shown. (**D**) Statistics of the overlap analysis at multiple thresholds. Fold difference and the statistical significance of the difference in percentage overlap between the two groups of animals increased as threshold increased. P-values were calculated from two-tailed t-tests. We did not implement multiple comparison corrections as values at different thresholds are derived from the same dataset and are not independent.

Figure S4. Inactivation of retinogeniculate synapses induces postsynaptic caspase-3 activity in dendritic compartments of dLGN relay neurons. (**A**) High-resolution images of three representative field-of-views (**i-iii**) in P5 TeTxLC-expressing dLGNs showing co-localization of punctate caspase-3 activity (magenta) and a dendritic marker, MAP2 (green). Dotted crosses mark x and y positions where the x-z and y-z cross-sections were generated. Scale-bars: 2 μm. (**B**) Example images showing caspase-3 activity in entire neurons (**i-iii**) or in multiple dendritic branches (but not the soma) of a neuron (**iv**) in P5 TeTxLC-expressing dLGNs. The neurons positive for active caspase-3 (**i-iii**) have relatively large and round somas and multipolar dendritic arbors that are characteristic of dLGN relay neurons. Scale-bars: 20 μm. (**C**) Example images of four representative neurons (**i-iv**) in P5 TeTxLC-expressing dLGNs that are positive for active caspase-3 (magenta) and a neuronal nuclear marker, NeuN (green). Scale-bars: 10 μm.

A Tomato (L) GFP (R)

B TeTxLC-Tomato (L) TeTxLC-GFP (R)

C

Percentage overlap (log scale) vs Threshold (%)

○ Control
○ Dual inactivation

D

Threshold (%)	Control mean	Dual inact. mean	P-value
5.0	0.1043	0.1732	0.005
7.5	0.0508	0.0940	0.0024
10.0	0.0265	0.0512	0.009
12.5	0.0131	0.0294	0.0044
15.0	0.0067	0.0177	0.0086
17.5	0.0033	0.0104	0.012
20.0	0.0018	0.0064	0.0072

Figure S5. Blocking spontaneous RGC activity in both eyes with TeTxLC disrupts eye-specific segregation in the retinogeniculate pathway. (**A** and **B**) Confocal images of the left dLGN (left panels) and right dLGN (right panels) of an animal receiving control injections in both eyes (**A**) and an animal receiving TeTxLC injection in both eyes (**B**). Eye-specific territories were labeled with tdTomato (magenta, left eye input) and eGFP (green, right eye input). In animals injected with TeTxLC in both eyes, territories of the two eyes are similar in size and are more dispersive compared to those in controls, leading to substantial overlap between eye-specific territories. Scale-bars: 100 μm. (**C**) Quantification of overlap between eye-specific territories in the dLGN of control and dual TeTxLC-injected animals. Analysis was done at multiple thresholds to avoid biases introduced with threshold selection. Note that percentage overlap is displayed on a log scale. For details on the analysis, see Fig. S2 and text associated with Fig. 3. n=8 for control animals and n=9 for dually TeTxLC-injected animals. Mean and S.D. are shown. (**D**) Statistics of the overlap analysis at multiple thresholds. Fold difference between the two groups of animals increases as the threshold increases. Statistical significance does not show clear increase with threshold as percentage overlap values are more variable at higher thresholds. P-values were calculated from two-tailed t-tests. We did not implement multiple comparison corrections as values at different thresholds are derived from the same dataset and are not independent.

Figure S6. Caspase-3 deficiency does not alter RGC density in the retina. (**A**) Images of whole-mount retinae from P10 *Casp3*[+/+] (left) and *Casp3*[-/-] (right) animals. RGCs were labeled in red by immunostaining against an RGC-specific marker, RBPMS (RNA-binding protein with multiple splicing) (*37*). The dark regions on the bottom of the left panel and on the left of the right panel are optic discs. Scale-bars: 200 μm. (**B**) Quantification of RGC densities in retinae of P10 *Casp3*[+/+] and *Casp3*[-/-] animals. Each point represents one retina. Left and right retinae were analyzed separately. n=4 animals for *Casp3*[+/+] mice and n=5 animals for *Casp3*[-/-] mice. Mean and S.D. are shown. P-values were calculated from unpaired two-tailed t-tests.

Figure S7. Measuring electrophysiological properties of retinogeniculate synapses. (**A**) Inferring the number of RGC inputs of dLGN relay neurons by measuring stimulation-response curves. To measure relay neuron responses, we prepared acute brain slices from a tilted parasagittal plane in P30 mice (*38*). These slice preparations preserved a long segment of the optic tract and a high level of connectivity between RGC axons and dLGN relay neurons (*38*). We placed a stimulating electrode on the optic tract and patch-clamped individual dLGN relay neurons with a recording electrode. We stimulated the optic tract with a series of gradually increasing currents and recorded excitatory postsynaptic currents (EPSCs) in relay neurons. At low stimulation intensity, no RGC axons were activated. As the stimulus increased, more and more RGC axons were recruited. As different RGC axons have different excitation thresholds, RGC axons were excited one at a time. Whenever a stimulus activated an RGC axon that did not respond to lower stimuli, a step increase in the relay neuron response was detected, and the increment corresponded to the EPSC evoked by the newly recruited RGC axon. We can therefore infer the number of RGC inputs innervating the relay neuron being recorded by counting the number of steps in the response curve of that relay neuron. t: time. Stim.: stimulation intensity. (**B**-**C**) Inferring the number of release sites from RGC inputs by measuring paired-pulse response ratio (PPR). We stimulated the optic tract with two stimuli separated by a short interval and recorded EPSCs in dLGN relay neurons. The stimulus intensity was chosen to evoke maximum response in the relay neuron. In wildtype mice (**B**), the first stimulus triggers the release of a significant fraction of the readily releasable pool (RRP) of neurotransmitters (black dots), evoking a strong response (left). At the time of the second stimulus, the RRP does not have sufficient time to recover and remains depleted, resulting in a weaker second response (right). PPR can be calculated from this experiment by dividing the peak amplitude of the second response with that of the first. PPR is small in wildtype animals. In caspase-3 deficient mice (**C**), if the number of release sites is increased compared to wildtype animals, the RRP should be larger (left), and a smaller fraction of the RRP is released during the first stimulus, leaving more neurotransmitters available for the second stimulus (right), thereby enhancing the second response and PPR. Note that in this model, we made two assumptions. The first assumption is that the average size of the RRP in one release site is similar in $Casp3^{+/+}$ and $Casp3^{-/-}$ mice, which is supported by the comparable fiber fractions (FFs) in the two groups of animals (see Fig. S8E). The second assumption is that the number of neurotransmitters released during the first stimulus is similar in $Casp3^{+/+}$ and $Casp3^{-/-}$ mice (4 dots in the illustration), which is supported by the comparable maximum EPSCs in the two groups of animals (see Fig. S8B-C).

Figure S8. Additional analyses of electrophysiological properties of retinogeniculate synapses in *Casp3*+/+ and *Casp3*-/- mice. (**A**) Distribution of RGC input numbers of individual dLGN relay neurons in wildtype and *Casp3*-/- mice measured by counting the number of steps in NMDAR-mediated EPSC response curves (upper right in Fig. 4A and 4B) while blind to genotypes. P-value was calculated from two-tailed t-test. n=22 cells for wildtype mice and n=24 cells for *Casp3*-/- mice. (**B-C**) Quantification of maximum amplitudes of AMPAR- (**B**) and NMDAR-mediated (**C**) EPSCs in wildtype and *Casp3*-/- dLGN relay neurons. (**D**) Quantification of the ratio between maximum amplitudes of AMPAR- and NDMAR-mediated EPSCs in wildtype and *Casp3*-/- dLGN relay neurons. (**E**) Quantification of fiber fractions in wildtype and *Casp3*-/- dLGN relay neurons. When stimulation on the optic tract is gradually increased, there will be a lowest stimulation intensity at which a non-zero response is first recorded in the relay neuron (see Fig. S8 for illustration and Fig. 4 for example data). This first response is presumed to be evoked by the activation of a single RGC axon fiber. Fiber fraction is defined as the ratio between this single-fiber response and the maximum response and ranges from 0 to 1. Only the AMPAR-mediated response is used to calculate fiber fractions. In **B-E**, mean and S.D. are shown. In **A-E**, p-values were calculated from two-tailed t-tests. In **A**, **C**, and **D**, n = 22 cells for wildtype mice and n=24 cells for *Casp3*-/- mice. In **B** and **E**, n=37 cells for wildtype mice and n=32 cells for *Casp3*-/- mice.

Figure S9. Caspase-3 deficiency does not cause microglia activation. (**A**) Representative images of P5 dLGNs in a *Casp3*[+/+] mouse (left) and a *Casp3*[-/-] mouse (right) showing Iba1 (green, a marker for microglia cell body) and Cd68 (magenta, a marker for microglia activation) staining. The dLGN areas are highlighted with dotted lines. Scale-bars represent 100 μm. (**B**) Quantification of microglia morphology in *Casp3*[+/+] and *Casp3*[-/-] mice using the sphericity metric. A sphericity of 1 corresponds to a perfect sphere. The smaller the sphericity is, the more ramified the cell is. (**C**-**D**) Quantification of microglia activation by normalizing the total volume of Cd68 signal (**C**) or the total intensity of Cd68 signal (**D**) in each microglia to the volume of that microglia. By both morphology and Cd68 signal, microglia in *Casp3*[-/-] mice show no evidence of activation. In **B**-**D**, each data point represents one microglia, and mean and S.D. are shown. n=46 microglia from 4 *Casp3*[+/+] mice and n=52 microglia from 4 *Casp3*[-/-] mice. P-values were calculated from two-tailed unpaired t-tests.

● *Casp3(+/-); Cx3cr1-Gfp(+/-)*   ● *Casp3(-/-); Cx3cr1-Gfp(+/-)*

Figure S10. Additional analyses of microglia-mediated engulfment of synaptic material. (**A-B**) Volume of contralateral (**A**) and ipsilateral (**B**) RGC axon terminals engulfed by individual microglia in the P5 dLGN of *Casp3*[+/-]; *Cx3cr1-Gfp*[+/-] and *Casp3*[-/-]; *Cx3cr1-Gfp*[+/-] mice. (**C**) Volume of individual microglia in the P5 dLGN of *Casp3*[+/-]; *Cx3cr1-Gfp*[+/-] and *Casp3*[-/-]; *Cx3cr1-Gfp*[+/-] mice. In **A-C**, each point represents one microglia. Mean and S.D. are shown. P-values were calculated from unpaired two-tailed t-tests. n=61 microglia from 8 *Casp3*[+/-]; *Cx3cr1-Gfp*[+/-] mice and n=54 microglia from 5 *Casp3*[-/-]; *Cx3cr1-Gfp*[+/-] mice.

Figure S11. Astrocyte-mediated synapse elimination does not appear to depend on caspase-3. (**A**) Representative 3D-reconstructed images of a P5 $Casp3^{+/-}$; $Aldh1l1$-$Gfp^{+/-}$ mouse dLGN with astrocytes displayed in green, contralateral RGC axon terminals in red, and ipsilateral RGC terminals in blue. The region from which astrocytes are selected for analysis is indicated with the dashed line in the merged image. Scale-bars represent 50 μm. (**B**) Representative surface rendering of astrocytes (green) from P5 dLGNs of $Casp3^{+/-}$; $Aldh1l1$-$Gfp^{+/-}$ and $Casp3^{-/-}$; $Aldh1l1$-$Gfp^{+/-}$ mice. Intracellular contralateral (red) and ipsilateral (blue) RGC axon terminals are shown. Only cell bodies and the base of processes were segmented. We made this segmentation choice because we observed that fine astrocytic processes are largely devoid of engulfed material. Scale-bars represent 5 μm. (**C**) Total volume of engulfed synaptic material in individual astrocytes from $Casp3^{+/-}$; $Aldh1l1$-$Gfp^{+/-}$ and $Casp3^{-/-}$; $Aldh1l1$-$Gfp^{+/-}$ mice. (**D**) Total volume of engulfed synaptic material in each astrocyte (from **C**) is normalized to the volume of that astrocytes. (**E**) Volume of individual astrocytes from the two groups of mice. In **C**-**E**, each point represents one astrocyte. Mean and S.D. are shown. P-values were calculated from unpaired two-tailed t-tests. n=595 astrocytes from 8 $Casp3^{+/-}$; $Aldh1l1$-$Gfp^{+/-}$ mice and n=517 astrocytes from 6 $Casp3^{-/-}$; $Aldh1l1$-$Gfp^{+/-}$ mice.

Figure S12. Additional analysis on activity-dependent microglia-mediated engulfment of synapses. (**A**-**D**) Volumes of synaptic material originating from right (**A** and **C**) or left (**B** and **D**) eyes engulfed by individual microglia from the left dLGN of *Casp3*$^{+/+}$ (**A**-**B**) or *Casp3*$^{-/-}$ (**C**-**D**) mice injected with AAV expressing mTurquoise2 (blue) or TeTxLC (red) in right eye were normalized to microglial volume and plotted. In *Casp3*$^{+/+}$ mice, inactivating right eye-originated retinogeniculate synapses specifically enhanced microglia-mediated engulfment of RGC axon terminals from right eyes (**A**) but not left eyes (**B**), resulting in significantly higher engulfment ratios (Fig. 6F). In *Casp3*$^{-/-}$ mice, inactivating right eye-originated synapses upregulated engulfment of RGC terminals from both eyes (**C** and **D**), resulting in no significant change in engulfment ratios (Fig. 6G). In **A**-**D**, 0, 25, 50, 75, and 100 percentiles are shown. p-values were calculated from unpaired two-tailed t-tests. n=52 microglia from 7 Turq-injected *Casp3*$^{+/+}$ mice, n=50 microglia from 8 TeTxLC-injected *Casp3*$^{+/+}$ mice, n=64 microglia from 5 Turq-injected *Casp3*$^{-/-}$ mice, and n=51 microglia from 6 TeTxLC-injected *Casp3*$^{-/-}$ mice. ***Notes on interpretation of C and D***: We observed that in *Casp3*$^{-/-}$ brains, particularly those injected with TeTxLC, microglia in dLGNs tended to increase in volume and have thicker processes. We therefore suspect the non-specific upregulation of synapse engulfment in TeTxLC-injected *Casp3*$^{-/-}$ mice is a consequence of microglial activation rather than synapse inactivation. This microglial activation is likely a ramification of intraocular surgeries, as changes in microglia morphology was more prominent in the left dLGN that received most of the inputs from AAV-injected right eyes, and microglia outside of dLGNs did not show characteristics of activation. We noticed that *Casp3*$^{-/-}$ brains tended to have weak structural integrity. It is likely that repetitive intraocular injections caused damage in the retinogeniculate pathway of *Casp3*$^{-/-}$ mice and activated microglia. Nevertheless, by calculating right-to-left engulfment ratios, substrate preference of microglia in TeTxLC-injected *Casp3*$^{-/-}$ mice can be deduced.

A

App/Ps1(-/-) Female 6 Months   *App/Ps1*(+/-) Female 5 Months   *App/Ps1*(+/-) Female 6 Months

Thioflavin S

i   ii   iii

App/Ps1(+/-) Male 5 Months   *App/Ps1*(+/-) Male 6 Months   *Casp3*(-/-); *App/Ps1*(+/-)
Female 6 Months

Thioflavin S

iv   v   vi

B

p = 0.73

#Plaque per section

App/Ps1(+/-)   *Casp3*(-/-);
App/Ps1(+/-)

Figure S13. Amyloid deposition APP/PS1 mouse lines. (**A**) Images of coronal brain sections of a female *App/Ps1*[-/-] mouse at 6 months (i), female *App/Ps1*[+/-] mice at 5 months (ii) or 6 months (iii), male *App/Ps1*[+/-] mice at 5 months (iv) or 6 months (v), and a female *Casp3*[-/-]; *App/Ps1*[+/-] mouse at 6 months, stained with Thioflavin S to reveal Aβ plaques (bright spots in hippocampus and cortex). Scale-bars represent 1 mm. (**B**) Quantification of the number of plaques per section in female 6 month-old *App/Ps1*[+/-] and *Casp3*[-/-]; *App/Ps1*[+/-] mice. Mean and S.D. are shown. p-values were calculated with unpaired two-tailed t-tests. n=6 for *App/Ps1*[+/-] mice and n=4 for *Casp3*[-/-]; *App/Ps1*[+/-] mice.

App/Ps1(-/-)   *App/Ps1(+/-)*   *Casp3(-/-); App/Ps1(-/-)*   *Casp3(-/-); App/Ps1(+/-)*

SV2 Homer1

Figure S14. Selection of field of interest for synapse loss analysis. To quantify synapse density, three regions of interest from each dentate gyrus on either side of the brain were chosen for each animal (total of 6 regions per animal). Regions of interest in one dentate gyrus of one animal from each genotype group are highlighted with dotted squares in the overviews above. High resolution images of SV2 (in green) and Homer1 (in red) stains were acquired for each region of interest and used for analysis. In sections from animals overexpressing mutant APP and PS1, we avoided choosing regions of interest from areas surrounding amyloid plaques (highlighted with arrows).

Figure S15. Additional analysis of synapse density in APP/PS1 mouse lines. (**A**-**D**) Quantification of presynaptic (**A** and **C**) and postsynaptic (**B** and **D**) puncta densities in the dentate gyrus of APP/PS1 mice in the caspase-3 wildtype (**A** and **B**) or deficient (**C** and **D**) background. Mean and S.D. are shown. p-values were calculated from unpaired two-tailed t-tests. n=5 for *App/Ps1*[-/-] mice, n=6 for *App/Ps1*[+/-] mice, n=6 for *Casp3*[-/-]; *App/Ps1*[-/-] mice, and n=4 for *Casp3*[-/-]; *App/Ps1*[+/-] mice.

Figure S16. Microgliosis in APP/PS1 mouse lines. Coronal sections from 6 month-old female *App/Ps1*$^{-/-}$ (left), *App/Ps1*$^{+/-}$ (middle), and *Casp3*$^{-/-}$; *App/Ps1*$^{+/-}$ (right) mice were stained for Iba1 (red, to detect microglia) and Cd68 (green, as a microglia activation marker). Dentate gyrus regions are shown in the overviews above. Clusters of reactive microglia surrounding amyloid plaques are highlighted with arrows.

Figure S17. Aβ-induced caspase-3 activity in APP/PS1 mice. (**A**-**C**) Density of activated caspase-3 puncti in the dentate gyrus of 4 month-old (**A**), 5 month-old (**B**), and 6 month-old (**C**) female *App/Ps1*[-/-] and *App/Ps1*[+/-] mice. Each column represents one animal, and each dot represent quantification from one dentate gyrus (either from left side or right side). Two out of four 4-month-old *App/Ps1*[+/-] mice showed robustly upregulation of caspase-3 activity in the dentate gyrus (**A**). No upregulation of caspase-3 activity was observed in 5 month-old or 6 month-old mice. (**D**) Images showing elevated caspase-3 activity in the molecular layer of the dentate gyrus of a 4 month-old female *App/Ps1*[+/-] mouse (A#5 in **A**). Images were set to the same intensity contrast. Upregulated caspase-3 activity remained in a punctate pattern and no apoptotic cell was observed. Scale-bar represents 10 μm. (**E**) Representative images of the dentate gyrus in 6 month-old female *App/Ps1*[-/-], *App/Ps1*[+/-], *Casp3*[-/-]; *App/Ps1*[-/-], and *Casp3*[-/-]; *App/Ps1*[+/-] mice that were stained with an anti-NeuN antibody to visualize neurons. The fields of interest used to quantify neuron density are highlighted with dotted rectangles. Scale-bars represent 100 μm. (**F**) Neuron density in the granule cell layer of the dentate gyrus in the four groups of mice shown in **E**. Neither caspase-3 deficiency nor APP/PS1 overexpression caused significant changes in neuron density. Mean and S.D. are shown. p-values were calculated from Tukey's multiple comparisons test. n=5 for *App/Ps1*[-/-] mice, n=6 for *App/Ps1*[+/-] mice, n=5 for *Casp3*[-/-]; *App/Ps1*[-/-] mice, and n=4 for *Casp3*[-/-]; *App/Ps1*[+/-] mice.

Material and methods

Mouse lines

The $Casp3^{-/-}$ mouse line was a generous gift from Dr. Richard Flavell. This mouse line was received as cryopreserved sperm and was rederived using C57Bl/6J oocytes. The resulting colony was maintained by continuous backcrossing to C57Bl/7J mice. We verified that mice in our $Casp3^{-/-}$ colony were congenic to the C57Bl/6J background (greater than 99% identity by SNP analysis). It is important to note that regular backcrossing is necessary to maintain phenotype stability in $Casp3^{-/-}$ mice. A small fraction of $Casp3^{-/-}$ mice displayed varying degrees of hydrocephalus. These animals were excluded from our analyses. *Cx3cr1-Gfp* mice (stock number 005582) were obtained from the Jackson laboratory. *Aldh1l1-Gfp* transgenic mice (stock number 011015-UCD) were obtained from Mutant Mouse Resource and Research Centers (MMRRC) as cryopreserved sperm and rederived using C57Bl/6J oocytes. *App/Ps1* mice were obtained from the Jackson laboratory (stock number 034832). Mice carrying *App/Ps1* transgenes were given DietGel® 76A (ClearH$_2$O, 72-07-5022) and were euthanized at the age of 6 months if not used in experiments. For experiments associated with Figure 1 and Figure 2, C57Bl/6NCrl mice were obtained from Charles Rive Laboratories (strain code 027). For experiments associated with Figure 6, C57Bl/6J mice were obtained from the Jackson Laboratory (stock number 000664). All animal procedures were carried out according to IACUC approved protocols.

Reagents

Reagents were obtained from the following sources: isoflurane liquid (NDC 50989-

150-15) was from Vedco; phosphate buffered saline without calcium and magnesium

(PBS⁻, BP2438) was from Fisher Scientific; bicuculline (0130) and cyclothiazide (0713)

were from Tocris Bioscience; PBS with calcium and magnesium (PBS⁺, SH30264) was

from Cytiva; dimethyl sulfoxide (DMSO, D2650), bovine serum albumin (BSA, A9647),

triton X-100 (T8787), Thioflavin S (T1892) were from Millipore-Sigma; paraformaldehyde

powder (19200) was from Electron Microscopy Science; RPMI 1640 medium (11875093),

DMEM medium (11965092), fetal bovine serum (FBS, 16000044), penicillin/streptomycin

(pen/strep, 15140122), CD Hybridoma medium (11279023), GlutaMAX supplement

(35050061), protein A/G agarose beads (20421), protein A/G IgG binding buffer (54200),

IgG elution buffer (21004), 1M Tris-HCl pH 8.0 (15568025), 10% normal goat serum

(50062Z), ProLong™ Gold mounting medium (P36934), Alexa Fluor 546 succinimidyl

ester (A20002), cholera toxin subunit beta(CTB) conjugated to AlexaFlour (AF) 488,

AF555, AF594 and AF647   (C22841, C22842, C22843, C34778), goat anti-rabbit

secondary antibodies conjugated to AF488plus (A32731), AF555plus (A32732),

AF594plus (A32740), and AF647plus (A32733), goat anti-chicken secondary antibody

conjugated to AF488plus (A32931), goat anti-guinea pig secondary antibody conjugated

to AF488 (A11073) and AF594 (A11076) were from ThermoFisher Scientific; anti-cleaved

caspase-3 antibody (9661) was from Cell Signaling Technologies; anti-GFP antibody

(GFP-1010) was from Aves Lab; anti-MAP2 antibody (NB300-213) was from Novus

Biologicals; anti-NeuN antibody (266004) was from Synaptic Systems; anti-RBPMS

antibody (1832-RBPMS) was from PhosphoSolutions; anti-Iba1 (ab178846) and anti-

Homer1 (Ab97593) antibodies were from Abcam; anti-Cd68 antibody (MCA1957) was

from Bio-Rad; hybridoma line producing anti-SV2 antibody (clone name SV2) was from

DSHB.

DNA cloning and AAV preparation

To make the pAAV-hSyn-mTurquoise2 plasmid, we ordered the DNA fragment for

mTurqouise2 (*1*) as a Gblock from IDT and cloned it via isothermal assembly into a

linearized AAV expression plasmid (derived from pAAV-Syn-iCre; Addgene #122518)

under the control of human synapsin promoter. To make the pAAV-hSyn-TeTxLC-P2A-

mTurqouise2 plasmid, the tetanus toxin light chain (TeTxLC) sequence along with the

P2A sequence (2A peptide from porcine teschovirus-1 polyprotein) at the 3' end was

obtained via PCR from Addgene plasmid #129102 and fused in-frame with the

mTurquoise2 DNA sequence in the same plasmid backbone as the pAAV-hSyn-

mTurquoise2 plasmid. Note that in the main text, we used pAAV-hSyn-TeTxLC to refer to

pAAV-hSyn-TeTxLC-P2A-mTurquoise2 because AAV carrying this plasmid induced very

low levels of mTurquoise2 expression *in vivo*. When we needed endogenous florescence

labeling, we chose to premix AAV carrying pAAV-hSyn-TeTxLC-P2A-mTurquoise2 with

other AAVs expressing fluorescent proteins (at 1:1 ratio).

The pAAV-hSyn-tdTomato plasmid was cloned and provided by Janelia Viral Tools. The

pAAV-CAG-GFP plasmid was previously made by Svoboda lab and is available as

Addgene #28014.

The recombinant AAVs were prepared by Viral Tools of Janelia Core Services. Briefly, two

days before transfection, 293T cells were seeded in three 150 mm T/C dishes at 1×10^7

cells/dish in DMEM medium supplemented with 10% FBS and cultured at 37 °C with 5%

CO_2. The cells were transfected with 84 μg of pAAV plasmid, DJ capsid plasmid (*2*), and

pHelper plasmid (Agilent 240071) at a ratio of 5:2:3 in polyethylenimine. After 6 to 8 hours,

the cells were replenished with serum-free DMEM and further incubated for 3 days. The

recombinant AAVs were collected from both the cells and supernatant and purified by two

rounds of continuous cesium chloride density gradient centrifugation, dialyzed,

concentrated, and filter sterilized. Titer (genome copies per ml) was determined by digital

PCR with primers targeting the ITR. For all in vivo experiments a titer of at least 10^{13}

genome copies/ml was used.

Intraocular injections

In utero intraocular injections were performed as previously described (*3*) with

modifications. Briefly, pregnant mouse females with embryos at E15 gestation were

anesthetized by 3% isoflurane inhalation, abdomens shaved, eye lube applied to each

eye, then transferred to a nose cone and set to maintenance level of 2% isofluorane

inhalation anesthesia. Warm fluids (lactated Ringer's solution + 5% dextrose, Strategic

Applications Inc.), Buprenorphine (0.1 mg/kg, Butler Animal Health), and Ketoprofen (5

mg/kg, Covetrus) were administered subcutaneously just prior to starting surgery. The

animal was placed in dorsal recumbency, and the abdomen was cleansed using

alternating swabs of 70% ethanol and 4% Chlorhexidine Gluconate (Molnlycke Health

Care). A sterile drape was placed over the abdomen and using aseptic technique, the

abdominal skin and wall were cut at midline, before gently removing the uterus from the

abdomen. While keeping the uterus/embryos hydrated with warm saline sterile solution,

the embryos were gently positioned with ring forceps to hold each embryo securely for

injection. A previously filled glass pipette (Drummond Scientific), beveled to 30-40 μm at

the opening was utilized to deliver constructs utilizing a pre-programmed Nanoject III

microinjector (Drummond Scientific). 200-250 nl of AAVs prep was injected through the

uterine wall and into each eye at a speed of 50nl per second. To confirm proper targeting

within the eye, the AAV solution was pre-mixed with 0.05% Fast Green (Sigma Aldrich).

After all injections were completed, the uterus/embryos were carefully placed back into

the abdomen and the abdomen was flushed with warm sterile saline. The abdominal wall

and skin were sutured, and Marcaine (Hospira) was applied subcutaneously below the

skin incision. A small drop of Vetbond (3M) was applied to seal the skin incision. The

animal was then removed from the anesthesia apparatus and allowed to recover on a

heating pad before being placed back into a clean cage.

For intraocular injection at P4 and P9, pups with desired genotypes were

anesthetized in an induction chamber filled with 2.5% - 3% isoflurane using a calibrated

vaporizer. After the onset of anesthesia, isoflurane is reduced to 1.25%, and the mouse

was removed from the chamber and placed under a dissection microscope. Proparacaine

ophthalmic solution (0.5%, Sandoz, NDC 61314-016-01) was applied to each eye to be

injected. A re-breathing nose cone was used to deliver 1.25% – 2% isoflurane throughout

the procedure. A gentle blunt dissection was performed at each eyelid junction utilizing

closed No.5 Dumont forceps (Fine Science Tools) to create a 2-3 mm opening. A 33-

gauge sharp beveled needle attached to a Hamilton microliter syringe (Hamilton

Company) was used to draw up CTB-AF solutions. Just prior to injecting, the eye was

proptosed and held securely in place by the Dumont forceps in preparation for injection.

While holding the eye securely in place, the sclera was punctured with the Hamilton

needle at the level of the ora serrata to inject 2.5 µL (for P9) or 2 µL (for P4) of desired

CTB-AF solution into each eye. After the injection, eye lube was applied, the eye was

allowed to move back into the socket, and the eyelids were gently pressed close to protect

the eye. Pups were allowed to recover from anesthesia on a heated pad and put back

into the cage with the dam.

Monoclonal antibody production and labeling

SV2 hybridoma cells were thawed and seeded in RPMI 1640 medium containing

10% FBS and 1% Pen/Strep at 0.3 million cells/ml and cultured in an incubator at 37 $^{\circ}$C

with 5% CO_2. After reaching $1x10^6$ viable cells/ml density, the cells were diluted 1:2 by

adding an equal volume of serum free CD Hybridoma medium supplemented with 4%

GlutaMax^{Tm} and returned to the incubator. This dilution process was repeated 2 more

times so that the cells reached $1x10^6$ viable cells/ml density in a mixed medium containing

~1% FBS. Cell viability obtained was at least 90%. The cells were then spun down at

150xG for 10 minutes and resuspended in complete CD Hybridoma medium at $5x10^5$

cells/ml density. The cells were further expanded by adding more complete CD

Hybridoma medium when cells reach $1x10^6$ cells/ml density. Once the desired total

volume is reached (typically 100 ml), cells were cultured for 2 weeks without medium

change.

Culture supernatant containing SV2 antibody was cleared of hybridoma cells by

centrifuging at 1000xG for 10 min. Cleared supernatant was diluted 1:1 with IgG binding

buffer and passed through 1 ml of protein A/G agarose bead slurry (ThermoFisher 20421,

corresponding to 500 ul of settled resin) packed in a gravity flow column. The column was

washed with IgG binding buffer, and SV2 antibody was eluted in five 1 ml fractions using

IgG elution buffer and immediately neutralized with 10% volume of 1M Tris-HCl, pH 8.0.

Fractions with highest antibody concentration were pooled, desalted with Zeba columns

(ThermoFisher 89891), and concentrated to ~1 mg/ml using Amicon centrifuge filters

(Millipore UFC801024). After adding glycerol (25% final), antibodies were flash frozen in

liquid nitrogen and kept at -80 °C.

To label the primary SV2 antibodies, we used Alexa Fluor 546 succinimidyl ester.

Briefly, we buffer-exchanged and concentrated SV2 IgG to 3 mg/ml using Amicon filters

(UFC501024 Millipore Sigma) in phosphate buffered saline that was brought to pH 8.3

with 100 mM NaHCO$_3$. Dimethyl sulfoxide-dissolved Alexa Flour 546 succinimidyl ester

was gently mixed with the SV2 IgG at 10:1 molar ratio and incubated for two hours at

room temperature on a rotating shaker. Alexa Fluor 546 labeled IgG were purified from

unreacted succinimidyl esters with the aid of PD MiniTrap Sephadex G-25 column (Cytiva

28918007) in PBS$^-$ by gravity flow following the manufacturer's instructions. The degree

of labeling was determined by spectrophotometry on Nanodrop One (ThermoFisher) by

accounting for the Alexa Fluor 546 extinction coefficient at 554 nm and the corrected

protein absorbance at 280 nm. Typically, a degree of labeling of 8-9 mols of dye per mol

of protein was obtained. Labeled SV2 were aliquoted and flash-frozen in 25% glycerol for

long term storage at -80 °C at a final concentration of approximately 0.75 mg/ml.

Immunohistochemistry

The mice were deeply anesthetized in a chamber filled with saturating isoflurane

gas and transcardially perfused first with PBS$^-$ and then with paraformaldehyde (PFA)

solution (4% w/v in 20.3 mM NaH$_2$PO$_4$ and 79.7 mM Na$_2$HPO$_4$, pH = 7.4). The brains

were removed and post-fixed in 4% PFA solution overnight at 4 °C with gentle shaking.

Fixed brains were washed 3x15 minutes in PBS⁻ at room temperature and stored in PBS⁻

at 4 °C. For long term storage, fixed brains were cryoprotected in 30% Sucrose (w/v in

PBS⁻) for 72 hours and frozen in Tissue-Tek® O.C.T. compound (Sakura) and stored at -

80 °C.

Brains stored in PBS⁻ were imbedded in 4% agarose, and coronal sections (50 um

thick for mice younger than P10 or 40 μm thick for adult mice) were prepared using a

vibratome (Leica VT 1200S). For brains frozen in O.C.T. compound, coronal sections

were prepared using a cryostat (Leica 3050S). Sections were floated in 24-well plates in

PBS⁻ and stored at 4 °C. For antibody staining, free-floating sections were first

permeabilized/blocked in 10% normal goat serum containing 0.3% (w/v) triton X-100 for

1 hour. The sections were then incubated with primary antibodies diluted in staining buffer

(PBS⁺ containing 2% BSA w/v and 0.3% triton X-100) at 4 °C overnight with gentle

agitation. Dilutions of primary antibodies were: anti-cleaved caspase-3, 1:500; anti-GFP,

1:2000; anti-MAP2, 1:1000; anti-NeuN, 1:500; anti-Iba1, 1:500; anti-Cd68, 1:500; anti-

SV2, dilute to 2.5 μg/ml; anti-Homer1, 1:400. The sections were then washed 3x15

minutes in washing buffer (PBS⁺ containing 0.1% triton X-100) and stained with

appropriate fluorophore-conjugated secondary antibodies diluted in staining buffer for 3

hours at room temperature with gentle agitation. Secondary antibodies were always used

at 1:500 dilution. The sections were washed 3x15 minutes in washing buffer, rinsed 3x5

minutes in PBS⁺, and mounted onto glass slides (Fisher, 12-550-15). After air-drying

completely, mounted sections were immersed in ProLong™ Gold and coverslipped

(Fisher, 12-541-024). The sections were imaged after 24 hours of curing at room

temperature. The sections were protected from light during the entire process.

To prepare whole-mount retinae, eyes were removed from PFA perfused mice and

placed in PBS⁻. The cornea was punctured with a #11 scalpel, and several anterior-to-

posterior cuts were made on the sclera starting from the puncture. Intact retinae were

isolated by tearing and peeling off the sclera along the cuts. 4 to 5 radial cuts were made

on each retina to flatten the tissue. Isolated retinae were post-fixed for 15 minutes in 4%

PFA solution at room temperature with gentle agitation and washed for 3x5 minutes in

PBS⁻ at room temperature with gentle agitation. Free-floating retinae were then stained

with primary (anti-RBPMS antibody, used at 1:200 dilution) and secondary (goat anti-

guinea pig Alexa Fluor 594plus) antibodies and mounted onto glass slides as described

above.

For staining of amyloid-β (Aβ) plaques, a 1% Thioflavin S stock solution (w/v,

dissolved in ddH$_2$O) was diluted 100-fold in 50% ethanol (prepared by mixing pure ethanol

and ddH$_2$O at 1:1 ratio) to create a 0.01% (w/v) working solution. The stock and working

solutions were discarded after each use. Free-floating brain sections were dehydrated by

washing 2x5 minutes in 50% ethanol, stained in 0.01% Thioflavin S solution for 8 minutes,

washed for 2x5 minutes in 50% ethanol, then re-hydrated by washing 3x5 minutes in

PBS⁺. All staining steps were performed at room temperature with gentle agitation. The

sections were protected from light throughout the procedure. Stained sections were then

mounted and cured as described above.

Measuring synapse inactivation-induced caspase-3 activity

For experiments associated with Figure 1 and Figure S4, E15 C57Bl/6NCrl mice

were injected in the right eyes either with AAV carrying hSyn-mTurquoise2 (control group),

or with a 1:1 mixture of AAV carrying hSyn-mTurquiose2 and AAV carrying hSyn-TeTxLC-

P2A-mTurquoise2 (synapse inactivation group). At the age of P5, brains of both groups

of mice were harvested and sectioned as described above. To quantify caspase-3 activity

in the dLGN and the localization of activated caspase-3 relative to TeTxLC-expressing

RGC axons (Figure 1), sections were stained with anti-cleaved caspase-3 antibody (using

goat anti-rabbit AF594plus as the secondary) and anti-GFP antibody (using goat anti-

chicken AF488plus as the secondary). Alternatively, sections were co-stained with anti-

cleaved caspase-3 (using goat anti-rabbit AF647plus as the secondary) and anti-MAP2

antibody (using goat anti-rabbit AF488plus as the secondary) (Figure S4A), or with anti-

cleaved caspase-3 (using goat anti-rabbit AF594plus as the secondary) and anti-NeuN

antibody (using goat anti-rabbit AF488plus as the secondary) (Figure S4C).

Immunostaining was performed as described above.

For experiments associated with Figure 2, E15 C57Bl/6NCrl mice were injected in

the left eyes with AAV carrying hSyn-tdTomato and in the right eyes with AAV carrying

CAG-eGFP (control group), or in the right eyes with AAV carrying hSyn-mTurquoise2 and

AAV carrying hSyn-TeTxLC-P2A-mTurquoise2 (single inactivation group), or in the left

eyes with AAV carrying hSyn-tdTomato and AAV carrying hSyn-TeTxLC-P2A-

mTurquoise2 and in the right eyes with AAV carrying CAG-eGFP and AAV carrying hSyn-

TeTxLC-P2A-mTurquoise2 (dual inactivation group). At the age of P5, brains of all three

groups of animals were harvested, sectioned, and stained with anti-cleaved caspase-3

antibody and goat antirabbit AF647plus secondary antibodies as described above.

To quantify caspase-3 activity in the dLGN (Figure 1B-E), confocal images of

dLGNs were acquired on a Leica Stellaris 8 microscope using a 10x/0.4 NA dry objective

(Leica HC PL Apo CS2 10x/0.4 dry). The channels were acquired sequentially (line-by-

line) with the anti-caspase-3 first and anti-mTurquoise2 second. A 590 nm laser and

emission read of 596 nm – 748 nm was used for the anti-cleaved caspase-3 channel, and

440 nm and 488 nm lasers and an emission read of 457 nm – 579 nm were used for anti-

mTurquoise2 channel. The imaging parameters (laser power, pixel dwell time, signal

averaging or accumulation) were set to maximize signal-to-background values while

avoiding detector (SiPM type Power HyD S) saturation. Photon detection was set in

counting mode and acquired as 12-bit images at 227 nm x 227 nm pixel size. All datasets

used for measurement of caspase-3 activity were acquired as Z-stacks (7 slices spaced

2.4 μm apart). To measure the total amount of active caspase-3 signal in the 3D stacks

of dLGN we used the Labkit (*4*) Fiji plugin to generate and train a pixel classifier through

Imaris v. 10 (Oxford Instruments). The same pixel classifier and surface creation settings

were applied to data from both the control and the synapse inactivation groups to

generate surface objects of active caspase-3 signal. All active caspase-3 objects found

within the dLGN area (manually delineated based on mTurquoise2 signal) were selected,

and the sum of voxel intensities the active caspase-3 objects was calculated and divided

by the area of dLGN. The final values plotted for each mouse are averages of two

consecutive sections.

To image active caspase-3 signals and TeTxLC-expressing RGC axons at high

resolution (Figure 1G), we used a Leica Stellaris 8 confocal microscope and a 63x/1.4 NA

oil objective (Leica HC PL APO 63x/1,40 OIL CS2) in "Lightning" mode (in LasX 4.6) set

at high resolution grade, which reduces the pinhole size to 0.5 Airy units. Laser and

emission settings were identical to those used the previous section that described

quantification of active caspase-3 signal in mTurquoise2-labeled dLGNs. 3D image

stacks (31.5 um x 31.5 um x 12 um) were acquired at high voxel density (43 nm x 43 nm

x 204 nm) in photon counting mode and 4x signal accumulation. Subsequently, the

images were processed by iterative deconvolution using a theoretical point spread

function informed by the imaging parameters of the microscope and constrained by an

"Adaptive" strategy that accounts for local background and signal-to-noise ratio

information.

Confocal images of MAP2 and active caspase-3 signals in dLGN (Figure S4A)

were similarly imaged on a Leica Stellaris 8 microscope with a 63x/1.4 NA oil objective

(Leica HC PL APO 63x/1,40 OIL CS2). A 653 nm laser line and an emission read of 662

nm – 829 nm were used for caspase-3 channel, and a 499 nm laser line and an emission

read of 504nm – 549 nm were used for Map2 channel. The channels were acquired

simultaneously at voxel densities of 65 nm x 65 nm x 299 nm. NeuN and active caspase-

3 signals in dLGN (Figure S4C) were imaged using a 20x/0.7 NA water objective (Leica

HC PL APO 20x/0,7 IMM CORR CS2). A 590 nm laser and an emission read of 595 nm

– 750 nm were used for imaging caspase-3 and a 499 nm laser with a 509 nm - 595 nm

emission read was used for the NeuN signal. The channels were acquired sequentially

(active caspase-3 first) at pixel densities of 142 nm x 142 nm.

Measuring overlap of eye-specific territories in dLGN

For comparison between Casp3$^{+/+}$ and Casp3$^{-/-}$ animals (Figure 3), Casp3$^{+/-}$ males

were bred to Casp3$^{+/-}$ females to generate Casp3$^{+/+}$ and Casp3$^{-/-}$ littermate pups. Pups

were genotyped at birth. Before intraocular injection, lyophilized CTB-AF488 and CTB-

AF594 powders were reconstituted in 1% DMSO in PBS^{-} to a final concentration of 5

mg/ml. CTB solutions were aliquoted and stored at -20 °C for later use. Thawed CTB

solutions were discarded after each surgery. Intraocular injection of CTB-AF solutions

was performed at the age of P9 as described above. CTB-AF488 was injected into left

eyes and CTB-AF594 into right eyes. 24 hours after the injection, brains of P10 pups were

harvested, and 50 um thick coronal sections prepared as described above.

Cured sections were imaged on a Zeiss AxioObserver Z.1 microscope equipped

with an LSM 880 Airy scan detector and an EC Plan-Neofluar 10x objective (NA = 0.3). A

488 nm laser and a 499-544 nm detection window were used for CTB-AlexaFlour488,

and a 594 nm laser and a 597-659 nm detection window were used for CTB-

AlexaFlour594. Signals from the two channels were acquired sequentially into 16-bit

images with a 0.4 µm x 0.4 µm pixel size. Laser power and detector gain settings were

maintained at the same level for most images, with occasional adjustment to avoid

saturation. One image per dLGN per section was acquired.

To quantify the overlap between contralateral-specific and ipsilateral-specific

territories in the dLGN, we selected 7-8 sections per animal that span the majority of left

dLGN of and analyzed their images in ImageJ. For each image, a mask for the entire

dLGN region and a mask for a background region in the thalamus with no labeling were

manually created. Background signal from each channel was estimated as the average

signal in the background region and was subtracted from signals in the dGLN region.

Then the background-subtracted signals in the dLGN region were normalized to between

0 and 1 for each image and each channel. We picked 7 increasing cutoff thresholds (0.1,

0.125, 0.15, 0.175, 0.2, 0.225, and 0.25), and for each threshold and each channel,

signals that are greater than or equal to the threshold were considered real signals. For

each animal (corresponding to 7-8 images of left dLGN) and each threshold, we

calculated percentage overlap as the ratio between the total number of pixels with real

signals in both channels in all images and the total number of pixels in the entire dLGN

region of all images. We plotted the percentage overlap as a measurement of segregation

of eye-specific territories in dLGN. Similar analyses were performed on right dGLN

images yielding similar results.

For overlap analysis presented in Figure S3, E15 mouse embryos were injected in

their right eyes either with AAV carrying hSyn-mTurqouise2 (control group) or a 1:1

mixture of AAV carrying hSyn-TeTxLC-P2A-mTurqouise2 and AAV carrying hSyn-

mTurquoise2 (inactivation group). The left eyes of both groups of mice were injected with

AAV carrying the hSyn-tdTomato. At P8, brains from both groups were harvested, and 50

μm thick coronal sections were prepared as described above. Only animals that had well

labeled dLGNs on both sides were selected for analysis. Mounted and cured sections

containing dLGN were imaged in confocal mode on a Leica Stellaris 8 microscope using

a 10x/0.4 NA dry objective (Leica HC PL Apo CS2 10x/0.4 dry). A 441 nm laser and a

445-548 nm detection window were used for the mTurquoise2 channel, and a 554 nm

laser and a 564-721 nm detection window were used for tdTomato channel. The images

were acquired at 8-bit depth with a 0.227 μm x 0.227 μm pixel size (at 1 Airy unit for

$\lambda = 580$ nm) and at a laser power, pixel dwell time, and signal accumulation sufficient to

avoid detector (Power HyD S in counting mode) saturation. One image per dLGN per

section was acquired. For each animal, overlap was measured as described above on

the left dLGN (the side where mTurquoise2 signal occupied the majority of the dLGN) by

averaging from 3 to 5 consecutive sections. The cutoff thresholds used were: 0.050, 0.075,

0.1, 0.125, 0.15, 0.175, 0.2.

For overlap analysis presented in Figure S5, embryos in the control group were
injected at E15 with AAV carrying tdTomato construct in the left eyes, and with AAV
carrying the CAG-GFP construct in the right eyes. Embryos in the dual inactivation group
were injected at E15 with a mixture (1:1) of AAV carrying hSyn-tdTomato construct and
AAV carrying hSyn-TeTxLC-P2A-mTurquoise2 construct in their left eyes, and with a
mixture (1:1) of AAV carrying the CAG-GFP construct and AAV carrying the hSyn-TeTxLC-
P2A-mTurquoise2 construct in their right eyes. At P10, the brains were harvested and
sectioned as described above. Confocal images of dLGN were acquired on a Leica
Stellaris 8 microscope using a 10x/0.4 NA dry objective (Leica HC PL Apo CS2 10x/0.4
dry). A 488 nm laser line with a 488 nm notch filter and a 485-559 nm detection window
were used for the GFP channel, and a 554 nm laser and a 559-701 nm detection window
were used for the tdTomato channel in sequential line mode to avoid any cross-excitation
(with tdTomato channel acquired first). The images were acquired at 12-bit depth with a
0.248 μm x 0.248 μm pixel size (at 1 Airy unit for $\lambda = 580$ nm) and at a laser power, pixel
dwell time, and signal accumulation that avoided detector (Power HyD S in counting mode)
saturation. One image per dLGN per section was acquired. For each animal overlap was
measured as described above on the right dLGN (the side where tdTomato signal
occupied the majority of the dLGN) by averaging from 5 consecutive sections.

Quantification of RGC densities in the retina

To estimate the density of RGCs in the retina (Figure S6), retinae of P10 *Casp3*[+/+]
and *Casp3*[-/-] mice were harvested and stained with anti-RBPMS antibody as described
above. We imaged 4 large square regions of interest (1.16 mm x 1.16 mm) in each retina
on four sides of the optic disc. The images were acquired with a Leica TCS SP8 Laser

Scanning Microscope using the 10x/0.4 NA dry objective (Leica HC PL Apo CS2 10x/0.4

dry), a 594 nm laser line, and an emission window of 607 nm to 694 nm, at a pixel

sampling of 302 nm x 302 nm. To account for tissue unevenness, we collected a stack of

3 Z-slices (2.4 μm apart). To count RGCs, we sampled 3 subregions of a size of 100 μm^2

from each large region of interest. The subregions were selected to be at similar distances

from the optic disc and in areas devoid of tissue tears. Cells were modeled to spots by

fitting them to ellipsoid objects (XY size = 8 μm; Z size = 16 μm, with background

subtraction) in Imaris 9.6 (Oxford Instruments). Densities of spots for each region were

averaged and reported as number of RGC per 100 μm^2 of retina.

Whole-cell patch-clamp recording in acute brain slices

To characterize electrophysiological properties of retinogeniculate synapses in

Casp3$^{+/+}$ and *Casp3$^{-/-}$* mice (Figure 4), the mice (aged p28 – p32) were decapitated under

deep isoflurane anesthesia, and the brain was removed and quickly transferred to an ice-

cold dissection solution containing (in mM): 194 sucrose, 30 NaCl, 2.5 KCl, 1.2 NaH2PO4,

26 NaHCO3, 10 D-Glucose, 1 MgCl2 (pH 7.4, oxygenated with 95% CO2 and 5% O2).

Parasagittal brain slices containing the optic tract and dLGN were prepared using a

vibratome (Leica VT 1200S) as previously described (*5, 6*). The slices were recovered at

35.5°C in an incubation chamber (BSC-PC, Warner Instrument, USA) filled with artificial

cerebrospinal fluid (ACSF) containing (in mM): 124 NaCl, 2.5 KCl, 1.2 NaH2PO4, 26

NaHCO3, 10 D-Glucose, 2 CaCl2, 1 MgCl2 (pH 7.4, oxygenated with 95% CO2 and 5%

O2, osmolarity ~310). After ~30 minutes of recovery, the chamber was maintained at room

temperature.

All recordings were performed with a MultiClamp 700B amplifier (Molecular

Devices) and signals were filtered at 2 kHz and digitized at 20 kHz with via USB-6343

(National Instruments) under the control of WaveSurfer software

(https://wavesurfer.janelia.org). All recordings were carried on slices submerged in the

recording chamber of an upright microscope (BX61WI; Olympus, Tokyo, Japan) equipped

with IR-DIC (infrared-differential interference contrast) microscopy and a water-immersion

objective lens (60X, 1.00 NA; Olympus). Slices were maintained under continuous

perfusion of oxygenated ACSF at 29-30°C. Patch pipettes were pulled from thin-wall

single-barrel borosilicate glass (TW150-6, WPI), resulting in electric resistance of 2 ~ 5

MΩ when filled with an intracellular solution contained the following (in mM): 120 cesium

methane sulfonate, 5 NaCl, 10 tetraethylammonium chloride, 10 HEPES, 4 lidocaine N-

ethyl bromide, 1.1 EGTA, 4 magnesium ATP, and 0.3 sodium GTP, with pH adjusted to

7.2 with CsOH and osmolality set to ~290 mOsm.

Whole-cell voltage-clamp recordings of relay neurons in the dLGN were performed

as previously described (Figure 4A-C) (*6*). The slices were submerged in ACSF

containing 20 uM bicuculline to inhibit $GABA_A$ receptors, and a bipolar concentric

stimulation electrode (FHC) was placed on the surface of the optic tract to deliver stimuli

with intensities ranged from 0 – 100 µA (200us duration) with an inter-trial interval of 40 –

60 s. In the whole-cell configuration (series resistance < 20 MΩ), the membrane potential

of the neuron was clamped at -70 mV to measure evoked excitatory postsynaptic current

(EPSC) mediated by activation of AMPA receptors, and then at +40 mV to measure

evoked EPSC mediated by activation of NMDA receptors. The stimulus intensity was

systemically increased until a response to the stimulus was observed, and then was

reduced to a previous level where no response was detected. From that level, the stimulus

intensity was increased by 0.5 µA steps to recruit single inputs which were putative single

fiber (SF) amplitudes. When the EPSC amplitude reached a plateau, it was considered a

putative maximum response. If the EPSC amplitude dropped while the stimulation

intensity increased, data from that relay neuron was discarded.

To measure paired pulse ratio (PPR) (Figure 4D-E), the slices were submerged in

the ACSF containing 20 uM bicuculline and 50 uM cyclothiazide to inhibit AMPAR

desensitization, and the stimulation electrode was placed on the optic tract. In the whole-

cell configuration, the membrane potential was clamped at -70 mV. After confirming the

stimulation intensity that evoked the maximum response, two consecutive electrical

stimuli of that intensity (300 µs duration) were delivered at 50, 150, 250, 500, and 1000

ms inter-stimulus intervals. Each pair of stimuli constitutes one trial, and two consecutive

trials were separated by an interval of 40 s. For each inter-stimulus interval, two trials

were performed and averaged. PPR was calculated as the maximum amplitude of the

second EPSC over the maximum amplitude of the first EPSC. If in any trial except for the

first trial the first EPSC dropped significantly in amplitude compared to the first EPSC of

the first trial, data from that dLGN relay neuron was discarded.

Measuring microglia- and astrocyte-mediated synapse engulfment *in vivo*

To measure microglia-mediated synapse engulfment *in vivo* (Figure 5 and Figure

S10), we bred *Casp3$^{-/-}$* males to *Casp3$^{+/-}$*; *Cx3ct1-Gfp$^{+/-}$* females to generate *Casp3$^{-/-}$*;

Cx3ct1-Gfp$^{+/-}$ and *Casp3$^{+/-}$*; *Cx3ct1-Gfp$^{+/-}$* littermate pups. To measure astrocyte-

mediated synapse engulfment *in vivo* (Figure S11), we bred *Casp3$^{-/-}$* males to *Casp3$^{+/-}$*;

Aldh1l1-Gfp$^{+/-}$ to generate *Casp3$^{-/-}$*; *Aldh1l1-Gfp$^{+/-}$* and *Casp3$^{+/-}$*; *Aldh1l1-Gfp$^{+/-}$* littermate

pups. P4 pups with desired genotypes were injected with 2 µL of CTB-AF555 in left eyes

and 2 µL of CTB-AF647 in right eyes as described above. 24 hours later, the brains were

harvested, sectioned, and mounted as described above.

To analyze engulfment of synapses, the sections were imaged using a Leica TCS

SP8 Laser Scanning Microscope equipped with an HC Plan Apo CS2 40x (NA = 1.3) oil

objective and a White Light Laser Unit. The 3D multichannel stacks with a voxel size of

0.075 x 0.075 x 0.312 µm for microglia samples or 0.048 x 0.048 x 0.2 µm for astrocytes

samples were acquired as 16-bit images with 4x line averages for microglia or 2x for

astrocytes at 400 Hz scanner speed. The stacks were set to collect a volume of 20

micrometers in the Z dimension and 387.5 micrometers in XY dimensions for microglia or

288 µm in XY for astrocytes. The pinhole was set at 1 Airy unit for 520 nm

wavelength. The excitation lasers and detection windows were set to minimize crosstalk

and bleed-through as well as the saturation of detectors while simultaneously imaging the

three channels. A 478 nm laser and a detection window of 488 nm – 537nm were used

for the GFP channel, a 550 nm laser and a detection window of 596 nm – 629 nm were

used for CTB-AF555, and a 651 nm laser and a detection window of 665 nm to 750 nm

were used for CTB-AF647. Imaris 9.6 (Oxford Instruments) was used to segment

microglia, astrocytes, ipsilateral RGC axon terminals, and contralateral RGC terminals

from the 3D stacks using the Surface Creation module with background subtraction. For

segmentation of RGC axon terminals and astrocytes, all parameters used were left at

default values (auto). For segmentation of microglia, we set the threshold manually at the

highest inflection curve in the histogram of voxel intensities, which in our experience

provided a good trade-off for good segmentation of cell bodies and major processes while

keeping the adjustment consistent across different image datasets. A background

subtraction ball size of 1 µm was used for microglia, a ball size of 1.4 µm was used for

astrocytes, and a ball size of 0.5 µm was used for AF555 and AF647 labeled axonal

materials. The segmented signal of AF555 and AF647 was each converted to a binary

mask of 0 and 1 voxels. Finally, the sum of the segmented volume (i.e. sum of ones) of

AF555 and or AF647 in each microglia or astrocyte was either directly plotted or

normalized to the microglia or astrocyte volume and plotted. To control the inherent signal

variability across LGN, only microglia and astrocytes present in the ipsilateral region of

the 3 middle sections of the left dLGN of each mouse were included in the analysis. Also,

only microglia or astrocytes whose soma and proximal processes were contained in the

volumetric stack were included in the final analysis. All littermate pups of the same

genotype were pooled together.

Measuring activity-dependent engulfment of synapses by microglia

To investigate the role of caspase-3 in activity-dependent synapse elimination

(Figure 6 and Figure S12), we used C57Bl/6J wildtype mice as controls and bred *Casp3*-

/- males with *Casp3*+/- females to generate *Casp3*-/- mice. Note that C57Bl/6J and *Casp3*-/-

mice used in this experiment are not littermates and should not be directly compared with

each other. C57Bl/6J and *Casp3*-/- embryos were injected at E15 in their right eye either

with AAV carrying hSyn-TeTxLC-P2A-mTurqouise2 or AAV carrying hSyn-mTurquoise2

as described above. To label the RGC axons, at P4, the pups were injected with 2 µL of

CTB-AF555 in left eyes and 2 µL of CTB-AF647 in right eyes as described above. At P5,

the brains were harvested and sectioned. Two 50 µm sections that spanned the central

part of the dLGN were stained with anti-Iba1 antibody and goat anti-rabbit AF488

secondary (as described above) to label microglial cell bodies.

Mounted and cured sections were then imaged on Leica Stellaris 8 confocal

microscope using a 63x/1.4 NA oil objective (Leica HC PL APO 63x/1,40 OIL CS2). Four

stitched 3D-stacks were acquired for each left dLGN (the side where CTF-AF647 and

mTurquoise2 occupied the majority of the dLGN) to generate a 3D volume of 366 μm x

366 μm in XY and approximately 20 μm in the Z dimension, with a voxel size of 64 nm x

64 nm x 299 nm. The channels were acquired sequentially: first using a 499 nm laser with

an emission window of 507-556 nm together with a 653 nm laser with an emission window

of 663-750 nm for Iba1-AF488 and CTB-AF647 signal, respectively, then using a 553 nm

laser with an emission window of 558-663 nm for the CTB-AF555 signal. Images acquired

were collected as 12-bit datasets, and care was taken to set the laser power, signal

accumulation, and pixel dwell time to avoid detector saturation. Only animals whose

dLGN areas were fully labeled with both AF555 and AF647 as well as the mTurquoise

signals were imaged and analyzed. Animals in which CTB signal appeared widespread

throughout the brain parenchyma were excluded from the analysis as these labeling

patterns were likely due to surgical mistargeting or issues with tissue integrity.

To segment volumes corresponding to microglia and CTB-AF555 and CTB-AF647

labeled RGC axon terminals, we utilized the Surface Creation module with machine

learning segmentation of Imaris 10.1.1 (Oxford Instruments). We first trained for several

rounds a pixel classifier for each channel using two independent image files and saved

the classifiers along with the other settings (smoothing grain size of 100 nm and

discarding volumes smaller than 10 voxels) as "Favorite Creation Parameter".

Subsequently, the same Favorite Creation Parameters were applied to all data sets, with

the rare instances when microglia segmentation was slightly tuned via the same pixel

classier to better capture some of the cell border areas. Similar to the previous analyses

(Figure 5), the segmented signals of CTB-AF647 and CTB-AF555 were converted to

binary images (0 or 1 voxels), and the total volume of CTB-AF647- or CTB-AF555-positive

voxels in each microglia was extracted, converted to fraction of the microglia volume, and

plotted. Only the cells whose body and major processes were bound within the 3D stack

and present within the middle area of dLGN (where the ipsilateral signal is present) were

included in the analysis. Data from the same genotype and treatment were pooled

together. Either one or two sections were analyzed for each animal.

Measuring microglia activation

To measure if lack of caspase 3 causes microglia activation (Figure S9), we bred

Casp3$^{+/-}$ males with *Casp3$^{+/-}$* females to generate *Casp3$^{-/-}$* and Casp3$^{+/+}$ littermates. After

genotyping, at P5, the brains were harvested and sectioned coronally as described above.

Two 50 μm sections that spanned the central part of the dLGN were stained with anti-

Iba1 antibody (detected by goat anti-rabbit AF488 secondary) to label microglial cell

bodies and anti-CD68 antibody (detected by donkey anti-rat AF555 secondary) to detect

CD68 protein. Mounted and cured sections were imaged on Leica Stellaris 8 confocal

microscope using a 63x/1.4 NA oil objective (Leica HC PL APO 63x/1,40 OIL CS2). A 3D

image stack of 185 μm x 185 μm in XY and approximately 20 μm in the Z dimension, with

a voxel size of 72 nm x 72 nm x 299 nm was acquired for each dLGN. The channels were

acquired in photon counting mode simultaneously: a 499 nm laser with an emission

window of 504-548 nm together with a 553 nm laser with an emission window of 560-650

nm for Iba1-AF488 and CD68-AF555. To eliminate any fluorescence emission bleed-

trough between channels, the photon arrival times were gated to 3 - 5 ns for Iba1-AF488

and to 0 - 1.5 ns for Iba1-AF555 channels, respectively. Images acquired were collected

as 8-bit datasets, and care was taken to set the laser power, signal accumulation, and

pixel dwell time to avoid fluorophore bleaching or detector saturation.

For segmentation of microglia and CD68 signals, we utilized the Surface Creation

module with machine learning segmentation of Imaris 10.1.1 (Oxford Instruments). We

first trained for several rounds a pixel classifier for each channel then saved the classifiers

along with the other settings (smoothing grain size of 145 nm and discarding volumes

smaller than 10 voxels) as "Favorite Creation Parameter". Subsequently, the same

Favorite Creation Parameters were applied to all data sets, with the rare instances when

microglia segmentation was slightly adjusted via the same pixel classier to better capture

some of the cell border areas. Similar to the previous analyses (Figure 5), the segmented

signals of CD68-AF555 were converted to binary images (0 or 1 voxels), and the total

volume CTB-AF555-positive voxels in each microglia was extracted, converted to fraction

of the microglia volume, and plotted. For measurement of relative CD68 intensity, the sum

of CD68 signal was divided by the microglia volume. Only the microglia whose body and

major processes were confined within the 3D stack were included in the analysis. All

staining, imaging and analysis parameters were identically applied to all sections. Two

sections were analyzed for each animal. Data from the same genotype and were pooled

together.

Characterization of *App/Ps1* transgenic mice

To detect Aβ deposition (Figure S13A), brains of 6 month-old female *App/Ps1^-/-*

mice, 5 month-old and 6 month-old male and female *App/Ps1^+/-* mice, and 6 month-old

female *Casp3^-/-*; *App/Ps1^+/-* mice were harvested, sectioned, and stained with Thioflavin

S as described above. Stained sections were imaged on a Leica Stellaris 8 microscope

using a 10x/0.4 NA dry objective (Leica HC PL Apo CS2 10x/0.4 dry). A 460 nm laser line

was used for excitation and a window of 469 nm - 600 nm was used for emission. Tile

images (pixel size: 284nm x 284 nm) were collected and merged into one image spanning

an entire coronal section in Las X 4.6 Navigator. To quantify the level of amyloid deposition

in 6 month-old female *App/Ps1^+/-* and *Casp3^-/-*; *App/Ps1^+/-* mice (Figure S13B), the

number of Thioflavin S-positive plaques was manually counted in a total of 3 sections per

animal and averaged.

To investigate the role of caspase-3 in Aβ-induced synapse loss (Figure 7), *Casp3^-*

^/- mice were bred to *App/Ps1^+/-* mice to create *Casp3^-/-*; *App/Ps1^-/-* and *Casp3^-/-*; *App/Ps1^+/-*

mice. Brains of 6 month-old female *App/Ps1^-/-* and *App/Ps1^+/-* littermates as well as *Casp3^-*

^/-; *App/Ps1^-/-* and *Casp3^-/-*; *App/Ps1^+/-* littermates were harvested, sectioned, and stained

with anti-SV2 antibody (conjugated to AF546) and anti-Homer1 antibody (using goat anti-

rabbit AF647 as the secondary) to label presynaptic and postsynaptic compartments,

respectively. Stained sections were imaged on a Leica Stellaris 8 confocal microscope

using a 63x/1.4 NA oil objective (Leica HC PL APO 63x/1,40 OIL CS2). To acquire a

representative dataset, we focused on the molecular layer of the dentate gyrus of the

hippocampus from both the left and right side of the brain and sampled 3 regions from

each side, as shown in Figure S14. 3D-stacks (61.5 μm x 61.5 μm x 20 μm) were acquired

with a voxel size of 64 nm x 64 nm x 298 nm. 561 nm and 638 nm laser lines were used

for excitation, 555 nm – 635 nm and 642 nm – 792 nm windows were used for emission

reading, and appropriate notch filters were used for blocking the exciting laser lines. The

channels were acquired as 8-bit data sets with detectors set in counting mode, and the

lasers power, signal accumulation, and pixel dwell time set at levels that avoided detector

saturation. To measure synapse number per unit volume, we first trimmed the raw 3D

stacks to a stack of approximately 3.5 μm thickness on the side closest to the coverslip

(to avoid any signal depletion the middle part of the section due to antibody penetration

issues) and cropped out regions where large gaps (corresponding to blood vessels)

devoid of Sv2 or Homer1 staining were present. The punctate signals of SV2 and Homer1

were fit in Imaris 10.1 (Oxford Instruments) to ellipsoids (500 nm in XY and 850 nm in Z

for SV2; 400 nm in XY and 800 nm in Z for Homer1) with background subtraction ("Spot

Creation"). Then we designated as "synapses" all Homer1 ellipsoids that were within 300

nm of a SV2 ellipsoid (center-to-center distance). For each animal, the total number of

synapses in all 3D-stacks analyzed was divided by the total volume of all stacks analyzed

and plotted as a single value.

To detect Aβ-induced microglia activation (Figure S16), brains of 6 month-old

female $App/Ps1^{-/-}$, $App/Ps1^{+/-}$, and $Casp3^{-/-}$; $App/Ps1^{+/-}$ mice were harvested, sectioned,

and stained with anti-Iba1 antibody (using goat anti-rabbit AF647plus as the secondary)

and anti-Cd68 (using goat anti-rat AF555plus as the secondary) as labels of microglia cell

body and microglial activation, respectively. Stained sections were imaged on a Leica

Stellaris 8 confocal microscope using a 20x/0.7 NA water objective (Leica HC PL APO

20x/0,7 IMM CORR CS2). Tile images spanning the hippocampus were acquired at voxel

sizes of 142 nm x 142 nm x 300 nm. A 548 nm laser line and a 554 nm – 634 nm emission

window were used for the CD68-AF555 channel, and a 653 laser line and a 663 nm –

748 nm emission window were used for the Iba1-AF647 channel. Microgliosis could be

detected as clusters of microglia with stubby processes and strong intracellular Cd68

signals (Figure S16).

To quantify Aβ-induced caspase-3 activity (Figure S17), brains of 4 month-old, 5

month-old and 6 month-old female *App/Ps1*$^{-/-}$ and *App/Ps1*$^{+/-}$ littermates were harvested,

sectioned, and stained with anti-cleaved caspase-3 antibody (using goat anti-rabbit

AF555plus as the secondary). Stained sections were imaged on a Leica Stellaris 8

confocal microscope using a 63x/1.4 NA oil objective (Leica HC PL APO 63x/1,40 OIL

CS2). We sampled at least 2 regions of interest from the molecular layer of the dentate

gyrus of the hippocampus on each side of the brain. 3D-stacks (92.26 μm x 92.26 μm x

22 μm) were acquired with a voxel size of 81 nm x 81 nm x 299 nm. A 561 laser line was

used for excitation and an emission window of 550 nm – 728 nm was used for the active

caspase-3 channel. To measure the number of caspase 3 objects in the dataset, we first

trimmed the 3D stacks to a volume where the signal is uniformly distributed (i.e. in focus)

and segmented the objects using the "Surface creation" analysis in Imaris 10.1 (Oxford

Instruments). The dataset was first smoothed by a two-pixel filter, then a ball size of 400

nm was used to find the objects with background subtraction. Only objects bigger than 10

and smaller than 2000 voxels were kept. Finally, the number of caspase 3 objects was

divided by the volume of the 3D stack analyzed and reported per brain side (left or right)

for each animal.

To measure the relative density of granule cells in the dentate gyrus (Figure S16), brains

of 6 month-old female *App/Ps1*$^{-/-}$, *App/Ps1*$^{+/-}$, *Casp3*$^{-/-}$; *App/Ps1*$^{-/-}$, and *Casp3*$^{-/-}$; *App/Ps1*$^{+/-}$

mice were harvested, sectioned, and stained with anti-NeuN antibody (using goat anti-

mouse AF594plus as the secondary) to label neuronal nuclei. Stained sections were

imaged on a Leica Stellaris 8 confocal microscope using a 20x/0.7 NA water objective

(Leica HC PL APO 20x/0,7 IMM CORR CS2). 3D stacks of the dentate gyrus (581 μm x

581 μm x 30 μm) were acquired at voxel sizes of 160 nm x 160 nm x 1000 nm for each

section and side (left and right) using a 594 nm laser and an emission window of 597 nm

to 750 nm. The channels were acquired as 8-bit images with the detector set in counting

mode, and the lasers power, signal accumulation, and pixel dwell time set at levels that

avoided detector saturation. To count the number of nuclei, images were first rotated to

position vertically the cortical side of the dentate gyrus. A maximum projection of the

brightest two Z-slices was extracted from each 3D stack, and NeuN+ nuclei were

manually counted within a rectangular region of interest (89 μm x 150 μm) that was

positioned on the granule cell layer at 300 μm away from the tip of the gyrus.

References:

1.   J. Goedhart *et al.*, Structure-guided evolution of cyan fluorescent proteins towards a

quantum yield of 93%. *Nat Commun* **3**, 751 (2012).

2.   D. Grimm *et al.*, In vitro and in vivo gene therapy vector evolution via multispecies

interbreeding and retargeting of adeno-associated viruses. *J Virol* **82**, 5887-5911

(2008).

3.   M. Yasuda, S. Nagappan-Chettiar, H. Umemori, In utero intraocular AAV injection for

early gene expression in the developing rodent retina. *STAR Protoc* **2**, 100742 (2021).

4.   J. D. M. Arzt, C. Schmied, T. Pietzsch, D. Schmidt, P. Tomancak, R. Haase, F. Jug,

LABKIT: Labeling and Segmentation Toolkit for Big Image Data. *Frontiers in Computer*

Science **4**, (2022).

5.   J. P. Turner, T. E. Salt, Characterization of sensory and corticothalamic excitatory

inputs to rat thalamocortical neurones in vitro. *J Physiol* **510 (Pt 3)**, 829-843 (1998).

6.   C. Chen, W. G. Regehr, Developmental remodeling of the retinogeniculate synapse.

Neuron **28**, 955-966 (2000).