1 **Assessing nest sites, survival rates and population dynamics of free-living**

2 **honeybee colonies in Germany: A comparative study using personally**

3 **collected and Citizen Science data**

Benjamin Rutschmann[1,2,3,§,*], Felix Remter[1,4,5,§,*], Sebastian Roth[1,§]

[1] BEEtree-Monitor, Munich, Germany

[2] Agroecology and Environment, Agroscope, Reckenholzstrasse 191, 8046, Zurich,

Switzerland

[3] Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg,

Würzburg, Germany

[4] Rachel Carson Center for Environmental Humanities, LMU Munich, Munich, Germany

[5] Department of Science, Technology and Society (STS) at the TUM School of Social

Sciences and Technology, TU Munich, Munich, Germany

[§] these authors contributed equally

[*] Correspondence: benjamin.rutschmann@agroscope.admin.ch,

felix.remter@primatevisions.de

ORCiD:
BR: 0000-0001-6589-6408
FR: 0000-0002-9781-3811
SR: 0000-0003-0576-8484

ABSTRACT

Our understanding of the western honeybee (*Apis mellifera*) predominantly stems from studies conducted within beekeeping environments, leaving the presence and characteristics of honeybees outside managed settings largely unexplored. This study focuses on free-living colonies, examining their habitats, nesting sites, and survival rates, thereby highlighting the liminal state of the species in Europe. Through the BEEtree-Monitor project, we personally monitored (PM) nest sites in Munich (N=107) and coordinated Citizen Science (CS) monitoring across Germany (N=423), resulting in 2,555 observations on 530 colonies from 311 participants over 7 years. While habitat preferences differed between urban, rural and forested areas, we found that 31% of the occupied nest sites were in buildings and 63% in mature trees with cavities, including lime (*Tilia* spp., 18%), beech (*Fagus sylvatica*, 14%), oak (*Quercus* spp., 13%), and ash (*Fraxin*us excelsior, 11%). On average, only 12% of the PM colonies in Munich survived annually, according to model predictions based on observed data. Consequently, we found a minimum free-living colony density per km² of 0.06 in spring but 0.42 in summer, accounting for at least 4% of the total Munich population during the summer. Comparing the two monitoring approaches (PM vs. CS) and published data revealed significant discrepancies: survival rates reported by CS were markedly higher (model prediction: 28%), than PM and other published studies. We found that CS not only yielded significantly fewer updates per colony, but also that 76% of CS reports noted active colonies compared to 42% from PM, indicating an underreporting of abandoned sites. To ensure the reliability of survival data in CS projects, regional swarming should be monitored, and the timing of reports needs particular attention as 46% of reports about overwintering were too late (i.e. after the onset of the swarming season; 11% in PM). CS data proved to be highly valuable for nest site and habitat analysis but require a strict validation protocol for survival statistics. This study enhances our understanding of the ecological dynamics and conservation needs of free-living honeybee cohorts, addresses potential monitoring biases, and suggests standardized data collection protocols for future monitoring projects.

KEYWORDS

wildlife monitoring, *Apis mellifera*, urban ecology, wild pollinators, liminal species, wildlife conservation, wild-living honeybees, multi-species studies

INTRODUCTION

The western honeybee (*Apis mellifera*) holds a significant place in the European entomofauna, facilitating the reproduction and genetic diversity of countless plant species, including many agricultural crops (Garibaldi et al., 2013; Breeze et al., 2014; Hung et al., 2018). Among researchers and beekeepers alike, *Apis mellifera* tends to be perceived solely as a domesticated animal, found and researched under managed conditions. Consequently, most of our comprehensive understanding of the western honeybee as a species, its behaviour, and its ecology predominantly stems from research conducted with colonies under beekeeping conditions, while wild honeybee populations have been neglected in the modern apidological tradition (Stoeckhert, 1954; Kohl and Rutschmann, 2018; Seeley, 2019; Requier et al., 2019). The introduction of the parasite *Varroa destructor* (Anderson and Trueman, 2000) from Asia to Europe in the 1970s intensified this oversight, as the mite's infestations suggested that only human-managed colonies receiving miticide treatment could survive (Rosenkranz et al., 2010; Meixner et al., 2015).

Yet, this belies the fact that the western honeybee is also present outside the realms of beekeeping and human husbandry (Grindrod and Martin, 2021; Visick and Ratnieks, 2023a) and how little is known about the formation and dynamics of this cohort. Currently, stable wild honeybee populations within their original range are known to exist in Africa and the Southern Ural, and outside their original range in the Americas and Australia (Schneider and Blyther, 1988; Moritz et al., 2007; Ilyasov et al., 2015b; Ratnieks et al., 1991; Seeley, 2007; Oldroyd et al., 1997). Studies on these populations usually focus on habitats where managed and non-managed colonies are relatively separated, or where the density of wild colonies matches or surpasses that of managed colonies (Visick and Ratnieks, 2023a). However, the situation in Europe is different: Due to the fragmented landscape (Ibisch et al., 2016; Lesiv et al., 2019) and the high density of managed colonies (Phiri et al., 2022; Jones, 2004), there likely isn't a spatial or genetic barrier between managed and free-living colonies in most parts of Europe. Hence, the key differences between free-living colonies and managed ones lie in their nest site ecology and their modus vivendi. Therefore we refer to both as *cohorts* of one local honeybee population, that is neither *wild* nor *domesticated*, but in a yet *liminal* state.

Recently, news about stable populations of western honeybee colonies outside beekeeping raised the interest of beekeepers in how to escape treatment (Seeley, 2019; Remter, 2021a) and spurred considerable research in this field. Free-living colonies have been discovered in

various environments, ranging from forests, to electric power poles in agricultural landscapes and rural and urban areas (Oleksa et al., 2013; Kohl and Rutschmann, 2018; Requier et al., 2020; Oberreiter et al., 2021; Kohl et al., 2022; Rutschmann et al., 2022; Hassett et al., 2018; Browne et al., 2020; Moro et al., 2021; Bila Dubaić et al., 2021; Lang et al., 2022; Visick and Ratnieks, 2023b; Cordillot, 2024). Due to their hidden locations in cavities high above the ground, free-living colonies lead secretive lives (Kohl and Rutschmann, 2018; Remter, 2021b): finding them and repeatedly collecting data in numbers high enough for statistical inference is extremely time-consuming and requires specialized skills and equipment (Kohl and Rutschmann, 2018; Kohl et al., 2022).

One approach that has garnered significant attention in the study of wild animals and biodiversity monitoring in general is the utilization of Citizen Science (Pocock et al., 2018; Fraisl et al., 2022; Koffler et al., 2021; Weissmann et al., 2023). As in the case of free-living honeybee colonies Citizen Science offers the advantage of enlisting the help of many individuals who share the task of finding and monitoring colonies, thus extending the geographic and temporal scope of the research beyond what researchers could achieve alone (Henneken et al., 2012; Lesiv et al., 2019; Hsing et al., 2022). To understand the free-living honeybee cohort more completely, it is crucial to examine not only their nesting sites but also the survival rates which are critical for assessing their genetic contribution to the local honeybee population. However, most of the studies on free-living honeybees in Europe have not systematically monitored individual colony survival, and have instead reported on nesting sites without comprehensive knowledge of the life histories of individual colonies (but see (Rutschmann et al., 2022; Kohl et al., 2022; Lang et al., 2022; Cordillot, 2024). Also, although previous studies have acknowledged the importance of Citizen Science in data collection (Moro et al., 2021; Bila Dubaić et al., 2021), none have yet investigated the quality of the data generated by Citizen Science and how to validate such reports.

We present data from a web-based monitoring scheme (BEEtree-Monitor) developed to study the habitats, nesting sites, and life histories of free-living honeybee colonies. In a parallel undertaking we collected data in the Munich region personally and via Citizen Science in Germany, both according to the same protocol. Over a span of seven years (2016-2023), we collected various parameters of 530 nest sites together with longitudinal occupation data: 107 nest sites monitored by ourselves and 423 by citizen scientists. We compare survival rates between these two groups to evaluate the validity and potential biases of in the Citizen

114 Science reports and methodology. Through this study, we aim to provide a broader

115 understanding of honeybees as a species existing outside human husbandry and offer

116 guidelines to leverage future Citizen Science projects for effective monitoring of free-living

117 honeybee colonies.

METHODS

<u>Data collection and data curation</u>

The monitoring and data collection process for this study was implemented through a

combination of personal monitoring, (PM) and reports by volunteering supporters with highly

diverging skill levels and knowledge sets (citizen science monitoring, CS). Leveraging our

own experiences in surveying free-living honeybee colonies and third-party reports, we

developed an advanced monitoring scheme tailored to our target groups. Initially, participants

were asked to fill in a designated form and send it to us via email (BEEtrees HOBOS form,

supplementary material (SM)). Later on, we constructed an online platform specifically

tailored for Citizen Science monitoring, launched as a website (BEEtree-Monitor;

www.beetrees.org) in 2018. This platform was created entirely from open-source software

(see SM for further information on the software and online form).

The CS recruiting was facilitated through social media outreach, presence in public media

and beekeeping journals. The community was maintained via regular newsletters with

guiding and motivating information. To enable as many volunteers as possible without

offering a special training, our protocol focused on location, easily measurable nest site

parameters, and continuous, repeated observations with specific date, time and focus. The

citizen scientists therefore contributed by collecting part of the data that we used but did not

contribute to the analysis and interpretation of that data. In total only 36 nesting sites were

found by the authors (and 494 by citizen scientists), however 107 locations were actively

monitored by us (and 423 by citizen scientists). Our primary analysis focused on colonies

located in Germany and several nearby Central European countries, including Switzerland

(N=9), Austria (N=3), Czechia (N=2), and Luxembourg (N=3). To mitigate the influence of

markedly different environmental conditions that could confound our study's findings, we

excluded observations of ten free-living colonies from geographically distant countries like

France, the UK, Italy, Spain, Norway, and Ukraine.

Precise GPS coordinates allowed us to investigate the nesting and foraging habitats of the colonies, and the nest site parameters were used to analyse the swarms' preferences for different nest types such as hollows in building structures (e.g., chimneys, window blind boxes or facade compound insulations) or hollows in different tree species. Additionally, we sought information on entrance directions and height, as well as the trunk diameter for trees. The online platform also featured an open-text/commentary field for participants to provide other relevant information.

To further understand the survival and behaviour of these colonies, we applied two distinct monitoring approaches (Figure 1A):

PM: This method entailed meticulous checks for visual signs of pollen import (indicating brood production, marking the colony as live) and frequent short-term observations of nest sites exhibiting ambiguous flight patterns or no observed foraging for pollen. We filmed some of the typical or ambiguous situations to collectively analyse and relate them to continuous observations at the site. This monitoring protocol substantiated our classification of flight patterns and the ideal standard we aimed to achieve with CS.

CS: This approach involved providing participants with protocols and guidance to e.g. prioritize pollen import observation. Accessible explanations in the online form and regular emails encouraged detailed reporting during key observation periods.

Analysis of nesting and foraging habitats of reported free-living colonies

To classify the nesting and foraging habitats of the reported free-living colonies, we imported the coordinates of each colony's location into QGIS version 3.16.2 (QGIS Development Team, 2020) and performed intersections (using point layers for nesting habitats and 2 km buffers for foraging habitats) with the land cover classes from the CORINE Land Cover map 2018 (European Environment Agency, 2018). We grouped different land cover types together and quantified the proportional contributions of five major land cover types: urban areas, cropland, grassland, deciduous forest, and coniferous forest (see SM for further details). These land cover types were for some of the analysis further grouped into three broader categories: Urban (urban areas), rural (cropland and grassland), and forest (deciduous and coniferous forests). In the case of the foraging habitat, we chose a radius of 2 km as approximately 80% of honeybee foraging occurs within this distance (Rutschmann et al. 2023). Moreover, the landscape within the 2 km scale has been shown to measurably

influence honeybee colony performance, affecting factors such as foraging rate, colony
growth, and winter survival (Steffan-Dewenter and Kuhn, 2003; Sponsler and Johnson, 2015;
Rutschmann et al., 2022, 2023).

<u>Observation scheme for colony survival statistics: Timing, pollen import and flight patterns</u>

We defined survival as an instance when a cavity occupied from summer (during or after the
swarming season) remained occupied until the following spring before the next swarming
season commenced (Figure 1B). One valid report of an active colony before the start of the
swarming season (1st Check) would be proof for an overwintering survival. Additionally, we
implemented and encouraged participants to conduct one or two more annual checks: right
after the end of swarming (2nd Check) and in Autumn before the winter (3rd Check). Post-
swarming checks at all known cavities (including the recently unoccupied ones) served
finding the new founder colonies for the further monitoring while the Autumn check (3rd
Check) detects summer deaths. In our study, colonies that died in summer/autumn were
attributed as perished to the survival statistics of the following year. For assessing
overwintering, observations typically take place in March or early April where it is crucial
that weather conditions are suitable for honeybee foraging (e.g. no rain and temperatures
above 12° C. [Kevan and Baker, 1983]). Consequently, observations without suitable weather
conditions for honeybee foraging were excluded. As only life histories recorded in the
database were considered for our analysis, oral reports or in retrospect reports (of colonies
living for "several years" in the same cavity) were considered hearsay and not included. Data
pre winter 2016/2017 were discarded due to small sample sizes , and nesting sites that were
destroyed were not considered for the survival statistics.

A potential pitfall of reporting mere flight activity at the entrance is that certain behaviours
such as robbing, or the presence of scout bees may be erroneously interpreted as signs of a
living colony by less experienced observers. Therefore pollen import into a colony was used
as an indicator of brood production, hence designating it as a live colony. However, if pollen
deposition is not visible, a death is not generally evident, either because pollen is scarce, not
needed for the bees or due to the difficulty in observing pollen import in colonies with nest
entrances that are high above ground level. Consequently, under certain conditions, even in
the absence of visible pollen import, we still considered the colony to be alive. These criteria
included:

Figure 1: A) Flowchart of the monitoring protocol for Citizen Science (CS) and personal monitoring in Munich (PM). The workflow details the steps for recording nest site parameters and colony status. The inclusion of location descriptions, weather comments, general comments, and supplementary materials like pictures and videos helps ensure comprehensive and accurate data collection in CS (light blue). Personal monitoring yielded reliable information regarding flight activity types, which were further analysed with video analysis (light green). B) Annual monitoring schedule for free-living honeybee colonies. The monitoring schedule outlines three critical observation windows: 1st Check Window (before the onset of the swarming season in late March and April), 2nd Check Window (post-swarming season in July), and 3rd Check Window (in autumn, September). These checks focus on different survival metrics: Spring survival and reoccupations, summer and winter survival, respectively. Citizen scientists (CS) are mobilized before each check window to ensure timely data collection. The onset of swarming shifts each year and must be monitored closely to ensure accurate data on colony survival. The inlet picture shows pollen import into the colony an indication for brood production (photo: Felix Remter).

1. Observation of foraging flight patterns: Colonies were classified as active if regular
and/or directional flight activity was reported convincingly in the commentary
section. These behaviours are differentiated from non-foraging activities based on
specific patterns observed in PM:
- Scout bees typically exhibit distinct behaviours such as taking time to land,
thoroughly exploring the entrance before entering the cavity, performing slow
orientation flights during departure, and might defend the entrance against
other scouts when near swarming.
- Robbing bees display violent interactions with defenders if attempting to rob a
living colony. Initially, robbers perform orientation flights akin to scouts but
may also demonstrate a distinct 'bouncing off' landing pattern. Although
robbers can temporarily mimic regular foraging patterns, these are
distinguishable by experienced observers.
2. Reliability of observer comments: Assessments were also based on the reliability of
observers' comments. If the observer was suspected to credibly discern differences in

flight patterns indicative of foraging rather than robbing or scouting due to a

competent comment given, the colony was considered active.

3. <u>Frequency and timing of observations</u>: Multiple observations made at short intervals

that consistently indicated foraging behaviour, as opposed to scouting or robbing,

supported the classification of a colony as active.

<u>Survival rates of free-living honeybee colonies and the impact of monitoring type and year</u>

To analyse colony survival statistics across different years in Germany we merged datasets

from this study (CS and PM) and from two published studies (Kohl et al., (2022) and Lang et

al., (2022)). Each dataset employed a distinct monitoring approach (see SM for further

details). To ascertain the influence of various predictors on the odds of colony survival, we

compared several mixed-effects logistic regression models. These models were constructed

using the glmmTMB package in R (Magnusson et al., 2017). Model 1 included 'monitoring

type' as a fixed effect (with the factor levels CS, PM , Kohl et al. 2022 and Lang et al. 2022)

and 'colony id' nested within 'year' as a random effect to account for repeated measures on the

same subjects across years. Model 2 included both 'monitoring type' and 'year' (from 2016 till

2023) as fixed effects, with 'colony id' as a random effect. Model 3, which did not converge,

included the interaction between 'monitoring type' and 'year' as fixed effects, with 'colony id'

as a random effect. The model selection process was guided by the Akaike Information

Criterion (AIC) and the 'emmeans' package (Lenth and Lenth, 2018) was used for post-hoc

comparisons. Residuals of the models were inspected with DHARMa package (Hartig and

Hartig, 2017).

<u>Exemplary estimation of free-living honeybee colony density</u>

Estimating the density of free-living honeybee colonies poses challenges due to the likelihood

of substantial underreporting. To address this, we concentrated our density estimation efforts

on the city of Munich, where the majority of our known free-living honeybee colony

locations are situated (further details on the density estimation can be found in the SM).

<u>Onset of regional swarming</u>

In 2019, two of the authors (SR and FR) established a website across Germany and a hotline

in Munich for the discovery of honeybee swarms and their potential capture. This enabled us

to amass substantial data (N=376) on the initiation of swarming activity in the years from

2019 to 2023 within the geographic extent of the study. These years also represent the period

during which most of the nest sites presented in this study were found (N=378 out of 530

colonies) and monitored (N=307 out of 350 life history reports) for of this study. For each

year, the first reported swarm served as a conservative estimate of the commencement of the

swarming season. We observed variability in the yearly onset of swarming over the five-year

period from 2019 to 2023. Specifically, swarming began on 17 April in 2019, 6 April in

2020, 8 May in 2021, 28 April in 2022, and 23 April in 2023 . Hence, the time difference

between the earliest and the latest recorded yearly swarming onset was more than one month

(32 days), suggesting it should be taken into account in planning of CS initiatives and

survival analyses of free-living honeybees. Especially critical are years where swarming

occurs unexpectedly early (e.g. year 2020). A small subset of the free-living colony reports in

this study stem from years before 2019 in which we lacked empirical data on the beginning of

the swarming season. Hence, we conservatively selected mid-April as the presumed start of

swarming for these years (Henneken et al., 2012).

<u>Statistical analysis of directional and height preferences in cavities</u>

To evaluate directional preferences of honeybees when selecting cavities, we recorded the

entrance orientation of nesting sites occupied by free-living colonies in the eight cardinal and

intercardinal directions (N, NE, E, SE, S, SW, W, NW) for both tree cavities and cavities in

buildings and assessed whether the distribution of orientations was non-random by

employing the Rayleigh test for circular statistics using Directional package in R (Tsagris et

al., 2016). The height of cavity entrances in trees and building structures was investigated

with a non-parametric Mann-Whitney U Test.

Further information on statistical analysis regarding number of reports per colony, reported

colony status and timing of the reports can be found in the supplementary material.

All statistical analyses were performed using R software (version 4.3.1; R Core Team, 2016).

For data wrangling and graphical representation of the results, we utilized 'tidyverse' and

'ggplot2' (Wickham, 2017, 2016).

RESULTS

Nesting and foraging habitat of reported free-living colonies

We garnered responses from 311 participants (including the authors), providing us with 2,555 observations on 530 colonies (Figure 2A). Using landcover data to assess the habitat of the free-living colonies we found 57 percent of the reported colonies to dwell in urban areas with high human density, 14 percent in deciduous forest, 14 percent in cropland, 9 percent in grassland and 5 percent in coniferous forest (Figure 2B). The distribution of nesting habitats for free-living honeybee colonies was significantly different from the proportional distribution based on land cover types in Germany[1] ($\chi^2 = 99.81$, df = 4, p < 0.001). Specifically, colonies were disproportionately more often found in urban areas relative to the habitat availability in Germany. Calculating the available foraging habitat 2 kilometres

Figure 2: A) Geographic distribution of the free-living colonies reported to the BEEtree-Monitor overlaid on the main land cover types within Germany (CORINE Land Cover map 2018; 3 colonies are hidden by the inlet). The inlet shows the free-living colonies in the Munich region. B) Proportional distribution of nesting habitats where free-living colonies have been reported in comparison to the relative distribution of different land cover types across Germany depicted by black diamond symbols. C) Foraging habitats that is available to the bees (habitat within a 2 km radius around each reported free-living honeybee colony). Again, black diamond symbols show the relative distribution of a land cover type across Germany.

---

[1] The proportional distribution assumes honeybees have no habitat preferences and their discovery is equally probable in all habitats.

around each nest site, where colonies are operating to find their food, we found similar

patterns. We estimated 41 percent of the available foraging habitat of the reported colonies to

be urban areas, 27 percent cropland, 12 percent deciduous forest, 11 percent grassland, and 9

percent coniferous forest (Figure 2C).

Nesting sites of the reported free-living honeybees

The ratio of tree cavities and nesting sites in buildings, as well as the occupation rate of

different tree species, differed between urban, rural and forest habitats. For the reported

colonies in urban areas (N=304), tree cavities comprised 52% of all cavities, while building

cavities accounted for 40% (other cavity types: 8%; Figure 3A). In rural areas (N=121), tree

Nest site characteristics

Figure 3: A) Proportional distribution of nesting cavity types (tree, building or other) across the three different habitats. While the x-axis gives the overall percentage, white numbers indicate the count (N) of trees, buildings, and other cavities. B) This panel shows the distribution of tree species used by reported free-living colonies as nesting sites in this study. Pattern fills illustrate the distribution across different habitats. Darker green fill represents coniferous tree species. C) Displays the preferred nesting entrance directions for the two different types of cavities - those in trees and buildings. D) Cavity height distribution: A violin plot with an embedded boxplot that demonstrates the range and distribution of the heights of cavity entrances occupied by free-living colonies, categorized by cavity type. E) Illustrates the distribution of tree trunk diameters (measured at breast height) for trees that hosted free-living honeybee colonies.

cavities were again the most common (68%), followed by building cavities (21%). In forest

areas (N=105), tree cavities dominated with 81% of all cavities, while building cavities made

up 13 %. Looking at the whole dataset 63% (N=324) of the colonies were found in trees, 31%

(N=161) in building structures, and the remaining 6% (N=32) in other types such as rock

crevices (N=3) and open nesting (N=15) (Figure SM). Among tree species, Lime (*Tilia* spp.)

was the most frequently occupied (N=59; 18%), followed by beech (*Fagus sylvatica*, N=45;

14%), oak (*Quercus* spp., N=41; 13%) and ash (*Fraxinus excelsior*, N=35; 11%) (Figure 3B).

Our study uncovered clear patterns in common flight entrance directions of nest sites used by

free-living honeybees. For colonies situated in tree cavities where the direction was reported,

the observed frequency distribution across the eight cardinal and intercardinal directions was

significantly non-uniform (Rayleigh Test Statistic = 20.23, Bootstrap p-value = 0.0010; N =

285). Specifically, there was a significant preference for southern directions, with the highest

frequency observed in the South direction (N = 58 or 20 %; Figure 3C). Similarly, for

colonies located in buildings, the distribution was tested to be marginally non-uniform

(Rayleigh Test Statistic = 4.85, Bootstrap p-value = 0.079; N = 146). Unlike tree cavities,

cavities in human-built structures showed no clear directional preferences, although the West

(N = 32 or 22%) and South (N = 24 or 16%) directions were observed most frequently

(Figure 3C).

Our observations indicate that free-living honeybee colonies predominantly use nesting sites

far from the ground (mean and median entrance height: 5.7 m and 4.5 m; Figure 3D). There

was a statistically significant difference in the height of the cavities entrance in trees and in

buildings (Mann-Whitney U Test W = 13482, p < 0.001). The entrance heights of cavities

occupied in trees ranged from 0.1 to 30 meters, with a median of 4 meters. In contrast, the

heights of cavities entrances in man-made structures ranged from 0.2 to 40 meters, with a

median of 6 meters.

Additionally, we found that the diameter at breast height (DBH) of trees harbouring free-

living honeybee colonies was on average 0.64 meters (median: 0.69 meters, Figure 3E),

suggesting that colonies are dependent on trees with a substantial trunk diameter.

Occupation rates and colony density across seasons in Munich

We investigated cavity occupation rates in the city of Munich during three distinct seasonal

periods (spring, summer and autumn, see SM for further details). Multiplying these

occupation rates with the density of 0.58 known cavities per square kilometre in Munich (92

nest sites on an area of 160 km², see SM), we estimated the minimum density of free-living

honeybee colonies in Munich to be approximately 0.06 colonies per square kilometre in

spring, 0.42 in summer, and 0.28 in fall. It is important to note that the reported densities

should be viewed as minimum estimates, given the likelihood that a significant number of

colonies remain undetected and unreported. Based on these occupation rates, we infer that the

number of colonies during summer is roughly seven-fold higher compared to spring.

Conversely, the number of colonies in fall is approximately 33% lower than in summer.

<u>Survival rates of free-living honeybee colonies and the impact of monitoring type and year</u>

We analysed the life histories of 343 free-living honeybee colonies over the period from 2016

to 2023. Of these, 151 survival reports were provided by citizen scientists, while 192 were

personally observed in Munich. It is important to note that one colony can have multiple

survival reports for different years. The yearly survival rates for the two monitoring types are

depicted in Figure 4. In most years, survival rates reported by citizen scientists were higher

than those observed through personal monitoring. Notably, in the spring of 2019, all PM

colonies (N=32) perished.

345
Figure 4: Comparison of observed colony survival rates from Citizen Science monitoring (CS, light blue) and
personal monitoring in the Munich region (PM, light green) across the years.

Additionally, we consulted data from two published studies to determine whether the

differences between CS and PM were likely due to biases in CS or actual differences. Kohl et

al. (2022) focused on 97 overwintering colonies in German managed forest landscapes from

2017 to 2021, while Lang et al. (2022) investigated 24 colonies in Dortmund, Germany from

2018 to 2022 (Figure 5A). More detailed information about the datasets and the reanalysis

with the swarming timepoints observed in this study is provided in the SM. To assess the

impact of different monitoring types and year on survival rates, we constructed a glmmTMB

model. Model selection, guided by AIC values and convergence considerations, favoured

including 'year' as a fixed effect alongside the 'type of monitoring' (CS, PM, Kohl et al.

(2022) and Lang et al., (2022)). The ANOVA results from the glmmTMB model indicated

that both 'monitoring type' (Chi-squared = 14.85, df = 3, p = 0.002) and 'year' (Chi-squared =

17.47, df = 6, p = 0.008) significantly contributed to the model. The estimated probability of

survival was notably higher in 'Citizen Science' (28%) than in personal monitoring in the

Munich region (p = 0.007) and in 'Kohl et al. 2022' (p = 0.007), but not significantly

different from 'Lang et al. 2022' (p = 0.33; probably due to the small number of colonies)

(Figure 4B). While minor variations may arise from regional or setting differences, the

significant discrepancies observed are likely due to reporting biases in CS compared to

systematic surveys by experts. Model estimates placed the survival probability at 12% for

PM, 11% for 'Lang et al. 2022', 9% for 'Kohl et al. 2022', and 28% for CS (Figure 5B). These

findings suggest that both the type of monitoring and the year of observation have a

statistically significant impact on the reported survival rates of free-living honeybee colonies.

Comparative analysis of survival rates across Germany

370   *Figure 5: (A) Geographic distribution of life-history reports of free-living colonies from this study (CS and PM) and*
371   *two published datasets (Kohl et al., 2022; Lang et al., 2022). (B) Model estimates and 95% confidence intervals*
372   *for colony survival rates for the different monitoring types and studies.*

Biases in the different monitoring types

Our dataset comprised 423 nest sites with 1,064 Citizen Science (CS) observations and 107 nest sites with 1,491 personal monitoring (PM) observations. We noted a significantly lower number of CS reports per colony compared to PM (Wilcoxon rank-sum test: W = 3950, p < 0.001; Figure 6A). Furthermore, we found a stark contrast in the distribution of "alive" versus "dead" colony status reports between CS and PM (Pearson's Chi-squared test: χ2 = 176, df = 1, p < 0.001; Figure 6B), where 76% of CS reports indicated alive colonies compared to 42% in PM.

Monitoring biases

Figure 6: Analysis of reporting biases between Citizen Science monitoring (CS, light blue) and personal monitoring in the Munich region (PM, light green). A) Violin plot with included boxplot illustrating the distribution of the number of observations per colony for the whole monitoring timespan. B) Proportional representation of reported colony statuses ("alive" in light grey, "dead" in dark grey) for CS and PM. C) Temporal variation in colony statuses reported post-winter for the two different monitoring approaches in relation to the swarm season (in yellow) from the years 2019-2023. Note: The dashed line marks the first observed swarm for each year, after which spring reports on colony status may not be reliable anymore (late reports in red colour). Bold horizontal lines indicate median values of the timing of CS and PM reports.

We evaluated the timelines of reports from Citizen Science and personal monitoring, with a particular focus on reports submitted after winter. To assess the effectiveness of the reporting, we quantified the proportion of reports considered 'late reports', defined as submissions after the swarming start date for that year. The median reporting date after winter for CS was consistently later than for PM, specifically 21 April versus 29 March, respectively (mean: 3 April for PM vs. 2 May for CS) (Figure 6C). Consequently, a substantial proportion of CS reports were rendered unusable annually, i.e. around 46 % (84 out of 184) reports of CS could not be used each year, as they were reported too late (compared to 11% in PM, 15 out of 131 reports). These findings suggest an urgent need for improved reporting intensity and timelines within Citizen Science programs to ensure data reliability.

DISCUSSION

To investigate claims of free-living survivor colonies in Germany, we launched the BEEtree-Monitor project with a low-tech monitoring scheme designed to collectively gather data on free-living honeybee colonies. This initiative resulted in the most extensive database on the topic to our knowledge to date, enabling a comprehensive analysis of the nesting habits and life histories of these colonies over a seven-year period.

Deciduous forests are widely seen as the original habitat of honeybees and offer cavities if mature trees are present. While the reported free-living honeybee colonies relied heavily on tree cavities as nesting sites, our data show that many of the tree species favoured by honeybees are rare in today's forests and are more likely to be found in settlement areas, along streets, in parks, or in cemeteries. When examining the available foraging habitat around the reported colonies, our findings indicate that forests play a minor role. Additionally, today's German beech forests, for instance, do not provide diverse and rich foraging opportunities throughout the year (Rutschmann et al., 2023). This points to the conclusion that there is no pristine habitat for free-living honeybee colonies in the sense of a "natural environment". From a conservation perspective, urban landscapes have become crucial refuges for species considered "wild" and those needing to escape rural environments dominated by intensive agricultural land use. These areas are often rich in floral diversity and offer an array of potential nesting sites, such as cavities in buildings and mature trees in parks and gardens (Angold et al., 2006; Threlfall and Kendal, 2018; Gathof et al., 2022). Additionally, urban areas provide an attractive foraging environment for both managed and

free-living honeybee cohorts (Ayers and Rehan, 2021; Baldock, 2020; Rutschmann et al., 2023; Young et al., 2021). Forest aren't "natural" spaces of "wilderness" and the urbanization has unexpected conservational potential. Having entered the Anthropocene, the old dichotomies of natural/artificial of wild/domestic become potentially misleading categorical foundations for research and conservation agendas.

As a case in point, our calculations reveal a notably high density of free-living honeybee colonies in the city of Munich in summer (0.42 colonies/km²), surpassing those in managed German woodlands (0.23 colonies/km²), Polish rural areas (0.1 colonies/km²), and agricultural landscapes in Spain (0.2 colonies/km²) (Kohl et al., 2022; Kohl and Rutschmann, 2018; Oleksa et al., 2013; Rutschmann et al., 2022). This accumulation of free-living colonies in Munich and in cities in general could be attributed to several factors, including those mentioned above, but also to the higher chances of discovering a colony and the prevalence of managed honeybee colonies in urban areas compared to rural areas and forests.

Honeybees living within tree cavities in forested areas rarely share contact zones with humans, while urban environments present a more complex scenario. Here, interactions with humans are more frequent, particularly when colonies establish within building structures. It is also known that higher human population density correlates with higher densities of beekeepers and consequently more escaping swarms searching for cavities (Oré Barrios et al., 2017; von Büren et al., 2019). For example, in Munich, the known density of managed hives is over 12 per km², surpassing the German average density by a factor of 4.3 (personal information from the veterinary office and data from the German Ministry of Food and Agriculture, both in 2022). The dominance of reports from urban areas introduces a bias, potentially skewing perceptions regarding the distribution and density of free-living colonies across different habitats. This issue can be addressed by incorporating systematic approaches with random sampling techniques, such as beelining (Seeley, 2016; Kohl and Rutschmann, 2018; Radcliffe and Seeley, 2018).

High managed honeybee densities in many European regions have sparked ongoing debates about food competition and pathogen transfer among pollinators (Geldmann and González-Varo, 2018; Ropars et al., 2019; Saunders et al., 2018; Herrera, 2020; Iwasaki and Hogendoorn, 2021; Ghazoul, 2005; Casanelles-Abella and Moretti, 2022; Weissmann et al., 2023; Egerer and Kowarik, 2020). Such concerns could be particularly pronounced in urban settings where some endangered solitary bees are present and the density of managed

honeybee colonies is already high (Mallinger et al., 2017; but see Harder and Miksha, 2022;
Steffan-Dewenter and Tscharntke, 2000). Yet, our findings suggest that the density of free-
living colonies in Munich remains relatively low, at approximately 4% of the density of
registered colonies in the city (see SM for more information). While the local densities of
managed colonies require careful consideration, free-living colonies should not be considered
problematic for urban pollinators nor for managed colonies from urban beekeepers (see Kohl
et al., (2023a) for an investigation on the parasite loads of free-living colonies).

Our findings reinforce the understanding that honeybees generally prefer elevated cavities for
nesting (Seeley and Morse, 1978; Seeley, 2019). The observed significant difference in the
height of nests between building and tree cavities may partially reflect a bias, where colonies
in buildings are more likely to be observed despite higher elevation from ground level.
Moreover, buildings in urban areas inherently offer more high-elevation spaces suitable for
bee colonies, unlike trees, where the likelihood of finding a cavity large enough to
accommodate a colony decreases with height.

Interestingly, our data indicate that free-living honeybees exhibit a pronounced directional
preference for southern or southwestern exposures when selecting cavities. This preference
aligns with the thermoregulatory benefits of southern orientations, which facilitate sun
exposure and warmth, particularly beneficial during spring when colonies are emerging from
winter. This finding is consistent with some previous studies that noted similar directional
preferences (Avitabile et al., 1978; Radcliffe and Seeley, 2018; Cordillot, 2024), though it
contrasts with others that reported a random distribution of entrance directions (Seeley and
Morse, 1976; Gambino et al., 1990). Additionally, in cities and forests, a high proportion of
bee-used hollows in trees and facade insulations are created by woodpeckers. While we
cannot exclude an underlying preference of woodpeckers for certain directions and heights,
the consistent directional preference observed in our study suggests a genuine selection by
the bees themselves.

In German forest environments, spruce (*Picea abies*, 25%), pine (*Pinus sylvestris*, 22%),
beech (*Fagus sylvatica*, 15%), and oak (*Quercus* spp., 10%) are the most common tree
species by area, collectively representing more than 70% of the trees (Schmitz, 2014). When
examining the tree species housing free-living honeybee colonies in German forests, these
four species were also the most common, but the distribution was significantly different.
While the deciduous species beech (35%) and oak (20%) were highly overused by bees,

colonies were greatly underrepresented in the two most important coniferous tree species in managed forests: pine (13%) and spruce (11%). Although coniferous species are still predominant in managed forests, they do not provide as many suitable cavities for bees and other animals (Requier et al., 2020). Lime trees (*Tilia* spp.), which are of minor importance in managed forests, dominate both the urban and rural landscapes as alley trees and are heavily utilized by honeybees as nesting sites and sources of food. Especially in urban areas, the diversity of tree species used by bees was much higher, with lime (23%), ash (*Fraxinus excelsior*, 16%), plane (*Platanus acerifolia*, 11%), and horse chestnut (*Aesculus hippocastanum*, 7%) being the most frequently occupied. Our observations that ash trees are disproportionately favoured as nesting sites, raises concerns in light of the ongoing threat of *Hymenoscyphus fraxineus*, the causal agent of ash dieback across Europe. In Munich, nearly half of the inhabited trees are ash, and with over half of the city's free-living colonies residing in buildings, the ash dieback could significantly increase human contact with these colonies.

Our study underscores the importance of mature, old-growth trees in providing suitable nesting sites for free-living honeybees (Bütler et al., 2013; Requier et al., 2020; Visick and Ratnieks, 2023b). With an average diameter at breast height (DBH) of 64 centimetres, these large trees in our study are vital for supporting diverse ecological communities. In general, conservation efforts should, prioritize the preservation of old-growth forests and the protection of long-lived tree species -across all habitats- to maintain crucial nesting habitats for many species including free-living honeybees.

To assess the self-sustainability of a population is important for the conservations of species with a clear status of being "wild" or "feral". For some species as for the western honeybees in Europe the situation is more complex. Local European honeybee populations consist of deeply entangled managed and free-living cohorts in varying proportions. For centuries, the breeding of the former and the natural selection of the latter have impacted the overall population. With the advent of *Varroa destructor* and the continuing use of acaricides in most beekeeping practices, the ratio might have shifted in some European regions, including Germany. Yet, the species remains in a liminal state, and depending on future conservation strategies, populations might either shift towards full dependency on human care or retain and even enhance their capability to self-sustain by naturally adapting to new environmental circumstances.

In regions such as Spain and the UK, survival rates of free-living honeybee cohorts suggest potential self-sustainability or near self-sustaining populations (Rutschmann et al., 2022; Visick and Ratnieks, 2023b). Notably, in Gwynedd, Wales, the use of acaricides in beekeeping is no longer necessary (Remter, 2015). Our personal monitoring analysis in Munich, however, indicates that the survival threshold required for self-sustaining cohorts— approximately one-third annual survival (Kohl et al., 2022)—is not being met in Germany. This finding aligns with other studies conducted in variable habitats in Germany and neighbouring countries (Kohl et al., 2022; Lang et al., 2022; Cordillot, 2024) , yet contrasts sharply with anecdotal claims from lay observers who report continuous and prolonged occupancy of cavities. These observations do not necessarily confirm extended lifespans of individual colonies; rather, it seems more plausible that many free-living colonies in Germany are recent escapees from managed apiaries, and that monitoring has not been conducted rigorously enough.

The factors contributing to the decline of free-living colonies are varied, encompassing both ecological and evolutionary aspects. From an ecological perspective, challenges such as a shortage of floral resources, lack of suitable and safe nesting sites, and parasite pressure can significantly impact colony viability (Kohl et al., 2023b; Rutschmann et al., 2023). Another aspect to consider is the evolutionary impact of modern beekeeping practices: They focus on breeding for desirable traits such as maximized honey production, reduced swarming tendencies, and docility (Seeley, 2019). Breeding efforts for varroa resistance exist, but they have not yet made a significant impact (Guichard et al., 2023), and the continuous medical treatment of managed colonies prevents tolerance traits from prevailing at the population level (Blacquière et al., 2019; Neumann and Blacquière, 2017). Additionally in Germany, the replacement of the native subspecies *Apis mellifera mellifera* with non-native subspecies might have profound implications on the genetic diversity and adaptability of honeybee populations in this area, potentially influencing their ability to thrive independently (Büchler et al., 2014). Understanding the complex interplay of these factors is crucial for developing reliable conservation strategies for free-living cohorts.

The issue of defining species as "wild" and self-sustaining is significant, especially as organizations like the IUCN assess species status based on these criteria. Emphasizing wilderness and self-sustainability can hinder conservation efforts for liminal species in danger of losing their ability to self-sustain. We suggest understanding honeybees in Europe

as a liminal species, with larger cohorts living under beekeeping conditions and smaller ones

living independently, both within a broad and shared geographic range. These cohorts are

unlikely to be found independently of each other and are deeply interconnected through

genetic exchange via mating and annual swarming. The survival rates of the free-living

cohort influence the overall population's genetic diversity in Europe, including the managed

cohort.

While the higher number of reports for colonies monitored by PM is partly due to the density

of observations applied in the PM to better understand the flight patterns of robbing, scouting

or survivor bees, there remains a concern with CS as many colonies in the CS database were

reported only once and could not be used for the survival analysis. In fact, it is plausible that

citizen scientists primarily report on active, living colonies, and may discontinue monitoring

once a colony is no longer present, thereby creating a biased dataset. Our model predictions,

incorporating monitoring type and data from CS, PM and two published datasets, also point

in this direction, reinforcing the potential bias introduced by CS reporting practices. In our

study, approximately 76% of reports from citizen scientists indicated the presence of living

colonies, contrasting with only about 42% in the personally monitored dataset meeting the

52% and 43% in two studies by Rutschmann et al. (2022) and Kohl et al. (2022), relying on

expert monitoring. This discrepancy between reports on occupied and unoccupied cavities

suggests that the ratio of reported colony status indicate inherent biases within the monitoring

methods of layperson in contrast to experts. Understanding and addressing these biases is

essential for accurately interpreting survival rates of free-living honeybees based on Citizen

Science data. Our analysis also revealed a significant delay in reporting by citizen scientists

during the spring. This delay rendered a significant proportion of the data unusable for

survival analysis. Despite accounting for this factor, the survival rates reported in CS data

remained much higher, suggesting a likely bias. These findings indicate that results from CS

should be interpreted with caution.

A key challenge of our study, similar to many Citizen Science projects, is its serendipitous

nature. While Citizen Science is extremely valuable for identifying the locations of free-

living honeybee colonies across huge geographic scales, it presents complexities and

significant efforts when it comes to monitoring these colonies coherently over time. Based on

our experience in the first years of the study we optimized the monitoring scheme for

ourselves and for the applicability in CS. To maximize the contribution of CS to bee research,

we recommend adopting a standardized monitoring scheme (Figure 1B). This approach can

be built on the successful protocol employed in our personal monitoring, which consisted of

at least three site visits per year to detect pollen import (or typologized flight-patterns) and

empirical data on the swarming season (especially its onset). The first visit should occur

before the swarming season begins, the second visit after the swarming season to confirm

newly established colonies and to include known but previously unoccupied sites to find new

founder colonies, and the third visit in autumn to assess summer-deaths and the number of

colonies going into winter. Citizen scientists need to be reminded early and often enough and

briefed on appropriate timing and weather conditions for observations. Given the average

height of the entrances, they should be equipped with sufficiently powerful binoculars or

spotting scopes to reliably observe pollen import at each of the three visits. Additionally,

observers could be trained to differentiate between various flight patterns, such as normal

traffic, robbing, or scouting along the criteria offered in this study. Including a commentary

section in the monitoring forms proved valuable, as it allowed observers to describe their

observations in detail. This additional information helped us assess the reliability of the data,

especially in cases where the data seemed suspicious or lacked information on pollen import.

While we could then decide whether to include or exclude such reports in the final dataset,

having consistent reports on pollen import is much more time-efficient and should be

prioritized.

In essence, structured and standardized monitoring projects are indispensable for thoroughly

understanding the mechanisms underlying survival of free-living colonies. Long-term

monitoring and more extensive geographic coverage would enhance our understanding of the

survival and reproductive success of free-living honeybee colonies in Europe. Collaborative

efforts combining scientific research with Citizen Science initiatives could prove beneficial in

achieving these objectives. In this sense, overwintering survival serves as a practical and

easily recordable indicator for identifying regions in Europe with high survival rates and

enlarged free-living cohorts. Such an approach requires minimal investment in monitoring

and does not rely on sophisticated kinship analysis, making it accessible for widespread

Citizen Science participation.

CONCLUSION

Given the growing trend of citizen science-based knowledge production, particularly for

monitoring free-living honeybee colonies, we critically compared the citizen science dataset

with data collected through our personal monitoring efforts. We identified epistemic pitfalls

in the collection and interpretation of citizen science data and outlined potential strategies to

optimize its use for monitoring purposes. Despite a reporting bias towards more densely

populated areas, the combined datasets provide a reliable and comprehensive foundation for

assessing nest site parameters of unmanaged colonies across various land cover types in

Germany. Importantly, the low-tech approach of our personal monitoring offered a valid

foundation for survival analysis. Our findings contribute to filling a gap in our knowledge

about the western honeybee and enhance our understanding of its free-living cohort in

Germany. Given the low survival rates of free-living colonies in Germany compared to other

regions in Europe and abroad, more detailed and comparative research is needed to develop

conservation strategies for free-living cohorts.

ACKNOWLEDGMENTS

625 We express our gratitude to the BEEtree-Monitor community and all the citizen scientists

626 who provided data for this study. A list of data contributors and those who helped shape the

627 monitoring protocol can be found in the Acknowledgment Appendix in the SM.

CONFLICTS OF INTEREST

629 The authors declare no conflict of interest.

AUTHORS' CONTRIBUTIONS

Conceptualization: SR-BR-FR
Methodology: SR-FR-BR
Software: SR
Formal analysis: BR
Investigation: SR-FR
Resources: FR-SR-BR
Data curation: SR-BR-FR
Writing - original draft: BR-FR
Writing - Review & editing: SR
Visualization: BR
Administration: SR-FR

FUNDING

643 No funding was received for conducting this study.

REFERENCES

Anderson, D.L., Trueman, J.W.H., 2000. Varroa jacobsoni (Acari: Varroidae) is more than one species. Experimental & Applied Acarology 24, 165–189. https://doi.org/10/c2263v

Angold, P.G., Sadler, J.P., Hill, M.O., Pullin, A., Rushton, S., Austin, K., Small, E., Wood, B., Wadsworth, R., Sanderson, R., Thompson, K., 2006. Biodiversity in urban habitat patches. Science of The Total Environment 360, 196–204. https://doi.org/10.1016/j.scitotenv.2005.08.035

Avitabile, A., Stafstrom, D.P., Donovan, K.J., 1978. Natural Nest Sites of Honeybee Colonies in Trees in Connecticut, USA. Journal of Apicultural Research 17, 222–226. https://doi.org/10.1080/00218839.1978.11099930

Ayers, A.C., Rehan, S.M., 2021. Supporting Bees in Cities: How Bees Are Influenced by Local and Landscape Features. Insects 12, 128. https://doi.org/10.3390/insects12020128

Baldock, K.C., 2020. Opportunities and threats for pollinator conservation in global towns and cities. Current Opinion in Insect Science 38, 63–71. https://doi.org/10.1016/j.cois.2020.01.006

Bila Dubaić, J., Simonović, S., Plećaš, M., Stanisavljević, L., Davidović, S., Tanasković, M., Ćetković, A., 2021. Unprecedented Density and Persistence of Feral Honey Bees in Urban Environments of a Large SE-European City (Belgrade, Serbia). Insects 12, 1127. https://doi.org/10.3390/insects12121127

Blacquière, T., Boot, W., Calis, J., Moro, A., Neumann, P., Panziera, D., 2019. Darwinian black box selection for resistance to settled invasive Varroa destructor parasites in honey bees. Biological Invasions 21, 2519–2528. https://doi.org/10.1007/s10530-019-02001-0

Breeze, T.D., Vaissière, B.E., Bommarco, R., Petanidou, T., Seraphides, N., Kozák, L., Scheper, J., Biesmeijer, J.C., Kleijn, D., Gyldenkærne, S., Moretti, M., Holzschuh, A., Steffan-Dewenter, I., Stout, J.C., Pärtel, M., Zobel, M., Potts, S.G., 2014. Agricultural Policies Exacerbate Honeybee Pollination Service Supply-Demand Mismatches Across Europe. PLoS ONE 9, e82996. https://doi.org/10.1371/journal.pone.0082996

Browne, K.A., Hassett, J., Geary, M., Moore, E., Henriques, D., Soland-Reckeweg, G., Ferrari, R., Mac Loughlin, E., O'Brien, E., O'Driscoll, S., Young, P., Pinto, M.A., McCormack, G.P., 2020. Investigation of free-living honey bee colonies in Ireland. Journal of Apicultural Research 1–12. https://doi.org/10.1080/00218839.2020.1837530

Büchler, R., Costa, C., Hatjina, F., Andonov, S., Meixner, M.D., Conte, Y.L., Uzunov, A., Berg, S., Bienkowska, M., Bouga, M., Drazic, M., Dyrba, W., Kryger, P., Panasiuk, B., Pechhacker, H., Petrov, P., Kezić, N., Korpela, S., Wilde, J., 2014. The influence of genetic origin and its interaction with environmental effects on the survival of *Apis mellifera* L. colonies in Europe. Journal of Apicultural Research 53, 205–214. https://doi.org/10.3896/IBRA.1.53.2.03

Bütler, R., Lachat, T., Larrieu, L., Paillet, Y., 2013. 2.1 Habitat trees: key elements for forest biodiversity, in: Kraus, D., Krumm, F. (Eds.), Integrative Approaches as an

Opportunity for the Conservation of Forest Biodiversity. Germany: European Forest
Institute, pp. 84–91.
Casanelles-Abella, J., Moretti, M., 2022. Challenging the sustainability of urban beekeeping
using evidence from Swiss cities. npj Urban Sustain 2, 3.
https://doi.org/10.1038/s42949-021-00046-6
Cordillot, F., 2024. Erste Suche nach wilden Honigbienen (Apis mellifera L., 1758) auf der
Schweizer Alpennordseite. Entomo Helvetica 17, 97–114.
Egerer, M., Kowarik, I., 2020. Confronting the Modern Gordian Knot of Urban Beekeeping.
Trends in Ecology & Evolution 35, 956–959.
https://doi.org/10.1016/j.tree.2020.07.012
European Environment Agency, 2018. CORINE Land Cover 2018.
Fraisl, D., Hager, G., Bedessem, B., Gold, M., Hsing, P.-Y., Danielsen, F., Hitchcock, C.B.,
Hulbert, J.M., Piera, J., Spiers, H., Thiel, M., Haklay, M., 2022. Citizen science in
environmental and ecological sciences. Nat Rev Methods Primers 2, 64.
https://doi.org/10.1038/s43586-022-00144-4
Gambino, P., Hoelmer, K., Daly, H.V., 1990. Nest sites of feral honey bees in California,
USA. Apidologie 21, 35–45. https://doi.org/10.1051/apido:19900105
Garibaldi, L.A., Steffan-Dewenter, I., Winfree, R., Aizen, M.A., Bommarco, R.,
Cunningham, S.A., Kremen, C., Carvalheiro, L.G., Harder, L.D., Afik, O.,
Bartomeus, I., Benjamin, F., Boreux, V., Cariveau, D., Chacoff, N.P., Dudenhoffer,
J.H., Freitas, B.M., Ghazoul, J., Greenleaf, S., Hipolito, J., Holzschuh, A., Howlett,
B., Isaacs, R., Javorek, S.K., Kennedy, C.M., Krewenka, K.M., Krishnan, S.,
Mandelik, Y., Mayfield, M.M., Motzke, I., Munyuli, T., Nault, B.A., Otieno, M.,
Petersen, J., Pisanty, G., Potts, S.G., Rader, R., Ricketts, T.H., Rundlof, M., Seymour,
C.L., Schuepp, C., Szentgyorgyi, H., Taki, H., Tscharntke, T., Vergara, C.H., Viana,
B.F., Wanger, T.C., Westphal, C., Williams, N., Klein, A.M., 2013. Wild Pollinators
Enhance Fruit Set of Crops Regardless of Honey Bee Abundance. Science 339, 1608–
1611. https://doi.org/10.1126/science.1230200
Gathof, A.K., Grossmann, A.J., Herrmann, J., Buchholz, S., 2022. Who can pass the urban
filter? A multi-taxon approach to disentangle pollinator trait–environmental
relationships. Oecologia 199, 165–179. https://doi.org/10.1007/s00442-022-05174-z
Geldmann, J., González-Varo, J.P., 2018. Conserving honey bees does not help wildlife.
Science 359, 392–393. https://doi.org/10.1126/science.aar2269
Ghazoul, J., 2005. Buzziness as usual? Questioning the global pollination crisis. Trends in
Ecology & Evolution 20, 367–373. https://doi.org/10.1016/j.tree.2005.04.026
Grindrod, I., Martin, S.J., 2021. Parallel evolution of *Varroa* resistance in honey bees: a
common mechanism across continents? Proc. R. Soc. B. 288, 20211375.
https://doi.org/10.1098/rspb.2021.1375
Guichard, M., Dainat, B., Dietemann, V., 2023. Prospects, challenges and perspectives in
harnessing natural selection to solve the 'varroa problem' of honey bees. Evolutionary
Applications 16, 593–608. https://doi.org/10.1111/eva.13533
Harder, L.D., Miksha, R.M., 2022. No statistical evidence that honey bees competitively
reduced wild bee abundance in the Munich Botanic Garden—a comment on Renner et
al. (2021). Oecologia 198, 337–341. https://doi.org/10.1007/s00442-022-05112-z

Hartig, F., Hartig, M.F., 2017. Package 'DHARMa.'

Hassett, J., Browne, K.A., McCormack, G.P., Moore, E., Society, N.I.H.B., Soland, G.,
Geary, M., 2018. A significant pure population of the dark European honey bee (*Apis*
mellifera mellifera) remains in Ireland. Journal of Apicultural Research 57, 337–350.
https://doi.org/10.1080/00218839.2018.1433949

Henneken, R., Helm, S., Menzel, A., 2012. Meteorological Influences on Swarm Emergence
in Honey Bees (Hymenoptera: Apidae) as Detected by Crowdsourcing. env. entom.
41, 1462–1465. https://doi.org/10.1603/EN12139

Herrera, C.M., 2020. Gradual replacement of wild bees by honeybees in flowers of the
Mediterranean Basin over the last 50 years. Proceedings of the Royal Society B:
Biological Sciences 287, 20192657. https://doi.org/10.1098/rspb.2019.2657

Hsing, P.-Y., Hill, R.A., Smith, G.C., Bradley, S., Green, S.E., Kent, V.T., Mason, S.S.,
Rees, J., Whittingham, M.J., Cokill, J., others, 2022. Large-scale mammal monitoring:
The potential of a citizen science camera-trapping project in the United Kingdom.
Ecological Solutions and Evidence 3, e12180. https://doi.org/10.1002/2688-
8319.12180

Hung, K.-L.J., Kingston, J.M., Albrecht, M., Holway, D.A., Kohn, J.R., 2018. The worldwide
importance of honey bees as pollinators in natural habitats. Proceedings of the Royal
Society of London B: Biological Sciences 285.
https://doi.org/10.1098/rspb.2017.2140

Ibisch, P.L., Hoffmann, M.T., Kreft, S., Pe'er, G., Kati, V., Biber-Freudenberger, L.,
DellaSala, D.A., Vale, M.M., Hobson, P.R., Selva, N., 2016. A global map of roadless
areas and their conservation status. Science 354, 1423–1427.
https://doi.org/10.1126/science.aaf7166

Ilyasov, R.A., Kosarev, M.N., Neal, A., Yumaguzhin, F.G., 2015b. Burzyan Wild-Hive
Honeybee *A.M. mellifera* in South Ural. Bee World 92, 7–11.
https://doi.org/10.1080/0005772X.2015.1047634

Iwasaki, J.M., Hogendoorn, K., 2021. How protection of honey bees can help and hinder bee
conservation. Current Opinion in Insect Science 46, 112–118.
https://doi.org/10.1016/j.cois.2021.05.005

Jones, R., 2004. European beekeeping in the 21st century: strengths, weaknesses,
opportunities, threats. Bee World 85, 77–80.
https://doi.org/10.1080/0005772X.2004.11099637

Kevan, P.G., Baker, H.G., 1983. Insects as Flower Visitors and Pollinators. Annu. Rev.
Entomol. 28, 407–453. https://doi.org/10.1146/annurev.en.28.010183.002203

Koffler, S., Barbiéri, C., Ghilardi-Lopes, N.P., Leocadio, J.N., Albertini, B., Francoy, T.M.,
Saraiva, A.M., 2021. A Buzz for Sustainability and Conservation: The Growing
Potential of Citizen Science Studies on Bees. Sustainability 13, 959.
https://doi.org/10.3390/su13020959

Kohl, P.L., D'Alvise, P., Rutschmann, B., Roth, S., Remter, F., Steffan-Dewenter, I.,
Hasselmann, M., 2023a. Reduced parasite burden in feral honeybee colonies. Ecol Sol
and Evidence 4, e12264. https://doi.org/10.1002/2688-8319.12264

Kohl, P.L., Rutschmann, B., 2018. The neglected bee trees: European beech forests as a home
for feral honey bee colonies. PeerJ 6, e4602. https://doi.org/10.7717/peerj.4602

Kohl, P.L., Rutschmann, B., Sikora, L.G., Wimmer, N., Zahner, V., D'Alvise, P.,
Hasselmann, M., Steffan-Dewenter, I., 2023b. Parasites, depredators, and limited
resources as potential drivers of winter mortality of feral honeybee colonies in
German forests. Oecologia 202, 465–480. https://doi.org/10.1007/s00442-023-05399-
6

Kohl, P.L., Rutschmann, B., Steffan-Dewenter, I., 2022. Population demography of feral
honeybee colonies in central European forests. R. Soc. open sci. 9, 220565.
https://doi.org/10.1098/rsos.220565

Lang, U., Albouy, V., Zewen, C., 2022. Comparative monitoring of free-living honey bee
colonies in three Western European regions. Natural Bee Husbandry Magazine.

Lenth, R., Lenth, M.R., 2018. Package 'lsmeans.' The American Statistician 34, 216–221.

Lesiv, M., Laso Bayas, J.C., See, L., Duerauer, M., Dahlia, D., Durando, N., Hazarika, R.,
Kumar Sahariah, P., Vakolyuk, M., Blyshchyk, V., Bilous, A., Perez-Hoyos, A.,
Gengler, S., Prestele, R., Bilous, S., Akhtar, I.U.H., Singha, K., Choudhury, S.B.,
Chetri, T., Malek, Ž., Bungnamei, K., Saikia, A., Sahariah, D., Narzary, W., Danylo,
O., Sturn, T., Karner, M., McCallum, I., Schepaschenko, D., Moltchanova, E., Fraisl,
D., Moorthy, I., Fritz, S., 2019. Estimating the global distribution of field size using
crowdsourcing. Global Change Biology 25, 174–186.
https://doi.org/10.1111/gcb.14492

Magnusson, A., Skaug, H., Nielsen, A., Berg, C., Kristensen, K., Maechler, M., van
Bentham, K., Bolker, B., Brooks, M., Brooks, M.M., 2017. Package 'glmmTMB.' R
Package Version 0.2. 0.

Mallinger, R.E., Gaines-Day, H.R., Gratton, C., 2017. Do managed bees have negative
effects on wild bees?: A systematic review of the literature. PLOS ONE 12,
e0189268. https://doi.org/10.1371/journal.pone.0189268

Meixner, M.D., Kryger, P., Costa, C., 2015. Effects of genotype, environment, and their
interactions on honey bee health in Europe. Social Insects * Vectors and Medical and
Veterinary Entomology 10, 177–184. https://doi.org/10/gcm437

Moritz, R.F.A., Kraus, F.B., Kryger, P., Crewe, R.M., 2007. The size of wild honeybee
populations (Apis mellifera) and its implications for the conservation of honeybees. J
Insect Conserv 11, 391–397. https://doi.org/10.1007/s10841-006-9054-5

Moro, A., Beaurepaire, A., Dall'Olio, R., Rogenstein, S., Blacquière, T., Dahle, B., de
Miranda, J.R., Dietemann, V., Locke, B., Licón Luna, R.M., Le Conte, Y., Neumann,
P., 2021. Using Citizen Science to Scout Honey Bee Colonies That Naturally Survive
Varroa destructor Infestations. Insects 12, 536.
https://doi.org/10.3390/insects12060536

Neumann, P., Blacquière, T., 2017. The Darwin cure for apiculture? Natural selection and
managed honeybee health. Evolutionary Applications 10, 226–230.
https://doi.org/10/f9zfsw

Oberreiter, H., Dünser, A., Schlagbauer, J., Brodschneider, R., 2021. Das Wildnisgebiet
Dürrenstein – ein Lebensraum für wildlebende Bienenvölker? 19.

Oldroyd, B.P., Thexton, E.G., Lawler, S.H., Crozier, R.H., 1997. Population demography of
Australian feral bees (Apis mellifera). Oecologia 111, 381–387.
https://doi.org/10.1007/s004420050249

Oleksa, A., Gawroński, R., Tofilski, A., 2013. Rural avenues as a refuge for feral honey bee population. Journal of Insect Conservation 17, 465–472. https://doi.org/10/f4xf44

Oré Barrios, C., Mäurer, E., Lippert, C., Dabbert, S., 2017. Eine ökonomische Analyse des Imkerei-Sektors in Deutschland.

Phiri, B.J., Fèvre, D., Hidano, A., 2022. Uptrend in global managed honey bee colonies and production based on a six-decade viewpoint, 1961–2017. Sci Rep 12, 21298. https://doi.org/10.1038/s41598-022-25290-3

Pocock, M.J.O., Chandler, M., Bonney, R., Thornhill, I., Albin, A., August, T., Bachman, S., Brown, P.M.J., Cunha, D.G.F., Grez, A., Jackson, C., Peters, M., Rabarijaon, N.R., Roy, H.E., Zaviezo, T., Danielsen, F., 2018. A Vision for Global Biodiversity Monitoring With Citizen Science, in: Advances in Ecological Research. Elsevier, pp. 169–223. https://doi.org/10.1016/bs.aecr.2018.06.003

QGIS Development Team, 2020. QGIS Geographic Information System. Open Source Geospatial Foundation.

R Core Team, 2016. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

Radcliffe, R., Seeley, T., 2018. Deep forest bee hunting: a novel method for finding wild colonies of honey bees in old-growth forests. American Bee Journal 158, 871–877.

Ratnieks, F.L., Piery, M.A., Cuadriello, I., 1991. The natural nest and nest density of the Africanized honey bee (Hymenoptera, Apidae) near Tapachula, Chiapas, Mexico. The Canadian Entomologist 123, 353–359. https://doi.org/10.4039/Ent123353-2

Remter, F., 2021a. Eine Frage der Haltung/ Beewildered Companions.

Remter, F., 2021b. Ecologizing Honeybee Care. Multi-species Bodies and Trust in the Varroa Pandemic. Theorizing the Contemporary, Fieldsights.

Remter, F., 2015. Treatment free Beekeeping in Gwynedd Wales.

Requier, F., Garnery, L., Kohl, P.L., Njovu, H.K., Pirk, C.W.W., Crewe, R.M., Steffan-Dewenter, I., 2019. The Conservation of Native Honey Bees Is Crucial. Trends in Ecology & Evolution 34, 789–798. https://doi.org/10.1016/j.tree.2019.04.008

Requier, F., Paillet, Y., Laroche, F., Rutschmann, B., Zhang, J., Lombardi, F., Svoboda, M., Steffan-Dewenter, I., 2020. Contribution of European forests to safeguard wild honeybee populations. CONSERVATION LETTERS 13. https://doi.org/10.1111/conl.12693

Ropars, L., Dajoz, I., Fontaine, C., Muratet, A., Geslin, B., 2019. Wild pollinator activity negatively related to honey bee colony densities in urban context. PLoS ONE 14, e0222316. https://doi.org/10.1371/journal.pone.0222316

Rosenkranz, P., Aumeier, P., Ziegelmann, B., 2010. Biology and control of Varroa destructor. Journal of Invertebrate Pathology 103, S96–S119. https://doi.org/10/dm3hk6

Rutschmann, B., Kohl, P.L., Machado, A., Steffan-Dewenter, I., 2022. Semi-natural habitats promote winter survival of wild-living honeybees in an agricultural landscape. Biological Conservation 266, 109450. https://doi.org/10.1016/j.biocon.2022.109450

Rutschmann, B., Kohl, P.L., Steffan-Dewenter, I., 2023. Foraging distances, habitat preferences and seasonal colony performance of honeybees in Central European forest

landscapes. Journal of Applied Ecology 60, 1056–1066. https://doi.org/10.1111/1365-2664.14389

Saunders, M.E., Smith, T.J., Rader, R., 2018. Bee conservation: Key role of managed bees. Science 360, 389–389. https://doi.org/10.1126/science.aat1535

Schmitz, F., 2014. Der Wald in Deutschland–ausgewählte Ergebnisse der dritten Bundeswaldinventur.

Schneider, S., Blyther, R., 1988. The habitat and nesting biology of the African honey beeApis mellifera scutellata in the Okavango River Delta, Botswana, Africa 15. https://doi.org/10.1007/BF02223930

Seeley, T.D., 2019. The Lives of Bees: The Untold Story of the Honey Bee in the Wild. Princeton University Press.

Seeley, T.D., 2016. Following the Wild Bees: The Craft and Science of Bee Hunting. Princeton University Press.

Seeley, T.D., 2007. Honey bees of the Arnot Forest: a population of feral colonies persisting with *Varroa destructor* in the northeastern United States. Apidologie 38, 19–29. https://doi.org/10.1051/apido:2006055

Seeley, T.D., Morse, R.A., 1978. Nest site selection by the honey bee,Apis mellifera. Ins. Soc 25, 323–337. https://doi.org/10.1007/BF02224297

Seeley, T.D., Morse, R.A., 1976. The nest of the honey bee (Apis mellifera L.). Ins. Soc 23, 495–512. https://doi.org/10.1007/BF02223477

Sponsler, D., Johnson, R., 2015. Honey bee success predicted by landscape composition in Ohio, USA. PeerJ 3, e838. https://doi.org/10.7717/peerj.838

Steffan-Dewenter, I., Kuhn, A., 2003. Honeybee foraging in differentially structured landscapes. Proceedings of the Royal Society of London B: Biological Sciences 270, 569–575. https://doi.org/10.1098/rspb.2002.2292

Steffan-Dewenter, I., Tscharntke, T., 2000. Resource overlap and possible competition between honey bees and wild bees in central Europe. Oecologia 122, 288–296. https://doi.org/10/bj7dwz

Stoeckhert, F.K., 1954. Fauna Apoideorum Germaniae.(Nachträge und Ergänzungen zu FK Stoeckhert, Die Bienen Frankens, Deutsche entomologische Zeitschrift, 1932, Beiheft, Berlin, 1933). Verlag der Bayerischen Akademie der Wissenschaften; in Kommission bei Beck.

Threlfall, C.G., Kendal, D., 2018. The distinct ecological and social roles that wild spaces play in urban ecosystems. Urban Forestry & Urban Greening 29, 348–356. https://doi.org/10.1016/j.ufug.2017.05.012

Tsagris, M., Athineou, G., Sajib, A., Amson, E., Waldstein, M.J., Tsagris, M.M., 2016. Package 'Directional.'

Visick, O.D., Ratnieks, F.L.W., 2023a. Density of wild honey bee, *Apis mellifera*, colonies worldwide. Ecology and Evolution 13, e10609. https://doi.org/10.1002/ece3.10609

Visick, O.D., Ratnieks, F.L.W., 2023b. Ancient, veteran and other listed trees as nest sites for wild-living honey bee, Apis mellifera, colonies. J Insect Conserv. https://doi.org/10.1007/s10841-023-00530-7

von Büren, R.S., Oehen, B., Kuhn, N.J., Erler, S., 2019. High-resolution maps of Swiss
apiaries and their applicability to study spatial distribution of bacterial honey bee
brood diseases. PeerJ 7, e6393. https://doi.org/10.7717/peerj.6393
Weissmann, J.A., Rader, S., Ritz, R., Walldorf, I.R.M., Probst, J., Szydlik, K.R., Schaefer,
H., 2023. Holistic wild bee management in urban spaces. Front. Sustain. Cities 5,
1155714. https://doi.org/10.3389/frsc.2023.1155714
Wickham, H., 2017. The tidyverse. R package ver 1, 836.
Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
York.
Young, A.M., Kohl, P.L., Rutschmann, B., Steffan-Dewenter, I., Brockmann, A., Dyer, F.C.,
2021. Temporal and spatial foraging patterns of three Asian honey bee species in
Bangalore, India. Apidologie 52, 503–523. https://doi.org/10.1007/s13592-020-
00839-1
917