ImPaqT - A Golden Gate-based Toolkit for Zebrafish Transgenesis

Saskia Hurst[1,2], Christiane Dimmler[2], Mark R. Cronan[2, *]

[1] Humboldt-Universität zu Berlin, Faculty of Life Sciences, Unter den Linden 6, 10117 Berlin,

Germany

[2] Max Planck Institute for Infection Biology, Charitéplatz 1, 10117 Berlin, Germany

[*]Correspondence: Mark Cronan (cronan@mpiib-berlin.mpg.de)

ORCID identifier: 0000-0002-3760-1649 (MRC)

Abstract

Transgenic animals continue to play an essential role in many aspects of zebrafish research, including the development of disease models. The most widely used system for zebrafish transgenesis is the Tol2 transposon system. Here, we have developed ImPaqT (**Im**munological toolkit for **Paq**CI-based Golden Gate Assembly of Tol2 **T**ransgenes), a new Tol2-based transgenesis system that utilizes Golden Gate assembly to facilitate the production of transgenic zebrafish lines. This system allows for rapid assembly of multiple fragments into a single transgene, facile swapping of individual sequences to generate new transgenes and an easy cloning workflow to incorporate new genetic elements into the existing kit. Within this toolkit framework, we have generated a number of reagents to enable gene expression within immune and non-immune cell types, an array of best-in-class fluorescent proteins to visualize cell populations and transgenes as well as tools to simplify genetic manipulation, purification and ablation of targeted cells. Unlike recombination-based systems, the Golden Gate approach is also expandable, allowing the incorporation of complex designs such as multi-fragment promoters within the established modular framework of ImPaqT. Here, we demonstrate the function of this new system by generating a number of novel transgenic immune reporter lines. While our toolkit is focused on the immune system as an emerging area of study within zebrafish research, the ImPaqT approach can be broadly adapted to the construction of almost any zebrafish transgene, offering new tools for the generation of transgenes within the zebrafish community.

Introduction

The zebrafish (*Danio rerio*) has emerged as a powerful model organism and is increasingly used in various research fields and disciplines. Zebrafish share 70% of their genome and more than 80% of disease genes with humans (Howe et al. 2013). This has led to the development of many distinct zebrafish disease models (Santoriello and Zon 2012) including models of hematopoietic and neurodegenerative diseases, cardiovascular diseases (Bowley et al. 2022) such as atherosclerosis (Tang et al. 2021), cancer (Liu and Leach 2011; Russo et al. 2022), and tuberculosis (Bouz and Al Hasawi 2018; Varela and Meijer 2022), among others. Transgenic animals continue to play a key role for many experimental strategies, including study of gene function, genetic screens, and development of disease models. Transgenic zebrafish lines have been widely generated by applying the Tol2 transposon system with a transgenesis rate of 50% (Kawakami et al. 2004). In this system, a transposon-donor plasmid containing a DNA sequence of interest flanked by *Tol2* sites and synthetic mRNA encoding the Tol2 transposase are introduced into the single cell of fertilized eggs by microinjection. Upon translation, the transposase protein integrates the *Tol2* flanked gene fragment randomly into the genome through a cut-and-paste mechanism without causing rearrangement or modifications at the target site (Kawakami and Shima 1999; Kawakami, Shima, and Kawakami 2000; Kawakami et al. 2004; Kawakami 2007).

A number of approaches have previously been used to generate Tol2 transgenesis constructs. Traditionally, multi-step restriction enzyme and ligase based subcloning was used. However, this has given way to many advanced cloning approaches, including Gateway cloning (Hartley, Temple, and Brasch 2000; Kwan et al. 2007; Katzen 2007), Gibson Assembly (Gibson et al. 2009), In-Fusion cloning (Zhu et al. 2007; Sleight et al. 2010) and Golden Gate cloning, which have been used to speed up and simplify the procedure of transgene assembly. Each of these approaches have different advantages and disadvantages. Gateway cloning allows for the establishment and reuse of a library of fragments (e.g. promoters, genes and fluorescent proteins) but requires the use of lambda phage-derived *att* sequences to facilitate insertion of

the desired sequences. These *att* sequences result in the addition of ~20-25 bp of additional sequence in between each element and expansion of the number of fragments to be assembled can require recloning of the sequence with new *att* ends (Katzen 2007; Engler, Kandzia, and Marillonnet 2008; Sone et al. 2008). By contrast, Gibson and In-Fusion cloning allow for joining of short (15-20 bp) regions of homology, enabling the seamless cloning of many transgenes. However, the requirement for homology for seamless cloning generally means that individual components have to be adapted to upstream and downstream elements to generate new transgenes and have an upper limit of approximately 5-6 fragments with currently available kits. Golden Gate cloning utilizes type IIS restriction enzymes to cleave and join multiple fragments in a one pot reaction (Padgett and Sorge 1996; Engler, Kandzia, and Marillonnet 2008). It balances the tradeoffs of the previous cloning systems, requiring inclusion of only a short 4 base pair sequence, while enabling researchers to generate a library of vectors that can be combinatorially assembled to generate distinct transgenes. Large numbers of DNA fragments can also be assembled by Golden Gate with reports of assembly of up to 52 fragments in a single reaction (Pryor et al. 2022). Despite the advantages of Golden Gate cloning, this approach has not been widely applied in zebrafish transgene assembly. Only a few examples of this approach have been described for Tol2 transgenesis including the recently described GoldenFish toolkit which uses Golden Gate cloning to gain more flexibility in transgene design (Jiang et al. 2023) and the Golden Gateway tool (Kirchmaier, Lust, and Wittbrodt 2013) which combines the well-established cloning tools Gateway and Golden Gate cloning to simplify creation of new components for Gateway assembly of transgenes.

Here, we have sought to apply Golden Gate cloning approaches to create a toolkit for zebrafish transgenesis. We have developed ImPaqT, a new **Im**munological toolkit for **Paq**CI-based Golden Gate Assembly of Tol2 **T**ransgenes, to facilitate the production of transgenic animals. Studies of immune function have been of growing interest in the zebrafish community due to the close genetic similarity of the immune systems of humans and zebrafish (Howe et al. 2013), but there are few compiled resources for these cell types. ImPaqT provides a collection of

commonly used immunological promoters targeting distinct lineages of vertebrate immunity, genetic tools and fluorophores supporting flexible generation of transgenes for immunological research. This toolkit utilizes Golden Gate cloning due to the advantages of this technique and incorporates a newly described type IIS restriction enzyme, PaqCI, with a 7 base recognition sequence to reduce the frequency of off target cutting (from 1 in 4,096 bases (4^6) for a 6-base cutter to 1 in 16,384 bases (4^7), assuming random cutting). We designed the kit to have consistent overhang sequences for each position in the backbone (e.g. 5', middle, and 3' element) in order to make the toolkit modular and simplify the construction of additional transgenes. As proof-of-concept, we show that this toolkit can facilitate the production of expression constructs and we demonstrate the generation of transgenic animals using the Tol2 transgenesis with these constructs. ImPaqT is modular, enabling easy swapping of gene sequences, easily expandable, allowing the introduction of multiple new genetic elements if needed, and relies only on the use of short (4 bp) overlaps that can also be designed to be scarless if desired. To establish an immunological toolkit that will be broadly useful to zebrafish researchers, we have generated a variety of immunological and non-immunological promoters, including *mfap4* (macrophages)*, irg1* (macrophages, induced by infection) *, lyz* (neutrophils), *lck* (T cells), *ubb* (ubiquitious) *and hsp70* (heat shock inducible), tools for genetic recombination (*icre*), cell killing (*kid, nitroreductase)* and cell isolation tools (*LNGFR*), and an array of fluorescence proteins including *mTurquoise2*, *tdStayGold*, *mCitrine*, *mNeonGreen*, *dLanYFP* and td*Tomato* that can be used to rapidly develop new transgenes and new transgenic lines.

Results

<u>PaqCI-based Golden Gate cloning enables the construction of Tol2 expression vectors</u>

We sought to create a toolkit for immune transgene construction for the zebrafish community. Ultimately, we decided to design this toolkit to use Golden Gate cloning, given the short regions of homology that this approach required, the ability to structure this toolkit in a modular fashion and the expandability of the Golden Gate cloning approach, enabling incorporation of additional genetic elements as needed. Golden Gate assembly uses type IIS restriction enzymes to generate a set of unique overhangs that are used to join fragments in a desired order. To create a system for modular cloning of transgenes, we chose consistent 4-base overhang sequences for each of the three elements such that DNA fragments amplified with these overhangs can be cloned in the same position in the final construct, allowing these elements to be used combinatorially. As zebrafish transgenes generally consist of a cell-type specific promoter, a gene of interest, and a fluorescent marker, we structured our toolkit to have three elements that would ultimately be joined within a backbone vector possessing Tol2 repeats for transgenesis. These elements were named based on their location within the final construct - 5' element (5E) for the first position, a middle element (ME) and a 3' element (3E) for the final position within the backbone.

To generate the overhangs required to join individual components for Golden Gate assembly, we chose the 7-base cutter PaqCI to limit off-target cutting of inserts used in assembly. Genes and regions of interest were generated by PCR amplification using primers containing the 7-base PaqCI recognition sequence along with a spacer consisting of any four nucleotides that will be excised from the final construct, and a 4-base overhang sequence specific to the location of the sequence of interest in the final construct (Fig. S1). These PCR products were cloned into vectors and sequence verified to generate a stable library of constructs for transgene assembly. Cleavage with PaqCI leads to the creation of specific 4-base overhangs for each position at their 5' and 3' end enabling proper assembly and single-step ligation into a backbone vector for transgene generation. These backbone vectors possess inverted *Tol2*

135 repeats for transgenesis surrounding a PaqCI-flanked LacZ gene for blue-white screening to

136 identify backbone constructs containing transgene components. A simplified schematic for the

137 assembly using one insert is illustrated in Fig. 1. As we demonstrated below, we have found

138 that this approach allows for efficient, combinatorial transgene generation.

Figure 1: Basic principles of Golden Gate cloning for Tol2 Transgenesis
Schematic of a Golden Gate cloning reaction using one insert. The insertion construct and backbone
vector are assembled in a single-tube reaction using PaqCI and ligase. PaqCI is a type IIS restriction
enzyme. Its recognition site and cleavage are indicated. The font color corresponds to the color of the
respective feature indicated in the cartoon. The backbone vector consists of Tol2 sites, PaqCI
recognition sites (PaqCI) along with 4-base spacer sequences (N$_4$) and two overhangs (O). The insertion
construct includes the gene of interest (GOI) flanked by PaqCI, N$_4$ and the same two overhangs.
Assembly involves digestion of backbone and insertion constructs by PaqCI producing matching
overhangs and ligation through the DNA ligase forming the final assembly product.

<u>ImPaqT facilitates the flexibility of transgenes design and the generation of transgenic</u>

<u>zebrafish</u>

Having established this Golden Gate cloning-based technique for the assembly of Tol2

transposon vectors, we sought to create an array of constructs that would be beneficial to the

immunology community to generate transgenic animals. To these ends, we have generated

constructs for expression in commonly studied zebrafish immune populations, including T cells

(*lck*) (Langenau et al. 2004)*,* neutrophils (*lyz*) (Hall et al. 2007) and macrophages (*mfap4* and

the infection-inducible *irg1*) (Walton et al. 2015; Hall et al. 2013). To enable lineage tracing

approaches for labeling of hematopoietic derived cells, we also included the hematopoietic

stem cell (HSC)-specific promoter *runx1 +23* (Tamplin et al. 2015) as well as tools for broader

expression including the ubiquitious promoter *ubiquitin B* (*ubb*) (Mosimann et al. 2011), and

the temperature inducible heat shock protein 70-like promoter (*hsp70*) (Shoji, Yee, and

Kuwada 1998; Halloran et al. 2000; Kwan et al. 2007). We have also integrated a number of

tools to allow for manipulation of these immune populations, including tools for genetic

manipulation (*icre* (Shimshek et al. 2002)), for rapid purification of cell populations (truncated

human *LNGFR* (Wattrus and Zon 2020)) and cell-specific ablation (*nitroreductase* (Curado et

al. 2007; Curado, Stainier, and Anderson 2008) and *kid* (Labbaf et al. 2022)). To visualize

labeled cell populations and genes, we provide a diverse array of best-in-class fluorescent

proteins in a range of colors to facilitate simultaneous imaging of distinct cell types. These

include *tdStayGold* (Hirano et al. 2022), *mTurquoise2* (Goedhart et al. 2012), *tdTomato*

(Shaner et al. 2004), m*Citrine* (Zacharias et al. 2002)*, mNeonGreen* (Shaner et al. 2013) and

dLanYFP (Shaner et al. 2013). A summary of all available constructs is shown in Table 1.

ImPaqT is also designed to be easily expandable through simple cloning approaches.

Table 1: Library of insertion constructs.
5' element (5E), middle element (ME), 3' element (3E).

Insertion Elements	Position	Function	Reference
irg1	5E	Infection-inducible macrophage-specific promoter	(Hall et al. 2013)
mfap4	5E	Macrophage-specific promoter	(Walton et al. 2015)
lyz	5E	Neutrophil-specific promoter	(Hall et al. 2007)
ubiquitin B (*ubb*)	5E	Promoter for ubiquitious expression	(Mosimann et al. 2011)
ubb:loxP GFP loxP	5E	Promoter for ubiquitious expression of floxed GFP for use in combination with Cre	(Mosimann et al. 2011)
hsp70	5E	heat shock protein 70-like promoter for inducible gene expression	(Shoji, Yee, and Kuwada 1998; Halloran et al. 2000; Kwan et al. 2007)
lck	5E	T cell-specific promoter	(Langenau et al. 2004)
runx1 + 23	5E	Promoter for expression in hematopoietic stem cells	(Tamplin et al. 2015)
icre	ME	Cre recombinase for inducible recombination of DNA between loxP sites	(Shimshek et al. 2002)
(p2A) dLanYFP	3E	Yellow fluorescent protein	(Shaner et al. 2013)
mNeonGreen	ME, 3E	Green/ yellow fluorescent protein	(Shaner et al. 2013)

LNGFR	ME	Low affinity nerve growth factor receptor for rapid cell enrichment	(Wattrus and Zon 2020)
nitroreductase	ME	Inducible ablation of specific cell lineages	(Curado et al. 2007; Curado, Stainier, and Anderson 2008)
kid	ME	Ablation of specific cell lineages	(Labbaf et al. 2022; de la Cueva-Mendez et al. 2003)
(p2A) tdTomato	ME, 3E	Red fluorescent protein	(Shaner et al. 2004)
tdTomato CAAX	ME	Red fluorescent protein with membrane localization protein	(Walton et al. 2015)
(p2A) mTurquoise2	ME, 3E	Cyan fluorescent protein	(Goedhart et al. 2012)
mCitrine	ME	Yellow fluorescent protein	(Zacharias et al. 2002)
tdStayGold	ME	Green fluorescent protein	(Hirano et al. 2022)
ubb:polyA	3E	Ubiquitious expression of a SV40 poly A	(Walton et al. 2015)

 Generation of new transgenic zebrafish lines using ImPaqT

To test our PaqCI-based Golden Gate assembly toolkit as well as generate new immunological

zebrafish lines for the community, we used ImPaqT to assemble a number of new transgenic

lines. Using the neutrophil-specific promoter *lyz*, we generated a T*g(lyz:mTurquoise2)*[ber2]

construct through Golden Gate assembly of 5E *lyz*, ME *mTurquoise2* and 3E *ubb:polyA* to

create the *lyz*:m*Turquoise2* transgene construct (Fig. 2A-B). The construct was coinjected with

Tol2 transposase mRNA into single cell of fertilized eggs and at 2 dpi we observed strong blue

fluorescence signal within likely neutrophil populations in the CHT (Fig. 2C). Positive animals

were raised to adulthood and screening of F1 progeny of these animals revealed robust

expression of mTurquoise2 within putative neutrophil populations throughout the animal (Fig.

2D). Crossing to the already established neutrophil reporter *Tg(lyz:EGFP)*[nz117] (Hall et al. 2007)

zebrafish confirmed specific expression of our transgene within neutrophils (Fig. S2).

Figure 2: Generation of *Tg(lyz:mTurquoise2)* zebrafish
A: Schematic of the Golden Gate assembly of *Tg(lyz:mTurquoise2)*. The backbone vector contains overhang sequences O1 (TTCC) and O4 (AGGA). *lyz* is a neutrophil-specific promoter containing overhang sequences O1 and O2 (CAAA). *mTurquoise2* encodes a bright cyan fluorescent protein with overhangs O2 and O3 (GCTA). *ubb:polyA* is assembled at the final position within the backbone with O3 and O4.B: Schematic of the final assembly product of *Tg(lyz:mTurquoise2)*. C-D: Images of a *Tg(lyz:mTurquoise2)* F0 (C) and F1 (D) larva showing neutrophil response in cyan at 2 days post fertilization.

Similarly, we used the ImPaqT toolkit to generate a new macrophage-specific transgene using

the infection-inducible macrophage-specific promoter *irg1* and the recently described bright

green fluorescent protein *tdStayGold* (Hirano et al., 2022). To generate the

Tg(irg1:tdStayGold)[ber3] construct, the 5E *irg1,* ME *tdStayGold* and 3E *ubb:polyA* elements were

combined into the backbone vector (Fig. 3 A-B). We injected zebrafish with this construct and

as *irg1* is an inflammation responsive promoter that is only weakly expressed in the absence

of immune stimulation, we injected larvae with LPS before screening. After LPS injection, we

were able to identify a number of F0 founders with strong green fluorescence within

subpopulations of their macrophages (Fig. 3C). These positive F0 larvae were raised to

adulthood and screening of F1 progeny revealed expression of tdStayGold throughout

macrophage populations upon stimulation with LPS (Fig. 3 D).

Figure 3: Generation of *Tg(irg1:tdStayGold)* zebrafish
A: Schematic of the Golden Gate assembly of *Tg(irg1:tdStayGold)*. The backbone vector contains
overhang sequences O1 (TTCC) and O4 (AGGA). *irg1* is a macrophage-specific promoter containing
overhang sequences O1 and O2 (CAAA). *tdStayGold* encodes a bright green fluorescent protein with
overhangs O2 and O3 (GCTA). *ubb:polyA* is assembled at the final position within the backbone with
O3 and O4. B: Illustration of the final assembly product of *Tg(irg1:tdStayGold)*. C-D: Images of a
Tg(irg1:tdStayGold) F0 (C) and F1 (D) larva showing macrophage-specific tdStayGold expression.

We also aimed to test the utility of the tools that we have designed within our kit. To these

ends, we created an infection-inducible, macrophage specific *icre* transgene construct,

Tg(irg1:icre-p2A mTurquoise2). This construct was generated by combining the 5E *irg1,*

ME *icre* and 3E *p2A mTurquoise2* (Fig. 4A and 4B)*. Animals injected with this transgene

showed macrophage-specific mTurquoise2 signal after injection with LPS (Fig. 4C). To further

confirm that this construct was able to drive rearrangement of floxed reporters, we injected

Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry)[cz1701] (Mosimann et al. 2011) zebrafish with this

construct. These animals were injected with LPS and imaged for 48 h to visualize

rearrangement of the floxed construct. LPS treatment drove the induction of *icre* and the co-

translationally cleaved *p2a mTurquoise2* transgene within macrophages starting ca. 4 h post

LPS injection. While the induction of the *irg1:icre-p2a mTurquoise2* transgene was transitory,

this short-term induction of Cre could induce permanent rearrangement of the *loxP-EGFP-*

loxP-mCherry construct as seen by induction of red fluorescent mCherry within macrophage

populations (Fig. 4D, S3, Movie S1). Control animals without the *icre*-construct neither showed

mTurquoise2 nor mCherry signal in macrophages (Fig. S3, Movie S 2). This experiment

showed not only the functionality of the individual insertion constructs but also the range of

assembly constructs that can be generated with them.

D

Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry) + Tg(irg1:icre-p2A mTurquoise2)

Figure 4: Generation of *Tg(irg1:icre-p2A mTurquoise2)* zebrafish for lineage tracing of individual macrophages

A: Schematic of the Golden Gate assembly of *Tg(irg1:icre-p2A mTurquoise2)*. The backbone vector contains overhang sequences O1 (TTCC) and O4 (AGGA). The macrophage promoter *irg1* contains overhang sequences O1 and O2 (CAAA). *icre* contains overhangs O2 and O3 (GCTA) for the middle position. *p2A mTurquoise2* is assembled at the final position within the backbone with O3 and O4. B: Illustration of the final assembly product of *Tg(irg1:icre-p2A mTurquoise2)*. C: Image of a *Tg(irg1:icre-p2A mTurquoise2)* F0 larva showing macrophage-specific mTurquoise2 signal after stimulation with LPS. D: Images of a *Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry)* larva that was introduced to the *Tg(irg1:icre-p2A mTurquoise2)* construct at single cell stage and was treated with PTU and LPS. The images show a macrophage-specific mTurquoise2 signal (e.g. 5 h post LPS injection) and a macrophage-specific mCherry signal (e.g. 33 h post LPS injection).

ImPaqT is expandable through the introduction of new overhang sequences

A limitation of many existing systems for transgenesis is that these systems have a fixed number of elements that can be combined to generate transgenes. Golden Gate methods however, are expandable through the addition of new overhangs and have been used to join 10-24 or even more than 50 fragments of DNA in a single reaction (Kucera 2021; Pryor et al. 2022). The ability to expand this kit to include larger numbers of constructs would simplify the generation of complex transgenes including multifragment promoters, loxP sites, localization tags (e.g NLS or prenylation signal) or multigene constructs for example. To provide proof of concept for the expandability of ImPaqT, vectors were designed to enable the construction of a *ubb* promoter-driven construct from three similar sized *ubb* promoter fragments, ubb_1, ubb_2 and ubb_3. These fragments were constructed such that the first fragment (ubb_1) and the last fragment (ubb_3) use the same O1 and O2 overhangs used by 5E constructs so that the array of promoter fragments can be cloned into the existing position of 5E elements and used alongside already existing ME and 3E constructs (Fig. 5A-B). New internal overhangs were designed for the ubb_1, ubb_2 and ubb_3 fragments to combine these distinct fragments (Fig. 5A-D). Individual *ubb* fragments were used for Golden Gate assembly in combination with ME *mTurquoise2* and 3E *ubb:polyA* (Fig. 5C) which led to the expected transgenesis construct *Tg(ubb:mTurquoise2)*[ber4] (Fig. 5D) that was injected into embryos at single cell stage. Screening of injected animals confirmed that transgenic embryos established with this construct had the expected expression pattern (Fig. 5E). We also found robust expression in the F1 progeny (Fig. 5F) after raising positive F0 animals, confirming that injected animals are viable and able to pass on the transgene to the next generation. Together, these findings demonstrated proof-of-concept for the expansion of this transgene assembly system enabling users to add fragments as needed through the addition of novel unique 4-base overhangs while maintaining compatibility with existing ImPaqT constructs.

Figure 5: ImPaqT allows for expandability of individual positions while retaining compatibility with already established transposon components.

A: Schematic of the 5E *ubiquitin B* (*ubb*) promoter construct containing PaqCI recognition sites (PaqCI) along with 4-base spacer sequences (N_4) and two overhangs (O1: TTCC, O2: CAAA) for positioning at the 5' end of the assembly construct. Primers were designed to break up *ubb* into three fragments with new overhang sequences. B: Schematic of PCR amplified *ubb* fragments *ubb$_{1-3}$* with PaqCI, N_4 and O1 and O1A (TAGC), O1A and O1B (TTGA), and O1B and O2, respectively. C: Schematic of the Golden Gate assembly of 5E *ubb$_{1-3}$*, ME *mTurquoise2* and 3E *ubb:polyA* into the backbone vector. D: Illustration of the final assembly product of *Tg(ubb$_{1-3}$:mTurquoise2)*. E-F: Images of a *Tg(ubb:mTurquoise2)* F0 (E) and F1 (F) larva showing ubiquitous mTurquoise2 signal.

Discussion

Transgenic animals remain essential for many research questions. They enable the visualization of cells and tissues, molecular pathways, immune responses and the perturbation of these pathways to identify the molecular mechanisms driving pathology. Use of the zebrafish as a model organism has been steadily increasing in many research disciplines due to the unique advantages of this model. This is true in immunology and host-pathogen interactions, where the optical transparency afforded by the zebrafish model have led to many research discoveries, but where, as a new immunology model, there are limited transgenic tools. Here, we demonstrate a new cloning toolkit for the generation of immunological transgenes that is based on Golden Gate cloning. The Golden Gate approach results in shorter overlap regions (4 bp) than recombination-based techniques like Gateway cloning, while compared to seamless, homology based cloning methods such as Gibson cloning, it allows for straightforward modular cloning approaches. The Golden Gate approach also takes advantage of off-the-shelf restriction enzymes and ligase, compared to other cloning approaches which use proprietary enzymes for cloning.

We have termed this toolkit ImPaqT - **Im**munological toolkit for **Paq**CI-based Golden Gate Assembly of Tol2 **T**ransgenes. It includes a range of immunologically focused promoters along with genetic tools and fluorescent proteins that can be assembled using PaqCI-based Golden Gate cloning. These tools will enable the flexible assembly of transgenes to drive expression and to genetically manipulate, purify and ablate immune cell populations of interest. While zebrafish possess the same immune lineages of other vertebrates, as a relatively new model for immunology studies, the zebrafish has a more limited array of cell type-specific promoters than mouse models. For instance, despite the identification of potential NK- and ILC-specific (Hernandez et al. 2018) as well as mast cell-specific (Dobson et al. 2008) genes, promoters specific for these cell populations have not been described. As new immune-specific promoters are identified, these can be readily added to this toolkit. While we have targeted the immune system as a particular area of need, this Golden Gate approach can also be adapted to any other tissue or cell lineages within the zebrafish.

While Golden Gate cloning enables the use of shorter homology stretches and the

combinatorial use of constructs for the assembly of diverse transgenes, as a restriction

enzyme-based system, a limitation of Golden Gate systems, compared to other assembly

approaches, has been the risk of off-target cutting within the cloned sequences. This off-target

cutting requires mutation of insert sequences to eliminate these sites (termed domestication),

(Marillonnet and Grutzner 2020; Bird, Marles-Wright, and Giachino 2022; Grützner and

Marillonnet 2020), which can potentially alter function of domesticated promoters and genes.

While domestication can result in changes in the sequence that may alter behavior of the

genetic element, these in principle can be managed for genes of interest using silent mutations

in critical bases. While promoters cannot be managed through the same approach, promoter

fragments used for zebrafish are frequently large (commonly 2-6 kb) and it is unlikely that the

mutation of a single base pair will inadvertently ablate the promoter activity. Consistent with

this, we have not seen any changes in promoter activity or localization in the two promoters

that we domesticated for the ImPaqT toolkit, although further cloning of additional promoters

will more completely test this idea. Our choice of PaqCI itself seeks to address the limitations

of domestication, as PaqCI uses a 7 base recognition site. In contrast to the 6 base-cutters

that have been described in previous zebrafish Golden Gate approaches (Kirchmaier, Lust,

and Wittbrodt 2013; Jiang et al. 2023), the rare cutting of PaqCI (if random, 1 in 16,384 bases)

makes this enzyme better suited for the large DNA fragments that are frequently used for

zebrafish promoters.

A limitation of existing systems has been the use of a fixed number of constructs for assembly

of the transgene. One benefit of the Golden Gate directed system is that it can be expanded

for cloning of complex transgenes, with some Golden Gate assemblies encompassing 20 or

even more than 50 fragments (Kucera 2021; Pryor et al. 2022). While we have designed the

system to incorporate 4 distinct DNA segments (3 inserts and a backbone vector) the flexibility

of Golden Gate cloning means that our system can be readily expanded and contracted while

maintaining compatibility with the other components of the kit. As proof-of-concept, we were

able to break a single 5E encoding the *ubb* promoter into 3 distinct plasmids that we could

subsequently assemble together with existing ME, 3E and backbone elements to generate a

functional *ubb* driven transgene. The flexibility of this system will simplify the construction of

large and complex transgenes. Potential applications would include breaking up larger

promoter fragments into smaller and easier to clone elements, insertion of recombination sites

for enzymes like Cre or Flp for rearrangement, assembly of multipromoter constructs encoding

genes such as Cas9 and CRISPR gRNAs for cell type-specific knockout, or the inclusion of

epitope or localization tags.

Lastly, while we have applied this Golden Gate method for the construction of Tol2 transgenes,

the advantages of the PaqCI-Golden Gate method will likely be useful for the editing of

zebrafish by other approaches as well. For instance, CRISPR insertion has been increasingly

used to generate knock-in zebrafish lines, via either homologous recombination, non-

homologous methods or by using short homology stretches. Knock-in approaches require the

construction of donor vectors which incorporate varied sequences such as homology regions,

fluorescent proteins and stretches of genomic DNA (Albadri, Del Bene, and Revenu 2017).

Similarly, many groups also use I-SceI insertion approaches to generate transgenic zebrafish

(Hoshijima, Jurynec, and Grunwald 2016). The PaqCI Golden Gate method could be adapted

to simplify the construction of transgenes and editing constructs for these other approaches as

well.

Materials and Methods

Ethics statement

Zebrafish husbandry and experiments were approved by the "Landesamt für Gesundheit und Soziales" (LaGeSo) Berlin (Germany).

Zebrafish husbandry and handling

Zebrafish (*Danio rerio*) embryos were raised in petri dishes containing E3 medium at 28°C for up to five days. Animals older than 5 days were maintained in a circulating system with constant water conditions of 28°C, pH 7.0 - 7.3, and a conductivity of 600-700 µS on a 12-hour light/ 12-hour dark cycle. They were fed twice a day with artemia and once a day with dry food.

Generation of transgenic zebrafish lines

Zebrafish strains

All zebrafish strains, including *Tg(lyz:mTurquoise2)^ber2^, Tg(irg1:tdStayGold)^ber3^,Tg(irg1:icre-p2A mTurquoise2)* and *Tg(ubb:mTurquoise2)^ber4^,* were in the *AB wildtype background and generated in this study by applying the Tol2 transgenesis system as previously described (Kawakami et al., 2004*). Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry)^cz1701^* zebrafish were obtained from the European Zebrafish Resource Center (EZRC). *Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry)^cz1701^* (Mosimann et al. 2011) and *Tg(lyz:EGFP)^nz117^* (Hall et al. 2007) have been previously described.

Preparation of the backbone vector and inserts

The backbone vector containing the LacZ gene flanked by PaqCI sites and *Tol2* arms in a vector backbone with a pMB1 origin and an AmpR gene for bacterial selection (pTWIST Amp High Copy) was purchased from Twist Bioscience. The complete sequence is shown in Figure S4. Primer sequences are shown in Table S1. Gene fragments for *runx1+23* (with the endogenous PaqCI site mutated), *tdStayGold* and truncated human *LNGFR* (zebrafish codon optimized and with endogenous PaqCI sites mutated) were purchased from Twist Bioscience

and TOPO cloned into pCRII TOPO (Thermofisher Zero Blunt TOPO). The complete

sequences are shown in Figure S5.

PaqCI recognition sites with 4-base spacer sequences and 4-base overhangs were added to

DNA sequences via PCR amplification. Primer were designed to add PaqCI sites such as

cleavage resulted in a gene fragment with a 4-base overhang that is compatible to the

overhang of another gene fragment that will be adjacent to it in the final assembly. Appropriate

4-base overhang sequences were confirmed using the online tool NEBridge Ligase Fidelity

Viewer™. PCR amplification was performed using Q5 polymerase and analyzed by gel

electrophoresis. The PCR product was cleaned up or, in case of multiple bands, the band of

correct size was excised under 70% UV light using a clean scalpel and DNA was extracted

using Nucleo Spin Gel and PCR Clean-up Kit (Macherey-Nagel). DNA sequences were

confirmed by sequencing. Gel purified gene fragments were subcloned into a Zero Blunt TOPO

vector (Invitrogen) and transformed into competent NEB 5α *E. coli* cells (C2987H). Cells were

cultivated in LB medium containing 50 µg/ml Kanamycin and plasmids were isolated using the

Nucleo Spin Plasmid Quick Pure protocol (Macherey-Nagel).

Both the *ubb* and *lck* promoters had endogenous PaqCI cleavage sites in the promoter

sequence. To domesticate these promoters, we mutated these PaqCI sites with the indicated

PCR primers (Table S1) and assembled them with In-Fusion cloning (Takara Bio).

<u>Golden Gate based assembly of expression constructs</u>

Golden Gate assembly was performed according to the NEB protocol. In detail, 75 ng of the

backbone vector and each insertion construct was mixed with 10 U PaqCI, 0.5 µl PaqCI

activator (20 µM) and 400 U T4 DNA ligase in T4 DNA ligase buffer to a final volume of 20 µl.

Final assembly product was obtained after 60 Cycles of incubation at 37°C for 1 min and

incubation at 16°C for 1 min, followed by 5 min incubation at 37°C and 5 min incubation at

60°C using a thermocycler. The final product was transformed into competent NEB 5α *E. coli*

cells (C2987H) and cells were cultivated in LB medium containing 20 µg/ml Ticarcillin.

Plasmids were isolated using the Nucleo Spin Plasmid Quick Pure protocol (Macherey-Nagel)

and PCR purified using the Nucleo Spin Gel and PCR Clean-up Kit (Macherey-Nagel) before

microinjection.

<u>Microinjections</u>

The injection mixture contained 25 ng/µL polyadenylated mRNA encoding the Tol2

transposase, 50 ng/µL transposon donor plasmid and 10% phenol red in a final volume of 10 µl

1 X Tango Buffer (Thermo Fisher). The mMessage mMachine kit (Life Technologies) was used

to transcribe the mRNA from T3TS-Tol2 (Balciunas et al. 2006). A volume of ~1 nL was injected

into the single cell of fertilized eggs by using a microinjector.

<u>Imaging</u>

Injected larvae were screened for insertion of the transgene 2 days post injection (dpi) using a

Nikon ECLIPSE Ti2 microscope (Nikon) with a Dragonfly 200 confocal system (Oxford

Instruments). Animals that were positive for the transgene were raised to adulthood and

outcrossed to the *AB wildtype strain for selection of transgenic founder zebrafish. Before

screening *Tg(irg1:icre-p2A mTurquoise2)* and *Tg(irg1:tdStayGold)* embryos, the activity of the

irg1 promoter was induced by injection of LPS. For that purpose, animals were anesthetized

with tricaine (MS-222) at a final concentration of 0.016% and injected into the caudal vein with

LPS (Sigma-Aldrich, F3665) (final: 0.8 mg/ml). Screening was performed at least 4 h after

injection. To reduce pigmentation, *Tg(-3.5ubb:LOXP-EGFP-LOXP-mCherry)* embryos were

treated with 45 µg/ml 1-phenyl-2-thiourea (PTU) starting ca. 20 h post fertilization. Image

analysis was performed using the Fiji ImageJ software (National Institute of Health, USA).

<u>Line and reagent availability</u>

All new lines and plasmids are available from the corresponding author upon contact and will

also be deposited to AddGene.

Acknowledgements

We would like to thank Miriam Herbert for sharing the ME *kid* construct. Additionally, we would

like to thank our animal welfare officer Antje Dreyer and our animal caretakers Silke Lehmann,

Marten Walk, Carsten Weiland, Ines Neumann, Jens-Phillip Otto und Janine Bleske for their

great work and support.

Competing interests

No competing interests declared.

Funding

This work was supported by the Max Planck Society.

Data availability

All relevant data can be found within the article and its supplementary information.

References

Albadri, S., F. Del Bene, and C. Revenu. 2017. 'Genome editing using CRISPR/Cas9-based knock-in approaches in zebrafish', *Methods*, 121-122: 77-85.

Balciunas, D., K. J. Wangensteen, A. Wilber, J. Bell, A. Geurts, S. Sivasubbu, X. Wang, P. B. Hackett, D. A. Largaespada, R. S. McIvor, and S. C. Ekker. 2006. 'Harnessing a high cargo-capacity transposon for genetic applications in vertebrates', *PLoS Genet*, 2: e169.

Bird, J. E., J. Marles-Wright, and A. Giachino. 2022. 'A User's Guide to Golden Gate Cloning Methods and Standards', *ACS Synth Biol*, 11: 3551-63.

Bouz, G., and N. Al Hasawi. 2018. 'The zebrafish model of tuberculosis - no lungs needed', *Crit Rev Microbiol*, 44: 779-92.

Bowley, G., E. Kugler, R. Wilkinson, A. Lawrie, F. van Eeden, T. J. A. Chico, P. C. Evans, E. S. Noel, and J. Serbanovic-Canic. 2022. 'Zebrafish as a tractable model of human cardiovascular disease', *Br J Pharmacol*, 179: 900-17.

Curado, S., R. M. Anderson, B. Jungblut, J. Mumm, E. Schroeter, and D. Y. R. Stainier. 2007. 'Conditional Targeted Cell Ablation inZebrafish: A New Tool for RegenerationStudies', *Developmental Dynamics*.

Curado, S., D. Y. Stainier, and R. M. Anderson. 2008. 'Nitroreductase-mediated cell/tissue ablation in zebrafish: a spatially and temporally controlled ablation method with applications in developmental and regeneration studies', *Nat Protoc*, 3: 948-54.

de la Cueva-Mendez, G., A. D. Mills, L. Clay-Farrace, R. Diaz-Orejas, and R. A. Laskey. 2003. 'Regulatable killing of eukaryotic cells by the prokaryotic proteins Kid and Kis', *EMBO*.

Dobson, J. T., J. Seibert, E. M. Teh, S. Da'as, R. B. Fraser, B. H. Paw, T. J. Lin, and J. N. Berman. 2008. 'Carboxypeptidase A5 identifies a novel mast cell lineage in the zebrafish providing new insight into mast cell fate determination', *Blood*, 112: 2969-72.

Engler, C., R. Kandzia, and S. Marillonnet. 2008. 'A one pot, one step, precision cloning method with high throughput capability', *PLoS One*, 3: e3647.

Gibson, D. G., L. Young, R. Y. Chuang, J. C. Venter, C. A. Hutchison, 3rd, and H. O. Smith. 2009.
'Enzymatic assembly of DNA molecules up to several hundred kilobases', *Nat Methods*, 6: 343-
5.
Goedhart, J., D. von Stetten, M. Noirclerc-Savoye, M. Lelimousin, L. Joosen, M. A. Hink, L. van Weeren,
T. W. Gadella, Jr., and A. Royant. 2012. 'Structure-guided evolution of cyan fluorescent
proteins towards a quantum yield of 93%', *Nat Commun*, 3: 751.
Grützner, R., and S. Marillonnet. 2020. 'Generation of MoClo Standard Parts Using Golden Gate
Cloning.' in Springer (ed.), *DNA Cloning and Assembly Methods and Protocols*.
Hall, C., M. V. Flores, T. Storm, K. Crosier, and P. Crosier. 2007. 'The zebrafish lysozyme C promoter
drives myeloid-specific expression in transgenic fish', *BMC Dev Biol*, 7: 42.
Hall, C. J., R. H. Boyle, J. W. Astin, M. V. Flores, S. H. Oehlers, L. E. Sanderson, F. Ellett, G. J. Lieschke, K.
E. Crosier, and P. S. Crosier. 2013. 'Immunoresponsive gene 1 augments bactericidal activity of
macrophage-lineage cells by regulating beta-oxidation-dependent mitochondrial ROS
production', *Cell Metab*, 18: 265-78.
Halloran, M. C., M. Sato-Maeda, Jr. Warren, J. T., F. Su, Z. Lele, P. H. Krone, J. Y. Kuwada, and W. Shoji.
2000. 'Laser-induced gene expression in specific cells of transgenic zebrafish', *Development*.
Hartley, J. L., G. F. Temple, and M. A. Brasch. 2000. 'DNA cloning using in vitro site-specific
recombination', *Genome Res*, 10: 1788-95.
Hernandez, P. P., P. M. Strzelecka, E. I. Athanasiadis, D. Hall, A. F. Robalo, C. M. Collins, P. Boudinot, J.
P. Levraud, and A. Cvejic. 2018. 'Single-cell transcriptional analysis reveals ILC-like cells in
zebrafish', *Sci Immunol*, 3.
Hirano, M., R. Ando, S. Shimozono, M. Sugiyama, N. Takeda, H. Kurokawa, R. Deguchi, K. Endo, K. Haga,
R. Takai-Todaka, S. Inaura, Y. Matsumura, H. Hama, Y. Okada, T. Fujiwara, T. Morimoto, K.
Katayama, and A. Miyawaki. 2022. 'A highly photostable and bright green fluorescent protein',
Nat Biotechnol, 40: 1132-42.
Hoshijima, K., M. J. Jurynec, and D. J. Grunwald. 2016. 'Precise Editing of the Zebrafish Genome Made
Simple and Efficient', *Dev Cell*, 36: 654-67.
Howe, K., M. D. Clark, C. F. Torroja, J. Torrance, C. Berthelot, M. Muffato, J. E. Collins, S. Humphray, K.
McLaren, L. Matthews, S. McLaren, I. Sealy, M. Caccamo, C. Churcher, C. Scott, J. C. Barrett, R.
Koch, G. J. Rauch, S. White, W. Chow, B. Kilian, L. T. Quintais, J. A. Guerra-Assuncao, Y. Zhou,
Y. Gu, J. Yen, J. H. Vogel, T. Eyre, S. Redmond, R. Banerjee, J. Chi, B. Fu, E. Langley, S. F. Maguire,
G. K. Laird, D. Lloyd, E. Kenyon, S. Donaldson, H. Sehra, J. Almeida-King, J. Loveland, S.
Trevanion, M. Jones, M. Quail, D. Willey, A. Hunt, J. Burton, S. Sims, K. McLay, B. Plumb, J.
Davis, C. Clee, K. Oliver, R. Clark, C. Riddle, D. Elliot, G. Threadgold, G. Harden, D. Ware, S.
Begum, B. Mortimore, G. Kerry, P. Heath, B. Phillimore, A. Tracey, N. Corby, M. Dunn, C.
Johnson, J. Wood, S. Clark, S. Pelan, G. Griffiths, M. Smith, R. Glithero, P. Howden, N. Barker,
C. Lloyd, C. Stevens, J. Harley, K. Holt, G. Panagiotidis, J. Lovell, H. Beasley, C. Henderson, D.
Gordon, K. Auger, D. Wright, J. Collins, C. Raisen, L. Dyer, K. Leung, L. Robertson, K. Ambridge,
D. Leongamornlert, S. McGuire, R. Gilderthorp, C. Griffiths, D. Manthravadi, S. Nichol, G.
Barker, S. Whitehead, M. Kay, J. Brown, C. Murnane, E. Gray, M. Humphries, N. Sycamore, D.
Barker, D. Saunders, J. Wallis, A. Babbage, S. Hammond, M. Mashreghi-Mohammadi, L. Barr,
S. Martin, P. Wray, A. Ellington, N. Matthews, M. Ellwood, R. Woodmansey, G. Clark, J. Cooper,
A. Tromans, D. Grafham, C. Skuce, R. Pandian, R. Andrews, E. Harrison, A. Kimberley, J. Garnett,
N. Fosker, R. Hall, P. Garner, D. Kelly, C. Bird, S. Palmer, I. Gehring, A. Berger, C. M. Dooley, Z.
Ersan-Urun, C. Eser, H. Geiger, M. Geisler, L. Karotki, A. Kirn, J. Konantz, M. Konantz, M.
Oberlander, S. Rudolph-Geiger, M. Teucke, C. Lanz, G. Raddatz, K. Osoegawa, B. Zhu, A. Rapp,
S. Widaa, C. Langford, F. Yang, S. C. Schuster, N. P. Carter, J. Harrow, Z. Ning, J. Herrero, S. M.
Searle, A. Enright, R. Geisler, R. H. Plasterk, C. Lee, M. Westerfield, P. J. de Jong, L. I. Zon, J. H.
Postlethwait, C. Nusslein-Volhard, T. J. Hubbard, H. Roest Crollius, J. Rogers, and D. L. Stemple.
2013. 'The zebrafish reference genome sequence and its relationship to the human genome',
Nature, 496: 498-503.

Iwanami, N. 2014. 'Zebrafish as a model for understanding the evolution of the vertebrate immune
system and human primary immunodeficiency', *Exp Hematol*, 42: 697-706.
Jiang, Z., L. Huang, J. Zhao, Y. Li, J. Ma, R. Ni, Q. Yang, L. Luo, Y. Yang, and J. Chen. 2023. 'GoldenFish: a
rapid and efficient system to customize constructs for zebrafish transgenesis', *J Mol Cell Biol*,
14.
Katzen, F. 2007. 'Gateway((R)) recombinational cloning: a biological operating system', *Expert Opin*
Drug Discov, 2: 571-89.
Kawakami, K. 2007. 'Tol2: a versatile gene transfer vector in vertebrates', *Genome Biol*, 8 Suppl 1: S7.
Kawakami, K. , and A. Shima. 1999. 'Identification of the Tol2 transposase of the medaka fish Oryzias
latipes that catalyzes excision of a nonautonomous Tol2 element in zebrafish Danio rerio',
Elsevier.
Kawakami, K., A. Shima, and N. Kawakami. 2000. 'Identification of a functional transposase of the Tol2
element, an Ac-like element from the Japanese medaka fish, and its transposition in the
zebrafish germ lineage', *Proc NatlAcad Sci USA*.
Kawakami, K., H. Takeda, N. Kawakami, M. Kobayashi, N. Matsuda, and M. Mishina. 2004. 'A
transposon-mediated gene trap approach identifies developmentally regulated genes in
zebrafish', *Dev Cell*, 7: 133-44.
Kirchmaier, S., K. Lust, and J. Wittbrodt. 2013. 'Golden GATEway cloning--a combinatorial approach to
generate fusion and recombination constructs', *PLoS One*, 8: e76117.
Kucera, R. 2021. "Using PaqCI™ for Golden Gate Assembly – What Makes it a Special Addition to NEB's
DNA Assembly Portfolio?" In.
Kwan, K. M., E. Fujimoto, C. Grabher, B. D. Mangum, M. E. Hardy, D. S. Campbell, J. M. Parant, H. J.
Yost, J. P. Kanki, and C. B. Chien. 2007. 'The Tol2kit: a multisite gateway-based construction kit
for Tol2 transposon transgenesis constructs', *Dev Dyn*, 236: 3088-99.
Labbaf, Z., K. Petratou, L. Ermlich, W. Backer, K. Tarbashevich, M. Reichman-Fried, S. Luschnig, S.
Schulte-Merker, and E. Raz. 2022. 'A robust and tunable system for targeted cell ablation in
developing embryos', *Dev Cell*, 57: 2026-40 e5.
Langenau, D. M., A. A. Ferrando, D. Traver, J. L. Kutok, J.-P. D. Hezel, J. P. Kanki, L. I. Zon, A. T. Look,
and N. S: Trede. 2004. 'In vivo tracking of T cell development, ablation, and engraftment in
transgenic zebrafish', *PNAS*.

Liu, S., and S. D. Leach. 2011. 'Zebrafish models for cancer', *Annu Rev Pathol*, 6: 71-93.
Marillonnet, S., and R. Grutzner. 2020. 'Synthetic DNA Assembly Using Golden Gate Cloning and the
Hierarchical Modular Cloning Pipeline', *Curr Protoc Mol Biol*, 130: e115.
Meeker, N. D., and N. S. Trede. 2008. 'Immunology and zebrafish: spawning new models of human
disease', *Dev Comp Immunol*, 32: 745-57.
Mosimann, C., C. K. Kaufman, P. Li, E. K. Pugach, O. J. Tamplin, and L. I. Zon. 2011. 'Ubiquitous transgene
expression and Cre-based recombination driven by the ubiquitin promoter in zebrafish',
Development, 138: 169-77.
Padgett, K. A., and J. A. Sorge. 1996. 'Creating seamless junctions independent of restriction sites in
PCR cloning', *Gene*.
Pryor, J. M., V. Potapov, K. Bilotti, N. Pokhrel, and G. J. S. Lohman. 2022. 'Rapid 40 kb Genome
Construction from 52 Parts through Data-optimized Assembly Design', *ACS Synth Biol*, 11:
2036-42.
Russo, I., E. Sartor, L. Fagotto, A. Colombo, N. Tiso, and M. Alaibac. 2022. 'The Zebrafish model in
dermatology: an update for clinicians', *Discov Oncol*, 13: 48.
Santoriello, C., and L. I. Zon. 2012. 'Hooked! Modeling human disease in zebrafish', *J Clin Invest*, 122:
2337-43.
Shaner, N. C., R. E. Campbell, P. A. Steinbach, B. N. Giepmans, A. E. Palmer, and R. Y. Tsien. 2004.
'Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma
sp. red fluorescent protein', *Nat Biotechnol*, 22: 1567-72.

Shaner, N. C., G. G. Lambert, A. Chammas, Y. Ni, P. J. Cranfill, M. A. Baird, B. R. Sell, J. R. Allen, R. N.
Day, M. Israelsson, M. W. Davidson, and J. Wang. 2013. 'A bright monomeric green fluorescent
protein derived from Branchiostoma lanceolatum', *Nat Methods*, 10: 407-9.
Shimshek, D. R., J. Kim, M. R. Hubner, D. J. Spergel, F. Buchholz, E. Casanova, A. F. Stewart, P. H.
Seeburg, and R. Sprengel. 2002. 'Codon-improved Cre recombinase (iCre) expression in the
mouse', *Genesis*, 32: 19-26.
Shoji, W., C. S. Yee, and J. Y. Kuwada. 1998. 'Zebrafish Semaphorin Z1a collapses specific growth cones
and alters their pathway in vivo', *Development*.

Sleight, Sean C., Bryan A. Bartley, Jane A. Lieviant, and Herbert M. Sauro. 2010. 'In-Fusion BioBrick
assembly and re-engineering', *Nucleic Acids Research*, 38: 2624-36.
Sone, T., K. Yahata, Y. Sasaki, J. Hotta, H. Kishine, J. D. Chesnut, and F. Imamoto. 2008. 'Multi-gene
gateway clone design for expression of multiple heterologous genes in living cells: modular
construction of multiple cDNA expression elements using recombinant cloning', *J Biotechnol*,
136: 113-21.
Tamplin, O. J., E. M. Durand, L. A. Carr, S. J. Childs, E. J. Hagedorn, P. Li, A. D. Yzaguirre, N. A. Speck, and
L. I. Zon. 2015. 'Hematopoietic stem cell arrival triggers dynamic remodeling of the perivascular
niche', *Cell*, 160: 241-52.
Tang, D., F. Geng, C. Yu, and R. Zhang. 2021. 'Recent Application of Zebrafish Models in Atherosclerosis
Research', *Front Cell Dev Biol*, 9: 643697.
Varela, M., and A. H. Meijer. 2022. 'A fresh look at mycobacterial pathogenicity with the zebrafish host
model', *Mol Microbiol*, 117: 661-69.
Walton, E. M., M. R. Cronan, R. W. Beerman, and D. M. Tobin. 2015. 'The Macrophage-Specific
Promoter mfap4 Allows Live, Long-Term Analysis of Macrophage Behavior during
Mycobacterial Infection in Zebrafish', *PLoS One*, 10: e0138949.
Wattrus, S. J., and L. I. Zon. 2020. 'A Transgenic System for Rapid Magnetic Enrichment of Rare
Embryonic Cells', *Zebrafish*, 17: 354-57.
Zacharias, D. A., J. D. Violin, A. C. Newton, and R.Y . Tsien. 2002. 'Partitioning of Lipid-Modified
Monomeric GFPs into Membrane Microdomains of Live Cells', *Science*.
Zhu, B., G. Cai, E. O. Hall, and G. J. Freeman. 2007. 'In-fusion assembly: seamless engineering of
multidomain fusion proteins, modular vectors, and mutations', *Biotechniques*, 43: 354-9.